Convergent evolution of desiccation tolerance in grasses

Rose A. Marks[1,2,*], Llewelyn Van Der Pas[3], Jenny Schuster[1,2], and Robert VanBuren[1,2,*]

[1] Department of Horticulture, Michigan State University, East Lansing, MI 48824, USA
[2] Plant Resilience Institute, Michigan State University, East Lansing, MI 48824, USA
[3] Department of Molecular and Cell Biology, University of Cape Town, Rondebosch 7701, South Africa

*Corresponding authors: Rose A. Marks, marksros@msu.edu; Robert VanBuren vanbur31@msu.edu

Abstract

Desiccation tolerance has evolved repeatedly in plants as an adaptation to survive extreme environments. Plants use similar biophysical and cellular mechanisms to survive life without water, but convergence at the molecular, gene, and regulatory levels remains to be tested. Here, we explore the evolutionary mechanisms underlying the recurrent evolution of desiccation tolerance across grasses. We observed substantial overlap in gene duplication and expression associated with desiccation, and syntenic genes of shared origin are activated across species, indicative of parallel evolution. In other cases, similar metabolic pathways are induced, but using different gene sets, pointing towards phenotypic convergence. Species-specific mechanisms supplement these shared core mechanisms, underlining the complexity and diversity of evolutionary adaptations. Our findings provide insight into the evolutionary processes driving desiccation tolerance and highlight the roles of parallel mutation and complementary pathway adaptation in response to environmental challenges.

Introduction

The ability to survive anhydrobiosis or life without water is rare, but widely distributed across life, spanning microbial, animal, and plant lineages. Plants that can tolerate desiccation are known as resurrection plants due to their dramatic ability to revive from an extremely dry state (water potential < -100 MPa or relative water content < 10%) [1]. Desiccation tolerance likely arose in plants during the Ordovician period and is thought to have played a critical role in facilitating the transition from aquatic to terrestrial environments by early land plants [2]. These ancestral mechanisms of anhydrobiosis were retained in many non-seed plants (e.g., mosses, liverworts, ferns, and fern allies) and there is a high frequency of vegetative desiccation tolerance among extant bryophytes and pteridophytes [3]. In contrast, vegetative desiccation tolerance was lost, or suppressed, in the common ancestor of seed plants, presumably in a tradeoff for other systems of drought avoidance and escape, such as annuity, and water transport and retention mechanisms including stomata, vasculature, and roots [4]. Desiccation tolerance then re-evolved convergently in a subset of vascular plants through the rewiring of ancestral anhydrobiosis pathways maintained in seeds, spores, and pollen [5–7]. The retention and re-evolution of desiccation tolerance was likely driven by selective pressures in habitats with extreme water limitation, seasonal drought, and sporadic water availability [8]. Consequently, diverse lineages of resurrection plants can often be found co-occurring in tightly intertwined communities on rocky outcroppings in arid tropical and subtropical regions of the world [3,9].

Despite more than 500 million years of evolution and divergence across extant resurrection plants, multiple biochemical and physiological mechanisms of desiccation tolerance are shared across distantly related species. For example, all surveyed resurrection plants accumulate small non-reducing sugars and other osmoprotectants to vitrify the cytoplasm and safeguard macromolecules during drying. Dramatic shifts in carbohydrate and lipid metabolism, and either degradation or protection of the photosynthetic apparatus are observed across all species during drying [10–13]. Desiccation tolerant plants also leverage robust anti-oxidant scavenging systems, mobilize numerous intrinsically disordered and protective proteins, and have specialized cell wall properties that maximize flexibility and mitigate the mechanical strain of shrinkage [3,13,14]. These broad features of anhydrobiosis are largely conserved across organisms and tissues, but the specific metabolic pathways, regulatory networks, and timing of recovery are notably variable among species [3,15,16] and tissues [17].

The recurrent evolution of desiccation tolerance offers an exciting opportunity to understand how complex traits evolve independently and convergently across broad phylogenetic distances. Convergent evolution can occur via multiple pathways [18], and it is often assumed that when closely related taxa evolve the same trait independently, they do so by leveraging the same genetic pathways. In contrast, distantly related taxa are expected to leverage divergent pathways, due to contrasting genetic starting points, yet arrive at the same convergent phenotype [18,19]. However, these patterns are not always observed in nature, and contradictory examples exist, where distantly related taxa evolve independent but identical mutations and closely related taxa do not [19]. Desiccation tolerance has evolved recurrently and independently across land plants, but the extent of shared genetic adaptations, overlapping pathways, and lineage specific processes across resurrection plants is not known.

Desiccation tolerance has received growing research attention in recent years and several resurrection plants have emerged as models for understanding this remarkable trait [20].

However, most genomic studies of resurrection plants have investigated only a single species in
isolation [17,21–25]. Sister taxa comparisons have explored differences between desiccation tolerant
and related sensitive species [7,26,27], but they have not identified core responses shared among
independent lineages of resurrection plants. Desiccation tolerance is found in ten angiosperms
families, and it is most common in Poaceae, where it evolved independently at least six times
across three subfamilies and dozens of species of grasses [3]. Thus, the grasses are an excellent
system to test if the same pathways, regulatory modules, and mechanisms were recruited
during the recurrent evolution of desiccation tolerance. Here, we explore the evolutionary
mechanisms underlying the recurrent evolution of desiccation tolerance across grasses and
investigate the roles of parallel mutation and complementary pathway adaptation in the
evolution of desiccation tolerance. We present genome assemblies of three resurrection
grasses native to Sub-Saharan Africa along with comprehensive gene expression datasets and
supporting physiological data. We leveraged comparative genomic and transcriptomic
approaches to identify patterns of convergence and divergence across these three species. We
expanded these analyses to other grasses and describe a core signature that defines
desiccation tolerance.
96
97
98 **Results**
99
Comparative genomics of desiccation tolerant grasses
We searched for signatures of convergent evolution across three grasses in separate
Chloridoideae subtribes with two independent origins of desiccation tolerance: *Microchloa caffra*
in Eleusininae and *Oropetium capense* and *Tripogon minimus* in the Tripogoninae subtribe.
These three species have overlapping distributions and tend to co-occur in shallow soils on
rocky outcroppings, locally known as ruwari, across Sub-Saharan Africa (Figure 1). *Microchloa*
caffra Nees, commonly known as pincushion grass, is distributed from Uganda to South Africa
and is the largest of the three species. *Oropetium capense Stapf* is smaller and grows as small,
densely packed tufts on exposed rock surfaces. *Tripogon minimus Steud* is a small but loosely
tufted grass that occurs in shallow soils in both western and southern Africa (Figure 1).
Microchloa caffra plants were collected from Buffelskloof Private Nature Reserve in
Mpumalanga and *O. capense* and *T. minimus* were collected from Swebe Swebe Private
Wildlife Reserve in Limpopo South Africa.
113
114

115
Figure 1. Estimated distribution of the three desiccation tolerant grasses: *Microchloa caffra, Oropetium*
capense, **and** *Tripogon minimus.* Distribution data is taken from GBIF.org (21 November 2023) GBIF Occurrence
Download https://doi.org/10.15468/dl.5jf47y. Collections for the current study were made in Mpumalanga and
Limpopo provinces of South Africa.

We generated reference genome assemblies for each of the three grasses using PacBio
HiFi data. *O. capense* and *T. minimus* are diploid with haploid genome sizes of ~195 Mb based
on flow cytometry, and *M. caffra* is hexaploid with a 1.25 Gb haploid genome. Sequencing reads
were assembled using Hifiasm, producing near complete reference assemblies for *O. capense*
and *T minimus* and a highly contiguous draft assembly of *M. caffra* (Table 1). Six and nine of the
ten chromosomes are assembled telomere to telomere for *T. minimus* and *O. capense*
respectively, and the remaining chromosomes are split into two contigs. The *M. caffra* genome
assembly is more fragmented, with 118 contigs spanning 968 Mb and a contig N50 of 16 Mb.
The monoploid genome size of *M. caffra* is 322 Mb, which is roughly 30% larger than *O.*
*capens*e and *T. minimus* (237 and 223 Mb, respectively), and this expansion is driven largely by
DNA transposons. All three species have a similar proportion of long terminal repeat
retrotransposons (22-27%), but 27% of the *M. caffra* genome is composed of DNA transposons
compared to 12% in *O. capense* and 16% in *T. minimus* (Table 1). Despite this expansion of
transposons in *M. caffra*, the three Chloridoid grasses have very compact genomes compared
to most grasses [28]. We used the MAKER-P pipeline to annotate these three genome

assemblies, with RNAseq data and protein homology as evidence. The *O. capense* and *T.*
*minimu*s genome assemblies have 28,826 and 26,527 gene models respectively, which is
comparable to the well-annotated model resurrection plant *Oropetium thomaeum* (28,835) [22,29].
The *M. caffra* genome assembly has 85,245 gene models, which matches the expectations for a
hexaploid genome (Table 1). We assessed annotation quality using the land plant
(Embryophyta) dataset of Benchmarking Universal Single-Copy Orthologs (BUSCO) and found
between 95.3-97.1% complete proteins across the three grasses, suggesting the genome
assemblies are largely complete and well-annotated (Table 1).

Table 1. Assembly stats of the three resurrection grasses

Assembly stats	*O. capense*	*T. minimus*	*M. caffra*
Ploidy	diploid	diploid	hexaploid
Total assembly size (Mb)	237	223	968
Number of contigs	14	57	118
Contig N50	27,924,228	19,548,099	16,141,787
Contig L50	4	5	22
Number of genes	28,826	26,527	85,245
Complete BUSCO (%)	97.1	95.3	96.2
LTR elements (% of genome):	27.6	22.4	27.6
Ty1/Copia (%)	4.0	5.2	5.5
Gypsy/DIRS1 (%)	21.4	13.2	15.1
DNA transposons (%)	12.2	15.9	27.6
Total repeats (%)	41.7	39.4	56.1

145
We leveraged comparative genomic approaches to identify evolutionary signatures
associated with desiccation tolerance and enable cross-species comparisons of gene
expression data. The three grass genomes are largely collinear with *O. thomaeum,* and have
considerable conserved gene content despite some notable structural rearrangements. Seven
pairs of *O. thomeaum* and *O. capense* chromosomes have near perfect synteny, with
chromosomes 8 and 9 showing a few large-scale inversions, and a telomeric translocation on
chromosome 2 (Supplemental Figure 1). *T. minimus* has similar macrosynteny with *O.*
thomaeum, but has no rearrangements in Chromosome 8. Synteny between *M. caffra* and *O.*
thomaeum is more fragmented because of phylogenetic divergence and polyploidy
(Supplemental Figure 2). Using MCScan with *O. thomaeum* as an anchor, we identified 18,428
syntenic orthologs (syntelogs) shared among the three grasses and *Eragrostis nindensis* [26].
These syntelogs were used to identify patterns of gene duplication associated with desiccation
tolerance across grasses and as anchor points to compare gene expression using genes
shared across species.
To investigate parallel evolution of desiccation tolerance we characterized patterns of
expansion and duplication in gene families with important roles in desiccation tolerance. The
genomes of all sequenced resurrection plants have an expansion of early light induced proteins
(ELIPs) [30], and we observed this same pattern across the desiccation tolerant grasses
investigated here. *O. capense*, *T. minimus*, and *M. caffra* all have massive tandem arrays of 39,
31, and 58 ELIPs respectively, compared to an average of 4 in the genomes of desiccation

sensitive grasses [30]. This expansion of ELIPs is similar to other chlorophyll retaining
(homiochlorophlylus) resurrection plants and is generally higher than chlorophyll degrading
(poikiochlorophyllus) species. We identified the origin of duplicated ELIPs to test if the same or
different ancestral copies were duplicated in each lineage. Tandem duplication of ELIPs within
the Tripogoninae occurred on Chromosome 8, and the Eleusininae and Eragrostidinae subtribe
species have no syntenic ELIPs in this region, despite otherwise high collinearity (Figure 2A).
Most ELIPs in Eleusininae and Eragrostidinae species are found in large tandem arrays on
Chromosome 7, compared to 4-5 ELIPS within Tripogoninae (Figure 2B). This suggests these
grass lineages duplicated ELIPs independently, supporting the convergent evolution of
desiccation tolerance within Chloridoideae.
176

177
Figure 2. Independent tandem gene duplication of ELIPs in different resurrection grass lineages.
Microsyntenic regions of the Chromosome 8 (A) and Chromosome 7 (B) ELIP tandem arrays is shown for
resurrection grasses in the Tropogoninae (*T. minimus*, *O. thomaeum*, and *O. capense*), Eleusininae (*M. caffra*), and
Eragrostidinae (*E. nindensis*) subtribes of Chloridoideae. Syntenic orthologs between the species are shown in beige
and the ELIPs are highlighted in blue. Only a single syntenic region for autopolyploids *M. caffra* (hexaploid) and *E.*
nindensis (tetraploid) is shown for simplicity, but each of the other haplotypes contain the same gene content in these
regions.
185
Searching for overlapping signatures of desiccation tolerance
We subjected replicated sets of *O. capense*, *T. minimus*, and *M. caffra* plants to a
desiccation and rehydration treatment. Plants reached desiccation after ~17-20 days of natural
drying, with a relative water content (RWC) < 10% and photosystem II efficiency (F_v/F_m)
approaching 0.0 (Figure 3A-C). RWC and F_v/F_m recovered within 12 hours of rehydration in *O.*
capense and *T. minimus*, but F_v/F_m took longer to recover in *M. caffra* (Figure 3A). We collected
gene expression data (RNAseq) at comparable timepoints of drying and recovery for each of the
three species.
194

195
Figure 3. Generating physiological comparable desiccation and rehydration timecourses for the three grasses. Relative water content and F_v/F_m of plants during dehydration and rehydration timecourses for (A) *Microchloa caffra,* (B) *Oropetium capense,* and (C) *Tripogon minimus*. Three biological replicates were sampled at each timepoint for each species. Error bars represent standard error of the mean. (D-F) Hierarchical clustering of gene expression across the dehydration timecourse for (D) *Microchloa caffra,* (E) *Oropetium capense,* and (F) *Tripogon minimus*. Samples cluster by sampling time, watering status, RWC, and F_v/F_m as expected.

We quantified RNA abundance and gene expression patterns across the dehydration-rehydration timecourse in each species individually. RNAseq reads were pseudo-aligned to the respective transcripts using Salmon and normalized counts were used for all downstream analyses. In general, gene expression profiles were tightly associated with the hydration status of the plants. Correlation matrices (Figure 3D-F) and principal component analysis (PCA) (Supplemental Figure 3) show tight clustering of samples by time and hydration status, with hydrated, desiccated, and rehydrated samples forming distinct clusters for each species.

Using RWC as a covariate, we identified genes that are significantly up- and down-regulated during dehydration and rehydration processes. Both dehydration and rehydration induced substantial changes in gene expression in all three grasses, with 35-52% of genes showing differential abundance during dehydration and 23-47% during rehydration (Figure 4A). *M. caffra* had significantly more differentially expressed (DE) genes given its hexaploidy (Supplemental Figure 4), but a lower proportion of DE genes compared to the other two grasses

216 (Figure 4A). Broadly, desiccation and rehydration have inverse expression profiles compared to
217 well-watered, and most genes that are upregulated during dehydration are downregulated
218 during rehydration and vice versa (Supplemental Figure 5B).
219

220
Figure 4. Overlapping expression dynamics of conserved genes across species. (A) Boxplot showing the
percentage of DEGs in each species for up- and down-regulated genes in dehydration and rehydration conditions. (B)
Venn diagram showing the number of syntenic orthologs that increased in abundance during dehydration and overlap
across species. The percentage and number of genes in each set are shown. (C) Principal component analysis of z-
score transformed expression values for conserved syntelogs across all three species. Samples are colored by
hydration status in the left panel and by species identity in the right panel.
227

To enable comparisons across species, we leveraged the 18,428 conserved syntelogs
and searched for patterns of convergence and divergence in the expression of these shared
genes. There was considerable overlap in cross-species gene expression, with ~18-24% of all
DE genes showing conserved expression across the three species (Figure 4B and
Supplemental Figure 5A). Dimensionality reduction also supports conserved expression during
desiccation and rehydration as samples cluster primarily by hydration status and secondarily by
species (Figure 4C). We also defined co-expression modules for each species and screened for
shared network level responses across co-expressed genes. Numerous high confidence
modules were defined for each species, and we grouped these into three broad classes based

on the expression pattern of each module: 1) elevated expression in hydrated conditions, 2) elevated expression during dehydration, and 3) elevated expression during rehydration (Figure 5B). We identified substantial overlap across species for modules characterized by elevated expression during dehydration hydrated conditions, but less overlap in modules with high expression during rehydration processes (Figure 5A). We then conducted gene ontology (GO) enrichment analyses of genes in each co-expression module and performed hierarchical clustering on the enrichment scores of these GO terms. Modules clustered by their expression profile rather than species identity, suggesting that hydration status is more predictive of gene expression than species identity (Figure 5C).

Figure 5. Comparative co-expression network dynamics across resurrection grasses. (A) UpSet plots showing the number of shared and unique syntenic orthologs among co-expression modules characterized by elevated expression in hydrated, dehydrated, or rehydrated conditions. (B) Co-expression modules for each species. The X-axis shows the approximate relative water content (RWC) of samples at each timepoint. The module name and total number of genes in the module are listed above. (C) Hierarchical clustering of enriched GO terms for each co-expression module. Secondary clustering performed on modules shows that modules are organized by expression profile rather than species.

Functional characterization of the shared signatures of desiccation tolerance

Our analyses of syntelog expression tested for ancestral conservation or convergence, but it is also possible that different lineages of resurrection plants recruit similar metabolic processes for desiccation with different underlying genes and pathways. To investigate this possibility, we used Kyoto Encyclopedia of Genes and Genomes (KEGG) annotations to assign each gene to a predicted metabolic function and compared the overlap in these functional predictions across species. We detected a substantially higher degree of overlap in KEGG terms across species (~30-40%) compared to only 18-24% of DE syntelogs (Figure 6B and Supplemental Figure 5C). This increased convergence at a metabolic level suggests that while these species do not always leverage parallel gene copies, they induce similar metabolic mechanisms to survive anhydrobiosis.

We further investigated the functional roles of shared gene expression via GO
enrichment analysis and analyses of KEGG annotations. These analyses revealed that multiple
hallmarks of desiccation tolerance were shared across the three resurrection grasses.
Numerous stress responsive genes were upregulated during drying and then dissipated during
rehydration, whereas photosynthetic genes exhibited general downregulation during
dehydration and reactivation during rehydration in all three species. Analysis of enriched GO
terms highlighted the inverse relationship between dehydration and rehydration process as
genes that increased during dehydration clustered tightly with genes that decreased during
rehydration and vice versa (Figure 6A and Supplemental Figure 5D). Enriched GO terms during
dehydration are related to signaling and stress responses (e.g., stress perception and ROS
scavenging activities), developmental regulation (e.g., photoperiodism and germination
processes), cellular reorganization (e.g., lipid droplet formation, vesicle fusion, endocytosis),
and modifications to transcription and translation (e.g., RNA modifications, splicing, and protein
degradation). In contrast, enriched GO terms during rehydration are related to photosynthesis
and metabolism (e.g., fructose biosynthesis, cellulose biosynthesis, and light harvesting),
pigment metabolism (e.g., chlorophyll biosynthesis and anthocyanin metabolism), protein
modification (e.g., protein phosphorylation and proteolysis), and some residual stress response
(e.g., response to cold and non photochemical quenching) (Figure 6A).
285

286
Figure 6. **Overlapping gene functions during desiccation and rehydration in resurrection grasses. (**A)
Hierarchical clustering of p-values for enriched GO terms for each species and condition. Selected GO terms are
highlighted for genes that increased in abundance during dehydration (decreased in abundance during rehydration)
and genes that increased in abundance during rehydration (decreased in abundance during dehydration). Points are
colored by the average enrichment p-value across all species and sized by the number of genes assigned to that GO
term. (B) Venn diagram showing the number of overlapping KEGG terms that increased in abundance during
dehydration. The percentage and number of KEGG terms in each set are shown.
294

KEGG annotations also revealed characteristic desiccation tolerance mechanisms
shared across species. Metabolic pathways associated with photosynthetic energy metabolism
were significantly down regulated in all three grasses. Interestingly, we observed an increase of
malate to pyruvate catalysis with concomitant regeneration of NADPH, which could be related to
NADPH's REDOX potential for antioxidant enzymes such as glutathione reductase. We also
detected noticeable changes to carbohydrate and energy metabolism, including a shift towards
the production of raffinose and stachyose under dehydrating conditions as seen in other
resurrection plants (reviewed in [16]. Central carbohydrate metabolism appeared operational
suggesting that at low water contents, other solvents, such as natural deep eutectic solvents
within the mitochondria may facilitate glycolysis, the TCA, and electron transport [31]. Amino acid
metabolism favored degradative pathways with an increase in endoplasmic reticulum-mediated
ubiquitination and proteolysis, which could be serving a glucogenic role, by converting amino
acids to pyruvate, or by generating an available amino acid pool for the rapid assembly of
thermo- and osmoprotective proteins. While amino acid metabolic pathways were generally
down-regulated, a few important pathways including glutathione metabolism were up-regulated.
Reduced glutathione (GSH) exerts numerous effects in the cell [32] from interaction with
hormones to acting as direct ROS quencher, and maintaining a steady supply of GSH is a
feature all three species share. Lipid metabolism showed a shift towards the production of
glycerolipids and glycerophospholipids which likely supports triacylglycerol phosphatidylcholine
production. The accumulation of phosphatidylcholine may further lead to phosphatidic acid
synthesis, which has been implicated in numerous plant processes from signaling to storage [33–
35]. Pathways involved in the transcription and translation of genetic information also showed an
up-regulation of transcription factors, RNA polymerase, and spliceosome activity, suggesting
that active transcription and RNA processing are still occurring. However, we observed
substantial downregulation of ribosome activity, suggesting that RNA is either differentially
translated or delayed. Upon rehydration, up-regulated processes involved in overall resumption
of normal metabolic activity such as several photosystem I and II proteins, light harvesting
complexes, starch synthesis, and cell wall remodeling such as xyloglucan O-acetyltransferase,
expansin, and pectinesterase were observed.
324
Desiccation tolerance mechanisms are broadly conserved across grasses
Desiccation tolerance evolved independently in at least four subtribes of Chloridiodeae
(Eleusininae, Eragrostidinae, Sporobolinae, and Tripogoninae), and we leveraged comparable
desiccation and rehydration expression datasets to test for patterns of convergence across
grasses more broadly. Building on our detailed comparisons across the three study species, we
expanded our analysis to include publicly available RNAseq samples from *O. thomaeum* [36] and
E. nindensis [26], leveraging syntelogs for cross-species comparisons. Similar to the three grass
comparisons described above, dimensionality reduction across the five species generally
separated samples by hydration status along PC1 and PC2 (Supplemental Figure 6). PCA
provided some degree of separation; however residual heterogeneity, experimental differences,
noise, or species level differences in the datasets might have obscured underlying conserved
biology. To account for this, we employed a topological data analysis (TDA) approach to discern
the underlying structure of the expression datasets. We utilized the Mapper algorithm, which
condenses the dataset into a scalable, navigable representation. The Mapper algorithm is
particularly well-suited for genome scale analyses, as the underlying datasets are often
characterized by high dimensionality and sparsity [37]. For our gene expression data, we

constructed Mapper graphs using a "stress lens" using the well-watered condition as a
reference point. This model represents the baseline for gene expression and we quantified the
residuals or deviation of each sample from the baseline, which represent the degree of water
stress or recovery.
The resultant Mapper graph illustrates a clear topological shape that delineates
desiccation processes across grasses. Each node on the graph represents a cluster of similar
RNAseq samples, and the node color depicts the identity of samples within that cluster.
Connections between nodes signify shared samples among the intersecting clusters.
These graphs reveal a compelling topological depiction of the gene expression variations
induced by water stress across different species. Similar topology was observed for both
targeted comparison of the three focal species (Figure 7A and B) and for the larger dataset
including *E. nindensis* and *O. thomaeum* (Figure 7C and D). In both instances, clear delineation
between samples of different hydration statuses are evident, while the species are intermixed.
Broadly, this supports our finding that the similar ancestral mechanisms are being recruited for
evolving desiccation tolerance across independent grass lineages.

356
Figure 7. Mapper graph showing the network of samples after TDA analysis. Nodes within the graph represent
clusters of RNAseq samples that are akin to one another, with the node color indicating the identity of the samples
contained within. Edges, or the connections between nodes, delineate shared samples across intersecting clusters.
Mapper graphs for the three species comparisons are shown in A and B, and Mapper graphs for the five species

comparisons are shown in C and D. Nodes within the graph are colored by species (A and C) or hydration status (B
and D).
363
Species specific mechanisms underlying desiccation tolerance

365   Despite the considerable overlap in gene expression across all three species, species-
specific processes were also evident. In *M. caffra,* unique antioxidant responses were induced
including glutathione biosynthetic processes, glutamate decarboxylation, and L-ascorbic acid
biosynthesis. Other processes enriched uniquely in *M. caffra* included seed-related terms such
as seed oil body biogenesis and seed maturation. Several GO terms associated with
phytohormones were also uniquely induced in *M. caffra*, including overall ethylene responses
such as S-adenosylmethionine metabolic process, ethylene-activated signaling pathway, and
response to 1-aminocyclopropane-1-carboxylic acid, suggesting that hormonal regulation might
be exerting an effect on the partial breakdown of thylakoids and photosynthetic machinery as
seen in classical senescence [38]. *M. caffra* also exhibited unique lipid, sphingolipid, riboflavin,
and selenocompound metabolism, as well as sesquiterpenoid and tripenoid biosynthesis. *M.*
caffra was the only species to have multiple pathways involved in signal transduction up-
regulated including phospholipase D and calcium signaling. Uniquely down-regulated processes
in *M. caffra* appear to center around arresting growth and development, such as phototropism,
gravitropism, leaf and root morphogenesis, cell wall biogenesis, and regulation of auxin polar
transport. There was also a down-regulation of general amino acid-tRNA aminoacylation and
nitrogen fixation and assimilation.
382   *O. capense* had fewer uniquely enriched processes compared to *M. caffra,* but some
notable patterns were detected. Uniquely upregulated processes in *O. capense* centered
around histone H3 and H4 acetylation, histone H3-K9 demethylation, and histone H2B
ubiquitination. The relative degree of acetylation of histones is directly related to the openness
of chromatin which impacts transcription in specific drought-responsive genes [39]. Histone
demethylation [40] and H2B ubiquitination also regulate drought responsive genes [41].
Interestingly, terms associated with chloroplast mRNA processing, poly(A)+ mRNA export from
the nucleus, ribosome assembly, and regulation of translation were also upregulated in *O.*
capense, suggesting continued translation and active processing of mRNA from both the
chloroplast and nucleus, presumably through increased transcriptional regulation due to histone
modifications. *O. capense* exhibited unique up-regulation of C5-branched dibasic metabolism
and down-regulation within galactose metabolism. Uniquely down-regulated processes in *O.*
capense were minimal, but included regulation of salicylic acid metabolic process and auxin
polar transport. Monoterpenoid biosynthesis was up-regulated for *O. capense* and fatty acid
degradation and steroid hormone biosynthesis was down-regualted. Ferroptosis, an iron-
dependent form of programmed cell death, was exclusively down-regulated in *O. capense*.
398   *T. minimus* also had fewer species-specific processes compared to *M. caffra*. Uniquely
up-regulated processes were centered around response to oxidative stress, peroxisome
organization, and removal of superoxide radicals. *T. minimus* was the only species to show
upregulation of anthocyanin-containing compound biosynthetic process which is a typical
response seen in the homoiochlorophyllous resurrection plants [42]. Similar to the other two
species, regulation of auxin-mediated signaling pathways were down-regulated as was cellular
response to salicylic acid stimulus. Other processes centered around mismatch repair,
chloroplast RNA processing, ribosome biogenesis, and plastid transcription. Phosphonate and

phosphinate, taurine and hypotaurine, and D-amino acid metabolism were exclusively down-
regulated in *T. minimus* whereas retinol metabolism was up-regulated.
Despite the unique GO terms and KEGG pathways identified in each of the focal
species, all three species are responding to drought and desiccation stress by leveraging similar
mechanisms. The processes uniquely enriched in each species are centered around defense
mechanisms, the induction of quiescence, and reduction of normal growth and metabolism
under desiccated conditions. Taken together, the three species appear to share a core set of
gene expression mechanisms which are then supplemented with species-specific modules.
While nuanced variation in metabolism and defense responses are evident, all species exhibit
well known mechanisms of desiccation tolerance.
416
417
Discussion
Our data suggest that the repeated evolution of desiccation tolerance within grasses
occurred by both parallel adaptations in the same genes and complementary modifications to
analogous pathways. We find evidence that core mechanisms of tolerance are shared across
species and are supplemented with species-specific adaptations that contribute to nuanced
variation. Phenotypic and metabolic similarities of anhydrobiosis across resurrection plants have
been observed for decades, but the genetic mechanisms of convergence have been obscured
by a lack of systems-level data and inconsistencies in experimental procedures. Here, we
leveraged large scale genomic and transcriptomic datasets in a replicated and standardized
framework to characterize signatures underlying the convergent evolution of desiccation
tolerance within chloridoid grasses.
Convergent evolution can occur when identical but independent mutations give rise to
the same adaptation (parallel evolution), or when mutations in different genes and pathways
lead to the same emergent phenotype [18,19]. While it is generally accepted that these processes
are driven by exposure to similar selective pressures, the precise evolutionary pathways
involved vary considerably and depend on the genetic constraints of the system. Numerous
convergently evolved traits such as C4 and CAM photosynthesis are highly complex, making
their repeated evolution surprising [43]. In the case of C4 photosynthesis, both parallel mutations
in the same genes and recruitment of unique pathways occurred in different lineages to enable
the emergent C4 phenotype [44]. Desiccation tolerance is similarly complex, involving the
synchronized orchestration of numerous pathways and genes. However, the adaptations
required for desiccation tolerance are sufficiently narrow, such that not any organism can, or
will, evolve desiccation tolerance [4]. The physiological changes that occur during desiccation are
dramatic, and specialized biochemistry and molecular mechanisms are required to survive life
without water. Achieving anhydrobiosis requires tight coordination and orchestration of multiple
physiological processes, and there may be only a few trajectories to evolve this trait. In plants,
desiccation tolerance is likely an ancestral adaptation that evolved during terrestrialization,
subsequently formed the basis of seed pathways, and was later rewired again in vegetative
tissues [5,23,36,45]. While previous studies have found surprisingly little overlap in gene expression
across desiccation tolerant plants [46,47], the specific genetic, biochemical, and physiological traits
required for anhydrobiosis, favor molecular convergence and parallelism, as observed here. We
suspect previous work failed to identify overlapping signatures because of inconsistencies in

experimental design and physiologically incomparable datasets that masked cross-species
conservation.
452   Our systems-level analyses add to the growing literature on the mechanisms of
desiccation tolerance, and many of the patterns observed here corroborate previous findings
[17,23,27,30,46]. We show that desiccation induces a major and reversible shift in gene expression
where normal growth and development are shut down and numerous protective mechanisms
are induced [12,13,48–50]. Similar to recent work in other chloridoid grasses [51], we detected
increased overlap in gene expression during desiccation relative to rehydration, which we
suspect is related to the intense stress of desiccation which requires coordinated response. In
contrast, rehydration leads to recovery and the resumption of species specific processes related
to growth and development. The shared pathways of anhydrobiosis observed in these grasses
represent the deeply conserved architecture of desiccation tolerance, and we suspect many of
these processes are conserved across all tolerant plant lineages. This reflects the relatively
narrow set of regulatory networks and pathways in plants that can enable the evolution of
desiccation tolerance.
465
466
467   **Methods**
468
469   *Field collections, plant growth, and maintenance*
470   Plants for the current study were collected from two research sites in South Africa:
Buffelskloof Nature Reserve in Mpumalanga (-25.30229 S, 030.50631 E) (*Microchloa caffra*)
and Swebe Swebe Private Wildlife Reserve in Limpopo (-23.7949 S, 028.0705 E) (*Oropetium*
capense and *Tripogon minimus*). Voucher specimens of each species were collected, pressed,
and deposited at the National Herbarium of South Africa in Pretoria (specimen numbers:
PRE1004810-0, PRE1004793-0, and PRE1004794-0). Seeds of each species were also
collected and transported to Michigan State University under United States Department of
Agriculture (USDA) permit #537-22-37-10071 and according to the specifications in a Material
Transfer Agreement established between Drs. Jill M. Farrant, Robert VanBuren, and Rose A.
Marks. Seeds were cold stratified at 4ºC for two weeks and then germinated on our standard
propagation mix (50:50 sure-mix:redi-earth) and grown in a climate controlled growth chamber
with a 16 hour photoperiod and internal temperature of 28/18ºC. Six weeks after germination,
individual seedlings were transplanted into separate pots and grown to maturity and a single
plant (genetic line) of each species was selected for downstream experimentation. Seeds of
each genetic line were collected, cold stratified, and ~70 seedlings from each species were
germinated. One seedling from each species was used for genome sequencing and was
transplanted into a larger pot. The remaining seedlings were used for the desiccation and
rehydration timecourses experiments and three seedlings were transplanted into 4" pots. The
three plants in each pot were pooled during sampling and treated as a single biological
replicate. These plants were grown for another two weeks prior to experimental treatments,
during which time they were maintained in constantly hydrated conditions in a growth chamber
set to the conditions described above.
492
Dehydration treatment and sample collection

After ~8 weeks of growth, plants were subjected to dehydration treatment. Prior to
treatment, any emerging reproductive tissues (e.g., panicles) were removed from plants. To
initiate dehydration treatment, plants were watered to full soil saturation and each pot was
weighed to ensure consistency across replicates. Water was then withheld until plants became
completely desiccated (between 2 and 3 weeks depending on the species). Plants were
sampled at targeted hydration states during the process of dehydration, including well watered,
partially dehydrated, fully desiccated, and rehydrated. We used visual cues to direct our
sampling and sampled plants at the first signs of visible leaf curling, partial pigmentation, deep
pigmentation, and full desiccation and validated the hydration status of tissues by measuring
relative water content (RWC). Plants were then rehydrated through a combination of watering
from the base and misting the aerial portions to simulate natural rainfall and sampled 24 and 48
hours post rehydration. We aimed to sample plants at biologically relevant water contents and
therefore directed our sampling by use of visual cues rather than a set number of hours. This
allowed us to compensate for different drying rates across species and plants due to subtle
variation in size, water use efficiency, and relative humidity in the growth chamber.
At each timepoint, we measured the photosynthetic efficiency (F_v/F_m) and RWC, and
harvested tissue for RNAseq. Briefly, F_v/F_m was measured on dark adapted leaves using a Opti-
Sciences OS30p+ chlorophyll fluorometer with the default test parameters. Relative water
content was measured using a set of 10-15 representative leaves from each pot / biological
replicate. Leaf mass was weighed immediately after collection (fresh weight), again after 48
hours submerged in dH20 in darkness at 4ºC (turgid weight), and finally after 48 hours in a 70ºC
drying oven (dry weight). RWC was calculated as (fresh weight - dry weight)/(turgid weight - dry
weight). Tissue for RNAseq was collected by harvesting all the vegetative tissue from each pot
and flash freezing in liquid nitrogen. Tissue samples were stored in a -80ºC freezer prior to
downstream processing.
519
RNA extraction and sequencing
Frozen leaf tissue was ground to a powder by hand in a mortar and pestle with liquid
nitrogen. RNA was extracted from each sample using Spectrum Plant Total RNA kit according
to the manufacturer instructions. Total RNA was then cleaned to remove impurities and
contaminants using Zymo Clean & Concentrator kit. DNAse treatment was carried out during
clean and concentration steps according to manufacturer instructions. Sample concentration
was assessed on a qubit using the RNA broad range reagent set, purity was assessed with a
nanodrop, and RNA integrity was visualized on an agarose gel. RNAseq libraries were
constructed by Novogene following a standard polyA+ enrichment strategy including
fragmentation and cDNA synthesis. The resulting libraries were sequenced on an Illumina
HiSeq 4000 under 150 bp paired end mode
531
High molecular weight DNA extraction, and sequencing
Tissue for whole genome sequencing was collected from a single mature plant of each
species. Healthy green tissue was harvested and flash frozen in liquid nitrogen. Tissue was
ground by hand in a mortar and pestle for >20 minutes to liberate nuclei. Pure, high molecular
weight genomic DNA was extracted by first isolating nuclei with the Circulomics Nuclei Isolation
kit and then extracting DNA with the Circulomics Nanobind Plant Nuclei Big DNA kit. HiFi

libraries were constructed from the Genomic DNA and sequenced at the University of Georgia
Sequencing Core on a PacBio Sequel II machine.
540
Genome assembly
We used flow cytometry to estimate genome sizes (2C DNA values) for the three
grasses. Healthy leaf tissue was harvested from each genotype. Nuclei were isolated and
stained according to standard protocols. The stained nuclei were then run on a BD Accuri™ C6
Plus Flow Cytometer at Plantploidy.com. *Hosta plantaginea* was used as an internal reference.
We built reference genomes for each species using high fidelity (HiFi) PacBio long read data. In
total, 70.1 Gb of HiFi reads were generated for *M. caffra*, 15.9 Gb for *O. capense,* and 20.2 Gb
for *T. minimus*, representing 56, 82, and 103 x genome coverage for each species respectively.
K-mer analysis revealed that *O. capense* and *T. minimus* have low within genome
heterozyogisity and *M. caffra* is a highly heterozygous autopolyploid [52]. PacBio reads were
assembled using hifiasm (v 0.18) [53,54] wth default settings for *O. capense* and *T. minimus* and
the number of haplotypes was set to 6 for *M. caffra* (flag: --n-hap 6). The resulting assemblies
were highly contiguous with six and nine of the ten chromosomes assembled telomere to
telomere for *T. minimus* and *O. capense* respectively, and 118 contigs across 968 Mb with an
N50 of 16 Mb for *M. caffra* (Table 1). Raw assemblies were filtered for non-plant contigs using a
representative microbial database with BLAST [55]. Full length chloroplast and mitochondrial
genomes were identified and retained, and any additional partial or rearranged organelle
genomes were removed.
559
Genome annotation
A library of repetitive elements was constructed for each of the three grass genomes
using the EDTA package (v2.0.0) [56]. EDTA comprehensively identifies DNA-based transposable
elements using HelitronScanner [57], and LTR retrotransposons using LTR_FINDER [58] and
LTRharvest [59]. Protein coding genes were annotated using the MAKER-P pipeline (v2.31.10) [60]
with the following sets of input data for training. Transcript evidence was generated using the
dehydration-rehydration timecourse RNAseq data from leaf tissue of each species described
below. Raw RNAseq reads were quality trimmed using fastp (v 0.23) [61] and aligned to the
unmasked genomes using the splice aware alignment program STAR (v2.6) [62]. A set of non-
overlapping transcripts was identified from the aligned data using StringTie (v1.3.4) [63] with
default parameters. The resulting gff files were used as transcript evidence for MAKER. The
same protein evidence was used as training for each of the three grasses and this includes the
full annotations of *Oryza sativa* [64], *Arabidopsis thaliana* [65], *Oropetium thomaeum* [22,29], and
Eragrostis tef [66]. These datasets were used as input for MAKER and we utilized SNAP [67] and
Augustus (version 3.0.2) [68] for *ab initio* gene prediction, performing two rounds of iterative
training to refine our models. To filter out repetitive element-derived proteins, we used BLAST
using a non-redundant transposase library against the raw gene models produced by MAKER.
We assessed the completeness of our assembly using the plant-specific embryophyte set of
Benchmarking Universal Single-Copy Orthologs (BUSCO v.2) [69]. These high-confidence gene
models were used for all downstream analyses.
580
Comparative genomics

The three desiccation tolerant grass genomes were compared to each other and other
Chloridoid grasses using the MCScan toolkit (v1.1) [70] implemented in python
[https://github.com/tanghaibao/jcvi/wiki/MCscan-(Python-version)]. Syntenic orthologs were
identified across species using the chromosome-scale *Oropetium thomeaum* genome as an
anchor. Syntenic blocks were identified using gene models aligned using LAST with a minimum
of five overlapping syntenic genes. The macrosyntenic dot plots, histograms of depth, and
microsynteny plots were generated using the python version of MCScan. A set of 18,428
conserved syntenic orthologs across all five desiccation tolerant grasses was created and used
for downstream comparative genomic and cross-species transcriptomic analyses.
591
Transcriptomic analyses
RNA sequencing reads were processed following a pipeline developed by the VanBuren
Lab (https://github.com/pardojer23/RNAseqV2). Briefly, sequence read quality was assessed
with fastQC (v 0.23) and reads were trimmed with trimmomatic (v 0.38) [71] to remove adapters
and low quality bases. Trimmed reads were sudo-aligned to reference genomes using Salmon
(v 1.9.0) [72], and the resulting quantification files were processed with tximport (v 3.18) [73] and to
generate normalized expression matrices of transcripts per million (TPM). A Principal
Component Analysis (PCA) was used to visualize replicate and sample relationships within
each species using the respective TPM expression values. A cross-species PCA was performed
using the TPM matrix of conserved syntenic orthologs across all species. For polyploid species
(*M. caffra* and *E. nindensis*) we summed the TPM values for all homeologs to each *O.*
thomaeum syntelog. We then computed the Z-score of the TPM for each syntenic ortholog in
each sample and ran PCA on the matrix of Z-scores of each syntenic ortholog across all
timepoints and species.
606
Differentially expressed genes
Differentially expressed genes (DEGs) were identified independently for each species
with DEseq2 R package (v 1.42.0) [74]. Briefly, transcript abundance estimates from Salmon were
imported into DEseq2 using tximport to generate counts matrices. Count matrices were
normalized, and hierarchical clustering was conducted for basic quality control and visualization
of relationships across experimental timepoints and biological replicates. We tested multiple
models for differential expression in DEseq2, including models that identified DEGs by pairwise
comparisons of each timepoint against well-watered, and models that used the continuous
variables of RWC or F_v/F_mas covariates. DEGs identified by pairwise comparisons were
summarized into a nonredundant list of up and down regulated genes during dehydration and
rehydration. DEGs identified using the continuous variables are based on a significant linear
association (positive or negative) with RWC or F_v/F_m. When identifying DEGs, we included the
term "process" in our model to differentiate between dehydration and rehydration processes. To
select the best performing model, we quantified similarities and differences in the number and
identity of DEGs defined by each model. There was a high degree of overlap in genes identified
by all three models. Ultimately, we selected the model based on RWC because it performed
well and is easily comparable across experiments regardless of sampling time, consistency
across replicates, or differences in experimental design. These analyses produced species-
specific lists of DEGs during dehydration and rehydration with significant (FDR adjusted P-value

<0.05) associations with RWC. Log2foldchange values are reported for one unit change in
RWC.
To gain insight into possible similarities and differences among the study species, we
looked at the overlap in DE syntenic orthologs. To do so, we used venn diagrams to identify the
shared syntelogs in up- and down-regulated genes during both dehydration and rehydration
across the three study species. We then conducted targeted analyses to look at the functional
roles of the shared DE syntelogs. We also investigated the functional signatures of differentially
abundant transcripts that were unique to each species. To do so we pulled the lists of DE
syntelogs that were only found in one of the focal species.
635
Functional annotation of DEGs
We annotated DE syntelogs with KEGG and GO terms to describe generalized
metabolic and cellular processes responses shared across in the three study species. KEGG
annotations were generated for each species using BLASTKoala
(https://www.kegg.jp/blastkoala/) on the complete set of annotated peptide sequences. These
KEGG terms were then assigned to syntenic orthologs and DE KEGG terms that were shared
across all three species during dehydration and rehydration were identified and plotted using
venn diagrams. These shared DE KEGG terms were used to generate metabolic pathway maps
using the reconstruct function of the KEGGmapper tool
(https://www.genome.jp/kegg/mapper/color.html) for up- and down-regulated terms in
dehydration and rehydration. This returned a list of syntelogs per metabolic pathway and Brite
descriptions. The difference between the number of syntelogs assigned to each pathway for up
and down regulated genes was computed. The list was then sorted to determine pathways that
were primarily up or down regulated. Next, the assigned KO numbers were paired with
quantitative data on gene expression to identify which pathways were active at various
timepoints. To summarize patterns, genetic information processes (transcription, translation,
folding, sorting, and degradation, replication and repair, and processing in viruses) were
grouped together. Similarly Environmental information processing (membrane transport and
signal transduction) and cellular processes (transport and catabolism, cell growth and death,
cellular community- prokaryote and eukaryotes, and cell motility) were grouped together.
Pathways assigned to Organismal systems and Human disease were ignored. KEGG
annotation is not without its limits as single KEGG identifiers can be present in multiple
pathways.
GO terms were assigned through homology with the well annotated genome of sister
species *Oropetium thomaeum* [24]. This was done through a BLASTP (v 2.14.0) [75] search of all *O.*
thomaeum protein sequences against the protein sequences of each study species. Parameters
were set to return the single best match for each peptide and an e-value cutoff of 1e-10. We
assigned the GO terms from *O. thomaeum* to the homologous genes in our target species. We
then used TopGO R package (v 2.54.0) to identify significantly enriched GO terms (P-
value<0.05) within sets of DEGs for up- and down-regulated genes during dehydration and
rehydration in each target species.
667
Co-expression analyses

To complement the above analyses, we generated co-expression networks using
Weighted Gene Co-expression Network Analysis (WGCNA) R package (v1.7) [76]. While DE
analyses can be informative to identify and describe overarching patterns and large shifts in the
data, more nuanced patterns of gene expression can be obscured. To investigate the more
subtle temporal changes in gene expression, we used co-expression analyses to identify
modules of co-expressed genes for each species.
For each species, we created a signed co-expression network using WGCNA. Each
dataset was filtered to remove genes with no expression. To construct a weighted co-
expression network, we determined a soft thresholding power for each dataset. This power was
chosen to satisfy WGCNA's assumption that a weighted co-expression network is scale-free. An
adjacency matrix, representing the strength of connections between genes in the network, was
constructed for each network using the soft thresholding power. For module detection, this
matrix was then converted to a topological overlap matrix (TOM) and hierarchal clustering was
used on the TOM to group genes into modules based on similar expression patterns.
Additionally, we calculated connectivity of each gene within its network and its assigned module
using WGCNA's network analysis functions.
We identified shared and species-specific co-expressed genes using UpSet plots [77]. For
all co-expressed genes, we identified the syntenic orthologs and computed the overlap across
species. We combined all modules for a species that showed increased expression during
dehydration, during rehydration, and under non-stressed conditions. We identified the sets of
shared syntelogs as well as those that were only found in a single species. We then ran GO
enrichment analysis of these sets of shared and unique co-expressed genes.
691
Topological data analysis
We employed a topological data analysis (TDA) approach following the pipeline
described at https://github.com/PlantsAndPython/plant-evo-mapper to discern the underlying
structure of the expression datasets. We utilized the Mapper algorithm, which condenses the
dataset into a scalable, navigable representation. The Mapper algorithm is particularly well-
suited for genome scale analyses, as the underlying datasets are often characterized by high
dimensionality and sparsity. For our gene expression data, we constructed Mapper graphs
using a "stress lens" formulated by applying a linear model using the well-watered condition as
a reference point. This model represents the baseline for leaf expression and we quantified the
residuals or deviation of each sample from the baseline, which represents the degree of water
stress or recovery. For the mapper graph, we specified different intervals and overlap for the 3
species comparisons and the 5 species comparisons. For the three species, we specified 110
intervals with a 90% overlap and for the 5 species comparison we specified 120 intervals with
95% overlap.
706

Acknowledgements: This work was funded by NSF IOS-PRFB-1906094 to RAM, DBI-
2213983 to RAM and RV, and MCB-1817347 to RV. We thank landowners and stewards Pieter
and Jennie Pretorius, Ken Maude, Pieter Vervoort, and Wayne and Messiah Mudenda for
assistance with field logistics and permission to collect plants. We thank the South African
National Herbarium, Pretoria for assistance identifying and vouchering of specimens and the
United States Department of Agriculture for providing import permits.

Author contributions: RAM and RV conceived of the study. RAM collected and curated data.
RAM, LVP, JS, and RV conducted data analyses and contributed to data interpretation and
conceptual framing of the manuscript. RAM and RV drew the figures. RAM, LVP, and RV wrote
the manuscript. All authors edited and reviewed the manuscript.

Data availability: Sequence data associated with this study are deposited at NCBI under
BioProject PRJNA1044305 and BioSamples SAMN38380430-92. Genome assemblies are
hosted on CoGe (https://genomevolution.org/) under the following IDs: 65089 (*T. minimus*),
65046 (*O. capense*), and 64494 (*M. caffra*). Metadata and other data summaries associated
with this study are provided at Dryad https://doi.org/10.5061/dryad.kh18932c4.

References:

1.   Bewley, J. D. Physiological Aspects of Desiccation Tolerance. *Annu. Rev. Plant Physiol.*

30, 195–238 (1979).

2.   Oliver, M. J., Tuba, Z. & Mishler, B. D. The evolution of vegetative desiccation tolerance in

land plants. *Plant Ecol.* **151**, 85–100 (2000).

3.   Marks, R. A., Farrant, J. M., Nicholas McLetchie, D. & VanBuren, R. Unexplored

dimensions of variability in vegetative desiccation tolerance. *Am. J. Bot.* (2021)

doi:10.1002/ajb2.1588.

4.   Alpert, P. Constraints of tolerance: why are desiccation-tolerant organisms so small or

rare? *J. Exp. Biol.* **209**, 1575–1584 (2006).

5.   VanBuren, R. Desiccation tolerance: Seedy origins of resurrection. *Nature Plants* **3**, 17046

(2017).

6.   Costa, M. C. D. *et al.* Key genes involved in desiccation tolerance and dormancy across life

forms. *Plant Sci.* (2016) doi:10.1016/j.plantsci.2016.02.001.

7.   VanBuren, R. *et al.* Desiccation Tolerance Evolved through Gene Duplication and Network

Rewiring in Lindernia. *Plant Cell* **30**, 2943–2958 (2018).

8.   Alpert, P. The limits and frontiers of desiccation-tolerant life. *Integr. Comp. Biol.* **45**, 685–

695 (2005).

9.   Porembski, S. & Barthlott, W. *Granitic and gneissic outcrops (inselbergs) as centers of*

diversity for desiccation-tolerant vascular plants. vol. 151 19–28

https://link.springer.com/content/pdf/10.1023%2FA%3A1026565817218.pdf (2000).

10. Crowe, J. H., Carpenter, J. F. & Crowe, L. M. THE ROLE OF VITRIFICATION IN

ANHYDROBIOSIS. *Annu. Rev. Physiol.* **60**, 73–103 (1998).

11. Hoekstra, F. A., Golovina, E. A. & Buitink, J. Mechanisms of plant desiccation tolerance.

Trends Plant Sci. **6**, 431–438 (2001).

12. Dinakar, C., Djilianov, D. & Bartels, D. Photosynthesis in desiccation tolerant plants: Energy

metabolism and antioxidative stress defense. *Plant Science* vol. 182 29–41 Preprint at

https://www.ncbi.nlm.nih.gov/pubmed/22118613 (2012).

13. Oliver, M. J. *et al.* Desiccation Tolerance: Avoiding Cellular Damage During Drying and

Rehydration. *Annu. Rev. Plant Biol.* (2020) doi:10.1146/annurev-arplant-071219-105542.

14. Moore, J. P., Le, N. T., Brandt, W. F., Driouich, A. & Farrant, J. M. Towards a systems-

based understanding of plant desiccation tolerance. *Trends in Plant Science* vol. 14 110–

117 Preprint at https://www.ncbi.nlm.nih.gov/pubmed/19179102 (2009).

15. Farrant, J. M. & Hilhorst, H. Crops for dry environments. *Curr. Opin. Biotechnol.* **74**, 84–91

(2021).

16. Dace, H. J. W. *et al.* A Horizontal View of Primary Metabolomes in Vegetative Desiccation

Tolerance. *bioRxiv* 2023.02.10.528018 (2023) doi:10.1101/2023.02.10.528018.

17. VanBuren, R. *et al.* Core cellular and tissue-specific mechanisms enable desiccation

tolerance in Craterostigma. *Plant J.* **114**, 231–245 (2023).

18. Stern, D. L. The genetic causes of convergent evolution. *Nat. Rev. Genet.* **14**, 751–764

(2013).

19. Arendt, J. & Reznick, D. Convergence and parallelism reconsidered: what have we learned

about the genetics of adaptation? *Trends Ecol. Evol.* **23**, 26–32 (2008).

20. Tebele, S. M., Marks, R. A. & Farrant, J. M. Two Decades of Desiccation Biology: A

Systematic Review of the Best Studied Angiosperm Resurrection Plants. *Plants* **10**, (2021).

21. Xiao, L. *et al.* The resurrection genome of *Boea hygrometrica* : A blueprint for survival of

dehydration. *Proceedings of the National Academy of Sciences* **112**, 5833–5837 (2015).

22. VanBuren, R. *et al.* Single-molecule sequencing of the desiccation-tolerant grass

Oropetium thomaeum. *Nature* **527**, 508–511 (2015).

23. Costa, M.-C. D. *et al.* A footprint of desiccation tolerance in the genome of Xerophyta

viscosa. *Nature Plants* **3**, 17038 (2017).

24. VanBuren, R., Wai, C. M., Keilwagen, J. & Pardo, J. A chromosome☐scale assembly of the

model desiccation tolerant grass Oropetium thomaeum. *Plant Direct* (2018).

25. VanBuren, R. *et al.* Extreme haplotype variation in the desiccation-tolerant clubmoss

Selaginella lepidophylla. *Nat. Commun.* **9**, 13 (2018).

26. Pardo, J. *et al.* Intertwined signatures of desiccation and drought tolerance in grasses.

Proc. Natl. Acad. Sci. U. S. A. **117**, 10079–10088 (2020).

27. Montes, R. A. C. *et al.* A comparative genomics examination of desiccation tolerance and

sensitivity in two sister grass species. *Proceedings of the National Academy of Sciences*

119, e2118886119 (2022).

28. Marks, R. A., Hotaling, S., Frandsen, P. B. & VanBuren, R. Representation and

participation across 20 years of plant genome sequencing. *Nat Plants* **7**, 1571–1578

(2021).

29. VanBuren, R., Wai, C. M., Keilwagen, J. & Pardo, J. A chromosome-scale assembly of the

model desiccation tolerant grass Oropetium thomaeum. *Plant Direct* **2**, e00096 (2018).

30. VanBuren, R., Pardo, J., Man Wai, C., Evans, S. & Bartels, D. Massive Tandem

Proliferation of ELIPs Supports Convergent Evolution of Desiccation Tolerance across Land

Plants. *Plant Physiol.* **179**, 1040–1049 (2019).

31. du Toit, S. F., Bentley, J. & Farrant, J. M. Chapter Nine - NADES formation in vegetative

desiccation tolerance: Prospects and challenges. in *Advances in Botanical Research* (eds.

Verpoorte, R., Witkamp, G.-J. & Choi, Y. H.) vol. 97 225–252 (Academic Press, 2021).

32. Hasanuzzaman, M., Nahar, K., Anee, T. I. & Fujita, M. Glutathione in plants: biosynthesis

and physiological role in environmental stress tolerance. *Physiol. Mol. Biol. Plants* **23**, 249–

268 (2017).

33. Gasulla, F. *et al.* The role of lipid metabolism in the acquisition of desiccation tolerance in

Craterostigma plantagineum: a comparative approach. *Plant J.* **75**, 726–741 (2013).

34. Sun, M. *et al.* Phosphatidylcholine Enhances Homeostasis in Peach Seedling Cell

Membrane and Increases Its Salt Stress Tolerance by Phosphatidic Acid. *Int. J. Mol. Sci.*

23, (2022).

35. Zhang, X., Gao, Y., Zhuang, L. & Huang, B. Phosphatidic acid priming-enhanced heat

tolerance in tall fescue (Festuca arundinacea) involves lipidomic reprogramming of lipids for

membrane stability and stress signaling. *Plant Growth Regul.* **99**, 527–538 (2023).

36. VanBuren, R. *et al.* Seed desiccation mechanisms co-opted for vegetative desiccation in

the resurrection grass *Oropetium thomaeum. Plant Cell Environ.* **40**, 2292–2306 (2017).

37. Palande, S. *et al.* The topological shape of gene expression across the evolution of

flowering plants. *bioRxiv* 2022.09.07.506951 (2022) doi:10.1101/2022.09.07.506951.

38. Lim, P. O., Kim, H. J. & Nam, H. G. Leaf senescence. *Annu. Rev. Plant Biol.* **58**, 115–136

(2007).

39. Li, S. *et al.* Histone Acetylation Changes in Plant Response to Drought Stress. *Genes* **12**,

(2021).

40. Wang, Q. *et al.* JMJ27-mediated histone H3K9 demethylation positively regulates drought-

stress responses in Arabidopsis. *New Phytol.* **232**, 221–236 (2021).

41. Ma, S. *et al.* Reversible Histone H2B Monoubiquitination Fine-Tunes Abscisic Acid

Signaling and Drought Response in Rice. *Mol. Plant* **12**, 263–277 (2019).

42. Farrant, J. M. A comparison of mechanisms of desiccation tolerance among three

angiosperm resurrection plant species. *Plant Ecol.* (2000).

43. Heyduk, K., Moreno-Villena, J. J., Gilman, I. S., Christin, P.-A. & Edwards, E. J. The

genetics of convergent evolution: insights from plant photosynthesis. *Nat. Rev. Genet.* **20**,

485–493 (2019).

44. Christin, P.-A., Salamin, N., Savolainen, V., Duvall, M. R. & Besnard, G. C4 Photosynthesis

evolved in grasses via parallel adaptive genetic changes. *Curr. Biol.* **17**, 1241–1247 (2007).

45. Illing, N., Denby, K. J., Collett, H., Shen, A. & Farrant, J. M. The signature of seeds in

resurrection plants: A molecular and physiological comparison of desiccation tolerance in

seeds and vegetative tissues. *Integr. Comp. Biol.* **45**, 771–787 (2005).

46. Alejo-Jacuinde, G., González-Morales, S. I., Oropeza-Aburto, A., Simpson, J. & Herrera-

Estrella, L. Comparative transcriptome analysis suggests convergent evolution of

desiccation tolerance in Selaginella species. *BMC Plant Biol.* **20**, 468 (2020).

47. Ostria-Gallardo, E. *et al.* A comparative gene co-expression analysis using self-organizing

maps on two congener filmy ferns identifies specific desiccation tolerance mechanisms

associated to their microhabitat preference. *BMC Plant Biol.* **20**, 56 (2020).

48. Vicré, M., Farrant, J. M. & Driouich, A. Insights into the cellular mechanisms of desiccation

tolerance among angiosperm resurrection plant species. *Plant, Cell and Environment* vol.

27 1329–1340 Preprint at (2004).

49. Gechev, T., Lyall, R., Petrov, V. & Bartels, D. Systems biology of resurrection plants. *Cell.*

Mol. Life Sci. (2021) doi:10.1007/s00018-021-03913-8.

50. Xu, X. *et al.* Molecular insights into plant desiccation tolerance: transcriptomics, proteomics

and targeted metabolite profiling in Craterostigma plantagineum. *Plant J.* (2021)

doi:10.1111/tpj.15294.

51. St Aubin, B., Wai, C. M., Kenchanmane Raju, S. K., Niederhuth, C. E. & VanBuren, R.

Regulatory dynamics distinguishing desiccation tolerance strategies within resurrection

grasses. *Plant Direct* **6**, e457 (2022).

52. Marks, R. A. *et al.* Polyploidy enhances desiccation tolerance in the grass Microchloa

caffra. *bioRxiv* 2023.06.20.545583 (2023) doi:10.1101/2023.06.20.545583.

53. Cheng, H., Concepcion, G. T., Feng, X., Zhang, H. & Li, H. Haplotype-resolved de novo

assembly using phased assembly graphs with hifiasm. *Nat. Methods* **18**, 170–175 (2021).

54. Cheng, H. *et al.* Haplotype-resolved assembly of diploid genomes without parental data.

Nat. Biotechnol. **40**, 1332–1335 (2022).

55. Wheeler, D. L. *et al.* Database resources of the National Center for Biotechnology

Information. *Nucleic Acids Res.* **36**, D13–21 (2008).

56. Ou, S. *et al.* Benchmarking transposable element annotation methods for creation of a

streamlined, comprehensive pipeline. *Genome Biol.* **20**, 275 (2019).

57. Xiong, W., He, L., Lai, J., Dooner, H. K. & Du, C. HelitronScanner uncovers a large

overlooked cache of Helitron transposons in many plant genomes. *Proc. Natl. Acad. Sci. U.*

S. A. **111**, 10263–10268 (2014).

58. Xu, Z. & Wang, H. LTR_FINDER: an efficient tool for the prediction of full-length LTR

retrotransposons. *Nucleic Acids Res.* **35**, W265–8 (2007).

59. Ellinghaus, D., Kurtz, S. & Willhoeft, U. LTRharvest, an efficient and flexible software for de

novo detection of LTR retrotransposons. *BMC Bioinformatics* **9**, 18 (2008).

60. Campbell, M. S. *et al.* MAKER-P: a tool kit for the rapid creation, management, and quality

control of plant genome annotations. *Plant Physiol.* **164**, 513–524 (2014).

61. Chen, S., Zhou, Y., Chen, Y. & Gu, J. fastp: an ultra-fast all-in-one FASTQ preprocessor.

Bioinformatics **34**, i884–i890 (2018).

62. Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21 (2013).

63. Pertea, M. *et al.* StringTie enables improved reconstruction of a transcriptome from RNA-

seq reads. *Nat. Biotechnol.* **33**, 290–295 (2015).

64. International Rice Genome Sequencing Project. The map-based sequence of the rice

genome. *Nature* **436**, 793–800 (2005).

65. Cheng, C.-Y. *et al.* Araport11: a complete reannotation of the Arabidopsis thaliana

reference genome. *Plant J.* **89**, 789–804 (2017).

66. VanBuren, R. *et al.* Exceptional subgenome stability and functional divergence in the

allotetraploid Ethiopian cereal teff. *Nat. Commun.* **11**, 884 (2020).

67. Korf, I. Gene finding in novel genomes. *BMC Bioinformatics* **5**, 59 (2004).

68. Stanke, M. & Waack, S. Gene prediction with a hidden Markov model and a new intron

submodel. *Bioinformatics* **19 Suppl 2**, ii215–25 (2003).

69. Simão, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V. & Zdobnov, E. M.

BUSCO: assessing genome assembly and annotation completeness with single-copy

orthologs. *Bioinformatics* **31**, 3210–3212 (2015).

70. Wang, Y. *et al.* MCScanX: a toolkit for detection and evolutionary analysis of gene synteny

and collinearity. *Nucleic Acids Res.* **40**, e49 (2012).

71. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina

sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

72. Patro, R., Duggal, G., Love, M. I., Irizarry, R. A. & Kingsford, C. Salmon provides fast and

bias-aware quantification of transcript expression. *Nat. Methods* **14**, 417–419 (2017).

73. Soneson, C., Love, M. I. & Robinson, M. D. Differential analyses for RNA-seq: transcript-

level estimates improve gene-level inferences. *F1000Res.* **4**, 1521 (2015).

74. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

75. Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. Basic local alignment

search tool. *J. Mol. Biol.* **215**, 403–410 (1990).

76. Langfelder, P. & Horvath, S. WGCNA: an R package for weighted correlation network

analysis. *BMC Bioinformatics* **9**, 559 (2008).

77. Conway, J. R., Lex, A. & Gehlenborg, N. UpSetR: an R package for the visualization of

intersecting sets and their properties. *Bioinformatics* **33**, 2938–2940 (2017).