TSAR, "Thermal Shift Analysis in R", identifies endogenous molecules that interact with HIV-1 capsid hexamers

William M. McFadden[1,2,†], Xinlin Gao[1,2,†], Zhijiang Ye[1,2,†], Xin Wen[1,2], Zach C. Lorson[1,2], Huanchun Zheng[1,2], Jessica Fahim[1,2], Andres E. Castaner[1,2], Karen A. Kirby[1,2], and Stefan G. Sarafianos[1,2,*]

[1]Center for ViroScience and Cure, Laboratory of Biochemical Pharmacology, Department of Pediatrics, Emory University School of Medicine, Atlanta, GA

[2]Children's Healthcare of Atlanta, Atlanta, GA

[†]These authors contributed equally

*Correspondence (stefanos.sarafianos@emory.edu)

Abstract

The thermal shift assay (TSA) is a versatile biophysical technique for studying protein-

ligand interactions *in vitro*. Here, we report a free, open-source software tool, TSAR ("Thermal

Shift Analysis in R"), to expedite the analysis of TSA data. The TSAR package incorporates

multiple workflows that facilitate TSA analyses, returns publication-ready graphics, and includes

an   optional   graphic   user   interface.   The   package   is   available   at

https://bioconductor.org/packages/TSAR/. Applying TSAR, we screened two chemical libraries

and found multiple molecules that potentially interact *in vitro* with the capsid protein (CA) of

human immunodeficiency virus type 1 (HIV-1). First, a library of vitamins exemplifies the

different graphic outputs of TSAR, and we report a change in the 50% melting temperature (ΔT_m)

for folic acid-treated CA hexamers (CA_{HEX}). Since HIV-1 CA_{HEX} interacts with host-derived acids

like inositol hexaphosphate (IP6) or dNTPs, a second library was screened containing 96 organic,

acidic metabolites; multiple anionic ligands caused a ΔT_m for CA_{HEX}. Subsequent investigation of

these interactions includes biolayer interferometry, antiviral activity against pseudotyped HIV-1,

and endogenous reverse transcriptase assays that were used to validate and investigate the

biological impact of these native ligands that thermally-stabilize CA_{HEX}. One compound hit, gallic

acid, exhibited anti–HIV-1 activity as previously reported, and we show interacts with CA_{HEX} as

a potentially novel mechanism. Overall, the TSAR package facilitated quick analysis of TSA data

from multiple libraries to help identify a biologically relevant hit, gallic acid, as a molecule that

can inhibit HIV-1 replication and targets CA_{HEX}.

Importance

The TSAR package is freely available (AGPL-3) and is designed for both experienced or

new R users, having command-line code for handling large and challenging datasets while also

including an optional GUI that enables easy use by non-programmers. This is the first TSA analysis

program written in R, a free and open-source language; TSAR simplifies TSA analysis while

maintaining diverse visualization options for small-to-large libraries and multidimensional

analysis. Additionally, we report multiple endogenous metabolites that potentially interact with

the HIV-1 capsid protein hexamers (CA_{HEX}) *in vitro*, including folic acid, gallic acid, and multiple

others, primarily from the tricarboxylic acid (TCA) cycle. Various methods validate gallic acid

interactions with CA_{HEX}, leading to a novel suggested mechanism of action, and in-line with

previous reports, this metabolite has potential for natural-based treatments of HIV-1. Further, we

advance the understanding of the potential mechanism(s) for HIV-1 inhibition by gallic acid

treatment.

Keywords

thermal shift assay, differential scanning fluorimetry, human immunodeficiency virus, capsid,

endogenous metabolites, gallic acid, folic acid

Introduction

The thermal shift assay (TSA), alternatively "differential scanning fluorimetry" (DSF), is an *in vitro* technique to study the thermal stability of a protein or biological complex with multiple applications (1-4). During TSA experiments, samples are heated at a constant rate to track the denaturation of protein(s) of interest (Supplementary Figure 1A). Proteins that contain internal aromatic residues, like tryptophan, can be tracked without a dye by measuring the change in residue fluorescence (4); however, TSA experiments often utilize dyes that fluoresce in hydrophobic environments, like SYPRO™ Orange, to follow the exposure of the internal, hydrophobic residues as denaturation progresses (2, 5). Plotting the temperature compared to the fluorescence reading gives an individual melting profile for the protein or complex measured. The melting curve of uncharacterized proteins can be unpredictable, and testing of various dyes can help validate the experiment (6). For folded and globular proteins, hydrophobic residues are typically internal and inaccessible to the solvent, making TSA a useful tool to assess if a batch of proteins is properly folded and consistently prepared (7, 8).

For most samples that are properly folded, the thermal profile is sigmoidal, and a melting temperature (T_m) value can be calculated, representing the temperature where 50% of the protein is unfolded (2, 7) (Supplementary Figure 1A). Curves that are not properly sigmoidal will need to be omitted or assessed independently for melting temperature(s). The T_m value is used to compare the thermal stability of proteins under various conditions, such as selecting optimal storage buffers and salts or for comparing the stability of mutant proteins (9, 10). Of note, intrinsically disordered proteins do not typically have a sigmoidal curve and lack T_m value at room temperature, as they are unfolded by definition (11, 12). TSA is also useful for screening chemical libraries to identify potential binding partners, since protein•ligand and protein•protein interactions that form

additional intramolecular bonds and thus require more energy, or a higher temperature, to denature

the sample compared to an unliganded sample (1, 2, 13-15). The "thermal shift", determined by

the change in the T_m (ΔT_m), is determined by calculating the difference between the T_m of treated

and untreated sample(s). Thus, TSA is used as a proxy for *in vitro* ligand binding to purified protein

targets within screens for molecular interactors such as inhibitors, natural ligands, and cofactors.

The TSA technique is used extensively in labs that focus on biological research, as it

requires only a qPCR machine for data acquisition and one of many commercially-available dyes,

like SYPRO™ Orange, once a protein or biological complex is purified (2, 6). The required

volume and concentration of the sample are quite low, typically under 25 μL and below 20 μM

per sample, although multiple replicates are needed for comparison and statistical analysis (1-4).

Some popular TSA analysis tools are Protein Thermal Shift™ Software, Serial Explorer in

MATLAB, and open-source workflows provided through the Konstanz Information Miner (16),

as well as some other free tools (17, 18). While both Protein Thermal Shift™ and MATLAB

provide high-throughput analysis, they are proprietary and require additional costs. Despite other

tools being free and open-source, they often offer limited visualization or analysis options. Given

the need to visualize the data in multiple dimensions and enhance accessibility of analysis tools,

we developed a software package written in R (19), a free programming language extensively used

for biological data analysis. 'TSAR', short for Thermal Shift Analysis in R, is a free and open-

source R package (Figure 1). TSAR is hosted by *Bioconductor* and distributed under an AGPL-3

license that offers a workflow as both command-line functions and an interactive ShinyR graphic

user interface (GUI) dashboard (20, 21). TSAR melting curve analysis allows users to select

Boltzmann-fit or derivative-modeling of data to identify either T_mB (Boltzmann-fit) or T_mD

(derivative-model) from melting curves that are smoothened with beta-knot fitting (Supplementary

Figure S1B-C) (22, 23). The TSAR package is a complete suite of tools from initial data

processing, T_m identification, multidimensional comparisons, and visualization of various

publication-quality graphics from individual experiments to large screens with any number of

replicates (Figure 1B-E). TSAR is a well-documented package and has many custom settings for

users experienced in R, while it also incorporates wrapping functions and a ShinyR dashboard to

minimize coding for easy use and quick implementation (20, 24). Overall, TSAR aims to simplify

TSA analysis yet produce diverse visualizations.

Here, we perform two example screens using TSAR to identify potential binding partners

to the human immunodeficiency virus type 1 (HIV-1) capsid protein (CA). The HIV-1 capsid core,

made of over 1500 copies of CA, is an essential macromolecular structure within the mature virus

that serves many roles during infection (25-28). The capsid core is the reaction chamber for reverse

transcription, it is a shield to hide the viral genome from cellular immune sensors, and it is the

vehicle that traffics the HIV-1 genetic material through the cytoplasm to, and likely through, the

nuclear pore complex (28-37). As it is essential in numerous steps of HIV-1 replication,

investigating the biology of the capsid is pivotal in the design of next-generation antiretroviral

therapies (ART) (38-40). In fact, when the CA-targeting compound Lenacapavir (LEN, formerly

GS-6207) was approved for use in highly-treatment experienced patients in 2022, CA became the

first new target for a clinically-used ART class in over a decade (38, 41). Recently, LEN was

additionally approved for use as pre-exposure prophylaxis (PrEP), reviewed in (42). Thus,

understanding the biology of this clinically-relevant target is of high importance.

One reason that the capsid core is a promising antiretroviral target is that CA is a highly

conserved domain and genetically fragile protein; the kinetics of CA•CA interactions are finely

tuned by evolution for balancing cellular trafficking and genome release, such that both increasing

and decreasing capsid stability has a negative impact on viral fitness (9, 28, 40, 43-46). The mature

HIV-1 core is comprised of primarily hexameric (CA_{HEX}) multimers of CA (Figure 1F), which

form a central pore with multiple basic residues like Arg18 and Lys25 (Figure 1G). In fact, reports

have demonstrated that CA_{HEX} interacts with host-derived acidic metabolites like dNTPs and

inositol hexaphosphate (IP6) at this pore to facilitate efficient reverse transcription and promote

stable capsid assemblies (Figure 1H-I) (28, 32, 47-51). Considering that the mature capsid

navigates through the cytoplasm during infection (27, 42, 47), in addition to reports that HIV-1

infection modulates cellular metabolism both in the presence and absence of ART (52-54), we

hypothesized that there might be additional endogenous interactions with other common protein

cofactors or naturally-present acidic small molecules. Thus, two example screens are presented to

identify potential interactions of vitamins or anionic metabolites with purified CA_{HEX} (55). First,

a small library of essential vitamins that are enzymatic cofactors finds a potential *in vitro*

interaction with folic acid (vitamin B9). Second, a library of 96 acidic metabolites finds many

anionic compounds that increase CA_{HEX} T_m. This library contained internal control of IP6 ("phytic

acid") that had the largest ΔT_m; thermally-stabilizing hits include gallic acid, xanthurenic acid,

metabolites from the tricarboxylic acid (TCA) cycle, and others. Both *in vitro* activity and cell-

based antiviral activity were tested for multiple hits, and gallic acid routinely affected HIV-1

replication or *in vitro* enzymatic activity when the mature capsid was present. Gallic acid has been

shown to have antiviral activity against HIV-1 previously, however these studies point to targets

like the reverse transcriptase (RT) enzyme (56-58); our analysis suggests an additional role for

capsid core that encapsulates the reverse transcription reaction. These results provide example

workflows for the TSAR package, a tool that can be broadly used in protein biology, and the

screens presented here have interesting implications for the stability of HIV-1 capsid in the context

of cellular metabolism.

Results

TSAR Package Accuracy

To assess the accuracy of TSAR, we found the T_mD and T_mB return similar values

(Supplementary Figure S1C). We benchmarked the determined T_m and ΔT_m shift from TSAR to

those from Protein Thermal Shift Software for the 12-component vitamin library. Using the Protein

Thermal Shift Software as the baseline value, TSAR T_mD error is centered at 0.048 with a variance

of 0.063 degree Celsius (Supplementary Figure S1D). Meanwhile, TSAR T_mB error is centered at

1.029 with a variance of 0.634 degree Celsius (Supplementary Figure S1E). While that error is

relatively greater than ΔT_mD, the error is systematic where computing ΔT_mB by using DMSO as

the baseline for ΔT_m, errors centering at 0.376 °C (Supplementary Figure S1F-G). Thus, we set

T_mD as the default modeling method in TSAR.

Vitamin library exemplifies TSAR output types

Testing the vitamin library, with IP6 as a positive control, yielded significant result for

folic acid with a ΔT_mD of 3.04 °C (Figure 2). Figure 2 exemplifies three methods of visualization

by TSAR: individual melting curves with an average (Figure 2A), compare averaged melting

curves (Figure 2B), and boxplots to show the average ΔT_m (Figure 2C). Derivative plots, which

show T_m as a maximum, are visualized also (Supplementary Figure 2A-C).

Testing at 50 μM, 25 μM, and 7.5 μM compound, the first derivative comparison graphs

demonstrate that the ΔT_m of folic acid with CA121 is dose dependent (Supplementary figure 2A-

D). At 7.5 μM, protein unfolding curves are distinguishable from DMSO control, but ΔT_m

experiences greater variance. Conducting one-tailed, paired T-tests show that all treatment

conditions are significantly different from the DMSO control, with 50 μM and 25 μM folic acid

(p≤5E-5), and 7.5 μM (p≤5E-3) (Supplementary Figure 2D). Structures of IP6 and Folic acid

shown (Supplementary Figure 2E). When breaking the CA_{HEX} into CA monomers using TCEP

(Supplementary Figure 2F), we find a loss of ΔT_m for both the IP6 control and folic acid, indicating

the importance of the multimeric CA (Supplementary figure 2G-H).

Multiple acidic metabolites cause a ΔT_m for CA_{HEX}

The second library contained 96 anionic compounds that are part of various metabolic

pathways and included an internal control of IP6. As expected, 25 μM IP6 provides an extremely

strong ΔT_m of 12°C, but multiple other metabolites also had statistically significant ΔT_m values

(Figure 3A-C, Supplementary Figure 3). Initial hits included gallic acid (ΔT_m=6.37°C), multiple

TCA cycle metabolites like α-ketoglutaric acid (αKg, ΔT_m=5.79°C), oxaloacetic acid

(ΔT_m=4.99°C), citric acid (ΔT_m=4.10°C), and others like xanthurenic acid (ΔT_m=2.28°C) and

indole-3-pyruvic acid (ΔT_m=1.65°C).

KEGG analysis mapped 53 metabolites in the acidic library to *H. sapiens* pathways (59),

and of them, over 60% of the represented TCA cycle metabolites caused a ΔT_m (5 hits of 8

possible, 5/8), especially those involved in the synthesis of αKg (Figure 3E, Supplementary Figure

4A). Other pathways with 50% or more matched metabolites that cause a ΔT_m at 25 μM include

amino acid biosynthesis (50%, 7/14) especially for tryptophan biosynthesis (66.7%, 4/6), cofactor

biosynthesis (58%, 7/12), lipoic acid metabolism (50%, 2/4), and inositol phosphate metabolism

(100%, 2/2) (Figure 3E).

To validate the binding of gallic acid to CA_{HEX}, we used BLI to determine binding kinetics.

We used a HIS-tagged CA_{HEX} ($CA_{HEX:HIS}$) to immobilize the proteins to the HIS-1K biosensor

(60). Using blocking buffer that contained Bovine Serum Albumin (BSA), which worked well for

other experiments (60-62) including compounds that bind to the phenylalanine-glycine (FG)

dipeptide binding site in CA_{HEX} (28, 60), did not return kinetic binding curves that could be

analyzed (data not shown). Thus, we optimized the blocking buffer by switching BSA to lysozyme,

as BSA is reported to interact with many metabolites including gallic acid (63, 64). We find PF74

binds to $CA_{HEX:HIS}$ with a K_D of 4.41±0.16 µM, k_{on} of 4.45E+03±1.44E+02 $M^{-1}s^{-1}$, and k_{off} of

1.96E-02±3.32E-04 s^{-1} (Figure 3F), slightly different than with the BSA-containing buffer (60,

65), and Gallic acid binds with K_D of 22.3±0.64 µM, k_{on} of 4.02E+02±9.58E+00 $M^{-1}s^{-1}$, and k_{off}

of 8.96E-03±1.42E-04 s^{-1} (Figure 3G). This confirms binding of these ligands to CA_{HEX}.

Gallic acid binds CA_{HEX} and has antiviral activity

We first checked the effect of adding these native metabolites in excess would influence

HIV-1 infection. No compound was cytotoxic at 100 µM, except for the antiretroviral control PF74

with a 50% cytotoxic concentration (CC_{50}) at 75.29 µM (14, 28, 40, 66). PF74 serves as a positive

control with a 50% effective concentration (EC_{50}) of 0.6±0.1 µM, aligning with literature values

(14, 28, 40, 66). Notably, gallic acid had antiviral activity with an EC_{50} of 19.2±0.1 µM for

pseudotyped HIV-$1_{NL4-3\Delta Env}$. This inhibition aligns with other reports of gallic acid antiretroviral

activity (56-58, 67).

While the PF74 control did increase CA assembly rate as reported, no metabolite tested

increased WT CA assembly in an assay that measures rates of assembly using turbidity

(Absorbance at 350 nm (A_{350})) as a proxy for lattice assembly (14, 68, 69) (Supplementary Figure

4B).

We further employed the ERT assay that tests the activity of RT within purified, native

capsid cores stabilized with IP6 (32, 49). This determines the effect on early, intermediate, and

late stages of reverse transcription (reported in (32, 49)), to systematically probe the inhibition of

RT. The early steps are somewhat impacted by PF74 treatment, but impacts intermediate and late

stages more, as damaging the capsid decreases the reaction efficiency of RT (49). Gallic acid has

increasing inhibition as the reaction proceeds: no determined effect on early reverse transcription,

a modest effect on intermediate steps (intermediate IC_{50}=17.21 µM), and a significant effect on

late step products (late IC_{50}=3.64 µM). Additionally, the metabolite xanthurenic acid consistently

impaired RT from early to late events (IC_{50}s = 9.65 µM - 11.99 µM). Other metabolites tested for

ERT and antiviral activity had no effect, including folic acid, oxalacetic acid, citric acid, isocitric

acid, tricarballylic acid, 2-methylcitric acid, αKg, and (Z)-aconitic acid (data not shown).

To further probe if the polymerase activity of RT is inhibited by gallic acid, we used a

reported primer-extension assay applying purified enzyme to a short template and treated with

inhibitors (70, 71). While RT inhibitors doravirine (DOR) and islatravir (EFdA) impair primer

extension (71), up to 1000 µM of gallic acid had no effect on the polymerase activity of RT, similar

to the other CA inhibitors tested, PF74 and LEN, that do not target RT (Supplementary Figure 4C).

Discussion

Here, we performed TSA screens to identify potential protein•ligand interactions that was

conducted with TSAR and found that multiple metabolites stabilize HIV-1 CA_{HEX} *in vitro*. TSAR

is a novel package written in the R language, which is a tool used by many biological research labs

and in higher-education coursework (72). There are other software tools for TSA data analysis,

but they are costly or have limited visualization options. Meltdown is a similar tool and is a free

and user-friendly python package (18), however, TSAR uniquely provides T_m estimation using the

boltzman-fit method and includes a GUI dashboard. MoltenProt is a free and useful online tool but

requires additional visualization software for high-resolution graphics (17); TSAR is free for

academics and can output publication-ready graphics.

There are reports identifying endogenous metabolites that interact with mature CA_{HEX}

being critical cofactors in HIV-1 replication, such as IP6 that is important for capsid stability as

well as dNTPs that are needed for genome replication (28-31, 47, 48). TSAs alone do not establish

the virological significance of these metabolites, only that stabilizing interactions are observed *in*

vitro for CA_{HEX}. By investigating the TSA hits, our studies that included gallic acid find antiviral

activity and inhibition of RT activity, aligned with some prior reports about the inhibition of

reverse transcription activity and HIV-1 infection by gallic acid (56-58, 67). Using ERT assays

where is RT inside the capsid core (32, 49), the intermediate and late reverse transcription products

were inhibited by gallic acid, but not if RT is alone in the *in vitro* primer-extension assay that tests

its polymerase activity (70, 71). CA_{HEX} are involved dNTP import that fuel RT; it is shown that

blocking the capsid pore with the flat, inorganic acid hexacarboxybenzene can impact reverse

transcription (47, 49, 73). Attempts to solve the atomic structure of CA with gallic acid by X-ray

crystallography were not successful (data not shown). Other functions of RT may be impacted,

such as RNase H activity or metal ion coordination (67, 74), as gallic acid is a divalent ion chelator

(75). If the enzymatic target is RT, as proposed by others (56, 57), our results indicate there is a

relationship with gallic acid, a non-nucleoside molecule that, in theory, would need to traverse the

mature capsid core and reach RT or impair its function distant from the enzyme.

We did not anticipate that supplementing cells with endogenous metabolites would impair

viral fitness, as most metabolites are readily converted to other molecules by cellular enzymes,

and the virus replicates within biologically active cells that typically contain these metabolites. No

observable effect was the case for almost all TSA hits, except for gallic acid. Gallic acid is a

secondary metabolite typically acquired through diet or gut microbiota; concentrations in human

cells are generally low and there are reports of gallic acid cytotoxicity in some cell-types (75, 76).

Gallic acid has been shown to have antiviral activity for multiple viruses, including HIV-1,

influenza, and encephalomyocarditis virus (56, 57, 76-78), indicating a complex and likely

multifaceted mechanism. Gallic acid is also a sticky molecule, and is known to interact with many

cellular and viral proteins, including a report that used surface plasmon resonance and found

interactions with HIV-1 proteins CA, RT and Gag (57, 63, 64, 79-81). ERT assays do not find an

impact on *in vitro* RT products by treatment with most of the tested metabolites, except gallic acid

and xanthurenic acid. Xanthurenic acid is a metabolite involved in the tryptophan oxidation

pathway with concentrations reportedly decreased in ART-naïve participants living with HIV-1

(54, 82); we found it impacts *in vitro* ERT but not an observable antiretroviral effect *in vivo* (Figure

4A-D).

Numerous studies have investigated the requisites for and changes in metabolism during

HIV-1 infection (53, 54, 82, 83). Of note, a report found that cells lacking mitochondria inhibit

HIV-1 post-entry and pre-integration (83), a phenotype typical with dysfunctional capsid; another

study found increased HIV-1 infection when the TCA cycle was highly active (53). While only

the TSAs showed an impact on CA thermal stability and not an effect in other virological assays,

the assays were tested with excess metabolites and not in the absence, which may have a different

effect. The role and relevance of these endogenous molecules, both the vitamin folic acid and the

organic acidic metabolites, in HIV-1 biology remains a topic of investigation, but the findings of

our screen reported here indicate the ability for CA$_{HEX}$ to form many stabilizing interactions with

endogenous polyanions. We report here that some CA-stabilizing molecules, identified by TSAR,

can impair HIV-1 reverse transcription inside capsid cores, and we further the mechanistic

knowledge of gallic acid antiretroviral activity.

Methods

Distribution and Package Information

TSAR is an open-source package written in R (Version (V)≥4.3) (19), distributed under an AGPL-

3 license in the peer-reviewed *Bioconductor* R package repository (V≥3.18) (21, 84) at

https://bioconductor.org/packages/TSAR. TSAR was coded using RStudio (85) and devtools (86).

TSAR contains well-documented functions and multiple vignettes (long-form documentation)

(24). TSAR depends on tidyverse packages (87) as well as: jsonlite (88), openxlsx (89), mgcv (90-

92), and minpack.lm (93) for analysis, and shiny (20), shinyWidgets (94), plotly (95), shinyjs (96),

rhandsontable (97), and ggpubr (98) packages for the GUI and data visualizations.

Package Structure

TSAR has three individual operating dashboards, "Data Preprocessing", "Data Analysis",

and "Data Visualization", each with one command line call (Figure 1B). The GUI dashboard is

optional, as TSAR also allows command-line functions. The package also enables "Data

Visualization"-only, if suppled with pre-determined T_m values to use TSAR to visualize previously

analyzed data. Mathematical modeling in TSAR relies on both generalized additive models (GAM)

cubic spline with beta-knots or Boltzmann fitting to compute T_mD and T_mB values (22, 23, 90, 92,

99, 100). See package documentation in TSAR vignettes (24).

Statistics and Analysis

TSAR utilizes GAM from the mgcv package to capture the derivatives of the curve and the

inflection point, T_mD (90-92). Alternatively, TSAR utilizes non-linear modeling from the

minpack.lm package (93) in R to capture Boltzmann curves with machine-learned minimum and

maximum, T_mB.

T_mB uses a kinetic equation describing the statistical behavior of a system of particles (99);

it is applied widely towards sigmoidal behaviors. Using parameter-based mathematical modeling,

individual melting curves are fit through a 5-parameter function (101) (Supplementary Figure

S1B).

T_mD, representing T_m values estimated using a derivative-model, utilizes smoothing to

remove noises from data and produces a polynomial function closely representing the trend and

shape of raw data. T_ms are then derived through the calculation of first derivatives and location

inflection point (101). Both methods are used in extant research protocols where T_mB are less

affected by trivial fluctuation and T_mD captures all levels of variations throughout the denaturation

process.

Protein Purification

The CA_{HEX} with A14C/E45C/W184A/M185A mutations (CA121) were cloned into a

pET11a expression plasmid (55), kindly provided by Dr. Owen Pornillos (University of Utah).

The expression and purification of CA121 were carried out in *E. coli* BL21(DE3)RIL, and

purification and cross-linking was performed as previously described to generate individual,

soluble, and disulfide-stabilized CA_{HEX} (14, 55). A C-terminal His-tag was cloned onto the CA121

construct (Genscript) for Biolayer Interferometry (BLI) experiments (60, 61), and this construct

was expressed *E. coli* NiCo21(DE3) and lysed and cross-linked as in (55), but purified by Ni^{2+}

chromatography before crosslinking and a final size-exclusion chromatography (SEC) step to

separate cross-linked CA_{HEX} from monomeric CA.

Additionally, monomeric full-length, wild-type CA (WT CA) was cloned into a pET24a

expression plasmid, kindly provided by Dr. Robert Dick (Emory University). The expression of

WT CA was performed in *E. coli* NiCo21(DE3) and purification involved ammonium sulfate

precipitation followed by desalting, cation-anion subtractive chromatography, and SEC as

previously described (50).

HIV-1 RT was expressed and purified as previously described (70). Briefly, pRT6H-PROT

(102) was expressed in *E. coli* BL21(DE3)RIL and purified by Ni^{2+} chromatography.

Compounds

The vitamin library of chemicals were purchased as the Vitamins Kit (Sigma Aldrich, V1-

1KT), resuspended in DMSO to a final concentration of 10-50 mM and stored at -20°C.

Compounds included: D-pantothenic acid hemicalcium salt (B5), biotin, nicotinamide (B3), 4-

aminobenzoic acid (4-ABA), pyridoxamine dihydrochloride (PyxAmine diHCl), pyridoxine

hydrochloride (PyxINE HCl), thiamine hydrochloride (Thiamine HCl), folic acid (B9), (±)-α-

lipoic acid (a-LA), pyridoxal hydrochloride (Pyxal HCl). 50% phytic acid solution (TCI) was used

for IP6 positive controls. The Organic Acid Metabolite Library of Standards (OAMLS) was used

(IROA Technologies), and DMSO was added to a final concentration of 5 mM. Phytic Acid is

within OAMLS and served as an internal control. After initial screening, select chemicals were

purchased: gallic acid [HY-N0523], xanthurenic acid [HY-W014666], dihydroferulic acid [HY-

N7080] (similar to 2-hydroxy-3-(4-hydroxy-3-methoxyphenyl)propanoic acid), indole-3-pyruvic

acid [HY-W028393], oxalacetic acid [HY-W010382], DL-isocitric acid trisodium salt [HY-

W009362], citric acid [HY-N1428], tricarballylic acid [HY-W020215], 2-methylcitric acid [HY-

113371], α-ketoglutaric acid [HY-W013636], and (Z)-aconitic acid [HY-W016814]

(MedChemExpress). PF74 powder [SML0835] was purchased from Sigma-Aldrich; EFdA-TP

tetraammonium [HY-138561A] and Doravirine [HY-16767] were purchased from

MedChemExpress. LEN (HRP-20266) was kindly provided by the NIH HIV Reagents Program.

dNTPs were purchased from Promega. Oligonucleotides were chemically synthesized and

purchased from Integrated DNA Technologies™.

Thermal Shift Assay (TSA)

TSAs were conducted using QuantStudio 3 qPCR machine (Thermo Fisher Scientific); the

final reaction was 20 µL per well of 7.5 µM CA in 50 mM Tris (pH 8.0) and 1x SYPRO™ Orange

Protein Gel Stain (Life Technologies) after incubation with compounds for 30 min on ice.

Fluorescence was measured while heating consistently increased 0.2°C/10s from 25–95°C (6, 9,

14, 103). For initial screening, CA121 was incubated with 25 µM of each compound. Some

previous reports use 50 mM sodium phosphate buffer (pH 8.0), however, the polyanionic

phosphates can potentially compete acidic metabolites for binding to the CA_{HEX}, thus, we opted

for 50 mM Tris (pH 8.0) instead. Thermal profiles were assessed using TSAR or Protein Thermal

Shift Software (V1.4, Applied Biosystems). The compounds that induced a significant ΔT_m (>1°C)

at 25 µM underwent further testing in a dose-response manner under concentrations of 7.5 µM and

50 µM.

Lastly, vitamins leading to significant thermal shifts were also tested in either WT CA

monomers or in the presence of the reducing agent tris(2-carboxyethyl)phosphine (TCEP). TCEP

breaks the CA121 disulfide-stabilizing bonds, likely affecting the stability of pockets formed by

multiple CA monomers. Comparing the fluorescence profile of TSA samples with and without

TCEP provides insight into the potential interactions of a compound with monomeric versus

hexameric CA.

Metabolic pathway analysis was conducted on the KEGG webserver (59). The KEGG/CSI

IDs were provided by the OMAL library manufacturer, and those without matches were manually

searched. Of the 96 compounds, 53 were successfully matched to *H. sapiens* pathways, including

the 13 of the 20 ΔT_m hits. Hits not mapped to human pathways are: gallic acid, 1,2,3,-

propanetricarboxylic acid, vanillactic acid, 3-Hydroxy-3-methylglutaric acid, indole-3-butyric

acid, indole-3-propionic acid, and 2,5-furandicarboxylic acid.

Biolayer Interferometry (BLI)

The BLI protocol was adapted from (60, 61) with minor modifications. Briefly, $CA_{HEX:HIS}$ was

immobilized onto a HIS1K biosensor at 100 µg/mL for 600 seconds. Loaded probes were

equilibrated for 120 seconds before being exposed to a series of association and dissociation

phases. Starting at the lowest analyte concentration in the series, $CA_{HEX:HIS}$ was dipped into a well

containing analyte (gallic acid, PF74 (60, 65), or vehicle control) for 100 seconds at concentrations

of 5, 10, 20, or 50 µM. Probes were then allowed to dissociate in a well containing blocking buffer

for 100 seconds. Blocking buffer contained: 50 mM Tris (pH 8.0), 1 mg/mL lysozyme, 0.6 M

Sucrose, and 20 mM imidazole. Three experiments were initially analyzed using the Octet software

where replicates were combined, and double reference subtraction was applied. Furthermore, the

Y-axis was aligned to the baseline step, the inter-step correction was set to baseline at time 0 s,

and Savitzky-Golay Filtering was applied.

Antiviral Activity and Cytotoxicity

Antiviral and cytotoxicity was determined as previously reported (14, 104). Pseudotyped HIV-1

virions (VSV-G-NL4-3ΔEnv) (105) were produced in HEK-293/17 cells that were cultured in

DMEM (Corning), and supplemented with 2 mM L-glutamine, 10% FBS, and 100 U/mL

penicillin/streptomycin and harvested 48 hpi. Infection was determined in TZM-GFP cells, a

reporter-cell line that expresses GFP induced by Tat and regulated by Rev, accompanied by an

internal ribosomal entry site (IRES) for translation, upon HIV-1 infection (106). To obtain the

EC_{50} values, dose-dependence data was plotted in GraphPad Prism and analyzed with the

log(compound concentration) compared to normalized fluorescence *via* a response–variable slope

equation and averaged. TZM-GFP cells were used for cytotoxicity, and cell viability was assessed

with an XTT kit (Roche) according to the manufacturer's instructions then CC_{50} values were

determined using the same method as the EC_{50} values. TZM-GFP cells (HRP-20041) and the HIV-

1_{NL4-3} virus (ARP-3418) were gifts from the NIH HIV Reagent Program.

399   **Endogenous Reverse Transcription (ERT) Assay**

The ERT assay was used to determine the levels of reverse transcription products within native

HIV-1 capsid cores, protocol adapted from (32, 49). Pseudotyped HIV-1 virions were produced in

HEK-293/17 cells, as mentioned above. As previously reported (14), virions were permeabilized

with melittin, then p24 levels for the sample was determined by ELISA using HIV-Ig (ARP-

3957) as the primary antibody and anti-human HRP (Rockland Immunochemicals) as the

secondary antibody, followed by quantification by a GloMax Navigator Plate Reader (Promega)

after the chemiluminescent substrate (ThermoFisher) was added.

Purified virions, normalized to 500 ng p24, were treated with TURBO DNA-*free*™ DNase

(AM1907), and various concentrations of metabolites were added in the presence of 10 mM Tris-

HCl (pH 7.8), 75 mM NaCl, 2 mM $MgCl_2$ 2.5 µg/mL melittin, 80 µM IP6, 0.5 mg/mL BSA, and

40 µM dNTPs at 37°C for 16 h. The reaction steps were quantified by qPCR with SYBR green

detection using the following primer sets: 5′-GAGCTCTCTGGCTAACTAG-3′ and 5′-

TGACTAAAAGGGTCTGAGGGATCT-3′ (minus-strand strong stop), 5′-

GAGCCCTCAGATGCTGCATAT-3′ and 5′-CCACACTGACTAAAAGGGTCTGAG-3′ (first-

strand transfer), 5′-TGTGTGCCCGTCTGTTGTGT-3′ and 5′-GAGTCCTGCGTCGAGAGAGC-

3′ (second-strand transfer) adapted from (49).

416   **Reverse Transcriptase Primer Extension**

Primer extension reactions were perform as previously described (70, 71). A Cy3-labled primer

was annealed to the template at 1:2 molar ratio. Gallic acid and control inhibitors were incubated

with RT for 30 minutes. Then, reactions were initiated by adding master mix containing 20 nM

annealed Primer-Template substrate, 15 µM dNTPs, 6 mM MgCl2 in RT buffer (50 mM Tris, pH

7.8 and 50 mM NaCl) to a final volume of 20 µL and final RT concentration of 100 nM. The

reactions were terminated after 60 minutes by adding an equal volume of 100% formamide

containing bromophenol blue. Reaction products were resolved on a 7 M urea / 15%

polyacrylamide gel, followed imaging on a phosphoimager (Typhoon FLA 9500).

The primer used are 5-Cy3-GTCACTGTTCGAGCACA-3' (18mer); template sequence is 5'-

CCATAGCTAGCATTGGTGCTCGAACAGTGAC-3' (31mer), from (70).

CA assembly assay and Transmission Electron Microscopy (TEM)

The *in vitro* CA assembly assay uses optical density as a proxy for lattice assembly (68), and these

were conducted as previously reported (14, 68). Concentrated WT CA was diluted to 100 µM in

50 mM Tris (pH 8.0), treated with 100 µM of compound or vehicle control (DMSO or dH2O), and

incubated on ice for at least 30 minutes. Then, to initiate assembly, high-salt buffer was added at

a 1:1 ratio to the compound-treated sample in a 96-well plate. Each well contained 100 µL of 50

µM WT CA, 50 µM compound, and 2 M NaCl in 50 mM Tris (pH 8.0). Data were collected on a

BioTek Neo plate reader, with A_{350} measured every 25 s for 2.5 h at room temperature.

Simultaneously, a background control was measured that contained 50 µM compound and 50 µM

WT CA in 50 mM Tris, with no NaCl; measurements of the protein assembly had the background

readings subtracted. GraphPad Prism was used for visualization and statistical analysis.

The negative stain TEM protocol was adapted from (48). Briefly, 500 µM WT CA in 50

mM MES (pH 6.2) was treated with 500 µM metabolite (1:1), then capsid-like particles (CLPs)

were assembled by mixing a 1:1 volume of the preincubated WT CA with a buffer that contains

an excess of IP6 at 37°C. See (14, 48, 50).

Acknowledgments

We would like to thank the members of the Sarafianos lab for feedback during TSAR package development. This project was supported in part by the National Institutes of Health (R01 AI120860 and U54 AI170855 to S.G.S.; F31 AI174951 to W.M.M.; F31 AI179424 to X.W.; W.M.M. and Z.C.L. were supported in part by T32 GM135060). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. S.G.S. acknowledges funding from the Nahmias-Schinazi Distinguished Chair in Research. The plasmid encoding the Disulfide Stabilized CA_{HEX} (CA121) was kindly provided by Dr. Owen Pornillos (University of Utah); the plasmid encoding WT CA monomers was kindly provided by Dr. Robert Dick (Emory University); the plasmid encoding the 6HRT construct was kindly provided by Dr. Michael Parniak (University of Pittsburgh). LEN, TZM-gfp, HIV-Ig, and plasmid for pseudotyped virions were gifts from the NIH HIV Reagent Program. Electron Microscopy was carried out by Emory University Robert P. Apkarian Integrated Electron Microscopy Core Facility (RRID: SCR_023537). For providing essential details in making an R package, we would like to acknowledge the book R Packages by Hadley Wickham (O'Reilly) ©2015 Hadley Wickham, ISBN: 978-1-491-91059-7.

Author contributions

Conceptualization, WMM, KAK, and SGS; Software, WMM, XG, and ZY; Methodology & Investigation, WMM, XG, ZY, XW, ZCL, HZ, JF and AE; Resources, WMM, ZCL, JF, AE and KAK; Visualization, WMM, XG, ZY, XW; Writing – Original Draft, WMM, XG, ZY and SGS; Writing – Review & Editing, WMM, XG, ZY, and SGS; Supervision, WMM, KAK, and SGS; Funding Acquisition, WMM, KAK and SGS.

Declaration of interests

466   The authors declare no competing interests.

Figure 1: TSAR (*Thermal Shift Analysis in R*) package overview and example protein target.
A) TSAR Logo. **B)** Workflow for TSAR, supporting data import and export at various stages. **C-E)** Screenshots of the TSAR program. **C)** shows the Shiny GUI for importing data, **D)** shows the interactive data analysis that TSAR performs, and **E)** is an example of a partial output from a library screen. **F)** Structure of WT HIV-1 CA$_{HEX}$, visualized with ChimeraX (PDB: 4XFX)(26), **G)** Partial side view of (F), showing the cationic central pore with Arg18 and Lys25 in stick representation. **H)** Structure of IP6 in the CA$_{HEX}$ central pore (PDB: 6BHT)(48). **I)** Structure of dATP in the CA$_{HEX}$ central pore (PDB: 5HGM)(47).

Figure 2: Example TSAR outputs and screening of CA$_{HEX}$ binding to a library of vitamins.

A) Folic acid thermal melting profiles by individual sample (n = 4). Each well is shown as a different color. Y is normalized relative fluorescence units (RFU). The grey line represents the modeled average between all qPCR wells **B)** Compare plot further showcases ΔT_m interactions as averaged RFU over temperature (°C), compared to DMSO control (grey). Only IP6 and folic acid

485   yield significant T_m change: IP6 $\Delta T_m D=13.75°C$ ($p<5E-11$), folic acid $\Delta T_m D=3.04°C$ ($p<5E-5$).

486   **C)** Screening vitamin library at 50 µM with CA121 (CA_{HEX}) at 7.5 µM. Compared to DMSO

487   vehicle.

A Initial ΔT_m Hits from Organic Acid Library

Phytic acid (IP6)
ΔT_m = 12°C

Gallic acid
ΔT_m = 6.37°C

2-Oxoglutaric acid
ΔT_m = 5.79°C

2-Methylcitric acid
ΔT_m = 5.77°C

Oxalacetic acid
ΔT_m = 4.99°C

Citric acid
ΔT_m = 4.10°C

1,2,3-Propanetricarboxylic
ΔT_m = 4.10°C

Isocitric acid
ΔT_m = 4.05°C

2-hydroxy-3-(4-hydroxy-3-methoxyphenyl)propanoic acid
ΔT_m = 3.04°C

2-oxoadipic acid
ΔT_m = 2.86°C

Xanthurenic acid
ΔT_m = 2.28°C

Aconitic acid
ΔT_m = 1.65°C

Indole-3-pyruvic acid
ΔT_m = 1.65°C

Indole-3-butyric acid
ΔT_m = 1.50°C

2-Isopropylmalic acid
ΔT_m = 1.40°C

3-hydroxy-3-methylglutaric acid
ΔT_m = 1.34°C

Quinolinic acid
ΔT_m = 1.34°C

Prephenic acid
ΔT_m = 1.13°C

Indole-3-propionic acid
ΔT_m = 1.12°C

2,5-furandicarboxylic acid
ΔT_m = 1.11°C

B CA121 + Phytic Acid

C CA121 + Gallic Acid

D CA121 + Citric Acid

E Pathway Analysis

Metabolic pathways — 24.5%
2–Oxocarboxylic acid metabolism — 46.7%
Biosynthesis of amino acids — 50.0%
Biosynthesis of cofactors — 58.3%
Citrate cycle (TCA cycle) — 62.5%
Butanoate metabolism — 14.3%
Lipoic acid metabolism — 50.0%
Pyrimidine metabolism — 0.0%
cAMP signaling pathway — 0.0%
GABAergic synapse — 33.3%
Inositol phosphate metabolism — 100.0%
Oxidative phosphorylation — 0.0%

Full Library
TSAR Hits

KEGG Match Count

F Biolayer Interferometry:
CA^HEX with PF74

Wavelength shift (nm)
Time (s)

— 5 μM — 10 μM — 20 μM

G CA^HEX with gallic acid

Wavelength shift (nm)
Time (s)

— 10 μM — 20 μM — 50 μM

Figure 3: TSA of organic metabolic acids find many potential interactions with CA$_{HEX}$. A)

Top 20 hits from the 96-compound library in 50 mM Tris (pH 8.0). 25 µM compound and 7.5 µM

CA$_{HEX}$ (CA121). Made in ChemDraw (V19.0). See Supplementary Figure 3. **B-C)** TSAR compare

plots to show the large ΔT_m caused by **B)** phytic acid (IP6, internal control), **C)** gallic acid, and **D)**

citric acid. **E)** *H. sapiens* pathway matches to metabolites in the library (in white) or those that

cause ΔT_m determined by TSAR (in red) in the KEGG Database (59). **F & G)** Biolayer

interferometry (BLI) validates interaction of CA$_{HEX:HIS}$ with **F)** PF74 or **G)** gallic acid.

A

Compound	EC$_{50}$ (μM)	CC$_{50}$ (μM)
PF74	0.605 ± 0.0313	75.29 ± 3.9†
Gallic acid	19.24 ± 0.1131	>100
Isocitric acid	>100	>100
Folic acid	>100	>100
2-methyl citric acid	>100	>100
Xanthurenic acid	>100	>100

B ERT: minus-strand strong stop (early)

	IC$_{50}$ (μM)
PF74	12.58
Gallic acid	ND
Xanthurenic acid	11.21

C ERT: First strand transfer (intermediate)

	IC$_{50}$ (μM)
PF74	1.95
Gallic acid	17.21
Xanthurenic acid	11.99

D ERT: Second strand transfer (Late)

	IC$_{50}$ (μM)
PF74	1.32
Gallic acid	3.64
Xanthurenic acid	9.65

E WT CA + IP6

F WT CA + Gallic Acid + IP6

G WT CA + Xanth. Acid + IP6

H WT CA + αKG + IP6

Figure 4: Testing the virological effects of gallic acid and other CA$_{HEX}$-stabilizing metabolites on HIV-1 replication. A) Antiviral activity (EC$_{50}$) using HIV-1$_{NL4-3}$ pseudotyped virus in TZM-GFP cells and cytotoxicity (CC$_{50}$) of select compounds. [†]PF74 CC$_{50}$ from (14). **B-D)** Endogenous reverse transcription (ERT), as previously described (32, 49), takes place inside an IP6-stabilized capsids requiring both CA and RT function: **B)** early, **C)** intermediate, and **D)** late RT products are determined by qPCR after incubation with compounds to determine inhibitory concentrations (IC$_{50}$), compared to DMSO control. ND=Not Determined. **E-H)** Negative stain transmission electron microscopy (TEM) of *in vitro* assembled capsid-like particles (CLPs). Monomeric CA mixed with an excess of IP6 form conical assemblies, from (48, 50). **E)** IP6 alone, or IP6 plus **F)** gallic acid, **G)** xanthurenic Acid, or **H)** or 2-oxoglutaric acid (αKG) are added to WT CA. Scale varies. Chemical structures made in ChemDraw (V.19.0) and protein structures in Chimera X (PDB: 4XFX)(26).

Supplementary Figure 1: TSAR models and accuracy. A) Cartoon of thermal shift curves and

protein denaturation to determine the 50% melting point (T_m), change in T_m (ΔT_m) from ligand-

bound (blue circle) bound to the protein, thus increasing the temperature needed to denature the

sample. Made with BioRender. **B)** Theoretical curves to exemplify the two modeling strategies

between derivative T_m estimation (T_mD), and Boltzmann T_m estimation (T_mB). T_mB uses the

equation $y=1/(1+e^{-k(x-x2)})$ (100). **C)** T_mD vs T_mB, showing overall correlation of the two values

near the diagonal. **D, F)** Derivative estimation errors for the vitamin library. (n = 54) is N~(0.048,

0.063). ΔT_mD estimation error is similarly distributed, N~(-0.001, 0.029). **E, G)** Boltzmann

estimation errors (n = 54) is approximately N~(1.029, 0.634). ΔT_mB estimation error is biased in

the positive direction, N~(0.376, 0.451).

Supplementary Figure 2: Dose dependence study of folic acid and CA_HEX shows ΔT_m is dosage-dependent. A-C) First derivative comparison graphs demonstrate that ΔT_m of folic acid with 7.5 μM CA121 (CA_HEX) is dose dependent at **A)** 50 μM, **B)** 25 μM, **C)** 7.5 μM compound.

D) TSAR-generated boxplot of (A-C), with one-tailed t-tests show that all treatment conditions

are significant (*=$p \leq$5E-3, **=$p \leq$5E-5, and ***=$p \leq$5E-9). **E)** Chemical Structures of hits, Folic

Acid and IP6. Made in ChemDraw (V19.0). **F-H)** Reducing cross-linked CA121 shows thermal

shift of folic acid requires hexameric status. **F)** TCEP was added as a reducing agent to break apart

CA121, theoretically eliminating the central pore of protein-ligand interaction. **G)** TSA performed

with ligands at 25 μM. Comparing first derivatives show that TCEP alters regular protein

fluorescence curve. **H)** Boxplot demonstrates adding TCEP leads to insignificant $\Delta T_m D$.

Supplementary Figure 3: All hits from the 96-well organic metabolic acid library.

Supplementary Figure 4: TCA cycle and *in vitro* HIV-1 assays testing metabolite hits. A)

The tricarboxylic acid (TCA) cycle is shown, with T_m determined by TSAR labeled below for

those tested against CA_{HEX}. Made in ChemDraw (V19.0). **B)** An *in vitro* assembly assay for CA

proteins that uses turbidity (A_{350}) as a proxy for determining the rate of lattice formations (14,

68). The presence of metabolites does not seem to impact the rate of assembly. **C)** Testing the *in*

vitro polymerase activity of HIV-1 RT using a primer-extension assay (70) finds that gallic acid

does not inhibit the elongation of a DNA template. PF74 and LEN are CA-targeting inhibitors,

and DOR and EFdA are RT-targeting inhibitors.

References

555
556

1.   Andreotti G, Monticelli M, Cubellis MV. 2015. Looking for protein stabilizing drugs with
thermal shift assay. Drug Testing and Analysis 7:831-834.
2.   Gao K, Oerlemans R, Groves MR. 2020. Theory and applications of differential scanning
fluorimetry in early-stage drug discovery. Biophys Rev 12:85-104.
3.   Huynh K, Partch CL. 2015. Analysis of Protein Stability and Ligand Interactions by
Thermal Shift Assay. Current Protocols in Protein Science 79:28.9.1-28.9.14.
4.   Magnusson AO, Szekrenyi A, Joosten H-J, Finnigan J, Charnock S, Fessner W-D. 2019.
nanoDSF as screening tool for enzyme libraries and biotechnology development. The
FEBS Journal 286:184-204.
5.   Lo M-C, Aulabaugh A, Jin G, Cowling R, Bard J, Malamas M, Ellestad G. 2004.
Evaluation of fluorescence-based thermal shift assays for hit identification in drug
discovery. Analytical Biochemistry 332:153-159.
6.   Niesen FH, Berglund H, Vedadi M. 2007. The use of differential scanning fluorimetry to
detect ligand interactions that promote protein stability. Nature Protocols 2:2212-2221.
7.   Wu T, Hornsby M, Zhu L, Yu JC, Shokat KM, Gestwicki JE. 2023. Protocol for performing
and optimizing differential scanning fluorimetry experiments. STAR protocols 4:102688.
8.   Biter AB, de la Peña AH, Thapar R, Lin JZ, Phillips KJ. 2016. DSF Guided Refolding As
A Novel Method Of Protein Production. Scientific Reports 6:18906.
9.   Miyazaki Y, Doi N, Koma T, Adachi A, Nomaguchi M. 2017. Novel In Vitro Screening
System Based on Differential Scanning Fluorimetry to Search for Small Molecules against
the Disassembly or Assembly of HIV-1 Capsid Protein. Front Microbiol 8:1413.
10. Layton CJ, Hellinga HW. 2011. Quantitation of protein–protein interactions by thermal
stability shift analysis. Protein Science 20:1439-1450.
11. Kazakov AS, Markov DI, Gusev NB, Levitsky DI. 2009. Thermally induced structural
changes of intrinsically disordered small heat shock protein Hsp22. Biophysical chemistry
145:79-85.
12. McFadden WM, Yanowitz JL. 2022. idpr: A package for profiling and analyzing
Intrinsically Disordered Proteins in R. PLOS ONE 17:e0266929.
13. Krishna SN, Luan C-H, Mishra RK, Xu L, Scheidt KA, Anderson WF, Bergan RC. 2013.
A Fluorescence-Based Thermal Shift Assay Identifies Inhibitors of Mitogen Activated
Protein Kinase Kinase 4. PLOS ONE 8:e81504.
14. McFadden WM, Casey-Moore MC, Bare GAL, Kirby KA, Wen X, Li G, Wang H, Slack
RL, Snyder AA, Lorson ZC, Kaufman IL, Cilento ME, Tedbury PR, Gembicky M, Olson
AJ, Torbett BE, Sharpless KB, Sarafianos SG. 2024. Identification of clickable HIV-1
capsid-targeting probes for viral replication inhibition. Cell Chemical Biology 31:477-
486.e7.
15. Schoenaers S, Lee HK, Gonneau M, Faucher E, Levasseur T, Akary E, Claeijs N, Moussu
S, Broyart C, Balcerowicz D, AbdElgawad H, Bassi A, Damineli DSC, Costa A, Feijó JA,
Moreau C, Bonnin E, Cathala B, Santiago J, Höfte H, Vissenberg K. 2024. Rapid
alkalinization factor 22 has a structural and signalling role in root hair cell wall assembly.
Nature Plants 10:494-511.
16. Samuel ELG, Holmes SL, Young DW. 2021. Processing binding data using an open-source
workflow. Journal of Cheminformatics 13:99.

17.   Kotov V, Mlynek G, Vesper O, Pletzer M, Wald J, Teixeira-Duarte CM, Celia H, Garcia-
Alai M, Nussberger S, Buchanan SK, Morais-Cabral JH, Loew C, Djinovic-Carugo K,
Marlovits TC. 2021. In-depth interrogation of protein thermal unfolding data with
MoltenProt. Protein Science 30:201-217.
18.   Rosa N, Ristic M, Seabrook SA, Lovell D, Lucent D, Newman J. 2015. Meltdown: A Tool
to Help in the Interpretation of Thermal Melt Curves Acquired by Differential Scanning
Fluorimetry. SLAS Discovery 20:898-905.
19.   R Core Team. 2025. R: A language and environment for statistical computing, v4.3.1. R
Foundation for Statistical Computing, Vienna, Austria. https://www.R-project.org/.
20.   Chang W, Cheng J, Allaire J, Sievert C, Schloerke B, Xie Y, Allen J, McPherson J, Dipert
A, Borges B. 2024. shiny: Web Application Framework for R, https://CRAN.R-
project.org/package=shiny.
21.   Gentleman RC, Carey VJ, Bates DM, Bolstad B, Dettling M, Dudoit S, Ellis B, Gautier L,
Ge Y, Gentry J, Hornik K, Hothorn T, Huber W, Iacus S, Irizarry R, Leisch F, Li C,
Maechler M, Rossini AJ, Sawitzki G, Smith C, Smyth G, Tierney L, Yang JYH, Zhang J.
2004. Bioconductor: open software development for computational biology and
bioinformatics. Genome Biology 5:R80.
22.   Lambers JV, Mooney AS, Montiforte VA. 2021. Explorations in Numerical Analysis
doi:10.1142/12031. World Scientific.
23.   Yeh R, Nashed YSG, Peterka T, Tricoche X. 2020. Fast Automatic Knot Placement
Method for Accurate B-spline Curve Fitting. Computer-Aided Design 128:102905.
24.   Gao X, McFadden WM, Ye Z, Sarafianos SG. 2023. TSAR, *on* Bioconductor.
https://doi.org/doi:10.18129/B9.bioc.TSAR. Accessed
25.   Ganser-Pornillos BK, Schwedler UKv, Stray KM, Aiken C, Sundquist WI. 2004.
Assembly Properties of the Human Immunodeficiency Virus Type 1 CA Protein. Journal
of Virology 78:2545-2552.
26.   Gres AT, Kirby KA, KewalRamani VN, Tanner JJ, Pornillos O, Sarafianos SG. 2015. X-
ray crystal structures of native HIV-1 capsid protein reveal conformational variability.
Science 349:99-103.
27.   Jang S, Engelman Alan N. 2023. Capsid–host interactions for HIV-1 ingress. Microbiology
and Molecular Biology Reviews 87:e00048-22.
28.   McFadden WM, Snyder AA, Kirby KA, Tedbury PR, Raj M, Wang Z, Sarafianos SG.
2021. Rotten to the core: antivirals targeting the HIV-1 capsid core. Retrovirology 18:41.
29.   Aiken C, Rousso I. 2021. The HIV-1 capsid and reverse transcription. Retrovirology 18:29.
30.   Müller TG, Zila V, Müller B, Kräusslich H-G. 2022. Nuclear Capsid Uncoating and
Reverse Transcription of HIV-1. Annual Review of Virology 9:261-284.
31.   Jennings J, Bracey H, Hong J, Nguyen DT, Dasgupta R, Rivera AV, Sluis-Cremer N, Shi
J, Aiken C. 2024. The HIV-1 capsid serves as a nanoscale reaction vessel for reverse
transcription. PLOS Pathogens 20:e1011810.
32.   Christensen DE, Ganser-Pornillos BK, Johnson JS, Pornillos O, Sundquist WI. 2020.
Reconstitution and visualization of HIV-1 capsid-dependent replication and integration in
vitro. Science 370:eabc8420.
33.   Scott TM, Arnold LM, Powers JA, McCann DA, Rowe AB, Christensen DE, Pereira MJ,
Zhou W, Torrez RM, Iwasa JH, Kranzusch PJ, Sundquist WI, Johnson JS. 2025. Cell-free
assays reveal that the HIV-1 capsid protects reverse transcripts from cGAS immune
sensing. PLOS Pathogens 21:e1012206.

34. Yoh SM, Mamede JI, Lau D, Ahn N, Sánchez-Aparicio MT, Temple J, Tuckwell A, Fuchs NV, Cianci GC, Riva L, Curry H, Yin X, Gambut S, Simons LM, Hultquist JF, König R, Xiong Y, García-Sastre A, Böcking T, Hope TJ, Chanda SK. 2022. Recognition of HIV-1 capsid by PQBP1 licenses an innate immune sensing of nascent HIV-1 DNA. Molecular Cell 82:2871-2884.e6.

35. Stephens C, Naghavi MH. 2024. The host cytoskeleton: a key regulator of early HIV-1 infection. The FEBS Journal 291:1835-1848.

36. Fu L, Weiskopf EN, Akkermans O, Swanson NA, Cheng S, Schwartz TU, Görlich D. 2024. HIV-1 capsids enter the FG phase of nuclear pores like a transport receptor. Nature 626:843-851.

37. Kreysing JP, Heidari M, Zila V, Cruz-León S, Obarska-Kosinska A, Laketa V, Rohleder L, Welsch S, Köfinger J, Turoňová B, Hummer G, Kräusslich H-G, Beck M. 2025. Passage of the HIV capsid cracks the nuclear pore. Cell 188:930-943.e21.

38. Ravichandran SM, McFadden WM, Snyder AA, Sarafianos SG. 2024. State of the ART (antiretroviral therapy): Long-acting HIV-1 therapeutics. Global Health & Medicine 6:285-294.

39. Sever B, Otsuka M, Fujita M, Ciftci H. 2024. A Review of FDA-Approved Anti-HIV-1 Drugs, Anti-Gag Compounds, and Potential Strategies for HIV-1 Eradication. International Journal of Molecular Sciences 25:3659.

40. Blair WS, Pickford C, Irving SL, Brown DG, Anderson M, Bazin R, Cao J, Ciaramella G, Isaacson J, Jackson L, Hunt R, Kjerrstrom A, Nieman JA, Patick AK, Perros M, Scott AD, Whitby K, Wu H, Butler SL. 2010. HIV capsid is a tractable target for small molecule therapeutic intervention. PLoS Pathog 6:e1001220.

41. Dvory-Sobol H, Shaik N, Callebaut C, Rhee MS. 2022. Lenacapavir: a first-in-class HIV-1 capsid inhibitor. Current Opinion in HIV and AIDS 17:15-21.

42. McFadden WM, Faerch M, Kirby KA, Dick RA, Torbett BE, Sarafianos SG. 2025. Considerations for capsid-targeting antiretrovirals in pre-exposure prophylaxis. Trends in Molecular Medicine 31:801-813.

43. Lamorte L, Titolo S, Lemke CT, Goudreau N, Mercier JF, Wardrop E, Shah VB, von Schwedler UK, Langelier C, Banik SS, Aiken C, Sundquist WI, Mason SW. 2013. Discovery of novel small-molecule HIV-1 replication inhibitors that stabilize capsid complexes. Antimicrob Agents Chemother 57:4622-31.

44. Rihn SJ, Wilson SJ, Loman NJ, Alim M, Bakker SE, Bhella D, Gifford RJ, Rixon FJ, Bieniasz PD. 2013. Extreme Genetic Fragility of the HIV-1 Capsid. PLOS Pathogens 9:e1003461.

45. Faysal KMR, Walsh JC, Renner N, Márquez CL, Shah VB, Tuckwell AJ, Christie MP, Parker MW, Turville SG, Towers GJ, James LC, Jacques DA, Böcking T. 2024. Pharmacologic hyperstabilisation of the HIV-1 capsid lattice induces capsid failure. eLife 13:e83605.

46. Gres AT, Kirby KA, McFadden WM, Du H, Liu D, Xu C, Bryer AJ, Perilla JR, Shi J, Aiken C, Fu X, Zhang P, Francis AC, Melikyan GB, Sarafianos SG. 2023. Multidisciplinary studies with mutated HIV-1 capsid proteins reveal structural mechanisms of lattice stabilization. Nature Communications 14:5614.

47. Jacques DA, McEwan WA, Hilditch L, Price AJ, Towers GJ, James LC. 2016. HIV-1 uses dynamic capsid pores to import nucleotides and fuel encapsidated DNA synthesis. Nature 536:349-353.

48. Dick RA, Zadrozny KK, Xu C, Schur FKM, Lyddon TD, Ricana CL, Wagner JM, Perilla JR, Ganser-Pornillos BK, Johnson MC, Pornillos O, Vogt VM. 2018. Inositol phosphates are assembly co-factors for HIV-1. Nature 560:509-512.

49. Jennings J, Shi J, Varadarajan J, Jamieson Parker J, Aiken C. 2020. The Host Cell Metabolite Inositol Hexakisphosphate Promotes Efficient Endogenous HIV-1 Reverse Transcription by Stabilizing the Viral Capsid. mBio 11:10.1128/mbio.02820-20.

50. Highland CM, Tan A, Ricaña CL, Briggs JAG, Dick RA. 2023. Structural insights into HIV-1 polyanion-dependent capsid lattice formation revealed by single particle cryo-EM. Proceedings of the National Academy of Sciences 120:e2220545120.

51. Renner N, Kleinpeter A, Mallery DL, Albecka A, Rifat Faysal KM, Böcking T, Saiardi A, Freed EO, James LC. 2023. HIV-1 is dependent on its immature lattice to recruit IP6 for mature capsid assembly. Nature Structural & Molecular Biology 30:370-382.

52. Castellano P, Prevedel L, Valdebenito S, Eugenin EA. 2019. HIV infection and latency induce a unique metabolic signature in human macrophages. Scientific Reports 9:3941.

53. Clerc I, Abba Moussa D, Vahlas Z, Tardito S, Oburoglu L, Hope TJ, Sitbon M, Dardalhon V, Mongellaz C, Taylor N. 2019. Entry of glucose- and glutamine-derived carbons into the citric acid cycle supports early steps of HIV-1 infection in CD4 T cells. Nature Metabolism 1:717-730.

54. Sebigi TW, Asia LK, January GG, Jansen van Vuren E, Williams ME. 2025. The Tryptophan-Kynurenine pathway in people living with HIV: a systematic review. Infection 53:1625-1644.

55. Pornillos O, Ganser-Pornillos BK, Kelly BN, Hua Y, Whitby FG, Stout CD, Sundquist WI, Hill CP, Yeager M. 2009. X-ray structures of the hexameric building block of the HIV capsid. Cell 137:1282-92.

56. Nutan, Modi M, Goel T, Das T, Malik S, Suri S, Rawat AKS, Srivastava SK, Tuli R, Malhotra S, Gupta SK. 2013. Ellagic acid & gallic acid from *Lagerstroemia speciosa* L. inhibit HIV-1 infection through inhibition of HIV-1 protease & reverse transcriptase activity. Indian Journal of Medical Research 137:540–548.

57. Zhang X, Xia Q, Yang G, Zhu D, Shao Y, Zhang J, Cui Y, Wang R, Zhang L. 2017. The anti-HIV-1 activity of polyphenols from Phyllanthus urinaria and the pharmacokinetics and tissue distribution of its marker compound, gallic acid. Journal of Traditional Chinese Medical Sciences 4:158-166.

58. Kratz JM, Andrighetti-Fröhner CR, Kolling DJ, Leal PC, Cirne-Santos CC, Yunes RA, Nunes RJ, Trybala E, Bergström T, Frugulhetti IC. 2008. Anti-HSV-1 and anti-HIV-1 activity of gallic acid and pentyl gallate. Memórias do Instituto Oswaldo Cruz 103:437-442.

59. Kanehisa M, Sato Y, Kawashima M, Furumichi M, Tanabe M. 2016. KEGG as a reference resource for gene and protein annotation. Nucleic Acids Research 44:D457-D462.

60. Lorson ZC, McFadden WM, Neilsen G, Castaner AE, Slack RL, Kirby KA, Sarafianos SG. 2025. Characterization of antiviral compounds using Bio-Layer Interferometry. bioRxiv doi:10.1101/2025.07.24.662752:2025.07.24.662752.

61. Bruce A, Adebomi V, Czabala P, Palmer J, McFadden WM, Lorson ZC, Slack RL, Bhardwaj G, Sarafianos SG, Raj M. 2024. A Tag-Free Platform for Synthesis and Screening of Cyclic Peptide Libraries. Angewandte Chemie International Edition 63:e202320045.

62.   Neilsen G, Lan S, Slack RL, Lorson ZC, Emanuelli Castaner A, Lee R, Edwards KG,
Zhang H, Lee J, Cantara WA, Cilento ME, Zhang H, De R, Amblard F, Tedbury PR, Kirby
KA, Schinazi RF, Sarafianos SG. 2025. Strategy to overcome a nirmatrelvir resistance
mechanism in the SARS-CoV-2 nsp5 protease. Science Advances 11:eadv8875.

63.   Precupas A, Leonties AR, Neacsu A, Sandu R, Popa VT. 2019. Gallic acid influence on
bovine serum albumin thermal stability. New Journal of Chemistry 43:3891-3898.

64.   Nikolaev YV, Kochanowski K, Link H, Sauer U, Allain FHT. 2016. Systematic
Identification of Protein–Metabolite Interactions in Complex Metabolite Mixtures by
Ligand-Detected Nuclear Magnetic Resonance Spectroscopy. Biochemistry 55:2590-
2600.

65.   McFadden WM, Kirby KA, Wang L, Du H, Lorson ZC, Zhang H, Castaner AE, Brancato
S, Hachiya A, Ravichandran SM, Highland CM, Tedbury PR, Dick RA, Wang Z,
Sarafianos SG. 2025. Modifying PF74 Improves Anti-HIV-1 Activity Against the
Resistance-Associated Capsid Mutation N74D. [CROI Abstract 727]. Topics in Antiviral
Medicine 33:208.

66.   Matreyek KA, Yücel SS, Li X, Engelman A. 2013. Nucleoporin NUP153 Phenylalanine-
Glycine Motifs Engage a Common Binding Pocket within the HIV-1 Capsid Protein to
Mediate Lentiviral Infectivity. PLOS Pathogens 9:e1003693.

67.   Sanna C, Marengo A, Acquadro S, Caredda A, Lai R, Corona A, Tramontano E, Rubiolo
P, Esposito F. 2021. In Vitro Anti-HIV-1 Reverse Transcriptase and Integrase Properties
of *Punica granatum* L. Leaves, Bark, and Peel Extracts and Their Main Compounds. Plants
10:2124.

68.   Lanman J, Sexton J, Sakalian M, Prevelige PE, Jr. 2002. Kinetic analysis of the role of
intersubunit interactions in human immunodeficiency virus type 1 capsid protein assembly
in vitro. J Virol 76:6900-8.

69.   Barklis E, Alfadhli A, McQuaw C, Yalamuri S, Still A, Barklis RL, Kukull B, López CS.
2009. Characterization of the in vitro HIV-1 capsid assembly pathway. Journal of
Molecular Biology 387:376-389.

70.   Michailidis E, Marchand B, Kodama EN, Singh K, Matsuoka M, Kirby KA, Ryan EM,
Sawani AM, Nagy E, Ashida N, Mitsuya H, Parniak MA, Sarafianos SG. 2009. Mechanism
of Inhibition of HIV-1 Reverse Transcriptase by 4′-Ethynyl-2-fluoro-2′-deoxyadenosine
Triphosphate, a Translocation-defective Reverse Transcriptase Inhibitor. Journal of
Biological Chemistry 284:35681-35691.

71.   Michailidis E, Huber AD, Ryan EM, Ong YT, Leslie MD, Matzek KB, Singh K, Marchand
B, Hagedorn AN, Kirby KA, Rohan LC, Kodama EN, Mitsuya H, Parniak MA, Sarafianos
SG. 2014. 4'-Ethynyl-2-fluoro-2'-deoxyadenosine (EFdA) Inhibits HIV-1 Reverse
Transcriptase with Multiple Mechanisms. Journal of Biological Chemistry 289:24533-
24548.

72.   Wright AM, Schwartz RS, Oaks JR, Newman CE, Flanagan SP. 2019. The why, when, and
how of computing in biology classrooms. F1000Res 8:1854.

73.   Xu C, Fischer DK, Rankovic S, Li W, Dick RA, Runge B, Zadorozhnyi R, Ahn J, Aiken
C, Polenova T, Engelman AN, Ambrose Z, Rousso I, Perilla JR. 2020. Permeability of the
HIV-1 capsid to metabolites modulates viral DNA synthesis. PLOS Biology 18:e3001015.

74.   Álvarez M, Sapena-Ventura E, Luczkowiak J, Martín-Alonso S, Menéndez-Arias L. 2021.
Analysis and Molecular Determinants of HIV RNase H Cleavage Specificity at the PPT/U3
Junction. Viruses 13:131.

75. Badhani B, Sharma N, Kakkar R. 2015. Gallic acid: a versatile antioxidant with promising
therapeutic and industrial applications. RSC Advances 5:27540-27557.
76. Hadidi M, Liñán-Atero R, Tarahi M, Christodoulou MC, Aghababaei F. 2024. The
Potential Health Benefits of Gallic Acid: Therapeutic and Food Applications. Antioxidants
13:1001.
77. Zhang Y, Wen Y, Xue W, Zhang T, Hou J, Chen X, Feng R, Tan C. 2025. Gallic acid
inhibits EMCV infection via targeting the interaction between TBK1 and IRF3 to promote
IFN-β expression. Microbiology Spectrum 0:e01524-25.
78. You H-L, Huang C-C, Chen C-J, Chang C-C, Liao P-L, Huang S-T. 2018. Anti-pandemic
influenza A (H1N1) virus potential of catechin and gallic acid. Journal of the Chinese
Medical Association 81:458-468.
79. Liu Y, Carver JA, Calabrese AN, Pukala TL. 2014. Gallic acid interacts with α-synuclein
to prevent the structural collapse necessary for its aggregation. Biochimica et Biophysica
Acta 1844:1481-1485.
80. Zhang Y, Dong L, Li J, Chen X. 2008. Studies on the interaction of gallic acid with human
serum albumin in membrane mimetic environments. Talanta 76:246-253.
81. Sang H, Jin H, Song P, Xu W, Wang F. 2024. Gallic acid exerts antibiofilm activity by
inhibiting methicillin-resistant Staphylococcus aureus adhesion. Scientific Reports
14:17220.
82. Frias MA, Pagano S, Bararpour N, Sidibé J, Kamau F, Fétaud-Lapierre V, Hudson P,
Thomas A, Lecour S, Strijdom H. 2024. People living with HIV display increased anti-
apolipoprotein A1 auto-antibodies, inflammation, and kynurenine metabolites: a case–
control study. Frontiers in Cardiovascular Medicine 11:1343361.
83. Lu G, Matsuura SE, Barrientos A, Scott WA. 2013. HIV-1 Infection Is Blocked at an Early
Stage in Cells Devoid of Mitochondrial DNA. PLOS ONE 8:e78035.
84. Morgan M, Ramos M. 2024. BiocManager: Access the Bioconductor Project Package
Repository, v1.30.25. https://CRAN.R-project.org/package=BiocManager.
85. Posit team. 2025. RStudio: Integrated Development Environment for R, Posit Software,
PBC, Boston, MA. http://www.posit.co/.
86. Wickham H, Hester J, Chang W, Bryan J. 2022. devtools: Tools to Make Developing R
Packages Easier, v2.4.5. https://CRAN.R-project.org/package=devtools.
87. Wickham H, Averick M, Bryan J, Chang W, McGowan LDA, François R, Grolemund G,
Hayes A, Henry L, Hester J. 2019. Welcome to the Tidyverse. Journal of Open Source
Software 4:1686.
88. Ooms J. 2014. The jsonlite Package: A Practical and Consistent Mapping Between JSON
Data and R Objects. arXiv:1403 2805 [stat CO].
89. Schauberger P, Walker A. 2025. openxlsx: Read, Write and Edit xlsx Files, v4.2.8.
https://CRAN.R-project.org/package=openxlsx.
90. Wood SN. 2011. Fast stable restricted maximum likelihood and marginal likelihood
estimation of semiparametric generalized linear models. Journal of the Royal Statistical
Society (B) 73:3-36.
91. Wood SN, Pya N, Saefken B. 2016. Smoothing parameter and model selection for general
smooth models. Journal of the American Statistical Association 111:1548-1575.
92. Wood SN. 2017. Generalized Additive Models: An Introduction with R.
93. Elzhov TV, Mullen KM, Spiess A-N, Bolker B. 2023. minpack.lm: R Interface to the
Levenberg-Marquardt Nonlinear Least-Squares

Algorithm Found in MINPACK, Plus Support for Bounds, v1.2-4. https://CRAN.R-
project.org/package=minpack.lm.
94. Perrier V, Meyer F, Granjon D. 2023. shinyWidgets: Custom Inputs Widgets for Shiny,
v0.9.0. https://CRAN.R-project.org/package=shinyWidgets.
95. Sievert C. 2020. Interactive Web-Based Data Visualization with R, plotly, and shiny.
Chapman and Hall/CRC.
96. Attali D. 2021. shinyjs: Easily Improve the User Experience of Your Shiny Apps in
Seconds, v2.1.0. https://CRAN.R-project.org/package=shinyjs.
97. Owen J. 2021. rhandsontable: Interface to the 'Handsontable.js' Library, https://CRAN.R-
project.org/package=rhandsontable.
98. Kassambara A. 2023. ggpubr: 'ggplot2' Based Publication Ready Plots, vR package version
0.6.0. https://CRAN.R-project.org/package=ggpubr.
99. Cercignani C. 1988. The Boltzmann Equation, p 40-103. *In* Cercignani C (ed), The
Boltzmann Equation and Its Applications doi:10.1007/978-1-4612-1039-9_2. Springer
New York, New York, NY.
100. Schroeder DV. 2020. An introduction to thermal physics. Oxford University Press.
101. Christopoulos D. 2016. On the Efficient Identification of an Inflection Point. International
Journal of Mathematics And Scientific Computing 6:13-20.
102. Le Grice SFJ, Grüninger-Leitch F. 1990. Rapid purification of homodimer and heterodimer
HIV-1 reverse transcriptase by metal chelate affinity chromatography. European Journal
of Biochemistry 187:307-314.
103. Wang L, Casey MC, Vernekar SKV, Do HT, Sahani RL, Kirby KA, Du H, Hachiya A,
Zhang H, Tedbury PR, Xie J, Sarafianos SG, Wang Z. 2020. Chemical profiling of HIV-1
capsid-targeting antiviral PF74. European Journal of Medicinal Chemistry 200:112427.
104. Cilento ME, Wen X, Reeve AB, Ukah OB, Snyder AA, Carrillo CM, Smith CP, Edwards
K, Wahoski CC, Kitzler DR, Kodama EN, Mitsuya H, Parniak MA, Tedbury PR,
Sarafianos SG. 2023. HIV-1 Resistance to Islatravir/Tenofovir Combination Therapy in
Wild-Type or NRTI-Resistant Strains of Diverse HIV-1 Subtypes. Viruses 15.
105. Connor RI, Chen BK, Choe S, Landau NR. 1995. Vpr Is Required for Efficient Replication
of Human Immunodeficiency Virus Type-1 in Mononuclear Phagocytes. Virology
206:935-944.
106. Gludish DW, Boliar S, Caldwell S, Tembo DL, Chimbayo ET, Jambo KC, Mwandumba
HC, Russell DG. 2020. TZM-gfp cells: a tractable fluorescent tool for analysis of rare and
early HIV-1 infection. Scientific Reports 10:19900.
863

A Modeled vs raw melting curves

Thermal Profile of CA121
With: Folic Acid

Tm=47.3C

B Compare Plots of Vitamins vs DMSO Melting Curves

CA121 + IP6
Tm = 14C

CA121 + PxyAmine diHCl
Tm = 0.3C

CA121 + Folic Acid
Tm = 3C

CA121 + Thiamine HCl
Tm = 0.1C

C Boxplot of Vitamin Library T_m

Ligand

- DMSO
- IP6
- B5
- Biotin
- B3
- 4-ABA
- PxyAmine diHCl
- PyxalHCl
- Folic Acid
- Thiamine HCl
- =+@LA
- PyxINE HCl

A Initial ΔT$_m$ Hits from Organic Acid Library

Phytic acid (IP6)
ΔT$_m$ = 12°C

Gallic acid
ΔT$_m$ = 6.37°C

2-Oxoglutaric acid
ΔT$_m$ = 5.79°C

2-Methylcitric acid
ΔT$_m$ = 5.77°C

Oxalacetic acid
ΔT$_m$ = 4.99°C

Citric acid
ΔT$_m$ = 4.10°C

1,2,3-Propanetricarboxylic
ΔT$_m$ = 4.10°C

Isocitric acid
ΔT$_m$ = 4.05°C

2-hydroxy-3-(4-hydroxy-3-methoxyphenyl)propanoic acid
ΔT$_m$ = 3.04°C

2-oxoadipic acid
ΔT$_m$ = 2.86°C

Xanthurenic acid
ΔT$_m$ = 2.28°C

Aconitic acid
ΔT$_m$ = 1.65°C

Indole-3-pyruvic acid
ΔT$_m$ = 1.65°C

Indole-3-butyric acid
ΔT$_m$ = 1.50°C

2-Isopropylmalic acid
ΔT$_m$ = 1.40°C

3-hydroxy-3-methylglutaric acid
ΔT$_m$ = 1.34°C

Quinolinic acid
ΔT$_m$ = 1.34°C

Prephenic acid
ΔT$_m$ = 1.13°C

Indole-3-propionic acid
ΔT$_m$ = 1.12°C

2,5-furandicarboxylic acid
ΔT$_m$ = 1.11°C

B CA121 + Phytic Acid

C CA121 + Gallic Acid

D CA121 + Citric Acid

E Pathway Analysis

Metabolic pathways — 24.5%
2−Oxocarboxylic acid metabolism — 46.7%
Biosynthesis of amino acids — 50.0%
Biosynthesis of cofactors — 58.3%
Citrate cycle (TCA cycle) — 62.5%
Butanoate metabolism — 14.3%
Lipoic acid metabolism — 50.0%
Pyrimidine metabolism — 0.0%
cAMP signaling pathway — 0.0%
GABAergic synapse — 33.3%
Inositol phosphate metabolism — 100.0%
Oxidative phosphorylation — 0.0%

Full Library
TSAR Hits

KEGG Match Count

F Biolayer Interferometry:
CA^{HEX} with PF74

Wavelength shift (nm)
Time (s)
— 5 µM — 10 µM — 20 µM

G CA^{HEX} with gallic acid

Wavelength shift (nm)
Time (s)
— 10 µM — 20 µM — 50 µM

A

Compound	EC$_{50}$ (µM)	CC$_{50}$ (µM)
PF74	0.605 ± 0.0313	75.29 ± 3.9†
Gallic acid	19.24 ± 0.1131	>100
Isocitric acid	>100	>100
Folic acid	>100	>100
2-methyl citric acid	>100	>100
Xanthurenic acid	>100	>100

B ERT: minus-strand strong stop (early)

	IC$_{50}$ (µM)
● PF74	12.58
■ Gallic acid	ND
▲ Xanthurenic acid	11.21

C ERT: First strand transfer (Intermediate)

	IC$_{50}$ (µM)
● PF74	1.95
■ Gallic acid	17.21
▲ Xanthurenic acid	11.99

D ERT: Second strand transfer (Late)

	IC$_{50}$ (µM)
● PF74	1.32
■ Gallic acid	3.64
▲ Xanthurenic acid	9.65

E WT CA + IP6

F WT CA + Gallic Acid + IP6

G WT CA + Xanth. Acid + IP6

H WT CA + αKG + IP6

A

Legend
- Control (black line)
- Treatment (purple line)

Stabilization ΔT$_m$

Denatured
Denatured
Partially denatured
Partially denatured
Folded
Folded

Fluorescence (RFU)

Temperature (25°C)   T$_m$ (Vehicle)   T$_m$ (Treatment)   Temperature (95°C)

B Example Boltzmann Fit

$$y = \frac{1}{1 + e^{(-k \cdot (x - T_mB))}}$$

Normalized

Temperature

T$_m$D
T$_m$B

- RFU
- Boltzmann-fit

C Melting Temperature Estimate Comparison by Model

Estimates by Boltzmann Model

Estiamtes by Derivative Model

D Distribution of T$_m$D Errors (n=54)

count

Error (°C)

E Distribution of T$_m$B Errors (n=54)

count

Error (°C)

F Distribution of ΔT$_m$D Errors (n=12)

count

Error (°C)

G Distribution of ΔT$_m$B Errors (n=12)

count

Error (°C)

A