The hydrogen isotope signatures of autotrophy versus heterotrophy recorded in archaeal tetraether lipids

Authors: Jeemin H. Rhim[1,*], Sebastian Kopf[2], Jamie McFarlin[2,3], Harpreet Batther[2], Carolynn M. Harris[1], Alice Zhou[1], Xiahong Feng[1], Yuki Weber[5], Shelley Hoeft-McCann[5], Ann Pearson[5], and William D. Leavitt[1,6]

[1] Department of Earth Sciences, Dartmouth College
[2] Department of Geological Sciences, University of Colorado Boulder
[3] Department of Geology & Geophysics, University of Wyoming
[4] Department of Earth & Planetary Sciences, University of Michigan
[5] Department of Earth & Planetary Sciences, Harvard University
[6] Department of Chemistry, Dartmouth College

* Corresponding author: jrhim@ucsb.edu

^Current address: Jeemin H. Rhim, Department of Ecology, Evolution, and Marine Biology, University of California Santa Barbara, Santa Barbara, CA, USA; TBD.

Keywords: Archaeal lipids, tetraether lipids, GDGTs, biphytanes, hydrogen isotope, biomarkers

Abstract

The stable hydrogen isotope composition of archaeal lipids is emerging as a potential paleoenvironmental proxy, adding to the well-established application of plant leaf wax-derived *n*-alkanes in paleohydrological reconstruction. A handful of studies reported relatively invariant and depleted hydrogen isotope compositions for archaeal lipids despite the range of different organisms and growth conditions explored. However, how modes of metabolism affect the hydrogen isotope signatures of archaeal lipids remains poorly understood, limiting our ability to interpret archaeal lipid biomarker records from the environment. Here we conducted water isotope label experiments with a metabolically flexible and well-studied model archaeon *Archaeoglobus fulgidus* and quantified the hydrogen isotope fractionation between lipids and water in response to different carbon substrates and electron donor-acceptor pairs. The $^2H/^1H$ fractionation between lipids and water ($\varepsilon_{L/W}$) was overall negative, ranging from –280 to –226 ‰, and overlapped with the ranges observed for other archaea in previous studies. Isotope flux-balance model results suggest that ≥80 % and ≥50 % of lipid-bound H in *A. fulgidus* cultures originate from water during autotrophy and heterotrophy, respectively. The model results also suggest the final saturation during isoprenoid lipid biosynthesis catalyzed by a flavoenzyme geranylgeranyl reductase (GGR) likely contributes to the large $^2H/^1H$ fractionation observed in this study, consistent with isotope flux-balance model results for a different archaeon. Finally, we synthesized available data to compare $\varepsilon_{L/W}$ patterns across all three domains of life: Bacteria, Archaea and Eukarya. Emerging patterns between archaeal and eukaryotic lipids are consistent with the notion of highly

41 fractionating GGR, and the patterns between archaeal and bacterial lipids suggest that the general
42 state of energy limitation may also contribute to large, negative values of $\varepsilon_{L/w}$ observed in
43 prokaryotic lipids. Altogether, these findings lend further support for the potential of archaeal lipid
44 $\varepsilon_{L/w}$ as a paleohydrological proxy and provide a broader insight into the $^2H/^1H$ fractionation
45 mechanisms potentially shared among prokaryotic and eukaryotic lipid biomarkers.

1. Introduction

Hydrogen (H) plays a crucial role in most metabolic processes and is incorporated into biological
molecules, often with distinct isotope fractionation patterns. Consequently, the ratio of stable
hydrogen isotopes ($^2H/^1H$) of certain biomolecules can be used to track the H isotope compositions
of source water, or the operation of specific biosynthetic pathways (Estep and Hoering, 1980;
Sessions et al., 1999; Hayes, 2001). Lipid-bound H in hydrocarbons may resist isotope exchange
over geologic time scales for up to 10^4 to 10^8 years (Sessions et al., 2004). Thus, sedimentary
record of lipid δ^2H can be applied to reconstruct past ecological or hydrological states (Sachse et
al., 2012; Sessions, 2016; McFarlin et al., 2019). The δ^2H of lipids produced by photoautotrophic
eukaryotes as well as chemoautotrophic and heterotrophic bacteria have been studied in detail for
this reason (e.g., Sessions et al., 1999; Chikaraishi and Naraoka, 2003; Zhang and Sachs, 2007;
Zhang et al., 2009; Sachse et al., 2012; Sachs, 2014; Dawson et al., 2015; Osburn et al., 2016;
Leavitt et al., 2016; Wijker et al., 2019; McFarlin et al., 2019). The H isotope data on lipids
produced by archaea, in contrast, are limited to only a handful of studies (e.g., Kaneko et al., 2011;
Dirghangi and Pagani, 2013; Wu et al., 2020; Lengger et al., 2021; Leavitt et al., 2023). As a result,
factors controlling the H isotope fractionation between lipid and water ($\varepsilon_{L/w}$) as well as the extent
and limits of proxy applications are relatively well understood for eukaryotic and bacterial lipids,
yet remain poorly understood for archaeal lipids.

The $\varepsilon_{L/w}$ values reflect a combination of source water δ^2H and the kinetic isotope effect (KIE) of
enzymes involved in lipid biosynthesis. For lipids produced by photoautotrophic eukaryotes (e.g.,
plants and algae), KIE is relatively invariant but can be altered by some environmental factors
(e.g., temperature, salinity, irradiance) and some factors intrinsic to an organism (e.g. growth rate
in response to different stimuli and underlying genetics) (Sachs, 2014; Van Der Meer et al., 2015;
Sachs and Kawka, 2015; Sachs et al., 2016, 2017; Maloney et al., 2016; Ladd and Sachs, 2017;
Wolfshorndl et al., 2019; Bender, n.d.). In general, these are well studied in eukaryotes, and have
been accounted for in hydroclimate proxy studies (Sachse et al., 2012; Sessions, 2016; McFarlin
et al., 2019).

Bacterial lipids show significantly more range in $\varepsilon_{L/w}$ than eukaryotic lipids. Depending on the
central metabolism a given bacteria operates, $\varepsilon_{L/w}$ can range from –400 to –200 ‰ (exclusively
negative values) for (an)aerobic chemoautotrophs to –154 to +380 ‰, associated with aerobic
heterotrophs (Sessions et al., 2002; Campbell et al., 2009; Zhang et al., 2009; Dawson et al., 2015;
Osburn et al., 2016; Leavitt et al., 2016; Wijker et al., 2019). For lipids produced by aerobic

heterotrophs, H fluxes through $NADP^+$-reducing and NADPH-balancing reactions provide a
quantitative explanation for the full range of $\varepsilon_{L/W}$ values observed (Wijker et al., 2019). Bacterial
lipids found in sediments, especially those produced by aerobic heterotrophs, may have more
useful applications as proxies for microbial metabolism and ecology rather than past hydroclimate
(Zhang et al., 2009; Wijker et al., 2019).

Less is known about the type(s) of information archaeal lipids record. Studies to-date indicate that
archaeal lipids are generally depleted in ^{2}H, with $\varepsilon_{L/W}$ values comparable to those observed in
eukaryotic photoautotrophs and bacterial chemoautotrophs (Kaneko et al., 2011; Dirghangi and
Pagani, 2013; Wu et al., 2020; Lengger et al., 2021; Leavitt et al., 2023). In one recent study,
continuous cultures of the obligate autotroph *Nitrosopumilus maritimus* revealed that $\varepsilon_{L/W}$ remains
nearly constant across a 3-fold range in growth rate (Leavitt et al., 2023). Other studies have
reported similarly large, negative values of $\varepsilon_{L/W}$ for archaea grown under heterotrophic conditions
(Kaneko et al., 2011; Dirghangi and Pagani, 2013; Wu et al., 2020; Lengger et al., 2021). However,
it remains to be determined whether and how carbon metabolism affects $\varepsilon_{L/W}$ within a single
organism. Testing this is important, considering that mixotrophy (using both autotrophic and
heterotrophic carbon sources) is a widespread mode of metabolism in nature and that many archaea
are adapted to fluctuating environments. For bacterial lipids, changes in carbon metabolism
resulted in significantly different $\varepsilon_{L/W}$ values within a single organism for some species (e.g.,
Cupriavidus oxalaticus and *Paracoccus denitrificans*; Zhang et al., 2009; Osburn et al., 2016), but
not in other species (sulfate-reducing bacteria; Dawson et al., 2015; Osburn et al., 2016; Leavitt et
al., 2016). As such, understanding the role of carbon metabolism in setting archaeal lipid-H isotope
compositions is important for interpreting archaeal lipids record from the environment.
Furthermore, our current understanding of broader patterns of $\varepsilon_{L/W}$ that emerge from comparison
across lipids produced by all three domains of life—Archaea, Bacteria, and Eukarya—is limited,
despite the similarities observed among certain subgroups of different domains.

In this study, we first conducted pure culture experiments to better understand factors influencing
$\varepsilon_{L/W}$ in archaeal lipids. A metabolically flexible model archaeon *Archaeoglobus fulgidus* was
grown in batch cultures on different carbon substrates and electron donor-acceptor pairs (lactate-
thiosulfate, lactate-sulfate, and H_2/CO_2-thiosulfate) to test the effect of metabolism on $\varepsilon_{L/W}$ within
a single organism. All conditions included replicates with different water isotope compositions to
track the fraction of water that gets incorporated into the final product lipid. The effect of growth
phase was also assessed by comparing biomass harvested at exponential and stationary growth
phases. We interpret the regression parameters derived from the water isotope label experiments
with an isotope flux-balance model adapted from Leavitt et al. (2023). The model is informed by
the enzymatic steps involved in the Wood-Ljungdahl (reductive acetyl-CoA) pathway used by *A.*
fulgidus as well as archaeal isoprenoid lipid biosynthesis used by all tetraether-producing archaea.
In addition to *A. fulgidus*, two thermoacidophilic archaea (*Acidianus* sp. DS80 and *Metallosphaera*
sedula DSM 5348T) were grown in batch cultures for additional $\varepsilon_{L/W}$ measurements. We present

a synthesis of representative $\varepsilon_{L/w}$ values observed in Archaea, Bacteria, and Eukarya and discuss
factors that may underlie some of the similarities observed across different domains of life.

2.   Materials and Methods

2.1.   Culture Strains and Growth

124
Archaeoglobus fulgidus str. VC-16 (DSM4304) was obtained from Deutsche Sammlung von
Mikroorganismen und Zellkulturen (Braunschweig, Germany) and grown in batch cultures on two
different carbon substrates and electron acceptors (Table 1). Culture media was prepared following
the DSMZ 399 recipe with modifications as follows (per liter): 0.34 g KCl, 4.00 g $MgCl_2$ x 6 H_2O,
3.45 g $MgSO_4$ x 7 H_2O, 0.25 g NH_4Cl, 0.14 g $CaCl_2$ x 2 H_2O, 0.11 g K_2HPO_4 x 3 H_2O, 0.20 g
KH_2PO_4, 18.00 g NaCl, 2.52 g $NaHCO_3$, 2 mL $Fe(NH_4)_2(SO_4)_2$ x 7 H_2O (0.1% w/v), and 10 mL
trace element (DSMZ 141). After autoclaving, sterile anoxic solutions were added to the medium
to a final concentration of 3.2 mM L-cysteine-HCl and 1% (v/v) vitamin (DSMZ 141). When
thiosulfate was used as an electron acceptor, 4.00 g/L $MgCl_2$ x 6 H_2O and 3.45 g/L $MgSO_4$ x 7
H_2O were replaced with 4.70 g/L $MgCl_2$ x 6 H_2O and 3.48 g/L $Na_2S_2O_3$ x 5 H_2O. For heterotrophic
growth, lactate was used as an electron donor and carbon source, 20 mM of Na-lactate solution
was added and incubated with approximately 1 bar of N_2/CO_2 (80:20, v/v) headspace; for
autotrophic growth, lactate was omitted and H_2 and CO_2 were used as an electron donor and carbon
source, respectively, with 21 psig H_2/CO_2 (80:20, v/v) in the headspace. The headspace of
autotrophic cultures were flushed and re-pressurized with H_2/CO_2 once per day after each OD
measurement. The δ^2H of culture media was manipulated by volumetrically diluting 99.9 % purity
^2H_2O.

142
All cultures of *A. fulgidus* were incubated at 82 °C. Heterotrophic cultures were incubated
statically, and autotrophic cultures were shaken at 200 rpm to increase gas exchange between the
H_2/CO_2 headspace and culture media. The optical density at 600 nm (OD_{600}) of samples was used
for monitoring growth, and linearity against direct cell counts was confirmed (depth 0.02 mm;
Petroff-Hausser Counter 3900) throughout the absorbance range. Biomass samples were harvested
twice per experiment, one during mid-exponential phase and the other during early stationary
phase. To harvest, 50 mL of culture was centrifuged for 15 minutes at 4 °C and 4500 x *g*, and
resulting wet cell pellets were stored at –80 °C until lipid extraction.

151
Batch cultures of *Acidianus* sp. DS80 and *M. sedula* DSM 5348T were grown for additional
archaeal lipid $\varepsilon_{L/w}$ measurements (not for water isotope labeling experiments or isotope flux-
balance model). *Acidianus* sp. DS80 was grown as previously described in Rhim et al. (2023), and
M. sedula was grown in Brock basal salts medium with Allen's trace element mixture (Brock et
al., 1972). See Supporting Information for additional details for growth conditions for *Acidianus*
sp. DS80 and *M. sedula*.
158

159   2.2.   Lipid Extraction, Ether Cleavage, and Quantification
160
Frozen cell pellets were lyophilized overnight, and dry cell pellets were extracted using acidic
hydrolysis-methanolysis. Briefly, dry cell pellets were physically disrupted in 1.5-mL
microcentrifuge tubes by vortexing with methanol (MeOH) and *ca.* 250 µL of 100 µm combusted
glass beads for 10 minutes at 3000 rpm using a Disruptor Genie (Scientific 126 Industries, SI-
DD38). After excess MeOH is evaporated, octacosanol (C_{28}) or lignocerol (C_{24}) and C_{46} GTGT
were added to all samples as internal extraction standards. Samples were incubated at 65 °C for 90
minutes with 3 N methanolic HCl (33 % water content) to hydrolyze tetraether headgroups. Methyl
tert butyl ether (MTBE) was added to the cooled samples at a ratio of 3:2 (acid:MTBE, v/v), and
samples were sonicated for 5 minutes (Qsonica Q500, Newtown, CT, USA). After sonication, n-
hexane was added at a 1:1 ratio (MTBE:hexane, v/v), vortexed, and centrifuged at 15,000 g for 3
minutes. The upper organic phase was collected three times with n-hexane for the two subsequent
rounds. The resulting total lipid extracts (TLEs) were dried under N_2 and proceeded to ether
cleavage or stored at –20 °C until ether cleavage.
174
Ether bonds were cleaved in 57% hydroiodic acid (HI) at 125 °C for 4 hours. Then the organic
phase was collected three times using n-hexane. The resulting alkyl iodides were reduced to
alkanes (biphytanes, BPs) with H_2 in the presence of $Pt^{(IV)}O_2$ (Kaneko et al., 2011). Hydrogenated
samples were passed through combusted pasteur pipettes with glass wool plugs to remove PtO_2
powder. Filtered samples were dried under N_2 and stored at –20 °C until analysis. The resulting
ether-cleaved BPs were analyzed on single quadrupole gas chromatography-mass spectrometer
(GC-MS; Thermo ISQ LT with TRACE 1310) for compound identification and on GC-flame
ionization detector (GC-FID; Thermo TRACE 1310) for quantification, both housed in the Organic
Geochemistry Laboratory at the University of Colorado Boulder.
184
185   2.3.   Hydrogen isotope analysis and data processing
186
The δ^2H of BPs was analyzed on a GC-pyrolysis-isotope ratio MS (GC-P-IRMS) on a GC IsoLink
II IRMS System (Thermo Scientific) in the Organic Geochemistry Laboratory at the University of
Colorado Boulder. The GC-P-IRMS system consists of a Trace 1310 GC fitted with a
programmable temperature vaporization (PTV) injector and either a 30 m ZB5HT column (i.d. =
0.25 mm, 0.25 µm, Phenomenex, Torrance, CA, USA) or a 60 m DB1 column (i.d. = 0.25 mm,
0.25 µm, Agilent, Santa Clara, CA, USA), ConFlo IV interface, and MAT 253 Plus mass
spectrometer (Thermo Scientific).
194
A. fulgidus has been observed to produce both tetraether and diether lipids, with the majority of
GDGT with no ring (GDGT-0) and a small fraction with up to two pentacyclic rings (Lai et al.,
2008). The tetraether-to-diether ratio for *A. fulgidus* is expected to be ~1:1 at the growth
temperature in this study (82 °C; Lai et al., 2008), and we focus on the tetraether fraction for the

rest of the discussion. In line with the previous observation, the majority of BPs derived from *A.*
fulgidus GDGT did not have rings (BP-0). All of the δ^2H values reported in this study are for BP-
0.
202
203   All hydrogen isotope ratios are reported in the standard delta notation (δ^2H) in permil (‰) units,
204   against the Vienna Standard Mean Ocean Water (VSMOW) international standard:
205

$$\delta\,2H \;=\; \left[\frac{(2H/\,1H)_{sample}}{(2H/\,1H)_{VSMOW}} - 1\right] \times 1000 \qquad \text{Eq. (1)}$$

206
207
208   The lipid-water hydrogen isotope fractionation factor is reported in alpha notation ($^2\alpha_{L/W}$), and
209   fractionation is reported in epsilon notation ($^2\varepsilon_{L/W}$) in ‰:
210

$$2\alpha_{L/W} \;=\; \frac{(2H/\,1H)_{lipid}}{(2H/\,1H)_{water}} \qquad \text{Eq. (2)}$$

211
212

$$2\varepsilon_{L/W} \;=\; \lfloor 2\alpha_{L/W} - 1\rfloor \times 1000 \qquad \text{Eq. (3)}$$

213
214
Values of δ^2H were first determined relative to H_2 reference gas (δ^2H_{raw}), and then calibrated
externally using a standard *n*-alkane mixture (A6, containing C_{15} through C_{30} n-alkanes spanning
from –9 to –263 ‰ vs. VSMOW; A. Schimmelmann, Indiana University). The A6 standard was
combined with a C_{36} n-alkane (nC_{36}, –259.2 ‰ vs. VSMOW; A. Schimmelmann, Indiana
University) and measured at regular intervals at different concentrations. The BP δ^2H calibration
was done in R, using the packages *isoreader* (Kopf et al., 2021) and *isoprocessor* available at
github.com/isoverse. Details of the data reduction and calibration methods are described in Leavitt
et al. (2022).
223
The δ^2H of the H_2 gas in H_2/CO_2 headspace of autotrophic cultures was analyzed on Thermo
Gasbench II in the Stable Isotope Laboratory at Northwestern University. The δ^2H of non-
exchangeable H in lactate used for heterotrophic cultures was assumed to be –60 ‰. The δ^2H of
natural abundance and deuterium-enriched medium water was measured at the Stable Isotope
Laboratory at Dartmouth College as previously described (Kopec et al., 2019). Briefly, water was
reduced with chromium at 850 °C using the H-Device, and the δ^2H of the resulting gas was
measured by an IRMS (Thermo Delta Plus XL). The measured δ^2H values were converted to the
water δ^2H equivalent by calibrating with water standards with known isotopic compositions
analyzed in the same way. For samples expected to be outside the range of the standards (e.g., the
deuterium-enriched medium), samples were diluted with water standards. The 1σ uncertainty for
δ^2H measurements is <0.5 ‰.
235
236   2.4.   Model implementation
237

An isotope flux-balance model was adapted from the conceptual framework described in Leavitt
et al. (2023). Our model considers the Wood-Ljungdahl (reductive acetyl-CoA) pathway used by
A. fulgidus for determining the sources and fluxes of H during acetyl-CoA (Ac-CoA) synthesis. It
also considers the general archaeal lipid biosynthesis pathway for the accounting of total H budget
in archaeal lipids. Analytical solutions for Eqn. 5–12 were obtained via R. Details of the model
design and outcomes are described in section 4.

3.   Results

3.1.   Cultures and growth rates
246

Archaeoglobus fulgidus is a metabolically flexible strain from the phylum Euryarchaeota. We
cultivated *A. fulgidus* in batch (closed-system) cultures on three different combinations of electron
donors and acceptors (Table 1; Fig. 1A–C). In two treatments, *A. fulgidus* was grown as a
heterotroph, using lactate as the carbon source and electron donor and sulfate or thiosulfate as the
terminal electron acceptor. In the third treatment, *A. fulgidus* was grown as an autotroph, using
CO_2 as the carbon source, H_2 as the electron donor, and thiosulfate as the terminal electron
acceptor. Cultures were acclimated to each treatment by growing on respective substrate
combinations for two or more passages before inoculation. Overall, heterotrophic conditions
resulted in faster growth (T-L, t_d = 6.2 ± 0.6 h; S-L, t_d = 10.3 ± 1.7 h) compared to autotrophic
growth (T-H_2/CO_2, t_d = 19.3 ± 3.9 h) (Table 1; Fig. 1A–B).
257

258 **Table 1.** Summary of culture experiments.
259

Experimental Condition	electron donor	electron acceptor	carbon source	t_d (h)
Thiosulfate-Lactate (T-L)	Lactate (20 mM)	Thiosulfate (14 mM)	Lactate (20 mM)	6.2 ± 0.6
Sulfate-Lactate (S-L)	Lactate (20 mM)	Sulfate (14 mM)	Lactate (20 mM)	10.3 ± 1.7
Thiosulfate-H_2/CO_2 (T-H_2/CO_2)	H_2 (g)* (80% v/v)	Thiosulfate (14 mM)	CO_2 (g)* (20% v/v)	19.3 ± 3.9

260
261 * The headspace of autotrophic cultures was filled with 21 psig H_2/CO_2 (80:20, v/v) and re-pressurized with H_2/CO_2
262 once per day after each OD measurement.
263

264 ### 3.2.   Carbon metabolism and growth phase affect lipid δ^2H and regression parameters
265

266 Different modes of carbon metabolism in *A. fulgidus* not only resulted in different growth rates
267 (Table 1) but also affected lipid δ^2H and regression parameters. The lipid isotopic composition
268 reflects the sum of distinct isotope fractionations between lipid and each external H source,
269 resulting in the isotopic mass balance (Zhang et al., 2009):
270

$$R_{BP} = f_W \cdot 2\alpha_{L/W} \cdot R_W + f_S \cdot 2\alpha_{L/S} \cdot R_S \qquad \text{Eq. (4)}$$

where R_{BP}, R_W, and R_S represent the $^2H/^1H$ ratios of BPs from GDGT-0 (BP-0), media water, and

substrate (lactate for the heterotrophic conditions; H_2 for the autotrophic condition), respectively.

f_W and f_S are the fractions of lipid-bound H originating from water and substrate, respectively. $^2\alpha$

represents the net isotope fractionation factor between lipid and water ($^2\alpha_{L/W}$) and between lipid

and substrate ($^2\alpha_{L/S}$), respectively. In parallel cultures where only one parameter (R_W) is varied

experimentally, measurements of R_L, R_W, and R_S provide the basis for regression of R_L on R_W.

The isotope analysis results and regression parameters are summarized in Table 2.

The δ^2H_L values were strongly correlated with δ^2H_W values in all experiments (Table 2; $R^2 \geq 0.993$

and $p \leq 0.05$ in every regression). The regression slopes varied as a function of carbon metabolism

and growth phase. Overall, the two heterotrophic conditions (T-L and S-L) and exponential growth

phase resulted in lower slopes compared to the autotrophic condition (T-H_2/CO_2) and stationary

growth phase. Slopes for the two heterotrophic conditions were comparable (0.481 for T-L, 0.477

for S-L during exponential phase; 0.548 for T-L, 0.519 for S-L during stationary phase; Table 2).

The autotrophic condition resulted in higher slopes (0.552, exponential phase; 0.628, stationary

phase). Notably, the higher slope observed during the stationary phase of the autotrophic culture

was the closest to the slope estimated assuming $f_W = 1$ (0.707) (Table 2; Fig. 1D).

Table 2. Data for tetraether-derived biphytanes produced by *Archaeoglobus fulgidus* grown on different
carbon substrates and electron donor-acceptor pairs.
294

Substrate [a]	Growth Phase [b]	Avg. OD [c]	δ^2H_W [d] (‰)	s.d. (‰)	δ^2H_L [e] (‰)	s.d. (‰)	$^2\varepsilon_{L/W}$ (‰)	s.d. (‰)	n [f]	Slope [g] ($^2\alpha_{L/W} \cdot f_W$)	R^2	*p*
T-L		0.147	362.4	0.2	−70.7	6.6	−317.9	29.4	3	0.481	0.996	1×10^{-7}
	Exp	0.135	176.0	0.2	−164.6	7.0	−289.6	12.4	3			
		0.141	−64.0	0.2	−275.8	7.1	−226.2	5.9	2			
		0.346	361.3	0.2	−52.7	6.7	−304.1	38.5	2	0.548	0.999	3×10^{-6}
	Exp+Sta	0.344	175.5	0.2	−152.4	7.5	−279.0	13.7	2			
		0.351	−63.7	0.2	−285.3	8.7	−236.7	7.2	2			
S-L		0.096	361.8	0.2	−87.5	9.7	−329.9	36.4	1	0.477	0.993	7×10^{-5}
	Exp	0.100	174.7	0.2	−163.5	4.8	−287.9	8.5	3			
		0.082	−61.7	0.2	−289.7	5.9	−243.0	5.0	2			
		0.307	361.4	0.2	−66.9	5.7	−314.6	26.9	2	0.519	0.998	4×10^{-7}
	Exp+Sta	0.295	175.4	0.2	−168.1	4.9	−292.3	8.5	2			
		0.282	−61.5	0.2	−287.1	10.2	−240.3	8.6	2			
T-H$_2$/CO$_2$		0.052	348.3	0.2	−72.8	5.0	−312.3	21.5	2	0.552	0.998	8×10^{-5}
	Exp	0.046	164.8	0.2	−176.8	7.0	−294.0	11.5	1			
		0.054	−66.5	0.2	−302.1	4.7	−252.4	4.0	1			
		0.137	348.5	0.2	−71.7	8.7	−311.7	37.8	3	0.628	0.997	1×10^{-5}
	Exp+Sta	0.132	166.0	0.2	−173.8	10.0	−291.4	16.8	1			
		0.134	−64.5	0.2	−326.4	10.2	−279.9	8.8	2			
T-H$_2$/CO$_2$ (calculated)	-	-	-	-	-	-	-	-	-	*0.707 ($f_W = 1$)*	-	-

295
[a] Thiosulfate-lactate (T-L); sulfate-lactate (S-L); thiosulfate-H$_2$/CO$_2$ (T-H$_2$/CO$_2$)
[b] Biomass harvested during mid-exponential growth phase (Exp); biomass harvested during early stationary phase,
including biomass from both exponential and stationary phase (Exp+Sta)
[c] Optical density (OD)
[d] δ^2H of culture medium at the time of biomass sampling. Average of biological and technical replicates ± 1 s.d.
[e] Weighted mean δ^2H of biphytane from GDGT-0
[f] Number of biological replicates used to calculate the average values of δ^2H_{Water}, δ^2H_{Lipid}, and $^2\varepsilon_{L/W}$
[g] Slope for the regression of δ^2H_{Lipid} against δ^2H_{Water} (*see* Figure 1D-F)
304

Figure 1. Growth curves and regression of lipid δ²H versus water δ²H for *Archaeoglobus fulgidus*. Growth curves of *A. fulgidus* grown autotrophically on (A) thiosulfate and H_2/CO_2 or heterotrophically on (B) thiosulfate and lactate or (C) sulfate and lactate. Regression of biphytane hydrogen isotopic composition (δ²H$_L$) on water isotopic composition (δ²H$_W$) grown on (D) thiosulfate and H_2/CO_2, (E) thiosulfate and lactate, and (F) sulfate and lactate.

4.   *A. fulgidus* H-isotope flux-balance model

To further interpret these findings, we modified the isotope flux-balance model recently developed for an ammonia-oxidizing archaeon *N. maritimus* SCM1 (Leavitt et al., 2023) to reflect the biochemistry underlying the carbon metabolism modes in *Archaeoglobus fulgidus* VC-16. The overall conceptual framework consists of two parts for the stoichiometric accounting of: 1) methyl-H in Ac-CoA and 2) the 80 H atoms in the C_{40} alkyl chain of BPs inherited from Ac-CoA. The former requires metabolism-specific information considering the mode(s) of carbon fixation and

320 experimental conditions such as substrate availability, while the latter is informed by more
321 universal information about archaeal isoprenoid biosynthesis. Below, we describe the details of
322 the two parts of our model.
323
324 4.1. Sources of lipid-H via the modified Wood-Ljungdahl pathway
325
326 *4.1.1. Heterotrophy*
327
A. fulgidus can grow autotrophically or heterotrophically via the modified reductive Ac-CoA or
Wood-Ljungdahl pathway (Möller-Zinkhan et al., 1989; Möller-Zinkhan and Thauer, 1990;
Hocking et al., 2014). During chemoorganoheterotrophic growth on lactate, the type species *A.*
fulgidus VC16 couples the complete oxidation of organic substrates to CO_2 with the reduction of
sulfate or thiosulfate to sulfide. In this study, the organic substrate was lactate. *A. fulgidus* oxidizes
lactate via pyruvate and Ac-CoA using enzymes and cofactors involved in the C_1-pathway, similar
to those found in methanogens, in reverse direction (Fig. 2A; Möller-Zinkhan et al., 1989; Möller-
Zinkhan and Thauer, 1990). A D-lactate dehydrogenase in *A. fulgidus* catalyzes the first step of
lactate oxidation to pyruvate coupled with the reduction of menaquinone (MQ), resulting in
menaquinol (MQH_2) (Step 1, Fig. 2A; Kunow et al., 1993; Reed and Hartzell, 1999). The next step
of pyruvate oxidation to Ac-CoA is catalyzed by pyruvate:ferredoxin oxidoreductase (Step 2, Fig.
2A; Kunow et al., 1993). The 3 H atoms in the methyl group of lactate most likely remain intact
and are inherited to Ac-CoA. Some of these Ac-CoA products are used for biomass synthesis (Step
3.2, Fig. 2A), including isoprenoid lipid biosynthesis, while some of them are degraded further for
the complete oxidation of lactate. For the latter, carbon monoxide dehydrogenase catalyzes both
the reversible carbon-carbon cleavage of Ac-CoA and the oxidation of carbonyl group to CO_2
coupled with ferredoxin (Fd) reduction (Step 3.1, Fig. 2A; Möller-Zinkhan and Thauer, 1990).
345
The coenzymes involved in the downstream oxidative branch of Ac-CoA degradation could
transfer hydrides (e.g., $F_{420}H_2$ in Steps 4 and 5, Fig. 2A). These fluxes of H are important to track,
because the hydrides from Ac-CoA (ultimately from lactate) can be recycled for NADP reduction,
the product of which (NADPH) can donate hydrides to isoprenoid precursors during lipid
biosynthesis (i.e., contribution of lactate-derived H to lipids). We assign F_{420} and Fd as the electron
acceptors for Steps 4–5 and Step 8, respectively (Fig. 2A), given that the genes relating to all steps
of the reductive Ac-CoA pathway are expressed in *A. fulgidus* (Hocking et al., 2014) and
considering the known mechanisms of how the enzymes of interest operate in methanogens. In
CO_2-reducing methanogens, both methylenetetrahydromethanopterin reductase (Mer) and
methylenetetrahydromethanopterin dehyrogenase (Mtd) oxidize $F_{420}H_2$ (Steps 4 and 5,
respectively, in reverse; Fig. 2A), and formylmethanofuran dehydrogenase (Fwd) requires Fd_{red}
(Step 8, in reverse; Fig. 2A) (Thauer et al., 2010).
358

4.1.2. Autotrophy

During chemolithoautotrophic growth, the type species *A. fulgidus* VC16 couples the oxidation of
H_2 or formate with the reduction of thiosulfate or sulfite (Stetter, 1988; Steinsbu et al., 2010). In
this study, H_2 and thiosulfate were used as electron donor and acceptor for the autotrophic
condition (T-H_2/CO_2). *A. fulgidus* reduces CO_2 to synthesize Ac-CoA via the C_1-pathway (Fig.
2B), in principle a reversal of the Ac-CoA degradation during heterotrophy (Fig. 2A). The same
enzymes involved in H additions used by CO_2-reducing methanogens catalyze the successive
addition of H during autotrophic carbon fixation in *A. fulgidus*: Fwd (Step 1), Mtd (Step 4), and
Mer (Step 5) (Fig. 2B). During chemolithoautotrophy, the H atoms in methyl precursors in theory
derive from intracellular water. However, one of the isoenzymes for
methylenetetrahydromethanopterin (CH_2-H_4MPT) dehydrogenase is H_2-dependent (Hmd) and
uses H_2 directly, as opposed to $F_{420}H_2$ used by the other isoenzyme Mtd, to reduce methyl-H_4MPT
(Hendrickson and Leigh, 2008). Given the similarities between methanogens and *A. fulgidus*, we
consider the possible flux of H from H_2 to Ac-CoA (Step 4–6, Fig. 2B) in one of the end-member
scenarios (see 4.2; Table 3).

Figure 2. Carbon metabolism by *Archaeoglobus fulgidus* via the modified reductive acetyl-CoA or Wood-Ljungdahl pathway. (A) Chemoorganoheterotrophic growth on lactate: lactate is oxidized via pyruvate and acetyl-CoA, which either serves as the building blocks for biosynthesis or is further degraded for the complete oxidation of lactate. (B) Chemolithoautotrophic growth on H_2/CO_2: CO_2 is reduced to produce acetyl-CoA via the C_1-pathway, with enzymes and cofactors similar to those involved in methanogenesis. Abbreviations: MQH_2, menaquinol or reduced menaquinone; Fd_{red}, reduced ferredoxin; Ac-CoA, acetyl-CoA; CH_3-H_4MPT, methyl-tetrahydromethanopterin (H_4MPT); CH_2=H_4MPT, methylene-H_4MPT; CH^+≡H_4MPT, methenyl-H_4MPT; CHO-H_4MPT, formyl-H_4MPT; CHO-MFR, formylmethanofuran; $F_{420}H_2$, reduced coenzyme F_{420}.

4.2.   Model schematics and stoichiometric accounting

Fractions of Ac-CoA produced via the modified Wood-Ljungdahl pathway described above are used for the synthesis of archaeal isoprenoids. Leavitt et al. (2023) provides a detailed overview of the sources of H and mechanisms of the biochemical steps involved in archaeal tetraether biosynthesis (Supporting Information, Fig. S1). In short, the mevalonate pathway synthesizes isopentenyl pyrophosphate (IPP), which are C_5 isoprene units, from three units of Ac-CoA. Then, three units of IPP and its isomer condense to yield the C_{20} precursor, geranylgeranyl diphosphate (GGPP). The geranylgeranyl chains are transferred around to yield di-O-geranylgenranylglyceryl phosphate (DGGGP) with the glycerol-1-phosphate (G1P) backbone. The tetraether synthase (Tes) enzyme catalyzes the condensation of two DGGGP units to yield the membrane-spanning tetraether. Finally, the geranylgeranyl reductase (GGR) enzyme catalyzes the saturation of tetraether to yield GDGT. We combine the stoichiometric accounting of the methyl-H in Ac-CoA for *A. fulgidus* (see 4.1) and of the 80 H atoms in the C_{40} alkyl chain of BPs inherited from Ac-CoA during tetraether biosynthesis described above (Leavitt et al., 2023) to build an overall schematics of the specific model scenarios explored in this study (Fig. 3).

Figure 3. Schematic diagram of the isotope flux-balance model for *Archaeoglobus fulgidus*. (A) Heterotrophic scenario for lactate oxidation; (B) autotrophic scenario for H_2 oxidation. Color codes for the different H fluxes (*f*) are shown in the legend. The subscript W stands for water as the ultimate H source; the subscript S stands for substrate as the ultimate H source, where the substrates are lactate and H_2 for heterotrophy and autotrophy, respectively. See 4.2.1 and Eq. 5–8 for detailed descriptions of individual fluxes. Abbreviations: LS, lipid synthesis; AS, acetyl-CoA synthesis; Cofa, cofactor; $F_{420}H_2$, reduced coenzyme F_{420}; Fd_{red}, reduced ferredoxin; Ac-CoA, acetyl-CoA; GGR, geranylgeranyl reductase.

4.2.1.   Empirically informed isotopic mass balance

With the fractions defined in the hydrogen budget (see 4.2.1), the isotopic mass balance in Eq. (4) can be further refined. Replacing the f_W and f_S terms in Eq. (4) with the sums of all water-derived and lactate-derived fractions in Eq. (5), isotopic mass balance for the heterotrophic case becomes:

$$R_{BP,het} = \left(f_{W,direct} \cdot \alpha_{W,direct} + f_{W,cofa-LS} \cdot \alpha_{W,cofa-LS}\right) \cdot R_W + \left(f_{S,cofa-LS} \cdot \alpha_{S,cofa-LS} + f_{S,AS} \cdot \alpha_{S,AS}\right) \cdot R_S$$

$$\text{Eq. (9)}$$

Similarly, isotopic mass balance for the autotrophic case becomes:

$$R_{BP,auto} = \left(f_{W,direct} \cdot \alpha_{W,direct} + f_{W,cofa-AS} \cdot \alpha_{W,cofa-AS} + f_{W,cofa-LS} \cdot \alpha_{W,cofa-LS}\right) \cdot R_W + f_{S,AS} \cdot \alpha_{S,AS} \cdot R_S$$

$$\text{Eq. (10)}$$

Now we can use the regression parameters estimated from the δ^2H_{H2O} label experiments (Fig. 1; Table 2) to calculate analytical solutions for reasonable ranges of individual terms in Eq. (9–10). In Eq. (9–10), the coefficient for R_W in the first term represents the slope, *m*, and the entire second term represents the intercept, *c*. For heterotrophy:

$$m_{het} = f_{W,net} \cdot \alpha_{W,net} = f_{W,direct} \cdot \alpha_{W,direct} + f_{W,cofa-LS} \cdot \alpha_{W,cofa-LS}$$

$$\varepsilon_{W,net} = \frac{m_{het}}{f_{W,net}} - 1 = \frac{f_{W,direct} \cdot \alpha_{W,direct} + f_{W,cofa-LS} \cdot \alpha_{W,cofa-LS}}{f_{W,net}} - 1 \qquad \text{Eq. (11)}$$

$$c_{het} = f_{S,net} \cdot \alpha_{S,net} \cdot R_S = f_{S,cofa-LS} \cdot \alpha_{S,cofa-LS} + f_{S,AS} \cdot \alpha_{S,AS}$$

$$\varepsilon_{S,net} = \frac{c_{het}}{f_{S,net} \cdot R_S} - 1 = \frac{f_{S,cofa-LS} \cdot \alpha_{S,cofa-LS} + f_{S,AS} \cdot \alpha_{S,AS}}{f_{S,net}} - 1 \qquad \text{Eq. (12)}$$

where $\varepsilon_{W,net}$ and $\varepsilon_{S,net}$ refer to the net isotope fractionation between water and lipid (combining

isotope effects associated with the water-derived fluxes, $f_{W,direct}$ and $f_{W,cofa\text{-}LS}$) and between

substrate and lipid (combining isotope effects associated with the substrate-derived fluxes, $f_{S,cofa\text{-}}$

$_{LS}$ and $f_{S,AS}$), respectively.

4.3.   Model results

Table 3, Fig. 4 and 5 shows the summary of model results. We first estimated $f_{W,net}$, which we

define as the net total contribution of H from water to lipids ($f_{W,net} = f_{W,direct} + f_{W,cofa}$), for a range

of x_{ex} from 0 % to 100 %. Overall, the autotrophic case had a higher range of $f_{W,net}$ compared to

the two heterotrophic cases (Table 3; Fig. 4A–C). Even for the minimal water exchange and

maximal substrate contribution scenario ($x_{ex} = 0$ %, $s = 100$ %), when the contribution from water

is expected to be the lowest, $f_{W,net}$ was 80 % (Table 3; dashed line, Fig. 4C). For the two

heterotrophic cases, the $f_{W,net}$ values were 31 and 70 % for the lowest ($x_{ex} = 0$ %, $s = 100$ %) and

highest ($x_{ex} = 100$ %, $s = 0$ %) water contribution scenarios, respectively (Table 3; Fig. 4A–B).

For the net isotope fractionation between lipid and water ($\varepsilon_{W,net}$), the two heterotrophic cases

showed similar ranges and patterns where the $\varepsilon_{W,net}$ values decrease with increasing x_{ex} (Fig. 4D–

E). The $\varepsilon_{W,net}$ range was relatively more positive during stationary growth phase (–218 to 776 ‰

for the T-L condition; –259 to 682 ‰ for the S-L condition) compared to exponential growth phase

(–312 to 561 ‰ for the T-L condition; –319 to 546 ‰ for the S-L condition) (Table 3; Fig. 4D–

E). For the autotrophic case, the growth phase-dependent trend was also observed (more positive

$\varepsilon_{W,net}$ during stationary phase; Table 3; Fig. 4F). Unlike in the heterotrophic cases, all $\varepsilon_{W,net}$ values

for the autotrophic case were negative and fell within a relatively narrow range for the entire range

of x_{ex} (–448 to –312 ‰ for exponential phase; –372 to –228 ‰ for stationary phase; Table 3). For

the autotrophic case, a theoretical end-member $\varepsilon_{W,net}$ with no substrate contribution can be

estimated by calculating the regression slope through the origin (i.e., no intercept, which represents

the substrate contribution term, $f_S \cdot {}^2\alpha_{L/W} \cdot R_S$; Eq. 4): –293 ‰ for the autotrophic case (magenta line;

Fig. 4F). This value falls within the range of $\varepsilon_{W,net}$ for stationary, rather than exponential, growth

phase (Fig. 4F). For the net isotope fractionation between lipid and substrate ($\varepsilon_{S,net}$), the two

heterotrophic cases showed similar ranges and patterns where $\varepsilon_{S,net}$ values increase with increasing

x_{ex} (Fig. 4G–I). The growth phase-dependent trend for $\varepsilon_{S,net}$ contrasted that of $\varepsilon_{W,net}$, where the $\varepsilon_{S,net}$

range was relatively more negative during stationary growth phase (–688 to –282 ‰ for the T-L

condition; –654 to –201 ‰ for the S-L condition) compared to exponential growth phase (–582 to

–36 ‰ for the T-L condition; –592 to –59 ‰ for the S-L condition) (Table 3; Fig. 4G–H). A similar

trend was observed for the autotrophic case (more negative $\varepsilon_{S,net}$ during stationary phase; Table 3;
Fig. 4I).
476
The net isotope fractionation factors ($\varepsilon_{W,net}$ and $\varepsilon_{S,net}$) can be further broken down into individual
fluxes associated with water-derived and substrate-derived H atoms, respectively. The net isotope
fractionation factor between lipid and water ($\varepsilon_{W,net}$) can be divided into: 1) $\varepsilon_{W,direct}$ associated with
water-derived H atoms that are transferred through the $f_{W,direct}$ flux (Fig. 3) via H exchange between
isoprenoid precursors and intracellular water or addition via GGR, and 2) $\varepsilon_{W,cofa}$ associated with
the $f_{W,cofa}$ flux ($f_{W,cofa-LS}$ for heterotrophy; $f_{W,cofa-LS}$ and $f_{W,cofa-AS}$ for autotrophy; Fig. 3) via
cofactors (Fd_{red} and $F_{420}H_2$ for autotrophy; Fd_{red} for heterotrophy) (Fig. 5A–C). The net isotope
fractionation factor between lipid and substrate ($\varepsilon_{S,net}$) can be divided into: 1) $\varepsilon_{S,AS}$ associated with
substrate-derived H atoms that are transferred through the $f_{S,AS}$ flux (Fig. 3) via Ac-CoA synthesis,
and 2) $\varepsilon_{S,cofa}$ associated with the $f_{S,cofa}$ flux ($f_{S,cofa-NADPH}$ and $f_{S,cofa-GGR}$ for heterotrophy; Fig. 3) via
cofactors ($F_{420}H_2$) (Fig. 5D–E). Interpretation of these results are discussed further in 5.2.1.
488

489
490 **Table 3.** Isotope flux-balance model results for *Archaeoglobus fulgidus*.

Growth phase	Scenario		$f_{W,net}$ (%)[c]			$\varepsilon_{W,net}$ (‰)[d]			$\varepsilon_{S,net}$ (‰)[e]		
	x_{ex} (%)[a]	s (%)[b]	T-L	S-L	T-H	T-L	S-L	T-H	T-L	S-L	T-H
Exp.	0	0	41	41	100	179	167	−448	−511	−523	n/a
	0	100	31	31	80	561	546	−312	−582	−592	−564
	100	0	70	70	100	−312	−319	−448	−36	−59	n/a
	100	100	60	60	90	−198	−206	−387	−277	−294	−139
Sta.	0	0	41	41	100	341	270	−372	−636	−595	n/a
	0	100	31	31	80	776	682	−218	−688	−654	−785
	100	0	70	70	100	−218	−259	−372	−282	−201	n/a
	100	100	60	60	90	−87	−136	−302	−461	−401	−577

491
[a] the degree of water exchange during lipid synthesis, which depends on the extent of re-equilibration during
isomerization steps (steps 2 and 6; Supporting Information, Fig. S1).
[b] the degree of H contribution to lipids from substrate (lactate for heterotrophy; H_2 for autotrophy)
[c] the net total contribution of H from water to lipids ($f_{W,net} = f_{W,direct} + f_{W,cofa}$)
[d] the net isotope fractionation between water and lipid (combining isotope effects associated with the water-derived
fluxes, $f_{W,direct}$ and $f_{W,cofa-LS}$)
[e] the net isotope fractionation between substrate and lipid (combining isotope effects associated with the substrate-
derived fluxes, $f_{S,cofa-LS}$ and $f_{S,AS}$)
500
501

Figure 4. Isotope flux-balance model results for *Archaeoglobus fulgidus*. (A–C) Net fraction of total hydrogen derived from water ($f_{W,net}$); (D–F) Net isotope fractionation between lipid and water ($\varepsilon_{W,net}$); (G–I) Net isotope fractionation between lipid and substrate ($\varepsilon_{S,net}$). Each column represents the metabolic mode and growth substrates: heterotrophy on thiosulfate and lactate (A, D and G); heterotrophy on sulfate and lactate (B, E and H); and autotrophy on thiosulfate and H_2/CO_2 (C, F and I). Lighter shaded areas refer to exponential growth phase; darker shades refer to stationary growth phase. Solid lines refer to the s = 0 % (minimal substrate contribution) scenarios; dashed lines refer to the s = 100 % (maximal substrate contribution) scenarios. See 4.2.1 and Supporting Information Table S1 for details of the conditions for each scenario.

Figure 5. Isotope flux-balance model results for *Archaeoglobus fulgidus*. (A–C) Breakdown of the net isotope fractionation between lipid and water ($\varepsilon_{W,net}$; Fig. 4D–F) into two different isotope fractionation factors associated with water-derived H: $\varepsilon_{W,direct}$ for water-derived H atoms that are transferred through exchange during isomerization or addition via GGR, corresponding to $f_{W,direct}$ in Fig. 3, and $\varepsilon_{W,cofa}$ for water-derived H atoms that are transferred through cofactors (Fd_{red} and $F_{420}H_2$ for autotrophy, panel A; Fd_{red} for heterotrophy, panels B and C), corresponding to $f_{W,cofa}$ in Fig. 3. (D–E) Breakdown of the net isotope fractionation between lipid and substrate ($\varepsilon_{S,net}$; Fig. 4G–I) into two different isotope fractionation factors associated with substrate-derived H: $\varepsilon_{S,AS}$ for substrate-derived H atoms that are inherited directly from substrate via Ac-CoA, corresponding to $f_{S,AS}$ in Fig. 3, and $\varepsilon_{S,cofa}$ for substrate-derived H atoms that are transferred through cofactors (only applicable for heterotrophy via $F_{420}H_2$, panels D and E), corresponding to $f_{S,cofa}$ in Fig. 3. The lighter shaded areas refer to the s = 0 % (minimal substrate contribution) scenarios; darker shaded areas refer to the s = 100 % (maximal substrate contribution) scenarios. Solid lines and brown colors refer to heterotrophic conditions; dashed lines and purple colors refer to autotrophic conditions. See 4.2.1 and Supporting Information Table S1 for details of the conditions for each scenario.

5. Discussion

5.1. Effects of carbon metabolism and growth phase on lipid H isotope composition

Isotope flux-balance model results suggest that a large fraction of lipid-bound H in *A. fulgidus* is sourced from water. The fraction was larger for the autotrophic case ($f_{W,net}$ = 80–100 %); the $f_{W,net}$

range for the heterotrophic cases was relatively lower ($f_{W,net}$ = 31–70 %; Table 3; Fig. 4A–C). However, considering that $\varepsilon_{W,net}$ is most likely a negative value (i.e., a normal kinetic isotope fractionation), we can constrain the $f_{W,net}$ range that corresponds to $\varepsilon_{W,net} < 0$ in Fig. 4D–E. This yields the $f_{W,net}$ ranges of: 38–52 % for T-L and 42–52 % for S-L during exponential growth phase, versus 50–56 % for T-L and 52–54 % S-L during stationary growth phase. The lowest $f_{W,net}$ were therefore recorded during heterotrophy at exponential phase growth, independent of the terminal electron acceptor. Moreover, ≥80 % of lipid-bound H during autotrophy and ≥50 % of lipid-bound H in stationary phase cultures of *A. fulgidus* during heterotrophy originate from water. The $f_{W,net}$ range for lipids produced during stationary phase alone is likely even larger, because lipids collected during stationary phase are a mixture of those produced during exponential phase and those newly produced during stationary phase. The higher $f_{W,net}$ range observed for autotrophy is reasonable, given that lactate is an additional source of H that contributes directly to Ac-CoA synthesis during heterotrophy. A similar observation was reported for bacterial lipids where, for instance, the mole fraction of lipid H derived from water ($\equiv X_W$ in Zhang et al., 2009; herein equivalent to $f_{W,net}$) was larger for *Escherichia coli* grown on glucose compared to the same organism grown on complex medium (Zhang et al., 2009). Though the lipid synthesis pathways are different for Bacteria versus Archaea, it appears that the fractional contribution of water to lipid H generally decreases when substrates that are more direct precursors for lipid biosynthesis are used.

The effects of carbon metabolism and growth phase on lipid isotope composition have not been investigated in detail for archaeal lipids. Our results indicate that both the mode of metabolism and growth phase affect the lipid isotope composition (Table 3; Fig. 4). A recent study reported $\varepsilon_{L/W}$ of BPs produced by *N. maritimus* in continuous cultures at a range of doubling time and observed little change in $\varepsilon_{W/L}$ (0.18 ± 0.09 ‰ increase in $\varepsilon_{L/W}$ per hour increase in doubling time; Leavitt et al., 2023). The relative variation in doubling time we observe for *A. fulgidus* grown on different substrates here in batch cultures (Table 1) is comparable (*ca.* 3-fold variation) to the variation observed for *N. maritimus* in Leavitt et al. (2023). A key difference here is that we do not strictly control growth and metabolic rates, as was the case for chemostat cultivation of *N. maritimus* (Leavitt et al., 2023), and for *Sulfolobus acidocaldarius* (Harris et al., 2022). Nonetheless, the range of $\varepsilon_{L/W}$ we observe across different metabolic modes by *A. fulgidus* (–231 ± 6 ‰ for T-L, –241 ± 4 ‰ for S-L, and –271 ± 18 ‰ for T-H_2/CO_2; Table 2) is significant compared to the entire range of $\varepsilon_{L/W}$ observed across a comparable variation in growth rate by *N. maritimus* (–272 ± 20 to –279 ± 8 ‰ for BP-0; Leavitt et al., 2023). *N. maritimus* is an obligate chemoautotroph, such that all BP-bound H is strictly biosynthetic, whereas we show here that patterns of $\varepsilon_{W/L}$ can change between heterotrophic and autotrophic growth conditions (Fig. 4 and 6). Our results support this hypothesis and show variation in $\varepsilon_{L/W}$ up to ~40 ‰ within the same metabolically versatile organism, in response to different electron donor-acceptor pairs and modes of carbon metabolism.

572 Even though the range of $\varepsilon_{L/w}$ in *A. fulgidus* is relatively larger compared to *N. maritimus*, the
573 overall variation of $\varepsilon_{L/w}$ across all archaeal lipid studies to-date is consistently small compared to
574 the variation observed across bacterial lipids (Fig. 6).
575
576 5.2.   Factors that affect $\varepsilon_{L/w}$ across the three domains of life
577
Here we present a synthesis of the broader $\varepsilon_{L/w}$ patterns observed in lipids from all three domains
of life. Archaeal lipids are characterized by a relatively narrow range of large and negative values
of $\varepsilon_{L/w}$ across a wide range of organisms, metabolisms, and growth conditions (Fig. 6; Kaneko et
al., 2011; Wu et al., 2020; Leavitt et al., 2023; this study). This is in contrast with the domain
Bacteria, where fatty acid $\varepsilon_{L/w}$ ranges a large span of >600 ‰ with both highly positive and
negative values (Fig. 6; Campbell et al., 2009; Zhang et al., 2009; Dawson et al., 2015; Osburn et
al., 2016; Wijker et al., 2019). Archaeal lipid $\varepsilon_{L/w}$ largely overlaps with that of eukaryotic lipid
$\varepsilon_{L/w}$, although eukaryotic lipid $\varepsilon_{L/w}$ spans a significant range depending on the types of organism
and classes of compound (Fig. 6; Sessions et al., 1999; Chikaraishi et al., 2004, 2009; Zhang and
Sachs, 2007; Sachs et al., 2016). Notably, there are subgroups of both domains Bacteria and
Eukarya that share similarities with Archaea in their $\varepsilon_{L/w}$. The shared similarities are interesting,
given the difference in lipid biosynthesis mechanism among the three domains. Below, we take a
closer look at the specific subsets of $\varepsilon_{L/w}$ values from the domains Euakrya and Bacteria that
resemble those from Archaea and discuss factors that may contribute to the large, negative values
of $\varepsilon_{L/w}$ observed across the three domains.
593

Figure 6. Summary of hydrogen isotope fractionation ($\varepsilon_{L/W}$) between lipids and water observed in culture studies. $\varepsilon_{L/W}$ values observed in all three domains of life: Archaea (circle), Bacteria (square), and Eukarya (diamond). Color codes represent different types of metabolism: aerobic heterotrophy (red), anaerobic heterotrophy (purple), aerobic photoautotrophy (green), aerobic chemoautotrophy (orange), and anaerobic chemoautotrophy (blue). Electron donors used for biosynthesis are shown on the horizontal axis: acetate (ac); succinate (su); pyruvate (py); lactate (la/ala); glucose (gl/agl); methane (me); H_2+CO_2 (h2/hc); formate (fo); oxalate (ox); H_2+Fe^{3+} (hf); H_2+S^0 (hs); ammonium (aa); yeast extract (ye); H_2, acetate or methanol for methanogenesis (mg); water for photoautotrophy (pa). Tested organisms for bacterial cultures include *C. oxalaticus, C. necator, R. palustris* (Zhang et al., 2009); *E. coli* (Zhang et al., 2009; Osburn et al., 2016; Wijker et al., 2019); *D. autotrophicum* (Campbell et al., 2009; Osburn et al., 2016); *M. capsulatus* (Sessions et al., 2002); *D. multivorans* (Dawson et al., 2015); *D. alaskensis* (Leavitt et al., 2016); *P. denitrificans, S. oneidensis, D. hydrogenophilus, D. alaskensis, D. propionicus* (Osburn et al., 2016); *B. subtilis, P. fluorescens, R. radiobacter, and E. meliloti* (Wijker et al., 2019). Tested organisms for eukaryotic cultures include *A. fundyense, I. galbana, Ascophyllum sp., F. vesiculosis, Z. marina, S. alterniflora* (Sessions et al., 1999); *C. japonica* (Chikaraishi, Naraoka and Poulton, 2004); *B. braunii, E. unicocca, V. aureus* (Zhang and Sachs, 2007); *C. sativuc* (Chikaraishi et al., 2009); *E. huxleyi* (Sachs et al., 2016). Tested organisms for archaeal cultures include *Acidianus* sp. DS80 ('hf' and 'hs', this study); *N. maritimus* ('aa', Leavitt, Kopf et al., 2023); *Sulfolobus* sp. ('ye', Kaneko et al., 2011); *A. fulgidus* ('ala' and 'hc', this study); *Metallosphaera sedula* ('hc',this study); and *M. barkeri* ('mg', Wu et al., 2020).

5.2.1.   Large $\varepsilon_{L/W}$ associated with isoprenoid saturation: Archaeal and Eukaryotic lipids

The consistently large, negative values of $\varepsilon_{W/L}$ for archaeal lipids are notable considering the
different types of organism and metabolic modes studied so far (Fig. 6). In Fig. 5, we present the
model solution spaces for individual H fluxes, within either net water-derived flux ($f_{W,net}$) or net
substrate-derived flux ($f_{S,net}$), to further investigate specific parts of archaeal lipid biosynthesis
pathway that may contribute to the observed large and negative values of $\varepsilon_{L/W}$. Prior work yielded
estimates of $\alpha_W \cong 0.9$ for isotope fractionation associated with direct water incorporation across
many bacteria (Zhang et al., 2009; Wijker et al., 2019), and $\varepsilon_{L/W}$ values modeled for *N. maritimus*
with this empirical α_W were consistent with experimental observations (Leavitt et al., 2023).
Assuming an equivalent value of $\varepsilon_{W,direct}$ (i.e., isotope fractionation associated with the $f_{W,direct}$ flux
is –100 ‰), we constrain plausible ranges of $\varepsilon_{W,cofa}$ (water-derived H fluxes that involve Fd_{red} and
$F_{420}H_2$) in *A. fulgidus* (Fig. 5A–C). The ranges of $\varepsilon_{W,cofa}$ at $\varepsilon_{W,direct} = -100$ ‰ are particularly well
constrained for the autotrophic case (–599 to –436 ‰ at $s = 0$ %; –519 to –276 ‰ at $s = 100$ %;
Fig. 5A).

It is plausible that both $\varepsilon_{W,cofa-AS}$ and $\varepsilon_{W,cofa-LS}$ have negative values that are comparable to the
combined value of $\varepsilon_{W,cofa}$, as the less likely alternative would be a negative $\varepsilon_{W,cofa}$ that consists of
very different $\varepsilon_{W,cofa-AS}$ and $\varepsilon_{W,cofa-LS}$. Our current model cannot distinguish between isotope
fractionation factors associated with different fluxes within $f_{W,cofa}$ (i.e., $f_{W,cofa-AS}$ through Ac-CoA
vs. $f_{W,cofa-LS}$ through NADPH), both of which can inherit water-derived H from cofactors $F_{420}H_2$
and/or Fd_{red} (Fig. 3B). However it is worth noting that, for both heterotrophic and autotrophic
cases, a fraction of the $f_{W,cofa-LS}$ flux represents hydride transfer during the final saturation (double
bond reduction) via GGR (Fig. 3; step 10, Fig., S1). Results of the isotope flux-balance model for
N. maritimus also indicated that the GGR reduction step is strongly fractionating ($\varepsilon_{GGR} = -690$ to
–820 ‰ if $\varepsilon_{NADPH} = -100$ to –300 ‰, depending on the degree of water exchange, or x_{ex} in this
study), regardless of how other model parameters are set (Leavitt et al., 2023). Given how *N.*
maritimus and *A. fulgidus* share little in common otherwise (metabolically, genetically, and
biochemically), large fractionation associated with the GGR reduction step provides a
parsimonious explanation for the similar $\varepsilon_{L/W}$ values observed in the two organisms, as it is the
final step of archaeal lipid biosynthesis shared across all tetraether-producing Archaea. This would
be consistent with the similarly large, negative values of $\varepsilon_{L/W}$ observed in other archaeal lipids
(Fig. 6); testing more Archaea with the combined approach of isotope labeling experiments and
biochemically informed isotope flux-balance model will help further confirm this idea.

A closer examination of the eukaryotic lipids that shares similar $\varepsilon_{L/W}$ values to archaeal lipids lends
further support to the notion that GGR reduction contributes to large H isotope fractionation
between lipids and water. Fig. 7A shows $\varepsilon_{L/W}$ for different subgroups of eukaryotic lipids,
categorized by three different biosynthetic pathways. In general, the two major families of lipids
are acetogenic and isoprenoid lipids which consist of acetate and isoprene monomeric units,
respectively (Pearson, 2014). Isoprenoid lipids are further divided into two groups, depending on
whether the isoprene (C_5) building blocks are synthesized via the mevalonic acid (MVA) pathway

(Lynen et al., 1958; Bloch, 1959) or the methylerythritol phosphate (MEP) pathway (Rohmer et
al., 1993). All Archaea discussed in this study synthesize isoprenoid lipids via the MVA pathway.
The MEP pathway is generally defined as bacterial and the MVA pathway as eukaryotic and
archaeal; both pathways often simultaneously occur in plants, where the MEP pathway is
expressed in chloroplasts (relics of the cyonobacterial plastid ancestor) and the MVA pathway is
expressed in the cytosol (Lichtenthaler et al., 1997; Lichtenthaler, 1999; Pearson, 2014). An
exception to this is the case of green algae, where both the cytosolic sterols and plastidic phytol
isoprenoids are synthesized via the MEP pathway (Schwender et al., 1996; Disch et al., 1998;
Pearson, 2014).

For eukaryotic lipids, it has been recognized for several decades that the three different
biosynthetic pathways (acetogenic, MVA, and MEP) are associated with distinct ranges of $\varepsilon_{L/W}$.
In general, isoprenoid lipids have more negative $\varepsilon_{L/W}$ compared to acetogenic lipids (Estep and
Hoering, 1980; Sessions et al., 1999). Among isoprenoid lipids, plastidic isoprenoids such as
phytols are more depleted in ^{2}H than the cytosolic counterparts (Sessions et al., 1999; Chikaraishi
et al., 2004, 2009). This general trend is captured in Fig. 7A, where acetogenic lipids (e.g., *n*-
alkanol and *n*-alkyl lipids including fatty acids) have more positive $\varepsilon_{L/W}$ compared to isoprenoid
lipids. The C_{15} and C_{30} isoprenoids synthesized via the MVA pathway (e.g., sterol, triterpenoids,
sesquiterpenes, squalene) are relatively more enriched in ^{2}H compared to the C_{20} isoprenoids (e.g.,
phytol, phytadiene, diterpenes) and botryococcenes (triterpenes) synthesized via the MEP pathway
(Fig. 7A).

While the relative difference in $\varepsilon_{L/W}$ between MVA and MEP pathways holds in general, the larger
and more negative values of $\varepsilon_{L/W}$ appear to correlate more with highly fractionating enzymatic
steps, rather than specific pathways. For example, Chikaraishi et al. (2004) observed in
Cryptomeria japonica (conifer) that $\varepsilon_{L/W}$ values for cytosolic lipids produced via the MVA
pathway, sesquiterpenes (–228 to –238 ‰) and squalene (–225 to –251 ‰), are comparable to
those for plastidic diterpenoids (–238 to –251 ‰) produced via the MEP pathway (Fig. 7A). Phytol
and botryococcenes, on the other hand, have particularly negative $\varepsilon_{L/W}$ values among eukaryotic
lipids produced via the MEP pathway. For example, Zhang and Sachs (2007) found that
botryococcenes with longer chain lengths had progressively lower δ^2H compared to the C_{30}
botryococcene (precursor to all other botryococcenes) in *Botryococcus braunii* race B. They
hypothesized that significantly ^{2}H-depleted hydrogen from methionine is incorporated into
botryococcene during the step-wise methylation (Zhang and Sachs, 2007). Similarly, phytol in *C.*
japonica had up to 65 ‰ more negative $\varepsilon_{L/W}$ compared to diterpenoids, another plastidic lipid
produced via the MEP pathway within the same organism (Chikaraishi et al., 2004). Later,
Chikaraishi et al. (2009) observed a gradual depletion in δ^2H from phytol precursors (starting from
geranylgeraniol) to phytol in cucumber, indicating that significantly ^{2}H-depleted hydrogen is
incorporated into phytol and its precursors during the step-wise hydrogenation catalyzed by GGR
($\varepsilon_{GGR} \cong$ –650 ‰; Chikaraishi et al., 2009).

The εL/w values of phytol (and its dehydration product phytadiene) investigated in a wide range of

eukaryotes so far are all significantly large and negative (e.g., –278 to –357 ‰ for phytol in

dinoflagellate, coccolithophore, marsh grass, cucumber, and conifer; Fig. 7A). Similarly, all values

of εL/w investigated in a range of different archaea are significantly large and negative (e.g., –341

to –148 ‰ for phytanes and biphytanes in *N. maritimus*, *Sulfolobus* sp., *M. sedula*, *Acidianus* sp.,

M. barkeri, and *A. fulgidus*; Fig. 7A). Taken together, empirical and model-based results suggest

that the GGR reduction is a highly fractionating step that contribute to the large fractionation

observed in both eukaryotic and archaeal lipid synthesis (Chikaraishi et al., 2009; Leavitt et al.,

2023; this study; Fig. 5 and 6).

Figure 7. Comparison of $\varepsilon_{L/W}$ values observed in eukaryotic *vs.* archaeal lipids and in bacterial *vs.* archaeal lipids. **(A)** $\varepsilon_{L/W}$ values observed in acetogenic and isoprenoid lipids from the domains Archaea and Eukarya. Different classes of compounds are shown on the horizontal axis. Compounds shown for acidogenic pathway: n-alkanol, *n*-alkane, *n*-alkyl (*n*-alkanoic acids and fatty acids), and alkadiene. For the MVA pathway: C_{15} and C_{30} isoprenoids (sterol, squalene, triterpenoids, and sesquiterpenes) and P+BP (phytanes and biphytanes). For the MEP pathway: C_{20} isoprenoids (diterpenes, sandaracopimarinol, ferruginol, diterpenol, and sugiol), phytol+ptd (phytol and phytadiene), and btc (botryococcene). **(B)** $\varepsilon_{L/W}$ values observed in fatty acids from anaerobic and/or autotrophic Bacteria and isoprenoid lipids from Archaea. Different organisms are shown on the horizontal axis. For Bacteria: *E. coli* (Ec), *Paracoccus denitrificans* (Pd), SRB (*Desulfobacterium autotrophicum*, *Desulfococcus multivorans*, *Desulfobacter hydrogenophilus*, *Desulfovibrio alaskensis*, *Desulfolobus propionicus*), and *Cupriavidus* sp. (Cu). For Archaea: *Acidianus* sp. DS80 (Ac), *Sulfolobus* sp. (Su), *Archaeoglobus fulgidus* (Af), *Metallosphaera sedula* (Ms), *Nitrosopumilus maritimus* (Nm), and *Methanosarcina barkeri* (Mb). A full list of references can be found in Supporting Information.

5.2.2.   Effect of energy limitation on $\varepsilon_{W/L}$: Archaeal and Bacterial lipids

Besides the shared biochemical mechanism underlying negative $\varepsilon_{L/W}$ in eukaryotic and archaeal lipids, another interesting pattern emerges from comparison of $\varepsilon_{L/W}$ across the three domains of life: a subset of bacterial lipids resembles archaeal lipids in its consistently negative $\varepsilon_{L/W}$ values (Fig. 6). For bacterial lipids, anaerobiosis and/or autotrophy appear to be associated with consistently large, negative values of $\varepsilon_{L/W}$. Examples include aerobic autotrophy (*Cupriavidus oxalaticus*, Zhang et al., 2009; *Paracoccus denitrificans*, Osburn et al., 2016), anaerobic autotrophy (*Desulfobacterium autotrophicum*, Campbell et al., 2009; *Desulfococcus multivorans*, Dawson et al., 2015; *P. denitrificans*, *Desulfobacterium autotrophicum*, *Desulfovibrio alaskensis*, Osburn et al., 2016), and anaerobic heterotrophy (*D. multivorans*, Dawson et al., 2015; *P. denitrificans*, *Escherichia coli*, *D. autotrophicum*, *Desulfobacter hydrogenophilus*, *D. alaskensis*, *Desulfolobus propionicus*, Osburn et al., 2016) (Fig. 6; Fig. 7B). Of these modes of metabolism, aerobic or anaerobic autotrophy has a more constrained range of $\varepsilon_{W/L}$ and more overlaps with $\varepsilon_{L/W}$ observed in archaeal lipids, compared to anaerobic heterotrophy (Fig. 6; Fig. 7B). We propose that NADPH flux imbalances associated with energy limitation that broadly characterize autotrophic and/or anaerobic modes of metabolism contribute to the apparent similarity in $\varepsilon_{L/W}$ observed between the domains of Bacteria and Archaea.

The concept of NADPH flux imbalance set forth by Wijker et al. (2019) provides a quantitative framework to relate metabolism-dependent patterns of $\varepsilon_{L/W}$ observed in aerobic heterotrophs to the fluxes of H through specific dehydrogenase and transhydrogenase enzymes. For the highly conserved fatty acid biosynthesis pathway, NADPH serves as the reducing agent for the reduction of acyl intermediates (Wakil et al., 1983). As a result, NADPH accounts for ~50% of fatty acid H (Sessions et al., 1999). Wijker et al. (2019) showed that $\varepsilon_{L/W}$ values of fatty acids produced by five wild-type species of aerobic heterotrophic bacteria are positively correlated with relative NADPH imbalance flux. Among the tested aerobic heterotrophs, two species (*E. coli* and *B. subtilis*) had

moderately negative values of NADPH flux imbalance (–32 and –11 %, respectively) (Wijker et
al., 2019). Here, negative values reflect underproduction of NADPH where catabolic production
of NADPH does not meet the anabolic requirement (Wijker et al., 2019). Furthermore, the results
from *E. coli* knockout mutants confirmed the direction of isotope fractionation for specific
transhydrogenases (e.g., ^{2}H-depletion by NADPH-producing PntAB and ^{2}H-enrichment by
NADPH-consuming UDhA) (Wijker et al., 2019).

For microbes adapted to chronic energy limitation, even more negative values of NADPH flux
imbalance would be expected where just enough NADPH is produced such that most to all of it is
used for biosynthesis (i.e., little to no flux from NADPH back to NADH). This prediction is
consistent with the general concept of energy spilling (also known as growth uncoupling or
overflow metabolism), which suggests that microbes under energy limitation tend to have more
efficient biomass production (Russell, 2007). For example, *E. coli* grown aerobically resulted in
more than 9-fold increase in ATP production per mole of glucose compared to the anaerobic
counterpart; however, the biomass yield was only 2-fold greater during aerobic respiration
(Stouthamer and Bettenhaussen, 1977; Gottschalk, 1986). Similar phenomenon of energy spilling
has also been observed in methanogenic archaea. For example, *Methanothermobacter*
thermautotrophicus and *Methanococcus maripaludis* produced less cells per mole of methane
when H_2 (electron donor) was present in excess compared to H_2-limiting condition (Morgan et al.,
1997; De Poorter et al., 2007; Costa et al., 2013).

In line with the observed and expected outcomes of energy limitation (negative NADPH flux
imbalance and tighter coupling between catabolism and anabolism), Leavitt et al. (2023) found
that $\varepsilon_{L/w}$ values for *N. maritimus* fall within the 95% confidence interval for the extension of
Wijker et al. (2019) regression when an extremely negative value (–100 %) of NADPH flux
imbalance was assigned. The apparent fit of *N. maritimus* $\varepsilon_{L/w}$ on the negative end of the
framework is reasonable, given that *N. maritimus* is an obligate autotroph adapted to oligotrophic
nitrification with energy-efficient CO_2 fixation pathway (Martens-Habbena et al., 2009; Könneke
et al., 2014). The fact that all archaeal lipids investigated for $\varepsilon_{L/w}$ so far fall within a relatively
narrow range of negative values (Fig. 6; Fig. 7B) may, at least partially, reflect the general concept
that Archaea are optimized for chronic energy limitation (Valentine, 2007). The results from this
study show that, while metabolic modes and growth rates have significant impacts on $\varepsilon_{L/w}$ within
a single organism (*A. fulgidus*), the observed variation in $\varepsilon_{L/w}$ still falls within the overall range
observed across archaeal lipids (e.g., heterotrophic and autotrophic *A. fulgidus* 'Af' data *vs*. other
archaeal lipids data; Fig. 7B).

The similarity between archaeal lipids and a subset of anaerobic and/or autotrophic bacterial lipids
is striking (Fig. 6; Fig. 7B) given the fundamental differences in fatty acid and isoprenoid lipid
biosynthesis pathways. This suggests that factors controlling microbial $\varepsilon_{L/w}$ may not be domain-
specific; instead, common challenges associated with anaerobic respiration and/or autotrophy (i.e.,

lower free energy yields and/or additional energetic costs for CO_2 fixation) may become primary
controlling factors of $\varepsilon_{L/w}$ for both Bacteria and Archaea. Notably, the range of $\varepsilon_{L/w}$ in fatty acids
produced by sulfate-reducing bacteria (SRB) is much narrower compared to that observed in
aerobic heterotrophs regardless of metabolic pathway or substrate (Fig. 7B).
795
One of the transhydrogenases that plays an important role in SRB is NfnAB, and it has been shown
that this enzyme is important for H isotope fractionation in *D. alaskensis* (Leavitt et al., 2016). The
details of the magnitude of $\varepsilon_{L/w}$ set by NfnAB under severe NADPH underproduction may be
complicated by other processes such as intracellular isotope distillation (i.e., more quantitative
transfer of NADPH hydride to lipids) as suggested by Leavitt et al. (2016). Regardless, given that
decoupling of NADPH production for anabolism from NADH for catabolism becomes more
important during energy limitation, transhydrogenases that are widely distributed among anaerobic
bacteria (namely NfnAB; Pereira et al., 2011; Leavitt et al., 2016; Buckel and Thauer, 2018) may
be key contributing factors for the negative $\varepsilon_{W/L}$ observed across these organisms (Fig. 6; Fig. 7B).
It is worth noting that *nfnAB* was not found in archaeal species investigated by Pereira et al. (2011),
including *A. fulgidus*.
807
A complex interplay among environmental, physiological, and enzymatic factors underlie net
isotope fractionation patterns observed in a biosynthetic product such as $\varepsilon_{L/w}$. For archaeal lipids
and a subset of bacterial lipids, however, two factors appear to play key roles in shaping the
consistently negative $\varepsilon_{L/w}$ patterns observed in the lipids from both domains: highly fractionating
enzymatic steps (e.g., GGR in archaeal isoprenoid lipid biosynthesis; transhydrogenases in
bacterial fatty acid biosynthesis) and energy limitation (i.e., NADPH flux imbalance). The
negative $\varepsilon_{L/w}$ commonly observed in archaea and a subset of bacteria is likely a convergent feature
resulting from a combination of both factors. A better mechanistic understanding of factors
conducive to similar patterns of $\varepsilon_{L/w}$ in bacterial and archaeal lipids could lead to biogeochemical
and/or ecological insights that are unique to microbial life. This would, in turn, significantly
expand the application of lipid biomarkers as paleoenvironmental or paleoecological proxy
currently possible with eukaryotic or bacterial lipids alone.

6.   Conclusion

In this study, we conducted water isotope label experiments with a metabolically flexible and well-
studied model archaeon *A. fulgidus* and quantified the hydrogen isotope fractionation between
lipids and water in response to different carbon substrates and electron donor-acceptor pairs. We
observed changes in regression parameters among experimental conditions, where higher slopes
($f_W \cdot {}^2\alpha_{L/w}$) were observed in autotrophic compared to heterotrophic conditions and during
stationary compared to exponential growth phase. The bio-isotopic model further constrains
individual parameters such as fractional contribution of water to lipids (f_W) and isotope
fractionation between lipids and water ($\varepsilon_{L/w}$) that together make up the slope. These results are
consistent with observations from regression parameters, where f_W is larger under autotrophic

condition and during stationary growth phase compared to the heterotrophic conditions and
exponential growth phase. The model results also allow us to constrain plausible ranges of isotope
fractionation associated with a specific route of hydrogen flux (e.g., flux for direct water
incorporation *vs*. flux from water through cofactors). The model results point toward the isotope
fractionation associated with the $f_{W,cofa}$ flux, which includes the hydride transfer during the final
saturation via GGR, as a highly fractionating step. This is consistent with recent bio-isotopic model
results for another archaeon *N. maritimus*. Finally, we synthesized a three-domain comparison of
$\varepsilon_{L/w}$ patterns which suggests that the highly fractionating double bond reduction via GGR may
also explain the similarly large and negative values of $\varepsilon_{L/w}$ observed between eukaryotic and
archaeal lipids. Furthermore, the shared negative $\varepsilon_{L/w}$ values between fatty acids in anaerobic
and/or autotrophic bacteria and isoprenoids in archaea may be associated with the general state of
energy limitation experienced by the different groups of organisms. Additional experimental work
with archaea that use carbon metabolism pathways that have not been examined with the bio-
isotopic model (i.e., 3HP/4HB and reductive Wood-Ljungdahl pathways) as well as more
systematic comparisons between bacterial and archaeal lipids will help determine the scope of
applicability for microbial lipid $\varepsilon_{L/w}$ in paleoenvironmental and paleoecological studies.

7.   Acknowledgements

This research was supported by funding from: collaborative research grant NSF EAR #1928303
(WDL, SHK); the American Chemical Society PRF #57209-DNI2 (WDL, YW); Simons
Foundation Award #623881 (WDL); Swiss National Science Foundation P2BSP2_168716 (YW);
NSF OCE-1843285 and 1702262 (AP). JHR was supported by the Dartmouth College Society of
Fellows. We thank Ashley Maloney (CU Boulder), Felix Elling (then at Harvard), and Beverly
Chiu (then at Dartmouth) for expert laboratory assistance.

8.   Appendix A. Overview of hydrogen budget

In our model for *A. fulgidus*, we consider two ultimate sources of H for both heterotrophic and
autotrophic model cases: water and substrate. Water refers to intracellular water for both cases,
and water-derived H can be conceptualized as either direct incorporation of intracellular water or
donation from reducing cofactors. Substrate refers to lactate and H_2, respectively, for the
heterotrophic (Fig. 3A) and autotrophic cases (Fig. 3B). The stoichiometric budget for H sources
in *A. fulgidus* can be simplified with the following fractions that make up either of the two passages
for water H incorporation and substrate H incorporation. For the 80 H atoms in BP produced via
heterotrophy (BP_{Het}): H derived directly from water via lipid biosynthesis ($f_{W,LS}$) and GGR
reduction ($f_{W,GGR}$) are combined to yield total direct water H ($f_{W,direct}$); H derived indirectly from
water via reducing cofactors NADPH ($f_{W,Cofa-NADPH}$) and GGR ($f_{W,Cofa-GGR}$) are combined to yield
total indirect water H during lipid synthesis ($f_{W,Cofa-LS}$); H derived indirectly from lactate (substrate)
via reducing cofactors NADPH ($f_{S,Cofa-NADPH}$) and GGR ($f_{S,Cofa-GGR}$) are combined to yield total

indirect water H during lipid synthesis ($f_{S,Cofa-LS}$); and H derived directly from lactate during Ac-
CoA synthesis ($f_{S,AS}$).

$BP_{Het} = f_{W,direct} + f_{W,Cofa-LS} + f_{S,Cofa-LS} + f_{S,AS}$ Eq. (5a)
$BP_{Het} = (f_{W,LS} + f_{W,GGR}) + (f_{W,Cofa-NADPH} + f_{W,Cofa-GGR}) + (f_{S,Cofa-NADPH} + f_{S,Cofa-GGR}) + f_{S,AS}$ Eq. (5b)

For the 80 H atoms in BP produced via autotrophy (BP_{Auto}): H derived directly from water via
lipid biosynthesis ($f_{W,LS}$) and GGR reduction ($f_{W,GGR}$) are combined to yield total direct water H
($f_{W,direct}$); H derived indirectly from water via reducing cofactors NADPH ($f_{W,Cofa-NADPH}$) and GGR
($f_{W,Cofa-GGR}$) are combined to yield total indirect water H during lipid synthesis ($f_{W,Cofa-LS}$); H
derived indirectly from water via reducing cofactors Fd_{red} ($f_{W,Cofa-Fdred}$) and $F_{420}H_2$ ($f_{W,Cofa-F420H2}$)
are combined to yield total indirect water H during Ac-CoA synthesis ($f_{W,Cofa-AS}$); and H derived
directly from H_2 (substrate) via Ac-CoA synthesis ($f_{S,AS}$).

$BP_{Auto} = f_{W,direct} + f_{W,Cofa-LS} + f_{W,Cofa-AS} + f_{S,AS}$ Eq. (6a)
$BP_{Auto} = (f_{W,LS} + f_{W,GGR}) + (f_{W,Cofa-NADPH} + f_{W,Cofa-GGR}) + (f_{W,Cofa-Fdred} + f_{W,Cofa-F420H2}) + f_{S,AS}$ Eq. (6b)

These fractions represent the relative fluxes of H (i.e., individual arrows either directly or indirectly
connecting the two ultimate H sources, water and substrate, to lipids) shown in Fig. 3. We explore
four scenarios for each case to explore the degree of water exchange, x_{ex}, ranging from 0% of
minimal exchange to 100% of maximal exchange and the degree of substrate contribution, ranging
from 0% of minimal contribution to 100% of maximal contribution. For the heterotrophic case,
the assigned fractions are as follows (Eq. 7):

$f_{W,LS} = \left[\dfrac{2}{3} \cdot (1 - x_{ex}) + 24 \cdot x_{ex}\right] \div 80$

$f_{W,GGR} = 8 \div 80$

$f_{W,direct} = \left[\dfrac{2}{3} \cdot (1 - x_{ex}) + 24 \cdot x_{ex} + 8\right] \div 80$

$f_{W,cofa-NADPH} = \left[16 \cdot (1 - \dfrac{1}{3} \cdot s)\right] \div 80$

$f_{W,cofa-GGR} = \left[8 \cdot (1 - \dfrac{1}{3} \cdot s)\right] \div 80$

$f_{W,cofa-LS} = \left[(16 + 8) \cdot (1 - \dfrac{1}{3} \cdot s)\right] \div 80$

$f_{S,cofa-NADPH} = \left[16 \cdot \dfrac{1}{3} \cdot s\right] \div 80$

$f_{S,cofa-GGR} = \left[8 \cdot \dfrac{1}{3} \cdot s\right] \div 80$

$f_{S,cofa-LS} = \left[(16 + 8) \cdot (\dfrac{1}{3} \cdot s)\right] \div 80$

$f_{S,AS} = \left[24 + \left(23 + \dfrac{1}{3}\right) \cdot (1 - x_{ex})\right] \div 80$ Eq. (7)

where x_{ex} is the degree of water exchange during lipid synthesis, which depends on the extent of
re-equilibration during isomerization steps (steps 2 and 6; Supporting Information, Fig. S1). The s
term is the degree of H contribution to lipids from substrate (lactate). Fractions that contain the x_{ex}
term ($f_{W,direct}$ and $f_{S,AS}$; Eq. 7) vary between the two end-member scenarios of $x_{ex} = 0$ % (minimal
exchange) and $x_{ex} = 100$ % (maximal exchange); fractions that contain the s term ($f_{W,cofa-LS}$ and
$f_{S,cofa-LS}$; Eq. 7) vary between the two end-member scenarios of $s = 0$ % (minimal contribution) and
$s = 100$ % (maximal contribution). At maximal substrate contribution ($s = 100$ %), ⅓ of $f_{W,cofa-LS}$
and $f_{S,cofa-LS}$ fractions derive from substrate H. During the complete oxidation of lactate to CO_2,
the average carbon oxidation state increases from 0 to +4 (i.e., 12 electrons transferred total for all
three carbons per molecule of lactate). The electron acceptors for the six oxidation steps are:
menaquinone, for lactate to pyruvate oxidation (step 1; Fig. 2A); ferredoxin, for pyruvate to Ac-
CoA oxidation and the final two oxidation steps to CO_2 (steps 3, 8 and 9; Fig. 2A); and F_{420}, for
methenyl-H₄MPT and methylene-H₄MPT oxidation (steps 4 and 5; Fig. 2A) (Möller-Zinkhan et
al., 1989; Schmitz et al., 1991; Klein et al., 1993; Kunow et al., 1993; Schwörer et al., 1993). None
of these redox reactions during lactate oxidation involve NAD or NADP as electron acceptors; in
A. fulgidus, reduced pyridine nucleotides (NADPH) required for biosynthesis is produced via the
reversible reduction of NADP with $F_{420}H_2$, similar to the case in methanogens (Kunow et al.,
1993). Therefore, in the maximal substrate contribution scenario ($s = 100$ %), up to ⅓ of H fluxes
through the cofactor NADPH and GGR can derive from the two (out of six, hence ⅓) lactate-
derived $F_{420}H_2$ that donate hydrides to NADP ($f_{S,cofa-LS}$; Fig. 3A).

For the autotrophic case, the assigned fractions are as follows (Eq. 8):

$f_{W,LS} = \left[\frac{2}{3} \cdot (1 - x_{ex}) + 24 \cdot x_{ex}\right] \div 80$

$f_{W,GGR} = 8 \div 80$

$f_{W,direct} = \left[\frac{2}{3} \cdot (1 - x_{ex}) + 24 \cdot x_{ex} + 8\right] \div 80$

$f_{W,Cofa-NADPH} = 16 \div 80$

$f_{W,Cofa-GGR} = 8 \div 80$

$f_{W,Cofa-LS} = (16 + 8) \div 80$

$f_{W,Cofa-Fdred} = \frac{1}{3} \cdot \left[24 + \left(23 + \frac{1}{3}\right) \cdot (1 - x_{ex})\right] \div 80$

$f_{W,Cofa-F420H2} = \left[\frac{1}{3} \cdot \left[24 + \left(23 + \frac{1}{3}\right) \cdot (1 - x_{ex})\right]\right] \cdot (2 - s) \div 80$

$f_{W,Cofa-AS} = \left[\left[\frac{1}{3} \cdot \left[24 + \left(23 + \frac{1}{3}\right) \cdot (1 - x_{ex})\right]\right] + \left[\frac{1}{3} \cdot \left[24 + \left(23 + \frac{1}{3}\right) \cdot (1 - x_{ex})\right]\right] \cdot (2 - s)\right] \div 80$

$f_{S,AS} = \frac{1}{3} \cdot s \cdot \left[24 + \left(23 + \frac{1}{3}\right) \cdot (1 - x_{ex})\right] \div 80$ Eq. (8)

where x_{ex} is the same as above (degree of water exchange during lipid synthesis, and s is the degree
of H contribution to lipids from substrate, H_2. Similar to the heterotrophic case above, the same
four scenarios with end-member values of x_{ex} and s are applied to the autotrophic case. Here, the

three reduction steps involved in autotrophic Ac-CoA synthesis (steps 1, 4 and 5; Fig. 2B) are the reverse of oxidative Ac-CoA degradation during heterotrophy (steps 4, 5 and 8; Fig. 2A). The first reduction step (step 1, Fig. 2B) involves Fd_{red}, and the next two reduction steps (steps 4 and 5, Fig. 2B) involve $F_{420}H_2$, where H in both of these electron donors derive from water ($f_{w,cofa-AS}$, Fig. 3B). For the autotrophic case, we assign the maximal substrate contribution ($s = 100\%$) when the H_2-dependent isoenzyme, Hmd, catalyzes the methenyl-H_4MPT reduction (step 4, Fig. 2B) under high partial pressure of H_2 (see 4.1.2). In this case, one of the three methyl-H in Ac-CoA derives directly from H_2 ($f_{S,AS}$, Fig. 3B).

9. References

Bender A. L. D. Genetic and Biological Studies on Solanum Plants: Insights into the Leaf Wax Alkane Paleoclimate Proxy.

Bloch K. (1959) Biogenesis and transformations of squalene. In *Biosynthesis of Terpenes and Sterols* Little, Brown and Company.

Brock T. D., Brock K. M., Belly R. T. and Weiss R. L. (1972) Sulfolobus: A new genus of sulfur-oxidizing bacteria living at low pH and high temperature. *Arch. F�r Mikrobiol.* **84**, 54–68.

Buckel W. and Thauer R. K. (2018) Flavin-Based Electron Bifurcation, Ferredoxin, Flavodoxin, and Anaerobic Respiration With Protons (Ech) or NAD+ (Rnf) as Electron Acceptors: A Historical Review. *Front. Microbiol.* **9**, 401.

Campbell B. J., Li C., Sessions A. L. and Valentine D. L. (2009) Hydrogen isotopic fractionation in lipid biosynthesis by H2-consuming Desulfobacterium autotrophicum. *Geochim. Cosmochim. Acta* **73**, 2744–2757.

Chikaraishi Y. and Naraoka H. (2003) Compound-specific δD–δ13C analyses of n-alkanes extracted from terrestrial and aquatic plants. *Phytochemistry* **63**, 361–371.

Chikaraishi Y., Naraoka H. and Poulson S. R. (2004) Carbon and hydrogen isotopic fractionation during lipid biosynthesis in a higher plant (Cryptomeria japonica). *Phytochemistry* **65**, 323–330.

Chikaraishi Y., Tanaka R., Tanaka A. and Ohkouchi N. (2009) Fractionation of hydrogen isotopes during phytol biosynthesis. *Org. Geochem.* **40**, 569–573.

Costa K. C., Yoon S. H., Pan M., Burn J. A., Baliga N. S. and Leigh J. A. (2013) Effects of H$_2$ and Formate on Growth Yield and Regulation of Methanogenesis in Methanococcus maripaludis. *J. Bacteriol.* **195**, 1456–1462.

Dawson K. S., Osburn M. R., Sessions A. L. and Orphan V. J. (2015) Metabolic associations with archaea drive shifts in hydrogen isotope fractionation in sulfate-reducing bacterial lipids in cocultures and methane seeps. *Geobiology* **13**, 462–477.

De Poorter L. M. I., Geerts W. J. and Keltjens J. T. (2007) Coupling of *Methanothermobacter*
thermautotrophicus Methane Formation and Growth in Fed-Batch and Continuous
Cultures under Different H_2 Gassing Regimens. *Appl. Environ. Microbiol.* **73**, 740–749.

Dirghangi S. S. and Pagani M. (2013) Hydrogen isotope fractionation during lipid biosynthesis by
Haloarcula marismortui. *Geochim. Cosmochim. Acta* **119**, 381–390.

Disch A., Schwender J., Müller C., Lichtenthaler H. K. and Rohmer M. (1998) Distribution of the
mevalonate and glyceraldehyde phosphate/pyruvate pathways for isoprenoid biosynthesis
in unicellular algae and the cyanobacterium Synechocystis PCC 6714. *Biochem. J.* **333**,
381–388.

Estep M. F. and Hoering T. C. (1980) Biogeochemistry of the stable hydrogen isotopes. *Geochim.*
Cosmochim. Acta **44**, 1197–1206.

Gottschalk (1986) Regulation of Bacterial Metabolism.

Harris C., Rhim J., Zhang Y., Cobban A., McFarlin J., Batther H., Kopf S. and Leavitt W. (2022)
Determining controls on hydrogen isotope fractionation in archaeal tetraether lipids in a
thermoacidophilic archaeal heterotroph. In *Goldschmidt2022 abstracts* Goldschmidt2022.
European Association of Geochemistry, Honolulu, HI, USA.

Hayes J. M. (2001) Fractionation of the Isotopes of Carbon and Hydrogen in Biosynthetic
Processes. *Stable Isot. Geochem.*, 225–277.

Hendrickson E. L. and Leigh J. A. (2008) Roles of Coenzyme F_{420}-Reducing Hydrogenases and
Hydrogen- and F_{420}-Dependent Methylenetetrahydromethanopterin Dehydrogenases in
Reduction of F_{420} and Production of Hydrogen during Methanogenesis. *J. Bacteriol.* **190**,
4818–4821.

Hocking W. P., Stokke R., Roalkvam I. and Steen I. H. (2014) Identification of key components
in the energy metabolism of the hyperthermophilic sulfate-reducing archaeon
Archaeoglobus fulgidus by transcriptome analyses. *Front. Microbiol.* **5**.

Kaneko M., Kitajima F. and Naraoka H. (2011) Stable hydrogen isotope measurement of archaeal
ether-bound hydrocarbons. *Org. Geochem.* **42**, 166–172.

Klein A. R., Breitung J., Linger D., Stetter K. O. and Thauer R. K. (1993) extremely thermophilic
sulfate reducing Archaeoglobusfulgidus: comparison of its properties with those of the
cyclohydrolase from the extremely thermophilic Methanopyrus kandleri.

Könneke M., Schubert D. M., Brown P. C., Hügler M., Standfest S., Schwander T., Schada Von
Borzyskowski L., Erb T. J., Stahl D. A. and Berg I. A. (2014) Ammonia-oxidizing archaea
use the most energy-efficient aerobic pathway for CO_2 fixation. *Proc. Natl. Acad. Sci.* **111**,
8239–8244.

Kopec B. G., Feng X., Posmentier E. S. and Sonder L. J. (2019) Seasonal Deuterium Excess
Variations of Precipitation at Summit, Greenland, and their Climatological Significance. *J.*
Geophys. Res. Atmospheres **124**, 72–91.

Kunow J., Schwörer B., Stetter K. O. and Thauer R. K. (1993) A F420-dependent NADP reductase
in the extremely thermophilie sulfate-reducing Archaeoglobusfulgidus.

Ladd S. N. and Sachs J. P. (2017) 2H/1H fractionation in lipids of the mangrove Bruguiera
gymnorhiza increases with salinity in marine lakes of Palau. *Geochim. Cosmochim. Acta*
204, 300–312.

Lai D., Springstead J. R. and Monbouquette H. G. (2008) Effect of growth temperature on ether
lipid biochemistry in Archaeoglobus fulgidus. *Extremophiles* **12**, 271–278.

Leavitt W. D., Kopf S. H., Weber Y., Chiu B., McFarlin J. M., Elling F. J., Hoeft-McCann S. and
Pearson A. (2023) Controls on the hydrogen isotope composition of tetraether lipids in an
autotrophic ammonia-oxidizing marine archaeon. *Geochim. Cosmochim. Acta* **352**, 194–
210.

Leavitt W. D., Venceslau S. S., Pereira I. A. C., Johnston D. T. and Bradley A. S. (2016)
Fractionation of sulfur and hydrogen isotopes in *Desulfovibrio vulgaris* with perturbed
DsrC expression ed. R. Boden. *FEMS Microbiol. Lett.* **363**, fnw226.

Lengger S. K., Weber Y., Taylor K. W. R., Kopf S. H., Berstan R., Bull I. D., Mayser J., Leavitt
W. D., Blewett J., Pearson A. and Pancost R. D. (2021) Determination of the δ^2H values
of high molecular weight lipids by high-temperature gas chromatography coupled to
isotope ratio mass spectrometry. *Rapid Commun. Mass Spectrom.* **35**, e8983.

Lichtenthaler H. K. (1999) THE 1-DEOXY-D-XYLULOSE-5-PHOSPHATE PATHWAY OF
ISOPRENOID BIOSYNTHESIS IN PLANTS. *Annu. Rev. Plant Physiol. Plant Mol. Biol.*
50, 47–65.

Lichtenthaler H. K., Rohmer M. and Schwender J. (1997) Two independent biochemical pathways
for isopentenyl diphosphate and isoprenoid biosynthesis in higher plants. *Physiol. Plant.*
101, 643–652.

Lynen F., Eggerer H., Henning U. and Kessel I. (1958) Farnesyl-pyrophosphat und 3-Methyl-Δ^3
-butenyl-1-pyrophosphat, die biologischen Vorstufen des Squalens. Zur Biosynthese der
Terpene, III. *Angew. Chem.* **70**, 738–742.

Maloney A. E., Shinneman A. L. C., Hemeon K. and Sachs J. P. (2016) Exploring lipid 2H/1H
fractionation mechanisms in response to salinity with continuous cultures of the diatom
Thalassiosira pseudonana. *Org. Geochem.* **101**, 154–165.

Martens-Habbena W., Berube P. M., Urakawa H., De La Torre J. R. and Stahl D. A. (2009)
Ammonia oxidation kinetics determine niche separation of nitrifying Archaea and
Bacteria. *Nature* **461**, 976–979.

McFarlin J. M., Axford Y., Masterson A. L. and Osburn M. R. (2019) Calibration of modern
sedimentary δ2H plant wax-water relationships in Greenland lakes. *Quat. Sci. Rev.* **225**,
105978.

Möller-Zinkhan D., Börner G. and Thauer R. K. (1989) Function of methanofuran,
tetrahydromethanopterin, and coenzyme F42oin Archaeoglobusfulgidus.

Möller-Zinkhan D. and Thauer R. K. (1990) Anaerobic lactate oxidation to 3 CO2 by
Archaeoglobus fulgidus via the carbon monoxide dehydrogenase pathway: demonstration
of the acetyl-CoA carbon-carbon cleavage reaction in cell extracts. *Arch. Microbiol.* **153**,
215–218.

Morgan R. M., Pihl T. D., Lling J. R. N. and Reeve J. N. (1997) Hydrogen Regulation of Growth,
Growth Yields, and Methane Gene Transcription in Methanobacterium
thermoautotrophicum ⬡H. *J BACTERIOL* **179**.

Osburn M. R., Dawson K. S., Fogel M. L. and Sessions A. L. (2016) Fractionation of Hydrogen
Isotopes by Sulfate- and Nitrate-Reducing Bacteria. *Front. Microbiol.* **7**.

Pearson A. (2014) Lipidomics for Geochemistry. In *Treatise on Geochemistry* Elsevier. pp. 291–
336.

Pereira I. A. C., Ramos A. R., Grein F., Marques M. C., Da Silva S. M. and Venceslau S. S. (2011)
A Comparative Genomic Analysis of Energy Metabolism in Sulfate Reducing Bacteria and
Archaea. *Front. Microbiol.* **2**.

Reed D. W. and Hartzell P. L. (1999) The Archaeoglobus fulgidus D-Lactate Dehydrogenase Is a
Zn2Ƨ Flavoprotein. *J BACTERIOL* **181**.

Rhim J. H., Zhou A., Amenabar M. J., Elling F. J., Weber Y., Pearson A., Boyd E. S. and Leavitt
W. D. (2023) *Mode of carbon and energy metabolism shifts lipid composition in the*
thermoacidophile Acidianus., Microbiology.

Rohmer M., Knani M., Simonin P., Sutter B. and Sahm H. (1993) Isoprenoid biosynthesis in
bacteria: a novel pathway for the early steps leading to isopentenyl diphosphate. *Biochem.*
J. **295**, 517–524.

Russell J. B. (2007) The Energy Spilling Reactions of Bacteria and Other Organisms. *Microb.*
Physiol. **13**, 1–11.

Sachs J. P. (2014) Hydrogen Isotope Signatures in the Lipids of Phytoplankton. In *Treatise on*
Geochemistry Elsevier. pp. 79–94.

Sachs J. P. and Kawka O. E. (2015) The Influence of Growth Rate on 2H/1H Fractionation in
Continuous Cultures of the Coccolithophorid Emiliania huxleyi and the Diatom
Thalassiosira pseudonana ed. G. P. Tochtrop. *PLOS ONE* **10**, e0141643.

Sachs J. P., Maloney A. E. and Gregersen J. (2017) Effect of light on 2H/1H fractionation in lipids
from continuous cultures of the diatom Thalassiosira pseudonana. *Geochim. Cosmochim.*
Acta **209**, 204–215.

Sachs J. P., Maloney A. E., Gregersen J. and Paschall C. (2016) Effect of salinity on 2H/1H
fractionation in lipids from continuous cultures of the coccolithophorid Emiliania huxleyi.
Geochim. Cosmochim. Acta **189**, 96–109.

Sachse D., Billault I., Bowen G. J., Chikaraishi Y., Dawson T. E., Feakins S. J., Freeman K. H.,
Magill C. R., McInerney F. A., van der Meer M. T. J., Polissar P., Robins R. J., Sachs J.
P., Schmidt H.-L., Sessions A. L., White J. W. C., West J. B. and Kahmen A. (2012)
Molecular Paleohydrology: Interpreting the Hydrogen-Isotopic Composition of Lipid
Biomarkers from Photosynthesizing Organisms. *Annu. Rev. Earth Planet. Sci.* **40**, 221–
249.

Schmitz R. A., Linder D., Stetter K. O. and Thauer R. K. (1991) N s,N1~ reductase (coenzyme
F42o-dependent) and formylmethanofurandehydrogenase from the
hyperthermophileArchaeoglobusfulgidus.

Schwender J., Seemann M., Lichtenthaler H. K. and Rohmer M. (1996) Biosynthesis of
isoprenoids (carotenoids, sterols, prenyl side-chains of chlorophylls and plastoquinone) via
a novel pyruvate/glyceraldehyde 3-phosphate non-mevalonate pathway in the green alga
Scenedesmus obliquus. *Biochem. J.* **316**, 73–80.

Schwörer B., Breitung J., Klein A. R., Stetter K. O. and Thauer R. K. (1993) Formylmethanofuran:
tetrahydromethanopterin formyltransferase and NS,N ~ dehydrogenase from the sulfate-
reducing Archaeoglobusfulgidus: similarities with the enzymes from methanogenic
Archaea.

Sessions A. L. (2016) Factors controlling the deuterium contents of sedimentary hydrocarbons.
Org. Geochem. **96**, 43–64.

Sessions A. L., Burgoyne T. W., Schimmelmann A. and Hayes J. M. (1999) Fractionation of
hydrogen isotopes in lipid biosynthesis. *Org. Geochem.* **30**, 1193–1200.

Sessions A. L., Jahnke L. L., Schimmelmann A. and Hayes J. M. (2002) Hydrogen isotope
fractionation in lipids of the methane-oxidizing bacterium Methylococcus capsulatus.
Geochim. Cosmochim. Acta **66**, 3955–3969.

Sessions A. L., Sylva S. P., Summons R. E. and Hayes J. M. (2004) Isotopic exchange of carbon-
bound hydrogen over geologic timescales 1 1Associate editor: J. Horita. *Geochim.*
Cosmochim. Acta **68**, 1545–1559.

Steinsbu B. O., Thorseth I. H., Nakagawa S., Inagaki F., Lever M. A., Engelen B., Ovreas L. and
Pedersen R. B. (2010) Archaeoglobus sulfaticallidus sp. nov., a thermophilic and
facultatively lithoautotrophic sulfate-reducer isolated from black rust exposed to hot ridge
flank crustal fluids. *Int. J. Syst. Evol. Microbiol.* **60**, 2745–2752.

Stetter K. O. (1988) Archaeoglobus fulgidus gen. nov., sp. nov.: a New Taxon of Extremely
Thermophilic Archaebacteria. *Syst. Appl. Microbiol.* **10**, 172–173.

Stouthamer A. H. and Bettenhaussen C. W. (1977) A continuous culture study of an ATPase-
negative mutant of Escherichia coli. *Arch. Microbiol.* **113**, 185–189.

Thauer R. K., Kaster A.-K., Goenrich M., Schick M., Hiromoto T. and Shima S. (2010)
Hydrogenases from Methanogenic Archaea, Nickel, a Novel Cofactor, and H_2 Storage.
Annu. Rev. Biochem. **79**, 507–536.

Valentine D. L. (2007) Adaptations to energy stress dictate the ecology and evolution of the
Archaea. *Nat. Rev. Microbiol.* **5**, 316–323.

Van Der Meer M. T. J., Benthien A., French K. L., Epping E., Zondervan I., Reichart G.-J., Bijma
J., Sinninghe Damsté J. S. and Schouten S. (2015) Large effect of irradiance on hydrogen
isotope fractionation of alkenones in Emiliania huxleyi. *Geochim. Cosmochim. Acta* **160**,
16–24.

Wakil S. J., Stoops J. K. and Joshi V. C. (1983) FATTY ACID SYNTHESIS AND ITS
REGULATION.

Wijker R. S., Sessions A. L., Fuhrer T. and Phan M. (2019) ^{2}H/^{1}H variation in microbial lipids
is controlled by NADPH metabolism. *Proc. Natl. Acad. Sci.* **116**, 12173–12182.

Wolfshorndl M., Danford R. and Sachs J. P. (2019) 2H/1H fractionation in microalgal lipids from
the North Pacific Ocean: Growth rate and irradiance effects. *Geochim. Cosmochim. Acta*
246, 317–338.

Wu W., Meador T. B., Könneke M., Elvert M., Wegener G. and Hinrichs K. (2020) Substrate-
dependent incorporation of carbon and hydrogen for lipid biosynthesis by *Methanosarcina*
barkeri. *Environ. Microbiol. Rep.* **12**, 555–567.

Zhang X., Gillespie A. L. and Sessions A. L. (2009) Large D/H variations in bacterial lipids reflect
central metabolic pathways. *Proc. Natl. Acad. Sci.* **106**, 12580–12586.

Zhang Z. and Sachs J. P. (2007) Hydrogen isotope fractionation in freshwater algae: I. Variations
among lipids and species. *Org. Geochem.* **38**, 582–608.

1142