Dual role of FOXG1 in regulating gliogenesis in the developing neocortex via the FGF signalling pathway.

Mahima Bose[1], Varun Suresh[1#], *Urvi Mishra[1#], *Ishita Talwar[1#], Shiona Biswas[1], Anuradha Yadav[1], Simon Hippenmeyer[2] Shubha Tole[1*]

[1] Tata Institute of Fundamental Research, Mumbai, India, [2] Institute of Science and Technology Austria (ISTA), Klosterneuburg, Austria
[#]*equally contributing co-authors;* Corresponding author email: stole@tifr.res.in

Abstract

In the developing vertebrate central nervous system, neurons and glia typically arise sequentially from common progenitors. Here, we report that the transcription factor Forkhead Box G1 (*Foxg1*) regulates gliogenesis in the mouse neocortex via distinct cell-autonomous roles in progenitors and in postmitotic neurons that regulate different aspects of the gliogenic FGF signalling pathway. We demonstrate that loss of *Foxg1* in cortical progenitors at neurogenic stages causes premature astrogliogenesis. We identify a novel FOXG1 target, the pro-gliogenic FGF pathway component *Fgfr3*, which is suppressed by FOXG1 cell-autonomously to maintain neurogenesis. Furthermore, FOXG1 can also suppress premature astrogliogenesis triggered by the augmentation of FGF signalling. We identify a second novel function of FOXG1 in regulating the expression of gliogenic ligand FGF18 in new born neocortical upper-layer neurons. Loss of FOXG1 in postmitotic neurons increases *Fgf18* expression and enhances gliogenesis in the progenitors. These results fit well with the model that new born neurons secrete cues that trigger progenitors to produce the next wave of cell types, astrocytes. If FGF signalling is attenuated in *Foxg1* null progenitors, they progress to oligodendrocyte production. Therefore, loss of FOXG1 transitions the progenitor to a gliogenic state, producing either astrocytes or oligodendrocytes depending on FGF signalling levels. Our results uncover how FOXG1 integrates extrinsic signalling via the FGF pathway to regulate the sequential generation of neurons, astrocytes, and oligodendrocytes in the cerebral cortex.

Significance Statement

Functional brain circuitry requires the correct numbers and subtypes of neurons and glia. A fundamental feature of the vertebrate central nervous system is that common progenitors produce first neurons and then glia. Cell-intrinsic factors modulate how a progenitor responds to cell-extrinsic cues to achieve this transition. Here, we report that the transcription factor Forkhead Box G1 (*Foxg1*) regulates gliogenesis in the mouse neocortex *in vivo* by regulating different aspects of the gliogenic FGF signalling pathway in progenitors and in postmitotic neurons. Loss of *Foxg1* results in premature gliogenesis via regulation of novel targets *Fgfr3* in progenitors and *Fgf18* in postmitotic neurons. Our findings offer mechanistic insight into FOXG1 syndrome, an Autism Spectrum Disorder caused by *Foxg1* mutations in humans.

Introduction

A fundamental feature of the developing vertebrate central nervous system is that common progenitors produce neurons and glia sequentially. Furthermore, two broad categories of glia, astrocytes and oligodendrocytes, each subserving distinct functions, are also produced sequentially. This temporal sequence of neurogenesis, followed by astrogliogenesis and then oligogenesis, is conserved in the developing mammalian neocortex (1–3). However, the molecular mechanisms instructing a cessation of neurogenesis and initiating gliogenesis within progenitors are poorly understood, particularly in terms of how cell-intrinsic factors interact with cell-extrinsic signalling cues.

Early *in vitro* (1, 4) and *in vivo (5)* studies demonstrated that cortical progenitors first produce neurons and later glia. Quantitative MADM lineage analyses *in vivo* further extended these findings and showed that neurons, astrocytes, and oligodendrocytes arise sequentially within a clone of cells from a single progenitor (5, 6). Molecular insights into this phenomenon identified several pro-gliogenic mechanisms. These include cell-intrinsic transcription factors *Nr2f1/2*, *Zbtb20*, *Sox9,* and *Nfia* (7–10), the Notch pathway (8, 11–14); and cell-extrinsic cues such as interleukins IL6 and Cardiotrophin1 (*Ct1*), Leukemia Inhibiting Factor (*Lif*), Ciliary Neurotrophic Factor (*Cntf*), Neuropoietin (*Np*), Cardiotrophin-like Cytokine (*Clc*), Epidermal Growth Factors (*Egf*) and Fibroblast Growth Factors (*Fgf* (15–20)) which act via the Jak/Stat or Mek/Mapk

signalling pathways *(12, 21, 22)*. In these studies, the primary readout of gliogenesis was astrogliogenesis, using either morphology or molecular markers to identify glial fate. Although a few factors, such as *Sox10,* have been reported to promote oligodendrocyte precursor cell (OPC) fate (23, 24), the progenitor-level mechanisms that govern astrocyte-to-oligodendrocyte transition remain poorly understood. Furthermore, how mechanisms that maintain ongoing neurogenesis crosstalk with those that promote gliogenesis remains to be understood. Another unexplored area is how cell-intrinsic and extrinsic factors interact with or regulate each other at the level of the individual progenitor.

Forkhead family transcription factor FOXG1 is a well-established regulator of the sequential production of neuronal subtypes in the mammalian neocortex (25–30)*.* Further support for a neurogenic role for this factor came from overexpression experiments *in vitro* or cultured progenitors transplanted *in vivo*, in which decreased gliogenesis and downregulation of progliogenic pathways and markers were seen (31–33). However, in these studies, *Foxg1* knockdown did not cause significant gliogenesis (31, 32). Furthermore, a role for FOXG1 in postmitotic neurons has not been examined, although these cells express *Foxg1* and are ideally positioned to signal progenitors to generate the next cell type. These are critical unanswered questions that we explored directly *in vivo*. We show that upon loss of *Foxg1*, progenitors autonomously produce glia at the expense of neurons. Using integrative analysis of the transcriptome, epigenome, and FOXG1 occupancy, we identified FGFR3, part of the established progliogenic FGF signalling pathway, as a FOXG1 target.

We show that the loss of *Foxg1* results in upregulating *Fgfr3* expression and FGF signalling. Whereas overexpression of FGF ligands induces premature gliogenesis (19), simultaneous overexpression of *Foxg1* restores neurogenesis, consistent with its established neurogenic role.

Furthermore, FOXG1 regulates the expression of *Fgf18*, a ligand of the gliogenic FGF8 sub-family, in newborn postmitotic neurons. Therefore, FOXG1 regulates both the availability of the ligand and the level of the receptor for Fgf signalling.

 Finally, attenuation of FGF signalling in a *Foxg1* loss-of-function (LOF) background leads to a premature production of OPCs, indicating that without FOXG1, the progenitor may progress through gliogenesis but cannot return to neurogenesis. Our results shed light on a novel

mechanism regulating the sequential generation of neurons, astrocytes, and oligodendrocytes in the cerebral cortex.

Results

We used a *Foxg1* conditional knockout mouse line with an intrinsic reporter (*Foxg1$^{lox/lox}$*; GFP^{FRT} (34)) to assess the cell-autonomous phenotypes of *Foxg1* disruption. In this line, Cre-mediated recombination of the floxed *Foxg1* allele results in the expression of flippase and consequently activates the expression of a GFP reporter inserted in the *Rosa26* locus. Introducing Cre recombinase into a progenitor results in GFP expression in all cells arising from it, serving as a label for its lineage.

Disrupting *Foxg1* during neurogenesis causes premature gliogenesis.
In wild-type mice, FOXG1 expression decreases in cortical progenitors from embryonic day (E) 15.5 to E18.5 (Supplementary Figure S1; (32)). We tested the consequences of a premature drop in FOXG1 at E15.5 when superficial layer neurons are generated in the neocortex. We introduced a plasmid encoding Cre recombinase at E15.5 using *in-utero* electroporation and examined the brains on postnatal day 14 (P14). In control brains, as expected, GFP+ cells with pyramidal neuron morphologies occupied layer 2/3 of the neocortex, whereas cells with glial morphologies were scattered throughout the tissue. 67% of the GFP+ cells also expressed neuronal marker NEUN and did not express mature astrocytic marker ALDH1L1. 16% of the GFP+ cells were also ALDH1L1+ (Figure 1A–D; Supplementary Figure S1). In contrast, Cre electroporation in *Foxg1$^{lox/lox}$* embryos results in GFP+ cells with astrocytic morphologies distributed all over the neocortex. None were NEUN+, but 56% were ALDH1L1+**,** confirming their astrocyte identity. We found similar results, i.e. premature gliogenesis, upon Cre electroporation at E14.5 (Figure 1E–H). This suggests that upon loss of FOXG1, progenitors switch from neurogenesis to gliogenesis, even if they have yet to generate the last cohorts of superficial layer neurons.

Enhanced gliogenesis upon loss of *Foxg1* is not due to the over-proliferation of mutant astrocytes.

Increased gliogenesis may arise from the over-proliferation of astrocytes and an accompanying death of neurons that have lost *Foxg1*. Alternatively, there could be a premature cell fate switch in the progenitors that should have produced neurons but instead produced astrocytes (Figure 2A, B). To distinguish between these scenarios, we examined the presence of KI67 in cells at E18.5 three days after Cre electroporation at E15.5. Similar numbers of electroporated (GFP+) cells displayed KI67 in control and *Foxg1lox/lox* brains, indicating that loss of *Foxg1* did not cause overproliferation (Figure 2B, B'). Furthermore, no difference was observed in the expression of a cell death marker Cleaved Caspase 3 (Supplementary Figure S2C, C'). Finally, consistent with a switch to gliogenesis, the *Foxg1* LOF cells were positive for gliogenic factor NFIA (Figure 2 C, C') and PAX6 (Supplementary Figure S2A, A') but not the neurogenic intermediate progenitor marker TBR2 (Supplementary Figure S2B, B').

We also employed a MADM (Mosaic Analysis with Double Markers)-based strategy to distinguish between these possibilities (Figure 2D) (35, 36). We used the MADM-12 (M12) line (36, 37) in which the MADM GT and TG cassettes are on chromosome 12 and crossed them onto a *Foxg1lox/+* background. In the resulting *Foxg1*-MADM (*M12GT/TG, Foxg1*) brains, a fraction of Cre-electroporated progenitors undergo recombination, resulting in progeny that is either *Foxg1−/−*, GFP+ (green) or *Foxg1+/+* tdTomato+ (red) cells. Since these red and green cells arise from single progenitors, the MADM system enables an accurate analysis of proliferation versus cell fate transformation upon disruption of *Foxg1*. A fraction of progenitors in the MADM paradigm undergoes a different recombination pattern, resulting in a yellow progeny due to the expression of both GFP and tdTomato (35, 36), which were not scored. We performed *in-utero* electroporation of Cre in *Foxg1*-MADM embryos in E14.5/E15.5 embryos. We analysed the brains at P14 (Figure 2D). Green and red cells were scored for neuronal or glial identity based on morphology (5, 20). Of the total control (red, +/+) cells, 61.9% were neurons, and 38.1% were glia. In contrast, of the total number of *Foxg1* mutant (green,–/–) cells, 16.7% were neurons, and 83.3% were glia.

In summary, the results demonstrate that loss of *Foxg1* results in a premature cell fate switch in neurogenic progenitors that makes them gliogenic.

Loss of *Foxg1* causes upregulation of *Fgfr3* in progenitors and enhanced FGF signalling.

We investigated how FOXG1 regulates the transcriptional landscape in cortical progenitors to maintain a temporal control of neurogenesis and gliogenesis. We used the inducible *hGFAPCreERT2* (hGCE) driver to achieve widespread recombination in apical progenitors at embryonic ages upon induction with tamoxifen (38) (39). First, we ascertained that hGCE-driven loss of *Foxg1* at E15.5 recapitulates the premature gliogenesis phenotype we described in Figure 1 (Supplementary Figure S3A). Also, two days after tamoxifen induction (E15.5 → E17.5), *Foxg1* LOF cells were positive for gliogenic factor NFIA, similar to that seen upon Cre electroporation (Supplementary Figure S3B; Figure 2C). To identify pathways that regulate the premature gliogenesis induced by the loss of *Foxg1*, we collected FACS-sorted control and mutant cells at E17.5, 48 hours post-tamoxifen administration at E15.5, for transcriptomic analysis (Supplementary Figure S3C). As expected, the loss of *Foxg1* caused the upregulation of known gliogenic factors such as *Nfia* and *Olig3* and the downregulation of neurogenic genes such as *Neurod2* and neuronal marker *Pou3f1* (Figure 3A).

We further filtered the list of genes upregulated upon loss of *Foxg1* using a strategy designed to narrow down candidates likely to drive gliogenesis. First, we examined a set of genes known to display FOXG1 occupancy at E15.5 (40). Second, we reasoned that cell fate transitions are driven by genes maintained in states poised for transcriptional activation or repression by displaying both epigenetic marks (41). Therefore, we analysed a dataset from E15.5 radial glial cells to identify genes that carry bivalent H3K4Me3 and H3K27Me3 marks in gene regulatory regions. Third, we obtained a list of genes enriched in cortical astrocytes (42). The overlap of these three datasets yielded a shortlist of 19 genes representing potential FOXG1 targets that may be suppressed during neurogenesis (Figure 3B, C). Finally, we compared this multimodal analysis with the list of genes we discovered to be upregulated upon loss of *Foxg1* and identified four targets: *Col2a1, Dpysl4, Fgfr3,* and *Pax6* (Figure 3D). The presence of *Fgfr3* in this set presented the exciting possibility that FOXG1 may function to suppress the potent progliogenic FGF signalling pathway (19) in neurogenic cortical progenitors.

In summary, we identified *Fgfr3* as a novel FOXG1 target. Upregulation of *Fgfr3* may mediate the gliogenic effects of loss of *Foxg1*. Consistent with this hypothesis, the MAPK/ERK

pathway, which mediates FGF signalling, was identified as a prominent upregulated pathway in a KEGG pathway analysis of the RNA-seq data (Figure 3E). Activation of this pathway results in the phosphorylation of ERK1/2 and its translocation to the nucleus to regulate the downstream targets of FGFR activation (43). To assess whether the loss of *Foxg1* indeed has functional consequences on FGF signalling, we examined FGFR3 and nuclear phospho-ERK1/2 levels in control and *Foxg1* mutant cortical progenitors. Loss of *Foxg1* at E15.5 led to an increase in FGFR3 labelling and nuclear localisation of phospho-ERK1/2 by E17.5 (Figure 3F–H). We quantified nuclear FOXG1 levels in wild-type apical progenitors. We found a significant decrease from E15.5 to E18.5, consistent with previous findings (32), suggesting an endogenous mechanism for *Fgfr3* upregulation with the initiation of gliogenesis (Supplementary Figure S4).

These results demonstrate that FOXG1 suppresses *Fgfr3* gene expression and the MAPK/ERK pathway, and loss of *Foxg1* results in an enhancement of this pro-gliogenic pathway. The decline in nuclear FOXG1 by E18.5 suggests an endogenous mechanism for the transition of neurogenesis to gliogenesis in apical progenitors.

Foxg1 **overexpression cell autonomously suppresses FGF-induced astrogliogenesis.**
Exogenous FGF8 is known to have potent progliogenic effects in the neocortex (19), which our results recapitulated (Figure 4A, B). As expected, electroporation at E15.5 of a control construct encoding EGFP did not impede neurogenesis. In wild-type mice, 100% of the electroporated GFP+ cells were NEUN+ and occupied the superficial layers of the neocortex by P7 (Figure 4A). Similar to the findings of (19), co-electroporation of constructs encoding *Fgf8*+*Egfp* induced premature gliogenesis, resulting in 85% of the GFP+ cells displaying astrocytic morphologies and glial marker SOX9 (Figure 4B). *Foxg1* overexpression alone did not affect neurogenesis but gave rise to neurons, some of which occupied the superficial layer and some displayed migration deficits, as previously shown ((34); Figure 4C).
We tested whether FOXG1 can override FGF8-induced premature gliogenesis by co-electroporating both constructs (*Foxg1-Egfp* and *Fgf8*, Figure 4D). This led to a recovery of the neuronal fate in a cell-autonomous manner, such that GFP+ cells also displayed NEUN but not SOX9 (Figure 4D). The electroporated cells also displayed migration deficits consistent with

Foxg1 overexpression (34). An interesting additional result was that while these FOXG1-EGFP+ cells themselves expressed neuronal marker NEUN, they were surrounded by SOX9+ cells consistent with the non-cell-autonomous gliogenic effects of *Fgf8* electroporation (Figure 4D).

In summary, FOXG1 is sufficient to cell-autonomously suppress the pro-gliogenic effects of FGF8, establishing it as a regulator of FGF signalling within cortical progenitors.

Postmitotic neuron-specific loss of *Foxg1* leads to premature gliogenesis in cortical progenitors.

Postmitotic neurons provide "feedback instruction" to progenitors via cues that modulate the cell types that arise subsequently (44, 45). We tested whether *Foxg1,* expressed in postmitotic neurons (29, 46), regulates gliogenic secreted factors in these cells. We used postmitotic neuron-specific *NexCre* to disrupt *Foxg1* and examined these brains at birth. We found a significant increase in the levels of gliogenic factor SOX9 in the ventricular zone progenitors compared to that in control brains, indicating a non-autonomous effect of neuron-specific loss of *Foxg1* on progenitors (Figure 5A–D).

To identify potential gliogenic molecules regulated by FOXG1 in postmitotic neurons, we performed transcriptomic analysis on cortical plate tissue isolated from control and *NexCre*-driven *Foxg1* LOF brains at birth (Figure 5E, Supplementary Table 4). *Fgf18,* a member of the Fgf8 family (47) and known to be expressed by neurons during the early postnatal window (48), was significantly upregulated upon postmitotic neuron-specific loss of *Foxg1* (Figure 5E). Consistent with enhanced gliogenesis, there was an apparent increase in staining of astrocyte marker ALDH1L1 in the entire cortical plate (Figure 5F). Furthermore, in these brains, phosphorylated p42/44-ERK1/2, indicative of active FGF signalling, is increased specifically in the SOX2+ progenitors in the ventricular zone (Figure 5G-I).

We examined whether the temporal expression of *Fgf18* correlates with the timing of astrogliogenesis in the cortex. We used a publicly available transcriptomic dataset from CUX2+ upper-layer neurons isolated at stages from E18.5 to P48 (49). In this dataset, *Fgf18* mRNA levels peaked in the first postnatal week and dropped thereafter, consistent with the temporal profile of astrogliogenesis in the cortex (Figure 5J, B; (50)).

Our results indicate that FOXG1 regulates the expression of *Fgf18* in postmitotic neurons and non-autonomously triggers a transition to gliogenesis in the progenitors residing in the ventricular zone.

Attenuation of FGF signalling does not restore neurogenesis in *Foxg1* mutant progenitors but causes premature oligogenesis.

Loss of *Foxg1* appears to promote astrogliogenesis via two distinct functions: enhancing FGFR3 expression in progenitors and increasing expression of *Fgf18* in postmitotic neurons. We, therefore, tested whether a reduction of available FGF ligands may be sufficient to restore neurogenesis in *Foxg1* LOF progenitors. We used a construct encoding soluble FGFR3c (sFGFR3c), which has been effectively used by other studies to sequester and limit the availability of FGFs using *in-utero* electroporation (19, 51).

As before (Figure 1), embryos were electroporated at E15.5, and the brains were harvested at P14 and examined for glial markers either common to glial precursor cells (OLIG2) or exclusive to either OPCs (PDGFRA) or astrocytes (ALDH1L1). For two of the conditions, control (Cre electroporation in *Foxg1^lox/+*) and *Foxg1* LOF alone (Cre electroporation in *Foxg1^lox/lox*), brains from the same set of experiments presented in Figure 1 were used from which additional sections were examined for OLIG2 and PDGFRA (Figure 6A and B). In controls, the majority of GFP⁺ cells were neurons that did not express any glial marker (Figure 1, 6A, 6D). Neurogenesis was significantly reduced upon loss of *Foxg1* alone, and there was a corresponding increase in both types of glia (Figure 6B, D). For the new experimental condition, which involves a combined loss of *Foxg1* and abrogation of FGF signalling, we co-electroporated *Cre* with *sFgfr3c* into *Foxg1^lox/lox* embryos. Strikingly, this condition did not restore neurogenesis but instead shifted the cell fate towards OLIG2+, PDGFRA+, and ALDH1L1⁻ OPCs (Figure 6C, D).

These results indicate that neurogenesis can neither proceed normally nor be restored by abrogating FGF signalling in the absence of FOXG1. Instead, the progenitors display a premature progression to an OPC fate, which appears to be a permissive cell fate in the context of the combined loss/decrease of FOXG1 and FGF signalling, respectively.

In summary, the results uncover a fundamental role of FOXG1 in regulating the transition of cortical progenitors from neurogenesis to gliogenesis and also highlight that neurogenesis is not possible in the absence of this critical factor.

Discussion

Cell type diversity arises due to genetic mechanisms that regulate how progenitors give rise to cells with unique identities. In the nervous system, neurons, astrocytes, and oligodendrocytes have distinct functions, and the production of the correct numbers of each of these three cell types must be carefully controlled. How these cells arise from common progenitors is one of the outstanding questions in developmental neuroscience. Progenitors regulate the cell fate of their progeny via cell-autonomous, *i.e.* intrinsic programs and cell non-autonomous, i.e. extrinsic signalling mechanisms. Understanding how these regulatory processes are orchestrated and how the progenitor integrates these effects is crucial in elucidating how these transitions in cell fate arise. Our study reveals two novel functions of transcription factor FOXG1 in controlling both the expression of an FGF family member *Fgf18* in upper layer cortical neurons, as well as the sensitivity of progenitors to FGF ligands via the expression of the receptor FGFR3. Thus, FOXG1 integrates the production and receptivity to extrinsic signals so progenitors may unfold cell-intrinsic programs to produce glia.

Whereas the different stages of neurogenesis have been well studied (52, 53), the progression from neurogenesis to gliogenesis and through the various stages of gliogenesis has been less well understood. *Foxg1* was previously identified as a regulator of sequential neurogenesis in the neocortex via controlling the timing and order of generation of specific neuronal subtypes (26, 29, 30, 40). *Foxg1* overexpression in E12 cortical progenitors *in vitro* was shown to suppress gliogenesis (32, 33). However, in these studies, the Fgf pathway was not identified as a possible FOXG1 target, and *Foxg1* knockdown did not result in gliogenesis. Therefore, the endogenous role of FOXG1 could not be established (32, 33).

Cortical progenitors express *Fgfr1, r2*, and *r3* from early stages (19, 54), and the expression levels increase during gliogenesis (19). Our work demonstrates that the timing of the cessation of neurogenesis and the subsequent emergence of glial lineages is controlled by FOXG1 in cortical progenitors *in vivo* and identifies a key target, *Fgfr3*, via which FOXG1 controls the response of the progenitor to extrinsic FGF signalling. Importantly, we find that progenitors at

E14.5 as well as E15.5 produce glia upon loss of *Foxg1*. This indicates that *Foxg1* mutant progenitors do not simply accelerate to the immediate next stage of cell type production, i.e. layer 2/3 instead of layer 4, but undergo a switch to producing a different class, astroglia. A decrease in FOXG1 in apical progenitors (Supplementary Figure S4 and (32) may be an endogenous mechanism for the transition of neurogenesis to gliogenesis in progenitors. Postmitotic neurons are known to provide cues to progenitors to initiate the production of the next wave of cells (44), (45), (48). In this context, we identify an additional function of FOXG1 in regulating the expression of the FGF18 ligand from postmitotic neurons, providing an elegant mechanism for timing the onset of astrogliogenesis after neurogenesis is complete. *In vivo*, *Foxg1* expression is dynamically downregulated in pyramidal neuronal precursors as they migrate in the intermediate zone (34), which provides a potential endogenous mechanism for the upregulation of *Fgf18* by postmitotic neurons. Additional regulatory mechanisms within maturing postmitotic neurons that remain unidentified may also participate in modulating such feedback signalling to progenitors.

FGF signalling has essential roles in the early patterning of the cortical primordium (51, 55, 56) and the guidance of major axon tracts in the brain (57). However, its importance during late embryonic and postnatal stages has yet to be fully explored. Recent evidence suggests that FGF signalling regulates neurogenesis and neuronal migration (19, 58, 59). Depletion of FGF from E15.5 progenitors leads to prolonged neurogenesis in the cortex (19). However, reducing FGF signalling in *Foxg1* mutant progenitors does not restore neurogenesis but causes a premature production of OPCs. These data indicate that FOXG1 is essential for neuronal production, and in its absence, the removal of FGF is insufficient to prolong the neurogenic programme in progenitors. Overexpression of *Foxg1*, however, does restore neurogenesis despite augmented FGF signalling. This establishes that FOXG1 is necessary and sufficient for neuronal fate regardless of FGF signalling levels.

A rich body of work has explored how FGF, SHH, BMP, and WNT signalling regulate the proliferation and differentiation of OPCs *in vitro* and the spinal cord (60–63). However, there is a lack of information on how OPCs arise from gliogenic progenitors and whether there are differences in gliogenic mechanisms between the spinal cord and cerebral cortex, given that the initial molecular context of progenitors in these two structures is vastly different (64, 65). Our data indicates that once a neocortical progenitor can no longer generate neurons due to

loss of *Foxg1*, it is forced into a gliogenic mode, producing astrocytes if FGF signalling is present and OPCs if FGF signalling is abrogated. *In vivo*, a decline in FGFs available to progenitors may initiate OPC production. This is consistent with a general mechanism in which the regulated secretion of the ligands from previously born cells governs the timing of generation and proliferation of later-born cells (44). In the particular context of astrogliogenesis and oligogenesis, a more in-depth analysis of the affinity of FGF-FGFR binding in progenitors and specified glial cells is required to understand the fine-tuning of these processes (66, 67). FOXG1 modulates the cell-intrinsic response of progenitors to cell-extrinsic FGF signalling, thereby regulating the production of neurons, astroglia, and oligodendrocytes, the three major components of brain circuitry. In parallel, FOXG1 also regulates FGF ligand expression in postmitotic neurons, which triggers gliogenesis in progenitors. In humans, either gain- or loss-of-function *Foxg1* mutations cause "Foxg1 syndrome", an Autism Spectrum Disorder (29, 68, 69) that results in a range of conditions that include microcephaly, seizures and may also increase the propensity for glioblastoma or myelination deficits (29). Our findings offer mechanistic insights into aspects of Foxg1 syndrome-associated neurodevelopmental deficits.

Materials and Methods

<u>Mice</u>

All procedures followed the Tata Institute of Fundamental Research Animal Ethics Committee (TIFR-IAEC) guidelines. The *Foxg1^{lox/lox}; FRT* mouse line used in this study is described in (34). MADM-12 GT/GT and MADM-12 TG/TG lines are described in (36). *hGFAPCreERT2* (Strain #:012849) and *Ai9* reporter mouse line (Strain #:007909) were obtained from Jackson Laboratory. The *NexCre/+* mouse line was obtained from Klaus Nave, Max Planck Institute for Experimental Medicine (70).

All animals were kept at an ambient temperature and humidity, with a 12h light-dark cycle and food available ad libitum. Noon of the day of the vaginal plug was designated as embryonic day 0.5 (E0.5). Both male and female animals were used for all experiments.

Primers for genotyping were (expected band sizes):

Foxg1cKO F: CCACTCCGAACCCGCTGG,

Foxg1cKO R: AGGCTGTTGATGCTGAACGA, (mutant: 190 bp, WT: 156 bp);

FRT Reporter:

RCE-Rosa1: CCCAAAGTCGCTCTGAGTTGTTATC,

RCE-Rosa2: GAAGGAGCGGGAGAAATGGATATG,

RCE-Cag3: CCAGGCGGGCCATTTACCGTAAG, (WT: 550bp, FRT: 350bp).

MADM 12 Cassette:

Chr 12 WT F: CACTAAGCTCCACTCGCACC,

Chr 12 WT R: TCCCTCATGATGTATCCCCT,

MADM R: TCA ATG GGC GGG GGT CGT T, (WT: 322bp, MADM Cassette: 200bp).

Cre F: ATTTGCCTGCATTACCGGTC.

Cre R: ATCAACGTTTTCTTTTCGG (Cre: 350bp)

NexCre Forward: GAGTCCTGGAATCAGTCTTTTTC

NexCre Reverse: AGAATGTGGAGTAGGGTGAC

NexCre Mutant Reverse: CCGCATAACCAGTGAAACAG (WT: 770 bp, NexCre: 525 bp)

In utero electroporation:

In utero, electroporation was performed as previously described (71). Embryos were injected with 1–2 μL of plasmid DNA solution dissolved in nuclease-free water with 0.1% fast green with plasmid DNA into the lateral ventricle through the uterine wall using a fine glass microcapillary (Sutter capillaries #B100-75-10). Constructs pCAGG-IRES-eGFP and pCAGG-IRES-FOXG1-EGFP were gifts from Gord Fishell, Harvard Medical School. pCAGG-FGF8 and pCAGG-sFGFR3c were gifts from Hiroshi Kawasaki, Kanazawa University.

Tissue Preparation: Embryos were isolated in ice-cold PBS. Embryonic brains were dissected and fixed overnight in 4% (wt/vol) paraformaldehyde at 4°C overnight and then cryoprotected by transferring to 30% (wt/vol) sucrose-PBS until sectioning. Postnatal mice were anaesthetised using Thiopentone and transcardially perfused with 4% (wt/vol) paraformaldehyde in phosphate buffer, followed by overnight fixation and then cryoprotected by transferring to 30% sucrose-PBS until sectioning. The brains were sectioned at 30μm and 40μm for MADM using a freezing microtome (Leica SM2000R).

Immunohistochemistry: Brains were sectioned (30μm), mounted on Superfrost plus glass microscope slides (Cat #71869-10), and dried for 2 h at room temperature. Three washes were given for 5 minutes each (3*5') in phosphate buffer. All antibodies except FGFR3 and PDGFRα required antigen retrieval at 90°C–95°C in 10 mM sodium citrate buffer (pH 6.0) for 10 min. Sections were immersed in blocking solution (5% (vol/vol) horse serum in phosphate buffer with 0.1% (vol/vol) Triton X-100 (Sigma; X100)) for 1 h at RT. Incubation with primary antibody was performed in phosphate buffer containing 0.1% (vol/vol) Triton X-100 and 2.5% (vol/vol) horse serum at 4°C overnight. For postnatal brain sections, free-floating immunohistochemistry was performed. Sections were given three washes for 5 minutes each (3*5') in phosphate buffer and then permeabilised with phosphate buffer containing 0.3% (vol/vol) Triton X-100 for 10 minutes. Blocking was done with 5% (vol/vol) horse serum in phosphate buffer with 0.3% (vol/vol) Triton X-100 for one hour at RT. This was followed by primary antibody treatment in phosphate buffer containing 0.3% (vol/vol) Triton X-100 and 2.5% (vol/vol) horse serum overnight at 4°C. The sections were washed in phosphate buffer, followed by the appropriate secondary antibody for two hours at room temperature.

This was followed by three washes for 5 minutes each (3*5') in phosphate buffer and DAPI staining for 10 minutes, after which the sections were washed with phosphate buffer for 15 minutes (3*5'). The slides were then mounted with Fluoroshield (Sigma cat no F6057 or F6182). Information on antibodies and their catalogue numbers are mentioned in SI Appendix Supplementary Table 1 and 2.

Image acquisition and analysis: Fluorescence images were taken using an Olympus FluoView 3000 confocal microscope with FluoView software. All the image analysis was done on Fiji-ImageJ. A nonlinear operation such as gamma correction was not performed in the figures. Brightness and contrast adjustments were performed identically for control and mutant conditions. Cell counting was performed using the Cell Counter plugin in Fiji.
Intensity quantifications for Phospho-p42/44-ERK1/2 and SOX9 were done by imaging Control and mutant samples at the same laser and gain settings on the FV3000 confocal microscope. Images were stacked based on one-cell thickness stacks, i.e. approximately 5µm. Intensity quantification was done by referencing the drawing ROIs of the nuclear areas referencing DAPI or SOX2, followed by intensity measurement of Phospho-p42/44-ERK1/2 and calculating mean intensity per unit area (ROI).

Histone and FOXG1 ChIP Seq analysis: FASTQ files deposited by previously published manuscripts (41) and (40) were obtained using SRA Toolkit's fastq-dump command and aligned to mouse reference genome mm10 using Bowtie2 (72). Peak calling and annotation were performed using Homer using default options and command style "-histone" for the H3K27Me3 and H3K4Me3 datasets. For the FOXG1 ChIP-seq analysis, peaks were called using all "-style" options and only the top 50% peaks called were considered for further analysis for all the ChIP-seq datasets. Peaks were intersected using BEDTools software (73) and annotated using Homer (74) annotatePeaks.pl function, and genome view plots were created using IGV (75).

Fluorescence-Assisted Cell Sorting:

Foxg1 mutant cells were obtained by administering tamoxifen (Cat #T5648) prepared in corn oil (Sigma; Cat #8267) to E15.5 *hGFAPCreERT2*; *Foxg1^lox/lox^; Ai9* dams, at 40 µg/g body weight dose and FACS-isolated using the Ai9 reporter at E17.5. We used FlashTag-labelled (Thermo Fisher Scientific SKU C34554) progenitors at E15.5 for controls and collected the cells at E17.5. Cortical tissue was dissected in HBSS without Calcium/Magnesium (Thermo Fischer Scientific Catalogue number: 14170112), and a single-cell suspension was prepared in HBSS with Calcium/Magnesium (Thermo Fischer Scientific Catalogue number: 14025092) using 0.25% Trysin (ThermoFischer Scientific Gibco Catalogue number: 15400054) and 70um cell strainer. FACS was performed using BD Aria Fusion (BD Biosciences) with the 568 and 488 lasers using an 85µm nozzle. Singlets were selected using forward scatter and side scatter. Cells were selected for collection based on their RFP signal (Mutant) or GFP signal (Control).

RNA seq - sample preparation and analysis: Cells obtained from FACS were stored in RNAlater until extraction. 400,000 cells were pooled from at least two brains to obtain a biological replicate. RNA extraction and sequencing were performed on three control replicates and four *Foxg1* mutant replicate cell suspensions. 1ug of RNA was used to obtain the cDNA library, and sequencing was performed on an Illumina platform to achieve 150 bp reads to generate 30 Million paired-end reads per sample. Fastq QC was performed as described in (https://www.bioinformatics.babraham.ac.uk/projects/fastqc/) and reads > 30 Phred scores were aligned using HISAT2 (76). Feature counts were used to quantify the number of reads per transcript. Differential expression analysis was performed using DESeq2 (77). A log2FoldChange cutoff of 0.58 and P-value < 0.05 was used to identify DEGs. GO analysis was performed using Shiny GO (78). Genes described as Astrocyte-enriched were obtained from (42).

Data availability: All RNAseq samples generated for this manuscript are deposited under PRJNA886320.

Reviewer's link:

https://dataview.ncbi.nlm.nih.gov/object/PRJNA886320?reviewer=mgjl04qlpcnfspn5t58g2p39n4

Acknowledgements

We thank Dr. Shital Suryavanshi and the animal house staff of the Tata Institute of Fundamental Research (TIFR) for their excellent support; Gord Fishell and Goichi Miyoshi for the *Foxg1* floxed mouse line; Hiroshi Kawasaki for the plasmids pCAG-FGF8 and pCAG-sFGFR3c. We thank Prof. S.K. Lee for the *Foxg1^{lox/lox}* genotyping primers and protocol. We thank Dr. Deepak Modi and Dr. Vainav Patel for allowing us to use the NIRRCH FACS Facility and the staff of the NIRRCH and TIFR FACS facilities for their assistance.

We thank Denis Jabaudon for his critical comments on the manuscript and members of the Jabaudon lab for helpful discussions. This work was funded by the Department of Atomic Energy (DAE), Govt. of India (Project Identification no. RTI4003, DAE OM no. 1303/2/2019/R&D-II/DAE/2079).

Figures

Figure 1: Loss of *Foxg1* causes premature gliogenesis

(A–D) Cre electroporation at E15.5 in control (A, *Foxg1*lox/+; *GFP* FRT) and *Foxg1* LOF (B, *Foxg1*lox/lox; *GFP* FRT) embryos, followed by analysis at P14. (C, D) 67.3% of GFP+ cells colocalised with NeuN in control brains and 1% in *Foxg1* LOF brains. 16% of GFP+ cells colocalised with ALDH1L1 in control brains and 56% in *Foxg1* LOF brains. n= 2151 (Control), 2761 (*Foxg1* LOF) cells from N=3 brains (biologically independent replicates). (E–H) Cre electroporation at E14.5 in control (E, *Foxg1*lox/+; *GFP* FRT) and *Foxg1* LOF (F, *Foxg1*lox/lox; *GFP* FRT) embryos, followed by analysis at P14. (G, H) 77.7% of GFP+ cells colocalised with NeuN in control brains and 0% in *Foxg1* LOF brains. 20.1% of GFP+ cells colocalised with ALDH1L1 in control brains and 55.7% in *Foxg1* LOF brains. n= 3160 (Control), 2978 (*Foxg1* LOF) cells from N=3 brains (biologically

independent replicates). The leftmost panels in each row are low-magnification images, as the scale bar indicates. Filled arrowheads depict colocalisation, and open arrowheads depict non-colocalisation of marker and electroporated cells. *Statistical test: two-tailed unpaired t-test. *(p<0.05), **(p<0.01), ***(p<0.001), ****(p<0.0001) . All scale bars are 50 μm.*

Figure 2: Foxg1 LOF leads to premature glial fate acquisition in progenitors but no proliferation defects.

(A) Schematic depicting the two alternative mechanisms that could result in enhanced gliogenesis upon loss of *Foxg1:* a change in cell type produced by the progenitor, i.e. "fate switch" (A) or increase in proliferation of astrocytes (B). (B, B', C, C') Images of the region of the ventricular and sub-ventricular zone (the dashed line indicates the ventricular boundary). Cre electroporation at E15.5 in control (*Foxg1*lox/+; *GFP* FRT) and *Foxg1* LOF (*Foxg1*lox/lox; *GFP* FRT) embryos, followed by analysis at E18.5. Proliferation marker KI67 (B, B') colocalises with similar numbers of GFP⁺ cells in control and *Foxg1* LOF brains. In contrast, glial progenitor markers NFIA (C, C') display increased colocalisation with GFP+ cells, n= 3590 (control), 2100 (mutant) cells from N=3 brains (biologically independent replicates). (D) Schematic depicting the genotype and corresponding fluorescent labels resulting from the MADM recombination. (E–F) Cre electroporation at E14.5/E15.5 in *Foxg1*-MADM brains (*M12*GT/TG, Foxg1) analysed at P14. Green (*Foxg1*⁻/⁻) and red (*Foxg1*+/+) cells were scored based on neuronal (open arrowheads, E) or glial (arrowheads, E) morphology. (F) represents the number of neurons or glia as a percentage of the total population of neurons+glia of each genotype: control (red; +/+) or *Foxg1* mutant (green;–/–) neurons. n= 354 cells from N=5 brains (biologically independent replicates). *Statistical test: two-tailed unpaired t-test. *(p<0.05), **(p<0.01), ***(p<0.001), ****(p<0.0001). All scale bars: 50µm.*

Figure 3: Foxg1 binds and regulates the expression of FGFR3

(A) RNA-seq analysis of FACS-purified Control and *Foxg1* LOF progenitors harvested two days after labelling at E15.5. Gliogenic factors such as *Nfia, Id3,* and *Olig3* are upregulated, and neuronal markers such as *Pou3f1* and *Robo4* are downregulated. (B–D) Multimodal analysis comparing FOXG1 occupancy (ChIP-seq) and bivalent epigenetic marks (H3K4Me3 and H3K27Me3) and astrocyte-enriched genes from (42) reveals a list of 19 genes common to each dataset (B). Four of these are upregulated upon loss of *Foxg1*, including the known gliogenic gene *Fgfr3* (D). FOXG1 occupies a -26kb enhancer region of *Fgfr3* (C). (E) KEGG analysis of the upregulated genes from (A) identifies the MAPK signalling pathway downstream of FGF signalling. (F) Loss of *Foxg1* from E15.5 progenitors (*hGFAPCreERT2*, tamoxifen at E15.5) causes upregulation of FGFR3 receptor by E17.5 (G; n=50 (Control and *Foxg1* LOF) ROIs from N=2 brains) and phosphorylated-ERK1/2 (H; n= 67 (Control) and 89 (*Foxg1* LOF) cells from N=3 brains (biologically independent replicates). *Statistical test: Mann-Whitney test *(p<0.05), **(p<0.01), ***(p<0.001), ****(p<0.0001). All scale bars: 50µm.*

Figure 4: Foxg1 suppresses FGF-induced gliogenesis.

(A–D) *In utero* electroporations were performed in wild-type embryos at E15.5, and the brains were analysed at P7. (A) GFP electroporation labels L2/3 cells that are NEUN+ (arrowheads) and SOX9⁻(open arrowheads). (B) Overexpression of *Fgf8* leads to premature gliogenesis, and the GFP+ cells are NEUN⁻ (open arrowheads) and SOX9⁺ (arrowheads). (C) Overexpression of *Foxg1* produced NEUN+ (arrowheads) and SOX9⁻ (open arrowheads) neurons, some of which displayed delayed migration (black asterisk), and others migrated to the cortical plate (white asterisk), as shown in (34). (D) Overexpression of *Foxg1* together with FGF8 partially rescued neuronal fate such that GFP+ cells also displayed NEUN (arrowheads) but not SOX9 (open arrowheads). In A—D, the leftmost panels in each set are low-magnification images, as indicated by the scale bar.

(E) Quantifications of GFP+ cells that are also NEUN+ in each condition: 98.6% (GFP); 1.8% (FGF8); 98.3% (Foxg1); 74.1% (*Foxg1+Fgf8*). (F) Quantifications of GFP+ cells that are also SOX9+ in each condition: 0% (*Egfp*); 87.7% (*Fgf8+Egfp*); 0% (*Foxg1-Egfp*); 24.2% (*Foxg1-Egfp+Fgf8*). n= 2123 (*Egfp*), 1643 (*Fgf8+Egfp*), 1357 (*Foxg1-Egfp*),1924 (*Foxg1-Egfp+Fgf8*) cells each from N=3 brains (biologically independent replicates).

Statistical test: Two-way ANOVA. *(p<0.05), **(p<0.01), ***(p<0.001), ****(p<0.0001). All scale bars: 50µm.

Figure 5: Postmitotic neuron *Foxg1* LOF leads to premature gliogenesis and upregulation of the MAPK pathway. (A) SOX9 staining in the control P0 cortex identifies gliogenic progenitors at the ventricular zone (white asterisk) and scattered cells throughout the cortical plate. (B-D) *NexCre*-driven loss of *Foxg1* is specific to postmitotic neurons, as seen by GFP reporter expression (C, white bars) and causes a non-autonomous upregulation of nuclear SOX9 in the ventricular zone progenitors (B; quantifications: D). B and C are images of the same section showing SOX9 alone (B) and together with the GFP reporter (C). (E) Transcriptomic analysis of cortical plate tissue from control and *NexCre/+; Foxg1^lox/lox; GFP^FRT* reveals a significant upregulation of *Fgf18* upon loss of *Foxg1*. (F- I) Analysis at P6 shows an increased ALDH1L1 in *NexCre/+; Foxg1^lox/lox* brains (F) and an increase in phosphorylated p42/44-ERK1/2 in SOX2+ progenitors at P0 (G–I), indicating enhanced FGF signalling. G and H are images of the same section showing pERK1/2 alone (G) and with SOX2 (H). Nuclear pERK1/2 levels in SOX2+ progenitors are quantified in (I). (J) *Fgf18* expression in CUX2+ upper layer cells peaks at P7, as seen in the RNA seq dataset from (49). Quantification of SOX9+ cells in each condition: Quantifications of SOX9+ in each condition: 137 (Control) and 198 (*NexCre/+; Foxg1* LOF) cells from N=3 brains (biologically independent replicates). Quantifications of phospho-ERK1/2 in SOX2+ cells in each condition: 207 (Control) and 189 (*NexCre/+; Foxg1* LOF) cells from N=3 brains (biologically independent replicates). *Statistical test: Mann-Whitney test. *(p<0.05), **(p<0.01), ***(p<0.001), ****(p<0.0001). All scale bars: 50µm.*

Figure 6: *Foxg1-Fgf* **Double LOF leads to premature oligogenesis**. (A–C) Cre electroporation at E15.5 in control (A, *Foxg1*^{lox/+}; *GFP* ^{FRT}) and *Foxg1* LOF (B, *Foxg1*^{lox/lox}; *GFP* ^{FRT}) embryos, followed by analysis at P14. GFP+ cells in control brains do not colocalise with ALDH1L1, OLIG2, and PDGFRA staining (A), whereas most GFP+ cells in *Foxg1* LOF brains display these markers (B). Co-electroporation of *Cre* together with a construct encoding secreted FGFR3c (an FGF chelator) in *Foxg1*^{lox/lox}; GFP^{FRT} causes a significant increase in the co-localization of GFP+ cells with Oligodendrocyte Precursor Cells (OPCs) markers such as OLIG2 and PDGFRA (C). In A–C, the leftmost panels in each set are low-magnification images, as indicated by the scale bar.A quantitative analysis reveals a drastic reduction of neurogenesis at the expense of gliogenesis (astrocytes + OPCs) upon loss of *Foxg1* and an additional increase in the percentage of OPCs with the additional abrogation of FGF signalling (D, D'). n= 4069 (Control), 3970 (*Foxg1* LOF), 3332 (*Foxg1* LOF + *sFGFR3c*) from N=3 brains (biologically independent replicates). *Statistical test: Two-way ANOVA. *(p<0.05), **(p<0.01), ***(p<0.001), ****(p<0.0001). All scale bars: 50µm.*

Figure 7: Schematic depicting the regulation of gliogenesis by FOXG1.

In neurogenic progenitors, FGFR3 levels are suppressed by FOXG1. As time progresses, FOXG1 levels decrease within progenitors, and FGFR3 levels increase, making progenitors more sensitive to FGF signalling. Concomitantly, postmitotic neurons secrete factors, including FGF18, which is also under FOXG1 regulation. FGF signalling drives progenitors towards astrogliogenesis. Later, when FOXG1 and FGF levels are low, the progenitor transitions to oligogenesis.

References

1. X. Qian, et al., Timing of CNS cell generation: a programmed sequence of neuron and glial cell production from isolated murine cortical stem cells. Neuron 28, 69–80 (2000).

2. Y. Hirabayashi, et al., Polycomb limits the neurogenic competence of neural precursor cells to promote astrogenic fate transition. Neuron 63, 600–613 (2009).

3. B. A. Reynolds, S. Weiss, Generation of neurons and astrocytes from isolated cells of the adult mammalian central nervous system. Science 255, 1707–1710 (1992).

4. S. Temple, Division and differentiation of isolated CNS blast cells in microculture. Nature 340, 471–473 (1989).

5. Z. Shen, et al., Distinct progenitor behavior underlying neocortical gliogenesis related to tumorigenesis. Cell Rep. 34, 108853 (2021).

6. P. Gao, et al., Deterministic progenitor behavior and unitary production of neurons in the neocortex. Cell 159, 775–788 (2014).

7. H. Naka, S. Nakamura, T. Shimazaki, H. Okano, Requirement for COUP-TFI and II in the temporal specification of neural stem cells in CNS development. Nat. Neurosci. 11, 1014–1023 (2008).

8. M. Namihira, et al., Committed neuronal precursors confer astrocytic potential on residual neural precursor cells. Dev. Cell 16, 245–255 (2009).

9. P. Kang, et al., Sox9 and NFIA coordinate a transcriptional regulatory cascade during the initiation of gliogenesis. Neuron 74, 79–94 (2012).

10. M. Nagao, T. Ogata, Y. Sawada, Y. Gotoh, Zbtb20 promotes astrocytogenesis during neocortical development. Nat. Commun. 7, 11102 (2016).

11. K. Tanigaki, et al., Notch1 and Notch3 instructively restrict bFGF-responsive multipotent neural progenitor cells to an astroglial fate. Neuron 29, 45–55 (2001).

12. Q. Cao, et al., Expression of Notch-1 receptor and its ligands Jagged-1 and Delta-1 in amoeboid microglia in postnatal rat brain and murine BV-2 cells. Glia 56, 1224–1237 (2008).

13. W. Ge, et al., Notch signaling promotes astrogliogenesis via direct CSL-mediated glial gene activation. J. Neurosci. Res. 69, 848–860 (2002).

14. Q. Cao, et al., Nuclear factor-κB/p65 responds to changes in the Notch signaling pathway in murine BV-2 cells and in amoeboid microglia in postnatal rats treated with the γ-secretase complex blocker DAPT. J. Neurosci. Res. 88, 2701–2714 (2010).

15. D. H. Rowitch, A. R. Kriegstein, Developmental genetics of vertebrate glial-cell specification. Nature 468, 214–222 (2010).

16. A. Mallamaci, Developmental control of cortico-cerebral astrogenesis. Int. J. Dev. Biol. 57, 689–706 (2013).

17. S. A. Sloan, B. A. Barres, Mechanisms of astrocyte development and their contributions to neurodevelopmental disorders. Curr. Opin. Neurobiol. 27, 75–81 (2014).

18. J. Takouda, S. Katada, K. Nakashima, Emerging mechanisms underlying astrogenesis in the developing mammalian brain. Proc Jpn Acad, Ser B, Phys Biol Sci 93, 386–398 (2017).

19. T. A. Dinh Duong, et al., FGF Signaling Directs the Cell Fate Switch from Neurons to Astrocytes in the Developing Mouse Cerebral Cortex. J. Neurosci. 39, 6081–6094 (2019).

20. X. Zhang, et al., Clonal analysis of gliogenesis in the cerebral cortex reveals stochastic expansion of glia and cell autonomous responses to egfr dosage. Cells 9 (2020).

21. K. Nakashima, M. Yanagisawa, H. Arakawa, T. Taga, Astrocyte differentiation mediated by LIF in cooperation with BMP2. FEBS Lett. 457, 43–46 (1999).

22. F. He, et al., A positive autoregulatory loop of Jak-STAT signaling controls the onset of astrogliogenesis. Nat. Neurosci. 8, 616–625 (2005).

23. C. C. Stolt, et al., Terminal differentiation of myelin-forming oligodendrocytes depends on the transcription factor Sox10. Genes Dev. 16, 165–170 (2002).

24. C. Bhattarai, P. P. Poudel, A. Ghosh, S. G. Kalthur, Comparative role of SOX10 gene in the gliogenesis of central, peripheral, and enteric nervous systems. Differentiation. 128, 13–25 (2022).

25. C. Hanashima, L. Shen, S. C. Li, E. Lai, Brain factor-1 controls the proliferation and differentiation of neocortical progenitor cells through independent mechanisms. J. Neurosci. 22, 6526–6536 (2002).

26. K. Toma, T. Kumamoto, C. Hanashima, The timing of upper-layer neurogenesis is conferred by sequential derepression and negative feedback from deep-layer neurons. J. Neurosci. 34, 13259–13276 (2014).

27. C. Hanashima, L. Suzanne, Shen Lijian, Lai, G. Fishell, Foxg1 Suppresses Early Cortical Cell Fate. science 303, 56–60 (2004).

28. T. Kumamoto, C. Hanashima, Evolutionary conservation and conversion of Foxg1 function in brain development. Dev. Growth Differ. 59, 258–269 (2017).

29. P.-S. Hou, D. Ó. hAilín, T. Vogel, C. Hanashima, Transcription and beyond: delineating FOXG1 function in cortical development and disorders. Front. Cell. Neurosci. 14, 35 (2020).

30. J. Liu, et al., FOXG1 sequentially orchestrates subtype specification of postmitotic cortical projection neurons. Sci. Adv. 8, eabh3568 (2022).

31. M. Brancaccio, C. Pivetta, M. Granzotto, C. Filippis, A. Mallamaci, Emx2 and Foxg1 inhibit gliogenesis and promote neuronogenesis. Stem Cells 28, 1206–1218 (2010).

32. C. Falcone, et al., Foxg1 antagonizes neocortical stem cell progression to astrogenesis. Cereb. Cortex 29, 4903–4918 (2019).

33. S. Frisari, et al., Multidimensional functional profiling of human neuropathogenic FOXG1 alleles in primary cultures of murine pallial precursors. Int. J. Mol. Sci. 23 (2022).

34. G. Miyoshi, G. Fishell, Dynamic FoxG1 expression coordinates the integration of multipolar pyramidal neuron precursors into the cortical plate. Neuron 74, 1045–1058 (2012).

35. H. Zong, J. S. Espinosa, H. H. Su, M. D. Muzumdar, L. Luo, Mosaic analysis with double markers in mice. Cell 121, 479–492 (2005).

36. X. Contreras, et al., A genome-wide library of MADM mice for single-cell genetic mosaic analysis. Cell Rep. 35, 109274 (2021).

37. S. Hippenmeyer, R. L. Johnson, L. Luo, Mosaic analysis with double markers reveals cell-type-specific paternal growth dominance. Cell Rep. 3, 960–967 (2013).

38. L. Zhuo, et al., hGFAP-cre transgenic mice for manipulation of glial and neuronal function in vivo. Genesis 31, 85–94 (2001).

39. Y. M. Ganat, et al., Early postnatal astroglial cells produce multilineage precursors and neural stem cells in vivo. J. Neurosci. 26, 8609–8621 (2006).

40. F. Cargnin, et al., FOXG1 Orchestrates Neocortical Organization and Cortico-Cortical Connections. Neuron 100, 1083-1096.e5 (2018).

41. J. Liu, X. Wu, H. Zhang, G. P. Pfeifer, Q. Lu, Dynamics of RNA Polymerase II Pausing and Bivalent Histone H3 Methylation during Neuronal Differentiation in Brain Development. Cell Rep. 20, 1307–1318 (2017).

42. L. Telley, et al., Temporal patterning of apical progenitors and their daughter neurons in the developing neocortex. Science 364 (2019).

43. H. Lavoie, J. Gagnon, M. Therrien, ERK signalling: a master regulator of cell behaviour, life and fate. Nat. Rev. Mol. Cell Biol. 21, 607–632 (2020).

44. E. Seuntjens, et al., Sip1 regulates sequential fate decisions by feedback signaling from postmitotic neurons to progenitors. Nat. Neurosci. 12, 1373–1380 (2009).

45. F. Barnabé-Heider, et al., Evidence that embryonic neurons regulate the onset of cortical gliogenesis via cardiotrophin-1. Neuron 48, 253–265 (2005).

46. S. G. Dastidar, P. M. Z. Landrieu, S. R. D'Mello, FoxG1 promotes the survival of postmitotic neurons. J. Neurosci. 31, 402–413 (2011).

47. Q. Hui, Z. Jin, X. Li, C. Liu, X. Wang, FGF family: from drug development to clinical application. Int. J. Mol. Sci. 19 (2018).

48. M. Hoshikawa, A. Yonamine, M. Konishi, N. Itoh, FGF-18 is a neuron-derived glial cell growth factor expressed in the rat brain during early postnatal development. Brain Res. Mol. Brain Res. 105, 60–66 (2002).

49. W. Yuan, et al., Temporally divergent regulatory mechanisms govern neuronal diversification and maturation in the mouse and marmoset neocortex. Nat. Neurosci. 25, 1049–1058 (2022).

50. C. L. Thompson, et al., A high-resolution spatiotemporal atlas of gene expression of the developing mouse brain. Neuron 83, 309–323 (2014).

51. T. Fukuchi-Shimogori, E. A. Grove, Neocortex patterning by the secreted signaling molecule FGF8. Science 294, 1071–1074 (2001).

52. D. Huilgol, J. B. Russ, S. Srivas, Z. J. Huang, The progenitor basis of cortical projection neuron diversity. Curr. Opin. Neurobiol. 81, 102726 (2023).

53. T. Mukhtar, V. Taylor, Untangling cortical complexity during development. J. Exp. Neurosci. 12, 1179069518759332 (2018).

54. R. Bansal, V. Lakhina, R. Remedios, S. Tole, Expression of FGF receptors 1, 2, 3 in the embryonic and postnatal mouse brain compared with Pdgfralpha, Olig2 and Plp/dm20: implications for oligodendrocyte development. Dev. Neurosci. 25, 83–95 (2003).

55. T. Fukuchi-Shimogori, E. A. Grove, Emx2 patterns the neocortex by regulating FGF positional signaling. Nat. Neurosci. 6, 825–831 (2003).

56. P. H. Crossley, S. Martinez, Y. Ohkubo, J. L. Rubenstein, Coordinate expression of Fgf8, Otx2, Bmp4, and Shh in the rostral prosencephalon during development of the telencephalic and optic vesicles. Neuroscience 108, 183–206 (2001).

57. S. Tole, G. Gutin, L. Bhatnagar, R. Remedios, J. M. Hébert, Development of midline cell types and commissural axon tracts requires Fgfr1 in the cerebrum. Dev. Biol. 289, 141–151 (2006).

58. E. Kon, et al., N-cadherin-regulated FGFR ubiquitination and degradation control mammalian neocortical projection neuron migration. eLife 8 (2019).

59. J. Szczurkowska, et al., Targeted in vivo genetic manipulation of the mouse or rat brain by in utero electroporation with a triple-electrode probe. Nat. Protoc. 11, 399–412 (2016).

60. A. McCaughey-Chapman, B. Connor, Cell reprogramming for oligodendrocytes: A review of protocols and their applications to disease modeling and cell-based remyelination therapies. J. Neurosci. Res. 101, 1000–1028 (2023).

61. A. Manzari-Tavakoli, et al., The Cross-Talks Among Bone Morphogenetic Protein (BMP) Signaling and Other Prominent Pathways Involved in Neural Differentiation. Front. Mol. Neurosci. 15, 827275 (2022).

62. B. El Waly, M. Macchi, M. Cayre, P. Durbec, Oligodendrogenesis in the normal and pathological central nervous system. Front. Neurosci. 8, 145 (2014).

63. M. A. Farreny, et al., FGFs are orchestra conductors of Shh-dependent oligodendroglial fate specification in the ventral spinal cord. Neural Development, 1–19 (2017).

64. A. Zeisel, et al., Molecular architecture of the mouse nervous system. Cell 174, 999-1014.e22 (2018).

65. J. C. Silbereis, S. Pochareddy, Y. Zhu, M. Li, N. Sestan, The cellular and molecular landscapes of the developing human central nervous system. Neuron 89, 248–268 (2016).

66. R. Bansal, M. Kumar, K. Murray, R. S. Morrison, S. E. Pfeiffer, Regulation of FGF receptors in the oligodendrocyte lineage. Mol. Cell. Neurosci. 7, 263–275 (1996).

67. R. Bansal, S. Magge, S. Winkler, Specific inhibitor of FGF receptor signaling: FGF-2-mediated effects on proliferation, differentiation, and MAPK activation are inhibited by PD173074 in oligodendrocyte-lineage cells. J. Neurosci. Res. 74, 486–493 (2003).

68. N. C. Hettige, C. Ernst, FOXG1 dose in brain development. Front. Pediatr. 7, 482 (2019).

69. C. Florian, N. Bahi-Buisson, T. Bienvenu, FOXG1-Related Disorders: From Clinical Description to Molecular Genetics. Mol. Syndromol. (2011) https:/doi.org/10.1159/000327329.

70. S. Goebbels, et al., Genetic targeting of principal neurons in neocortex and hippocampus of NEX-Cre mice. Genesis 44, 611–621 (2006).

71. S. Pal, et al., An Early Cortical Progenitor-Specific Mechanism Regulates Thalamocortical Innervation. J. Neurosci. 41, 6822–6835 (2021).

72. B. Langmead, S. L. Salzberg, Fast gapped-read alignment with Bowtie 2. Nat. Methods 9, 357–359 (2012).

73. A. R. Quinlan, I. M. Hall, BEDTools: a flexible suite of utilities for comparing genomic features. Bioinformatics 26, 841–842 (2010).

74. S. Heinz, et al., Simple combinations of lineage-determining transcription factors prime cis-regulatory elements required for macrophage and B cell identities. Mol. Cell 38, 576–589 (2010).

75. J. T. Robinson, et al., Integrative genomics viewer. Nat. Biotechnol. 29, 24–26 (2011).

76. D. Kim, J. M. Paggi, C. Park, C. Bennett, S. L. Salzberg, Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. Nat. Biotechnol. 37, 907–915 (2019).

77. M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol. 15, 550 (2014).

78. S. X. Ge, D. Jung, R. Yao, ShinyGO: a graphical gene-set enrichment tool for animals and plants. Bioinformatics 36, 2628–2629 (2020).