1

Histone deacetylases regulate organ-specific growth in a horned beetle

Authors:

Yonggang Hu[1,2] [*], Jordan R. Crabtree[1], Anna L. M. Macagno[1,3], Armin P. Moczek[1]

Affiliations:

[1] Department of Biology, Indiana University, 915 East 3rd Street, Bloomington, IN 47405, USA

[2] State Key Laboratory of Resource Insects, Institute of Sericulture and Systems Biology, Southwest University, Chongqing, 400715, China

[3] Biostatistics Consulting Center, Department of Epidemiology and Biostatistics, School of Public Health Bloomington, Indiana University, 2719 E. 10th Street, Bloomington, IN 47405, USA

* corresponding author details:

State Key Laboratory of Resource Insects, Institute of Sericulture and Systems Biology, Southwest University, Chongqing, 400715, China

email: yongganghu@swu.edu.cn

Short running title: HDAC function in horned beetle development

Keywords: *Onthophagus*, chromatin, plasticity, nutrition

ABSTRACT

Background

Nutrient availability is among the most widespread means by which environmental variability affects developmental outcomes. Because almost all cells within an individual organism share the same genome, structure-specific growth responses must result from changes in gene regulation. Earlier work suggested that *histone deacetylases* (*HDACs*) may serve as epigenetic regulators linking nutritional conditions to trait-specific development. Here we expand on this work by assessing the function of diverse *HDACs* in the structure-specific growth of both sex-shared and sex-specific traits including evolutionarily novel structures in the horned dung beetle *Onthophagus taurus.*

Results

We identified five *HDAC* members whose down-regulation yielded highly variable mortality depending on which *HDAC* member was targeted. We then show that *HDAC1*, *3*, and *4* operate in both a gene- and trait-specific manner in the regulation of nutrition-responsiveness of appendage size and shape. Specifically, *HDAC 1, 3,* or *4* knockdown diminished wing size similarly while leg development was differentially affected by RNAi targeting *HDAC3* and *HDAC4.* In addition, depletion of *HDAC3* transcript resulted in a more rounded shape of genitalia at the pupal stage and decreased the length of adult aedeagus across all body sizes. Most importantly, we find that *HDAC3* and *4* pattern the morphology and regulate the scaling of evolutionarily novel head and thoracic horns as a function of nutritional variation.

Conclusion

Collectively, our results suggest that both functional overlap and division of labor among *HDAC* members contribute to morphological diversification of both conventional and recently evolved appendages. More generally, our work raises the possibility that *HDAC*-mediated scaling relationships and their evolution may underpin morphological diversification within and across insect species broadly.

BACKGROUND

Phenotypic plasticity is the ability of an organism to change its phenotype in response to environmental stimuli (Schlichting and Pigliucci 1998; West-Eberhard 2003), a universal phenomenon in the living world. Diverse abiotic (e.g. temperature, photoperiod) and biotic (e.g. conspecific density) may influence growth and differentiation (Régnière et al., 2012; Prudic et al., 2011; O'Leary et al., 2017). Among those, nutrition is one of the most widespread means by which environmental variability affects developmental outcomes. Lack of essential nutrients can slow or arrest development, and sometimes trigger alternative developmental programs, such as the diapause of insects and worms (Lafuente and Beldade 2019). Nutrition also serves as a major determinant of animal size and shape (Andersen et al. 2013; Rohner et al. 2022), with poor nutrition generally yielding reduced growth and final adult body size in animals with determinate growth such as insects and mammals. However, different body parts within an individual typically differ in their response to nutrient availability. For instance, brain size in mammals and male genital size in arthropods are relatively nutrition-insensitive (Koh et al. 2005; Eberhard 2009), whereas secondary sexual traits such as male horns of dung and rhinoceros beetles (Moczek 1998; Ito et al. 2013), or mandibles of stag and broad-horned flour beetles (Gotoh et al. 2014; Okada et al. 2019) are exquisitely sensitive to nutritional variation during development. Such trait-specific scaling relationships therefore contribute in important ways to shape morphological diversity within and among taxa. Here we investigate the role of epigenetic modifications via organ-specific growth responses in an insect model.

Because all cells within an individual organism essentially share the same genome, organ- or structure-specific growth must result from changes in gene regulation. Much recent work has described changes in transcription profiles in response to environmental modifications such as nutrient availability, and has begun to identify key regulators of condition-responsive growth (e.g. *insulin*/*IIS* (Emlen et al. 2012; Snell-Rood and Moczek 2012; Xu et al. 2015),

doublesex (Kijimoto et al. 2012), and *hedgehog* (Kijimoto and Moczek 2016)). However, how

such variation in gene expression is achieved in the first place, and then subsequently

transduced into organ-specific growth is much less well understood. Here we investigate the

role of epigenetic modifications in enabling organ-specific growth, with a particular emphasis on

histone modifications. Histone acetylation/deacetylation is crucial in the organization of

euchromatin (which enables transcription) and heterochromatin (which inhibits transcription),

thereby mediating changes in gene expression. Histone deacetylases (HDACs) are members of

an ancient enzyme family that reverses the acetylation of protein substrates. *HDAC*-mediated

removal of acetylation from histone tail lysines generally correlates with gene silencing by

decreasing the ability of transcription factors to access DNA (Chen et al. 2015). In insects,

studies have confirmed that *HDACs* play a role in various developmental processes, such as

growth (Ozawa et al. 2016), metamorphosis (George et al. 2019; George and Palli 2020a;

George and Palli 2020b), long-term memory (Fitzsimons et al. 2013), reproduction (Zhang et al.

2018), longevity (Frankel et al. 2015), caste differentiation (Spannhoff et al. 2011; Suzuki et al.

2019), diapause (Cui et al. 2022), and immunity (Mukherjee et al. 2012). Furthermore, in the

broad-horned flour beetle *Gnatocerus cornutus*, *HDAC1* and *HDAC3* differentially participate in

the nutrition-dependent growth of wings and male-exaggerated mandibles, suggesting that

HDACs may serve as epigenetic regulators linking nutritional conditions to trait-specific

development (Ozawa et al. 2016). Here we expand on this work by assessing the function of

diverse histone deacetylases in the structure-specific growth of both sex-shared and sex-

specific traits including evolutionarily novel structures in a horned dung beetle.

Horned dung beetles (genus *Onthophagus*) have emerged as promising model systems

to investigate the development and diversification of scaling relationships. Here we employ one

such model, the bull-headed dung beetle *O. taurus*, to investigate the function of five different

HDAC members in the development of four different morphological structures. We selected hind

legs as examples of traits that exhibit moderate nutrition-responsiveness and therefore scale

roughly isometrically with body size. We also investigated male genitalia because of their

relatively muted nutrition response and corresponding hypoallometric scaling. Finally, we

assessed thoracic horns and head horns because of their highly sex-specific growth and scaling

relationships (Casasa et al. 2017). Thoracic horns are observed only in the pupal stage of both

sexes where they function as molting devices in the shedding of the larval head capsule during

the larval-pupal molt, and exhibit exaggerated growth in males (Moczek et al. 2006; Moczek

2006). In partial contrast, head horns are found only in male pupae as well as adults and exhibit

extreme nutrition-responsive growth resulting in hyper-allometric scaling with body size. Head

horns function as weapons in competition between adult males over reproductive access to

females. While thoracic horns have recently been identified as partial wing serial homologs (Hu

et al. 2019), head horns lack any obvious homology with other structures and are thus

considered evolutionary novelties even by the strictest of definitions (Müller and Wagner 1991;

Moczek 2005). Below we detail our results and discuss them in the light of the developmental

regulation of growth and plasticity in horned beetles in particular and insects broadly.

RESULTS

We sought to characterize the presence and function of *HDACs* during the development of the

bull-headed dung beetle *O. taurus*. We identified five HDACs in the annotated genome, which,

based on phylogenetic analysis, could be classified into three classes: class I (HDAC1 and

HDAC3), class II (HDAC4 and HDAC6), and class IV (HDAC11) (Fig. S1). It is worth noting that

a HDAC protein that was predicted as HDAC5 for *O. taurus* in NCBI (reference number:

XP_022905538.1) was found to be nested within the cluster of HDAC4 proteins across insect

species (Fig. S1). As a result, we have re-annotated this protein as Ot-HDAC4. We then

performed RNAi experiments by injecting double-stranded RNA (dsRNA) corresponding to each

of the five HDACs into newly molted final-instar *O. taurus* larvae and assessed their influence

on pupal and adult morphologies. Wildtype morphology for each focal phenotype is shown in

Figure 1.

HDAC-RNAi resulted in highly variable mortality depending on the target *HDAC*

RNAi-mediated knockdown of *HDAC1* resulted in 100% larval mortality at the initial 1 µg/µl

dsRNA injection dosage, as well as subsequent dosages as low as 0.25 µg/µl (Table S1). Most

larvae exhibited arrested development at the pre-pupal stages and eventually died with larval-

pupal intermediate phenotypes with pupal traits, such as compound eyes, observed underneath

the larval integument. A mass of fat body accumulated underneath the posterior region of the

developing pupal abdomen, resulting in a cavity filled with hemolymph between the larval

integument and newly-formed pupal cuticle (Fig. 2A and B). When dsRNA concentration was

decreased to 0.01 µg/µl, very few individuals succeeded to develop to pupal (3/44) and adult

(2/44) stages (Table S1) amenable to phenotyping. In marked contrast, RNAi targeting *HDAC 3*,

4, *6*, or *11* resulted in mortalities ranging from 16.7% to 86.7% depending on dosage and

permitted more nuanced and quantitative analysis of phenotypic effects (Table S1). Because no

observable phenotypes were found following *HDAC6*[RNAi] or *HDAC11*[RNAi] even at high dsRNA

dosage, we focus on *HDAC 1*, *3*, and *4* in the remainder.

HDACs function during appendage development

To determine affected traits, we measured the scaling relationship of each trait to the cubic root

of pupal mass as a proxy of body size since the more commonly used measure of thorax width

was affected by *HDAC3* knockdown (treatment: P < 0.001, Table S2). *HDAC1*[RNAi] resulted in

curtailment of both forewings (i.e. elytra) and hindwings at the pupal stage, which was retained

into the adult (Fig. 2C). Similarly, both *HDAC3* or *HDAC4* knockdown diminished wing size (Fig.

2C). Thus, *HDAC 1, 3,* and *4* appear to regulate wing development in similar ways. RNAi

targeting *HDAC3* and *HDAC4* also affected leg development, but specific effects diverged. To

assess leg phenotypes quantitatively we selected the femur, which is especially amenable to

width and length measurement, for morphometric analyses. $HDAC4^{RNAi}$ led to a reduction in

femur length (treatment: $P < 0.001$, Table S2) while the slope of the body size/femur length

allometry was decreased in $HDAC3^{RNAi}$ animals (treatment: $P = 0.163$; treatment × body size: P

$= 0.004$, Fig. 2D and G, and Table S2). In contrast, whereas $HDAC3^{RNAi}$ increased femur width

compared to control individuals (treatment: $P < 0.001$, Table S2), the same measure was

reduced in *HDAC4* knockdown animals (treatment: $P = 0.001$, Fig. 2D and G, and Table S2).

Lastly, we found that knockdown of *HDAC3* also affected the development of the male

reproductive organ, the aedeagus, itself composed of the more proximal phallobase and the

more distal parameres. Specifically, genital width increased and attained a more rounded shape

at the pupal stage (treatment: $P < 0.001$, Fig. E and G, and Table S2), while adult genitalia

exhibited a deformation and overall shortening of both parameres and phallobase, which

combined yielded a shortening of overall aedeagus length across all body sizes (treatment: $P <$

0.001, Fig. F and G, and Table S2).

***HDACs* function during thoracic and head horn formation**

Head and thoracic horns are textbook examples of evolutionary novelties, and we sought to

determine whether *HDAC* function may have been co-opted during the evolution of one or both

horn types. $HDAC1^{RNAi}$ resulted in a reduction in thoracic horn length and a split tip at the pupal

stage (Fig. 3A and C), whereas the corresponding area in the adult exhibited a broad

indentation (compared to the smoothly convex outline observed in wildtype or buffer control

injected individuals) and small bilateral projections at the respective edge of the indentation (Fig.

3F and G). Effects on head horns could not be quantified with certainty due to the high degree

of natural variability of the trait and the very low number of surviving males, which in addition

were too small to develop fully formed head horns. However, *HDAC3* knockdown caused

measurable shape and scaling changes in thoracic horns. Specifically, $HDAC3^{RNAi}$ increased

pupal thoracic horn width (treatment: $P < 0.001$, Fig. 3D, E, L, and Table S3), but decreased

pupal thoracic horn length (treatment: $P < 0.001$, Fig. 3L and Table S3). In addition, we found a

bilateral indentation to the distal region of the thoracic horn, causing the thoracic horns of large

$HDAC3^{RNAi}$ pupae to attain a more conical shape (Fig. 3B). In contrast to thoracic horn

phenotypes, $HDAC3^{RNAi}$ yielded drastically enlarged head horns, in particular concerning head

horn *width* across the entire range of male body sizes (Fig. 3E, H-K, and Fig. S2). However, due

to the highly varied nature of these phenotypes, we were unable to arrive at reliable landmarks

for quantitative measure, hence this observation could only be made qualitatively. To determine

whether head horn *length* was also affected, we further measured the scaling relationship of

head horn length to body size. $HDAC3^{RNAi}$ steepened the slope of the head horn length

allometry at both pupal (treatment: $P = 0.025$, Fig. 3L and Table S3) and adult stages (treatment:

$P = 0.012$, Fig. 3L and Table S3). Notably, $HDAC3^{RNAi}$ also reduced the maximum asymptotic

horn length in adults (treatment: $P < 0.001$, Fig. 3L and Table S3), which was not detected at

the pupal stage (treatment: $P = 0.107$, Fig. 3L and Table S3). The quantification of head horn

length reduction from pupa to adult further confirmed this observation (treatment: $P < 0.05$,

Table S2), and this effect was enhanced with increasing body size (treatment × body size effect:

$P = 0.010$, Fig. 3L and Table S2). These results suggest that $HDAC3^{RNAi}$ affects head horn *width*

during the horn growth phase (which takes place during the larval–to–pupal transition, thereby

resulting in visible *pupal* phenotypes), but affects horn *length* during the pupal remodeling phase

of ontogeny (thus becoming apparent in adults only). $HDAC4^{RNAi}$ in turn not only decreased

maximum asymptotic horn length in adults (treatment: $P = 0.027$, Table S3), but also altered the

threshold body size of sigmoidal allometry separating small hornless from large and fully horned

males, such that large head horns now also presented in males of relatively small body sizes

(treatment: $P < 0.001$, Fig. 3L and Table S3). In contrast, we did not find abnormal phenotypes

with respect to thoracic horns in $HDAC4^{RNAi}$ individuals.

DISCUSSION

The significance of *HDACs* in horned beetle development and evolution

Earlier work on the broad-horned flour beetle *Gnatoceros cornutus* was the first to document the role of *HDACs* in the regulation of nutrition-responsive plasticity in insects (Ozawa et al. 2016). Large males in this species develop conspicuous mandibular projections (called mandibular horns) which were reduced following *HDAC1*[RNAi], whereas *HDAC3*[RNAi] led to hypertrophy. Opposite effects were observed with respect to wing size, yet none in genitalia (Ozawa et al. 2016). These results were the first to suggest that *HDACs* operate in a trait-specific manner, and in particular contribute to the plastic and sex-specific expression of exaggerated male mandibles. Our results presented here further support a role of *HDACs* in the regulation of trait-specific plasticity, as well as add important new aspects to our understanding of HDAC function in insect development.

First, similar to the horns of rhinoceros beetle *Trypoxylus dichotomus*, *Onthophagus* horns frequently exhibit pronounced nutrition-dependent plasticity, in contrast to the more isometric growth typical of wings and legs, or the hypo-allometric growth of genitalia (Emlen et al. 2012; Casasa et al. 2017). Several mechanisms have been proposed as possibly underlying module-specific conditional growth (e.g. insulin/insulin-like growth factor (Emlen et al. 2012; Okada et al. 2019), FOXO (Tang et al. 2011; Snell-Rood and Moczek 2012; Casasa and Moczek 2018), HDAC (Ozawa et al. 2016)). Ozawa *et al.* (2016), in particular, proposed a mechanistic explanation of epigenetic flexibility in which developmentally plastic organs (e.g. mandibular horns in *G. cornutus*) are more susceptible to epigenetic (i.e. *HDAC*) perturbation, whereas developmentally robust organs (e.g. genitalia) are non-responsive to *HDAC* perturbation. However, results presented here are at odds with this model. Specifically, even though head and thoracic horn development in *O. taurus* exhibit exaggerated nutritional plasticity, the effects of *HDAC3*[RNAi] were considerably more pronounced in genitalia and wings.

Genitalia in particular exhibited a considerable reduction of size relative to body size across all

body sizes following *HDAC3*[RNAi], thereby highlighting a previously unexpected role of *HDAC* in

regulating the development of traits generally assumed to be robust to nutritional variation.

Intriguingly, a similar outcome was observed following *insulin receptor* (*InR1/2*) transcript

depletion in *O. taurus* (Casasa and Moczek 2018). Hence, the epigenetic flexibility hypothesis

proposed in *Gnatoceros* beetles is unlikely to explain the findings seen here in *Onthophagus*,

consistent with divergences in *HDAC* function across the Coleoptera, again similar to what has

recently been reported for the insulin signaling pathway (Rohner et al. 2023). This in turn raises

the possibility that, in addition to its primary function in regulating epigenetic status, *HDAC3* may

also function in aspects of trait morphogenesis not related to nutritional status and

developmental plasticity.

Second, we found that *HDAC1*[RNAi] induced developmental arrest at the prepupal stages

in line with previous studies in *Tribolium castaneum* (George et al. 2019), which suggests that

HDAC1 expression is required for suppressing the juvenile hormone (JH) response gene

Krüppel homolog 1 (*Kr-h1*). In *Tribolium*, *HDAC1* knockdown prevents larval-to-pupal transition

via derepressing *Kr-h1* expression, thereby influencing JH actions and thus halting

metamorphosis. Similarly, high dose of *HDAC1* knockdown caused developmental failures

during the pupal stage in *Gnatoceros*, indicating a possibly conserved role in basic

developmental process mediating metamorphosis. However, despite this putative conservation

of *HDAC1* function across the Coleoptera assessed to date, we also found that the precise

nature and direction of *HDAC*[RNAi] effects on appendage formation diverged between *Gnatoceros*

and *Onthophagus* beetles even beyond those already noted above: for example, in

Onthophagus downregulation of *HDAC1*, *3*, and *4* all appear to affect wing size similarly,

whereas in *Gnatoceros* *HDAC1*[RNAi] and *HDAC3*[RNAi] yield opposite effects. Likewise, in

Onthophagus, *HDAC3*[RNAi] increased femur width, but *HDAC4*[RNAi] decreased it, whereas leg

morphology was generally unaffected in *Gnatoceros* beetle. Lastly, our results document the

recruitment of *HDAC* function into the formation of an evolutionarily novel structure - head and

thoracic horns - suggesting that *HDAC* function is not just evolutionarily labile among conserved

insect traits but also contributed to the comparatively recent evolution of *Onthophagus*

weaponry, including the regulation of size, shape, and key components of scaling.

Development and evolution of pupal remodeling

The horns of adult beetles are the product of developmental processes operating at at least two

distinct stages of development, a rapid growth phase approximately 48h immediately prior to the

larval-to-pupal molt and a remodeling phase during the pupal stage (Moczek 2006). While

generally given less attention, pupal remodeling can be quite extensive and fully formed pupal

horns may be subject to considerable reduction and even complete resorption in many species.

Thus, the morphological diversity of adult horns is not only influenced by the differential

regulation of growth during the prepupal stage, but also by the developmental processes

underlying the differential resorption of horn tissue during the pupal stage (Moczek 2006;

Moczek 2007; Morita et al. 2019). Previous work identified that differential programmed cell

death facilitates species, sex, and body-region specific resorption of horn primordia (Kijimoto et

al. 2010). However, the mechanisms regulating horn resorption during the pupal stage remain

largely unknown. Our results implicate *HDAC3* as a regulator of both prepupal growth and pupal

remodeling of horn primordia. Specifically, we show that $HDAC3^{RNAi}$ increased head horn width,

that this phenotype was already prominently visible at the pupal stage, and must therefore have

resulted from modifications to the prepupal growth phase of horn formation (Fig. 3). In addition,

however, we also find that $HDAC3^{RNAi}$ altered horn *length* in a manner not evident at the pupal

stage but clearly discernible in the resulting adults, and thus a consequence of $HDAC3^{RNAi}$

effects on the pupal remodeling phase of horn formation. As such, *HDAC3* is one of relatively

few genes identified to date to be involved in horn remodeling during the pupal stage of horned

beetles (Morita et al. 2019; Matsuda et al. 2023).

Chromatin modifications and developmental plasticity

This work is the first to implicate chromatin modifications in the regulation of development and

plasticity in horned dung beetles. While earlier work documented the existence of the complete

methylation machinery in the *O. taurus* genome alongside sex- and nutrition-dependent

differences in methylation signatures, the functional significance of chromatin modifications, if

any, had remained unknown (Choi et al. 2010; Snell-Rood and Moczek 2013). We now show

that downregulation of *HDAC3* and *HDAC4* affect critical aspects of horn formation including

size, shape, and the location of the inflection point separating alternate male morphs. Future

work will need to explore if and how *HDAC* functions may be contributing to the genome-wide

remodeling patterns, and more generally the roles of *cis*-regulatory elements in the development

and evolution of plasticity in insects.

METHODS

Insects

Adult *O. taurus* were collected, courtesy of John Allen, from Paterson Farm near Ravenswood,

Western Australia. A laboratory population was maintained at 25°C in a sand/soil mixture and

fed cow manure twice a week. Larvae used for injection were collected and prepared as

described previously (Shafiei et al. 2001).

Identification of *O. taurus* orthologs for HDACs

The *Onthophagus* orthologs of *HDAC* genes were identified via reciprocal BLAST to *T.*

castaneum, *G. cornutus*, *Drosophila melanogaster*, and *Homo sapiens* in NCBI databases.

Amino acid sequences of *HDAC* genes from the above species were aligned with ClustalW

algorithm implemented in MEGA X (Kumar et al. 2018). Neighbor-Joining trees were

constructed using MEGA X.

Double-stranded RNA (dsRNA) synthesis and injection

Total RNA was extracted with RNeasy Mini Plus Kit (QIAGEN) and reverse transcribed with

iScript cDNA Synthesis kit (Bio-Rad). Partial fragments of each genes were amplified with PCR

by using gene-specific primers and cloned into pCR4-TOPO TA vector (Invitrogen, Thermo

Fisher Scientific). DNA templates for *in vitro* transcription were produced with PCR by using

TOPO RNAi primer (Supplementary table S4) (Philip and Tomoyasu 2011). PCR products were

purified and concentrated using the QIAquick PCR Purification kit (QIAGEN) and subjected to *in*

vitro transcription (MEGAscript T7 Transcription Kit, Thermo Fisher Scientific) and dsRNA

purification (MEGAclear Transcription Clean-Up Kit, Thermo Fisher Scientific), according to the

manufacturer's protocol. DsRNA was quantified and stored at -80°C until use. Injection was

carried out at the early stage of the last larval instar (i.e. the third larval instar, L3). Control

animals were injected with injection buffer (1.4 mM NaCl, 0.07mM Na_2HPO_4, 0.03mM KH_2PO_4,

and 4mM KCl) kept at the same condition as dsRNA injected animals (See Supplementary table

S1 for detailed injection information).

Effect of *HDAC3*- and *HDAC4*-RNAi on the scaling relationships between several

morphological traits and body size

Since the usual measure of *Onthophagus* body size - thorax width (e.g. Casasa and Moczek

2018; Macagno et al. 2021) - was clearly affected by our RNAi treatments (Fig. 3B and Table

S2), we measured the cube root of pupal mass as a proxy for individual body size (Marchini et

al. 2014; Fleming et al. 2021). We used t-tests to compare body sizes between each RNAi-

treated and control group.

Consistent with previous studies we analyzed nonlinear horn allometries using

untransformed data (e.g. Casasa and Moczek 2018; Macagno et al. 2021). We used the

package drc (Ritz et al. 2015) in R 3.5.2 (R Core Team 2018) to fit the body size/head horn

length distribution a four-parameter log-logistic (Hill) function in the form

$$y = c + \frac{d-c}{1+\exp(b\,(\log(x) - \log\,(a)))},$$

with x = body size, y = head horn length, a = body size at the point of inflection of the sigmoid

curve, b = slope of the curve, c = minimum and d = maximum asymptotic horn lengths

(Macagno et al. 2021). We then inferred whether a complex model including a sigmoidal

regression per treatment (i.e. control and RNAi treatments) fitted our data better than a simpler

model with one sigmoidal regression including the whole sample by means of the Akaike

Information Criterion (AIC) (Akaike 1974). The AIC measures relative model fit - the lower its

value, the better the model fits to the experimental data (Burnham et al. 2011). Upon finding the

complex model more fitting, we used Welch's t-tests (with Holm-Bonferroni sequential correction

where applicable) to compare parameter means (a, b, c, d) between control-injected and RNAi

treatment groups (Moczek et al. 2002; Casasa and Moczek 2018). We compared control

individuals to $HDAC3^{RNAi}$ and $HDAC4^{RNAi}$ individuals in the case of the adult horn allometry. As

for pupal horn allometry, we compared $HDAC3^{RNAi}$ to control individuals.

To inspect the effect of RNAi manipulations on all other morphological traits, we

modeled trait size as a function of body size, treatment ($HDAC3^{RNAi}$ or $HDAC4^{RNAi}$ vs control-

injected) and their interaction using general linear models in SPSS Statistics 25 (IBM Corp.,

2017). Models were simplified by removing non-significant interactions. Data were log-

transformed prior to analyses (Huxley 1924). The reduction in horn size during metamorphosis

(defined as pupal horn length - adult horn length) was analyzed similarly with a general linear

model, using untransformed data as in other analyses of horn morphology. Statistics for

morphometric analyses are provided in Supplementary Table S2 and S3.

Image Processing

All images were captured with a digital camera (Scion) mounted to a dissecting microscope

(Leica MZ16, Germany). Brightness and contrast of images were adjusted across the entire

image with Adobe Photoshop CC 2017 (Adobe, USA).

DECLARATIONS

Ethical Approval

Not applicable.

Funding

Support for this study was provided by awards from the National Science Foundation

(IOS1256689) and John Templeton Foundation (61369) to APM. Additional support was

provided by the Natural Science Foundation of Chongqing for Distinguished Young Scholars

award to YH (CSTB2022NSCQ-JQX0025). The opinions, interpretations, conclusions and

recommendations are those of the authors and are not necessarily endorsed by the National

Science Foundation, the John Templeton Foundation, or the Natural Science Foundation of

Chongqing.

Availability of data and materials

All data are available in the manuscript and supplementary materials.

Authors' contributions

Y.H. conceived the initial idea, Y.H. and A.P.M. designed the research, Y.H. and J.R.C.

performed the experiments, Y.H. and A.L.M.M. analyzed the data, Y.H., A.P.M., and A.L.M.M.

wrote the manuscript with contribution from J.R.C.

Competing interests

395 All authors declare no competing interests.

Acknowledgements

398 We would like to thank Kayla Copper and Levi Burdine for help with beetle care, and Patrick

399 Rohner for discussion of statistical analyses.

References

Akaike H. 1974. A new look at the statistical model identification. *IEEE Trans. Autom. Control* 19:716–

723.

Andersen DS, Colombani J, Léopold P. 2013. Coordination of organ growth: principles and outstanding

questions from the world of insects. *Trends Cell Biol.* 23:336–344.

Burnham KP, Anderson DR, Huyvaert KP. 2011. AIC model selection and multimodel inference in

behavioral ecology: some background, observations, and comparisons. *Behav. Ecol. Sociobiol.*

65:23–35.

Casasa S, Moczek AP. 2018. Insulin signalling's role in mediating tissue-specific nutritional plasticity and

robustness in the horn-polyphenic beetle Onthophagus taurus. *Proc. R. Soc. B Biol. Sci.*

285:20181631.

Casasa S, Schwab DB, Moczek AP. 2017. Developmental regulation and evolution of scaling: novel

insights through the study of Onthophagus beetles. *Curr. Opin. Insect Sci.* 19:52–60.

Chen HP, Zhao YT, Zhao TC. 2015. Histone deacetylases and mechanisms of regulation of gene

expression. *Crit. Rev. Oncog.* 20:35–47.

Choi J-H, Kijimoto T, Snell-Rood E, Tae H, Yang Y, Moczek AP, Andrews J. 2010. Gene discovery in the

horned beetle Onthophagus taurus. *BMC Genomics* 11:703.

Cui J, Lin K, Xu L, Yue F, Yu L, Zhang Q. 2022. Transcriptome Analysis of Beet Webworm Shows That

Histone Deacetylase May Affect Diapause by Regulating Juvenile Hormone. *Insects* 13:835.

Eberhard WG. 2009. Static Allometry and Animal Genitalia. *Evolution* 63:48–66.

Emlen DJ, Warren IA, Johns A, Dworkin I, Lavine LC. 2012. A mechanism of extreme growth and reliable

signaling in sexually selected ornaments and weapons. *Science* 337:860–864.

Fitzsimons HL, Schwartz S, Given FM, Scott MJ. 2013. The Histone Deacetylase HDAC4 Regulates Long-

Term Memory in Drosophila. *PLOS ONE* 8:e83903.

Fleming MJ, Carter AW, Sheldon KS. 2021. Dung beetles show metabolic plasticity as pupae and smaller

adult body size in response to increased temperature mean and variance. *J. Insect Physiol.*

131:104215.

Frankel S, Woods J, Ziafazeli T, Rogina B. 2015. RPD3 histone deacetylase and nutrition have distinct but

interacting effects on Drosophila longevity. *Aging* 7:1112–1129.

George S, Gaddelapati SC, Palli SR. 2019. Histone deacetylase 1 suppresses Krüppel homolog 1 gene
expression and influences juvenile hormone action in Tribolium castaneum. *Proc. Natl. Acad. Sci.*
116:17759–17764.

George S, Palli SR. 2020a. Histone deacetylase 3 is required for development and metamorphosis in the
red flour beetle, Tribolium castaneum. *BMC Genomics* 21:420.

George S, Palli SR. 2020b. Histone Deacetylase 11 Knockdown Blocks Larval Development and
Metamorphosis in the Red Flour Beetle, Tribolium castaneum. *Front. Genet.* 11:683.

Gotoh H, Miyakawa H, Ishikawa A, Ishikawa Y, Sugime Y, Emlen DJ, Lavine LC, Miura T. 2014.
Developmental Link between Sex and Nutrition; doublesex Regulates Sex-Specific Mandible
Growth via Juvenile Hormone Signaling in Stag Beetles. *PLOS Genet.* 10:e1004098.

Hu Y, Linz DM, Moczek AP. 2019. Beetle horns evolved from wing serial homologs. *Science* 366:1004–
1007.

Huxley JS. 1924. Constant Differential Growth-Ratios and their Significance. *Nature* 114:895–896.

Ito Y, Harigai A, Nakata M, Hosoya T, Araya K, Oba Y, Ito A, Ohde T, Yaginuma T, Niimi T. 2013. The role
of doublesex in the evolution of exaggerated horns in the Japanese rhinoceros beetle. *EMBO*
Rep. 14:561–567.

Kijimoto T, Andrews J, Moczek AP. 2010. Programed cell death shapes the expression of horns within
and between species of horned beetles. *Evol. Dev.* 12:449–458.

Kijimoto T, Moczek AP. 2016. Hedgehog signaling enables nutrition-responsive inhibition of an
alternative morph in a polyphenic beetle. *Proc. Natl. Acad. Sci.* 113:5982–5987.

Kijimoto T, Moczek AP, Andrews J. 2012. Diversification of doublesex function underlies morph-, sex-,

and species-specific development of beetle horns. *Proc. Natl. Acad. Sci.* 109:20526–20531.

Koh I, Lee MS, Lee NJ, Park KW, Kim KH, Kim H, Rhyu IJ. 2005. Body size effect on brain volume in Korean

youth. *Neuroreport* 16:2029–2032.

Kumar S, Stecher G, Li M, Knyaz C, Tamura K. 2018. MEGA X: Molecular Evolutionary Genetics Analysis

across Computing Platforms. *Mol. Biol. Evol.* 35:1547–1549.

Lafuente E, Beldade P. 2019. Genomics of Developmental Plasticity in Animals. *Front. Genet.* 10:720.

Macagno ALM, Edgerton TJ, Moczek AP. 2021. Incipient hybrid inferiority between recently introduced,

diverging dung beetle populations. *Biol. J. Linn. Soc.* 132:931–944.

Marchini M, Sparrow LM, Cosman MN, Dowhanik A, Krueger CB, Hallgrimsson B, Rolian C. 2014. Impacts

of genetic correlation on the independent evolution of body mass and skeletal size in mammals.

BMC Evol. Biol. 14:258.

Matsuda K, Adachi H, Gotoh H, Inoue Y, Kondo S. 2023. Mechanical models affecting beetle horn

remodeling. :2023.01.16.524208. Available from:

https://www.biorxiv.org/content/10.1101/2023.01.16.524208v1

Moczek A, Hunt J, Emlen D, Simmons L. 2002. Threshold Evolution in Exotic Populations of a Polyphenic

Beetle. *Evol. Ecol. Res.* 4:587–601.

Moczek AP. 1998. Horn polyphenism in the beetle Onthophagus taurus: larval diet quality and plasticity

in parental investment determine adult body size and male horn morphology. *Behav. Ecol.*

9:636–641.

Moczek AP. 2005. The Evolution and Development of Novel Traits, or How Beetles Got Their Horns.

BioScience 55:937–951.

Moczek AP. 2006. Pupal Remodeling and the Development and Evolution of Sexual Dimorphism in

Horned Beetles. *Am. Nat.* 168:711–729.

Moczek AP. 2007. Pupal remodeling and the evolution and development of alternative male

morphologies in horned beetles. *BMC Evol. Biol.* 7:151.

Moczek AP, Cruickshank TE, Shelby A. 2006. When Ontogeny Reveals What Phylogeny Hides: Gain and

Loss of Horns During Development and Evolution of Horned Beetles. *Evolution* 60:2329–2341.

Morita S, Ando T, Maeno A, Mizutani T, Mase M, Shigenobu S, Niimi T. 2019. Precise staging of beetle

horn formation in Trypoxylus dichotomus reveals the pleiotropic roles of doublesex depending

on the spatiotemporal developmental contexts. *PLOS Genet.* 15:e1008063.

Mukherjee K, Fischer R, Vilcinskas A. 2012. Histone acetylation mediates epigenetic regulation of

transcriptional reprogramming in insects during metamorphosis, wounding and infection. *Front.*

Zool. 9:25.

Müller GB, Wagner GP. 1991. Novelty in Evolution: Restructuring the Concept. *Annu. Rev. Ecol. Syst.*

22:229–256.

Okada Y, Katsuki M, Okamoto N, Fujioka H, Okada K. 2019. A specific type of insulin-like peptide

regulates the conditional growth of a beetle weapon. *PLOS Biol.* 17:e3000541.

Ozawa T, Mizuhara T, Arata M, Shimada M, Niimi T, Okada K, Okada Y, Ohta K. 2016. Histone

deacetylases control module-specific phenotypic plasticity in beetle weapons. *Proc. Natl. Acad.*

Sci. U. S. A. 113:15042–15047.

Philip BN, Tomoyasu Y. 2011. Gene Knockdown Analysis by Double-Stranded RNA Injection. In: Orgogozo

V, Rockman MV, editors. Molecular Methods for Evolutionary Genetics. Methods in Molecular

Biology. Totowa, NJ: Humana Press. p. 471–497. Available from: https://doi.org/10.1007/978-1-

61779-228-1_28

R Core Team. 2018. R: A language and environment for statistical computing. Available from:

https://www.R-project.org

Ritz C, Baty F, Streibig JC, Gerhard D. 2015. Dose-Response Analysis Using R. *PloS One* 10:e0146021.

Rohner PT, Casasa S, Moczek AP. 2023. Assessing the evolutionary lability of insulin signalling in the

regulation of nutritional plasticity across traits and species of horned dung beetles. *J. Evol. Biol.*

36:1641–1648.

Rohner PT, Hu Y, Moczek AP. 2022. Developmental bias in the evolution and plasticity of beetle horn

shape. *Proc. R. Soc. B Biol. Sci.* 289:20221441.

Schlichting C, Pigliucci M. 1998. Phenotypic Evolution: A Reaction Norm Perspective. Sinauer

Shafiei M, Moczek AP, Nijhout HF. 2001. Food availability controls the onset of metamorphosis in the

dung beetle Onthophagus taurus (Coleoptera: Scarabaeidae). *Physiol. Entomol.* 26:173–180.

Snell-Rood EC, Moczek AP. 2012. Insulin Signaling as a Mechanism Underlying Developmental Plasticity:

The Role of FOXO in a Nutritional Polyphenism. *PLOS ONE* 7:e34857.

Snell-Rood EC, Moczek AP. 2013. Horns and the role of development in the evolution of beetle contests.

In: Hardy ICW, Briffa M, editors. Animal Contests. Cambridge: Cambridge University Press. p.

178–198. Available from: https://www.cambridge.org/core/books/animal-contests/horns-and-

the-role-of-development-in-the-evolution-of-beetle-

contests/9384532AAA46CD20CF0846294A02DCD4

Spannhoff A, Kim YK, Raynal NJ-M, Gharibyan V, Su M-B, Zhou Y-Y, Li J, Castellano S, Sbardella G, Issa J-

PJ, et al. 2011. Histone deacetylase inhibitor activity in royal jelly might facilitate caste switching

in bees. *EMBO Rep.* 12:238–243.

Suzuki R, Yaguchi H, Maekawa K. 2019. Histone modifying genes are involved in the molting period

during soldier differentiation in Zootermopsis nevadensis. *J. Insect Physiol.* 117:103892.

West-Eberhard MJ. 2003. Developmental Plasticity and Evolution. Oxford University Press

Xu H-J, Xue J, Lu B, Zhang X-C, Zhuo J-C, He S-F, Ma X-F, Jiang Y-Q, Fan H-W, Xu J-Y, et al. 2015. Two

insulin receptors determine alternative wing morphs in planthoppers. *Nature* 519:464–467.

Zhang J-L, Yuan X-B, Chen S-J, Chen H-H, Xu N, Xue W-H, Fu S-J, Zhang C-X, Xu H-J. 2018. The histone

deacetylase NlHDAC1 regulates both female and male fertility in the brown planthopper,

Nilaparvata lugens. *Open Biol.* 8:180158.

IBM Corp (2017). IBM SPSS Statistics for Windows, Version 25.0. Armonk, NY: IBM Corp.

Fig. 1. Wildtype morphology and morphometric landmarks used for morphological

quantification. (**A-F**) Morphology of the male pupal head (A), male adult head (B), pupal

pronotum from dorsal view (C), pupal reproductive organ (D), adult aedeagus (E), and hind leg

(F), respectively, and the morphometric landmarks used for measurements (purple and yellow

line). TW, thoracic width; TL, thoracic length; GW, genital width; GL, genital length; FL, femur

length; FW, femur width.

Fig. 2. *HDAC^RNAi* **effects on molting and appendage development.**

(**A** and **B**) Larval-pupal intermediate induced by *HDAC1* knockdown. The fat body

accumulated outside of newly-formed pupa is outlined (red dotted line) before (A) and

after (B) peeling away the larval cuticle, respectively. Hemolymph in the cavity between

larval and newly-formed pupal cuticle is indicated (red arrow in A). (**C**) Representative

wing phenotypes are shown as follows: buffer injection, $HDAC1^{RNAi}$, $HDAC3^{RNAi}$, and

$HDAC4^{RNAi}$. (**D-F**) Morphology of the hind leg (D), pupal genitalia (E), and adult

aedeagus (F) compared to buffer injected (left column) and $HDAC3^{RNAi}$ (right column)

individuals, respectively. (**G**) Knockdown of *HDAC3* (orange dots) or *HDAC4* (blue dots)

on femur length and width compared to buffer-injected control (gray dots). Effects of

$HDAC3^{RNAi}$ on the male pupal and adult reproductive organ (bottom row in **G**). The

RNAi phenotypes and their corresponding negative controls are shown at the same

magnification. Scale bars: 1 mm.

Fig. 3. $HDAC^{RNAi}$ effects on thoracic and head horn formation.

(**A-C**) The pronotum in buffer injected (A), *HDAC3*^{RNAi} (B), and *HDAC1*^{RNAi} (C)

individuals. Inset in C shows the prothoracic horn. The furrow between paired horn

vestiges is indicated by red arrowheads. (**D** and **E**) Representative head horn of

negative control (D) and *HDAC3*^{RNAi} (E) individuals, respectively. The right head horn is

colored magenta. (**F-K**) The morphology of the adult pronotum in front view (F and G),

front view of head horns (H and I), as well as head horn viewed from lateral (J and K) in

negative control (third row) individuals and following *HDAC*^{RNAi} (bottom row),

respectively. (**L**) Changes in thoracic and head horns resulting from *HDAC3*^{RNAi} and

HDAC4^{RNAi}. Reduction in head horn length during the transition from pupa to adult is

increased by *HDAC3*^{RNAi} compared to buffer-injections (measured as the difference

between pupal head horn length and adult head horn length in the same individual).

RNAi phenotypes and their corresponding negative controls are shown at the same

magnification. Scale bars: 1 mm.