A chromosome-level, haplotype-resolved genome assembly and annotation for the Eurasian minnow (Leuciscidae - *Phoxinus phoxinus*) provide evidence of haplotype diversity

Oriowo Temitope Opeyemi[1#], Chrysostomakis Ioannis[1#], Martin Sebastian[1], Kukowka Sandra[1], Brown Thomas[2], Winkler Sylke[2], Myers Eugene W[2,3], Böhne Astrid[1], Stange Madlen*[1]

[1] Leibniz Institute for the Analysis of Biodiversity Change – Bonn, Germany

[2] Max Planck Institute of Molecular Cell Biology and Genetics, Pfotenhauerstr. 108, 01307 Dresden, Germany

[3] Okinawa Institute of Science and Technology, Okinawa, Japan

shared first authorship

*Corresponding author

Correspondence: m.stange@leibniz-lib.de

ABSTRACT

In this study we present an in-depth analysis of the Eurasian Minnow (*Phoxinus phoxinus*) genome, highlighting its genetic diversity, structural variations, and evolutionary adaptations. We generated an annotated haplotype-phased, chromosome-level genome assembly (2n = 25) by integrating high-fidelity (HiFi) long reads and chromosome conformation capture data (Hi-C). We achieved a haploid length of 940 Mbp for haplome one and 929 Mbp for haplome two with high N50 values of 36.4 Mb and 36.6 Mb and BUSCO scores of 96.9% and 97.2%, indicating a highly complete genome.

We detected notable heterozygosity (1.43%) and a high repeat content (approximately 54%), primarily consisting of DNA transposons, which contribute to genome rearrangements and variations. We found substantial structural variations within the genome, including insertions, deletions, inversions, and translocations. These variations affect genes enriched in functions such as dephosphorylation, developmental pigmentation, phagocytosis, immunity, and stress response.

Protein annotation identified 30,980 mRNAs and 23,497 protein-coding genes with a high completeness score, providing further support for our genome's high contiguity. We performed a gene family evolution analysis by comparing our proteome to ten other teleost species, which identified immune system gene families that prioritise histone-based disease prevention over NLR-based immune responses.

Additionally, demographic analysis indicates historical fluctuations in the effective population size of *P. phoxinus*, likely correlating with past climatic changes.

This annotated, phased reference genome provides a crucial resource for resolving the taxonomic complexity within the genus *Phoxinus* and highlights the importance of haplotype-phased assemblies in understanding haplotype diversity in species characterised by high heterozygosity.

Keywords: *Phoxinus, Haplotype-phased genome assembly, comparative genomics, structural variants, immune system, annotation, transposable elements*

Introduction

Species of the genus *Phoxinus* show considerable morphologic and genetic differences (Kottelat & Freyhof, 2007). For a long time, *Phoxinus phoxinus* (Linnaeus, 1758) - as its common name Eurasian minnow suggests - was thought to be a widespread species throughout Europe. However, the taxonomy of the genus was revealed to be more complex (Kottelat, 2007; Kottelat & Freyhof, 2007) including substantial cryptic species diversity (Palandačić et al., 2015, 2017, 2020). At present, 22 distinct genetic *Phoxinus* lineages have been distinguished, comprising at least 13 recognised species (Palandačić et al., 2020). The neotype locality for *P. phoxinus* was placed on the Agger stream (Kottelat, 2007), a tributary of the river Sieg in Germany that feeds into the Rhine river. The designation of a neotype locality is used when the holotype material and sources used to describe a species are missing. Prior to 2012, minnows were a rare occurrence in that area (Landesamt für Natur, Umwelt und Verbraucherschutz Nordrhein-Westfalen, 2020). Taxonomically problematic is the fact that the connection between the species name *P. phoxinus* and its assigned lineage "10" (Knebelsberger et al., 2015) is uncertain. This is due to the lack of genetic analysis of the neotype (NRM-55108) designated by Kottelat, (2007), which comes from an area with multiple present lineages (5b, 10, 12; defined by a combination of morphological and single-gene markers) (Palandačić et al., 2020, 2022). Several studies and researchers are currently dealing with resolving the taxonomic and phylogenetic conundrum this genus poses (Bogutskaya et al., 2023; Corral-Lou et al., 2019; Denys et al., 2020; Palandačić et al., 2017, 2020; Reier et al., 2022; Vucić et al., 2018). Until now, these studies suffer from the lack of a high quality reference genome, which would make it possible to go beyond the analysis of single genes.

Due to the previously unrecognised species diversity, most of what is known about the natural history and behaviour of Eurasian minnows is attributed to *P. phoxinus* or its synonyms. Certain presumptions about minnows are assumed to be universally applicable for the entire genus. In the following we will refer to Eurasion minnows including all cryptic species in the *Phoxinus* genus. For instance, Eurasian minnows are small, schooling freshwater fishes that occupy oxygen-rich, cold streams of flowing waters, lake-shores with wave-movements, or highland lakes (Frost, 1943; Mills, 1987, 1988). They assume a complex role in the river or lake ecosystem as they feed on algae, plant debris, molluscs, crustaceans, and insects (Frost, 1943). *Phoxinus* species interact with smaller animals such as water snails and insect larvae, but also larger fishes such as co-occurring salmonids (Museth et al., 2010). However, seemingly contradictory behaviours have been reported as well. According to Frost (1943), *Phoxinus* spp. spawn over clean gravel and shallow, flowing waters. However, populations from the Baltic Sea reportedly reside and spawn in brackish habitats (Svirgsden et al., 2018). While Eurasian Minnows are sometimes reported to be sensitive to pollution (Bagge & Hakkari, 1992), they are also tolerant of polluted waters (King et al., 2011). They might even benefit from increased primary production in eutrophic waters (King et al., 2011; Moiseenko et al., 2009). In terms of dispersal and migration behaviour, Eurasian minnows from the Volga were less frequently found in migratory habitats (Pavlov & Mikheev, 2017), or demonstrated a tendency for downstream movement in a mesocosm experiment conducted with specimens from Wales (Jones et al., 2021). In contrast, Eurasian minnows were found to migrate upstream during breeding season (Frost, 1943; Svirgsden et al., 2018) as well as through fishways (Jansen et al., 1999) in populations from England, Estonia, and Germany, respectively. The aforementioned reported differences or the seemingly broad occurrence pattern could be due to the fact that studies reporting on *P. phoxinus* or "the Eurasian Minnow" were conducted on different *Phoxinus* species.

A crucial step towards informed and state-of-the-art biodiversity, phylogenetic, and comparative genomic research is the generation of haplotype-resolved, chromosome-level reference genomes. Phasing of haplomes, the haplotype-resolved elements in reference genome assemblies, is now easily facilitated with the latest development of chromosome conformation capture, namely Hi-C (Belton et al., 2012; Lin et al., 2018) and its integration into assembly pipelines, such as Hifiasm (H. Cheng et al., 2021). Resultantly, the number of studies acknowledging genetic variation between haplomes, is rapidly

increasing including e.g. Cavendish banana (Huang et al., 2023), African cassava (Mansfeld et al., 2021),
Australian lime (Nakandala et al., 2023), carnation (H. Jiang et al., 2023), cattle (Low et al., 2020), pearl
oyster (Takeuchi et al., 2022), and ocellated puffer fish (Zeng et al., 2023).

We conducted comparisons between the two *Phoxinus* haplomes to identify both sequence and
structural variations present in a *P. phoxinus* genome. When examining sequence variation,
heterozygosity, polymorphism, and allelic diversity were considered (Carley et al., 2022; DeWoody et al.,
2021). Structural variation can be examined within and among genomes by investigating the copy
number, orientation and location of coding and non-coding sequences on the chromosome
(Wellenreuther et al., 2019). Copy number variation comprises deletions, insertions, and duplications,
while orientational variation encompasses inversions. Local variation can be attributed to translocations.
Structural variation at the genome level has been associated with variation in both phenotypes and
complex traits (Mérot et al., 2020; Weischenfeldt et al., 2013).

Ultimately, this reference genome serves two purposes. First, it is to aid the *Phoxinus* community
with the necessary basis for phylogenetic unravelling. Second, this haplotype-resolved, chromosome-
level reference genome demonstrates once more that genetic diversity contains an individual layer,
namely, the intra-individual diversity on the haplome level that needs to find attention in the scientific
community.

 Methods

Specimen Collection and Sampled Tissues

Two *P. phoxinus* specimens originally from the river Sieg close to the neotype locality, North-Rhine
Westphalia, Germany, were sacrificed using Ethyl 3-aminobenzoate methanesulfonate salt (MS-222;
Sigma-Aldrich) under permission per § 11 Abs. 1 Nr. 1 b, Sa and 8d Tierschutzgesetz (TierSchG) issued by
the Amt für Umwelt, Verbraucherschutz und Lokale Agenda, Lebensmittelüberwachung und
Veterinärdienste, under Zeichen 56.2. Tissues from fin, brain, liver, spleen, muscle, heart, skin, and testis
were immediately sampled and flash-frozen in liquid nitrogen (individual identifiers ZFMK-TIS-60719
[male] and ZFMK-TIS-60720 [male]. Tissues (brain, liver, spleen, muscle, heart, gonad) from individual
ZFMK-TIS-60720 were used for the genome assembly; tissues (brain, gill, gonad, liver, muscle, skin,
spleen) from ZFMK-TIS-60719 were used for RNA-sequencing and annotation.

Extraction of Long Genomic DNA

Long genomic DNA (gDNA) of *P. phoxinus* was extracted with the circulomics Nanobind Tissue Big
DNA kit (part number NB-900-701-01, protocol version Nanobind Tissue Big DNA Kit Handbook v1.0
(11/19)) according to the manufacturer's instructions. In brief, 11 mg of spleen tissue were minced into
small slices on a clean and cold surface. Tissues were finally homogenised with the TissueRuptor II
device (Qiagen) making use of its maximal settings. After complete tissue lysis, the remaining cell debris
were removed, and the gDNA was bound to circulomics Nanobind discs in the presence of Isopropanol.
High molecular weight (HMW) gDNA was eluted from the nanobind discs in EB. The integrity of the
HMW gDNA was determined by pulse field gel electrophoresis using the Pippin Pulse™ device (SAGE
Science) and the Agilent Femtopulse. The majority of the gDNA was between 20 and more than 400 kb
in length. All pipetting steps of long gDNA have been done very carefully with wide-bore pipette tips.
HMW gDNA was further purified after extraction with 1x AMPure beads.

PacBio HiFi Library Preparation and Sequencing

A long insert HiFi library was prepared as recommended by Pacific Biosciences according to the
"Guidelines for preparing HiFi SMRTbell libraries using the SMRTbell Express Template Prep Kit 2.0" (PN
101-853-100, version 03). In summary, 10 µg of purified HMW gDNA was sheared twice, aiming for 25
and 20 kb fragments, respectively, with the MegaRuptor™ device (Diagenode). The sheared gDNA was
enriched with 1x AMPure beads according to the manufacturer's instructions. 6 µg sheared gDNA have
been used for Pacbio HiFi library preparation. The PacBio SMRTbell™ library was size-selected for
fragments larger than 6 kb with the BluePippin™ device according to the manufacturer's instructions.
The final HiFi library had a fragment size of 12.8 kb on the Agilent Fragment Analyzer on the High
sensitivity 50kb large fragment kit.

60 pM of the size selected library were run on two Sequel II SMRT cells with the SEQUEL II
sequencing kit 2.2 for 30 hours on the SEQUEL II; pre-extension time was 2 hours. PacBio CCS reads
(read quality > 0.99) were created from the subreads.bam files using PacBio's ccs command line tool
(version 6.3.0) and further refined by applying the tool DeepConsensus (v0.2, on PacBio reads within
98.0-99.5% read accuracy) (Baid et al., 2023). Finally, reads containing the PacBio adapter sequence
were filtered out by applying a Blastn search (v2.9) (Altschul et al., 1990) providing the PacBio adapter
sequence and the following arguments: ``reward 1 -penalty -5 -gapopen 3 -gapextend 3 -dust no -
soft_masking false -evalue 700 -searchsp 1750000000000 -outfmt 7". This resulted in a total yield of
37,3 Gb data with a mean read length of 10.7 kb and a read N50 of 10.2 kb.

ARIMA HiC Chromatin Conformation Capturing

Chromatin conformation capturing was done making use of the HiC+ Kit (Material Nr. A410110) and
followed the user guide for animal tissues (ARIMA-HiC 2.0 kit Document Nr: A160162 v00). In brief, 40
mg flash-frozen powdered muscle tissue was cross-linked chemically. The cross-linked chromatin was
digested with a restriction enzyme cocktail consisting of four restriction enzymes. The 5'-overhangs
were filled in and labelled with biotin. Spatially proximal digested DNA ends were ligated, and, finally,
the ligated biotin-containing fragments were enriched and went for Illumina library preparation, which
followed the ARIMA user guide for Library preparation using the Kapa Hyper Prep kit (ARIMA Document
Part Number A160139 v00). The barcoded HiC libraries run on an S4 v1.5 XP flow cell of a NovaSeq6000
with 300 cycles.

RNA Extraction and Sequencing

RNA was extracted from nitrogen flash-frozen brain, gill, gonad, liver, muscle, skin, and spleen
(specimen identifier ZFMK-TIS-60719) tissues. All tissues except for spleen and liver were extracted
using the RNeasy Mini Kit (Qiagen, Cat. No. 74104) suitable for tissue amounts of 0.5 to 30 mg. Spleen
and liver were extracted using QIAamp DNA Micro Kit (Qiagen) suitable for starting material of less than
5 mg. All tissues were homogenised using PowerBead Tubes Ceramic (Qiagen, diameter 2.8mm; Cat. No.
13114-50) on a PowerLyzer™ 24 Bench Top Bead-Based Homogenizer (MO BIO Laboratories, Inc). Before
use, 180 µL 1 M dithiothreitol (DTT) was added to 4.5 ml lysis buffer RLT. 350 µL prepared RLT buffer
was added to each tissue in a PowerBead Tube (Qiagen) and one cycle of 45 seconds at 3500 rpm and
another for 45 seconds at 4200 rpm on the PowerLyzer™ Homogenizer with a 5-minute pause at -20°C
were performed. After homogenisation, the solution was centrifuged for 1 minute at 14800 rpm. The
lysate including tissue debris was transferred into a fresh 1.5 mL tube (LoBind®) and again centrifuged at
14800 rpm for 3 minutes at room temperature (RT). Finally, the lysate was transferred to a gDNA
Eliminator column (provided in the QIAamp Kits) and centrifuged at 10,000 rpm for 30 sec at RT.
Washing and elution steps followed the standard protocol of the Micro and Mini Kit, respectively.
Samples were eluted into 14 µL (spleen, liver) and 30 µL (all other tissues) RNAse-free water during 1
minute at 10,000 rpm centrifugation. RNA library preparation and sequencing were performed at
Biomarker Technologies (BMKGene) GmbH on an illumina Novaseq 6000 S4 chip.

K-mer based Genome Size and Heterozygosity Estimation

To estimate the genome size and heterozygosity of the genome, a k-mer-based approach was used.
K-mers were counted in the HiFi reads using Jellyfish (v2.3.0) (Marçais & Kingsford, 2011) with
parameters count *-C -m*, with *-C* indicating that the input reads from both the forward and reverse
strand are being used, and -m specifying a k-mer length; we specified k-mer lengths of 19, 23, 25 and 30.
The resulting k-mer counts were exported into a k-mer count histogram using the *histo* command of
Jellyfish for subsequent analysis in GenomeScope (v2.0) (Ranallo-Benavidez et al., 2020) for estimation
of genome size and heterozygosity.

De novo **genome Assembly and Scaffolding**

Initial contigs for two representative Haplomes were generated using HiFiasm (v0.16.1-r375) (H.
Cheng et al., 2021) with parameters *--hic -l2* and then purge-dups (v1.2.3) (Guan et al., 2020) was run on
each produced haplotype assembly.

Initial scaffolding of the primary assembly was performed by mapping HiC reads to the primary
contigs using bwa-mem (v0.7.17-r1198-dirty) (H. Li, 2013) and mappings were filtered following the VGP
arima mapping pipeline (https://github.com/VGP/vgp-assembly/tree/master/pipeline/salsa). The final
bed file was given to yahs (v1.1a) (Zhou et al., 2022) for scaffolding. Scaffolds were then manually
curated into chromosomes using higlass (v2.1.11) (Kerpedjiev et al., 2018) with Cooler (v0.9.1)
(Abdennur & Mirny, 2020) to visualise the HiC data (Supplementary Figure S1 - S2).

Finally, the two haplotype assemblies were polished by mapping the HiFi reads to the assemblies
using pbmm2 (v1.3.0) [https://github.com/PacificBiosciences/pbmm2.git] with arguments --preset CCS -
N 1 and variants called using deepvariant (v0.2.0) with --model_type=PACBIO. Errors were corrected in
the assembly by filtering the vcf file given by deepvariant (v0.2.0) (Poplin et al., 2018) with bcftools view
(v1.12) (Danecek et al., 2021) with arguments *-i 'FILTER=\"PASS\" && GT=\"1/1\"'* and a consensus
called with bcftools consensus. Assembly completeness was assessed using BUSCO (v5.4.7) (Simão et al.,
2015) ran in genome mode with the actinopterygii_odb10 database.

Annotation of Repetitive Elements

Repeat sequences were identified and annotated in both haplomes of the *P. phoxinus* reference
genome by combining homology-based and *de novo* approaches. We identified low-complexity repeats
and tandem repeats with DustMasker from BLAST+ (v.2.9) from the NCBI C++ toolkit (Morgulis et al.,
2006) and Tandem Repeat Finder (v4.10.0) (Benson, 1999), respectively. We de novo identified
Transposable Elements (TEs) with Repeatmodeler (v2.0.1) (Smit and Hubley, 2008) to create a species-
specific library, and we combined the identified species-specific transposable elements (TEs) with the
Danio rerio (zebrafish - Cypriniformes) repeats library from Dfam 3.1 (Hubley et al., 2016) to generate a
custom combined repeats library.

Using this custom combined library, repetitive elements were identified and annotated in our
genome with RepeatMasker (v.4.1.0) (https://repeatmasker.org/). Divergence of our repeat annotation
from the consensus repeats sequence was estimated with the *calcDivergenceFromAlign.pl* script
implemented in RepeatMasker using the *.align* output file of the previous masking step. The repeats
landscape was plotted with the *createRepeatLandscape.pl* script, also in RepeatMasker. Output files of
Duskmasker, Tandem Repeats Finder and RepeatMasker were converted to *gff3* format. We merged the
positions of repeats with Bedtools (v2.29.2) (Quinlan, 2014) before our genome was softmasked with
these identified repeats using Bedtools (v2.29.2) in preparation for protein prediction.

Protein Annotation

Structural annotation of the protein-coding genes in the P. phoxinus genome was carried out with a
combination of ab initio, homology, and transcriptome-based protein prediction models. The
sequencing reads (NCBI accession numbers SRR26699630 to SRR26699636) from the seven RNA
samples from brain, gill, gonad, liver, muscle, skin, and spleen were mapped to the reference genome
with STAR (v2.7.10b) (Dobin et al., 2013), the resulting mapping files (.bam) were merged with samtools
(v1.10) (H. Li et al., 2009).

BRAKER3 (Brůna et al., 2021; Gabriel et al., 2023; Hoff et al., 2016), the latest genome annotation
pipeline in the BRAKER (Hoff et al., 2019) suite was used for protein annotation. Using our assembled
transcripts as input, BRAKER3 predicts protein-coding genes with GeneMarkS-T (Tang et al., 2015) and
then selects proteins with the best similarity scores and the quality of *ab initio* predictions with
GeneMark-ETP (Brůna et al., 2020). GeneMark-ETP then produces three groups of hints, i) hints with
transcript and protein similarity support, ii) hints with transcript and *ab initio* support and finally, iii)
hints with only protein similarity support using ProtHint. All these hints are used to create a set of high-
confidence genes. AUGUSTUS (Stanke et al., 2006) is trained with these gene sets and does an *ab initio*
prediction of another genome-wide gene set before TSEBRA (Gabriel et al., 2021) combines the results
of AUGUSTUS and GeneMark-ETP to produce a final set of high-confidence genes. The final set was used
as input to BUSCO (v5.4.7), which was run in protein mode with the actinopterygii_odb10 database.

Structural annotation of the resulting predicted genes was done with the *blastp* algorithm of
DIAMOND (v2.1.8) (Buchfink et al., 2021). Our predicted proteins were blasted against protein
databases of Swissprot edition 2023-05-13 (Boeckmann et al., 2003), The Protein Data Bank (Berman et
al., 2000) Pdbaa database edition 2023-05-13 and TrEMBL edition 2023-05-14 (Boeckmann et al., 2003).

The blast was done with an e-value cut-off of < $1*10^{-5}$. Functional annotation of these protein domains
was done with eggNOG-mapper (v2.0) (Cantalapiedra et al., 2021).

Protein sequences of *D. rerio* (zebrafish, Cypriniformes), GCF_000002035.6 (Howe et al., 2013);
Sinocyclocheilus rhinocerous (rhinoceros golden-line barbel, Cypriniformes), GCF_001515625.1 (J. Yang
et al., 2016); *Sinocyclocheilus grahami* (golden-line barbel, Cypriniformes), GCF_001515645.1 (J. Yang et
al., 2016); *Cyprinus carpio* (common carp, Cypriniformes), GCF_000951615.1 (Xu et al., 2014); *Carassius*
auratus (goldfish, Cypriniformes), GCF_003368295.1 (Z. Chen et al., 2019); *Pimephales promelas*
(fathead minnow, Cypriniformes), GCF_016745375.1 (Martinson et al., 2022); *Megalobrama*
amblycephala (Wuchang bream, Cypriniformes), GCF_018812025.1(H. Liu et al., 2021);
Ctenopharyngodon idella (grass carp, Cypriniformes)*,* GCF_019924925.1 (Wu et al., 2022); *Myxocyprinus*
asiaticus (Chinese high-fin banded shark, Cypriniformes)*,* GCF_019703515.2 (Krabbenhoft et al., 2021);
and the outgroup, *Ictalurus punctatu*s (channel catfish, Siluriformes), GCF_001660625.3 (Waldbieser et
al., 2023) were retrieved from NCBI RefSeq database (O'Leary et al., 2016) on 2023-07-12. We
considered *Pimephales promelas* (Fathead minnow) as a representative of the most closely related
taxonomic species to *P. phoxinus,* for which a well-documented whole genome assembly and
annotation is readily accessible. AGAT (v1.1.0) (Dainat, 2021) was used to extract the longest gene
isoforms against which the protein sequences of the *P. phoxinus* were blasted with DIAMOND (v2.1.8)
(Buchfink et al., 2021), to identify homologous sequences.

Genome-wide Comparison between Haplomes

Sequence Variation
Genome-wide average heterozygosity was estimated for both haplomes to assess sequence variation
using the 25 largest scaffolds representing the 25 chromosomes. To do so, raw HiFi reads were mapped
to both haplomes with minimap2 (v2.1) (H. Li, 2018) with preset *-ax asm20*, which optimises alignments
between sequences with less than 20% divergence. The resulting alignment files were filtered for
mapping quality lower than 20 with the view *-q 20* command and sorted with the *sort* command using
Samtools (v1.10) (H. Li et al., 2009). Duplicates were removed with the markdup -r command as
implemented in Sambamba (v0.7.1) (Tarasov et al., 2015). Furthermore, we generated windows of 100
kb with the *makewindows* command implemented in Bedtools (v2.31.0) with *-w 100000.* Then, we
estimated depth in 100 kb for both haplomes using *coverage -mean* as implemented in Bedtools
(v2.31.0) (Quinlan, 2014) in combination with the file specifying coordinates of 100kb windows created.

To estimate heterozygosity, site allele frequency likelihood was estimated using the GATK model for
genotype likelihoods in combination with the *-doSAF 1* parameter in ANGSD v.0.940 (Korneliussen et al.,
2014), the site frequency spectrum was estimated with the *realSFS* parameter also in ANGSD. Finally,
heterozygosity was estimated from sample allele frequencies in non-overlapping windows of 1 Mb. The
number of heterozygous alleles was counted within each Mb and then converted to the number of
heterozygous alleles per kb and plotted per chromosome across the genome using ggplot2 (v3.1.3), to
visually inspect patterns of heterozygosity across the entire genome. Patterns of heterozygosity often
reflect standing genetic variation which may be linked to demographic history of a species (Guevara et
al., 2021; Miller et al., 2014; Nadachowska-Brzyska et al., 2016).

Demographic History of P. phoxinus
The demographic history of *P. phoxinus* was reconstructed using Pairwise Sequentially Markovian
Coalescent (PSMC) (v0.6.5) as implemented by Li & Durbin, (2011). Variants were called from both
haplomes with a combination of samtools (v1.17) *mpileup* (H. Li et al., 2009) and Bcftools (v1.17) *call*
(Danecek et al., 2021) commands. The resulting variant file was converted to a diploid consensus file
with vcfutils.pl vcf2fq accessory implemented in the PSMC package. The resulting fastq files were
converted into the input fasta format for PSMC using the fqpsmcfa accessory. The PSMC model was run
with the following parameters: -N25 -t15 -r5 -p "4+25*2+4+6" and 100 bootstraps, a generation time of

3 years (Museth et al., 2010), which is the maturation age of the *P. phoxinus* and an assumed mutation
rate of 3.51 x10^{-9} (Tian et al., 2022). Results of the run were plotted using ggplot2 (v3.1.3).

Structural Variation

We aligned both haplomes with minimap2 (v2.26) (H. Li, 2018) using the *-asm5* preset setting.
Alignments were processed with Synteny and Rearrangements Identifier (SyRI) (v.1.6.3) (Goel et al.,
2019) to detect structural variation between both haplomes and the results were visualised using the R
package plotsr (v.1.1.1) (Goel & Schneeberger, 2022). Genes overlapping Insertions, deletions and
Inversions extracted using bedtool (v2.29.2) (Quinlan, 2014). Subsequently, functional enrichment of
these genes was achieved with g:profiler (Raudvere et al., 2019) using the zebrafish database option.
Finally, the results were visualised by go-figure (v1.0.2) (Reijnders & Waterhouse, 2021).

Identification of Orthologs and Gene Family Evolution

From the above described proteomes, amino acids shorter than 30 bp were filtered out, and
Orthofinder (v2.5.4) (Emms & Kelly, 2019) with default settings was run to infer orthologous gene
families. Orthofinder created orthogroup gene trees and then inferred a rooted species tree from gene
duplication events with the STRIDE approach (Emms & Kelly, 2017) using just single-copy orthologs. The
species tree underwent conversion into an ultrametric tree utilising Orthofinder's *make_ultrametric.py*
script, employing the parameter -r 142. This value corresponds to approximately 142 million years,
which aligns with the estimated median divergence time between Siluriformes and Cypriniformes, as
suggested by TimeTree (Kumar et al., 2017). This estimate is calculated by considering the median of
divergence times that have been publicly estimated using molecular sequence data analysis.
Orthologous gene counts from the Orthofinder run were filtered to remove ortholog groups that are
overrepresented in a particular species; this was done with the *clade_and_size_filter.py* script in CAFE
(v5.0.0) (Mendes et al., 2021). Gene family evolution analysis was carried out in CAFE5 (v5.0.0) using the
ultrametric tree and filtered orthologous gene counts as input. Genes that were confirmed to be
species-specific, expanding and contracting in *P. phoxinus,* were blasted against the TrEMBL database.
The GO terms derived from the TrEMBL database were then used as input into REViGO (Supek et al.,
2011) to remove redundant terms and summarise GO results.

Results and Discussion

K-mer Based Genome Size and Heterozygosity Estimation

We estimated the haploid length of the *P. phoxinus* genome to be 805.8 Mbp with a unique content of 62.1% and a heterozygosity of 1.43%. This estimate deviates from the determined C-value of 1.15 pg (corresponding to 1.13 Gbp) (genomesize database; https://genomesize.com/). Running genome size estimates using 19, 23 ,25 and 30 k-mer lengths, showed slight variation in the estimated genome sizes and heterozygosities (Supplementary Table S1, Figure S3). We chose the 19-mer length due to a lower error rate in comparison to other k-mer lengths. The estimated heterozygosity of 1.43% is high compared to other published k-mer-based heterozygosity estimates in cyprinid fishes. For example, *Onychostoma macrolepis* has an estimated heterozygosity of 0.29% (L. Sun et al., 2020), *Myxocyprinus asiaticus* (Chinese high-fin banded shark) of 0.20% (Krabbenhoft et al., 2021), *Gymnocypris przewalskii* (Przewalskii's naked carp) of 0.96% (Tian et al., 2022), and *Poropuntius huangchuchieni* of 0.68% (L. Chen et al., 2021). To the best of our knowledge, the highest k-mer-based heterozygosity estimate for a cyprinid is 1.82% estimated in *Gymnocypris eckloni* (F. Wang et al., 2022), while the lowest estimate of 0.1% was found in *Triplophysa tibetana* (X. Yang et al., 2019). This suggests the estimated k-mer-based heterozygosity of *P. phoxinus* is on the high side within cyprinids.

The clear bimodal coverage distribution visible from the k-mer plot (Figure 1) is indicative of the diploidy of the *P. phoxinus* genome, with the first peak at around 18-fold coverage corresponding to heterozygous sites and the second peak at around 36-fold coverage corresponding to homozygous genome positions.

Figure 1: K-mer spectra profile of *Phoxinus phoxinus*. Generated from raw PacBio HiFi reads with GenomeScope2. The y-axis shows the k-mer counts and the x-axis shows sequencing depth. The clear bimodal pattern observed indicates a diploid genome with high heterozygosity of 1.43%. The haploid genome size was estimated to be around 805 Mb.

Genome Assembly and Scaffolding

We generated a haplotype resolved assembly of *P. phoxinus*, with a total assembly size of 940 Mbp and 929 Mbp for haplotype 1 (Hap1) and haplotype 2 (Hap2), respectively. The initial assembly of each haplotype contained 394 primary contigs, which were further placed into 101 (N50: 36.4 Mb) and 81 (N50: 36.6 Mb) scaffolds, respectively, by mapping HiC reads to the primary contigs. Integrating Hi-C data, we ultimately scaffolded 99.6% of the Hap1 assembly and 99.5% of the Hap2 assembly each into 25 chromosomes (Supplementary Table S2). BUSCO analysis revealed the *P. phoxinus* assemblies had only 2.3% and 2.1% missing genes. The assembly statistics for both haplomes (Table 1) denote high contiguity and high completeness. The 11 Mb difference in size between both haplomes is a result of structural variations between haplomes.

Table 1. Summary of *P. phoxinus* haplotype 1 and haplotype 2 genome assembly statistics.

Assembly Statistics	Haplotype 1	Haplotype 2
Assembly Size	940 Mb	929 Mb
Contigs	394	394
Scaffolds	101	81
N50 Contigs	3.85 Mb	4.59 Mb
N90 Contigs	869 Kb	1.05 Mb
N50 Scaffolds	36.4 Mb	36.6 Mb
N90 Scaffolds	30.7 Mb	30.3 Mb
% of Assembly in Scaffolds	99.6%	99.5%
Complete BUSCOs	3528 (96.9%)	3540 (97.2%)
Complete and Single-copy BUSCOs	3488 (95.8%)	3495 (96.0%)
Complete and duplicated BUSCOs	40 (1.1%)	45 (1.2%)
Fragmented BUSCOs	30 (0.8%)	25 (0.7%)
Missing BUSCOs	82 (2.3%)	75 (2.1%)
Total BUSCO Search	3640	3640

Repeats Annotation

Teleost fish genomes are known for their highly diverse repeat content (Gao et al., 2016) and the assembly of the *P. phoxinus* genome follows this precedent. Annotation of the assembly using our custom combined repeats library revealed that around 506 Mb (53.86 %) of the *P. phoxinus* genome is made up of repetitive elements. This is higher than reported to date from other species of the family cyprinids like fathead minnow (43.27%) (Martinson et al., 2022), grass carp (43.26%) (Wu et al., 2022), common carp (31.3%) (Xu et al., 2014) and goldfish (39.6%) (Z. Chen et al., 2019). The repeat content is slightly lower than in zebrafish, which contains around 54.47% repetitive elements (Howe et al., 2013).

From the identified TEs, DNA transposons were the most abundant repeat elements in the genome, making up 21.25% (199 Mb) of the genome. SINEs were relatively poorly represented at 1.03%, in line with the expectations derived from other published fish genomes (Shao et al., 2019; Sotero-Caio et al., 2017). DNA transposon content is similar to that of the fathead minnow genome, which contains an

371 estimated 21.47% (Martinson et al., 2022). Retroelements made up 11%; 6.16% LTRs and 4.15% LINEs.
372 Overall, we annotated 46.59% of the *P. phoxinus* genome to be total interspersed repeats, the
373 remaining repetitive elements were annotated as 0.47% small RNA, 1.50% satellites, 2.66% simple
374 repeats, and 0.17% low complexity repeats (full details of all repetitive elements are described in Table
375 S3 and S4).

Figure 2: Repeat landscape of the *P. phoxinus* genome. Distribution of transposable elements (TEs) in the *Phoxinus phoxinus*
genome across different Kimura substitution levels (in %). TEs on the left side of the histogram represent recently active TEs
with low divergence from the consensus sequence of TEs. TEs towards the right side of the histogram represent ancient TEs
with higher degrees of divergence from the consensus TE sequences.

Repeat landscapes illustrate how TEs cluster in correlation to the Kimura substitution rate, i.e. how
quickly TEs in a genome have diverged from the consensus TE sequence (Shao et al., 2019). The Kimura
substitution level is correlated with the age of transposition activity: low substitution levels indicate
recent transposition events, higher substitution levels suggest old transposition events. The repeat
landscape of the annotated *P. phoxinus* genome (Figure 2) indicates a high level of active, i.e., recent,
transposition bursts (Watanabe et al., 2014). In fish genomes, active transposons accumulate at a faster
rate than they decline (Shao et al., 2019), leading to an abundance of active transposons with low
kimura substitution levels (Böhne et al., 2012; Gao et al., 2016; Sotero-Caio et al., 2017), an exact
pattern observed in TEs in the *P. phoxinus* genome. The repeat landscape for haplotype 2 can be found
in Supplementary Figure S4 .

Active transposition and high repeat content in the *P. phoxinus* genome may result in
rearrangements and variations in genome size (Kojima, 2019). Transposable elements can also function
as recombination agents, generating structural variations such as deletions, insertions, translocations
and inversions (Carvalho & Lupski, 2016; Cerbin & Jiang, 2018; Deininger et al., 2003), which can be
sources of standing genetic variation present in an individual.

Protein Annotation

Protein prediction, based solely on transcripts derived from our RNA-seq data mapped to the Hap1 and Hap2 genome assembly (Supplementary Table S5), generated 40,364 gene models and a mRNA count of 42,408, which covered 47.7% and 53.6% of the genome, respectively. The *ab-initio* prediction resulted in 24,420 genes, covering 49.9% of the genome. After filtering and generating a final high-confidence gene set, the final annotation contained 30,980 mRNAs, of which 23,397 were marked as high-confidence genes in Hap 1, and 29,614 mRNAs, with 23,191 high-confidence genes in Hap 2 (full summary statistics of protein annotation for Hap1 and Hap2 in supplementary Table S6 and S7). Using the predicted gene content of Hap1 of our *P. phoxinus* assembly in comparison to its closest relative with available whole-genome resources, the fathead minnow genome contains more predicted genes (26,150), (Martinson et al., 2022). We ran BUSCO with the actinopterygii_odb10 database and obtained a total of 3,484 (95.7%) complete BUSCOs, 3,435 (94.4%) complete and single-copy BUSCOS, 49 (1.3%) complete and duplicated BUSCOs, 20 (0.5%) fragmented BUSCOs and 136 (3.8%) missing BUSCOs. Structural annotation of the predicted proteins resulted in 22,761 (97.28%) hits in at least 1 database, with 19,871 (84.56%) proteins shared across all four investigated databases (Table 2). These statistics hint at a well-assembled and annotated *P. phoxinus* genome. Homology search of genes in *P. phoxinus* against zebrafish, fathead minnow, goldfish, common carp and grass carp had 21,206 (90.6%) genes annotated across all 5 species (Supplementary Figure S5)

Table 2. Functional annotation of genes in *P.phoxinus* genome across four different databases: TrEMBL, Swissprot, PDBAA, and eggNOG

Database	Number	Percentage (%)
Swissprot	19,978	85.39
TrEMBL	22,761	97.28
egg-NOG	21,296	91.02
PDBAA	19,978	85.39
Gene overlap	19,871	84.93
Total Annotated genes	22,761	97.28

Genome-wide Comparison of Structural and Sequence Variation between Haplomes

Sequence variation follows universal pattern of eukaryotes

Genome-wide estimates and patterns of mean heterozygosity were similar between both haplomes, with an estimate of 0.7866% in Hap1 and 0.7926% in Hap2. Genome-wide heterozygosity estimates were much lower than k-mer-based estimates and this may be due to the difference between the two approaches, with the genome-wide approach being more accurate as it makes estimates from direct observation of different loci that have been confidently mapped, and the k-mer-based approach being more sensitive to low coverage and error-prone regions. The distribution of heterozygosity across the chromosomes varied, but the pattern found on each chromosome was similar (Figure 3). The regions of highest heterozygosity were always found at both ends of each chromosome and the lowest heterozygosity was found in the central region of each chromosome.

It is possible that the regions of high heterozygosity are linked to telomeric regions of the chromosome, which generally have an elevated recombination rate (Nachman, 2002), which in turn can be linked to higher genetic diversity (Jaramillo-Correa et al., 2010). In fishes, this was evidenced, e.g., in the Atlantic silverside (*Menidia menidia*) (Tigano et al., 2021) and brown trout (*Salmo trutta)* (Leitwein et al., 2016) genomes. This contrasts with regions of reduced heterozygosity found at central locations potentially linked to centromeres, which in eukaryotes are known as "cold-spots" for recombination, hence have low levels of recombination (Yan & Yu, 2022).

Genomes with high heterozygosity can pose assembly challenges due to multiple scaffold formation from the same sequence (Kajitani et al., 2014; Pryszcz & Gabaldón, 2016). However, implementation of long range information such as Hi-Fidelity long reads and Hi-C contact maps can aid in overcoming these difficulties by reducing sequence redundancy (Kronenberg et al., 2021). Our high-quality assembly, indicating augmented levels of standing genetic variation, is a testament to this approach. Previous studies have also indicated abundant levels of genetic variation in the *Phoxinus* genus (Bogutskaya et al., 2020; Reier et al., 2022), potentially enabling more expeditious responses to environmental changes and even greater potential for adaptation across diverse environments (Lai et al., 2019; Matuszewski et al., 2015).

Demographic History of *P. phoxinus*

Climatic fluctuations have always had an effect on the distribution and survival of species (Hewitt, 2000; J. Li et al., 2021). For aquatic species in the temperate zone, effects of climate fluctuations are even more pronounced as a result of glaciation cycles, which led to fluctuating sea water levels (Munnecke et al., 2010). A PSMC analysis enabled us to track fluctuations in effective population size *(Ne)* of *P. phoxinus* from more than 10 Million years ago (Mya) until around 10 thousand years ago (kya).

Figure 4 shows a gradual and continual rise in *Ne* with time, from roughly 20,000 individuals about 10 million years ago to around 25,000 individuals approximately 800,000 years ago. This incremental increase coincides with the Neogene period, which is divided into the Miocene and the Pliocene epochs

- both known for their global warmth and minimum ice volumes (Böhme et al., 2011; Haywood et al.,
2000; Hui et al., 2018; Steinthorsdottir et al., 2021). The climate conditions prevailing during this
timeframe suggest the existence of an environment favourable for the propagation and development of
freshwater species. Following a small increase in *Ne*, the population declined to fewer than 20,000
individuals between 800 kya and 100 kya, during the mid-Pleistocene period, which was marked by
severe quaternary glaciation and low global sea levels. This period also saw changes in global ice volume,
sea levels, and temperatures (Ehlers et al., 2006; Ruddiman et al., 1989; Y. Sun et al., 2019). During this
epoch, temperatures were on average 11°C lower than current values (Hughes et al., 2007) and lowland
glaciation periods were frequent (Rose, 2009). As a result, species either became extinct, migrated to
southern Europe seeking refuge, or underwent adaptation (Feliner, 2011).

This reduction in *Ne* during the Last Glacial Maximum is generally observed in fish (Jardim de
Queiroz et al., 2022; J. Li et al., 2021) and in birds (Nadachowska-Brzyska et al., 2015), especially in
European species (Yuan et al., 2018). In *P. phoxinus*, the end of Last Glacial Maximum was followed by a
very sharp increase in *Ne* from around 70 kya, peaking at nearly a million estimated individuals,
indicating that *P. phoxinus* was able to repopulate previously covered freshwater habitats across Europe.
The high *Ne* also corroborates the high heterozygosity estimates for this genome.

474
475
476 **Figure 4. Inferred demographic history of *P. phoxinus* from PSMC analysis**. The time ranges cover the Miocene, Pliocene, Early
477 to Late Pleistocene and Holocene, shown in different colours. The x-axis shows years before present on a logarithmic scale and
478 the y-axis shows the estimated effective population size. Bootstrap results are shown as transparent lines. Haplome 2 showed a
479 similar curve.
480

Large non-syntenic regions between P. phoxinus haplomes

A comparison of both haplomes of *P. phoxinus* with SyRI revealed around 815 to 820 Mb of syntenic
regions between both haplomes (Table 3, Figure 5). Our SyRI analysis also identified 1,572 translocations
of between 5.34 Mb and 5.36 Mb in both haplomes; 214 duplications totalling 571 kb in Hap1 and 667
duplications of around 1.8 Mb total length in Hap2; 215 inversions with a total length of around 10.8 Mb
in Hap1 and 11.1 Mb in Hap2 (Table 3, Figure 5).

Table 3. Identity**,** number, and total lengths of identified structural variations between haplomes 1 and
2 in the *P. phoxinus* reference genome.

Structural Variation	Count	Length haplotype 1 [bases]	Length haplotype 2 [bases]
Syntenic regions	1457	813,594,094	820,681,234
Inversions	215	10,843,815	11,161,414
Translocations	1572	5,342,064	5,361,801
Duplications (Haplotype 1)	214	57,1045	-
Duplications (Haplotype 2)	667	-	1,776,709
Insertions	447,578	-	8,108,033
Deletions	447,699	8,113,615	-

Inversions represented the dominant structural variation within the *P. phoxinus* genome in terms of
total size in either haplome (Table 3). A total of 215 inversions were identified, with the largest eight
inversions making up 72% (8 Mb) of the estimated 11 Mb of inversions present in the *P. phoxinus*
genome. Using Hap1 as a reference for inversion size, chromosome 6 contained the largest inversion
with a size of 1.8 Mb; closely followed by a 1.7 Mb inversion on chromosome 2; a 1.5 Mb inversion on
chromosome 21; a 1.1 Mb inversion on chromosomes 9 and 19, each; and finally an 800 kb inversion on
chromosome 3 (Figure 7).

We investigated genes enriched in regions of copy number structural variants, specifically deletions
and insertions, which can also be characterised as presence-absence variations (PAVs). GO terms
enriched in this region of PAVs between both haplotypes include terms such as "dephosphorylation",
"peptidyl-serine phosphorylation", and "protein processing", which have been linked with the activation
of sperm motility in numerous fish species (Dumorné et al., 2018; Zilli et al., 2008, 2017). Some GO
terms related to development were also enriched within PAVs, such as "regulation of developmental
growth" and "developmental pigmentation" (Supplementary Tables S8-S9 and Figure S6).

Genes enriched within locational structural variants, in this instance inversions, were primarily
associated with immunity and transposable elements in hAT, DDE, and L1 families . These GO terms are
"phagocytosis", "phagosome maturation involved in apoptotic cell clearance" and "phagolysosome
assembly involved in apoptotic cell clearance". Phagocytosis is a conserved immune defence mechanism
across multicellular organisms that prevents infection and invasion of the body by pathogens (Esteban
et al., 2015; Neumann et al., 2001) (Supplementary Tables S10-S11 and Figure S7).

Our haplotype-resolved genome allows for the identification of structural variants, such as inversions,
translocations, duplications, deletions, and insertions, between both haplomes of our reference genome.
To explore intra-individual structural variation, haplotype-resolved reference genomes have been
created in various organisms, particularly in plant species (S.-P. Cheng et al., 2021; L. Jiang et al., 2022; H.
Sun et al., 2022). It has also been utilised to a lesser degree to uncover structural variation in animals,
for example, oysters (Takeuchi et al., 2022), where transposable elements played a large role in the
structural variation between both haplomes, Additionally, in turkeys, structural variations have affected
genes involved in disease susceptibility and fertility (Barros et al., 2023).

Figure 5: **Synteny plot of all 25 chromosomes of Hap1 (top, blue line) and Hap2 (bottom, orange line) of the *P. phoxinus* genome assembly**. Aligned regions between both haplomes are shown in blocks of grey, unaligned regions are shown by gaps of white. Inversions are represented in orange, translocations in green, duplications in light-blue.

Haplotype-resolved genome assembly revealed high level of within-genome variation in P. phoxinus

Analysis of mean genome wide heterozygosity and structural variation, as outlined in the preceding sections, revealed a high level of variation present in the *P. phoxinus* haplomes. This is a possible reflection of high fecundity and large population size, expected in most freshwater fish species (Karjalainen et al., 2016; Röpke et al., 2021). The detection of genetic diversity in highly heterozygous species can be compromised by either un-phased genome-assembly approaches or haplotype-merged assemblies, which would ultimately lead to a significant underestimation of true genetic diversity (Nakandala et al., 2023; Takeuchi et al., 2022). Collapsed reference genomes restrict assembly accuracy and exploration of heterozygous regions of the genome, missing out on potentially interesting intra-individual variation (Low et al., 2020).

Gene Family Evolution - Histone Gene Families Retracted, NLRC Genes Expanded in *P. phoxinus*

Inferring gene orthology from sequence data is a fundamental approach for understanding evolutionary relationships between genes and species through their evolutionary history (Tekaia, 2016). A total of 396,455 genes across all 11 species described in the protein annotation section were included and 381,263 (96.12%) of these genes were successfully assigned to 27,672 orthogroups, i,e. gene-sets in those species or in a subset of these species, that derived from their respective last common ancestor (Supplementary Table S12). The total number of orthogroups comprising proteins from all species was 14,914 (Table 4). Single-copy orthogroups, also known as 1-to-1 orthogroups, contain genes that have evolved from a common ancestral gene but are present in a single copy across all proteomes used in an ortholog analysis (Creevey et al., 2011; Simão et al., 2015; Tekaia, 2016). Our orthology inference analysis revealed the presence of 301 single-copy orthogroups which provided a solid basis for phylogenetic studies.

Table 4. Summary statistics of orthology inference analysis of *P. phoxinus* and 10 other teleost species.

Overall Summary Statistic	
Number of species	11
Number of genes	396,455
Number of genes in orthogroups	381,263
Number of unassigned genes	15,192
Percentage of genes in orthogroups	96.2
Percentage of unassigned genes	3.8
Number of orthogroups	27,672
Number of species-specific orthogroups	2319
Number of genes in species-specific orthogroups	7975
Percentage of genes in species-specific orthogroups	2.0
Mean orthogroup size	13.8
Median orthogroup size	15.0
Number of single-copy orthogroups	301
Number of orthogroups with all species present	14,914

There were 273 genes distributed across 72 *P. phoxinus*-specific orthogroups; a homology blast of
these genes in the TrEMBL database returned only results for 130 genes, many of which were
uncharacterised proteins (Supplementary Table S13). Only 54 of these genes annotated from the
TrEMBL database returned GO annotations and these annotations were clustered into eight major GO
terms (Figure 6). The presence of GO terms "response to oxidative stress" and "regulation of response
to reactive oxygen", suggests that *P. phoxinus* is particularly sensitive to changes in water quality.
Observational studies have found conflicting evidence, with Eurasian minnows ranging from sensitive
(Bagge & Hakkari, 1992) to insensitive to intermediate pollution (King et al., 2011). Those findings may
be attributed to the fact that different species of *Phoxinus* were studied. Cyprinoidea in general have
been thought to be sensitive to oxidative stress, demonstrated in Amur minnow (*Phoxinus lagowskii*) (Q.
Wang et al., 2019; Yao et al., 2022) and Chinese rare minnow (*Gobiocypris rarus*) (Z.-H. Li et al., 2014),
while the fathead minnow (*Pimephales promelas*) has been used as a bioindicator of water quality
(Elizalde-Velázquez et al., 2020).

Figure 6: Gene ontology network of 54 functionally annotated *P. phoxinus* specific genes into eight major GO terms: "DNA
replication", "DNA repair", "regulation of DNA-templated transcription", "response to oxidative stress", "defense response to
virus", "cellular response to stimulus", "regulation of response to reactive oxygen species" and "negative regulation of TORC1
signalling". Bubble size indicates the frequency of the GO term in the gene ontology annotation database i.e larger bubbles
correspond to more general terms. Edges on the graph connect highly similar or linked GO terms to each other.

Gene family changes are expressed as duplication and loss of genes among related species and are
based on mutation, fixation, and retention rates of the genes in question (Demuth & Hahn, 2009). Gene
family expansion might be an indication of positive selection (Demuth & Hahn, 2009). An example is the
expansion of Myxovirus resistance (Mx) genes in the rainbow trout, providing special antiviral resistance
for their anadromous lifestyle (T. Wang et al., 2019). While the immune system is relatively conserved
across different jawed vertebrates (Flajnik & Kasahara, 2010; Litman et al., 2010), a wide array of
species-specific modifications in the underlying immune pathway were observed in other fishes, such as
cods, coelacanths, and sharks (Buonocore & Gerdol, 2016; Malmstrøm et al., 2016; Stein et al., 2007).

We analysed gene family expansions and contractions in *P. phoxinus* relative to 10 other teleost
species (Supplementary Figure S8). Since the split of *P. phoxinus* and the fathead minnow (Figure 7)

from their last common ancestor, *P. phoxinus* counted more significant gene contractions (309 gene families) than significant gene expansions (55 gene families) relative to the fathead minnow, which had 132 gene family expansions and 129 gene family contractions. The significantly expanded gene families contained genes such as *HIST1, H2B, CENP-T, GCRV, HIST2H2A, HIST2H3A, H3F3A, TRBV25-1*, which mostly belong to histone PFAM domains (Supplementary Figure S9). Immune genes belonging to the NACHT PFAM domain, were also more abundant in the set of contracted gene families (Supplementary Figure S10, genes like *NLRP12, NLRP3, CD22, CLEC17A, GIMAP7*. The expanding immune gene families mostly belonged to histone gene families, whereas the contracted gene families were mostly represented by NLRC gene families (Supplementary Tables S14 and S15).

Figure 7: **Ultrametric phylogenetic tree of selected teleost species and *P. phoxinus*.** Numbers on the branches represent counts for expanded (+) and contracted (-) gene families. Positive numbers indicate significant gene families that have been expanded since the split from the last common ancestor and negative numbers indicate significant gene families that have been contracted since the split from the last common ancestor.

Evolution of the immune system has greatly contributed to speciation in teleosts and a lot of diversity in immune system pathways has been identified in teleosts, like the expanded MHC1 and TLE genes in the Atlantic cod (*Gadus morhua*) (Star et al., 2011) and the absence of MHC class II-mediated immunity in the broad-nosed pipefish (*Syngnathus typhle)* (Haase et al., 2013). In the *P. phoxinus* genome*,* we observed an expansion in immunity-boosting histones *H1, H2B, H3*, and *H4,* particularly *H2A* and *H2B.* Histones are highly conserved proteins involved in the innate host defence against microbes (X. Li et al., 2022), they include core histones like *H2A, H2B, H3*, and *H4*. The final histone type, called *H1,* is generally referred to as a linker histone (X. Li et al., 2022; Westman et al., 2015). Histones are known to play a key role in antibacterial immunity across multiple species (X. Li et al., 2022; Pasupuleti et al., 2012). Histone 2A and 2B have antibacterial activity against different types of bacteria in fish. Histones are particularly implicated in fish mucosal immunity (Ángeles Esteban & Cerezuela, 2015), acting as a strong barrier to microbial invasion. This has been confirmed in the Indian major carp (*Labeo catla*) (Nigam et al., 2017), lumpfish (*Cyclopterus lumpu*s) (Patel & Brinchmann, 2017), gilthead seabream (*Sparus aurata*) (Jurado et al., 2015), and rainbow trout (*Oncorhynchus mykiss*) (Fernandes et al., 2002; Oscoz et al., 2005). The expansion of histone genes in *P. phoxinus* suggests a strong focus on mucosal protection against bacteria and other pathogens. Assuming, that similar evolutionary trends can be found in other *Phoxinus* species, this is a particularly interesting find, given that there are

populations of *Phoxinus* spp. in the Baltic Sea, some of which are anadromous and some of which are
entirely resident in the brackish water (Svirgsden et al., 2018). Such migration habits demand an
efficient immune system to cope with the different pathogens and parasite communities present in
brackish and freshwater.

In contrast, NLRC genes were found to be reduced in the *P. phoxinus* genome, particularly *NLRP3* and
NLRP12, which are important inflammatory genes (Morimoto et al., 2021). NLRC genes are majorly
involved in immune response to exogenous pathogens like bacterial and parasitic infections (Chang,
2023; Morimoto et al., 2021). NLRC genes are involved in immune response in Nile tilapia (*Oreochromis*
niloticus) in *Streptococcus agalactiae* and *Aeromonas hydrophila* infections (Q. Li et al., 2022). NLRC
genes were also overexpressed in response to *V. anguillarum* infection in the Mi-iuy croaker (*Miichthys*
miiuy) (G. Liu et al., 2023). In summary, the increased presence of genes involved in preventive mucosal
defence and decreased amount of genes involved in immune responsiveness, might hint to a change in
immune response strategy towards infection prevention.

Conclusion

In this paper, we present a highly contiguous and complete Eurasian minnow, *Phoxinus phoxinus,*
genome achieved by combining Pacbio Hifi reads and Hi-C data, along with comprehensive repeats and
gene annotation. Demographic history analysis and high observed heterozygosity seem to suggest
substantial population and individual diversity, respectively. While the identified diversity could pose a
challenge for a collapsed assembly, our phased genome enables a more accurate depiction of intra-
individual variation. Consequently, we anticipate this genome will prove an invaluable resource for
forthcoming research investigating the intricate taxonomic and demographic complexities of the
Phoxinus genus. For future studies, it will be of great importance to determine the genetic identity of
the *Phoxinus* spp. studied before associating natural history traits with the species name. Similarly, a
comprehensive understanding of the behaviour of Eurasian Minnows requires tracing the possible
genetic lineages in historical literature, in order to obtain a clearer picture.

Acknowledgments

635 We thank the Long Read Team of the DRESDEN Concept Genome Center, part of the MPI-CBG and the
636 technology platform of the CMCB at the TU Dresden, supported by DFG (INST 269/768-1).

Author Contributions
MS devised the project. TOO and MS lead the writing and devising of the manuscript. EWM, SW, and
TB performed genome sequencing and assembly. IC established the genome repeat and protein
annotation pipeline with support from SM. TOO performed genome-wide comparison, gene family
evolution and demographic history analyses. SM and AB provided TOO and IC with bioinformatic
support. AB contributed with sex-determination and genome assembly expertise. All authors
contributed to revising the manuscript.

Data, scripts, code, and supplementary information availability

Scripts used for genome assembly by EMW, SW and TB, scripts used for genome-wide comparison, gene family evolution and demographic history by TOO are available here: https://doi.org/10.5281/zenodo.10210241.

The repeat and protein annotation was implemented using a snakemake pipeline. The code is available here: https://doi.org/10.5281/zenodo.10220349. RNA data for annotation is available under NCBI accession numbers SRR26699630 to SRR26699636. *P. phoxinus* haplomes are available under NCBI Bioproject numbers PRJNA1030283 and PRJNA1030284 both of which are linked to the Euro-fish project under umbrella bioproject number PRJNA768423.

Conflict of interest disclosure

The authors declare that they comply with the PCI rule of having no financial conflicts of interest about the content of the article.

Funding

This research was funded by the Leibniz Association under project SAW-J96/2020 awarded to MS, and by the German Research Foundation under INST 269/768-1 awarded to the technology platform of the CMCB at the TU Dresden.

References

Abdennur, N., & Mirny, L. A. (2020). Cooler: Scalable storage for Hi-C data and other genomically labeled arrays. *Bioinformatics*, *36*(1), 311–316. https://doi.org/10.1093/bioinformatics/btz540

Altschul, S. F., Gish, W., Miller, W., Myers, E. W., & Lipman, D. J. (1990). Basic local alignment search tool. *Journal of Molecular Biology*, *215*(3), 403–410. https://doi.org/10.1016/S0022-2836(05)80360-2

Ángeles Esteban, M., & Cerezuela, R. (2015). 4 - Fish mucosal immunity: Skin. In B. H. Beck & E. Peatman (Eds.), *Mucosal Health in Aquaculture* (pp. 67–92). Academic Press. https://doi.org/10.1016/B978-0-12-417186-2.00004-2

Bagge, P., & Hakkari, L. (1992). Effects of paper mill effluents on the fish fauna of stony shores of Lake Päijänne. In V. Ilmavirta & R. I. Jones (Eds.), *The Dynamics and Use of Lacustrine Ecosystems* (pp. 413–420). Springer Netherlands. https://doi.org/10.1007/978-94-011-2745-5_43

Baid, G., Cook, D. E., Shafin, K., Yun, T., Llinares-López, F., Berthet, Q., Belyaeva, A., Töpfer, A., Wenger, A. M., Rowell, W. J., Yang, H., Kolesnikov, A., Ammar, W., Vert, J.-P., Vaswani, A., McLean, C. Y., Nattestad, M., Chang, P.-C., & Carroll, A. (2023). DeepConsensus improves the accuracy of sequences with a gap-aware sequence transformer. *Nature Biotechnology*, *41*(2), Article 2. https://doi.org/10.1038/s41587-022-01435-7

Barros, C. P., Derks, M. F. L., Mohr, J., Wood, B. J., Crooijmans, R. P. M. A., Megens, H.-J., Bink, M. C. A. M., & Groenen, M. A. M. (2023). A new haplotype-resolved turkey genome to enable turkey genetics and genomics research. *GigaScience*, *12*, giad051. https://doi.org/10.1093/gigascience/giad051

Belton, J.-M., McCord, R. P., Gibcus, J. H., Naumova, N., Zhan, Y., & Dekker, J. (2012). Hi–C: A comprehensive technique to capture the conformation of genomes. *Methods*, *58*(3), 268–276. https://doi.org/10.1016/j.ymeth.2012.05.001

Benson, G. (1999). Tandem repeats finder: A program to analyze DNA sequences. *Nucleic Acids Research*,

27(2), 573–580. https://doi.org/10.1093/nar/27.2.573

Berman, H. M., Westbrook, J., Feng, Z., Gilliland, G., Bhat, T. N., Weissig, H., Shindyalov, I. N., & Bourne,

P. E. (2000). The Protein Data Bank. *Nucleic Acids Research*, *28*(1), 235–242.

Boeckmann, B., Bairoch, A., Apweiler, R., Blatter, M.-C., Estreicher, A., Gasteiger, E., Martin, M. J.,

Michoud, K., O'Donovan, C., Phan, I., Pilbout, S., & Schneider, M. (2003). The SWISS-PROT

protein knowledgebase and its supplement TrEMBL in 2003. *Nucleic Acids Research*, *31*(1), 365–

370. https://doi.org/10.1093/nar/gkg095

Bogutskaya, N. G., Diripasko, O. A., & Palandačić, A. (2023). Novel data support validity of Phoxinus

chrysoprasius (Pallas, 1814) (Actinopterygii, Leuciscidae). *European Journal of Taxonomy*, *861*,

1–20. https://doi.org/10.5852/ejt.2023.861.2061

Bogutskaya, N. G., Jelić, D., Vucić, M., Jelić, M., Diripasko, O. A., Stefanov, T., & Klobučar, G. (2020).

Description of a new species of Phoxinus from the upper Krka river (Adriatic Basin) in Croatia

(Actinopterygii: Leuciscidae), first discovered as a molecular clade. *Journal of Fish Biology*, *96*(2),

378–393. https://doi.org/10.1111/jfb.14210

Böhme, M., Winklhofer, M., & Ilg, A. (2011). Miocene precipitation in Europe: Temporal trends and

spatial gradients. *Palaeogeography, Palaeoclimatology, Palaeoecology*, *304*(3), 212–218.

https://doi.org/10.1016/j.palaeo.2010.09.028

Böhne, A., Zhou, Q., Darras, A., Schmidt, C., Schartl, M., Galiana-Arnoux, D., & Volff, J.-N. (2012).

Zisupton—A Novel Superfamily of DNA Transposable Elements Recently Active in Fish.

Molecular Biology and Evolution, *29*(2), 631–645. https://doi.org/10.1093/molbev/msr208

Brůna, T., Hoff, K. J., Lomsadze, A., Stanke, M., & Borodovsky, M. (2021). BRAKER2: Automatic

eukaryotic genome annotation with GeneMark-EP+ and AUGUSTUS supported by a protein

database. *NAR Genomics and Bioinformatics*, *3*(1), lqaa108.

https://doi.org/10.1093/nargab/lqaa108

Brůna, T., Lomsadze, A., & Borodovsky, M. (2020). GeneMark-EP+: Eukaryotic gene prediction with self-

training in the space of genes and proteins. *NAR Genomics and Bioinformatics*, *2*(2), lqaa026.

https://doi.org/10.1093/nargab/lqaa026

Buchfink, B., Reuter, K., & Drost, H.-G. (2021). Sensitive protein alignments at tree-of-life scale using

DIAMOND. *Nature Methods*, *18*(4), Article 4. https://doi.org/10.1038/s41592-021-01101-x

Buonocore, F., & Gerdol, M. (2016). Alternative adaptive immunity strategies: Coelacanth, cod and shark

immunity. *Molecular Immunology*, *69*, 157–169. https://doi.org/10.1016/j.molimm.2015.09.003

Cantalapiedra, C. P., Hernández-Plaza, A., Letunic, I., Bork, P., & Huerta-Cepas, J. (2021). eggNOG-

mapper v2: Functional Annotation, Orthology Assignments, and Domain Prediction at the

Metagenomic Scale. *Molecular Biology and Evolution*, *38*(12), 5825–5829.

https://doi.org/10.1093/molbev/msab293

Carley, L. N., Morris, W. F., Walsh, R., Riebe, D., & Mitchell-Olds, T. (2022). Are genetic variation and

demographic performance linked? *Evolutionary Applications*, *15*(11), 1888–1906.

https://doi.org/10.1111/eva.13487

Carvalho, C. M. B., & Lupski, J. R. (2016). Mechanisms underlying structural variant formation in genomic

disorders. *Nature Reviews Genetics*, *17*(4), Article 4. https://doi.org/10.1038/nrg.2015.25

Cerbin, S., & Jiang, N. (2018). Duplication of host genes by transposable elements. *Current Opinion in*

Genetics & Development, *49*, 63–69. https://doi.org/10.1016/j.gde.2018.03.005

Chang, M. X. (2023). Emerging mechanisms and functions of inflammasome complexes in teleost fish.

Frontiers in Immunology, *14*.

https://www.frontiersin.org/articles/10.3389/fimmu.2023.1065181

Chen, L., Li, B., Chen, B., Li, C., Zhou, Z., Zhou, T., Yang, W., & Xu, P. (2021). Chromosome-level genome

of Poropuntius huangchuchieni provides a diploid progenitor-like reference genome for the

allotetraploid Cyprinus carpio. *Molecular Ecology Resources*, *21*(5), 1658–1669.

https://doi.org/10.1111/1755-0998.13365

Chen, Z., Omori, Y., Koren, S., Shirokiya, T., Kuroda, T., Miyamoto, A., Wada, H., Fujiyama, A., Toyoda, A.,

Zhang, S., Wolfsberg, T. G., Kawakami, K., Phillippy, A. M., NISC Comparative Sequencing

Program, Mullikin, J. C., & Burgess, S. M. (2019). De novo assembly of the goldfish (Carassius

auratus) genome and the evolution of genes after whole-genome duplication. *Science Advances*,

5(6), eaav0547. https://doi.org/10.1126/sciadv.aav0547

Cheng, H., Concepcion, G. T., Feng, X., Zhang, H., & Li, H. (2021). Haplotype-resolved de novo assembly

using phased assembly graphs with hifiasm. *Nature Methods*, *18*(2), Article 2.

https://doi.org/10.1038/s41592-020-01056-5

Cheng, S.-P., Jia, K.-H., Liu, H., Zhang, R.-G., Li, Z.-C., Zhou, S.-S., Shi, T.-L., Ma, A.-C., Yu, C.-W., Gao, C.,

Cao, G.-L., Zhao, W., Nie, S., Guo, J.-F., Jiao, S.-Q., Tian, X.-C., Yan, X.-M., Bao, Y.-T., Yun, Q.-Z., …

Mao, J.-F. (2021). Haplotype-resolved genome assembly and allele-specific gene expression in

cultivated ginger. *Horticulture Research*, *8*, 188. https://doi.org/10.1038/s41438-021-00599-8

Corral-Lou, A., Perea, S., Aparicio, E., & Doadrio, I. (2019). Phylogeography and species delineation of

the genus Phoxinus Rafinesque, 1820 (Actinopterygii: Leuciscidae) in the Iberian Peninsula.

Journal of Zoological Systematics and Evolutionary Research, *57*(4), 926–941.

https://doi.org/10.1111/jzs.12320

Creevey, C. J., Muller, J., Doerks, T., Thompson, J. D., Arendt, D., & Bork, P. (2011). Identifying Single

Copy Orthologs in Metazoa. *PLOS Computational Biology*, *7*(12), e1002269.

https://doi.org/10.1371/journal.pcbi.1002269

Danecek, P., Bonfield, J. K., Liddle, J., Marshall, J., Ohan, V., Pollard, M. O., Whitwham, A., Keane, T.,

McCarthy, S. A., Davies, R. M., & Li, H. (2021). Twelve years of SAMtools and BCFtools.

GigaScience, *10*(2), giab008. https://doi.org/10.1093/gigascience/giab008

Deininger, P. L., Moran, J. V., Batzer, M. A., & Kazazian, H. H. (2003). Mobile elements and mammalian

genome evolution. *Current Opinion in Genetics & Development*, *13*(6), 651–658.

https://doi.org/10.1016/j.gde.2003.10.013

Demuth, J. P., & Hahn, M. W. (2009). The life and death of gene families. *BioEssays*, *31*(1), 29–39.

https://doi.org/10.1002/bies.080085

Denys, G. P. J., Dettai, A., Persat, H., Daszkiewicz, P., Hautecœur, M., & Keith, P. (2020). *Revision of*

Phoxinus in France with the description of two new species (Teleostei, Leuciscidae).

https://doi.org/10.26028/CYBIUM/2020-443-003

DeWoody, J. A., Harder, A. M., Mathur, S., & Willoughby, J. R. (2021). The long-standing significance of

genetic diversity in conservation. *Molecular Ecology*, *30*(17), 4147–4154.

https://doi.org/10.1111/mec.16051

Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., & Gingeras,

T. R. (2013). STAR: Ultrafast universal RNA-seq aligner. *Bioinformatics*, *29*(1), 15–21.

https://doi.org/10.1093/bioinformatics/bts635

Dumorné, K., Figueroa, E., Cosson, J., Lee-Estevez, M., Ulloa-Rodríguez, P., Valdebenito, I., & Farías, J. G.

(2018). Protein phosphorylation and ions effects on salmonid sperm motility activation. *Reviews*

in Aquaculture, *10*(3), 727–737. https://doi.org/10.1111/raq.12198

Ehlers, J., Astakhov, V., Gibbard, P., Mangerud, J., & Svendsen, J. (2006). Late Pleistocene Glaciations in

Europe. *Encyclopedia of Quaternary Science*, 1085–1095.

Elizalde-Velázquez, A., Crago, J., Zhao, X., Green, M. J., & Cañas-Carrell, J. E. (2020). In vivo effects on the

immune function of fathead minnow (Pimephales promelas) following ingestion and

intraperitoneal injection of polystyrene nanoplastics. *Science of The Total Environment*, *735*,

139461. https://doi.org/10.1016/j.scitotenv.2020.139461

Emms, D. M., & Kelly, S. (2017). STRIDE: Species Tree Root Inference from Gene Duplication Events.

Molecular Biology and Evolution, *34*(12), 3267–3278. https://doi.org/10.1093/molbev/msx259

Emms, D. M., & Kelly, S. (2019). OrthoFinder: Phylogenetic orthology inference for comparative

genomics. *Genome Biology*, *20*(1), 238. https://doi.org/10.1186/s13059-019-1832-y

Esteban, M. Á., Cuesta, A., Chaves-Pozo, E., & Meseguer, J. (2015). Phagocytosis in Teleosts. Implications

of the New Cells Involved. *Biology*, *4*(4), Article 4. https://doi.org/10.3390/biology4040907

Feliner, G. N. (2011). Southern European glacial refugia: A tale of tales. *TAXON*, *60*(2), 365–372.

https://doi.org/10.1002/tax.602007

FERNANDES, J. M. O., KEMP, G. D., MOLLE, M. G., & SMITH, V. J. (2002). Anti-microbial properties of

histone H2A from skin secretions of rainbow trout, Oncorhynchus mykiss. *Biochemical Journal*,

368(2), 611–620. https://doi.org/10.1042/bj20020980

Flajnik, M. F., & Kasahara, M. (2010). Origin and evolution of the adaptive immune system: Genetic

events and selective pressures. *Nature Reviews Genetics*, *11*(1), Article 1.

https://doi.org/10.1038/nrg2703

Frost, W. E. (1943). The Natural History of the Minnow, Phoxinus phoxinus. *Journal of Animal Ecology*,

12(2), 139–162. https://doi.org/10.2307/1374

Gabriel, L., Brůna, T., Hoff, K. J., Ebel, M., Lomsadze, A., Borodovsky, M., & Stanke, M. (2023). *BRAKER3:*

Fully Automated Genome Annotation Using RNA-Seq and Protein Evidence with GeneMark-ETP,

AUGUSTUS and TSEBRA (p. 2023.06.10.544449). bioRxiv.

https://doi.org/10.1101/2023.06.10.544449

Gabriel, L., Hoff, K. J., Brůna, T., Borodovsky, M., & Stanke, M. (2021). TSEBRA: Transcript selector for

BRAKER. *BMC Bioinformatics*, *22*(1), 566. https://doi.org/10.1186/s12859-021-04482-0

Gao, B., Shen, D., Xue, S., Chen, C., Cui, H., & Song, C. (2016). The contribution of transposable elements

to size variations between four teleost genomes. *Mobile DNA*, *7*(1), 4.

https://doi.org/10.1186/s13100-016-0059-7

Goel, M., & Schneeberger, K. (2022). plotsr: Visualizing structural similarities and rearrangements

between multiple genomes. *Bioinformatics*, *38*(10), 2922–2926.

https://doi.org/10.1093/bioinformatics/btac196

Goel, M., Sun, H., Jiao, W.-B., & Schneeberger, K. (2019). SyRI: Finding genomic rearrangements and

local sequence differences from whole-genome assemblies. *Genome Biology*, *20*(1), 277.

https://doi.org/10.1186/s13059-019-1911-0

Guan, D., McCarthy, S. A., Wood, J., Howe, K., Wang, Y., & Durbin, R. (2020). Identifying and removing

haplotypic duplication in primary genome assemblies. *Bioinformatics*, *36*(9), 2896–2898.

https://doi.org/10.1093/bioinformatics/btaa025

Guevara, E. E., Webster, T. H., Lawler, R. R., Bradley, B. J., Greene, L. K., Ranaivonasy, J., Ratsirarson, J.,

Harris, R. A., Liu, Y., Murali, S., Raveendran, M., Hughes, D. S. T., Muzny, D. M., Yoder, A. D.,

Worley, K. C., & Rogers, J. (2021). Comparative genomic analysis of sifakas (Propithecus) reveals

selection for folivory and high heterozygosity despite endangered status. *Science Advances*,

7(17), eabd2274. https://doi.org/10.1126/sciadv.abd2274

Haase, D., Roth, O., Kalbe, M., Schmiedeskamp, G., Scharsack, J. P., Rosenstiel, P., & Reusch, T. B. H.

(2013). Absence of major histocompatibility complex class II mediated immunity in pipefish,

Syngnathus typhle: Evidence from deep transcriptome sequencing. *Biology Letters*, *9*(2),

20130044. https://doi.org/10.1098/rsbl.2013.0044

Haywood, A. M., Sellwood, B. W., & Valdes, P. J. (2000). Regional warming: Pliocene (3 Ma) paleoclimate

of Europe and the Mediterranean. *Geology*, *28*(12), 1063–1066. https://doi.org/10.1130/0091-

7613(2000)28<1063:RWPMPO>2.0.CO;2

Hewitt, G. (2000). The genetic legacy of the Quaternary ice ages. *Nature*, *405*(6789), Article 6789.

https://doi.org/10.1038/35016000

Hoff, K. J., Lange, S., Lomsadze, A., Borodovsky, M., & Stanke, M. (2016). BRAKER1: Unsupervised RNA-

Seq-Based Genome Annotation with GeneMark-ET and AUGUSTUS. *Bioinformatics*, *32*(5), 767–

769. https://doi.org/10.1093/bioinformatics/btv661

Hoff, K. J., Lomsadze, A., Borodovsky, M., & Stanke, M. (2019). Whole-Genome Annotation with BRAKER.

In M. Kollmar (Ed.), *Gene Prediction: Methods and Protocols* (pp. 65–95). Springer.

https://doi.org/10.1007/978-1-4939-9173-0_5

Howe, K., Clark, M. D., Torroja, C. F., Torrance, J., Berthelot, C., Muffato, M., Collins, J. E., Humphray, S.,

McLaren, K., Matthews, L., McLaren, S., Sealy, I., Caccamo, M., Churcher, C., Scott, C., Barrett, J.

C., Koch, R., Rauch, G.-J., White, S., … Stemple, D. L. (2013). The zebrafish reference genome

sequence and its relationship to the human genome. *Nature*, *496*(7446), Article 7446.

https://doi.org/10.1038/nature12111

Huang, H.-R., Liu, X., Arshad, R., Wang, X., Li, W.-M., Zhou, Y., & Ge, X.-J. (2023). Telomere-to-telomere

haplotype-resolved reference genome reveals subgenome divergence and disease resistance in

triploid Cavendish banana. *Horticulture Research*, uhad153.

https://doi.org/10.1093/hr/uhad153

Hubley, R., Finn, R. D., Clements, J., Eddy, S. R., Jones, T. A., Bao, W., Smit, A. F. A., & Wheeler, T. J.

(2016). The Dfam database of repetitive DNA families. *Nucleic Acids Research*, *44*(D1), D81–D89.

https://doi.org/10.1093/nar/gkv1272

Hughes, P. D., Woodward, J. C., & Gibbard, P. L. (2007). Middle Pleistocene cold stage climates in the

Mediterranean: New evidence from the glacial record. *Earth and Planetary Science Letters*,

253(1), 50–56. https://doi.org/10.1016/j.epsl.2006.10.019

Hui, Z., Zhang, J., Ma, Z., Li, X., Peng, T., Li, J., & Wang, B. (2018). Global warming and rainfall: Lessons

from an analysis of Mid-Miocene climate data. *Palaeogeography, Palaeoclimatology,*

Palaeoecology, *512*, 106–117. https://doi.org/10.1016/j.palaeo.2018.10.025

Jansen, W., Kappus, B., Böhmer, J., & Beiter, T. (1999). Fish communities and migrations in the vicinity of

fishways in a regulated river (Enz, Baden-Württemberg, Germany). *Limnologica*, *29*(4), 425–435.

https://doi.org/10.1016/S0075-9511(99)80050-4

Jaramillo-Correa, J. P., Verdú, M., & González-Martínez, S. C. (2010). The contribution of recombination

to heterozygosity differs among plant evolutionary lineages and life-forms. *BMC Evolutionary*

Biology, *10*(1), 22. https://doi.org/10.1186/1471-2148-10-22

Jardim de Queiroz, L., Doenz, C. J., Altermatt, F., Alther, R., Borko, Š., Brodersen, J., Gossner, M. M.,

Graham, C., Matthews, B., McFadden, I. R., Pellissier, L., Schmitt, T., Selz, O. M., Villalba, S.,

Rüber, L., Zimmermann, N. E., & Seehausen, O. (2022). Climate, immigration and speciation

shape terrestrial and aquatic biodiversity in the European Alps. *Proceedings of the Royal Society*

B: Biological Sciences, *289*(1980), 20221020. https://doi.org/10.1098/rspb.2022.1020

Jiang, H., Zhang, X., Leng, L., Gong, D., Zhang, X., Liu, J., Peng, D., Wu, Z., & Yang, Y. (2023). A

chromosome-scale and haplotype-resolved genome assembly of carnation (Dianthus

caryophyllus) based on high-fidelity sequencing. *Frontiers in Plant Science*, *14*, 1230836.

https://doi.org/10.3389/fpls.2023.1230836

Jiang, L., Lin, M., Wang, H., Song, H., Zhang, L., Huang, Q., Chen, R., Song, C., Li, G., & Cao, Y. (2022).

Haplotype-resolved genome assembly of Bletilla striata (Thunb.) Reichb.f. To elucidate

medicinal value. *The Plant Journal*, *111*(5), 1340–1353. https://doi.org/10.1111/tpj.15892

Jones, P. E., Champneys, T., Vevers, J., Börger, L., Svendsen, J. C., Consuegra, S., Jones, J., & Garcia de

Leaniz, C. (2021). Selective effects of small barriers on river-resident fish. *Journal of Applied*

Ecology, *58*(7), 1487–1498. https://doi.org/10.1111/1365-2664.13875

Jurado, J., Fuentes-Almagro, C. A., Guardiola, F. A., Cuesta, A., Esteban, M. Á., & Prieto-Álamo, M.-J.

(2015). Proteomic profile of the skin mucus of farmed gilthead seabream (Sparus aurata).

Journal of Proteomics, *120*, 21–34. https://doi.org/10.1016/j.jprot.2015.02.019

Kajitani, R., Toshimoto, K., Noguchi, H., Toyoda, A., Ogura, Y., Okuno, M., Yabana, M., Harada, M.,

Nagayasu, E., Maruyama, H., Kohara, Y., Fujiyama, A., Hayashi, T., & Itoh, T. (2014). Efficient de

novo assembly of highly heterozygous genomes from whole-genome shotgun short reads.

Genome Research, *24*(8), 1384–1395. https://doi.org/10.1101/gr.170720.113

Karjalainen, J., Urpanen, O., Keskinen, T., Huuskonen, H., Sarvala, J., Valkeajärvi, P., & Marjomäki, T. J.

(2016). Phenotypic plasticity in growth and fecundity induced by strong population fluctuations

affects reproductive traits of female fish. *Ecology and Evolution*, *6*(3), 779–790.

https://doi.org/10.1002/ece3.1936

Kerpedjiev, P., Abdennur, N., Lekschas, F., McCallum, C., Dinkla, K., Strobelt, H., Luber, J. M., Ouellette, S.

B., Azhir, A., Kumar, N., Hwang, J., Lee, S., Alver, B. H., Pfister, H., Mirny, L. A., Park, P. J., &

Gehlenborg, N. (2018). HiGlass: Web-based visual exploration and analysis of genome

interaction maps. *Genome Biology*, *19*(1), 125. https://doi.org/10.1186/s13059-018-1486-1

King, J. L., Marnell, F., Kingston, N., Rosell, R., Boylan, P., Caffrey, J. M., FitzPatrick, Ú., Gargan, P. G.,

Kelly, F. L., O'Grady, M. F., Poole, R., Roche, W. K., & Cassidy, D. (2011). *Ireland Red List No. 5:*

Amphibians, Reptiles & Freshwater Fish (National Parks and Wildlife Service, Department of Arts,

Heritage and the Gaeltacht). https://archive.nationalredlist.org/amphibians-reptiles-freshwater-

fish-in-ireland/

Knebelsberger, T., Dunz, A. R., Neumann, D., & Geiger, M. F. (2015). Molecular diversity of Germany's

freshwater fishes and lampreys assessed by DNA barcoding. *Molecular Ecology Resources*, *15*(3),

562–572. https://doi.org/10.1111/1755-0998.12322

Kojima, K. K. (2019). Structural and sequence diversity of eukaryotic transposable elements. *Genes &*

Genetic Systems, *94*(6), 233–252. https://doi.org/10.1266/ggs.18-00024

Korneliussen, T. S., Albrechtsen, A., & Nielsen, R. (2014). ANGSD: Analysis of Next Generation

Sequencing Data. *BMC Bioinformatics*, *15*(1), 356. https://doi.org/10.1186/s12859-014-0356-4

Kottelat, M. (2007). Three new species of Phoxinus from Greece and southern France (Teleostei:

Cyprinidae). *Ichthyological Exploration of Freshwaters*, *18*(2), 145–162.

Kottelat, M., & Freyhof, J. (2007). *Handbook of European freshwater fishes* (Kottelat, Cornol, & Freyhof,

Eds.). Kottelat.

Krabbenhoft, T. J., MacGuigan, D. J., Backenstose, N. J. C., Waterman, H., Lan, T., Pelosi, J. A., Tan, M., &

Sandve, S. R. (2021). Chromosome-Level Genome Assembly of Chinese Sucker (Myxocyprinus

asiaticus) Reveals Strongly Conserved Synteny Following a Catostomid-Specific Whole-Genome

Duplication. *Genome Biology and Evolution*, *13*(9), evab190.

https://doi.org/10.1093/gbe/evab190

Kronenberg, Z. N., Rhie, A., Koren, S., Concepcion, G. T., Peluso, P., Munson, K. M., Porubsky, D., Kuhn,

K., Mueller, K. A., Low, W. Y., Hiendleder, S., Fedrigo, O., Liachko, I., Hall, R. J., Phillippy, A. M.,

Eichler, E. E., Williams, J. L., Smith, T. P. L., Jarvis, E. D., … Kingan, S. B. (2021). Extended

haplotype-phasing of long-read de novo genome assemblies using Hi-C. *Nature Communications*,

12(1), Article 1. https://doi.org/10.1038/s41467-020-20536-y

Kumar, S., Stecher, G., Suleski, M., & Hedges, S. B. (2017). TimeTree: A Resource for Timelines,

Timetrees, and Divergence Times. *Molecular Biology and Evolution*, *34*(7), 1812–1819.

https://doi.org/10.1093/molbev/msx116

Lai, Y.-T., Yeung, C. K. L., Omland, K. E., Pang, E.-L., Hao, Y., Liao, B.-Y., Cao, H.-F., Zhang, B.-W., Yeh, C.-F.,

Hung, C.-M., Hung, H.-Y., Yang, M.-Y., Liang, W., Hsu, Y.-C., Yao, C.-T., Dong, L., Lin, K., & Li, S.-H.

(2019). Standing genetic variation as the predominant source for adaptation of a songbird.

Proceedings of the National Academy of Sciences, *116*(6), 2152–2157.

https://doi.org/10.1073/pnas.1813597116

Landesamt für Natur, Umwelt und Verbraucherschutz Nordrhein-Westfalen. (2020).

Fachinformationssystem FischInfo [dataset].

Leitwein, M., Gagnaire, P.-A., Desmarais, E., Guendouz, S., Rohmer, M., Berrebi, P., & Guinand, B. (2016).

Genome-wide nucleotide diversity of hatchery-reared Atlantic and Mediterranean strains of

brown trout Salmo trutta compared to wild Mediterranean populations. *Journal of Fish Biology*,

89(6), 2717–2734. https://doi.org/10.1111/jfb.13131

Li, H. (2013). Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. *arXiv*,

1303.3997 [q-bio.GN].

Li, H. (2018). Minimap2: Pairwise alignment for nucleotide sequences. *Bioinformatics (Oxford, England)*,

34(18), 3094–3100. https://doi.org/10.1093/bioinformatics/bty191

Li, H., & Durbin, R. (2011). Inference of human population history from individual whole-genome

sequences. *Nature*, *475*(7357), Article 7357. https://doi.org/10.1038/nature10231

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., Durbin, R., &

1000 Genome Project Data Processing Subgroup. (2009). The Sequence Alignment/Map format

and SAMtools. *Bioinformatics*, *25*(16), 2078–2079.

https://doi.org/10.1093/bioinformatics/btp352

Li, J., Bian, C., Yi, Y., Yu, H., You, X., & Shi, Q. (2021). Temporal dynamics of teleost populations during

the Pleistocene: A report from publicly available genome data. *BMC Genomics*, *22*(1), 490.

https://doi.org/10.1186/s12864-021-07816-7

Li, Q., Jiang, B., Zhang, Z., Huang, Y., Xu, Z., Chen, X., Huang, Y., Jian, J., & Yan, Q. (2022). Involvement

and characterization of NLRCs and pyroptosis-related genes in Nile tilapia (Oreochromis niloticus)

immune response. *Fish & Shellfish Immunology*, *130*, 602–611.

https://doi.org/10.1016/j.fsi.2022.09.041

Li, X., Ye, Y., Peng, K., Zeng, Z., Chen, L., & Zeng, Y. (2022). Histones: The critical players in innate

immunity. *Frontiers in Immunology*, *13*.

https://www.frontiersin.org/articles/10.3389/fimmu.2022.1030610

Li, Z.-H., Chen, L., Wu, Y.-H., Li, P., Li, Y.-F., & Ni, Z.-H. (2014). Effects of Mercury on Oxidative Stress and

Gene Expression of Potential Biomarkers in Larvae of the Chinese Rare Minnow Gobiocypris

Rarus. *Archives of Environmental Contamination and Toxicology*, *67*(2), 245–251.

https://doi.org/10.1007/s00244-014-0034-6

Lin, D., Hong, P., Zhang, S., Xu, W., Jamal, M., Yan, K., Lei, Y., Li, L., Ruan, Y., Fu, Z. F., Li, G., & Cao, G.

(2018). Digestion-ligation-only Hi-C is an efficient and cost-effective method for chromosome

conformation capture. *Nature Genetics*, *50*(5), 754–763. https://doi.org/10.1038/s41588-018-

0111-2

Litman, G. W., Rast, J. P., & Fugmann, S. D. (2010). The origins of vertebrate adaptive immunity. *Nature*

Reviews Immunology, *10*(8), Article 8. https://doi.org/10.1038/nri2807

Liu, G., Xin, S., Geng, S., Zheng, W., Xu, T., & Sun, Y. (2023). Identification of a novel fusion gene NLRC3-

NLRP12 in miiuy croaker (Miichthys miiuy). *Fish & Shellfish Immunology*, *136*, 108697.

https://doi.org/10.1016/j.fsi.2023.108697

Liu, H., Chen, C., Lv, M., Liu, N., Hu, Y., Zhang, H., Enbody, E. D., Gao, Z., Andersson, L., & Wang, W.

(2021). A Chromosome-Level Assembly of Blunt Snout Bream (Megalobrama amblycephala)

Genome Reveals an Expansion of Olfactory Receptor Genes in Freshwater Fish. *Molecular*

Biology and Evolution, *38*(10), 4238–4251. https://doi.org/10.1093/molbev/msab152

Low, W. Y., Tearle, R., Liu, R., Koren, S., Rhie, A., Bickhart, D. M., Rosen, B. D., Kronenberg, Z. N., Kingan,

S. B., Tseng, E., Thibaud-Nissen, F., Martin, F. J., Billis, K., Ghurye, J., Hastie, A. R., Lee, J., Pang, A.

W. C., Heaton, M. P., Phillippy, A. M., … Williams, J. L. (2020). Haplotype-resolved genomes

provide insights into structural variation and gene content in Angus and Brahman cattle. *Nature*

Communications, *11*(1), Article 1. https://doi.org/10.1038/s41467-020-15848-y

Malmstrøm, M., Matschiner, M., Tørresen, O. K., Star, B., Snipen, L. G., Hansen, T. F., Baalsrud, H. T.,

Nederbragt, A. J., Hanel, R., Salzburger, W., Stenseth, N. C., Jakobsen, K. S., & Jentoft, S. (2016).

Evolution of the immune system influences speciation rates in teleost fishes. *Nature Genetics*,

48(10), Article 10. https://doi.org/10.1038/ng.3645

Mansfeld, B. N., Boyher, A., Berry, J. C., Wilson, M., Ou, S., Polydore, S., Michael, T. P., Fahlgren, N., &

Bart, R. S. (2021). Large structural variations in the haplotype-resolved African cassava genome.

The Plant Journal, *108*(6), 1830–1848. https://doi.org/10.1111/tpj.15543

Martinson, J. W., Bencic, D. C., Toth, G. P., Kostich, M. S., Flick, R. W., See, M. J., Lattier, D., Biales, A. D.,

& Huang, W. (2022). De Novo Assembly of the Nearly Complete Fathead Minnow Reference

Genome Reveals a Repetitive but Compact Genome. *Environmental Toxicology and Chemistry*,

41(2), 448–461. https://doi.org/10.1002/etc.5266

Matuszewski, S., Hermisson, J., & Kopp, M. (2015). Catch Me if You Can: Adaptation from Standing

Genetic Variation to a Moving Phenotypic Optimum. *Genetics*, *200*(4), 1255–1274.

https://doi.org/10.1534/genetics.115.178574

Mendes, F. K., Vanderpool, D., Fulton, B., & Hahn, M. W. (2021). CAFE 5 models variation in evolutionary

rates among gene families. *Bioinformatics*, *36*(22–23), 5516–5518.

https://doi.org/10.1093/bioinformatics/btaa1022

Mérot, C., Oomen, R. A., Tigano, A., & Wellenreuther, M. (2020). A Roadmap for Understanding the

Evolutionary Significance of Structural Genomic Variation. *Trends in Ecology & Evolution*, *35*(7),

561–572. https://doi.org/10.1016/j.tree.2020.03.002

Miller, J. M., Malenfant, R. M., David, P., Davis, C. S., Poissant, J., Hogg, J. T., Festa-Bianchet, M., &

Coltman, D. W. (2014). Estimating genome-wide heterozygosity: Effects of demographic history

and marker type. *Heredity*, *112*(3), Article 3. https://doi.org/10.1038/hdy.2013.99

Mills, C. A. (1987). The life history of the minnow Phoxinus phoxinus (L.) in a productive stream.

Freshwater Biology, *17*(1), 53–67. https://doi.org/10.1111/j.1365-2427.1987.tb01028.x

Mills, C. A. (1988). The effect of extreme northerly climatic conditions on the life history of the minnow,

Phoxinus phoxinus (L.). *Journal of Fish Biology*, *33*(4), 545–561. https://doi.org/10.1111/j.1095-

8649.1988.tb05498.x

Moiseenko, T. I., Sharov, A. N., Vandish, O. I., Kudryavtseva, L. P., Gashkina, N. A., & Rose, C. (2009).

Long-term modification of Arctic lake ecosystems: Reference condition, degradation under toxic

impacts and recovery (case study Imandra Lakes, Russia). *Limnologica*, *39*(1), 1–13.

https://doi.org/10.1016/j.limno.2008.03.003

Morgulis, A., Gertz, E. M., Schäffer, A. A., & Agarwala, R. (2006). A Fast and Symmetric DUST

Implementation to Mask Low-Complexity DNA Sequences. *Journal of Computational Biology*,

13(5), 1028–1040. https://doi.org/10.1089/cmb.2006.13.1028

Morimoto, N., Kono, T., Sakai, M., & Hikima, J. (2021). Inflammasomes in Teleosts: Structures and

Mechanisms That Induce Pyroptosis during Bacterial Infection. *International Journal of*

Molecular Sciences, *22*(9), Article 9. https://doi.org/10.3390/ijms22094389

Munnecke, A., Calner, M., Harper, D. A. T., & Servais, T. (2010). Ordovician and Silurian sea–water

chemistry, sea level, and climate: A synopsis. *Palaeogeography, Palaeoclimatology,*

Palaeoecology, *296*(3), 389–413. https://doi.org/10.1016/j.palaeo.2010.08.001

Museth, J., Borgstrøm, R., & Brittain, J. E. (2010). Diet overlap between introduced European minnow

(Phoxinus phoxinus) and young brown trout (Salmo trutta) in the lake, Øvre Heimdalsvatn: A

result of abundant resources or forced niche overlap? *Hydrobiologia*, *642*(1), 93–100.

https://doi.org/10.1007/s10750-010-0162-6

Nachman, M. W. (2002). Variation in recombination rate across the genome: Evidence and implications.

Current Opinion in Genetics & Development, *12*(6), 657–663. https://doi.org/10.1016/S0959-

437X(02)00358-1

Nadachowska-Brzyska, K., Burri, R., Smeds, L., & Ellegren, H. (2016). PSMC analysis of effective

population sizes in molecular ecology and its application to black-and-white Ficedula flycatchers.

Molecular Ecology, *25*(5), 1058–1072. https://doi.org/10.1111/mec.13540

Nadachowska-Brzyska, K., Li, C., Smeds, L., Zhang, G., & Ellegren, H. (2015). Temporal Dynamics of Avian

Populations during Pleistocene Revealed by Whole-Genome Sequences. *Current Biology*, *25*(10),

1375–1380. https://doi.org/10.1016/j.cub.2015.03.047

Nakandala, U., Masouleh, A. K., Smith, M. W., Furtado, A., Mason, P., Constantin, L., & Henry, R. J.

(2023). Haplotype resolved chromosome level genome assembly of Citrus australis reveals

disease resistance and other citrus specific genes. *Horticulture Research*, *10*(5), uhad058.

https://doi.org/10.1093/hr/uhad058

Neumann, N. F., Stafford, J. L., Barreda, D., Ainsworth, A. J., & Belosevic, M. (2001). Antimicrobial

mechanisms of fish phagocytes and their role in host defense. *Developmental & Comparative*

Immunology, *25*(8), 807–825. https://doi.org/10.1016/S0145-305X(01)00037-4

Nigam, A. K., Kumari, U., Mittal, S., & Mittal, A. K. (2017). Evaluation of antibacterial activity and innate

immune components in skin mucus of Indian major carp, Cirrhinus mrigala. *Aquaculture*

Research, *48*(2), 407–418. https://doi.org/10.1111/are.12889

O'Leary, N. A., Wright, M. W., Brister, J. R., Ciufo, S., Haddad, D., McVeigh, R., Rajput, B., Robbertse, B.,

Smith-White, B., Ako-Adjei, D., Astashyn, A., Badretdin, A., Bao, Y., Blinkova, O., Brover, V.,

Chetvernin, V., Choi, J., Cox, E., Ermolaeva, O., … Pruitt, K. D. (2016). Reference sequence

(RefSeq) database at NCBI: Current status, taxonomic expansion, and functional annotation.

Nucleic Acids Research, *44*(Database issue), D733–D745. https://doi.org/10.1093/nar/gkv1189

Oscoz, J., Leunda, P. M., Campos, F., Escala, M. C., García-Fresca, C., & Miranda, R. (2005). Spring diet

composition of Rainbow Trout, Oncorhynchus mykiss (Walbaum, 1792) in the Urederra river

(Spain). *Annales de Limnologie - International Journal of Limnology*, *41*(1), 27–34.

https://doi.org/10.1051/limn/2005003

Palandačić, A., Bravničar, J., Zupančič, P., Šanda, R., & Snoj, A. (2015). Molecular data suggest a

multispecies complex of Phoxinus (Cyprinidae) in the Western Balkan Peninsula. *Molecular*

Phylogenetics and Evolution, *92*, 118–123. https://doi.org/10.1016/j.ympev.2015.05.024

Palandačić, A., Kruckenhauser, L., Ahnelt, H., & Mikschi, E. (2020). European minnows through time:

Museum collections aid genetic assessment of species introductions in freshwater fishes

(Cyprinidae: Phoxinus species complex). *Heredity*. https://doi.org/10.1038/s41437-019-0292-1

Palandačić, A., Naseka, A., Ramler, D., & Ahnelt, H. (2017). Contrasting morphology with molecular data:

An approach to revision of species complexes based on the example of European Phoxinus

(Cyprinidae). *BMC Evolutionary Biology*, *17*(1), 184. https://doi.org/10.1186/s12862-017-1032-x

Palandačić, A., Witman, K., & Spikmans, F. (2022). Molecular analysis reveals multiple native and alien

Phoxinus species (Leusciscidae) in the Netherlands and Belgium. *Biological Invasions*, *24*(8),

2273–2283. https://doi.org/10.1007/s10530-022-02784-9

Pasupuleti, M., Schmidtchen, A., & Malmsten, M. (2012). Antimicrobial peptides: Key components of the

innate immune system. *Critical Reviews in Biotechnology*, *32*(2), 143–171.

https://doi.org/10.3109/07388551.2011.594423

Patel, D. M., & Brinchmann, Monica. F. (2017). Skin mucus proteins of lumpsucker (Cyclopterus lumpus).

Biochemistry and Biophysics Reports, *9*, 217–225. https://doi.org/10.1016/j.bbrep.2016.12.016

Pavlov, D. S., & Mikheev, V. N. (2017). Downstream migration and mechanisms of dispersal of young fish

in rivers. *Canadian Journal of Fisheries and Aquatic Sciences*, *74*(8), 1312–1323.

https://doi.org/10.1139/cjfas-2016-0298

Poplin, R., Chang, P.-C., Alexander, D., Schwartz, S., Colthurst, T., Ku, A., Newburger, D., Dijamco, J.,

Nguyen, N., Afshar, P. T., Gross, S. S., Dorfman, L., McLean, C. Y., & DePristo, M. A. (2018). A

universal SNP and small-indel variant caller using deep neural networks. *Nature Biotechnology*,

36(10), Article 10. https://doi.org/10.1038/nbt.4235

Pryszcz, L. P., & Gabaldón, T. (2016). Redundans: An assembly pipeline for highly heterozygous genomes.

Nucleic Acids Research, *44*(12), e113. https://doi.org/10.1093/nar/gkw294

Quinlan, A. R. (2014). BEDTools: The Swiss-Army Tool for Genome Feature Analysis. *Current Protocols in*

Bioinformatics, *47*(1), 11.12.1-11.12.34. https://doi.org/10.1002/0471250953.bi1112s47

Ranallo-Benavidez, T. R., Jaron, K. S., & Schatz, M. C. (2020). GenomeScope 2.0 and Smudgeplot for

reference-free profiling of polyploid genomes. *Nature Communications*, *11*(1), Article 1.

https://doi.org/10.1038/s41467-020-14998-3

Raudvere, U., Kolberg, L., Kuzmin, I., Arak, T., Adler, P., Peterson, H., & Vilo, J. (2019). g:Profiler: A web

server for functional enrichment analysis and conversions of gene lists (2019 update). *Nucleic*

Acids Research, *47*(W1), W191–W198. https://doi.org/10.1093/nar/gkz369

Reier, S., Kruckenhauser, L., Snoj, A., Trontelj, P., & Palandačić, A. (2022). The minnow *Phoxinus*

lumaireul (Leuciscidae) shifts the Adriatic–Black Sea basin divide in the north-western Dinaric

Karst region. *Ecohydrology*, *15*(6). https://doi.org/10.1002/eco.2449

Reijnders, M. J. M. F., & Waterhouse, R. M. (2021). Summary Visualizations of Gene Ontology Terms

With GO-Figure! *Frontiers in Bioinformatics*, *1*.

https://www.frontiersin.org/articles/10.3389/fbinf.2021.638255

Röpke, C., Pires, T. H. S., Zuanon, J., Freitas, C. E. C., Hernandes, M. C., Souza, F., & Amadio, S. (2021).

Growth–reproduction trade-off and fecundity regulate population stability in Amazon floodplain

fishes. *Freshwater Biology*, *66*(6), 1101–1109. https://doi.org/10.1111/fwb.13702

Rose, J. (2009). Early and Middle Pleistocene landscapes of eastern England. *Proceedings of the*

Geologists' Association, *120*(1), 3–33. https://doi.org/10.1016/j.pgeola.2009.05.003

Ruddiman, W. F., Raymo, M. E., Martinson, D. G., Clement, B. M., & Backman, J. (1989). Pleistocene

evolution: Northern hemisphere ice sheets and North Atlantic Ocean. *Paleoceanography*, *4*(4),

353–412. https://doi.org/10.1029/PA004i004p00353

Shao, F., Han, M., & Peng, Z. (2019). Evolution and diversity of transposable elements in fish genomes.

Scientific Reports, *9*(1), Article 1. https://doi.org/10.1038/s41598-019-51888-1

Simão, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V., & Zdobnov, E. M. (2015). BUSCO:

Assessing genome assembly and annotation completeness with single-copy orthologs.

Bioinformatics, *31*(19), 3210–3212. https://doi.org/10.1093/bioinformatics/btv351

Sotero-Caio, C. G., Platt, R. N., II, Suh, A., & Ray, D. A. (2017). Evolution and Diversity of Transposable

Elements in Vertebrate Genomes. *Genome Biology and Evolution*, *9*(1), 161–177.

https://doi.org/10.1093/gbe/evw264

Stanke, M., Keller, O., Gunduz, I., Hayes, A., Waack, S., & Morgenstern, B. (2006). AUGUSTUS: Ab initio

prediction of alternative transcripts. *Nucleic Acids Research*, *34*(suppl_2), W435–W439.

https://doi.org/10.1093/nar/gkl200

Star, B., Nederbragt, A. J., Jentoft, S., Grimholt, U., Malmstrøm, M., Gregers, T. F., Rounge, T. B., Paulsen,

J., Solbakken, M. H., Sharma, A., Wetten, O. F., Lanzén, A., Winer, R., Knight, J., Vogel, J.-H., Aken,

B., Andersen, Ø., Lagesen, K., Tooming-Klunderud, A., … Jakobsen, K. S. (2011). The genome

sequence of Atlantic cod reveals a unique immune system. *Nature*, *477*(7363), 207–210.

https://doi.org/10.1038/nature10342

Stein, C., Caccamo, M., Laird, G., & Leptin, M. (2007). Conservation and divergence of gene families

encoding components of innate immune response systems in zebrafish. *Genome Biology*, *8*(11),

R251. https://doi.org/10.1186/gb-2007-8-11-r251

Steinthorsdottir, M., Coxall, H. K., de Boer, A. M., Huber, M., Barbolini, N., Bradshaw, C. D., Burls, N. J.,

Feakins, S. J., Gasson, E., Henderiks, J., Holbourn, A. E., Kiel, S., Kohn, M. J., Knorr, G., Kürschner,

W. M., Lear, C. H., Liebrand, D., Lunt, D. J., Mörs, T., … Strömberg, C. a. E. (2021). The Miocene:

The Future of the Past. *Paleoceanography and Paleoclimatology*, *36*(4), e2020PA004037.

https://doi.org/10.1029/2020PA004037

Sun, H., Jiao, W.-B., Krause, K., Campoy, J. A., Goel, M., Folz-Donahue, K., Kukat, C., Huettel, B., &

Schneeberger, K. (2022). Chromosome-scale and haplotype-resolved genome assembly of a

tetraploid potato cultivar. *Nature Genetics*, *54*(3), Article 3. https://doi.org/10.1038/s41588-

022-01015-0

Sun, L., Gao, T., Wang, F., Qin, Z., Yan, L., Tao, W., Li, M., Jin, C., Ma, L., Kocher, T. D., & Wang, D. (2020).

Chromosome-level genome assembly of a cyprinid fish Onychostoma macrolepis by integration

of nanopore sequencing, Bionano and Hi-C technology. *Molecular Ecology Resources*, *20*(5),

1361–1371. https://doi.org/10.1111/1755-0998.13190

Sun, Y., Yin, Q., Crucifix, M., Clemens, S. C., Araya-Melo, P., Liu, W., Qiang, X., Liu, Q., Zhao, H., Liang, L.,

Chen, H., Li, Y., Zhang, L., Dong, G., Li, M., Zhou, W., Berger, A., & An, Z. (2019). Diverse

manifestations of the mid-Pleistocene climate transition. *Nature Communications*, *10*(1), Article

1. https://doi.org/10.1038/s41467-018-08257-9

Supek, F., Bošnjak, M., Škunca, N., & Šmuc, T. (2011). REVIGO Summarizes and Visualizes Long Lists of

Gene Ontology Terms. *PLOS ONE*, *6*(7), e21800. https://doi.org/10.1371/journal.pone.0021800

Svirgsden, R., Rohtla, M., Albert, A., Taal, I., Saks, L., Verliin, A., & Vetemaa, M. (2018). Do Eurasian

minnows (Phoxinus phoxinus L.) inhabiting brackish water enter fresh water to reproduce:

Evidence from a study on otolith microchemistry. *Ecology of Freshwater Fish*, *27*(1), 89–97.

https://doi.org/10.1111/eff.12326

Takeuchi, T., Suzuki, Y., Watabe, S., Nagai, K., Masaoka, T., Fujie, M., Kawamitsu, M., Satoh, N., & Myers,

E. W. (2022). A high-quality, haplotype-phased genome reconstruction reveals unexpected

haplotype diversity in a pearl oyster. *DNA Research*, *29*(6), dsac035.

https://doi.org/10.1093/dnares/dsac035

Tang, S., Lomsadze, A., & Borodovsky, M. (2015). Identification of protein coding regions in RNA

transcripts. *Nucleic Acids Research*, *43*(12), e78. https://doi.org/10.1093/nar/gkv227

Tarasov, A., Vilella, A. J., Cuppen, E., Nijman, I. J., & Prins, P. (2015). Sambamba: Fast processing of NGS

alignment formats. *Bioinformatics*, *31*(12), 2032–2034.

https://doi.org/10.1093/bioinformatics/btv098

Tekaia, F. (2016). Inferring Orthologs: Open Questions and Perspectives. *Genomics Insights*, *9*,

GEI.S37925. https://doi.org/10.4137/GEI.S37925

Tian, F., Liu, S., Zhou, B., Tang, Y., Zhang, Y., Zhang, C., & Zhao, K. (2022). Chromosome-level genome of

Tibetan naked carp (Gymnocypris przewalskii) provides insights into Tibetan highland

adaptation. *DNA Research*, *29*(4), dsac025. https://doi.org/10.1093/dnares/dsac025

Tigano, A., Jacobs, A., Wilder, A. P., Nand, A., Zhan, Y., Dekker, J., & Therkildsen, N. O. (2021).

Chromosome-Level Assembly of the Atlantic Silverside Genome Reveals Extreme Levels of

Sequence Diversity and Structural Genetic Variation. *Genome Biology and Evolution*, *13*(6),

evab098. https://doi.org/10.1093/gbe/evab098

Vucić, M., Jelić, D., Žutinić, P., Grandjean, F., & Jelić, M. (2018). Distribution of Eurasian minnows

(Phoxinus: Cypriniformes) in the Western Balkans. *Knowledge & Management of Aquatic

Ecosystems*, *419*, 11. https://doi.org/10.1051/kmae/2017051

Waldbieser, G. C., Liu, S., Yuan, Z., Older, C. E., Gao, D., Shi, C., Bosworth, B. G., Li, N., Bao, L., Kirby, M.

A., Jin, Y., Wood, M. L., Scheffler, B., Simpson, S., Youngblood, R. C., Duke, M. V., Ballard, L.,

Phillippy, A., Koren, S., & Liu, Z. (2023). Reference genomes of channel catfish and blue catfish

reveal multiple pericentric chromosome inversions. *BMC Biology*, *21*(1), 67.

https://doi.org/10.1186/s12915-023-01556-8

Wang, F., Wang, L., Liu, D., Gao, Q., Nie, M., Zhu, S., Chao, Y., Yang, C., Zhang, C., Yi, R., Ni, W., Tian, F.,

Zhao, K., & Qi, D. (2022). Chromosome-level assembly of Gymnocypris eckloni genome.

Scientific Data, *9*(1), Article 1. https://doi.org/10.1038/s41597-022-01595-w

Wang, Q., Luo, S., Ghonimy, A., Chen, Y., Guo, Z., Liu, H., & Zhang, D. (2019). Effect of dietary l-carnitine

on growth performance and antioxidant response in Amur minnow (Phoxinus lagowskii

Dybowskii). *Aquaculture Nutrition*, *25*(4), 749–760. https://doi.org/10.1111/anu.12892

Wang, T., Liu, F., Tian, G., Secombes, C. J., & Wang, T. (2019). Lineage/species-specific expansion of the

Mx gene family in teleosts: Differential expression and modulation of nine Mx genes in rainbow

trout Oncorhynchus mykiss. *Fish & Shellfish Immunology*, *90*, 413–430.

https://doi.org/10.1016/j.fsi.2019.04.303

Watanabe, K., Koga, H., Nakamura, K., Fujita, A., Hattori, A., Matsuda, M., & Koga, A. (2014).

Spontaneous germline excision of Tol1, a DNA-based transposable element naturally occurring

in the medaka fish genome. *Genome*, *57*(4), 193–199. https://doi.org/10.1139/gen-2014-0011

Weischenfeldt, J., Symmons, O., Spitz, F., & Korbel, J. O. (2013). Phenotypic impact of genomic structural

variation: Insights from and for human disease. *Nature Reviews Genetics*, *14*(2), Article 2.

https://doi.org/10.1038/nrg3373

Wellenreuther, M., Mérot, C., Berdan, E., & Bernatchez, L. (2019). Going beyond SNPs: The role of

structural genomic variants in adaptive evolution and species diversification. *Molecular Ecology*,

28(6), 1203–1209. https://doi.org/10.1111/mec.15066

Westman, J., Papareddy, P., Dahlgren, M. W., Chakrakodi, B., Norrby-Teglund, A., Smeds, E., Linder, A.,

Mörgelin, M., Johansson-Lindbom, B., Egesten, A., & Herwald, H. (2015). Extracellular Histones

Induce Chemokine Production in Whole Blood Ex Vivo and Leukocyte Recruitment In Vivo. *PLoS*

Pathogens, *11*(12), e1005319. https://doi.org/10.1371/journal.ppat.1005319

Wu, C.-S., Ma, Z.-Y., Zheng, G.-D., Zou, S.-M., Zhang, X.-J., & Zhang, Y.-A. (2022). Chromosome-level

genome assembly of grass carp (Ctenopharyngodon idella) provides insights into its genome

evolution. *BMC Genomics*, *23*(1), 271. https://doi.org/10.1186/s12864-022-08503-x

Xu, P., Zhang, X., Wang, X., Li, J., Liu, G., Kuang, Y., Xu, J., Zheng, X., Ren, L., Wang, G., Zhang, Y., Huo, L.,

Zhao, Z., Cao, D., Lu, C., Li, C., Zhou, Y., Liu, Z., Fan, Z., … Sun, X. (2014). Genome sequence and

genetic diversity of the common carp, Cyprinus carpio. *Nature Genetics*, *46*(11), Article 11.

https://doi.org/10.1038/ng.3098

Yan, A., & Yu, H. (2022). New insights into centromeres from Arabidopsis Col-CEN assembly. *Trends in*

Genetics, *38*(5), 416–418. https://doi.org/10.1016/j.tig.2022.02.001

Yang, J., Chen, X., Bai, J., Fang, D., Qiu, Y., Jiang, W., Yuan, H., Bian, C., Lu, J., He, S., Pan, X., Zhang, Y.,

Wang, X., You, X., Wang, Y., Sun, Y., Mao, D., Liu, Y., Fan, G., … Shi, Q. (2016). The

Sinocyclocheilus cavefish genome provides insights into cave adaptation. *BMC Biology*, *14*(1), 1.

https://doi.org/10.1186/s12915-015-0223-4

Yang, X., Liu, H., Ma, Z., Zou, Y., Zou, M., Mao, Y., Li, X., Wang, H., Chen, T., Wang, W., & Yang, R. (2019).

Chromosome-level genome assembly of Triplophysa tibetana, a fish adapted to the harsh high-

altitude environment of the Tibetan Plateau. *Molecular Ecology Resources*, *19*(4), 1027–1036.

https://doi.org/10.1111/1755-0998.13021

Yao, T., Wang, S., Xu, K., Zhang, X., Zhang, T., Wang, J., Wang, Z., & Mu, W. (2022). Biochemical,

histological, and gene expression analysis in brain and heart in Phoxinus lagowskii under

sustained and diel-cycling hypoxia. *Journal of the World Aquaculture Society*, *53*(4), 860–878.

https://doi.org/10.1111/jwas.12901

Yuan, Z., Huang, W., Liu, S., Xu, P., Dunham, R., & Liu, Z. (2018). Historical demography of common carp

estimated from individuals collected from various parts of the world using the pairwise

sequentially markovian coalescent approach. *Genetica*, *146*(2), 235–241.

https://doi.org/10.1007/s10709-017-0006-7

Zeng, Q., Zhou, Z., He, Q., Li, L., Pu, F., Yan, M., & Xu, P. (2023). Chromosome-level haplotype-resolved

genome assembly for Takifugu ocellatus using PacBio and Hi-C technologies. *Scientific Data*,

10(1), Article 1. https://doi.org/10.1038/s41597-023-01937-2

Zhou, C., McCarthy, S., & Durbin, R. (2022). *YaHS: Yet another Hi-C scaffolding too*.

https://doi.org/10.1101/2022.06.09.495093

Zilli, L., Schiavone, R., Storelli, C., & Vilella, S. (2008). Molecular Mechanisms Determining Sperm Motility

Initiation in Two Sparids (Sparus aurata and Lithognathus mormyrus). *Biology of Reproduction*,

79(2), 356–366. https://doi.org/10.1095/biolreprod.108.068296

Zilli, L., Schiavone, R., & Vilella, S. (2017). Role of protein phosphorylation/dephosphorylation in fish

sperm motility activation: State of the art and perspectives. *Aquaculture*, *472*, 73–80.

https://doi.org/10.1016/j.aquaculture.2016.03.04

1220

1221

<p align="center">**Appendix**</p>

1223 Supplementary figures S1 - S8 and tables S1 - S5: https://doi.org/10.5281/zenodo.10221159

1224

Supplementary tables S6 - S15: https://doi.org/10.5281/zenodo.10220266

Supplementary figures

- Figure S1: Heatmap of haplome 1 Hi-C assembly with darker blocks indicating higher intensity of
sequence interaction
- Figure S2: Heatmap of haplome 2 Hi-C assembly with darker blocks indicating higher intensity of
sequence interaction -
- Figure S3: GenomeScope plots of 19, 23, 25 and 30 mer analysis
- Figure S4: Repeats landscape of annotated repeats in Haplotype 2
- Figure S5: Shared homologs between Phoxinus phoxinus and zebrafish, fathead minnow,
goldfish, grass carp and common carp
- Figure S6: Gene ontology of genes in Insertions/Deletions
- Figure S7: Gene ontology of genes in Inversions
- Figure S8: Expansion/contraction profile including all gene families (significant and non-
significant families)

1239

Supplementary tables
- Table S1: Summary statistics of 19, 23, 25 and 30 mer genome profiles
- Table S2: Genome assembly statistics of 25 chromosomes of both haplomes
- Table S3: Statistics of repeat sequences annotated in haplome 1
- Table S4: Statistics of repeat sequences annotated in haplome 2
- Table S5: Mapping statistics of RNA sequence data used for protein annotation in both
haplomes
- Table S6: Summary statistics of annotated proteins in haplome 1
- Table S7: Summary statistics of annotated proteins in haplome 2
- Table S8: List of genes present in insertion/deletions
- Table S9: Gene ontology of genes present in insertions/deletions
- Table S10: List of genes present in inversions
- Table S11: Gene ontology of genes present in inversions
- Table S12: Summary statistics of orthology between *Phoxinus phoxinus* proteomes and selected
teleosts
- Table S13: List of genes identified by orthology analysis as *Phoxinus phoxinus*-specific
- Table S14: List of genes significantly expanding in *Phoxinus phoxinus* genome
- Table S15: List of genes significantly contracting in *Phoxinus phoxinus* genome

1258