Time-of-day effects of drugs revealed by high-throughput deep phenotyping

Running title: Time-of-day effects of drugs

Authors: Carolin Ector[1,2], Christoph Schmal[3], Jeff Didier[4], Sébastien De Landtsheer[4], Anna-Marie Finger[5,6], Francesca Müller-Marquardt[1,7], Johannes Schulte[8,9], Thomas Sauter[4], Ulrich Keilholz[1,10], Hanspeter Herzel[3,11], Achim Kramer[5], Adrián E. Granada*[1,10]

First author: Carolin Ector || **Corresponding author:** adrian.granada@charite.de

Affiliations:
1. Charité Comprehensive Cancer Center, Charité – Universitätsmedizin Berlin, Charitéplatz 1, 10117 Berlin, Germany.
2. Faculty of Life Sciences, Humboldt-Universität zu Berlin, Invalidenstraße 42, 10115 Berlin, Germany.
3. Institute for Theoretical Biology, Humboldt-Universität zu Berlin, Philippstraße 13, 10115 Berlin, Germany.
4. Department of Life Sciences and Medicine, University of Luxembourg, 2, Avenue de l'Université, L-4365 Esch-sur-Alzette, Luxembourg.
5. Institute for Medical Immunology, Charité – Universitätsmedizin Berlin, Charitéplatz 1, 10117 Berlin, Germany.
6. Present address: Department of Anatomy, University of California, San Francisco, 513 Parnassus Avenue HSW 1330, San Francisco, CA 94143-0452, USA.
7. Present address: Institute of Research for Development, University of Montpellier, 911 Av. Agropolis, 34394 Montpellier, France.
8. Department of Pediatric Oncology, Hematology and Stem Cell Transplantation, Charité – Universitätsmedizin Berlin, Charitéplatz 1, 10117 Berlin, Germany.
9. Present address: Clinic for Pediatrics and Adolescent Medicine, Universitätsklinikum Tübingen, Hoppe-Seyler-Str. 1, 72076 Tübingen, Germany.
10. German Cancer Consortium (DKTK), Berlin, Germany.
11. Charité – Universitätsmedizin Berlin, Charitéplatz 1, 10117 Berlin, Germany.

Author contributions:
C.E., A.K., and A.E.G. conceived and planned the experiments.
C.E. performed the experiments. F.M.M. performed bioluminescence recordings of three cell lines.
C.E., C.S., J.D., and S.D.L. analyzed the data.
A.M.F. and A.K. supported in the planning and implementation of the bioluminescence recordings.
C.E., C.S., A.M.F., J.D., S.D.L., J.S., T.S., U.K, H.H., A.K., and A.E.G. contributed to the interpretation of the results.
C.E. and A.E.G wrote the manuscript. All authors provided critical feedback and helped shape the research and manuscript.

Keywords: circadian clock, drug sensitivity, systems biology, chronopharmacology.

Abbreviations:
5-FU, 5-fluorouracil; AC, autocorrelation; ATCC, American type culture collection; BL1, basal-like1; BL2, basal-like2; BMAL1, brain and muscle arnt-like protein-1; CI, confidence interval; CLOCK, circadian locomotor output cycles protein kaput; CRY, cryptochrome; CWT, continuous wavelet transform; dKO, double-knockout; DMEM, Dulbecco's Modified Eagle Medium; DMSO, dimethyl sulfoxide; DWT, discrete wavelet transform; EGF, epidermal growth factor; GR, growth rate inhibition; LDA, linear discriminant analysis; Luc, luciferase; MES, mesenchymal; mTOR, mammalian target of rapamycin; NLS, nuclear localization sequence; PARP, poly ADP ribose polymerase; PCA, principal component analysis; PER2, period2; PI3K, phosphatidylinositol 3-kinase/alpha serine/ threonine-protein kinase; ROR, RAR-related orphan receptor; RPMI, Roswell Park Memorial Institute medium; SCN, suprachiasmatic nucleus; sKO, single-knockout; TNBC, triple-negative breast cancer; ToD, time-of-day; TPM, transcripts per million.

Abstract

The circadian clock, a fundamental biological regulator, governs essential cellular processes in health and disease. Circadian-based therapeutic strategies are increasingly gaining recognition as promising avenues. Aligning drug administration with the circadian rhythm can enhance treatment efficacy and minimize side effects. Yet, uncovering the optimal treatment timings remains challenging, limiting their widespread adoption. In this work, we introduce a novel high-throughput approach integrating live-imaging and data analysis techniques to deep-phenotype cell models, evaluating their circadian rhythms, growth, and drug responses. We devised a streamlined process for profiling drug sensitivities across different times of the day, identifying optimal treatment windows and responsive cell types and drug combinations. Finally, we implement multiple computational tools to uncover cellular and genetic factors shaping time-of-day drug sensitivity. Our versatile approach is adaptable to various biological models, facilitating its broad application and relevance. Ultimately, this research leverages circadian rhythms to optimize drug treatments, promising improved outcomes and transformative treatment strategies.

Introduction

The circadian clock is a central regulator of multiple physiological and behavioral processes found in cyanobacteria, plants, fungi, and animals. In mammals, the hierarchical organization of the circadian system ensures coordinated biological rhythms from the level of the individual cell to whole organism level[1]. Primate and mouse studies showed that up to 40% of all protein-coding genes are rhythmically expressed in a tissue-specific manner[2,3]. These clock-controlled genes regulate key biological processes such as metabolism[4,5], cell proliferation[6], immune response[7], DNA repair and apoptosis[8].

Disruption of the circadian system is classified as a carcinogen and is associated with multiple cancer subtypes[9-12]. Cancer hallmarks such as sustained proliferation and metastasis[13,14] have been linked to the circadian clock[15,16] and patients with mutations in circadian clock genes exhibit lower survival rates[17-19]. Beyond its role in cancer development, the circadian clock directly interacts with therapeutic targets that affect drug responses[19-21]. Consistent with these observations, recent works have shown that administration of chemotherapeutic agents aligned with the circadian rhythm change their degree of efficacy throughout the day[20,22-24]. Despite the broad recognition of the benefits of circadian-based drug treatments[22,25,26], an efficient strategy to identify optimal treatment times remains elusive, creating a bottleneck in the implementation. In addition, the mechanisms shaping time-of-day (ToD) sensitivity profiles remain widely unknown.

Here, we introduce a novel method for the thorough characterization of time-of-day responses in tumor and healthy tissue cell models (**Fig.1**). Using an array of experimental and data analysis methods we perform a deep-phenotyping of the critical cellular factors underlying time-of-day responses, i.e., the circadian clock strength, growth dynamics and drug response features. We then deploy a new high-throughput strategy to obtain ToD profiles in a panel of drugs and cell line models. Comparing tumor versus non-tumor ToD profiles provides candidate treatment timings to increase efficacy and reduce toxicity. We subsequently integrate our dataset with publicly available gene-expression databases to rigorously address three fundamental questions in the field of circadian pharmacology, known as chronopharmacology. These questions are: (A) what is the optimal time-of-day for drug treatment? (B) which cell subtype benefits the most from a circadian-aligned drug treatment? and (C) how do cellular features shape time-of-day profiles? Finally, we define a chronotherapeutic index, ranking

95 cellular models and drug agents that stands to gain the most benefit from circadian-based
96 treatments.
97

Results

Deep circadian phenotyping in cancer cell models

Despite the increasing recognition of the circadian clock in cancer, the extent to which different
cancer subtypes maintain circadian rhythmicity remains poorly understood. To address this,
we integrate recordings of circadian clock activity with comprehensive time-series analysis
techniques, as depicted in **Fig. 2a**. To robustly characterize the circadian clock network, we
monitored the positive and negative feedback arms of the molecular clock using a combination
of two circadian luciferase reporters for *Bmal1* and *Per2* (**Fig. 2b**). In **Fig. 2c** we show
representative raw luciferase signals from the breast cancer cell line MDAMB468, and the
corresponding detrended and normalized signals[27]. As expected for a robust functional clock
network, *Bmal1* and *Per2* signals show stable anti-phasic expression patterns throughout the
recording (**Fig. 2c** and **Supplementary Fig. 1a,b**).

Circadian clock strength varies in cancer and healthy tissue models

Following the acquisition of *Bmal1* and *Per2* expression dynamics, we conducted a
comprehensive circadian phenotyping of the signals. For this we implemented a strategy that
integrates three complementary time-series analysis techniques, i.e., autocorrelation (AC),
continuous wavelet transform (CWT), and multiresolution analysis (MRA). Using this
approach we screened a broad panel of cell models, including non-malignant breast epithelial
MCF10A, cancer cell models of various entities (breast cancer, neuroblastoma, and sarcoma)
as well as two knockout variants of osteosarcoma U-2 OS cells with a single deletion in the
circadian clock gene *Cry1* (U-2 OS sKO) or paired with a deletion in the *Cry2* locus[28] (U-2 OS
dKO). Calculating the autocorrelation function from each recording provides the strength and
period values, obtained from the second peak ordinate and abscissa (lag), respectively (**Fig.**
2d). This analysis showed a wide range of circadian strengths across models with the U-2 OS
wild-type ranking highest with a median autocorrelation value of 0.74, whereas U-2 OS sKO
and U-2 OS dKO variants ranked lowest with median correlation values of -0.04 ($p=3.5\times10^{-10}$)

and 0.09 ($p=1.3\times10^{-10}$), respectively (**Fig. 2e**). Heterogeneity between and within cancer entities was further observed for the oscillation period ranging from short periods of ~21 h in HCC1806 and U-2 OS sKO cells ($p=2.1\times10^{-3}$) to longer periods of ~26 h in MDAMB468 and MCF10A cells. U-2 OS dKO (34.8 h, $p=2.6\times10^{-4}$) and the neuroblastoma cell line SH-SY5Y (39.2 h) showed periods well above the circadian range and were excluded from this analysis (**Fig. 2f**). Consistent with these results, detrended signal traces of the different cell models and additional breast cancer cell lines indicate high variability in circadian clock signals across the models tested (**Supplementary Fig. 1c**).

To capture the non-stationary aspects of circadian signals, we employed continuous wavelet transform, a prevalent technique for analyzing dynamic temporal signals[29]. **Fig. 2g** shows a CWT power spectrum heatmap of *Bmal1* signal from the MDAMB468 example. The heatmap displays signal components (0–40 h period) over the 120-h recording, color-coded by relative power (see **Methods**). The most important high-power component is marked by a persistent ridge for each time point. Strong signals exhibit continuous long ridges whereas weaker signals have short and discontinuous ridges. As a complementary measure of clock strength, we quantified the ridge length from all our recordings. Our analysis shows that U-2 OS cells have a well-maintained clock with a median ridge length of 4.1 days, closely followed by MCF7, MDAMB468 and MCF10A cells (3.4–3.8 days). U-2 OS sKO and dKO cells showed ridges shorter than two days (1.9 days [$p=6.7\times10^{-4}$] and 1.5 days [$p=5.5\times10^{-5}$], respectively) (**Fig. 2h**).

For a more comprehensive understanding of the signal components, we next implemented multiresolution analysis. MRA involves decomposing the detrended signal into four distinct component bands, namely the noise (1–4 h), ultradian (4–16 h), circadian (16–32 h) and infradian (32–48 h) components (see **Methods**). In **Fig. 2i** we present an example of MDAMB468 *Bmal1*-Luc where 75.9% of the signal is in the circadian range, 4.7% in the noise and the remaining 19.4% in the ultradian or infradian range. To obtain an analogous signal-to-noise measure of all recorded cell models, we plotted the circadian component ("circadianicity") versus the noise component and observed a broad range of ratios (**Fig. 2j**). Consistent with our previous analysis, U-2 OS cells show strong circadian signal with the lowest noise levels (<1%) and the highest proportion of circadian components (~93%) for both reporters. Knockout of *Cry1* or *Cry1/Cry2* reduced the circadian component to 59% ($p=1\times10^{-6}$) and 40% ($p=5.6\times10^{-7}$), respectively, while increasing the noise component by 5.8-

fold in the single knockout and by 209-fold in the double knockout. This signal-to-noise analysis map indicates that signals from the *Per2*-reporter were slightly more circadian and less noisy than those from the *Bmal1*-reporter (**Fig. 2j** and **Supplementary Fig. 1d**). Finally, to obtain a global strength metric, we normalized each circadian parameter to its respective maximum value across all tested cell line models and computed the mean, facilitating a gradual ranking of cell models from low to high circadian strength (**Fig. 2k**, see **Methods**).

In summary, we identified heterogeneous circadian clock phenotypes in the different models tested, suggesting a strong clock in U-2 OS, MCF7, MDAMB468 and MCF10A cells and an impaired but present clock in GIMEN and HCC1806 cells. These results challenge the common expectation that most cancer cells have a weak clock and underscore the significance of defining gradual metrics of circadian strength in a model-specific manner.

Growth and drug response dynamics in cancer cell models

Together with circadian potency, cell growth dynamics and how cells respond to drug treatment in time is expected to influence responses along the day. Thus, we next evaluated growth characteristics and drug sensitivities across a spectrum of drug and cell line models. To showcase our approach's ability in detecting within-tissue differences, we examined nine cell lines of the triple-negative breast cancer (TNBC) subtype alongside the non-malignant MCF10A breast cell model. To capture the dynamics of growth and drug response, we implemented a live-cell imaging setup using two imaging channels, the brightfield for confluency and the fluorescence channel for nuclear detection (**Fig. 3a**). **Fig. 3b** shows representative snapshots of both imaging channels from MDAMB468 cells under unperturbed conditions. Frame-by-frame quantification provides growth trajectories for each cell model tested (**Supplementary Fig. 2a**), with MDAMB468 cells showing a 5.3-fold increase in growth and reaching approximately 100% confluency after 4 days (**Fig. 3c**). Quantification of doubling times (DTs) calculated from confluency and cell numbers correlated well across the 10 cell line models tested (R^2=0.69) (**Supplementary Fig. 2b**). Upon examining the normalized cell number trajectories, we observed substantial variability in growth across the cell line models, ranging from approximately 2- to 16-fold over the 4-day recordings (**Fig. 3d**). To capture growth signatures from the entire recording, we fitted an exponential function to each time series and obtained the growth rate k for each trajectory (**Fig. 3e**). As expected, cell number

growth rates and doubling times exhibit a strong anticorrelation (R^2=0.83) (**Fig. 3f** and

Supplementary Fig. 2c).

Next, we studied the response of the cancer cell models upon treatment with a panel of seven

drugs targeting a broad range of mechanisms and pathways, i.e., the DNA synthesis inhibitors

5-FU and doxorubicin, mitosis-inhibiting alisertib and paclitaxel, the PI3K/AKT/mTOR

inhibitor torin2, as well as cisplatin and olaparib which target the DNA damage response

(DDR) pathway (**Fig. 3g**). We then tracked responses to a broad range of drug concentrations,

as exemplified in **Fig. 3h** by MDAMB468's olaparib treatment. Here, doses up to 1.5 µM

resulted in slightly slower growth relative to the control, while 6.3 µM of olaparib significantly

hindered growth, and ≥25 µM led to total inhibition and cell death. Fitting an exponential

function to growth curves yielded positive growth rate values for weak doses while higher

doses resulted in negative growth rates, indicating population decline. Using this approach, we

next quantified the response and stratified all tested cell lines and drugs.

Multi-parametric evaluation of drug sensitivity reveals heterogeneity between and within models

Accurate assessment of drug effects is essential for identifying exploitable weaknesses in

cancer treatments. To obtain robust drug sensitivity metrics, we employed the normalized

growth rate inhibition (GR) curve approach as described by Hafner $et\ al.$[30]. Here, GR values

are calculated as the ratio between growth rates under treated and untreated conditions and

normalized to a single cell division. Fitting a dose-response equation to GR values yields five

drug sensitivity parameters, namely, the concentrations at half-maximal effect (GEC_{50}) and at

which GR=0.5 (GR_{50}), the drug effect at infinite concentration (GR_{inf}), the Hill coefficient of

the sigmoidal fit (*Hill coeff.*), and the area over the curve (GR_{AOC}) (**Fig. 3i**). Interestingly, when

the same cell line was treated with different cancer drugs at their respective concentration

evoking a saturating GR_{inf} response distinct drug-dependent trajectories emerged (**Fig. 3j**). This

underscores the importance of a time-resolved approach for characterizing drug sensitivity

across different drugs and, accordingly, across distinct cell line models.

We next compared a subset of drug-sensitivity metrics and cell line models for the different

drugs tested (**Fig. 3k–m**). While most drugs were able to induce death in the subset of cell lines

to varying degrees, we identified three drugs that resulted in partial growth inhibition of

HCC1806, namely olaparib, 5-FU and alisertib (**Fig. 3k**). Hierarchical clustering of GR_{inf} values revealed a grouping of the two TNBC cell lines and while this clustering was maintained for the GEC_{50} values (**Fig. 3l**), the Hill coefficient resulted in a different clustering of the cell lines (**Fig. 3m**). To explore the potential relationships between various sensitivity metrics, we combined data from multiple drugs and cell models and computed cross-correlations, revealing significant associations among GEC_{50}, GR_{50}, and EC_{50} values ($p \leq 0.0001$) (**Fig. 3n**). Additionally, significant yet less pronounced correlations emerged between the Hill coefficient and different drug sensitivity parameters, whereas the GR_{inf} and GR_{AOC} values showed minimal correlation with other metrics. For complete GR curves and information on additionally tested cell lines and drugs, see **Supplementary Fig. 3** and **Supplementary Table 1**.

Variability throughout the day depends on the drug and tissue model

The presence of a robust circadian clock combined with drugs that efficiently affect growth in a tumor model are promising prerequisites for identifying drug variability throughout the day. However, these prerequisites do not provide a priori insights into the specific time-of-day response profile. To systematically screen ToD drug sensitivities, we developed a new experimental strategy designed to significantly reduce the investigator's workload and number of consecutive drug perturbations, thereby increasing throughputness, reproducibility and accuracy (**Fig. 4a**). In this new approach, cell populations are seeded 24 h prior to the start of continuous live-cell imaging followed by performing a 3-step circadian clock resetting protocol in which separate cell populations receive a dexamethasone pulse at three different times (0, 4, 8 h). To test cells in later stages of the circadian cycle, drugs are administered at their estimated half effective dose 32 and 48 h after the first resetting step, creating a range of time differences between reset and treatment of 0, 4, 8, 16, 20 and 24 h in relative circadian time (**Fig. 4a**, *right panel*). The effects of the different treatment times on cell growth were monitored by live-cell imaging up to day 6, enabling the evaluation of drug responses for 4 days in both treatment groups, as shown for the alisertib-treated TNBC cell line HCC1937 in **Fig. 4b**. In typical ToD assays, cells are constantly proliferating during the 24 hours of ToD treatments, resulting in varying cellular densities at the time of the drug treatments. These varying cell densities have the potential to influence drug responses in vitro, possibly concealing or introducing bias when determining ToD-specific drug effects. To account for different cell densities at the time of

treatment, drug responses are determined as the ratio of the number of cells at the time of each

treatment to the number of cells 96 h after each treatment, keeping the time-window from

treatment to evaluation identical across conditions (**Fig. 4c**, *left panel*). To highlight relative

response differences within a day, results are presented relative to the circadian time of 0 h.

Values >1 indicate increased resistance, while values <1 indicate higher sensitivity compared

to treatment at time 0 h (**Fig. 4c**, *right panel*). We quantify the maximum variability in relative

responses as the ToD Maximum Range (*ToD*$_{MR}$).

Our next objective was to explore the variability of ToD profiles of different drugs within a

subset of TNBC cells. To accomplish this, we screened ten cell line models shown in **Fig. 3**

treated with eight different drugs, generating approximately 80 ToD profiles (**Fig. 4d,e** and

Supplementary Fig. 4a,b). Comparing examples from individual cell lines, namely HCC1937

and MCF10A, revealed that HCC1937 exhibited relatively conserved ToD profiles, whereas

the non-malignant cell line showed greater ToD variability for the different drugs (**Fig. 4d**).

Furthermore, ToD sensitivity profiles varied significantly from drug to drug (**Fig. 4e** and

Supplementary Fig. 4b), suggesting that ToD profiles may be highly dependent on the cancer

cell model and drug mechanism of action.

To assess varying time-of-day sensitivities across all tested cell line models and drugs, we

calculated the corresponding *ToD*$_{MR}$ value for each drug-cell combination (**Fig. 4f**). Averaging

the *ToD*$_{MR}$ values per drug and cell line revealed a gradual ranking of ToD variability for each

drug and cell model tested (**Fig. 4g**). Within the drug panel, the highest and lowest ToD

variability was observed for cisplatin and alisertib, respectively, with a ~2-fold difference in

the average *ToD*$_{MR}$ value (**Fig. 4g,** *top panel*). Similarly, the ToD variability of the tested cell

lines varied by ~2-fold, with MCF10A showing the highest and HCC38 the lowest average

variability. Considering only cancer cell lines, SUM149PT ranked highest and showed a

similar degree of ToD sensitivity variability as the non-malignant cell line (**Fig. 4g**, *bottom*

panel).

Together, our findings reveal distinct ToD profiles across most tested drugs and models,

highlighting individualized ToD sensitivity within distinct drugs and cell models despite the

common TNBC categorization and shared drug-target pathways.

Determining treatment times for maximum drug effect

The critical factor in the design of future chronotherapeutic treatments may be not only be the sensitivity of cancer cells, but rather the differential time-of-day sensitivity between cancer and non-malignant tissues[26]. To illustrate this, we compared the ToD response profiles of TNBC cancer cell models with the profile of MCF10A, as shown for HCC1937 and alisertib in **Fig. 4h**. In this case, the cancer and non-malignant tissue models show an antiphasic ToD response profile, with an almost inverted profile. By calculating the greatest response differences between the two cell models, we determined the treatment times of maximum and minimum benefit (**Fig. 4h–j**). Analyzing the entire panel of tested drugs per cell line model and plotting polar density histograms of the times of maximum and minimum benefit, we found that highest overall benefit is achieved at 10–12 h and 18–20 h in the treatment of HCC1937, while earlier treatment times of the day yield minimum benefit. In contrast, SUM149PT and MDAMB231 show a single prominent time window throughout the day that provides maximum treatment benefit (**Fig. 4i**). In addition to cell model-to-model variability in the benefit times, we explored the drug-to-drug variability by calculating the polar density histogram for the individual drugs tested across all cell line models (**Fig. 4j**). 5-FU showed a clear preference for administration between 8 and 10 h, while torin2 and paclitaxel showed more variability in maximum and minimum benefit times, indicating different response relationships to the non-malignant tissue model across the cancer cell lines tested.

Finally, we quantified the extent of maximum and minimum treatment benefit by calculating the corresponding fold changes in ToD responses between the cancer and non-malignant tissue model (**Fig. 4k**). By averaging these fold changes across cell lines and drugs, we established a ranking referred to here as the "chronotherapeutic index". While the average ToD variability highlights the benefits of a single model (**Fig. 4g**), the chronotherapeutic index reveals distinctions between cancer and non-malignant models, thereby it is offering valuable insights into the potential advantages of adopting a chronotherapeutic-based schedule. It's worth noting that the rankings based solely on average ToD variability within a single model or drug didn't align completely with the ranking of the chronotherapeutic index. This emphasizes the importance of evaluating effects in relation to healthy tissues when considering chronotherapy applications (**Fig. 4g,k**).

To sum up, our findings emphasize distinct chronotherapeutic response dynamics in vitro between cancer and healthy tissue models, underscoring two key objectives of chronotherapy:

311 pinpointing optimal treatment times to maximize cancer toxicity while minimizing impacts on
312 healthy tissues.
313

The relationship between time-of-day profiles and clock, growth, and drug sensitivity metrics

The time-of-day sensitivity in cancer models results from complex interactions involving the
circadian clock, cancer cell growth, and drug responses. However, the specific mechanisms
governing these interactions remain largely unknown. To address this, we employ statistical
tools, including linear regression, dominance analysis, and determinant ranking, to uncover the
key parameters influencing a key aspect of ToD sensitivity curves, namely the maximum range
in responses (ToD_{MR}) (**Fig. 5a**). To explore how individual metrics relate to ToD_{MR} values, we
first performed pairwise linear regression analysis (**Fig. 5b–f**). We considered each circadian
channel individually or combined and found the strongest and most significant correlations for
clock metrics from the *Bmal1* channel (**Fig. 5d** and **Supplementary Fig. 5a**). ToD_{MR} values of
5-FU were significantly correlated with amplitudes ($r=0.8$, $R^2=0.6$) and the prominence of
circadian component (circadianicity) of the *Bmal1* signal ($r=0.7$, $R^2=0.43$), whereas other
circadian clock metrics were poorly associated (**Fig. 5b,d**). On the other hand, we observed no
significant and rather low associations for ToD_{MR} values and growth metrics as well as for
ToD_{MR} values and drug sensitivity metrics (**Fig. 5c,e,f**). To identify the best associated
individual metric and drug, we integrated all correlations and generated a ranking as shown in
Fig. 5g. For the different metrics, we observed the highest correlation for the amplitude
($r=0.43\pm0.02$, mean $\pm$ s.e.m.). Among the individual drugs, adavosertib ranked highest in
overall correlation ($r=0.39\pm0.03$, mean $\pm$ s.e.m.) between ToD_{MR} values and the different
metrics. Relative average associations between all metrics and ToD_{MR} values were
predominantly positive, with only five metrics demonstrating inverse relationships
(**Supplementary Fig. 5b**). Notably, only one drug, paclitaxel, displayed average negative
associations across all metrics.
While linear regression approaches are robust for examining individual metrics, they do not
provide relative information about which metrics have the most significant impact on ToD_{MR}
values. To address this, we used dominance analysis and systematically tested all possible
combinations of metrics. By measuring how much each metric improved the accuracy of the

model when added or removed, we identified the most important predictors of ToD_{MR} values to the different drugs. For alpelisib our analysis identified high predictive power for the amplitude of *Bmal1* signals, contributing 42% of the observed ToD sensitivity variability (**Fig. 5h**). In contrast, paclitaxel and adavosertib showed the most homogeneous distribution of individual contributions, with no metric appearing to have a dominant predictive power. Considering the shares of each determinant combined for all drugs led to a ranking of determinants as shown in **Fig. 5i**. Here, we observed the largest contributions for the *Bmal1* amplitude (mean = 21.9%) and smallest for the Hill coefficient of drug response curves (mean = 9.9%).

In summary, we were able to identify systematic dependencies of the maximal time-of-day drug sensitivities and different circadian clock, growth and drug sensitivity parameters. We have further shown that the predictive power of the different determinants varies substantially depending on the specific drug, highlighting the importance of considering multiple factors in understanding the time-of-day dependent variability of drug response in a cell model.

Differential impact of core clock genes in shaping time-of-day sensitivity of cancer models

Beyond a cancer model's circadian clock, growth, and drug features, the expression levels of core circadian clock genes are likely to contribute the time-of-day sensitivity profiles. Thus, we explored the connection between the expression patterns of 16 essential clock genes and the maximum range of ToD sensitivity profiles (ToD_{MR}). We defined core clock genes as those that directly control the transcriptional-translational feedback loops of the molecular clock network and whose dysregulation or mutations result in disrupted circadian rhythms[31,32]. The selected gene panel displayed distinct expression patterns across the TNBC cell lines tested (**Supplementary Fig. 6a**). To examine potential relationships between core clock genes and ToD_{MR} values, we used three different methods, namely linear correlation analysis, dimensionality reduction by linear discriminant analysis (LDA) and principal component analysis (PCA) (**Fig. 6a**).

Linear correlation analysis revealed significant correlations between the ToD_{MR} values from our panel of tested drugs and several of the core clock genes, suggesting potential links between the circadian clock machinery and drug sensitivity (**Fig. 6b**). Focusing on the mitosis inhibitor

paclitaxel, we identified particularly strong anticorrelations with the expression levels of *Per3*

(r=-0.88, R^2=0.78) and *Dbp* (r=-0.72, R^2=0.51) (**Fig. 6b,c**). Accumulating absolute

correlations of each drug for all circadian clock genes revealed highest overall correlation of

ToD$_{MR}$ values for cisplatin (r=0.40±0.21, mean ± s.d.), while *Per2* was most associated with

ToD$_{MR}$ values among the circadian clock gene panel (r=0.48±0.20, mean ± s.d.) (**Fig. 6d**).

To test if our gene set could broadly categorize the strength of time-of-day effects as indicated

by the *ToD*$_{MR}$ values, we next turned to LDA analysis in a drug-by-drug basis. To do so, we

first categorized cell models into two groups based on the median ToD$_{MR}$ value for each drug.

Fig. 6e provides an example using paclitaxel, effectively separating the data into groups with

high and low ToD$_{MR}$ values. Implementing LDA to examine the individual linear discriminant

components of each circadian clock gene revealed a clear ranking of genes in terms of their

importance in discriminating the cell lines into the two *ToD*$_{MR}$ groups. For paclitaxel, the core

clock genes *Cry2*, *Dbp*, and *Per3* were the most important contributors to the specific

discrimination, collectively accounting for approximately 46% of the discriminative

information (**Fig. 6e**, *right panel*). Importantly, the contribution of each gene in discriminating

between *ToD*$_{MR}$ responses varied for the different drugs, suggesting that each drug interacts

differently with the molecular components of the circadian clock (**Supplementary Fig. 6b–h**).

To explore the circadian clock's impact on ToD sensitivity across various drug treatments, we

assessed the overall contribution of each gene by cumulating their effects across all drugs.

Here, *Rorβ* exhibited the overall highest contribution, accounting for 10.5% ± 9.6% (mean ±

s.d.) of the discriminative information (**Fig. 6f**).

While LDA is a powerful method to determine the discriminatory contribution of the individual

core clock genes, PCA allows for identifying underlying patterns and relationships among the

different genes. For paclitaxel, PCA distributed the cell models well along the first two

principal components which combined explained 73.7% of the variability among the different

ToD sensitivity profiles (**Supplementary Fig. 6i**, *left panel*). Ranking of the PC loadings

showed that the degree of contribution of each circadian clock gene in shaping the PCA

outcome was highly individual. For paclitaxel, *Csnk1d* was the major contributor to the ToD

sensitivity variability within the first principal component, while along the second principal

component, *Clock* ranked as main contributor (**Supplementary Fig. 6i**, *right panels*).

Collectively, our findings suggest associations between the expression levels of core clock

genes and the observed ToD maximum ranges within the examined cell lines. These results

underscore potential connections between the molecular circadian clock and time-of-day drug responses.

Discussion

While the fundamental role of the circadian clock in influencing disease progression and treatment response is widely recognized[22,33,34], our understanding of the underlying biological mechanisms remains fragmented. Approaches that enable the precise study of drug-perturbations and treatment responses within controlled circadian-timed environments play a pivotal role in advancing our understanding of the effects regulated by the circadian clock. To date, there has been a notable absence of standardized methodologies for the comprehensive exploration of the chronotherapeutic potential of drugs and cancer models. In this work we present an integrative approach for the high-throughput quantification of time-of-day response profiles and the elucidation of circadian, growth and drug sensitivity factors that shape these profiles (**Fig. 1**). This innovative method holds significant promise for advancing biological discovery, offering a robust platform to explore the intricate interplay between circadian rhythms, cellular growth, and drug timing efficacy.

Despite the general expectation that the circadian clock is likely dysregulated in highly transformed cancers[19,33], our approach reveals robust rhythms in numerous cancer cell models, whose circadian properties were previously unknown (**Fig. 2**). Furthermore, our assessment of circadian rhythms in non-malignant epithelial cells, luminal breast cancer cells, and osteosarcoma cells aligns with prior research[28,35-37], underscoring the versatility of our approach in elucidating circadian clock properties across various cancer and tissue types.

Among these, we focused on triple-negative breast cancer, a highly aggressive and heterogeneous subtype, representing about 15% of diagnosed breast tumors. Current therapies are ineffective, with cytotoxic chemotherapy still being an integral part of treatment despite limited efficacy and severe side effects[38]. Our investigation into the growth dynamics and drug responses of multiple TNBC cell models has generated a diverse array of metrics. Notably, in agreement with recent studies[39], our results indicate that the sensitivity metrics exhibit milder correlations, challenging the conventional binary classification of sensitive versus resistant models and suggesting that drug sensitivity rankings are rather metric-specific and drug-dependent (**Fig. 3**).

By implementing our screening approach for the profiling of drug sensitivities across different times of the day, we generated approximately 80 distinct time-of-day profiles, unveiling a wide range of sensitivity variations across drugs and cell line models (**Fig. 4** and **Supplementary Fig. 4**). We then ranked models by their time-of-day sensitivity and observed up to 30% response differences throughout the day for the top scoring models. We further detected substantial variation in time-of-day sensitivity among the panel of drugs tested, revealing previously unknown time-of-day dependent drug effects. Consistent with previous findings, we show time-of-day dependent variability in responses to the DNA synthesis inhibitors 5-FU and doxorubicin as well as to the DNA intercalator cisplatin[22]. Furthermore, we have introduced a novel chronotherapeutic index, which gauges the relative advantages of adopting a circadian-based drug schedule. Various non-malignant tissue models may be employed for calculating this index, tailored to the particular disease model under investigation. For example, fibroblast models, whose strong circadian rhythms have been described in several works[40-42], are expected to show unique relationships with cancer models in terms of sensitivity patterns at different times of the day.

Delving into the intricate molecular mechanisms that potentially govern time-of-day sensitivity, our novel method has unraveled key relationships between circadian clock dynamics, growth patterns, and drug sensitivities, all of which significantly impact time-of-day drug responses. It is noteworthy that the specific contribution of these parameters varies depending on the particular drug under consideration, underscoring the multifactorial nature of time-of-day dependent variability in drug reactions (**Fig. 5**). Furthermore, our exploration has shed light on the involvement of core circadian clock genes in this context, hinting at potential connections between the molecular circadian clock network and ToD drug sensitivity (**Fig. 6**). While our findings are primarily based on in vitro models and are not directly applicable for treatment recommendations, our framework can be directly applied to model organisms of higher complexity, such as 3D patient-derived organoids and animal models where the circadian clock can be tracked. Furthermore, additional research is needed to discern the relative contributions of the cell cycle and circadian clock to the observed ToD sensitivity[40,43-45].

Altogether, we provide a comprehensive method to determine optimal drug treatment times of complex diseases and to identify the cell subtypes that will most benefit from a time-of-day based treatment. Leveraging innovative experimental and computational techniques our

468 approach further elucidates the specific biological features that shape time-of-day response
469 profiles, thereby advancing our understanding of the cellular mechanisms that drive time-of-
470 day sensitivity. As a result, our method provides tools to uncover the cellular mechanisms that
471 govern time-of-day sensitivity, paving the way for new biological discoveries.
472

Acknowledgements

We would like to thank Malthe Skytte Nordentoft Nielsen as well as the laboratories of Michela
Di Virgilio and Ingeborg Tinhofer-Keilholz for their valuable feedback to our project. We
would also like to thank Annika Winkler for her assistance with sample preparation. The results
are part of a project funded by the German Federal Ministry of Education and Research
(BMBF) through the e:Med Juniorverbund DeepLTNBC TP3-01ZX1917C. C.E. was partially
supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation)–
RTG2424/CompCancer – project number: 377984878 and is enrolled in the doctoral program
of the Berlin School of Integrative Oncology (BSIO). C.S. acknowledges support from the
DFG–SCHM 3362/4–1 project number: 511886499.

Materials and Methods

Experimental methods

Cell culture

HCC1143, HCC1806, HCC1937, HCC38, and MDAMB468 cells were obtained from the
American Type Culture Collection. CAL51, MDAMB231, MDAMB436 and SUM149PT cells
were kindly provided by the Sorger lab (Harvard Medical School, Ludwig Cancer Center,
Boston, USA). MCF10A and MCF7 cells were kindly gifted by the Brugge lab (Harvard
Medical School, Ludwig Cancer Center, Boston, USA). GIMEN and SH-SY5Y cells were
provided by the Schulte lab (Universitätsklinikum Tübingen, Clinic for Pediatrics and
Adolescent Medicine, Thübingen, Germany) and the U-2 OS reporter cell lines by the Kramer
lab (Charité, Institute for Medical Immunology, Berlin, Germany). MCF10A cells were
maintained according to the Brugge lab's media recipe based on DMEM/F12 media (Gibco)

supplemented with 5% horse serum (Gibco), 20 ng/ml EGF (Peprotech), 0.5 mg/ml
Hydrocortisone (Sigma), 100 ng/ml Cholera Toxin (Sigma), 10 µg/ml Insulin (Sigma) and 1%
penicillin-streptomycin (Pen-Strep, Gibco). All other cell lines were maintained in RPMI 1640
(Gibco) supplemented with 10% fetal bovine serum (FBS) (Gibco) and 1% Pen-Strep. For
bioluminescence recordings and long-term imaging, cells were cultured in FluoroBrite DMEM
medium (Gibco) supplemented with 10% FBS, 300 mg L-Glutamine (Gibco) and 1% Pen-
Strep. Cells were kept at 37 °C in a humidified 5% CO_2 environment and regularly tested for
mycoplasma.

Generation of reporter cell lines

To produce lentivirus carrying constitutive red-fluorescent nuclear reporters, HEK293T cells
at 80% confluence were transfected with a mix of 1.8 µg gag/pol packaging plasmid (Addgene
#14887), 0.7 µg pRev packaging plasmid (Addgene #12253), 0.3 µg VSV-G envelope plasmid
(Addgene #14888) and 3.2 µg of a plasmid with a EF1α-mKate2-NLS sequence. Lentivirus
expressing either *Bmal1*- or *Per2*-promoter driven luciferase reporter was produced by
transfecting HEK293T cells with 6 µg psPAX2 (Addgene #12260), 3.6 µg pMD2G (Addgene
#12259) and 8.4 µg lentiviral expression plasmid (pAB-mBmal1:Luc-Puro or plenti6-
mPer2:Luc-Blast, respectively). Transfections were conducted using Lipofectamine 3000
(Invitrogen) according to the manufacturer's instructions and media was replaced by RMPI
1640 medium supplemented with 10 mM HEPES before adding the transfection mixture.
Lentiviral supernatant was collected after 48 h and 72 h and passed through a 0.45 µm filter
(Millipore). For lentiviral transduction, receiver cells at 70% confluence were incubated with
a mix of 1 ml lentivirus-containing supernatant, 8 µg/ml protamine sulfate (Sigma) and 10 µM
HEPES (Gibco) for 6 h. Ensuing, cells were washed with PBS (Gibco) and cultured in their
regular culture medium for 2 days before starting antibiotic selection of transduced cells. To
select for transduced cells, cells were grown in medium containing 5 µg/ml blasticidin (Adooq)
or 2 µg/ml puromycin (Gibco) according to the resistance cassette on the lentiviral expression
plasmid until non-transduced control cells died. Details about the generation of the U-2 OS-
Cry1-sKO and U-2 OS-*Cry1*/*Cry2*-dKO cell lines can be found in the original publication by
Börding *et al.*[28].

Bioluminescence recordings

Cells expressing *Bmal1*- or *Per2*-promoter driven luciferase reporters were seeded in 35-mm dishes (Nunc) to reach approximate confluence on the following day. To account for potentially different phases of single-cell circadian clocks in cell populations, we collectively reset the circadian clocks by administering a standard dose of 1 µM dexamethasone[46,47] (Sigma, dissolved in EtOH). After 30 min incubation cells were washed once with PBS and imaging medium supplemented with 250 µM D-Luciferin (Abmole) was added. To avoid evaporation of the media throughout the bioluminescence recordings, the dishes were sealed with parafilm as described in Finger *et al.*[46]. Luminescence intensity was monitored at 10-min intervals for 5 days in an incubator-embedded luminometer (LumiCycle, Actimetrics).

Long-term live-cell imaging

All live-cell imaging experiments were performed with cells expressing the fluorescent mKate2-NLS nuclear reporter, seeded in 48-well plates (Falcon) at a density that saturates at unperturbed growth conditions towards the end of each experiment. Long-term live-cell imaging was conducted using an incubator-embedded Incucyte live-cell widefield microscope (Essen BioScience). Cells were imaged in the brightfield and for nuclei segmentation and cell counting in the red channel (excitation: 567–607 nm, emission: 622–704 nm). Images were analyzed by frame-by-frame nuclei counting with the in-built Incucyte software and results were further processed in MATLAB. For experiments that involved drug treatments, cells were seeded 1 day before starting the live recordings and images were taken using a 4x magnification lens in 2 fields-of-views per well every 1–2 h for a total duration of 4–6 days. Two independent plates per condition were assayed. For drug response experiments with cisplatin a single plate per condition was assayed, imaged in 9 fields-of-views using a 10x magnification lens. For experiments capturing unperturbed growth dynamics, 300–1000 cells, corresponding to ~10% confluency, were seeded 2 days before starting the live recordings. Using a 10x magnification lens, 9 images per well were acquired in 1–2 h intervals for a total duration of 4 days.

Dose-response curves

Drug stock solutions (100–10 mM) were prepared in DMSO and preserved at -20°C. Cisplatin (Sigma) stock solution of 3.33 mM was prepared in 0.9% NaCl and stored at room temperature.

For dose-response assays, a serial 5–6 point log4 dilution of each drug was freshly prepared in

its solvent prior to treatment. Following concentration ranges were tested: 100–0.1 μM for 5-

fluorouracil (5-FU, Sigma), alpelisib (Biozol) and olaparib (Adooq); 10–0.01 μM for torin2

(Sigma) and alisertib (Hölzel); 10–0.04 μM for adavosertib (Biocat); 1–0.004 μM for

doxorubicin (Hölzel); and 0.4–0.0004 μM for paclitaxel (Hölzel). Cisplatin was tested in a

serial 10-point log2 dilution with doses ranging from 70–0.14 μM. Compounds were added to

the cells one day after cell seeding in a drug-media mixture of 9% of total well-volume.

Solvent-only treated control cells were assayed along the treated conditions. Each tested

condition contained equal amounts of the solvent. Cell growth was monitored by live-cell

imaging for at least 4 days as described above.

Time-of-day treatments

Cells were seeded and allowed to attach overnight. The next day, live recordings commenced

as described earlier. We performed independent resetting steps every 4 h over an 8-h period,

creating distinct cell populations at 0, 4, or 8 h of circadian time. Subsequently, we

administered the same drug concentration, which corresponded to the estimated half-maximal

effective concentration, at either 32 or 48 h after the initial resetting step. This allowed us to

simultaneously test six different circadian stages (0, 4, 8, 16, 20, and 24 h). Drug concentrations

were determined separately for each cell line and drug combination. For cisplatin, the

treatments we performed slightly different, using 4 independent resetting steps every 3 hours

over a 9-h period followed by drug treatments 32 or 48 h post initial resetting. In all cases, cell

growth was continuously monitored through long-term live-cell imaging for a total of 6 days.

Computational methods

Time-series analysis of circadian signals

Detrending. Raw time-series data were detrended by applying a sinc filter with a 48-h cut-off

period using the open-source software package pyBOAT[27] (v0.9.1) within the Anaconda

Navigator (v1.10.0).

Amplitude envelope and normalization. Continuous amplitude envelope calculation was

obtained using continuous wavelet transform implemented in pyBOAT with a time window of

48 hours. The amplitude normalization was done by taking the inverse of the envelope of the

detrended signal as describe in Mönke G, *et al.* [27] .

Autocorrelation analysis. Periodicity data was assessed by calculating its autocorrelation and

abcissa at the second peak using the 'autocorr' and 'findpeaks' MATLAB functions[48] from the

detrended time-series.

Continuous wavelet transform. The main oscillatory component, known as the ridge

component, was obtained using a wavelet-based spectral analysis from amplitude-normalized

and detrended signals. For the ridge detection we used both an adaptable-threshold and a fixed-

threshold. The adaptable-threshold was based on each signal's half-maximal spectral power

and was implemented for the "ridge length" metric shown in Fig. 2. The instantaneous phase

difference between the *Bmal1* and *Per2* signals was calculated using the MATLAB 'atan2'

function (Supplementary Fig. 1a) and for the polar histogram representation, we deployed the

'polarhistogram' function from MATLAB (Supplementary Fig. 1b). For the analysis of

determinants for time-of-day sensitivity shown in Fig. 5, the metrics of the main oscillatory

component were derived using a fixed-ridge threshold of 40. This ridge threshold value was

chosen as it offered a well-balanced threshold suitable for comparing signals of varying

strengths.

Multiresolution analysis. Detrended time-series data were decomposed into a set of different

wavelet details D_j representing distinct disjoint frequency bands and a final smooth using a

discrete wavelet transform based multiresolution analysis[49]. The algorithm has been

implemented using the 'PyWavelets' python package[50], with a *db20* wavelet of the Daubechies

wavelet family as previously described in Myung, Schmal and Hong *et al.*[51]. Since each

wavelet detail D_j represents a period range between $2^j\Delta t$ and $2^{j+1}\Delta t$ (for j=1, 2, 3, …) we down-

sampled the time series from a $\Delta t=10$ min to a $\Delta t=30$ min sampling frequency to obtain a

circadian period band between 16–32 h for further analysis. Since the MRA decomposes the

variance of the detrended signal with respect to the different disjoint period bands it can be

used to determine the rhythmicity of the signal in the circadian period range[52].

Global circadian strength. To calculate the global circadian strength (*GCS*) of a cell line

model *i*, the autocorrelation peak value (peak), the wavelet-based continuous ridge length

(ridge) and the discrete circadianicity component (circadianicity) were normalized to the

respective maximum (max) value measured among all tested cell line models and averaged

according to the following equation:

$$614 \quad GCS_i = \text{mean}\left(\frac{\text{peak}_i}{\text{peak}_\text{max}}, \frac{\text{ridge}_i}{\text{ridge}_\text{max}}, \frac{\text{circadianicity}_i}{\text{circadianicity}_\text{max}}\right)$$

615 **Statistical analysis.** Linear regression model fitting of *Bmal1* and *Per2* circadianicity

616 components obtained by MRA was done with the 'fitlm' MATLAB function. Significant

617 differences in circadian parameters between knockout and wild-type U-2 OS cells were

618 calculated with a two-sample *t*-test using the 'ttest2' MATLAB function.

Multi-parametric analysis of growth dynamics

620 Growth data obtained from long-term live-cell imaging was smoothed using a robust local

621 regression approach of weighted linear least squares and a 2nd degree polynomial model

622 ('rloess' MATLAB function). Doubling times of smoothed cell numbers or confluency were

623 calculated according to the following equation:

$$624 \quad \text{Doubling time}(t) = t * \frac{\log(2)}{\log(y_0/y_t)}$$

625 where t refers to the time of assessment, in our case 96 h, and y_0 refers to the cell number at

626 timepoint 0. To calculate the exponential growth rate k per unit of time t, we normalized cell

627 numbers to the initial timepoint 0 (y_0) and fitted an exponential function to the growth curves:

$$628 \quad \text{Growth}(t) = y_0 * e^{(k*t)}$$

629 Exponential function fitting was done with the MATLAB 'fit' function[48]. Linear regression

630 model fitting to different combinations of growth parameters was done with the 'fitlm'

631 MATLAB function.

Estimation of drug sensitivity parameters

633 Drug response data obtained from long-term live-cell imaging was smoothed using a robust

634 local regression approach of weighted linear least squares and a 2nd degree polynomial model

635 ('rloess' MATLAB function). Following the method established by Hafner *et al.*[30], we

636 computed the growth rate inhibition (GR) at time t and for each dose c as follows:

$$637 \quad GR(c,t) = 2^{k(c,t)/k(0)} - 1$$

638   where $k(c, t)$ is the growth rate under drug treatment and $k(0)$ is the growth rate of untreated

639   cells. Drug response parameters were retrieved by fitting the dose-dependent GR values to a

640   sigmoid curve using the following equation:

641
$$GR(c) = GR_{\text{inf}} + \frac{1 - GR_{\text{inf}}}{1 + (c/GEC_{50})^{h_{\text{GR}}}}$$

642   where the fitted parameters are as described in Hafner *et al.*[30]. Standard EC_{50} values were

643   calculated by fitting final nucleus counts, normalized to the respective count of the control, to

644   the following sigmoidal function:

645
$$f(c) = E_{\text{min}} + \frac{1 - E_{\text{min}}}{1 + (c/EC_{50})^{h}}$$

where E_{min} corresponds to the minimum response, restricted to values between 0 and 1, and h

is the hill slope of the response curve, constrained to 0.5–10. Sigmoidal function fitting steps

were done with the MATLAB functions 'fit' (GR metrics) or 'lsqnonlin' (EC_{50} value)[48].

Hierarchical clustering analysis was implemented with the MATLAB 'clustergram' function

using an euclidian distance and average linkage method[48]. To account for the different tested

dose ranges across drugs for clustering, GEC_{50} values were normalized to the dose at GR_{inf}.

Pearson's linear correlation coefficients across drug sensitivity metrics were computed using

the MATLAB 'corr' function. For the comparison of cellular growth dynamics to drug doses

evoking the maximum tested effect, we chose growth curves corresponding to the doses closest

to the determined GR_{inf} values.

Time-of-day sensitivity evaluation

Drug response data of time-of-day treatment experiments was smoothed using a moving

average ('smoothdata' MATLAB function). Final nucleus counts were normalized to the

nucleus count at the respective time of treatment. Time-of-day response curves were generated

from the relative final nucleus counts of each treatment timepoint to the final nucleus count at

timepoint 0 and interpolated using the 'smoothingspline' function in MATLAB. The

smoothing parameter was set to 0.7. The maximum range across smoothed time-of-day

responses (ToD_{MR}) was calculated by subtraction of the minimum from the maximum relative

response. To assess whether the smoothing of response data introduces artifacts into ToD_{MR}

estimates, we performed linear regression analysis between ToD_{MR} values from raw and spline

smoothed data using the 'fitlm' MATLAB function and confirmed a high positive correlation

(**Supplementary Fig. 7**). ToD_{MR} values were clustered as described for the drug sensitivity

parameters, with missing data points substituted by values from the closest relevant column,

adhering to the 'nearest-neighbor' principle. Polar histograms of the treatment times with

maximum and minimum benefit were generated using the 'polarhistogram' MATLAB function

with 'probability' normalization to depict each bar's height as the fraction of observations

within its bin relative to the total observations.

Correlation and shapley value regression analysis

674 Linear regressions of cell-intrinsic features and drug-dependent ToD sensitivity have been

675 obtained using the 'stats' and 'optimize' modules of the 'SciPy' and the 'uncertainties' package

676 of the Python programming language. The relative importance of the different predictors in a

677 multiple linear regression model are obtained by Shapley value regression via the Python

678 'dominance-analysis' package.

Circadian clock gene expression analysis

Gene expression data of circadian clock genes were obtained from the Cancer Cell Line

Encyclopedia Dependency Map (CCLE DepMap, https://sites.broadinstitute.org/ccle/datasets,

Q4 of 2022)[53]. Circadian clock genes of the individual breast cancer cell lines were clustered

using the 'clustermap' plotting function of 'seaborn' with the euclidian distance metric and

complete linkage method. Spearman rank correlation coefficients of circadian clock genes and

drug-specific ToD sensitivity values were computed using the spearman rank correlation

algorithm of the 'SciPy stats' python language module and hierarchically clustered with the

correlation distance metric and single linkage method. Regression plots were obtained using

the 'jointplot' plotting function from 'seaborn', showing drug-specific correlation between the

ToD sensitivities and gene expression of each cell line, including their distribution, squared

spearman rank correlation coefficient, p-value, and a confidence interval of 95%.

Supervised and unsupervised dimensionality reduction

Supervised dimensionality reduction by linear discriminant analysis was performed with the 'LinearDiscriminantAnalysis' function of the 'sklearn discriminant_analysis' python language module. The default parameters were retained, using the exact full singular value decomposition solver and 1 component. Median-based binarization of the drug-dependent ToD sensitivity values were used as target and the resulting linear discriminant vector was plotted using random y-axis units to avoid overlapping of the data points. The contribution to the obtained discriminative information is shown in percentage for each of the circadian clock genes. Unsupervised dimensionality reduction by principal component analysis (PCA) was performed with the 'PCA' function of the 'sklearn decomposition' python language module using the exact full singular value decomposition solver and number of components equivalent to the sample size of the corresponding drug. Further analysis of the most informative PCA components was done by drawing the biplots of the first two principal components and showing the loadings of each circadian clock gene for both components.

Data availability

The experimental time series data and data tables for all results of this study are available upon request.

Code availability

Code used for the data analysis in this work is available upon request.

Competing interests

The authors declare no competing interests.

References

Chaix, A., Zarrinpar, A. & Panda, S. The circadian coordination of cell biology. *J Cell Biol* **215**, 15-25, doi:10.1083/jcb.201603076 (2016).

Zhang, R., Lahens, N. F., Ballance, H. I., Hughes, M. E. & Hogenesch, J. B. A circadian gene expression atlas in mammals: implications for biology and medicine. *Proc Natl Acad Sci U S A* **111**, 16219-16224, doi:10.1073/pnas.1408886111 (2014).

Mure, L. S. *et al.* Diurnal transcriptome atlas of a primate across major neural and peripheral tissues. *Science* **359**, doi:10.1126/science.aao0318 (2018).

Saini, C. *et al.* A functional circadian clock is required for proper insulin secretion by human pancreatic islet cells. *Diabetes, Obesity and Metabolism* **18**, 355-365, doi:https://doi.org/10.1111/dom.12616 (2016).

Neufeld-Cohen, A. *et al.* Circadian control of oscillations in mitochondrial rate-limiting enzymes and nutrient utilization by PERIOD proteins. *Proc Natl Acad Sci U S A* **113**, E1673-1682, doi:10.1073/pnas.1519650113 (2016).

Chakrabarti, S. & Michor, F. Circadian clock effects on cellular proliferation: Insights from theory and experiments. *Current Opinion in Cell Biology* **67**, 17-26, doi:https://doi.org/10.1016/j.ceb.2020.07.003 (2020).

Scheiermann, C., Kunisaki, Y. & Frenette, P. S. Circadian control of the immune system. *Nature Reviews Immunology* **13**, 190-198, doi:10.1038/nri3386 (2013).

Sancar, A. *et al.* Circadian clock control of the cellular response to DNA damage. *FEBS Lett* **584**, 2618-2625, doi:10.1016/j.febslet.2010.03.017 (2010).

Schernhammer, E. S. *et al.* Rotating night shifts and risk of breast cancer in women participating in the nurses' health study. *J Natl Cancer Inst* **93**, 1563-1568, doi:10.1093/jnci/93.20.1563 (2001).

Puram, Rishi V. *et al.* Core Circadian Clock Genes Regulate Leukemia Stem Cells in AML. *Cell* **165**, 303-316, doi:10.1016/j.cell.2016.03.015 (2016).

Papagiannakopoulos, T. *et al.* Circadian Rhythm Disruption Promotes Lung Tumorigenesis. *Cell Metab* **24**, 324-331, doi:10.1016/j.cmet.2016.07.001 (2016).

Chun, S. K. *et al.* Disruption of the circadian clock drives Apc loss of heterozygosity to accelerate colorectal cancer. *Science Advances* **8**, eabo2389, doi:doi:10.1126/sciadv.abo2389 (2022).

Diamantopoulou, Z. *et al.* The metastatic spread of breast cancer accelerates during sleep. *Nature* **607**, 156-162, doi:10.1038/s41586-022-04875-y (2022).

14   Wang, J. *et al.* Circadian protein BMAL1 promotes breast cancer cell invasion and metastasis
by up-regulating matrix metalloproteinase9 expression. *Cancer Cell International* **19**, 182,
doi:10.1186/s12935-019-0902-2 (2019).

15   Lee, Y. *et al.* G1/S cell cycle regulators mediate effects of circadian dysregulation on tumor
growth and provide targets for timed anticancer treatment. *PLOS Biology* **17**, e3000228,
doi:10.1371/journal.pbio.3000228 (2019).

16   Chen, W.-D. *et al.* The circadian rhythm controls telomeres and telomerase activity.
Biochemical and Biophysical Research Communications **451**, 408-414,
doi:https://doi.org/10.1016/j.bbrc.2014.07.138 (2014).

17   Rida, P., Syed, M. I. & Aneja, R. Time will tell: Circadian clock dysregulation in triple negative
breast cancer. *Front Biosci (Schol Ed)* **11**, 178-192 (2019).

18   Innominato, P. F. *et al.* Prediction of overall survival through circadian rest-activity monitoring
during chemotherapy for metastatic colorectal cancer. *International Journal of Cancer* **131**,
2684-2692, doi:https://doi.org/10.1002/ijc.27574 (2012).

19   Ye, Y. *et al.* The Genomic Landscape and Pharmacogenomic Interactions of Clock Genes in
Cancer Chronotherapy. *Cell Systems* **6**, 314-328.e312,
doi:https://doi.org/10.1016/j.cels.2018.01.013 (2018).

20   Lee, Y. *et al.* Time-of-day specificity of anticancer drugs may be mediated by circadian
regulation of the cell cycle. *Science Advances* **7**, eabd2645, doi:10.1126/sciadv.abd2645
(2021).

21   Dallmann, R., Brown, S. A. & Gachon, F. Chronopharmacology: New Insights and Therapeutic
Implications. *Annual Review of Pharmacology and Toxicology* **54**, 339-361,
doi:10.1146/annurev-pharmtox-011613-135923 (2014).

22   Lévi, F., Okyar, A., Dulong, S., Innominato, P. F. & Clairambault, J. Circadian Timing in
Cancer Treatments. *Annual Review of Pharmacology and Toxicology* **50**, 377-421,
doi:10.1146/annurev.pharmtox.48.113006.094626 (2010).

23   Wood, P. A., Du-Quiton, J., You, S. & Hrushesky, W. J. M. Circadian clock coordinates cancer
cell cycle progression, thymidylate synthase, and 5-fluorouracil therapeutic index. *Molecular
Cancer Therapeutics* **5**, 2023-2033, doi:10.1158/1535-7163.Mct-06-0177 (2006).

24   Damato, A. R. *et al.* A randomized feasibility study evaluating temozolomide circadian
medicine in patients with glioma. *Neuro-Oncology Practice* **9**, 193-200,
doi:10.1093/nop/npac003 (2022).

25   Ruben, M. D., Smith, D. F., FitzGerald, G. A. & Hogenesch, J. B. Dosing time matters. *Science*
365, 547-549, doi:doi:10.1126/science.aax7621 (2019).

26   Kramer, A. *et al.* Foundations of circadian medicine. *PLOS Biology* **20**, e3001567,
doi:10.1371/journal.pbio.3001567 (2022).

27   Mönke, G., Sorgenfrei, F. A., Schmal, C. & Granada, A. E. Optimal time frequency analysis
for biological data - pyBOAT. *bioRxiv*, 2020.2004.2029.067744,
doi:10.1101/2020.04.29.067744 (2020).

28   Börding, T., Abdo, A. N., Maier, B., Gabriel, C. & Kramer, A. Generation of Human CRY1
and CRY2 Knockout Cells Using Duplex CRISPR/Cas9 Technology. *Front Physiol* **10**, 577,
doi:10.3389/fphys.2019.00577 (2019).

29   Leise, T. L. Analysis of Nonstationary Time Series for Biological Rhythms Research. *J Biol*
Rhythms **32**, 187-194, doi:10.1177/0748730417709105 (2017).

30   Hafner, M., Niepel, M., Chung, M. & Sorger, P. K. Growth rate inhibition metrics correct for
confounders in measuring sensitivity to cancer drugs. *Nature Methods* **13**, 521-527,
doi:10.1038/nmeth.3853 (2016).

31   Ko, C. H. & Takahashi, J. S. Molecular components of the mammalian circadian clock. *Hum*
Mol Genet **15 Spec No 2**, R271-277, doi:10.1093/hmg/ddl207 (2006).

32   Takahashi, J. S. Transcriptional architecture of the mammalian circadian clock. *Nature Reviews*
Genetics **18**, 164-179, doi:10.1038/nrg.2016.150 (2017).

33   Mormont, M.-C. & Levi, F. Cancer chronotherapy: Principles, applications, and perspectives.
Cancer **97**, 155-169, doi:https://doi.org/10.1002/cncr.11040 (2003).

34   Sulli, G., Lam, M. T. Y. & Panda, S. Interplay between Circadian Clock and Cancer: New
Frontiers for Cancer Treatment. *Trends in Cancer* **5**, 475-494,
doi:https://doi.org/10.1016/j.trecan.2019.07.002 (2019).

35   Lin, H.-H., Qraitem, M., Lian, Y., Taylor, S. R. & Farkas, M. E. Analyses of BMAL1 and PER2
Oscillations in a Model of Breast Cancer Progression Reveal Changes With Malignancy. *Integr*
Cancer Ther **18**, 1534735419836494-1534735419836494, doi:10.1177/1534735419836494
(2019).

36   Lellupitiyage Don, S. S. *et al.* Circadian oscillations persist in low malignancy breast cancer
cells. *Cell Cycle* **18**, 2447-2453, doi:10.1080/15384101.2019.1648957 (2019).

37   Baggs, J. E. *et al.* Network Features of the Mammalian Circadian Clock. *PLOS Biology* **7**,
e1000052, doi:10.1371/journal.pbio.1000052 (2009).

38   Gupta, G. K. *et al.* Perspectives on Triple-Negative Breast Cancer: Current Treatment
Strategies, Unmet Needs, and Potential Targets for Future Therapies. *Cancers (Basel)* **12**, 2392,
doi:10.3390/cancers12092392 (2020).

39   Fallahi-Sichani, M., Honarnejad, S., Heiser, L. M., Gray, J. W. & Sorger, P. K. Metrics other
than potency reveal systematic variation in responses to cancer drugs. *Nat Chem Biol* **9**, 708-
714, doi:10.1038/nchembio.1337 (2013).

40   Nagoshi, E. *et al.* Circadian gene expression in individual fibroblasts: cell-autonomous and self-
sustained oscillators pass time to daughter cells. *Cell* **119**, 693-705,
doi:10.1016/j.cell.2004.11.015 (2004).

41   Izumo, M., Johnson, C. H. & Yamazaki, S. Circadian gene expression in mammalian fibroblasts
revealed by real-time luminescence reporting: Temperature compensation and damping.
Proceedings of the National Academy of Sciences **100**, 16089-16094,
doi:doi:10.1073/pnas.2536313100 (2003).

42   Balsalobre, A., Damiola, F. & Schibler, U. A Serum Shock Induces Circadian Gene Expression
in Mammalian Tissue Culture Cells. *Cell* **93**, 929-937, doi:https://doi.org/10.1016/S0092-
8674(00)81199-X (1998).

43   Bieler, J. *et al.* Robust synchronization of coupled circadian and cell cycle oscillators in single
mammalian cells. *Mol Syst Biol* **10**, 739, doi:10.15252/msb.20145218 (2014).

44   Gérard, C. & Goldbeter, A. Entrainment of the Mammalian Cell Cycle by the Circadian Clock:
Modeling Two Coupled Cellular Rhythms. *PLOS Computational Biology* **8**, e1002516,
doi:10.1371/journal.pcbi.1002516 (2012).

45   Bernard, S., Čajavec Bernard, B., Lévi, F. & Herzel, H. Tumor Growth Rate Determines the
Timing of Optimal Chronomodulated Treatment Schedules. *PLOS Computational Biology* **6**,
e1000712, doi:10.1371/journal.pcbi.1000712 (2010).

46   Finger, A.-M. *et al.* Intercellular coupling between peripheral circadian oscillators by TGF-
β signaling. *Science Advances* **7**, eabg5174, doi:doi:10.1126/sciadv.abg5174 (2021).

47   Balsalobre, A. *et al.* Resetting of Circadian Time in Peripheral Tissues by Glucocorticoid
Signaling. *Science* **289**, 2344-2347, doi:10.1126/science.289.5488.2344 (2000).

48   version 9.11.0.1769968 (R2021b) (The MathWorks Inc., 2021).

49   Leise, T. L. & Harrington, M. E. Wavelet-based time series analysis of circadian rhythms. *J*
Biol Rhythms **26**, 454-463 (2011).

50   Lee, G. R., Gommers, R., Waselewski, F., Wohlfahrt, K. & O'Leary, A. PyWavelets: A Python
package for wavelet analysis. *Journal of Open Source Software* **4**.

51   Myung, J. *et al.* The choroid plexus is an important circadian clock component. *Nature*
Communications **9**, 1062, doi:10.1038/s41467-018-03507-2 (2018).

52   Leise, T. L. Wavelet analysis of circadian and ultradian behavioral rhythms. *Journal of*
Circadian Rhythms, doi:10.1186/1740-3391-11-5 (2013).

53   Barretina, J. *et al.* The Cancer Cell Line Encyclopedia enables predictive modelling of
anticancer drug sensitivity. *Nature* **483**, 603-607, doi:10.1038/nature11003 (2012).

Fig. 1 | Framework for identifying optimal treatment times in cancer and healthy tissue models. Schematic overview of the experimental and computational framework to thoroughly characterize time-of-day drug responses in a variety of cell subtypes, such as cancer and non-malignant cell models. A combination of live recordings is implemented for the deep phenotyping of circadian strength, growth dynamics and drug responses that shape time-of-day profiles. Using a novel streamlined experimental approach, time-of-day sensitivity profiles are obtained in tumor and non-malignant cell models, providing best and worst timings for increased efficacy and reduced toxicity (*top panel*). A tandem computational pipeline integrates the deep phenotyping metrics as well as gene expression data of circadian clock genes to quantitatively address three fundamental questions in chronopharmacology (*bottom panel*). Combining multiple signatures, we define a chronotherapeutic index, ranking cellular models and drug agents by their size-effect gains from drug treatments aligned with the circadian clock (*bottom right panel*).

Fig. 2 | Determining circadian clock strength in cancer and healthy tissue cell models. a, Schematic of the experimental setup to detect circadian oscillations. **b,** Circadian feedback loops involving the core clock genes *Bmal1* and *Per2*, used here as circadian Luc-reporters. **c,** Raw (*left*) and processed (*right*) signals from MDAMB468-*Bmal1/Per2*-Luc cells. **d,** Autocorrelation (AC) of signals shown in **c**. Arrow indicates 2nd peak and corresponding abscissa (lag). Dashed lines=95.4% CI. **e,** Boxplot of AC values and **f,** corresponding lags at the 2nd peak of signals from various cell models. **g,** Detrended and amplitude-normalized signal (*top*) and corresponding wavelet power spectrum (*bottom*) obtained from continuous wavelet transform (CWT) showing time-resolved periods of signals from MDAMB468-*Bmal1*-Luc cells. Main oscillatory component (ridge) marked as red line. **h,** Boxplot of CWT ridge lengths of signals from the indicated cell models. Bottom and top edges of the boxes in **e**, **f**, and **h** indicate the 25th and 75th percentiles, respectively. Red horizontal lines denote the data's median and red crosses outliers. *n*=6–17 technical replicates, collected from two Luc-reporters. **i,** Multiresolution analysis (MRA) of detrended MDAMB468-*Bmal1*-Luc signals. %=fraction of respective component to total signal. **j,** Scatterplot of normalized noise versus circadianicity of signals from the indicated cell models, determined by MRA. Shaded area covers unattainable range. Data represents the mean ± s.d. *n*=3–9 technical replicates. **k,** Bar diagram ranking cell models by their global circadian strength integrating parameters obtained from AC (peak), CWT (ridge) and MRA (circadianicity). Data represents the mean ± s.d. For representation purposes only the positive s.d. is shown. *n*=3–6. Color coding of data shown in **b–i** and **k** according to the corresponding Luc-reporter for *Bmal1* (yellow) and *Per2* (blue). Color coding of different cell models in **e**, **f**, and **h–k** according to the tissue of origin. A two-sample *t*-test assuming equal variances was used to evaluate statistical significance, where **, *** and ****, indicate *p*-values<0.01, 0.001, and 0.0001, respectively.

Fig. 3 | Unravelling growth and drug response dynamics through long-term live-cell imaging. a, Schematic of the setup for capturing cell growth using fluorescent nuclear reporters. **b,** Bi-daily snapshots of MDAMB468 cell growth in the brightfield (*top*) and red fluorescent channel (*bottom*). Ruler = 400 μM. **c,** MDAMB468 confluency (*black line*) and cell numbers (*red line*) over time. **d,** Normalized growth curves of the indicated cell models. **e,** Exponential fit for MDAMB468 growth curves, yielding the growth rate (*k*), and fit accuracy (R^2). **f,** Bar diagrams ranking cell models by doubling times and growth rates. Data in **c–f** represents the mean ± s.e.m. from 9 snapshots. *k*-value errors reflect the 95% CI. For clarity in **f**, one-sided error bars are shown, with the arrow indicating the top range of the respective error bar. **g,** Schematic of the pathways targeted by the drugs used in this study. **h,** Cell numbers and corresponding *k*-values of MDAMB468 cells treated with varying olaparib doses (color-coded) or solvent (dashed line). Data represents the mean ± s.d. of two plates. **i,** Dose-response curve of *GR* values corresponding to **h**, highlighting various drug sensitivity metrics. Error bars = 95% CI. **j,** Normalized numbers of MDAMB468 cells treated with the GR_{inf} dose of the indicated drugs. Data represents the mean ± s.d. of two plates, or ± s.e.m of 9 snapshots (cisplatin). **k–m,** Hierarchical clustering of drug sensitivity parameters across cell-drug combinations. GEC_{50} values in **l** are shown relative to the GR_{inf} dose. **n,** Pearson's linear correlation coefficients of sensitivity parameters shown in **k–m** and additional cell-drug combinations (see Supplementary Table 1; *n*=49–50). A two-sample *t*-test assuming equal variances was used to evaluate statistical significance; *, **, *** and ****, indicate *p*-values<0.05, 0.01, 0.001, and 0.0001, respectively. Denoted are significant correlation coefficients. Data in **k–n** based on the mean of two plates or 9 snapshots (cisplatin).

Fig. 4 | Time-of-day drug sensitivity is drug and tissue model dependent. a, Schematic of the experimental setup to screen for time-of-day (ToD) responses. Three clock-resetting steps are performed in 4-h intervals. Drugs are administered 32 or 48 h post-initial reset, resulting in six circadian times (0, 4, 8, 16, 20 and 24 h). Growth is monitored by long-term live-cell imaging. Timelines on the right depict experimental (*top*) and relative circadian times (*bottom*), color-coded by each ToD tested. **b,** Cell counts of HCC1937 treated with alisertib at different times of the day. Normalization to respective time of treatment. **c,** Cell numbers 4 days post-treatment versus treatment times, corresponding to **b** (*left*). ToD response curve (ToD-RC) depicting relative responses to ToD 0 h. Blue arrows mark the maximum ToD response range (ToD_MR). **d,** ToD-RCs for HCC1937 or MCF10A cells treated with different drugs (color-coded). **e,** ToD-RCs for 10 cell models treated with paclitaxel or doxorubicin. Color coding of cell models according to tissue of origin. **f,** Hierarchical clustering of ToD_MR values across drug-cell combinations. Values above 0.2 are shown. Data clustering with the UPGMA method and euclidian distance. **g,** Bar diagrams ranking ToD_MR values per drug and cell model. **h,** ToD-RCs for alisertib-treated HCC1937 tumor and MCF10A non-tumor cells overlaid, yielding maximum and minimum benefit times. Data shown in **b–h** represents mean (± s.d.) of two plates. **i,** Polar histograms for benefit times across cell models (*n*=7–9 drugs) and j, drugs (*n*=9 cell models). **k,** Butterfly charts depicting fold changes relative to MCF10A at benefit times, averaged per cancer cell model (*left*) and drug (*right*). Color-coding of maximum and minimum benefit times shown in **h–k** in green and red, respectively. Data shown in **g** and **k** represents mean ± s.d. of tested cell models per drug and vice versa (*see* **f**). For clarity, one-sided error bars are shown.

Fig. 5 | Clock and drug sensitivity metrics shape ToD curves. a, Computational approach to investigate how circadian rhythms, cell growth dynamics, and drug sensitivity factors influence the time-of-day (ToD) drug efficacy. **b,** and **c,** Example of linear correlations between ToD_{MR} and circadian (**b**) or drug sensitivity parameters (**c**), for 5-FU (*n*=10 cell models) and cisplatin (*n*=5 cell models), respectively. Color coding of cell models according to tissue of origin. Grey-shaded area=95% CI. Model accuracy indicated by R^2-values. **d–f,** Hierarchical clustering of correlation coefficients (*r*) between ToD_{MR} values of different drugs (*rows*) and the respective metric (*columns*) for clock strength (**d**, *n*=10 cell models), growth (**e**, *n*=10 cell models) or drug sensitivity (**f**, *n*=5 cell models). *r*-values ≥ 0.5 and ≤ -0.5 are shown. Black boxes indicate examples shown in **b** and **c**. A two-sample *t*-test assuming equal variances was used to evaluate statistical significance; *, and **, indicate *p*-values <0.05, and 0.01, respectively. *Note*: for alpelisib in **d–f** sample size=(*n*-1); for cisplatin in **d** and **e**, *n*=9 and in **f**, *n*=9. **g,** Bar diagrams ranking the absolute correlation between ToD_{MR} values and each metric depicted in **d–f**, ranked by metric (*n*=8 drugs) or by drug (*n*=14 metrics). Data represents the mean ± s.e.m. For clarity, one-sided error bars are shown. **h,** Hierarchical clustering of the dominance analysis matrix showing the individual contribution of circadian clock, growth and drug sensitivity parameters (*columns*) in predicting drug dependent ToD_{MR} values (*rows*). Colors indicate the percentage contribution, as detailed in the color bar. See key for **d–f** for sample sizes (*n*). **i,** Boxplot of overall contribution of cellular metrics to predict ToD_{MR} values, corresponding to **h**. Black data points indicate the individual result from each drug (*n*=8 drugs). Red horizontal lines denote the median, white circles denote the mean. Bottom and top edges of the boxes indicate the 25th and 75th percentiles, respectively.

Fig. 6 | Gene expression analysis unveils diverse roles of circadian genes in determining time-of-day patterns. a, Approach to identify the role of core clock genes that determine the strength of time-of-day (ToD) drug sensitivity. **b,** Hierarchical clustering of spearman rank correlation coefficients between drug-dependent ToD_{MR} values (*rows*) and core clock genes (*columns*). Black boxes indicate examples shown in **c**. Statistical significance of the correlations were calculated, where *, and **, indicate *p*-values<0.05, and 0.01, respectively. Cisplatin and alpelisib: *n*=8 cell lines, else *n*=9 cell lines. **c,** Example of linear correlation analysis between paclitaxel ToD_{MR} values and *Per3* or *Dbp* expression levels measured as log2(TPM). Color-coded data points indicate different tumor cell models. Grey-shaded area=95% CI. Model accuracy indicated by R^2-values. Top and lateral histograms indicate the count of samples along the range of gene expression, and ToD_{MR} values, respectively. The grey continuous line indicates the normal distribution fit of the samples. *n*=9 cell lines. **d,** Boxplot showing the mean-based ranking of absolute spearman rank correlation values between circadian clock gene expression levels and ToD_{MR} values, categorized by drug (*n*=16 circadian clock genes) or by circadian clock gene (*n*=8 drugs). **e,** Linear discriminant analysis (LDA) on median-based binarized ToD_{MR} values for paclitaxel. Cell models with ToD_{MR} values below or above the median (M) are colored in blue and red, respectively (*middle and left panel*). The contribution to the obtained discriminative information is shown in percentage for each of the circadian clock genes (*right panel*). **f,** Boxplot showing the mean-based ranking of overall discriminative contributions of each circadian clock gene across all tested drugs. *n*=8 drugs. Bottom and top edges of the boxes shown in **d** and **f** indicate the 25th and 75th percentiles, respectively. Red horizontal lines denote the median, white circles denote the mean, and red crosses indicate outliers.

Supplementary Fig. 1. a, Example of absolute phase difference evolution analysis between *Bmal1*- and *Per2*-Luc signals here shown for single samples of MDAMB468 cells. **b,** Polar histogram depicting the phase difference between *Bmal1* and *Per2* signals in MDAMB468 cells over time. 2π denotes one full circadian cycle. Median and mean phase difference are indicated as pink circle and black star, respectively. **c,** Detrended *Bmal1*- (*yellow*) and *Per2*-Luc (*blue*) signals of all cell lines used in this study. **d,** Linear regression analysis of the MRA circadian component ('circadianicity') of *Bmal1*- versus *Per2*-Luc signals. Markers indicate different cell models. Data represents the mean ± s.d. (*n*=3–9 technical replicates). Grey-shaded area=95% CI. Model accuracy indicated by R^2-values. Color coding of cell models in **c** and **d** according to their tissue of origin. Data depicted in **c** and **d** represents the mean ± s.d. of 3–9 technical replicates.

Supplementary Fig. 2. a, Growth curves of the indicated cell lines monitored by live-cell imaging for 4 days quantified by cell numbers (*red line*) or confluency (*black line*). Data represents the mean ± s.e.m. of 9 images. **b,** Linear regression analysis of doubling times (DT's) calculated from confluency versus cell numbers. **c,** Linear regression analysis of the doubling time and growth rate (*k*) calculated from cell numbers of various cell lines.
Color coding of cell models in **b** and **c** according to their tissue of origin. Data shown in **b** and **c** represents the mean ± s.d. (DT's), or ± CI of the exponential fit (*k*-values). Grey-shaded area=95% CI. Model accuracy indicated by R^2-values.

Supplementary Fig. 3. Dose-response curves of *GR* values for the specified cell models and drugs. The data is based on the mean of two plates or 9 images (cisplatin), assayed by live-cell imaging for 4 days. Error bars indicate the 95% CI of the exponential fit.

Supplementary Fig. 4. a, Overlay of ToD response curves (ToD-RCs) for individual cell models treated with different drugs (color-coded). **b,** Overlay of ToD-RCs for multiple cell models, grouped by the administered drugs. Cell line models are color-coded. Data in **a** and **b** represents the mean for each drug-cell combination tested on two plates or 9 images (cisplatin).

Supplementary Fig. 5. a, Hierarchical clustering of correlation coefficients between ToD$_{MR}$ values for different drugs (*rows*) and clock strength metrics (*columns*) obtained from the *Per2*-reporter (*left*) or calculated as a mean from the *Bmal1*- and *Per2*-reporter (*right*). A two-sample *t*-test assuming equal variances was used to evaluate statistical significance; *, and **, indicate *p*-values <0.05, and 0.01, respectively. *n*=10 or 9 (cisplatin and alpelisib) cell models. **b,** Bar plots ranking relative overall correlation between ToD$_{MR}$ values and individual cellular metrics (depicted in **Fig. 5d–f**), averaged by metric (*left*, *n*=8 drugs) or by drug (*right*, *n*=14 metrics). Positive and negative correlation values are color-coded in light red and blue, respectively. Data represents the mean ± s.e.m.

Supplementary Fig. 6. a, Gene expression clustermap of the selected list of core clock genes (*rows*) for the different cell models (*columns*) used in this work. **b-h,** Linear discriminant analysis on median-based binarized ToD$_{MR}$ values for the indicated drugs. ToD$_{MR}$ values below or above the median (M) are color-coded as blue and red, respectively (*middle and left panel of each subfigure*). The contribution to the obtained discriminative information is shown in percentage for each of the circadian clock genes (*right panel of each subfigure*). **i,** Principal component analysis (PCA) biplot of the first two principal component scores on ToD sensitivity groups related to paclitaxel. The principal components for the circadian clock genes are super-imposed with the PCA biplot scores at the similar scale that reflects the contribution of the variables in determination of ToD sensitivity groups (color-coded, *left panel*). The variance (PC loadings) explained by each component is expressed in percentage (*middle and right panel*).

Supplementary Fig. 7. Linear regression analysis of the maximum range in ToD responses (ToD$_{MR}$) determined from discrete measured data points or from continuous spline smoothed curves. Drugs are color-coded. Dotted lines=95% CI. Model accuracy indicated by R^2-values. Data represents the mean ± s.d. of 4-day live-cell imaging recordings for each drug tested on two plates or 9 images (cisplatin). n=80 drug-cell combinations.