Time-of-day effects of cancer drugs revealed by high-throughput deep phenotyping

Running title: Time-of-day effects of drugs

Authors: Carolin Ector[1,2], Christoph Schmal[3], Jeff Didier[4], Sébastien De Landtsheer[4], Anna-Marie Finger[5,6], Francesca Müller-Marquardt[1,7], Johannes Schulte[8,9], Thomas Sauter[4], Ulrich Keilholz[1,10], Hanspeter Herzel[3,11], Achim Kramer[5], Adrián E. Granada*[1,10]

First author: Carolin Ector || **Corresponding author:** adrian.granada@charite.de

Affiliations:
1. Charité Comprehensive Cancer Center, Charité – Universitätsmedizin Berlin, Charitéplatz 1, 10117 Berlin, Germany.
2. Faculty of Life Sciences, Humboldt-Universität zu Berlin, Invalidenstraße 42, 10115 Berlin, Germany.
3. Institute for Theoretical Biology, Humboldt-Universität zu Berlin, Philippstraße 13, 10115 Berlin, Germany.
4. Department of Life Sciences and Medicine, University of Luxembourg, 2, Avenue de l'Université, L-4365 Esch-sur-Alzette, Luxembourg.
5. Institute for Medical Immunology, Charité – Universitätsmedizin Berlin, Charitéplatz 1, 10117 Berlin, Germany.
6. Present address: Department of Anatomy, University of California, San Francisco, 513 Parnassus Avenue HSW 1330, San Francisco, CA 94143-0452, USA.
7. Present address: Institute of Research for Development, University of Montpellier, 911 Av. Agropolis, 34394 Montpellier, France.
8. Department of Pediatric Oncology, Hematology and Stem Cell Transplantation, Charité – Universitätsmedizin Berlin, Charitéplatz 1, 10117 Berlin, Germany.
9. Present address: Clinic for Pediatrics and Adolescent Medicine, Universitätsklinikum Tübingen, Hoppe-Seyler-Str. 1, 72076 Tübingen, Germany.
10. German Cancer Consortium (DKTK), Berlin, Germany.
11. Charité – Universitätsmedizin Berlin, Charitéplatz 1, 10117 Berlin, Germany.

Author contributions:
C.E., A.K., and A.E.G. conceived and planned the experiments. C.E. performed the experiments. F.M.M. performed bioluminescence recordings of three cell lines. C.E., C.S., J.D., and S.D.L. analyzed the data. A.M.F. and A.K. supported in the planning and implementation of the bioluminescence recordings. C.E., C.S., A.M.F., J.D., S.D.L., J.S., T.S., U.K, H.H., A.K., and A.E.G. contributed to the interpretation of the results. C.E. and A.E.G wrote the manuscript. All authors provided critical feedback and helped shape the research and manuscript.

Keywords: circadian clock, drug sensitivity, systems biology, chronopharmacology.

Abbreviations:
5-FU, 5-fluorouracil; AC, autocorrelation; ATCC, American type culture collection; BL1, basal-like1; BL2, basal-like2; BMAL1, brain and muscle arnt-like protein-1; CI, confidence interval; CLOCK, circadian locomotor output cycles protein kaput; CRY, cryptochrome; CWT, continuous wavelet transform; dKO, double-knockout; DMEM, Dulbecco's Modified Eagle Medium; DMSO, dimethyl sulfoxide; DWT, discrete wavelet transform; EGF, epidermal growth factor; EP, epithelial; GR, growth rate inhibition; LDA, linear discriminant analysis; Luc, luciferase; MES, mesenchymal; mTOR, mammalian target of rapamycin; NLS, nuclear localization sequence; PARP, poly ADP ribose polymerase; PCA, principal component analysis; PER2, period2; PI3K, phosphatidylinositol 3-kinase/alpha serine/ threonine-protein kinase; ROR, RAR-related orphan receptor; RPMI, Roswell Park Memorial Institute medium; SCN, suprachiasmatic nucleus; sKO, single-knockout; TNBC, triple-negative breast cancer; ToD, time-of-day; TPM, transcripts per million.

Abstract

The circadian clock, a fundamental biological regulator, governs essential cellular processes in health and disease. Circadian-based therapeutic strategies are increasingly gaining recognition as promising avenues. Aligning drug administration with the circadian rhythm can enhance treatment efficacy and minimize side effects. Yet, uncovering the optimal treatment timings remains challenging, limiting their widespread adoption. In this work, we introduce a novel high-throughput approach integrating live-imaging and data analysis techniques to deep-phenotype cancer cell models, evaluating their circadian rhythms, growth, and drug responses. We devised a streamlined process for profiling drug sensitivities across different times of the day, identifying optimal treatment windows and responsive cell types and drug combinations. Finally, we implement multiple computational tools to uncover cellular and genetic factors shaping time-of-day drug sensitivity. Our versatile approach is adaptable to various biological models, facilitating its broad application and relevance. Ultimately, this research leverages circadian rhythms to optimize anti-cancer drug treatments, promising improved outcomes and transformative treatment strategies.

Introduction

The circadian clock is a central regulator of multiple physiological and behavioral processes found in cyanobacteria, plants, fungi, and animals. In mammals, the hierarchical organization of the circadian system ensures coordinated biological rhythms from the level of the individual cell to whole organism level[1]. Primate and mouse studies showed that protein-coding genes are rhythmically expressed in a tissue-specific by up to 80% and 40%, respectively[2,3]. These clock-controlled genes regulate key biological processes such as metabolism[4,5], cell proliferation[6], immune response[7], DNA repair and apoptosis[8].

Disruption of the circadian system is classified as a carcinogen and is associated with multiple cancer subtypes[9-12]. Cancer hallmarks such as sustained proliferation and metastasis[13,14] have been linked to the circadian clock[15,16] and patients with mutations in circadian clock genes exhibit lower survival rates[17-19]. Beyond its role in cancer development, the circadian clock directly interacts with therapeutic targets that affect drug responses[19-21]. Consistent with these observations, recent works have shown that administration of chemotherapeutic agents aligned with the circadian rhythm change their degree of efficacy throughout the day[20,22-24]. Despite the broad recognition of the benefits of circadian-based drug treatments[22,25,26], an efficient strategy to identify optimal treatment times remains elusive, creating a bottleneck in the implementation. In addition, the mechanisms shaping time-of-day (ToD) sensitivity profiles remain widely unknown.

Here, we introduce a novel method for the thorough characterization of time-of-day responses in tumor and healthy tissue cell models (**Fig.1**). Using an array of experimental and data analysis methods we perform a deep-phenotyping of the critical cellular factors underlying time-of-day responses, i.e., the circadian clock strength, growth dynamics and drug response features. We then deploy a new high-throughput strategy to obtain ToD profiles in a panel of drugs and cell line models. Comparing tumor versus non-tumor ToD profiles provides candidate treatment timings to increase efficacy and reduce toxicity. We subsequently integrate our dataset with publicly available gene-expression databases to rigorously address three fundamental questions in the field of circadian pharmacology, known as chronopharmacology. These questions are: (A) what is the optimal time-of-day for drug treatment? (B) which cell subtype benefits the most from a circadian-aligned drug treatment? and (C) how do cellular features shape time-of-day profiles? Finally, we define a chronotherapeutic index, ranking

93 cellular models and drug agents that stands to gain the most benefit from circadian-based
94 treatments.
95

Results

Deep circadian phenotyping in cancer cell models

Despite the increasing recognition of the role of the circadian clock in cancer progression and
treatment response, the extent to which different cancer subtypes maintain circadian
rhythmicity remains poorly understood. To address this and quantitatively characterize the
degree of rhythmicity of cancer cell models, we implement an approach that integrates
recordings of circadian clock activity with comprehensive time-series analysis techniques, as
depicted in **Fig. 2a**. To robustly characterize the circadian clock molecular network, we
monitored the positive and negative feedback arms of the molecular clock using a combination
of two circadian luciferase reporters for *Bmal1* and *Per2* (**Fig. 2b**). In **Fig. 2c** we show
representative raw luciferase signals from the breast cancer cell line MDAMB468, and the
corresponding detrended and normalized signals[27]. As expected for a robust functional clock
network, *Bmal1* and *Per2* signals show stable anti-phasic expression patterns throughout the
recording (**Fig. 2c** and **Supplementary Fig. 1a,b)**.

Circadian clock strength varies in cancer and healthy cell models

Following the acquisition of *Bmal1* and *Per2* expression dynamics, we conducted a
comprehensive circadian phenotyping of the signals. To accurately capture different aspects of
the circadian clock dynamics, we implemented a strategy that integrates three complementary
time-series analysis techniques, i.e., autocorrelation (AC), continuous wavelet transform
(CWT), and multiresolution analysis (MRA). The rationale behind using this complementary
analysis is to harness the strengths of each technique; AC for identifying stable temporal
features, CWT for revealing time-dependent amplitude and period changes, and MRA for
extracting multi-scale features, ensuring a comprehensive understanding of the signal
dynamics. Using this approach we screened a broad panel of cell models, including non-
malignant breast epithelial MCF10A, cancer cell models of various entities (breast cancer,
neuroblastoma, and sarcoma) as well as two knockout variants of osteosarcoma U-2 OS cells

with a single deletion in the circadian clock gene *Cry1* (U-2 OS sKO) or paired with a deletion

in the *Cry2* locus[28] (U-2 OS dKO).

Autocorrelation is a robust method for estimating the periodic quality of a signal, particularly

for time-series whose properties remain stable over time, known as stationary signals.

Calculating the autocorrelation function from each recording provides the strength and period

values, obtained from the second peak ordinate and abscissa (lag), respectively (**Fig. 2d**). This

analysis showed a wide range of circadian strengths across models with the U-2 OS wild-type

ranking highest with a median autocorrelation value of 0.74, whereas U-2 OS sKO and U-2 OS

dKO variants ranked lowest with median correlation values of -0.04 ($p= 7.7 \times 10^{-12}$) and 0.09

($p=1.4\times10^{-10}$), respectively (**Fig. 2e**). Heterogeneity between and within cancer types was

further observed for the oscillation period ranging from short periods of ~21 h in HCC1806

and U-2 OS sKO cells (p=n.s.) to longer periods of ~26 h in MDAMB468 and MCF10A cells.

U-2 OS dKO (34.8 h, $p=8.2\times10^{-5}$) and the neuroblastoma cell line SH-SY5Y (39.2 h) showed

periods well above the circadian range and were excluded from this analysis (**Fig. 2f**).

Consistent with these results, detrended signal traces of the different cell models and additional

breast cancer cell lines indicate high variability in circadian clock signals across the models

tested (**Supplementary Fig. 1c**).

To capture non-stationary features of the circadian signals, such as unstable periods and

fluctuating amplitudes, we implemented continuous wavelet transform, a prevalent technique

for analyzing dynamic temporal signals[29]. **Fig. 2g** shows a CWT power spectrum heatmap of

Bmal1 signal from the MDAMB468 example. The heatmap displays signal components (0–40

h period) over the 120-h recording, color-coded by relative power (see **Methods**). Time-

connected regions of high relative power mark the signal's main oscillatory component,

referred to as ridge. Strong signals exhibit continuous long ridges whereas weaker signals have

short and discontinuous ridges. As a complementary measure of clock strength, we quantified

the ridge length from all our recordings. Our analysis shows that U-2 OS cells have a well-

maintained clock with a median ridge length of 4.1 days, closely followed by MCF7,

MDAMB468 and MCF10A cells (3.4–3.8 days). U-2 OS sKO and dKO cells showed ridges

shorter than two days (1.9 days [$p=5.7\times10^{-3}$] and 1.5 days [$p=4.8\times10^{-4}$], respectively) (**Fig.

2h**).

While both, autocorrelation and continuous wavelet transform, provide insights into the most

significant signal component, they do not quantify how the signal is distributed among non-

circadian frequencies. To obtain more comprehensive signal information and an analogous signal-to-noise metric, we next implemented multiresolution analysis. MRA involves decomposing the detrended signal into four distinct component bands, namely the noise (1–4 h), ultradian (4–16 h), circadian (16–32 h) and infradian (32–48 h) components (see **Methods**). In **Fig. 2i** we present an example of MDAMB468 *Bmal1*-Luc where 75.9% of the signal is in the circadian range, 4.7% in the noise and the remaining 19.4% in the ultradian or infradian range. To obtain an analogous signal-to-noise measure of all recorded cell models, we plotted the circadian component ("circadianicity") versus the noise component and observed a broad range of ratios (**Fig. 2j**). Consistent with our previous analysis, U-2 OS cells show strong circadian signal with the lowest noise levels (<1%) and the highest proportion of circadian components (~93%) for both reporters. Knockout of *Cry1* or *Cry1/Cry2* reduced the circadian component to 59% ($p=5.8\times10^{-6}$) and 40% ($p=2.0\times10^{-8}$), respectively, while increasing the noise component by 5.8-fold in the single knockout and by 209-fold in the double knockout. This signal-to-noise analysis map indicates that signals from the *Per2*-reporter were slightly more circadian and less noisy than those from the *Bmal1*-reporter (**Fig. 2j** and **Supplementary Fig. 1d**). *Bmal1* and *Per2* exhibit unique oscillatory patterns, reflecting distinct biological pathways within the circadian clock system. Their individual behaviors might offer valuable insights into the functionality of the circadian rhythm. To streamline our analysis and facilitate comparison, we averaged the values of *Bmal1* and *Per2* in **Fig. 2b, e, h, and k**. However, for a more detailed examination, separate analyses for each can be found in **Supplementary Figure 1e and f**.

Finally, to obtain a global strength metric, we normalized each circadian parameter to its respective maximum value across all tested cell line models and computed the mean, facilitating a gradual ranking of cell models from low to high circadian strength (**Fig. 2k**, see **Methods**).

In summary, by using a multi-faceted approach to characterize circadian rhythms in the different models tested, we identified heterogeneous circadian clock phenotypes, suggesting a strong clock in U-2 OS, MCF7, MDAMB468 and MCF10A cells and an impaired but present clock in GIMEN and HCC1806 cells. These results challenge the common expectation that most cancer cells have a weak clock and underscore the significance of defining gradual metrics of circadian strength in a model-specific manner.

Growth and drug response dynamics in cancer cell models

Together with circadian potency, cell growth dynamics and how cells respond to drug treatment in time is expected to influence responses along the day. Thus, we next evaluated growth characteristics and drug sensitivities across a spectrum of drug and cell line models. To showcase our approach's ability in detecting within-subtype differences, we examined nine cell lines of the triple-negative breast cancer (TNBC) subtype alongside the non-malignant MCF10A breast cell model. The TNBC cell lines were analyzed across several molecular subtypes as classified by Lehmann *et al.*, specifically the basal-like 1 (BL1), basal-like 2 (BL2), and mesenchymal (MES) TNBC subtypes. Growth and drug sensitivity assays in cancer models often rely on single time point measurements (e.g., ATP-based viability and replating assays) that provide a time-averaged snapshot. However, cancer cells respond dynamically to drugs, with temporally evolving effects, so conclusions drawn from single time point data can potentially mask the true effects of drugs. To accurately capture the dynamics of growth and drug response, we implemented a time-resolved live-cell imaging setup, directly counting cell nuclei in a fluorescence channel, while simultaneously evaluating confluency in the complementary brightfield channel (**Fig. 3a**). **Fig. 3b** shows representative snapshots of both imaging channels from MDAMB468 cells under untreated conditions. Frame-by-frame quantification provides growth trajectories for each cell model tested (**Supplementary Fig. 2a)**, with MDAMB468 cells showing a 5.3-fold increase in growth and reaching approximately 100% confluency after 4 days (**Fig. 3c**). Quantification of doubling times (DTs) calculated from confluency and cell numbers correlated well across the 10 cell line models tested (R^2=0.69) (**Supplementary Fig. 2b**). Upon examining the normalized cell number trajectories, we observed substantial variability in growth across the cell line models, ranging from approximately 2- to 16-fold over the 4-day recordings (**Fig. 3d**). To capture growth signatures from the entire recording, we fitted an exponential function to each time series and obtained the growth rate k for each trajectory (**Fig. 3e**). As expected, cell number growth rates and doubling times exhibit a strong anticorrelation (R^2=0.83) (**Fig. 3f** and **Supplementary Fig. 2c**). Next, we studied the response of the cancer cell models upon treatment with a panel of seven drugs targeting a broad range of mechanisms and pathways, i.e., the DNA synthesis inhibitors 5-FU and doxorubicin, mitosis-inhibiting alisertib and paclitaxel, the PI3K/AKT/mTOR inhibitor torin2, as well as cisplatin and olaparib which target the DNA damage response (DDR) pathway (**Fig. 3g**). We then tracked responses to a broad range of drug concentrations,

as exemplified in **Fig. 3h** by MDAMB468's olaparib treatment. Here, doses up to 1.5 µM
resulted in slightly slower growth relative to the control, while 6.3 µM of olaparib significantly
hindered growth, and ≥25 µM led to total inhibition and cell death. Fitting an exponential
function to growth curves yielded positive growth rate values for weak doses while higher
doses resulted in negative growth rates, indicating population decline. Using this approach, we
next quantified the response and stratified all tested cell lines and drugs.

Multi-parametric evaluation of drug sensitivity reveals heterogeneity between and within models

Accurate assessment of drug effects is essential for identifying exploitable weaknesses in
cancer treatments. Traditional drug sensitivity metrics, like the IC_{50} (the drug concentration
that reduces cell counts by 50% relative to the control), can be greatly influenced by factors
such as the assay duration and the number of cell divisions between drug administration and
final evaluation of drug sensitivity. These factors can inadvertently introduce artefactual
correlations and lead to misinterpretations of drug sensitivity results. To obtain robust drug
sensitivity metrics, we employed the normalized growth rate inhibition (*GR*) approach as
described by Hafner *et al.*[30]. Here, growth rates under treated and untreated conditions are
compared and normalized to a single cell division. Fitting a dose-response equation to *GR*
values yields five drug sensitivity parameters, namely, the concentrations at half-maximal
effect (GEC_{50}) and at which $GR=0.5$ (GR_{50}), the drug effect at infinite concentration (GR_{inf}),
the steepness of the sigmoidal fit (*Hill coefficient*), and the area over the curve (GR_{AOC}) (**Fig.
3i**). Moreover, the *GR* value directly reflects cellular response phenotypes, indicating partial
growth inhibition for values between 0 and 1, complete cytostasis for a value of 0, and
signifying cell death in the range from 0 to -1. Interestingly, when the same cell line was treated
with different cancer drugs at their respective concentration evoking a saturating GR_{inf} response
distinct drug-dependent trajectories emerged (**Fig. 3j**). This underscores the importance of a
time-resolved approach for characterizing drug sensitivity across different drugs and,
accordingly, across distinct cell line models.
We next compared a subset of drug-sensitivity metrics and cell line models for the different
drugs tested (**Fig. 3k–m**). While most drugs were able to induce death in the subset of cell lines
to varying degrees, we identified three drugs that resulted in partial growth inhibition of

HCC1806, namely olaparib, 5-FU and alisertib (**Fig. 3k**). Hierarchical clustering of GR_{inf}

values revealed a grouping of the two TNBC cell lines and while this clustering was maintained

for the GEC_{50} values (**Fig. 3l**), the Hill coefficient resulted in a different clustering of the cell

lines (**Fig. 3m**). To explore the potential relationships between various sensitivity metrics, we

combined data from multiple drugs and cell models and computed cross-correlations, revealing

significant associations among GEC_{50}, GR_{50}, and EC_{50} values ($p{\leq}0.0001$) (**Fig. 3n**).

Additionally, significant yet less pronounced correlations emerged between the Hill coefficient

and different drug sensitivity parameters, whereas the GR_{inf} and GR_{AOC} values showed minimal

correlation with other metrics. For complete GR curves and information on additionally tested

cell lines and drugs, see **Supplementary Fig. 3** and **Supplementary Table 1**.

Variability throughout the day depends on the drug and cell model

The presence of a robust circadian clock combined with drugs that efficiently affect growth in

a tumor model are promising prerequisites for identifying drug variability throughout the day.

However, these prerequisites do not provide a priori insights into the specific time-of-day

response profile. To systematically screen ToD drug sensitivities, we developed a new

experimental strategy designed to significantly reduce the investigator's workload and number

of consecutive drug perturbations, thereby increasing throughputness, reproducibility and

accuracy (**Fig. 4a**). In this new approach, cell populations are seeded 24 h prior to the start of

continuous live-cell imaging followed by performing a 3-step circadian clock resetting protocol

in which separate cell populations receive a dexamethasone pulse at three different times (0, 4,

8 h). To test cells in later stages of the circadian cycle, drugs are administered at their estimated

half effective dose 32 and 48 h after the first resetting step, creating a range of time differences

between reset and treatment of 0, 4, 8, 16, 20 and 24 h in relative circadian time (**Fig. 4a**, *right*

panel). The effects of the different treatment times on cell growth were monitored by live-cell

imaging up to day 6, enabling the evaluation of drug responses for 4 days in both treatment

groups, as shown for the alisertib-treated TNBC cell line HCC1937 in **Fig. 4b**. In typical ToD

assays, cells are constantly proliferating during the 24 hours of ToD treatments, resulting in

varying cellular densities at the time of the drug treatments. These varying cell densities have

the potential to influence drug responses in vitro, possibly concealing or introducing bias when

determining ToD-specific drug effects. To account for different cell densities at the time of

treatment, drug responses are determined as the ratio of the number of cells at the time of each
treatment to the number of cells 96 h after each treatment, keeping the time-window from
treatment to evaluation identical across conditions (**Fig. 4c**, *left panel*). To highlight relative
response differences within a day, results are presented relative to the circadian time of 0 h.
Values >1 indicate increased resistance, while values <1 indicate higher sensitivity compared
to treatment at time 0 h (**Fig. 4c**, *right panel*). We quantify the maximum variability in relative
responses as the ToD Maximum Range (*ToD*$_{MR}$).

Our next objective was to explore the variability of ToD profiles of different drugs within a
subset of TNBC cells. To accomplish this, we screened ten cell line models shown in **Fig. 3**
treated with eight different drugs, generating approximately 80 ToD profiles (**Fig. 4d,e** and
Supplementary Fig. 4a,b). Comparing examples from individual cell lines, namely HCC1937
and MCF10A, revealed that HCC1937 exhibited relatively conserved ToD profiles, whereas
the non-malignant cell line showed greater ToD variability for the different drugs (**Fig. 4d**).
Furthermore, ToD sensitivity profiles varied significantly from drug to drug (**Fig. 4e** and
Supplementary Fig. 4b), suggesting that ToD profiles may be highly dependent on the cancer
cell model and drug mechanism of action.

To assess varying time-of-day sensitivities across all tested cell line models and drugs, we
calculated the corresponding *ToD*$_{MR}$ value for each drug-cell combination (**Fig. 4f**). Averaging
the *ToD*$_{MR}$ values per drug and cell line revealed a gradual ranking of ToD variability for each
drug and cell model tested (**Fig. 4g**). Within the drug panel, the highest and lowest ToD
variability was observed for cisplatin and alisertib, respectively, with a ~2-fold difference in
the average *ToD*$_{MR}$ value (**Fig. 4g,** *top panel*). Similarly, the ToD variability of the tested cell
lines varied by ~2-fold, with MCF10A showing the highest and HCC38 the lowest average
variability. Considering only cancer cell lines, SUM149PT ranked highest and showed a
similar degree of ToD sensitivity variability as the non-malignant cell line (**Fig. 4g**, *bottom*
panel).

To assess the impact of circadian clock disruptions on ToD-dependent drug sensitivity, we
evaluated circadian-perturbed U-2 OS *Cry1/2*-dKO cells, alongside wild-type cells that
demonstrated the strongest circadian rhythms in our assessments (see **Fig. 2**). Using our ToD
treatment approach (see **Fig. 4a**), we tested three drugs which elicited high ToD response
variations in our breast cancer panel (see **Fig. 4g**), and which target distinct molecular
pathways. Our results demonstrate substantial ToD-dependent drug sensitivity in WT cells and

markedly reduced responses in *Cry1/2*-dKO cells (**Supplementary Fig. 4c**). Specifically,
ToD$_{MR}$ values decreased by 62% for cisplatin, 58% for paclitaxel, and 40% for 5-FU in *Cry1/2*-
dKO compared to WT cells (**Supplementary Fig. 4d**), highlighting a critical role of the
circadian clock in influencing drug sensitivity throughout the day.
Together, our findings reveal distinct ToD profiles across most tested drugs and models,
highlighting individualized ToD sensitivity within distinct drugs and cell models despite the
common TNBC categorization and shared drug-target pathways.
319

Determining treatment times for maximum drug effect

The critical factor in the design of future chronotherapeutic treatments may be not only be the
sensitivity of cancer cells, but rather the differential time-of-day sensitivity between cancer and
non-malignant tissues[26]. To illustrate this, we compared the ToD response profiles of TNBC
cancer cell models with the profile of MCF10A, as shown for HCC1937 and alisertib in **Fig.
4h**. In this case, the cancer and non-malignant cell models show an antiphasic ToD response
profile, with an almost inverted profile. By calculating the greatest response differences
between the two cell models, we determined the treatment times of maximum and minimum
benefit (**Fig. 4h–j**). Analyzing the entire panel of tested drugs per cell line model and plotting
polar density histograms of the times of maximum and minimum benefit, we found that highest
overall benefit is achieved at 10–12 h and 18–20 h in the treatment of HCC1937, while earlier
treatment times of the day yield minimum benefit. In contrast, SUM149PT and MDAMB231
show a single prominent time window throughout the day that provides maximum treatment
benefit (**Fig. 4i**). In addition to cell model-to-model variability in the benefit times, we explored
the drug-to-drug variability by calculating the polar density histogram for the individual drugs
tested across all cell line models (**Fig. 4j**). 5-FU showed a clear preference for administration
between 8 and 10 h, while torin2 and paclitaxel showed more variability in maximum and
minimum benefit times, indicating different response relationships to the non-malignant cell
model across the cancer cell lines tested.
Finally, we quantified the extent of maximum and minimum treatment benefit by calculating
the corresponding fold changes in ToD responses between the cancer and non-malignant cell
model (**Fig. 4k**). By averaging these fold changes across cell lines and drugs, we established a
ranking referred to here as the "chronotherapeutic index". While the average ToD variability

highlights the benefits of a single model (**Fig. 4g**), the chronotherapeutic index reveals distinctions between the cancer models and the non-malignant MCF10A cell model, thereby it is offering valuable insights into the potential advantages of adopting a chronotherapeutic-based schedule. It's worth noting that the rankings based solely on average ToD variability within a single model or drug didn't align completely with the ranking of the chronotherapeutic index. This emphasizes the importance of evaluating effects in relation to healthy tissues when considering chronotherapy applications (**Fig. 4g,k**).

To sum up, our findings emphasize distinct chronotherapeutic response dynamics in vitro between cancer and healthy cell models, underscoring two key objectives of chronotherapy: pinpointing optimal treatment times to maximize cancer toxicity while minimizing impacts on healthy tissues.

The relationship between time-of-day profiles and clock, growth, and drug sensitivity metrics

The time-of-day sensitivity in cancer models results from complex interactions involving the circadian clock, cancer cell growth, and drug responses. However, the specific mechanisms governing these interactions remain largely unknown. To address this, we employ statistical tools, including linear regression, dominance analysis, and determinant ranking, to uncover the main parameters influencing a key aspect of ToD sensitivity curves, namely the maximum range in responses (ToD_{MR}) (**Fig. 5a**). To explore how individual metrics relate to ToD_{MR} values, we first performed pairwise linear regression analysis (**Fig. 5b–f**). We considered each circadian channel individually or combined and found the strongest and most significant correlations for clock metrics from the *Bmal1* channel (**Fig. 5d** and **Supplementary Fig. 5a,** see **Supplementary Table 2** for a complete set of circadian metrics across the cell models tested). ToD_{MR} values of 5-FU were significantly correlated with amplitudes ($r=0.8$, $R^2=0.6$) and the prominence of circadian component (circadianicity) of the *Bmal1* signal ($r=0.7$, $R^2=0.43$), whereas other circadian clock metrics were poorly associated (**Fig. 5b**,**d**). On the other hand, we observed no significant and rather low associations for ToD_{MR} values and growth metrics as well as for ToD_{MR} values and drug sensitivity metrics (**Fig. 5c,e,f**). To identify the best associated individual metric and drug, we integrated all correlations and generated a ranking as shown in **Fig. 5g**. For the different metrics, we observed the highest

correlation for the amplitude (r=0.43±0.02, mean±s.e.m.). Among the individual drugs, adavosertib ranked highest in overall correlation (r=0.39±0.03, mean ± s.e.m.) between ToD_{MR} values and the different metrics. Relative average associations between all metrics and ToD_{MR} values were predominantly positive, with only five metrics demonstrating inverse relationships (**Supplementary Fig. 5b**). Notably, only one drug, paclitaxel, displayed average negative associations across all metrics. To further test the associations between individual cellular parameters and ToD_{MR} values, we applied a linear regression model and compared calculated versus actual ToD_{MR} values of up to five new cell line models in Bland-Altman plots (**Supplementary Fig. 5c**, see **Supplementary Table 3** for the complete new dataset). This revealed mean biases of -0.04, -0.05, and 0.06 between calculated and actual ToD_{MR} values for the clock, growth, and drug sensitivity metrics, respectively. As also shown in **Supplementary Fig. 5c**, mean biases are near zero and most predicted data points lie within the limits of agreement, which indicates minimal overall bias and good agreement between the predicted and observed data points.

While linear regression approaches are robust for examining individual metrics, they do not provide relative information about which metrics have the most significant impact on ToD_{MR} values. To address this, we used dominance analysis and systematically tested all possible combinations of metrics. By measuring how much each metric improved the accuracy of the model when added or removed, we identified the highest contributing ones to the ToD_{MR} values of the different drugs. For alpelisib our analysis identified the amplitude of *Bmal1* signals as most important factor, accounting for 42% of the observed ToD sensitivity variability (**Fig. 5h**). In contrast, paclitaxel and adavosertib showed the most homogeneous distribution of individual contributions, with no single metric appearing especially important. Considering the shares of each determinant combined for all drugs led to a ranking of determinants as shown in **Fig. 5i**. Here, we observed the largest contributions for the *Bmal1* amplitude (mean = 21.9%) and smallest for the Hill coefficient of drug response curves (mean = 9.9%).

In summary, we were able to identify systematic dependencies of the maximal time-of-day drug sensitivities and different circadian clock, growth and drug sensitivity parameters. We have further shown that the relative importance of the different determinants varies substantially depending on the specific drug, highlighting the importance of considering multiple factors in understanding the time-of-day dependent variability of drug response in a cell model.

Differential impact of core clock genes in shaping time-of-day sensitivity of cancer models

Beyond a cancer model's circadian clock, growth, and drug features, the expression levels of core circadian clock genes are likely to contribute to the time-of-day sensitivity profiles. Thus, we explored the connection between the expression patterns of 16 essential clock genes and the maximum range in ToD sensitivity profiles (ToD_{MR}). We defined core clock genes as those that directly control the transcriptional-translational feedback loops of the molecular clock network and whose dysregulation or mutations result in disrupted circadian rhythms[31,32]. The selected gene panel displayed distinct expression patterns across the TNBC cell lines tested (**Supplementary Fig. 6a**). To examine potential relationships between core clock genes and ToD_{MR} values, we used three different methods, namely linear correlation analysis, dimensionality reduction by linear discriminant analysis (LDA) and principal component analysis (PCA) (**Fig. 6a**).

Linear correlation analysis revealed significant correlations between ToD_{MR} values of five tested compounds from our drug panel and selected core clock genes (**Fig. 6b**). Focusing on the mitosis inhibitor paclitaxel, we identified particularly strong anticorrelations with the expression levels of *Per3* ($r=-0.88$, $R^2=0.78$) and *Dbp* ($r=-0.72$, $R^2=0.51$) (**Fig. 6b,c**). Yet, overall correlation between circadian clock genes and ToD_{MR} values were relatively weak, which got also apparent when accumulating absolute correlations per drug or per metric (**Supplementary Fig. 6b**). Here, highest overall correlation of ToD_{MR} values were found for cisplatin ($r=0.40\pm0.21$, mean±s.d.), while *Per2* was most associated with ToD_{MR} values among the circadian clock gene panel ($r=0.48\pm0.20$, mean ± s.d.) (**Supplementary Fig. 6b**).

To assess the cumulative impact of circadian gene expression on the strength of time-of-day effects as indicated by the ToD_{MR} values, aiming to uncover more complex patterns that might not be explainable through linear correlations alone, we proceeded to apply LDA on a drug-by-drug basis. To do so, we first categorized cell models into two groups based on the median ToD_{MR} value for each drug. **Fig. 6d** provides an example using paclitaxel, effectively separating the data into groups of high and low ToD_{MR} values. Implementing LDA to examine the individual linear discriminant components of each circadian clock gene revealed a clear ranking of genes in terms of their importance in discriminating the cell lines into the two ToD_{MR}

groups. For paclitaxel, the core clock genes *Cry2*, *Dbp*, and *Per3* were the most important

contributors to the specific discrimination, collectively accounting for approximately 46% of

the discriminative information (**Fig. 6d**, *right panel*). Importantly, the contribution of each gene

in discriminating between ToD_{MR} response groups varied for the tested drugs, suggesting that

either the drug or the drug targets may interact differently with the molecular components of

the circadian clock (**Supplementary Fig. 6c–i**). To explore the circadian clock's impact on

ToD sensitivity across various drug treatments, we assessed the overall contribution of each

gene by cumulating their effects across all drugs. This revealed modest overall contributions,

with the highest ranking *Rorβ* accounting for 10.5%±9.6% (mean±s.d.) of the discriminative

information (**Fig. 6e**).

While LDA is a powerful method to determine the discriminatory contribution of the individual

core clock genes, PCA allows for identifying underlying patterns and relationships among the

different genes. For paclitaxel, PCA distributed the cell models well along the first two

principal components which combined explained 73.7% of the variability among the different

ToD sensitivity profiles (**Supplementary Fig. 6j**, *left panel*). Ranking of the PC loadings

showed that the degree of contribution of each circadian clock gene in shaping the PCA

outcome was highly individual. For paclitaxel, *Csnk1d* was the major contributor to the ToD

sensitivity variability within the first principal component, while along the second principal

component, *Clock* ranked as main contributor (**Supplementary Fig. 6j**, *right panels*).

Collectively, our findings suggest limited associations between individual expression levels of

core clock genes and the observed ToD maximum ranges within the examined cell lines.

However, we found interesting relationships when considering the circadian clock gene panel

as a collective, indicating potential connections between the molecular circadian clock and

drug-dependent time-of-day sensitivity.

Discussion

While the fundamental role of the circadian clock in influencing disease progression and

treatment response is widely recognized[22,33,34], our understanding of the underlying biological

mechanisms remains fragmented. Approaches that enable the precise study of drug-

perturbations and treatment responses within controlled circadian-timed environments play a

pivotal role in advancing our understanding of the effects regulated by the circadian clock. To

date, there has been a notable absence of standardized methodologies for the comprehensive

exploration of the chronotherapeutic potential of drugs and cancer models. In this work we

present an integrative approach for the high-throughput quantification of time-of-day response

profiles and the elucidation of circadian, growth and drug sensitivity factors that shape these

profiles (**Fig. 1**). This innovative method holds significant promise for advancing biological

discovery, offering a robust platform to explore the intricate interplay between circadian

rhythms, cellular growth, and drug timing efficacy.

Despite the general expectation that the circadian clock is likely dysregulated in highly

transformed cancers[19,33], our approach reveals robust rhythms in numerous cancer cell models,

whose circadian properties were previously unknown (**Fig. 2**). Furthermore, our assessment of

circadian rhythms in non-malignant epithelial cells, luminal breast cancer cells, and

osteosarcoma cells aligns with prior research[28,35-37], underscoring the versatility of our

approach in elucidating circadian clock properties across various cancer and tissue types.

Among these, we focused on triple-negative breast cancer, a highly aggressive and

heterogeneous subtype, representing about 15% of diagnosed breast tumors. Current therapies

are ineffective, with cytotoxic chemotherapy still being an integral part of treatment despite

limited efficacy and severe side effects[38]. Our investigation into the growth dynamics and drug

responses of multiple TNBC cell models has generated a diverse array of metrics. Notably, our

results indicate that our set of drug sensitivity metrics, namely the GR_{50}, GEC_{50}, GR_{AOC}, GR_{inf}

and the Hill coefficient, exhibit milder correlations to each other, which is in agreement with

recent studies[39]. This challenges the conventional binary classification of sensitive versus

resistant models and suggests that drug sensitivity rankings are rather drug-dependent and

sensitivity metric-specific (**Fig. 3**).

By implementing our screening approach for the profiling of drug sensitivities across different

times of the day, we generated approximately 80 distinct time-of-day sensitivity profiles,

unveiling a wide range of sensitivity variations across drugs and cell line models (**Fig. 4** and

Supplementary Fig. 4). We then ranked models by their time-of-day sensitivity and observed

up to 30% response differences throughout the day for the top scoring models. We further

detected substantial variation in time-of-day sensitivity among the panel of drugs tested,

revealing previously unknown time-of-day dependent drug effects. Consistent with previous

findings, we show time-of-day dependent variability in responses to the DNA synthesis

inhibitors 5-FU[22,23,40] and doxorubicin[22,41] as well as to the DNA intercalator cisplatin[22,41].

Furthermore, we have introduced a novel chronotherapeutic index, which gauges the relative

advantages of adopting a circadian-based drug schedule. Various non-malignant cell models

may be employed for calculating this index, tailored to the particular disease model under

investigation. For example, fibroblast models, whose strong circadian rhythms have been

described in several works[42-44], are expected to show unique relationships with cancer models

in terms of sensitivity patterns at different times of the day.

Delving into the molecular mechanisms that potentially govern time-of-day sensitivity, our

novel method has unraveled key relationships between circadian clock dynamics, growth

patterns, and drug sensitivities, all of which significantly impacting time-of-day drug

responses. It is noteworthy that the specific contribution of these parameters varies depending

on the drug under consideration, underscoring the multifactorial nature of time-of-day

dependent variability in drug reactions (**Fig. 5**). Aiming to simplify the circadian analysis and

to provide an overarching view on circadian clock dynamics, we initially combined *Bmal1-*

and *Per2*-signals in Fig. 2. However, separate evaluations of the circadian genes, as presented

in **Supplementary Fig. 1e and f**, proved crucial to elucidate the significant impact of

oscillatory *Bmal1* dynamics on ToD_{MR} values.

In an exploratory approach, we identified limited associations between expression levels of

core circadian clock genes and ToD sensitivities (**Fig. 6**). This may hint to a more complex role

of the molecular circadian clock network in this context, where distinct associations might be

masked by high redundancy and interconnection between core clock genes. Additionally, our

results could indicate a discrete role of the cell cycle in shaping our observed ToD sensitivity

profiles[45-47], which also invites for further research in that context.

Assessing the cumulative impact of circadian gene expression on ToD_{MR} values, we found

differential contributions of each circadian gene in discriminating between high and low ToD-

dependent sensitivity based on the drug tested. To investigate whether this observation

underlies variations in the expression levels of the drug targets, follow-up explorations

involving e.g. mRNA or protein quantification techniques performed at different times of the

day could be conducted in future studies focusing on the molecular mechanisms of ToD

sensitivity.

Given their recognized influence on pharmacodynamics[22,33,34], our study primarily focused on

circadian rhythms. Nevertheless, our multiresolution analysis of *Bmal1* and *Per2* oscillatory

signals uncovered distinct alternate rhythms of shorter and longer periods, as illustrated in **Fig.**

2i and **j**. The significant impact of independent 12-hour ultradian rhythms on stress responses has been previously documented for mammalian cells[48], and constitutes an interesting field for future research in the scope of our time-of-day drug sensitivity profiles.

Despite employing a robust method to introduce luciferase reporters and validating several circadian results, as well as using population-based recordings of non-clonal cell lines, our approach does not rule out the potential impact of random reporter insertions on circadian clock estimation. Moreover, while our findings are primarily based on in vitro models and are not directly applicable for treatment recommendations, our framework can be directly applied to model organisms of higher complexity, such as 3D patient-derived organoids and animal models where the circadian clock can be tracked.

Altogether, we provide a comprehensive method to determine optimal drug treatment times of complex diseases and to identify the cell subtypes that will most benefit from a time-of-day based treatment. Leveraging innovative experimental and computational techniques our approach further elucidates the specific biological features that shape time-of-day response profiles, thereby advancing our understanding of the cellular mechanisms that drive time-of-day sensitivity. As a result, our method provides tools to uncover the cellular mechanisms that govern time-of-day sensitivity, paving the way for new biological discoveries.

Acknowledgements

We thank Malthe Skytte Nordentoft Nielsen as well as the laboratories of Michela Di Virgilio and Ingeborg Tinhofer-Keilholz for their valuable feedback to our project. We would like to express our gratitude to Annika Winkler and Marie Möser for their assistance with sample preparation, as well as to Sofía Peso-García and Franziska Reiher for their experimental support. We further thank Christian Gabriel and Valentina Alejandra Balde Araya for valuable feedback during the review of this manuscript. Lastly, we sincerely thank the editor and anonymous reviewers for their comprehensive review of our manuscript and for providing constructive feedback which allowed us to significantly improve our manuscript. The results are part of a project funded by the German Federal Ministry of Education and Research (BMBF) through the e:Med Juniorverbund DeepLTNBC TP3-01ZX1917C. C.E. was partially supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation)– RTG2424/CompCancer – project number: 377984878 and is enrolled in the doctoral program

563 of the Berlin School of Integrative Oncology (BSIO). C.S. acknowledges support from the
564 DFG–SCHM 3362/4–1 project number: 511886499.

Materials and Methods

Experimental methods

Cell culture

569 HCC1143, HCC1806, HCC1937, HCC38, and MDAMB468 cells were obtained from the
570 American Type Culture Collection. BT549, CAL51, MDAMB231, MDAMB436 and
571 SUM149PT cells were kindly provided by the Sorger lab (Harvard Medical School, Ludwig
572 Cancer Center, Boston, USA). MCF10A and MCF7 cells were kindly gifted by the Brugge lab
573 (Harvard Medical School, Ludwig Cancer Center, Boston, USA). GIMEN and SH-SY5Y cells
574 were provided by the Schulte lab (Universitätsklinikum Tübingen, Clinic for Pediatrics and
575 Adolescent Medicine, Thübingen, Germany) and the U-2 OS reporter cell lines by the Kramer
576 lab (Charité, Institute for Medical Immunology, Berlin, Germany). MCF10A cells were
577 maintained according to the Brugge lab's media recipe based on DMEM/F12 media (Gibco)
578 supplemented with 5% horse serum (Gibco), 20 ng/ml EGF (Peprotech), 0.5 mg/ml
579 Hydrocortisone (Sigma), 100 ng/ml Cholera Toxin (Sigma), 10 µg/ml Insulin (Sigma) and 1%
580 penicillin-streptomycin (Pen-Strep, Gibco). All other cell lines were maintained in RPMI 1640
581 (Gibco) supplemented with 10% fetal bovine serum (FBS) (Gibco) and 1% Pen-Strep. For
582 bioluminescence recordings and long-term imaging, cells were cultured in FluoroBrite DMEM
583 medium (Gibco) supplemented with 10% FBS, 300 mg L-Glutamine (Gibco) and 1% Pen-
584 Strep. Cells were kept at 37 °C in a humidified 5% CO_2 environment and regularly tested for
585 mycoplasma.

Generation of reporter cell lines

587 To produce lentivirus carrying constitutive red-fluorescent nuclear reporters, HEK293T cells
588 at 80% confluence were transfected with a mix of 1.8 µg gag/pol packaging plasmid (Addgene
589 #14887), 0.7 µg pRev packaging plasmid (Addgene #12253), 0.3 µg VSV-G envelope plasmid
590 (Addgene #14888) and 3.2 µg of a plasmid with a EF1α-mKate2-NLS sequence. Lentivirus

expressing either *Bmal1*- or *Per2*-promoter driven luciferase reporter was produced by

transfecting HEK293T cells with 6 μg psPAX2 (Addgene #12260), 3.6 μg pMD2G (Addgene

#12259) and 8.4 μg lentiviral expression plasmid (pAB-mBmal1:Luc-Puro or plenti6-

mPer2:Luc-Blast, respectively). Transfections were conducted using Lipofectamine 3000

(Invitrogen) according to the manufacturer's instructions and media was replaced by RMPI

1640 medium supplemented with 10 mM HEPES before adding the transfection mixture.

Lentiviral supernatant was collected after 48 h and 72 h and passed through a 0.45 μm filter

(Millipore). For lentiviral transduction, receiver cells at 70% confluence were incubated with

a mix of 1 ml lentivirus-containing supernatant, 8 μg/ml protamine sulfate (Sigma) and 10 μM

HEPES (Gibco) for 6 h. Ensuing, cells were washed with PBS (Gibco) and cultured in their

regular culture medium for 2 days before starting antibiotic selection of transduced cells. To

select for transduced cells, cells were grown in medium containing 5 μg/ml blasticidin (Adooq)

or 2 μg/ml puromycin (Gibco) according to the resistance cassette on the lentiviral expression

plasmid until non-transduced control cells died. Details about the generation of the U-2 OS

Cry1-sKO, U-2 OS *Cry2*-sKO, and U-2 OS *Cry1/Cry2*-dKO cell lines can be found in the

original publication by Börding *et al.*[28].

Bioluminescence recordings

Cells expressing *Bmal1*- or *Per2*-promoter driven luciferase reporters were seeded in 35-mm

dishes (Nunc) to reach approximate confluence on the following day. To account for potentially

different phases of single-cell circadian clocks in cell populations, we collectively reset the

circadian clocks by administering a standard dose of 1 μM dexamethasone[49,50] (Sigma,

dissolved in EtOH). After 30 min incubation cells were washed once with PBS and imaging

medium supplemented with 250 μM D-Luciferin (Abmole) was added. To avoid evaporation

of the media throughout the bioluminescence recordings, the dishes were sealed with parafilm

as described in Finger *et al.*[49]. Luminescence intensity was monitored at 10-min intervals for 5

days in an incubator-embedded luminometer (LumiCycle, Actimetrics). Bioluminescence

recordings were conducted using two biological replicates. For each individual experiment,

three, in some cases only two, technical replicates were assessed. MCF10A cells have been

tested on a single day using three technical replicates.

Long-term live-cell imaging

All live-cell imaging experiments were performed with cells expressing the fluorescent
mKate2-NLS nuclear reporter, seeded in 48-well plates (Falcon) at a density that saturates at
unperturbed growth conditions towards the end of each experiment. Long-term live-cell
imaging was conducted using an incubator-embedded Incucyte live-cell widefield microscope
(Essen BioScience). Cells were imaged in the brightfield and for nuclei segmentation and cell
counting in the red channel (excitation: 567–607 nm, emission: 622–704 nm). Images were
analyzed by frame-by-frame nuclei counting with the in-built Incucyte software and results
were further processed in MATLAB. For experiments that involved drug treatments, cells were
seeded 1 day before starting the live recordings and images were taken using a 4x magnification
lens in 2 fields-of-views per well every 1–2 h for a total duration of 4–6 days. Two independent
plates per condition were assayed in a single experiment. For drug response experiments with
cisplatin a single plate per condition was assayed in a single experiment, imaged in 9 fields-of-
views per well using a 10x magnification lens. For experiments capturing unperturbed growth
dynamics, 300–1000 cells, corresponding to ~10% confluency, were seeded 2 days before
starting the live recordings. Using a 10x magnification lens, 9 images per well were acquired
in 1–2 h intervals for a total duration of 4 days.

Dose-response curves

Drug stock solutions (100–10 mM) were prepared in DMSO and preserved at -20°C. Cisplatin
(Sigma) stock solution of 3.33 mM was prepared in 0.9% NaCl and stored at room temperature.
For dose-response assays, a serial 5–6 point log4 dilution of each drug was freshly prepared in
its solvent prior to treatment. Following concentration ranges were tested: 100–0.1 μM for 5-
fluorouracil (5-FU, Sigma), alpelisib (Biozol) and olaparib (Adooq); 10–0.01 μM for torin2
(Sigma) and alisertib (Hölzel); 10–0.04 μM for adavosertib (Biocat); 1–0.004 μM for
doxorubicin (Hölzel); and 0.4–0.0004 μM for paclitaxel (Hölzel). Cisplatin was tested in a
serial 10-point log2 dilution with doses ranging from 70–0.14 μM. Compounds were added to
the cells one day after cell seeding in a drug-media mixture of 9% of total well-volume.
Solvent-only treated control cells were assayed along the treated conditions. Each tested
condition contained equal amounts of the solvent. Cell growth was monitored by live-cell
imaging for at least 4 days as described above.

Time-of-day treatments

Cells were seeded and allowed to attach overnight. The next day, live recordings commenced as described earlier. We performed independent resetting steps every 4 h over an 8-h period, creating distinct cell populations at 0, 4, or 8 h of circadian time. Subsequently, we administered the same drug concentration, which corresponded to the estimated half-maximal effective concentration, at either 32 or 48 h after the initial resetting step. This allowed us to simultaneously test six different circadian stages (0, 4, 8, 16, 20, and 24 h). Drug concentrations were determined separately for each cell line and drug combination. For cisplatin, the treatments we performed slightly different, using 4 independent resetting steps every 3 hours over a 9-h period followed by drug treatments 32 or 48 h post initial resetting. In all cases, cell growth was continuously monitored through long-term live-cell imaging for a total of 6 days.

Computational methods

Time-series analysis of circadian signals

Detrending. Raw time-series data were detrended by applying a sinc filter with a 48-h cut-off period using the open-source software package pyBOAT[27] (v0.9.1) within the Anaconda Navigator (v1.10.0).

Amplitude envelope and normalization. Continuous amplitude envelope calculation was obtained using continuous wavelet transform implemented in pyBOAT with a time window of 48 hours. The amplitude normalization was done by taking the inverse of the envelope of the detrended signal as described in Mönke G, *et al.* [27] .

Autocorrelation analysis. Periodicity data was assessed by calculating its autocorrelation and abcissa at the second peak using the 'autocorr' and 'findpeaks' MATLAB functions[51] from the detrended time-series.

Continuous wavelet transform. The main oscillatory component, known as the ridge component, was obtained using a wavelet-based spectral analysis from amplitude-normalized and detrended signals. For the ridge detection we used both an adaptable-threshold and a fixed-threshold. The adaptable threshold was based on each signal's half-maximal spectral power and was implemented for the "period" and "phase difference" metric shown in Fig. 2 and Fig. S1. The instantaneous phase difference between the *Bmal1* and *Per2* signals was calculated using the MATLAB 'atan2' function (Supplementary Fig. 1a) and for the polar histogram

representation, we deployed the 'polarhistogram' function and the 'Circular Statistics Toolbox'
(v1.21.9.0. by Philipp Behrens) from MATLAB (Supplementary Fig. 1b). For the analysis of
"amplitudes" and "ridge lengths" shown in Fig. 2 and the determinants for time-of-day
sensitivity shown in Fig. 5, the metrics of the main oscillatory component were derived using
a fixed-ridge threshold of 40. This ridge threshold value was chosen as it offered a well-
balanced threshold suitable for comparing signals of varying strengths.

Multiresolution analysis. Detrended time-series data were decomposed into a set of different
wavelet details D_j representing distinct disjoint frequency bands and a final smooth using a
discrete wavelet transform based multiresolution analysis[52]. The algorithm has been
implemented using the 'PyWavelets' python package[53], with a *db20* wavelet of the Daubechies
wavelet family as previously described in Myung, Schmal and Hong *et al.*[54]. Since each
wavelet detail D_j represents a period range between $2^j\Delta t$ and $2^{j+1}\Delta t$ (for j=1, 2, 3, …) we down-
sampled the time series from a $\Delta t=10$ min to a $\Delta t=30$ min sampling frequency to obtain a
circadian period band between 16–32 h for further analysis. Since the MRA decomposes the
variance of the detrended signal with respect to the different disjoint period bands it can be
used to determine the rhythmicity of the signal in the circadian period range[55].

Global circadian strength. To calculate the global circadian strength (*GCS*) of a cell line
model *i*, the autocorrelation peak value (peak), the wavelet-based continuous ridge length
(ridge) and the discrete circadianicity component (circadianicity) were normalized to the
respective maximum (max) value measured among all tested cell line models and averaged
according to the following equation:

701
$$GCS_i = \text{mean}\left(\frac{\text{peak}_i}{\text{peak}_{\text{max}}}, \frac{\text{ridge}_i}{\text{ridge}_{\text{max}}}, \frac{\text{circadianicity}_i}{\text{circadianicity}_{\text{max}}}\right)$$

702 **Statistical analysis.** Linear regression model fitting of *Bmal1* and *Per2* circadianicity
703 components obtained by MRA was done with the 'fitlm' MATLAB function. Significant
704 variances in circadian parameters between wild-type U-2 OS cells and both, *Cry1*-sKO and
705 *Cry1/2*-dKO cells, were calculated with a one-way ANOVA and Tukey's post-hoc test using
706 the 'anova1' and 'multcompare' MATLAB functions.

Multi-parametric analysis of growth dynamics

Growth data obtained from long-term live-cell imaging was smoothed using a robust local regression approach of weighted linear least squares and a 2^{nd} degree polynomial model ('rloess' MATLAB function). Doubling times of smoothed cell numbers or confluency were calculated according to the following equation:

$$\text{Doubling time}(t) = t * \frac{\log(2)}{\log(y_0/y_t)}$$

where t refers to the time of assessment, in our case 96 h, and y_0 refers to the cell number at timepoint 0. To calculate the exponential growth rate k per unit of time t, we normalized cell numbers to the initial timepoint 0 (y_0) and fitted an exponential function to the growth curves:

$$\text{Growth}(t) = y_0 * e^{(k*t)}$$

Exponential function fitting was done with the MATLAB 'fit' function[51]. Linear regression model fitting to different combinations of growth parameters was done with the 'fitlm' MATLAB function.

Estimation of drug sensitivity parameters

Drug response data obtained from long-term live-cell imaging was smoothed using a robust local regression approach of weighted linear least squares and a 2^{nd} degree polynomial model ('rloess' MATLAB function). Following the method established by Hafner et $al.$[30], we computed the growth rate inhibition (GR) at time t and for each dose c as follows:

$$GR(c,t) = 2^{k(c,t)/k(0)} - 1$$

where $k(c,t)$ is the growth rate under drug treatment and $k(0)$ is the growth rate of untreated cells. Drug response parameters were retrieved by fitting the dose-dependent GR values to a sigmoid curve using the following equation:

$$GR(c) = GR_{\text{inf}} + \frac{1 - GR_{\text{inf}}}{1 + (c/GEC_{50})^{h_{\text{GR}}}}$$

730   where the fitted parameters are as described in Hafner *et al.*[30]. Standard EC_{50} values were

731   calculated by fitting final nucleus counts, normalized to the respective count of the control, to

732   the following sigmoidal function:

$$f(c) = E_{\min} + \frac{1 - E_{\min}}{1 + (c/EC_{50})^h}$$

where $E_{\min}$ corresponds to the minimum response, restricted to values between 0 and 1, and h

is the hill slope of the response curve, constrained to 0.5–10. Sigmoidal function fitting steps

were done with the MATLAB functions 'fit' (GR metrics) or 'lsqnonlin' (EC_{50} value)[51].

Hierarchical clustering analysis was implemented with the MATLAB 'clustergram' function

using an euclidian distance and average linkage method[51]. To account for the different tested

dose ranges across drugs for clustering, GEC_{50} values were normalized to the dose at $GR_{\inf}$.

Pearson's linear correlation coefficients across drug sensitivity metrics were computed using

the MATLAB 'corr' function. For the comparison of cellular growth dynamics to drug doses

evoking the maximum tested effect, we chose growth curves corresponding to the doses closest

to the determined $GR_{\inf}$ values.

Time-of-day sensitivity evaluation

Drug response data of time-of-day treatment experiments was smoothed using a moving

average ('smoothdata' MATLAB function). Final nucleus counts were normalized to the

nucleus count at the respective time of treatment. ToD response data from U-2 OS WT and

Cry1/2-dKO cell lines were obtained from confluency readouts in the brightfield channel.

Time-of-day response curves were generated from the relative final responses of each treatment

timepoint to the final response at timepoint 0 and interpolated using the 'smoothingspline'

function in MATLAB. The smoothing parameter was set to 0.7. The maximum range across

smoothed time-of-day responses (ToD_{MR}) was calculated by subtraction of the minimum from

the maximum relative response. To assess whether the smoothing of response data introduces

artifacts into ToD_{MR} estimates, we performed linear regression analysis between ToD_{MR} values

from raw and spline smoothed data using the 'fitlm' MATLAB function and confirmed a high

positive correlation (**Supplementary Fig. 7a**). ToD_{MR} values were clustered as described for

the drug sensitivity parameters, with missing data points substituted by values from the closest

relevant column, adhering to the 'nearest-neighbor' principle. Polar histograms of the treatment

times with maximum and minimum benefit were generated using the 'polarhistogram'
MATLAB function with 'probability' normalization to depict each bar's height as the fraction
of observations within its bin relative to the total observations.

Correlation and shapley value regression analysis

Linear regressions of cell-intrinsic features and drug-dependent ToD sensitivity have been
obtained using the 'stats' and 'optimize' modules of the 'SciPy' and the 'uncertainties' package
of the Python programming language. Predictions of ToD_{MR} values from new data was based
on fitting a linear regression model to the original data that was associated with Pearson
correlation coefficients ≥ 0.5, utilizing the 'fitlm' MATLAB function. Bland-Altman plots
comparing predicted and actual ToD_{MR} values were generated using the 'Bland-Altman and
Correlation Plot' package (v1.12.0.0) from MATLAB Central File Exchange (2017, Ran
Klein). Live-imaging data yielding growth, drug sensitivity, and ToD sensitivity metrics from
the new cell lines, utilized for the predictions, were based on the confluency readout due to
unviability of nuclear reporter cell lines for most models. Only the drug sensitivity metrics for
MCF7 were based on nuclear counts. The relative importance of the different parameters in a
multiple linear regression model are obtained by Shapley value regression via the Python
'dominance-analysis' package.

Circadian clock gene expression analysis

Gene expression data of circadian clock genes were obtained from the Cancer Cell Line
Encyclopedia Dependency Map (CCLE DepMap, https://sites.broadinstitute.org/ccle/datasets,
Q4 of 2022)[56]. Circadian clock genes of the individual breast cancer cell lines were clustered
using the 'clustermap' plotting function of 'seaborn' with the euclidian distance metric and
complete linkage method. Spearman rank correlation coefficients of circadian clock genes and
drug-specific ToD sensitivity values were computed using the spearman rank correlation
algorithm of the 'SciPy stats' python language module and hierarchically clustered with the
correlation distance metric and single linkage method. Regression plots were obtained using
the 'jointplot' plotting function from 'seaborn', showing drug-specific correlation between the
ToD sensitivities and gene expression of each cell line, including their distribution, squared
spearman rank correlation coefficient, p-value, and a confidence interval of 95%.

Supervised and unsupervised dimensionality reduction

Supervised dimensionality reduction by linear discriminant analysis was performed with the 'LinearDiscriminantAnalysis' function of the 'sklearn discriminant_analysis' python language module. The default parameters were retained, using the exact full singular value decomposition solver and 1 component. Median-based binarization of the drug-dependent ToD sensitivity values were used as target and the resulting linear discriminant vector was plotted using random y-axis units to avoid overlapping of the data points. The robustness of the discriminative information from the LDA was tested using a leave-one-out cross-validation (LOOCV) strategy (**Supplementary Figure 7b**). LOOCV was conducted using the 'model_selection' module from 'sklearn' in Python (v3.9.7) and implemented via the PyCharm Community Edition IDE (v2021.2.2). The contribution to the obtained discriminative information is shown in percentage for each of the circadian clock genes. Unsupervised dimensionality reduction by principal component analysis (PCA) was performed with the 'PCA' function of the 'sklearn decomposition' python language module using the exact full singular value decomposition solver and number of components equivalent to the sample size of the corresponding drug. Further analysis of the most informative PCA components was done by drawing the biplots of the first two principal components and showing the loadings of each circadian clock gene for both components.

Data availability

The experimental data and data tables for all results of this study are available in Figshare with the identifier https://figshare.com/projects/Time-of-Day-Drug-Response/180916.

Code availability

Code used for the data analysis in this work is available on GitHub: https://github.com/Granada-Lab/Time-of-Day-Drug-Response.

Competing interests

The authors declare no competing interests.

References

Chaix, A., Zarrinpar, A. & Panda, S. The circadian coordination of cell biology. *J Cell Biol* **215**, 15-25 (2016). https://doi.org/10.1083/jcb.201603076

Zhang, R., Lahens, N. F., Ballance, H. I., Hughes, M. E. & Hogenesch, J. B. A circadian gene expression atlas in mammals: implications for biology and medicine. *Proc Natl Acad Sci U S A* **111**, 16219-16224 (2014). https://doi.org/10.1073/pnas.1408886111

Mure, L. S. *et al.* Diurnal transcriptome atlas of a primate across major neural and peripheral tissues. *Science* **359** (2018). https://doi.org/10.1126/science.aao0318

Saini, C. *et al.* A functional circadian clock is required for proper insulin secretion by human pancreatic islet cells. *Diabetes, Obesity and Metabolism* **18**, 355-365 (2016). https://doi.org/doi.org/10.1111/dom.12616

Neufeld-Cohen, A. *et al.* Circadian control of oscillations in mitochondrial rate-limiting enzymes and nutrient utilization by PERIOD proteins. *Proc Natl Acad Sci U S A* **113**, E1673-1682 (2016). https://doi.org/10.1073/pnas.1519650113

Chakrabarti, S. & Michor, F. Circadian clock effects on cellular proliferation: Insights from theory and experiments. *Current Opinion in Cell Biology* **67**, 17-26 (2020). https://doi.org/https://doi.org/10.1016/j.ceb.2020.07.003

Scheiermann, C., Kunisaki, Y. & Frenette, P. S. Circadian control of the immune system. *Nature Reviews Immunology* **13**, 190-198 (2013). https://doi.org/10.1038/nri3386

Sancar, A. *et al.* Circadian clock control of the cellular response to DNA damage. *FEBS Lett* **584**, 2618-2625 (2010). https://doi.org/10.1016/j.febslet.2010.03.017

9   Schernhammer, E. S. *et al.* Rotating night shifts and risk of breast cancer in women participating
in the nurses' health study. *J Natl Cancer Inst* **93**, 1563-1568 (2001).
https://doi.org/10.1093/jnci/93.20.1563

10   Puram, Rishi V. *et al.* Core Circadian Clock Genes Regulate Leukemia Stem Cells in AML.
Cell **165**, 303-316 (2016). https://doi.org/10.1016/j.cell.2016.03.015

11   Papagiannakopoulos, T. *et al.* Circadian Rhythm Disruption Promotes Lung Tumorigenesis.
Cell Metab **24**, 324-331 (2016). https://doi.org/https://doi.org/10.1111/dom.12616

12   Chun, S. K. *et al.* Disruption of the circadian clock drives Apc loss of heterozygosity to
accelerate colorectal cancer. *Science Advances* **8**, eabo2389 (2022).
https://doi.org/doi:10.1126/sciadv.abo2389

13   Diamantopoulou, Z. *et al.* The metastatic spread of breast cancer accelerates during sleep.
Nature **607**, 156-162 (2022). https://doi.org/10.1038/s41586-022-04875-y

14   Wang, J. *et al.* Circadian protein BMAL1 promotes breast cancer cell invasion and metastasis
by up-regulating matrix metalloproteinase9 expression. *Cancer Cell International* **19**, 182
(2019). https://doi.org/10.1186/s12935-019-0902-2

15   Lee, Y. *et al.* G1/S cell cycle regulators mediate effects of circadian dysregulation on tumor
growth and provide targets for timed anticancer treatment. *PLOS Biology* **17**, e3000228 (2019).
https://doi.org/10.1371/journal.pbio.3000228

16   Chen, W.-D. *et al.* The circadian rhythm controls telomeres and telomerase activity.
Biochemical and Biophysical Research Communications **451**, 408-414 (2014).
https://doi.org/https://doi.org/10.1016/j.bbrc.2014.07.138

17   Rida, P., Syed, M. I. & Aneja, R. Time will tell: Circadian clock dysregulation in triple negative
breast cancer. *Front Biosci (Schol Ed)* **11**, 178-192 (2019).

18   Innominato, P. F. *et al.* Prediction of overall survival through circadian rest-activity monitoring
during chemotherapy for metastatic colorectal cancer. *International Journal of Cancer* **131**,
2684-2692 (2012). https://doi.org/https://doi.org/10.1002/ijc.27574

19   Ye, Y. *et al.* The Genomic Landscape and Pharmacogenomic Interactions of Clock Genes in
Cancer Chronotherapy. *Cell Systems* **6**, 314-328.e312 (2018).
https://doi.org/https://doi.org/10.1016/j.cels.2018.01.013

20   Lee, Y. *et al.* Time-of-day specificity of anticancer drugs may be mediated by circadian
regulation of the cell cycle. *Science Advances* **7**, eabd2645 (2021).
https://doi.org/10.1126/sciadv.abd2645

21   Dallmann, R., Brown, S. A. & Gachon, F. Chronopharmacology: New Insights and Therapeutic
Implications. *Annual Review of Pharmacology and Toxicology* **54**, 339-361 (2014).
https://doi.org/10.1146/annurev-pharmtox-011613-135923

22 Lévi, F., Okyar, A., Dulong, S., Innominato, P. F. & Clairambault, J. Circadian Timing in
Cancer Treatments. *Annual Review of Pharmacology and Toxicology* **50**, 377-421 (2010).
https://doi.org/10.1146/annurev.pharmtox.48.113006.094626

23 Wood, P. A., Du-Quiton, J., You, S. & Hrushesky, W. J. M. Circadian clock coordinates cancer
cell cycle progression, thymidylate synthase, and 5-fluorouracil therapeutic index. *Molecular*
Cancer Therapeutics **5**, 2023-2033 (2006). https://doi.org/10.1158/1535-7163.MCT-06-0177

24 Damato, A. R. *et al.* A randomized feasibility study evaluating temozolomide circadian
medicine in patients with glioma. *Neuro-Oncology Practice* **9**, 193-200 (2022).
https://doi.org/10.1093/nop/npac003

25 Ruben, M. D., Smith, D. F., FitzGerald, G. A. & Hogenesch, J. B. Dosing time matters. *Science*
365, 547-549 (2019). https://doi.org/doi:10.1126/science.aax7621

26 Kramer, A. *et al.* Foundations of circadian medicine. *PLOS Biology* **20**, e3001567 (2022).
https://doi.org/10.1371/journal.pbio.3001567

27 Mönke, G., Sorgenfrei, F. A., Schmal, C. & Granada, A. E. Optimal time frequency analysis
for biological data - pyBOAT. *bioRxiv*, 2020.2004.2029.067744 (2020).
https://doi.org/10.1101/2020.04.29.067744

28 Börding, T., Abdo, A. N., Maier, B., Gabriel, C. & Kramer, A. Generation of Human CRY1
and CRY2 Knockout Cells Using Duplex CRISPR/Cas9 Technology. *Front Physiol* **10**, 577
(2019). https://doi.org/10.3389/fphys.2019.00577

29 Leise, T. L. Analysis of Nonstationary Time Series for Biological Rhythms Research. *J Biol*
Rhythms **32**, 187-194 (2017). https://doi.org/10.1177/0748730417709105

30 Hafner, M., Niepel, M., Chung, M. & Sorger, P. K. Growth rate inhibition metrics correct for
confounders in measuring sensitivity to cancer drugs. *Nature Methods* **13**, 521-527 (2016).
https://doi.org/10.1038/nmeth.3853

31 Ko, C. H. & Takahashi, J. S. Molecular components of the mammalian circadian clock. *Hum*
Mol Genet **15 Spec No 2**, R271-277 (2006). https://doi.org/10.1093/hmg/ddl207

32 Takahashi, J. S. Transcriptional architecture of the mammalian circadian clock. *Nature Reviews*
Genetics **18**, 164-179 (2017). https://doi.org/10.1038/nrg.2016.150

33 Mormont, M.-C. & Levi, F. Cancer chronotherapy: Principles, applications, and perspectives.
Cancer **97**, 155-169 (2003). https://doi.org/https://doi.org/10.1002/cncr.11040

34 Sulli, G., Lam, M. T. Y. & Panda, S. Interplay between Circadian Clock and Cancer: New
Frontiers for Cancer Treatment. *Trends in Cancer* **5**, 475-494 (2019).
https://doi.org/https://doi.org/10.1016/j.trecan.2019.07.002

35   Lin, H.-H., Qraitem, M., Lian, Y., Taylor, S. R. & Farkas, M. E. Analyses of BMAL1 and PER2
Oscillations in a Model of Breast Cancer Progression Reveal Changes With Malignancy. *Integr*
Cancer   Ther   **18**,   1534735419836494-1534735419836494   (2019).
https://doi.org/10.1177/1534735419836494

36   Lellupitiyage Don, S. S. *et al.* Circadian oscillations persist in low malignancy breast cancer
cells. *Cell Cycle* **18**, 2447-2453 (2019). https://doi.org/10.1080/15384101.2019.1648957

37   Baggs, J. E. *et al.* Network Features of the Mammalian Circadian Clock. *PLOS Biology* **7**,
e1000052 (2009). https://doi.org/10.1371/journal.pbio.1000052

38   Gupta, G. K. *et al.* Perspectives on Triple-Negative Breast Cancer: Current Treatment
Strategies, Unmet Needs, and Potential Targets for Future Therapies. *Cancers (Basel)* **12**, 2392
(2020). https://doi.org/10.3390/cancers12092392

39   Fallahi-Sichani, M., Honarnejad, S., Heiser, L. M., Gray, J. W. & Sorger, P. K. Metrics other
than potency reveal systematic variation in responses to cancer drugs. *Nat Chem Biol* **9**, 708-
714 (2013). https://doi.org/10.1038/nchembio.1337

40   Lévi, F. A. *et al.* Chronomodulated Versus Fixed-Infusion—Rate Delivery of Ambulatory
Chemotherapy With Oxaliplatin, Fluorouracil, and Folinic Acid (Leucovorin) in Patients With
Colorectal Cancer Metastases: a Randomized Multi-institutional Trial. *JNCI: Journal of the*
National Cancer Institute **86**, 1608-1617 (1994). https://doi.org/10.1093/jnci/86.21.1608

41   Sothern, R. B., Lévi, F., Haus, E., Halberg, F. & Hrushesky, W. J. Control of a murine
plasmacytoma with doxorubicin-cisplatin: dependence on circadian stage of treatment. *J Natl*
Cancer Inst **81**, 135-145 (1989). https://doi.org/10.1093/jnci/81.2.135

42   Nagoshi, E. *et al.* Circadian gene expression in individual fibroblasts: cell-autonomous and self-
sustained   oscillators   pass   time   to   daughter   cells.   *Cell*   **119**,   693-705   (2004).
https://doi.org/10.1016/j.cell.2004.11.015

43   Izumo, M., Johnson, C. H. & Yamazaki, S. Circadian gene expression in mammalian fibroblasts
revealed by real-time luminescence reporting: Temperature compensation and damping.
Proceedings   of   the   National   Academy   of   Sciences   **100**,   16089-16094   (2003).
https://doi.org/doi:10.1073/pnas.2536313100

44   Balsalobre, A., Damiola, F. & Schibler, U. A Serum Shock Induces Circadian Gene Expression
in   Mammalian   Tissue   Culture   Cells.   *Cell*   **93**,   929-937   (1998).
https://doi.org/https://doi.org/10.1016/S0092-8674(00)81199-X

45   Bieler, J. *et al.* Robust synchronization of coupled circadian and cell cycle oscillators in single
mammalian cells. *Mol Syst Biol* **10**, 739 (2014). https://doi.org/10.15252/msb.20145218

46   Gérard, C. & Goldbeter, A. Entrainment of the Mammalian Cell Cycle by the Circadian Clock:
Modeling Two Coupled Cellular Rhythms. *PLOS Computational Biology* **8**, e1002516 (2012).
https://doi.org/10.1371/journal.pcbi.1002516

47   Bernard, S., Čajavec Bernard, B., Lévi, F. & Herzel, H. Tumor Growth Rate Determines the
Timing of Optimal Chronomodulated Treatment Schedules. *PLOS Computational Biology* **6**,
e1000712 (2010). https://doi.org/10.1371/journal.pcbi.1000712

48   Zhu, B. *et al.* A Cell-Autonomous Mammalian 12 hr Clock Coordinates Metabolic and Stress
Rhythms.   *Cell   Metabolism   **25**,   1305-1319.e1309   (2017).
https://doi.org/https://doi.org/10.1016/j.cmet.2017.05.004

49   Finger, A.-M. *et al.* Intercellular coupling between peripheral circadian oscillators by TGF-
β   signaling.   *Science   Advances*   **7**,   eabg5174   (2021).
https://doi.org/doi:10.1126/sciadv.abg5174

50   Balsalobre, A. *et al.* Resetting of Circadian Time in Peripheral Tissues by Glucocorticoid
Signaling. *Science* **289**, 2344-2347 (2000). https://doi.org/10.1126/science.289.5488.2344

51   version 9.11.0.1769968 (R2021b) (The MathWorks Inc., 2021).

52   Leise, T. L. & Harrington, M. E. Wavelet-based time series analysis of circadian rhythms. *J
Biol Rhythms* **26**, 454-463 (2011).

53   Lee, G. R., Gommers, R., Waselewski, F., Wohlfahrt, K. & O'Leary, A. PyWavelets: A Python
package for wavelet analysis. *Journal of Open Source Software* **4**

54   Myung, J. *et al.* The choroid plexus is an important circadian clock component. *Nature
Communications* **9**, 1062 (2018). https://doi.org/10.1038/s41467-018-03507-2

55   Leise, T. L. Wavelet analysis of circadian and ultradian behavioral rhythms. *Journal of
Circadian Rhythms* (2013). https://doi.org/10.1186/1740-3391-11-5

56   Barretina, J. *et al.* The Cancer Cell Line Encyclopedia enables predictive modelling of
anticancer drug sensitivity. *Nature* **483**, 603-607 (2012). https://doi.org/10.1038/nature11003

Fig. 1 | Framework for identifying optimal treatment times in cancer and healthy tissue models. Schematic overview of the experimental and computational framework to thoroughly characterize time-of-day drug responses in a variety of cell subtypes, such as cancer and non-malignant cell models. A combination of live recordings is implemented for the deep phenotyping of circadian strength, growth dynamics and drug responses that shape time-of-day profiles. Using a novel streamlined experimental approach, time-of-day sensitivity profiles are obtained in tumor and non-malignant cell models, providing best and worst timings for increased efficacy and reduced toxicity (*top panel*). A tandem computational pipeline integrates the deep phenotyping metrics as well as gene expression data of circadian clock genes to quantitatively address three fundamental questions in chronopharmacology (*bottom panel*). Combining multiple signatures, we define a chronotherapeutic index, ranking cellular models and drug agents by their size-effect gains from drug treatments aligned with the circadian clock (*bottom right panel*).

Fig. 2 | Determining circadian clock strength in cancer and healthy tissue cell models. a, Schematic of the experimental setup to detect circadian oscillations. **b,** Circadian feedback loops involving the core clock genes *Bmal1* and *Per2*, used here as circadian Luc-reporters. **c,** Raw (*left*) and processed (*right*) signals from MDAMB468-*Bmal1/Per2*-Luc cells. **d,** Autocorrelation (AC) of signals shown in **c**. Arrow indicates 2nd peak and corresponding abscissa (lag). Dashed lines=95% CI. **e,** Boxplot of AC values and **f,** corresponding lags at the 2nd peak of signals from various cell models. **g,** Detrended and amplitude-normalized signal (*top*) and corresponding wavelet power spectrum (*bottom*) obtained from continuous wavelet transform (CWT) showing time-resolved periods of signals from MDAMB468-*Bmal1*-Luc cells. Main oscillatory component (ridge) marked as red line. **h,** Boxplot of CWT ridge lengths of signals from the indicated cell models. Bottom and top edges of the boxes in **e**, **f**, and **h** indicate the 25th and 75th percentiles, respectively. Red horizontal lines denote the data's median and red crosses outliers. *n*=6–17 samples, collected from each reporter cell line. **i,** Multiresolution analysis (MRA) of detrended MDAMB468-*Bmal1*-Luc signals. %=fraction of respective component to total signal. **j,** Scatterplot of normalized noise versus circadianicity of signals from the indicated cell models, determined by MRA. Shaded area covers unattainable range. Data represents the mean±s.d. *n*=3–9 samples per reporter cell line. **k,** Bar diagram ranking cell models by their global circadian strength integrating min-max scaled parameters obtained from AC (peak), CWT (ridge) and MRA (circadianicity) for *Bmal1* and, where applicable, *Per2*. Data represents the mean±s.d of the scaled parameters. For representation purposes, only the positive s.d. is shown. *n*=3–6 input parameters. Color coding of data shown in **b–i** and **k** according to the corresponding Luc-reporter for *Bmal1* (yellow) and *Per2* (blue). Color coding of different cell models in **e**, **f**, **h**, **j**, and **k** according to the tissue of origin. Samples were acquired on two independent days, with each day including 2-3 technical replicates per reporter cell line. MCF10A cells and MDAMB468-*Per2*-Luc cells have been assayed in 3 technical replicates on a single day. One-way ANOVA and Tukey's post-hoc test was applied to compare U-2 OS WT and KO cell lines, where *, **, *** and ****, indicate *p*-values ≤ 0.05, 0.01, 0.001, and 0.0001, respectively. n.s.=non-significant.

Fig. 3 | Unravelling growth and drug response dynamics through long-term live-cell imaging. a, Schematic of the setup for capturing cell growth using fluorescent nuclear reporters. **b,** Bi-daily snapshots of MDAMB468 cell growth in the brightfield (*top*) and red fluorescent channel (*bottom*). Ruler = 400 μM. **c,** MDAMB468 confluency (*black line*) and cell numbers (*red line*) over time. **d,** Normalized growth curves of the indicated cell models. BL1, BL2 and MES refer to the TNBC subtypes basal-like 1, basal-like 2, and mesenchymal-like, respectively. EP is short for for epithelial. **e,** Exponential fit (blacl solid line) for MDAMB468 growth curves, yielding the growth rate (k), and fit accuracy (R^2). The dots represent mean normalized cell numbers calculated from two independent assay plates, the shaded area represents the corresponding standard deviation. **f,** Bar diagrams ranking cell models by doubling times and growth rates. Data in **c–f** represents the mean ± s.e.m. from 9 images taken per cell line on a single plate. k-value errors reflect the 95% CI. For clarity in **f**, one-sided error bars are shown, with the arrow indicating the top range of the respective error bar. **g,** Schematic of the pathways targeted by the drugs used in this study. **h,** Cell numbers and corresponding k-values of MDAMB468 cells treated with varying olaparib doses (color-coded) or solvent (dashed line). Data represents the mean ± s.d. of two plates. **i,** Dose-response curve of GR values corresponding to **h**, highlighting various drug sensitivity metrics. Error bars = 95% CI. **j,** Normalized numbers of MDAMB468 cells treated with the GR_{inf} dose of the indicated drugs. Data represents the mean ± s.d. of two plates, or ± s.e.m of 9 images taken per cell line on a single plate (cisplatin). **k–m,** Hierarchical clustering of drug sensitivity parameters across cell-drug combinations. GEC_{50} values in **l** are shown relative to the GR_{inf} dose. **n,** Pearson's linear correlation coefficients of sensitivity parameters shown in **k–m** and additional cell-drug combinations (see Supplementary Table 1; n=49–50). Denoted are significant correlation coefficients, where *, **, ***, and **** indicate p-values ≤ 0.05, 0.01, 0.001, and 0.0001, respectively. Data in **k–n** based on the mean of two plates or 9 images taken per well on a single plate (cisplatin).

Fig. 4 | Time-of-day drug sensitivity is drug and tissue model dependent. a, Schematic of the experimental setup to screen for time-of-day (ToD) responses. Three clock-resetting steps are performed in 4-h intervals. Drugs are administered 32 or 48 h post-initial reset, resulting in six circadian times (0, 4, 8, 16, 20 and 24 h). Growth is monitored by long-term live-cell imaging. Timelines on the right depict experimental (*top*) and relative circadian times (*bottom*), color-coded by each ToD tested. **b,** Cell counts of HCC1937 treated with alisertib at different times of the day. Normalization to respective time of treatment. **c,** Cell numbers 4 days post-treatment versus treatment times, corresponding to **b** (*left*). ToD response curve (ToD-RC) depicting relative responses to ToD 0 h. Blue arrows mark the maximum ToD response range (ToD$_{MR}$) (*right*). **d,** ToD-RCs for HCC1937 (*left*) or MCF10A (*right*) cells treated with different drugs (color-coded). **e,** ToD-RCs for 10 cell models treated with paclitaxel (*left*) or doxorubicin (*right*). Color coding of cell models according to tissue of origin. **f,** Hierarchical clustering of ToD$_{MR}$ values across drug-cell combinations. Values above 0.2 are shown. Data clustering with the UPGMA method and euclidian distance. **g,** Bar diagrams ranking ToD$_{MR}$ values per drug (*top*) and cell model (*bottom*). **h,** ToD-RCs for alisertib-treated HCC1937 tumor and MCF10A non-tumor cells overlaid, yielding maximum and minimum benefit times. Data shown in **b–h** represents mean (± s.d.) of two plates. **i,** Polar histograms for benefit times across cell models (*n*=7–9 drugs) and **j,** drugs (*n*=9 cell models). **k,** Butterfly charts depicting fold changes relative to MCF10A at benefit times, averaged per cancer cell model (*left*) and drug (*right*). Color-coding of maximum and minimum benefit times shown in **h–k** in green and red, respectively. Data shown in **g** and **k** represents mean ± s.d. of tested cell models per drug and vice versa (*see* **f**). For clarity, one-sided error bars are shown.

Fig. 5 | Clock and drug sensitivity metrics shape ToD curves. a, Computational approach to investigate how circadian rhythms, cell growth dynamics, and drug sensitivity factors influence the time-of-day (ToD) drug efficacy. **b,** and **c,** Example of linear correlations between $_{MR}$ and circadian (**b**) or drug sensitivity parameters (**c**), for 5-FU (*n*=10 cell models) and cisplatin (*n*=5 cell models), respectively. Color coding of cell models according to tissue of origin. Grey-shaded area=95% CI. Model accuracy indicated by R^2-values. **d–f,** Hierarchical clustering of correlation coefficients (*r*) between ToD_{MR} values of different drugs (*rows*) and the respective metric (*columns*) for clock strength (**d**, *n*=10 cell models), growth (**e**, *n*=10 cell models) or drug sensitivity (**f**, *n*=5 cell models). *r*-values ≥ 0.5 and ≤ -0.5 are shown. Black boxes indicate examples shown in **b** and **c**. Statistical significance of pairwise correlations is indicated by stars, where *, and **, denote *p*-values ≤ 0.05 and 0.01, respectively. *Note*: sample size in **d–f** for alpelisib=*n*-1; and for cisplatin=9. **g,** Bar diagrams ranking the absolute correlation between ToD_{MR} values and each metric depicted in **d–f**, ranked by metric (*n*=8 drugs) or by drug (*n*=14 metrics). Data represents the mean ± s.e.m. For clarity, one-sided error bars are shown. **h,** Hierarchical clustering of the dominance analysis matrix showing the individual contribution of circadian clock, growth and drug sensitivity parameters (*columns*) in predicting drug dependent ToD_{MR} values (*rows*). Colors indicate the percentage contribution, as detailed in the color bar. See key for **d–f** for sample sizes (*n*). **i,** Boxplot overall contribution of cellular metrics to predict ToD_{MR} values, corresponding to **h**. Black data points indicate the individual result from each drug (*n*=8 drugs). Red horizontal lines denote the median, white circles denote the mean. Bottom and top edges of the boxes indicate the 25th and 75th percentiles, respectively.

Fig. 6 | Gene expression analysis unveils diverse roles of circadian genes in determining time-of-day patterns. a, Approach to identify the role of core clock genes in determining the strength of time-of-day (ToD) drug sensitivity. **b,** Hierarchical clustering of spearman rank correlation coefficients between drug-dependent ToD_{MR} values (*rows*) and core clock genes (*columns*). Black boxes indicate examples shown in **c**. Statistical significance of the correlations are indicated as stars, where * and **, denote *p*-values ≤ 0.05 and 0.01, respectively. Cisplatin and alpelisib: *n*=8 cell lines, else *n*=9 cell lines. **c,** Example of linear correlation analysis between paclitaxel ToD_{MR} values and *Per3* or *Dbp* expression levels measured as log2(TPM). Color-coded data points indicate different cancer cell models. Grey-shaded area=95% CI. Model accuracy indicated by R^2-values. Top and lateral histograms indicate the count of samples along the range of gene expression, and ToD_{MR} values, respectively. The grey continuous line indicates the normal distribution fit of the samples. *n*=9 cell lines. **d,** Linear discriminant analysis (LDA) on median-based binarized ToD_{MR} values for paclitaxel. Cell models with ToD_{MR} values below or above the median (M) are colored in blue and red, respectively (*middle and left panel*). The contribution to the obtained discriminative information is shown in percentage for each of the circadian clock genes (*right panel*). **e,** Boxplot showing the mean-based ranking of overall discriminative contributions of each circadian clock gene across all tested drugs. *n*=8 drugs. Bottom and top edges of the boxes indicate the 25th and 75th percentiles, respectively. Red horizontal lines denote the median, white circles the mean, and red crosses indicate outliers.

Supplementary Fig. 1. a, Example of absolute phase difference evolution between *Bmal1-* and *Per2*-Luc signals, here shown for single samples of MDAMB468 cells. **b,** Polar histogram depicting the phase difference between *Bmal1* and *Per2* signals in MDAMB468 cells over time. 2π denotes one full circadian cycle. Median and mean phase difference are indicated as pink circle and black star, respectively. **c,** Detrended *Bmal1-* (*yellow*) and *Per2*-Luc (*blue*) signals of all cell lines used in this study. Data represents the mean±s.d. of 3 technical replicates recorded on the same day. **d,** Linear regression analysis of the MRA circadian component ('circadianicity') of *Bmal1-* versus *Per2*-Luc signals. Markers indicate different cell models. Grey-shaded area=95% CI. Model accuracy indicated by R^2-values. Data represents the mean±s.d. of 3–9 samples. **e,** *Bmal1-* and **f,** *Per2*-dependent circadian analysis corresponding to the rankings of the combined data shown **Fig. 2e, f, h, k**. Bottom and top edges of the boxes in the box plots indicate the 25th and 75th percentiles, respectively. Red horizontal lines denote the data's median and red crosses outliers. n=3–9 samples. Color-coding of cell models in **c-f** according to their tissue of origin. Samples in **d-f** were acquired on two independent days, with each day including 2-3 technical replicates per reporter cell line. MCF10A cells and MDAMB468-*Per2*-Luc cells have been assayed in 3 technical replicates on a single day.

Supplementary Fig. 2. a, Growth curves of the indicated cell lines monitored by live-cell imaging for 4 days and quantified by either cell numbers (*red line*) or confluency (*black line*). Data represents the mean±s.e.m. of 9 images taken per cell line on a single plate . **b,** Linear regression analysis of doubling times (DT's) calculated from confluency versus cell numbers. **c,** Linear regression analysis of the doubling time and growth rate (*k*) calculated from cell numbers of various cell lines. Color coding of cell models in **b** and **c** according to their tissue of origin. Data shown in **b** and **c** represents the mean±s.d. (DT's), or ±CI of the exponential fit (*k*-values). Grey-shaded area=95% CI. Model accuracy indicated by R^2-values.

Supplementary Fig. 3. Dose-response curves of *GR* values for the specified cell models and drugs. The data is based on the mean responses of technical replicates, live-imaged on independent assay plates for 4 days. For cisplatin, 9 images were taken per cell line on a single plate. Error bars indicate the 95% CI of the exponential fit.

Supplementary Fig. 4. a, Overlay of ToD response curves (ToD-RCs) for individual cell models treated with different drugs (color-coded). **b,** Overlay of ToD-RCs for multiple cell models, grouped by the administered drugs. Cell line models are color-coded. Data in **a** and **b** represents the mean for each drug-cell combination tested on two plates or 9 images (cisplatin). **c,** Overlay of ToD-RCs for U-2 OS wildtype and circadian *Cry1/Cry2*-double knock-out cell lines, grouped by the administered drugs. Cell line models are color-coded. Data represents the mean (± s.d.) of two plates. **d,** Ranking of ToD$_{MR}$ values for each drug-cell combination.

Supplementary Fig. 5. a, Hierarchical clustering of correlation coefficients between *ToD*_{MR} values for different drugs (*rows*) and clock strength metrics (*columns*) obtained from the *Per2*-reporter (*left*) or calculated as a mean from both, *Bmal1*- and *Per2*-reporters (*right*). Statistical significance of pairwise correlations is indicated by stars, where * and **, denote *p*-values ≤ 0.05 and 0.01, respectively. *n*=10 or 9 (cisplatin and alpelisib) cell models. **b,** Bar plots ranking relative overall correlation between *ToD*_{MR} values and individual cellular metrics (depicted in **Fig. 5d–f**), averaged by metric (*left*, *n*=8 drugs) or by drug (*right*, *n*=14 metrics). Positive and negative correlation values are color-coded in light red and blue, respectively. Data represents the mean±s.e.m. **c,** Bland-Altman plots comparing predicted and actual *ToD*_{MR} values of new data that was not used to derive the original statistical associations shown in **Fig 5d-f**, including 26, 8, and 24 new datapoints for the clock, growth, and drug sensivity datasets, respectively (refer to **Supplementary Table 3** for full new datatset). The predictions were based on fitting a linear regression model to the original data associated with pearson correlation coefficients ≥ 0.5. The individual parameters considered for each dataset are distinguished by marker, different drugs are uniformly color-coded across all three datasets. The central solid line indicates the mean bias between the predicted and actual *ToD*_{MR} values of the new data and the outer lines mark the upper and lowers limits of agreement (±1.96*s.d.).

Supplementary Fig. 6. a, Gene expression clustermap of the selected list of core clock genes (*rows*) for the different cell models (*columns*) used in this work. **b,** Boxplot showing the mean-based ranking of absolute spearman rank correlation values between circadian clock gene expression levels and ToD$_{MR}$ values, categorized by drug (*n*=16 circadian clock genes) or by circadian clock gene (*n*=8 drugs). Bottom and top edges of the boxes indicate the 25[th] and 75[th] percentiles, respectively. Red horizontal lines denote the median, white circles the mean, and red crosses indicate outliers. **c-i,** Linear discriminant analysis on median-based binarized ToD$_{MR}$ values for the indicated drugs. ToD$_{MR}$ values below or above the median (M) are color-coded as blue and red, respectively (*middle and left panel of each subfigure*). The contribution to the obtained discriminative information is shown in percentage for each of the circadian clock genes (*right panel of each subfigure*). **j,** Principal Component Analysis (PCA) biplot of the first two principal components on ToD sensitivity groups related to paclitaxel, with ToD sensitivity groups color-coded. Overlaid are the principal component scores for circadian clock genes, scaled to demonstrate their relative contribution to determining ToD sensitivity. The percentage of variance explained by each principal component is denoted by their respective loadings (*left panel*). The bar charts detail the ranked contribution of individual circadian clock genes to PC1 and PC2, presented in descending order (*middle and right panels, respectively*).

Supplementary Fig. 7. a, Linear regression analysis of the maximum range in ToD responses (ToD$_{MR}$), determined from the discrete measured data points or from continuous spline smoothed ToD response curves. Drugs are color-coded. Dotted lines=95% CI. Model accuracy indicated by R^2-values. n=80 drug-cell combinations. **b,** Leave-one-out cross-validation test (LOOCV) on LDA results shown in **Fig. 6d** and **Supplementary Fig. 6d-h**, where the potential of circadian gene expression levels to discriminate between high and low ToD sensitivity groups (based on median-binarized ToD$_{MR}$ values) is investigated. The color coding reflects the accuracy of each cell line model in being correctly assigned to its ToD-sensitivity group, where green and red indicate true and false categorization, respectively. The rightmost column denotes the summed LOOCV accuracy per drug, where 1 refers to 100% accuracy and 0 to 0%. n=9 cell line models; cisplatin and alpelisib: n=8 cell line models.