Tradeoff Between Speed and Robustness in Primordium Initiation Mediated by

Auxin-CUC1 Interaction

Shuyao Kong,[1,2] Mingyuan Zhu,[1,2,3] David Pan,[1,2] Brendan Lane,[4] Richard S. Smith,[4] Adrienne H.

K. Roeder[1,2,5,*]

[1] Weill Institute for Cell and Molecular Biology, Cornell University, Ithaca, NY 14853, USA

[2] Section of Plant Biology, School of Integrative Plant Science, Cornell University, Ithaca, NY

14853, USA

[3] Present address: Department of Biology, Duke University, Durham, NC 27708, USA

[4] Department of Computational and Systems Biology, John Innes Centre, Norwich NR4 7UH, UK.

[5] Lead Contact

[*] Correspondence: ahr75@cornell.edu

ABSTRACT

Robustness is the reproducible development of a phenotype despite stochastic noise. It often

involves tradeoffs with other performance metrics, but the mechanisms underlying such tradeoffs

were largely unknown. An *Arabidopsis* flower robustly develops four sepals from four precisely

positioned auxin maxima. The *development related myb-like 1* (*drmy1)* mutant generates

stochastic noise in auxin signaling that disrupts both the robust position and number of sepal

primordia. Here, we found that increased expression of *CUP-SHAPED COTYLEDON1* (*CUC1*),

a boundary specification transcription factor, in the *drmy1* mutant underlies this loss of

robustness. CUC1 surrounds and amplifies stochastic auxin patches in *drmy1* to form variably

positioned auxin maxima and sepal primordia. Removing *CUC1* from *drmy1* provides time for the

noise in auxin signaling to resolve into four precisely positioned auxin maxima, restoring robust

sepal initiation. However, removing CUC1 decreases auxin maxima intensity and slows down

sepal initiation. Thus, CUC1 increases morphogenesis speed but impairs robustness against

auxin noise. Further, using a computational model, we found that the observed phenotype can be

explained by the effect of CUC1 in repolarizing PIN FORMED1 (PIN1), a polar auxin transporter.

Thus, our study illustrates a tradeoff between speed and robustness during development.

Keywords: Robustness, tradeoff, primordium initiation, sepal, Arabidopsis, auxin, CUC1

INTRODUCTION

Cells within an organism experience a multitude of noise, such as stochastic gene

expression[1] and heterogenous growth rate[2,3]. Despite such noise, organisms often develop

invariantly and reproducibly, a phenomenon termed developmental robustness[4]. How

developmental robustness is achieved is one of the most intriguing open questions in cell biology[5],

which has attracted increasing research efforts in recent years[2,3,6–18]. It was shown that noise in

morphogen signaling can be buffered by certain gene regulatory network structures[13,14] and self-

organized cell sorting[11,15] to achieve robust patterning. Noise in growth rate can be buffered by

averaging growth among neighboring cells[2,3] or in the same cell over time[2] to achieve robust

organ size and shape. Notably, robustness is often involved in tradeoffs with other important

aspects of development[19–21]. For example, during porcine embryogenesis, manual removal of

zona pellucida speeds up development by a few hours but significantly reduces the robustness

of blastocysts development in terms of symmetry and cell size uniformity[21]. How such tradeoffs

are mediated remains largely unknown.

In plants, developmental robustness has been studied in sepals[2,3,16–18,22]. Sepals are the

outermost floral organs that enclose and protect the immature bud before the flower opens. To

achieve this protection, each flower robustly develops four sepals of constant size, positioned

evenly around the bud typical of a cruciferous flower. This robustness in size, number, and

position ensures tight closure critical for protection (Fig. 1a). In contrast, the *development related*

myb-like 1 (*drmy1*) mutant produces 3-5 sepals of different sizes, unevenly positioned, leaving

gaps that expose the inner floral organs (Fig. 1a)[16,17]. This loss of developmental robustness

originates during the initiation of sepals from the floral meristem, where the sepal primordia are

robust in size, number, and position in wild type (WT) but variable in *drmy1* (Fig. 1b)[16]. We have

been studying how DRMY1 maintains robust sepal development. We previously showed that

DRMY1 maintains robustness by increasing TARGET OF RAPAMYCIN (TOR) signaling and

protein translation, which supports the rapid synthesis of A-type ARABIDOPSIS RESPONSE

REGULATOR (ARR) proteins to dampen cytokinin signaling. A proper level of cytokinin signaling

ensures robust patterning of auxin signaling and sepal initiation. In *drmy1*, lack of A-type ARR

proteins and the consequent upregulation of cytokinin signaling increases stochastic noise in

auxin patterning, underlying variable sepal initiation[17]. However, whether there are any tradeoffs

between robustness and other properties of development in this system remains unknown.

Here, we found that a tradeoff exists between robustness and speed of sepal initiation

from the floral meristem. In WT, strong, robust auxin maxima restricts the expression of *CUP-*

SHAPED COTYLEDON1 (*CUC1*), encoding a boundary-specifying transcription factor[23,24], to

precise boundary domains immediately outside the auxin maxima. CUC1 increases the intensity

of auxin maxima it surrounds, and promotes rapid sepal initiation. In *drmy1*, lack of robustness in

auxin patterning causes an expansion of *CUC1* expression. CUC1 amplifies stochastic auxin

noise in the *drmy1* floral meristem, forming variably positioned auxin maxima and sepal primordia.

Removing CUC1 slows down sepal initiation but provides robustness against noise. Thus, the

feedback interactions between auxin and CUC1 promotes rapid organogenesis under low noise

conditions, but disrupts robustness under high noise conditions. Our study thus illuminates the

mechanism behind the tradeoff between robustness and speed during organ initiation.

RESULTS

***CUC1* is upregulated in *drmy1* mutant**

To gain insights into key mechanisms controlling developmental robustness, we

previously performed RNA-seq on inflorescence tissue of *drmy1* vs. WT at the stage of sepal

primordium initiation[17]. We found a 2.4-fold increase in the expression of *CUC1* in *drmy1*, but not

for its paralogs *CUC2* and *CUC3* (Fig. 1c). *CUC* genes encode NAC (NAM, ATAF1,2, CUC2)

family transcription factors important for boundary specification[23,24]. To investigate where *CUC1*

is upregulated within the floral meristem, we imaged the *CUC1* transcriptional reporter. In WT,

CUC1 is expressed in four precisely specified boundaries separating the incipient sepal primordia

from the center of floral meristem and each other, first appearing in the lateral boundaries followed

by the outer and inner boundaries (Fig. 1d). In contrast, in *drmy1*, *CUC1* expression is expanded

and localized to the bud periphery in early stage 2 meristems (Fig. 1e, Supplementary Movie 1).

As the bud develops, this broadened expression coalesces into narrower, WT-like boundary

domains in some parts of the bud (Fig. 1e, arrowheads), but remains in the bud periphery in other

parts (Fig. 1e, brackets), correlated with the presence or absence of sepal outgrowth in stage 3.

drmy1 shows a similar disruption in the protein accumulation pattern of CUC1 (Fig. 1f, g,

Supplementary Movie 2), as well as the expression and protein accumulation patterns of CUC2

(Supplementary Fig. 1).

It was previously shown that auxin inhibits the expression of *CUC* genes and restricts them

to organ boundaries[25–27]. In the WT floral meristem, CUC1 accumulates in four boundaries

immediately outside the four auxin maxima (Supplementary Fig. 2a). In *drmy1*, diffuse, bud

periphery-localized CUC1 colocalizes with diffuse, weak bands of auxin signaling. As these auxin

bands concentrate into variably positioned auxin maxima, CUC1 domains retreat from the bud

periphery and refine into boundaries around the auxin maxima (Supplementary Fig. 2b). These

observations led us to hypothesize that the broadened *CUC1* expression in *drmy1* is due to lack

of robust, concentrated auxin maxima. Consistent with this idea, buds treated with L-Kynurenine

(L-Kyn, inhibitor of auxin synthesis) or Naphthylphthalamic acid (NPA, inhibitor of polar auxin

transport), both of which reduce auxin maxima (Supplementary Fig. 2c, d), show an expansion of

CUC1 expression into the bud periphery (Supplementary Fig. 2e, f). Treatment with both 1-

Naphthaleneacetic acid (NAA) and NPA, which uniformly increases auxin signaling around the

bud periphery (Supplementary Fig. 2d), largely represses *CUC1* expression, and only a weak ring

of *CUC1* expression immediately inside the bud periphery remains (Supplementary Fig. 2f).

These results support the idea that robust, concentrated auxin maxima are required for the

precise boundary expression of *CUC1*, and that *drmy1* shows expanded *CUC1* expression due

to diffuse auxin signaling.

CUC1 upregulation is necessary and sufficient for variable sepal initiation

We next tested the phenotypic consequence of _CUC1_ upregulation, by imaging plants in

which the repression of _CUC1_ by miR164[28–30] has been removed. Specifically, we imaged _miR164_

mutants (_eep1_[30], _mir164abc_[29]), and plants carrying _CUC1_ expression constructs in which the

miR164 target sequence has been mutated (_5mCUC1_[28] and _CUC1m-GFP_[30]). In WT, buds

robustly develop four sepal primordia that are evenly spaced (Fig. 1b, 2a)[16,17]. We found that this

robustness is disrupted in buds upregulating _CUC1_, which, similar to _drmy1_, produce a range of

2-6 sepal primordia that are unevenly spaced and of different sizes (Fig. 2b-e, Supplementary

Fig. 3a-h). In WT, uniform sepal size within each flower is achieved by coordinated initiation timing

between sepal primordia, where inner and lateral sepals initiate within 12 hours of the outer sepal

(Supplementary Fig. 4a, c)[16]. In contrast, in _drmy1_, variability of sepal size originates from the

disorganized initiation timing, where the initiation of the inner and lateral sepals are severely

delayed (Supplementary Fig. 4f, h)[16]. Similar to _drmy1_, we found that in buds upregulating _CUC1_,

the initiation of inner and lateral sepals are greatly delayed relative to the outer sepal, underlying

the variability in size (Supplementary Fig. 4a-c). In addition, the time difference between outer,

inner, and lateral sepal initiation events is more variable between buds (Supplementary Fig. 4c).

Overall, these results suggest that _CUC1_ overexpression is sufficient to disrupt the robustness in

number, position, and coordinated initiation timing of sepal primordia.

Our results show that _CUC1_ overexpression is sufficient for disrupting robustness in sepal

initiation, but is it also necessary? We found that the _drmy1 cuc1_ double mutant often robustly

develops four sepal primordia that are evenly spaced, rescuing the variability in sepal number

and position in _drmy1_ (Fig. 2f-k). The result is specific to _cuc1_ since the _drmy1 cuc2_ double mutant

exhibits disrupted robustness similar to _drmy1_ (Supplementary Fig. 3i-n). While the _cuc1_ mutation

restores robustness in sepal primordia number and position in _drmy1_, it does not restore

robustness in sepal primordia size and coordination of initiation timing (Supplementary Fig. 4d-

h), suggesting that other mechanisms also contribute to these defects. Overall, these results show

that *CUC1* upregulation causes the variability of sepal primordium number and position in *drmy1*.

CUC1 increases auxin maxima intensity and facilitates rapid sepal initiation

CUC1 functions redundantly with CUC2 and CUC3 in organ separation[31,32], and the *cuc1*

cuc2 double mutant shows sepal fusion in stage 8 flowers[31]. However, *cuc1 cuc2* flowers robustly

initiate four sepal primordia at stage 4, suggesting that they are dispensable for sepal initiation.

Then why is *CUC1* still expressed so early on in the floral meristem, when it has the potential to

disrupt robustness when dysregulated (Fig. 2)? Comparing buds of similar size (as an indicator

of similar developmental progression), we found that overexpression of *CUC1* makes the outer

sepal primordium initiate earlier from the floral meristem (Fig. 3a-c). This increased speed in outer

sepal initiation correlated with an increase in auxin signaling (Fig. 4a, b). Mutation of *CUC1* delays

the initiation of all four sepals relative to bud size (Fig. 3d-f), correlated with weaker auxin signaling

maxima (Fig. 4a, b). These results suggest that CUC1 increases auxin maxima intensity, which

in turn promotes rapid sepal initiation to promptly cover and protect the developing floral meristem.

This beneficial role may explain why *CUC1* is expressed in the early-stage floral meristem despite

its potential in reducing developmental robustness when dysregulated, and suggests a potential

conflict between speed and robustness in sepal initiation.

CUC1 amplifies sporadic auxin noise into variably positioned auxin maxima

How does *CUC1* mislocalization disrupt developmental robustness in *drmy1*? There is

evidence that feedback interaction between CUC and auxin plays an important role in plant

morphogenesis[25], and we previously showed that the disrupted pattern of auxin signaling

underlies variable sepal initiation in *drmy1*[16]. Our results illustrated the first half of the feedback

loop where auxin regulates CUC1, by showing that diffuse auxin signaling underlies the

mislocalization of CUC1 expression in *drmy1* (Supplementary Fig. 2). Here, we hypothesized the

second half of the feedback loop where CUC1 regulates auxin, and tested whether CUC1

mislocalization disrupts robustness in auxin pattern in *drmy1*. In WT, four auxin maxima form

robustly, marking four incipient sepal primordia (Fig. 4a, Outer, Inner, Lateral, Lateral). We found

that this robustness is disrupted in *5mCUC1* (CUC1 overexpression), as additional auxin maxima

form (Fig. 4a, arrow). The *cuc1* single mutant forms four auxin maxima robustly, although weaker

(Fig. 4a, b). The *drmy1* single mutant, where CUC1 is upregulated and mislocalized, shows

diffuse, noisy bands of auxin signaling (Fig. 4a, brackets). Removing CUC1 from *drmy1* restores

four robustly positioned auxin maxima (Fig. 4a, c-d). Overall, these results are consistent with the

second half of the feedback loop where CUC1 mislocalization disrupts robustness in auxin

pattern.

How is robust auxin pattern disrupted when *CUC1* is upregulated, and restored when

CUC1 is removed? To address this question, we live imaged buds of WT, *5mCUC1*, *cuc1*, *drmy1*,

and *drmy1 cuc1* every 6 hours from late stage 1 to early stage 3 (Fig. 4e-n). Initially, WT buds

show three robustly positioned auxin maxima. One of them is in the cryptic bract (a suppressed

inflorescence leaf), a remnant of floral meristem initiation[33–36]. The other two auxin maxima appear

in the incipient lateral sepals. These auxin maxima are later followed by two more in the incipient

outer and inner sepals (Fig. 4e, j, Supplementary Movie 3). In *5mCUC1*, some buds initially form

WT-like pattern of four auxin maxima, followed by additional ones in between (Fig. 4f, k, arrows,

Supplementary Movie 4). Other buds display more dynamic spatiotemporal changes in auxin

maxima localization (Supplementary Fig. 5). This suggests that *CUC1* overexpression disrupts

auxin patterning by inducing the formation of new auxin maxima and making them more dynamic.

In *cuc1*, auxin maxima sequentially and robustly form like in WT, although weaker (Fig. 4g, l,

Supplementary Movie 5). In *drmy1*, auxin signaling initially appears as diffuse, noisy bands with

sporadic patches of cells having stronger signal than neighboring cells which fluctuates over time

(Fig. 4h, m, brackets). These sporadic patches seed the subsequent formation of variably sized

and positioned auxin maxima (Fig. 4h, m, arrowheads). As the bud further expands, additional

auxin maxima form, similar to *5mCUC1* (Fig. 4h, m, arrows, Supplementary Movie 6). In *drmy1*

cuc1, auxin signaling initially accumulates in sporadic patches, but unlike *drmy1*, they fade away,

allowing the subsequent formation of the robust auxin pattern (Fig. 4i, n, asterisk, Supplementary

Movie 7). Our results suggest that increased *CUC1* expression disrupts robust auxin patterning

in *drmy1* by amplifying sporadic auxin noise to form variably positioned auxin maxima.

We next tested whether CUC1 amplifies sporadic auxin noise from sources other than the

drmy1 mutation. It was previously shown that exogenous cytokinin alters patterns of polar auxin

transport[37,38], causing sporadic patches of PIN convergence and auxin signaling[16]. We

hypothesized that the cytokinin induced sporadic patches of auxin signaling would not be

amplified in the *cuc1* mutant, and over-amplified in *CUC1* overexpression (*5mCUC1*) buds.

Indeed, WT buds treated with the synthetic cytokinin 6-Benzylaminopurine (BAP) amplifies

sporadic auxin patches to form variably positioned auxin maxima and sepal primordia (Fig. 5a, b,

g-k). While BAP induces sporadic auxin noise in early stage 2 buds of *cuc1*, it quickly fades away,

and most buds robustly form four sepal primordia (Fig. 5c, d, g-k). In contrast, *5mCUC1* buds

treated with BAP form numerous auxin maxima that often connect into a ring, which grow into 3-

8 primordia, more variable than the mock-treated, and often with mixed sepal-petal identity (Fig.

5e-k). Overall, these data support the idea that, under conditions that increase noise in auxin

patterning, such as *drmy1* or under exogenous cytokinin treatment, CUC1 amplifies the noise to

form variably positioned auxin maxima, which in turn disrupts robustness in sepal initiation. In the

absence of CUC1, these noisy patches are not amplified and quickly fade away, leaving four

robust auxin maxima.

Modeling predicts CUC1 amplifies auxin noise by repolarizing PIN

It was previously reported that *CUC* genes promote PIN polarity[25,39]. We wondered

whether an increase in PIN repolarization could explain the noise-amplifying effect of *CUC1*. To

test this, we implemented a computational model of auxin pattern formation (Supplementary

Dataset 2). The floral meristem was modeled using a 2D growing disk of cells, where cells divide

when a size threshold is met. In each cell, PIN1 transports auxin to neighboring cells with highest

auxin concentration. This up-the-gradient auxin transport has been shown to generate auxin

maxima patterns from a homogeneous initial condition with small perturbations[40–42] and is a

potential mechanism of auxin noise amplification. CUC1 is produced in cells with low auxin

concentration. The sole function of CUC1 in the model is to increase PIN repolarization. Below a

CUC1 concentration threshold, PIN repolarizes linearly according to the auxin concentration of

neighboring cells; above the threshold, PIN repolarizes quadratically, and is thus more sensitive

to auxin concentration differences among neighboring cells (Fig. 6a). We hypothesize that this

function of CUC1 in increasing PIN repolarization can by itself promote auxin maxima formation

while amplifying auxin noise. Starting with a patternless disk of cells with small fluctuations in

auxin production rate, auxin maxima form, which are then extracted and quantified (Fig. 6b). We

modeled the *drmy1* mutation by increasing the amplitude of fluctuation in auxin production rate,

which recreates the sporadic auxin patches and PIN convergence points observed in *drmy1*[16]

(Fig. 6c). We modeled the *cuc1* mutant by eliminating CUC1 production (Fig. 6c). We found that

in both WT and *drmy1* models, having CUC1 results in stronger, more concentrated auxin maxima

that form more rapidly compared to *cuc1* and *drmy1 cuc1* respectively (Fig. 6d-h). These modeling

results are similar to real buds (Fig. 4a, b). In both the model and the data, while *drmy1* amplifies

auxin production noise, creating variability in the final auxin pattern (Fig. 6f, arrowheads), *drmy1*

cuc1 shows a relatively robust final pattern despite initial noise (Fig. 6g, arrows, 6i, Supplementary

Movie 8). These simulations suggest that the previously demonstrated function of CUC1 in

increasing PIN repolarization is sufficient for its role in promoting rapid, strong auxin maxima

formation while disrupting robustness against high auxin noise.

In addition to an increase in PIN repolarization which transports more auxin up the

concentration gradient, noise may also be amplified if a cell rapidly grows and divides, forming

two daughter cells that inherit the same auxin noise, before noise is dampened by transport or

decay (Supplementary Fig. 6a). To test this idea, we examined the effects of tissue growth rate

(which correlates with cell division rate in the model) on robustness of auxin patterning. We found

that increasing growth rate increases variability of auxin maxima in both WT and *drmy1*, while

reducing growth rate restores robust auxin patterning in *drmy1* (Supplementary Fig. 6b, d). In

drmy1 simulations under reduced growth rate, the initial stochastic noise in auxin concentration

fades, allowing the formation of robust auxin pattern (Supplementary Fig. 6c), similar to

simulations of the *drmy1 cuc1* double mutant (Fig. 6g). This finding supports the idea that speed

and robustness can be conflicting sides of pattern formation during development.

It was previously shown that spatiotemporal averaging of cell heterogeneity can underlie

tissue-wide developmental robustness[2,43,44]. We hypothesize that reduced PIN sensitivity to

fluctuating auxin levels in neighboring cells (Fig. 6g) or reduced growth rate (Supplementary Fig.

6b, c) restores robust auxin patterning because they allow more time for auxin noise to average

to concentrations similar to nearby cells. We deduce that setting auxin noise temporally (but not

spatially) unchanging would eliminate this averaging (Supplementary Fig. 7a), and thus mutating

cuc1 or reducing growth rate would no longer rescue the *drmy1* patterning defect. Indeed, when

noise is set temporally unchanging, both *drmy1* and *drmy1 cuc1* shows stabilization of initial auxin

noise into variably positioned auxin maxima, as does *drmy1* under reduced growth rate

(Supplementary Fig. 7b-d). Overall, these results support our idea that increased CUC1 disrupts

robustness in auxin patterning by increasing the sensitivity of PIN to fluctuating auxin levels, which

hinders temporal noise averaging and promotes noise amplification.

DISCUSSION

Developmental robustness has fascinated biologists for over 80 years[45], yet the underlying

mechanisms have just begun to be explored[2–4,6–18]. Here, we elucidated a mechanism through

which the developmental robustness of sepals is shaped by DRMY1-auxin-CUC1 interaction (Fig.

6j). In the floral meristem, DRMY1 maintains robust auxin patterning[16] which in turn restricts the

expression of *CUC1* to precise boundary domains adjacent to auxin maxima (Fig. 1d, f,

Supplementary Fig. 2). CUC1 increases the intensity of auxin maxima and promotes rapid sepal

initiation (Fig. 3, 4a, b). In the *drmy1* mutant, diffuse, noisy auxin signaling causes an upregulation

and expansion of the *CUC1* expression domain (Fig. 1e, g, Supplementary Fig. 2). CUC1

amplifies sporadic auxin patches to form variably positioned auxin maxima and sepal primordia

(Fig. 4, 5). In the *drmy1 cuc1* double mutant, auxin fluctuations have time to average out, allowing

robust auxin pattern formation (Fig. 6, Supplementary Fig. 7) and robust sepal initiation (Fig. 2).

In an Arabidopsis flower, sepals enclose and protect the inner, developing floral organs

before the flower opens. To achieve this function, they need to not only rapidly initiate from the

floral meristem to promptly cover it, but also develop robustly so as to not leave any gaps (Fig.

1a). We found that these two traits, speed and robustness, are conflicting sides of sepal

development, and that this tradeoff is mediated by auxin-CUC1 interaction. CUC1 promotes

strong auxin maxima formation and rapid sepal initiation, but also stabilizes auxin noise and can

therefore disrupt robustness under high auxin noise (Fig. 6j, left). On the other hand, lack of CUC1

slows down sepal development, but also allows time for noisy auxin signaling to robustly converge

(Fig. 6j, right).

Our computational modeling suggest that this speed-robustness tradeoff can be fully

explained by the previously reported function of CUCs in increasing PIN polarity[25,39]. How CUCs

increase PIN polarity remains unknown. It was shown that PIN polarity can be regulated by

phosphorylation (e.g., PID[46], D6PK[47], and PP2A[48]) or membrane trafficking (e.g., ABCB19[49] and

ROP2[50]). Thus, CUCs may increase PIN polarity by changing the expression of these important

PIN regulators. Alternatively, CUCs may also inhibit growth, causing mechanical conflict with

adjacent fast-growing regions which alters PIN polarity[51]. Further study is needed to test whether

CUCs increase PIN polarity by any of these mechanisms.

Is morphogenesis speed always involved in a tradeoff with robustness? Can a system

achieve both aspects simultaneously? Earlier theoretical studies on the mammalian olfactory

epithelium suggest that tissue regeneration regulated by a feedback loop cannot achieve both

speed and robustness, unless a second feedback loop is involved[19,20]. Similarly, a recent study

on Hedgehog signaling shows that coupled extracellular and intracellular feedback loops

mediated by PTCH provides both robustness and speed of signaling gradient formation,

compared to single or uncoupled feedback loops[13]. In summary, at the cost of additional

regulatory mechanisms, morphogenesis speed and robustness may be both achieved

simultaneously.

MATERIALS AND METHODS

Plant material

Most Arabidopsis plants were in Col-0 background (WT). Plants presented in

Supplementary Fig. 1, 3a-c, 3e-f, and 7e were in Ler background. *mir164abc* was in a mixed Ler-

Col background[29]. *drmy1* (Ler) was generated by backcrossing *drmy1* (Col-0) with Ler twice. The

following lines were provided by the Arabidopsis Biological Resource Center: *pCUC2::3xVENUS-*

N7 (CS23891)[52], *pCUC2::CUC2-VENUS pPIN1::PIN1-GFP* (CS67929)[52], *pCUC1::CUC1m-GFP*

(CS65830)[30], *cuc1-13* (SALK_006496C), *cuc2-3* (CS875298), *eep1* (CS65826)[30], and *mir164a-4*

mir164b-1 mir164c-1 (CS65828)[29]. In addition, the following lines were previously described:

drmy1-2[16], *5mCUC1*[28], *pCUC1::3xVENUS-N7*[29], *pCUC1::CUC1-GFP*[53], *DR5rev::3xVENUS-N7*[52],

DR5rev::ER-mRFP1.2[54], *35S::mCitrine-RCI2A*[16]. *pCUC1::3xVENUS-N7* in Col-0 background

(Fig. 1d-e) was generated by transforming Col-0 plants with the *pCUC1::3xVENUS-N7* construct.

Plant growth conditions

Seeds were sown in wetted Lambert LM-111 soil and stratified at 4°C for 3-5 days. Plants

were grown under 16 h – 8 h light-dark cycles (fluorescent light, 100 µmol m^{-1} s^{-1}) at 22°C in a

Percival walk-in growth chamber.

Flower staging

Flower buds were staged as previously described[55].

Confocal microscopy

Confocal imaging was done as previously described[16,17]. Briefly, inflorescences were cut

and dissected with a Dumont tweezer (Electron Microscopy Sciences, style 5, no. 72701-D) down

to stage 9, inserted upright into a small petri dish (VWR, 60 x 15 mm) containing inflorescence

culture medium (1/2 MS, 1% (w/v) sucrose, 1x Gamborg vitamin mixture, 0.1% (v/v) plant

preservative mixture (Plant Cell Technology), 1% (w/v) agarose, pH 5.8), further dissected down

to stage 6 (for static imaging) or stage 2 (for live imaging), immersed with water, and imaged

under a Zeiss710 upright confocal microscope with a 20x Plan-Apochromat water-dipping lens

(1.0 NA). For static imaging, to visualize tissue morphology, samples were stained with 0.1 mg/ml

propidium iodide (PI) for 5 minutes before imaging. For live imaging, dissected samples were put

in a 24 h-light growth chamber (fluorescent light, 100 μmol m^{-1}s^{-1}) between time points. To prevent

bacterial growth, every 2-3 days, samples were transferred onto fresh media and treated with 100

μg/ml Carbenicillin (GoldBio, C-103-5, lot # 0129.091814A).

The following lasers and wavelengths were used. Chlorophyll, excitation 488 or 514 nm,

emission 660-722 nm (when also imaging mRFP1.2) or 647-721 nm (others). Propidium iodide,

excitation 514 nm, emission 590-660 nm. mRFP1.2, excitation 561 nm, emission 582-657 nm.

mCitrine, excitation 514 nm, emission 519-580 nm. For VENUS, in *DR5::3xVENUS-N7* and

pCUC2::CUC2-VENUS, excitation 514 nm, emission 519-558 nm; in *pCUC1::3xVENUS-N7*,

excitation 514 nm, emission 518-578 nm; in *pCUC2::3xVENUS-N7*, excitation 488 nm, emission

493-550 nm. GFP, excitation 488 nm, emission 493-556 nm.

Image processing

Image processing was done as previously described[16,17] and also briefly described below.

Tissue morphology was visualized by taking screenshots of the Chlorophyll, PI, or

35S::mCitrine-RCI2A channels in MorphoGraphX[56] or by 3D rendering in the ZEN software

(Processing → 3D). For Fig. 3 and Supplementary Fig. 4, to aid visualization of sepal initiation,

Gaussian curvature heatmaps were calculated in MorphoGraphX as follows: Gaussian blur (X/Y/Z

sigma = 1 μm twice and then X/Y/Z sigma = 2 μm once), edge detection (threshold = 2000-8000

depending on brightness, multiplier = 2.0, adapt factor = 0.3, fill value = 30000), marching cube

surface (cube size = 8 μm, threshold = 20000), change lookup table to "jet", subdivide mesh,

smooth mesh (passes = 5), subdivide mesh, smooth mesh (passes = 5), and project mesh

curvature (type = Gaussian, neighborhood = 10 µm, autoscale = no, min curv = -0.002, max curv

= 0.002). A sepal primordium is considered initiated when it emerges as a deep red band (positive

curvature) separated from the floral meristem by a deep blue band (negative curvature).

For circular histograms and kymographs of DR5 signal, each stack was cropped in ImageJ

and trimmed using the Voxel Edit function in MorphographX[56] so that only the focal bud remained

in the stack. The bud was positioned so that it was centered, facing the Z direction, and the

incipient outer sepal was at 45°. A circular histogram was calculated using the function Export

Histogram Circular, summing signal (in voxel intensity units, 0-255) in each 1° sector around the

Z axis starting from 0° (between the incipient outer and lateral sepals) counterclockwise. These

histograms were then 4°-binned and used for plotting mean ± SD and kymographs. Total signal

was calculated by summing all the bins.

Quantification of developmental robustness

For variability in sepal primordium position, within each bud, an angle was measured

between each pair of adjacent sepal primordia with respect to the bud center. CV was calculated

within each bud as a measurement of how evenly sepal primordia are distributed around the bud.

For relative initiation timing of sepals within each bud, we considered that the robust

temporal sepal initiation pattern in WT consists of the inner sepal initiating within 6 hours of the

outer sepal, and the lateral sepal initiating within 12 hours of the outer sepal[16]. Severely delated

inner and lateral sepal initiation indicate loss of robustness. Thus, for each genotype, we

calculated, among all buds, the mean and SD of initiation timing difference between inner and

outer sepals, and between lateral and outer sepals. Robustness is considered lost if sepal

initiation does not follow the WT temporal pattern (mean is greater than 6 and 12 respectively),

and also if different buds have different temporal patterns (SD is large).

For sepal initiation timing relative to bud size, bud area (μm^2) in maximum intensity

projection images were measured in ImageJ[57,58] and used as a proxy for bud size. Representative

images of WT vs. *cuc1* (both carrying 35S::mCitrine-RCl2A, a plasma membrane marker) that

are of similar area and growth rate throughout the time series were shown. Similarly for WT vs.

5mCUC1, although they did not have a membrane marker and thus the Chlorophyll channel was

used for area measurement. Bud size at the time of outer, inner, and lateral (WT vs *cuc1*) sepal

initiation was plotted. For *5mCUC1*, not enough samples initiated lateral sepals, and thus bud

size at the time of lateral sepal initiation was not analyzed. We reasoned that buds whose sepals

initiate rapidly should be smaller when they initiate sepals, compared to buds whose sepals take

longer to initiate. To control for the gradual reduction in size of successive buds during prolonged

in vitro culture, only buds that did not have any sepals at the first time point and produced at least

one sepal at or before the fourth time point (within 18 hours of the first time point) are analyzed.

***In vitro* drug treatments on inflorescence samples**

For BAP treatment in Fig. 5, inflorescences were dissected, put onto an inflorescence

culture medium (see above) containing 0.05% DMSO and 1 µM BAP (6-Benzylaminopurine, Alfa

Aesar, A14678). Mock contained just 0.05% DMSO. They were left in a growth chamber for 32

hours and then transferred onto inflorescence culture medium without treatments, and imaged at

0 h, 24 h, 48 h, and 72 h after the transfer. For BAP treatment in Supplementary Fig. 7e,

inflorescences were cultured on media containing 0.01% DMSO and 5 µM BAP (mock contained

just 0.01% DMSO) for 96 h, and PI-stained and imaged.

For L-Kyn treatment, inflorescences were dissected and put onto an inflorescence culture

medium containing 0.02% DMSO and 80 µM L-Kyn (l-kynurenine, Sigma, K8625). Mock

contained just 0.02% DMSO. They were left in a growth chamber for 4 days before imaging.

For NPA and NPA+NAA treatment, inflorescences were dissected and put onto

inflorescence culture medium. The following solutions were made in water: for mock, 0.05%

DMSO, 0.01% Silwet L-77; for NPA, 100 µM NPA (Naptalam, Sigma, 33371), 0.05% DMSO,

0.01% Silwet L-77; for NPA+NAA, 100 µM NPA, 20 µM NAA (1-Naphthaleneacetic acid, Sigma,

N0640), 0.05% DMSO, 0.01% Silwet L-77. These solutions were applied on top of the dissected

inflorescences for 24 hours. Then, the solutions were discarded, and samples were washed three

times with sterile water. Images were taken 48 hours (for *DR5::3xVENUS-N7*) or 72 hours (for

pCUC1::3xVENUS-N7) after the end of the treatment.

Mass-spring model of a growing floral meristem

We modeled the stage 2 floral meristem as a 2D disk of growing and dividing cells. The

simulation begins as a round disk of 73 cells and 348 walls with no auxin, no CUC1, and apolar

PIN. As detailed below, in each iteration, the following processes are run in order: deformation of

cell walls under turgor pressure (repeated until convergence); dilution of auxin and CUC due to

changes in cell size; updating noise in auxin production; 10 steps of chemical interactions; cell

division; splitting of cell walls longer than 1 µm; cell growth; reinitialization; data output. The result

of each iteration was used as the starting point of the next. Simulations were run for the desired

number of iterations. Screenshots and data were saved every 10 iterations.

For tissue mechanics, a mass-spring model was used. Cell walls are subdivided to have

a maximum length of 1 µm. Wall segments are represented by springs in the simulation, with an

initial resting length the same as in the starting configuration. Uniform turgor pressure is simulated

by assigning a normal force to the boundary walls, as it cancels out on interior walls. The force

acting on a vertex v due to the springs was calculated as:

423
$$F_v = \sum_{n \in N_v} k \left(\frac{\|p_n - p_v\|}{L_{v:n}} - 1 \right) \frac{p_n - p_v}{\|p_n - p_v\|}$$

where N_v are neighboring vertices of vertex v, p_v is the position of vertex v, p_n is the position of a

neighboring vertex n, k is the spring constant (stiffness), and $L_{v:n}$ is the resting length of the spring

joining v and n[59,60]. Stiffness is set uniform and constant for all the cell walls. The resulting system

of equations is solved using the backward Euler method with the GPU based stabilized bi-

conjugate gradient solver available in MorphoDynamX. Growth was simulated by increasing the

rest length of the springs based on their relative stretch, multiplied by an extensibility factor g:

430
$$\frac{dL_{v:n}}{dt} = g\left(\frac{\|p_n - p_v\| - L_{v:n}}{L_{v:n}}\right)$$

Within each simulation, extensibility were set constant and uniform for all cell walls. Extensibility

was changed when modeling meristems with increased or decreased growth rate.

The molecular interactions are that (1) auxin represses CUC1 expression, and (2) CUC1

increases the sensitivity of PIN repolarization to auxin concentration differences among

neighboring cells. To simulate them, each cell is assigned two fields, auxin concentration (aux)

and CUC concentration (cuc). Each wall segment is assigned two fields, pin_P and pin_N, denoting

the amount of PIN protein on each side that mediates polar auxin transport in either direction. At

each chemical step, three calculations take place concomitantly:

(a) For each cell i at time step t, the amount of PIN on its wall facing neighboring cell j is

calculated as:

441
$$pin_{i \to j}(t) = (1 - \alpha)pin_{i \to j}(t - 1) + \alpha \frac{aux_j^n L_j}{\sum_{k \in N_i} aux_k^n L_k}$$

442
$$pin_{i \to j}(0) = \frac{L_j}{\sum_{k \in N_i} L_k}$$

443   where $\alpha = 0.01$ is the PIN repolarization speed, N_i is all the neighboring cells of cell i, L_j and L_k

444   are lengths of cell walls of cell i facing neighboring cells j and k respectively, and aux_j and aux_k

445   are the auxin concentrations of cell j and k respectively. n is the PIN sensitivity factor dependent

446   on the CUC concentration of cell i, cuc_i, and the threshold CUC concentration, $cuc_{thres} = 2$:

447
$$n = \begin{cases} 1 \; (cuc_i < cuc_{thres}) \\ 2 \; (cuc_i > cuc_{thres}) \end{cases}$$

Note that the total amount of PIN in each cell sums to 1 over all its cell walls, unaffected by auxin

concentration. Also note that a zero-flux boundary condition was used, i.e., no PIN is polarized

towards the boundary of the modeled tissue, and there is no flux of auxin across the boundary.

(b) Change in auxin concentration in cell i due to production, decay, and transport

$$\frac{d(aux_i)}{dt} = Prod_{aux} \times \delta_{aux,i} - Dec_{aux} \times aux_i + \frac{Tran_{aux} \sum_{j \in N_i}(aux_j \times pin_{j \rightarrow i} - aux_i \times pin_{i \rightarrow j})}{Area_i}$$

where $Prod_{aux} = 1$ is the auxin production coefficient, $Dec_{aux} = 0.2$ is the auxin decay coefficient,

$Tran_{aux} = 400$ is the polar auxin transport coefficient, N_i are the neighboring cells of i, $Area_i$ is

the area of cell i, $pin_{i \rightarrow j}$ is the amount of PIN on the wall of cell i facing neighbor j, and $pin_{j \rightarrow i}$ is

the amount of PIN on the wall of neighbor j facing cell i. $\delta_{aux,i}$ is the auxin production noise of cell

i, drawn during initialization and at each iteration from a Gaussian distribution $N(1, SD_{aux})$ where

$SD_{aux} = 0.1$ for WT and *cuc1*, and $SD_{aux} = 1$ for *drmy1* and *drmy1 cuc1*, and negative values are

set to 0. If the noise is set temporally unchanging (Supplementary Fig. 7), δ_{aux} values are drawn

only during initialization but not at each iteration.

(c) Change in CUC concentration in cell i due to production and decay

$$\frac{d(cuc_i)}{dt} = Prod_{cuc} \times \frac{1}{1 + (\frac{aux_i}{K_{aux}})^{hill}} - Dec_{cuc} \times cuc_i$$

where $Prod_{cuc}$ is the CUC production coefficient (1 for WT and *drmy1*, and 0 for *cuc1* and *drmy1*

cuc1), $Dec_{cuc} = 0.2$ is the CUC decay coefficient, $K_{aux} = 5$ is the concentration of auxin at which

CUC production is halved, and $hill = 4$ is an arbitrary Hill coefficient.

For cell division, a cell divides when its area passes a threshold of 50 µm^2. Position of the

new wall follows the minimal wall length principle, with noise parameters as follows (constant for

all simulations): cell division noise 2.0, cell center noise 2.0, wall junction noise 2.0. Daughter

cells inherit the same aux, cuc, and δ_{aux}. Split cell walls inherit pin_P and pin_N proportional to their

new lengths. New cell walls are assigned infinitesimal (non-zero) starting values of pin_P and pin_N.

Software

Image processing was done in ImageJ (version 2.14.0/1.54f, build c89e8500e4)[57,58] and

MorphoGraphX (version 2.0, revision 1-354, CUDA version 11.40)[56]. Modeling was implemented

in C++ using vlab (version 5.0, build #3609)[61] and MorphoDynamX (version 2.0, revision 2-1395,

CUDA version 11.40; www.MorphoDynamX.org) in an Ubuntu 20.04.6 LTS system, equipped with

an Intel Core i9-10900X 3.7 GHz 10-Core Processor, G.Skill Trident Z RGB 256 GB DDR4-3600

CL18 Memory, and a NVIDIA GeForce RTX 4090 24 GB Graphics Card. Data processing was

done in RStudio (R version 4.3.1 (2023-06-16) -- "Beagle Scouts")[62]. Graphs were made using

the package ggplot2 (version 3.4.2)[63]. Fisher's contingency table tests were done using

fisher.test. Wilcoxon rank sum tests were done using wilcox.test. Levene's tests of

homoscedasticity were done using leveneTest in package "car" (version 3.1-2). Data fitting with

ANOVA was done using the function aov. Figures were assembled in Adobe Illustrator (version

27.8.1). An RGB color profile "Image P3" was used for all the figures.

Statistical analysis

In most cases, each bud, either from the same inflorescence, a different inflorescence

from the same plant, or a different plant, is considered a biological replicate. For RNA-seq

presented in Fig. 1c, each RNA sample from 5-10 inflorescences of *ap1 cal AP1-GR* background

(either WT or *drmy1*), extracted separately, is considered a biological replicate.

For bar plots of sepal primordium number or auxin maxima number, all buds (or

simulations) were groups by genotype, and then grouped by number, and plotted as stacked bars.

For bar-and-whisker plots in Fig. 1c, 4b, 6h, Supplementary Fig. 4c, 4h, bar shows mean, and

whiskers show mean ± SD. For violin plots in Fig. 2e, 2k, Supplementary Fig. 3h, 3n, horizontal

lines show the quartiles. For circular histograms in Fig. 4d and 5g, lines show mean, and shaded

area shows mean ± SD.

Student's t-tests in Fig. 4b and Wilcoxon's rank sum tests in Fig. 2e, 2k, Supplementary

Fig. 3h, 3n, 4c, 4h are two-tailed, and compare only the indicated pairs of genotypes. For Levene's

test of variability in Supplementary Fig. 4c, 4h, center of each group were calculated as mean.

For ANOVA, formulae, degree of freedom, F, and p values are indicated in the Figure legends.

For Tukey's HSD in Fig. 6h, 6i, Supplementary Fig. 6d, 7d, all groups (combinations of genotype

and growth rate) were fit using a linear model, Estimated Marginal Means (EMM) were calculated,

and significant differences were indicated using compact letter display.

For RNA-seq, an FDR of 0.05 was used. For Tukey's HSD, family-wise type I error rate

was 0.05. For all other analyses, a p-value threshold of 0.05 was used.

Data availability

Data behind all the graph quantifications are available in Supplementary Dataset 1. RNA

seq data are available in GEO under project number PRJNA957462 and dataset number

GSE230100.

Code availability

Source code for the computational model, a tutorial, and demo data are available in

Supplementary Dataset 2 and also on GitHub (https://github.com/RoederLab/MassSpringAuxin).

REFERENCES

1. Araújo, I. S. *et al.* Stochastic gene expression in *Arabidopsis thaliana*. *Nat. Commun.* **8**, 2132 (2017).

2. Hong, L. *et al.* Variable Cell Growth Yields Reproducible Organ Development through Spatiotemporal Averaging. *Dev. Cell* **38**, 15–32 (2016).

3. Le Gloanec, C. *et al.* Cell type-specific dynamics underlie cellular growth variability in plants. *Dev.* **149**, dev200783 (2022).

4. Lachowiec, J., Queitsch, C. & Kliebenstein, D. J. Molecular mechanisms governing differential robustness of development and environmental responses in plants. *Ann. Bot.* **117**, 795–809 (2016).

5. Roeder, A. H. K. *et al.* Fifteen compelling open questions in plant cell biology. *Plant Cell* **34**, 72–102 (2022).

6. Hintze, M., Katsanos, D., Shahrezaei, V. & Barkoulas, M. Phenotypic Robustness of Epidermal Stem Cell Number in *C. elegans* Is Modulated by the Activity of the Conserved N-acetyltransferase nath-10/NAT10. *Front. Cell Dev. Biol.* **9**, 640856 (2021).

7. Garelli, A., Gontijo, A. M., Miguela, V., Caparros, E. & Dominguez, M. Imaginal discs secrete insulin-like peptide 8 to mediate plasticity of growth and maturation. *Science* **336**, 579–582 (2012).

8. Folta, A. *et al.* Over-expression of Arabidopsis AtCHR23 chromatin remodeling ATPase results in increased variability of growth and gene expression. *BMC Plant Biol.* **14**, 76 (2014).

9. Cassidy, J. J. *et al.* MiR-9a minimizes the phenotypic impact of genomic diversity by buffering a transcription factor. *Cell* **155**, 1556–1567 (2013).

10. Maugarny, A. *et al.* Compensation of compromised PRC2 regulation by a miRNA ensures robustness of Arabidopsis leaf development. *Preprint at bioRxiv:* https://doi.org/10.1101/2023.10.11.561475 (2023).

11. Tsai, T. Y. C. *et al.* An adhesion code ensures robust pattern formation during tissue
morphogenesis. *Science* **370**, 113–116 (2020).

12. Besnard, F. *et al.* Cytokinin signalling inhibitory fields provide robustness to phyllotaxis.
Nature **505**, 417–421 (2014).

13. Li, P. *et al.* Morphogen gradient reconstitution reveals Hedgehog pathway design
principles. *Science* **360**, 543–548 (2018).

14. Winter, K. U. *et al.* Evolution of class B floral homeotic proteins: Obligate heterodimerization
originated from homodimerization. *Mol. Biol. Evol.* **19**, 587–596 (2002).

15. Xiong, F. *et al.* Specified neural progenitors sort to form sharp domains after noisy Shh
signaling. *Cell* **153**, 550–561 (2013).

16. Zhu, M. *et al.* Robust organ size requires robust timing of initiation orchestrated by focused
auxin and cytokinin signalling. *Nat. Plants* **6**, 686–698 (2020).

17. Kong, S. *et al.* DRMY1 promotes robust morphogenesis by sustaining translation of a
hormone signaling protein. *Preprint at bioRxiv:* https://doi.org/10.1101/2023.10.11.561475
(2023).

18. Trinh, D.-C. Increased gene expression variability hinders the formation of regional
mechanical conflicts leading to reduced organ shape robustness. *Proc. Natl. Acad. Sci.*
120, 2017 (2023).

19. Lander, A. D., Gokoffski, K. K., Wan, F. Y. M., Nie, Q. & Calof, A. L. Cell lineages and the
logic of proliferative control. *PLoS Biol.* **7**, e1000015 (2009).

20. Lander, A. D. Pattern, growth, and control. *Cell* **144**, 955–969 (2011).

21. Li, R., Liu, Y., Pedersen, H. S., Kragh, P. M. & Callesen, H. Development and quality of
porcine parthenogenetically activated embryos after removal of zona pellucida.
Theriogenology **80**, 58–64 (2013).

22. Roeder, A. H. K. Arabidopsis sepals: A model system for the emergent process of
morphogenesis. *Quant. Plant Biol.* **2**, e14 (2021).

23.   Olsen, A. N., Ernst, H. A., Leggio, L. Lo & Skriver, K. NAC transcription factors: Structurally

distinct, functionally diverse. *Trends Plant Sci.* **10**, 79–87 (2005).

24.   Aida, M. & Tasaka, M. Genetic control of shoot organ boundaries. *Curr. Opin. Plant Biol.*

9, 72–77 (2006).

25.   Bilsborough, G. D. *et al.* Model for the regulation of *Arabidopsis thaliana* leaf margin

development. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 3424–3429 (2011).

26.   Furutani, M. *et al.* PIN-FORMED1 and PINOID regulate boundary formation and cotyledon

development in Arabidopsis embryogenesis. *Development* **131**, 5021–5030 (2004).

27.   Aida, M., Vernoux, T., Furutani, M., Traas, J. & Tasaka, M. Roles of PIN-FORMED1 and

MONOPTEROS in pattern formation of the apical region of the Arabidopsis embryo.

Development **129**, 3965–3974 (2002).

28.   Mallory, A. C., Dugas, D. V., Bartel, D. P. & Bartel, B. MicroRNA regulation of NAC-domain

targets is required for proper formation and separation of adjacent embryonic, vegetative,

and floral organs. *Curr. Biol.* **14**, 1035–1046 (2004).

29.   Sieber, P., Wellmer, F., Gheyselinck, J., Riechmann, J. L. & Meyerowitz, E. M. Redundancy

and specialization among plant microRNAs: Role of the MIR164 family in developmental

robustness. *Development* **134**, 1051–1060 (2007).

30.   Baker, C. C., Sieber, P., Wellmer, F. & Meyerowitz, E. M. The early extra petals1 mutant

uncovers a role for microRNA miR164c in regulating petal number in Arabidopsis. *Curr.*

Biol. **15**, 303–315 (2005).

31.   Aida, M., Ishida, T., Fukaki, H., Fujisawa, H. & Tasaka, M. Genes involved in organ

separation in Arabidopsis: An analysis of the *cup-shaped cotyledon* mutant. *Plant Cell* **9**,

841–857 (1997).

32.   Hibara, K. I. *et al.* Arabidopsis CUP-SHAPED COTYLEDON3 regulates postembryonic

shoot meristem and organ boundary formation. *Plant Cell* **18**, 2946–2957 (2006).

33.   Dinneny, J. R., Yadegari, R., Fischer, R. L., Yanofsky, M. F. & Weigel, D. The role of

JAGGED in shaping lateral organs. *Development* **131**, 1101–1110 (2004).

34. Long, J. & Barton, M. K. Initiation of axillary and floral meristems in Arabidopsis. *Dev. Biol.*

218, 341–353 (2000).

35. Peng, Z. *et al.* Differential growth dynamics control aerial organ geometry. *Curr. Biol.* **32**,

4854-4868.e5 (2022).

36. Ohno, C. K., Reddy, G. V., Heisler, M. G. B. & Meyerowitz, E. M. The Arabidopsis *JAGGED*

gene encodes a zinc finger protein that promotes leaf tissue development. *Development*

131, 1111–1122 (2004).

37. Růžička, K. *et al.* Cytokinin regulates root meristem activity via modulation of the polar

auxin transport. *Proc. Natl. Acad. Sci. U. S. A.* **106**, 4284–4289 (2009).

38. Marhavý, P. *et al.* Cytokinin controls polarity of PIN1-dependent Auxin transport during

lateral root organogenesis. *Curr. Biol.* **24**, 1031–1037 (2014).

39. Galbiati, F. *et al.* An integrative model of the control of ovule primordia formation. *Plant J.*

76, 446–455 (2013).

40. Reinhardt, D. *et al.* Regulation of phyllotaxis by polar auxin transport. *Nature* **426**, 255–260

(2003).

41. Smith, R. S. *et al.* A plausible model of phyllotaxis. *Proc. Natl. Acad. Sci. U. S. A.* **103**,

1301–1306 (2006).

42. Jönsson, H., Heisler, M. G., Shapiro, B. E., Meyerowitz, E. M. & Mjolsness, E. An auxin-

driven polarized transport model for phyllotaxis. *Proc. Natl. Acad. Sci. U. S. A.* **103**, 1633–

1638 (2006).

43. Gregor, T., Tank, D. W., Wieschaus, E. F. & Bialek, W. Probing the Limits to Positional

Information. *Cell* **130**, 153–164 (2007).

44. Erdmann, T., Howard, M. & Ten Wolde, P. R. Role of spatial averaging in the precision of

gene expression patterns. *Phys. Rev. Lett.* **103**, 2–5 (2009).

45. Waddington, C. H. Canalization of Development and the Inheritance of Acquired

Characters. *Nature* **150**, 563–656 (1942).

46.   Friml, J. *et al.* A PINOID-dependent binary switch in apical-basal PIN polar targeting directs

auxin efflux. *Science* **306**, 862–865 (2004).

47.   Zourelidou, M. *et al.* Auxin efflux by PIN-FORMED proteins is activated by two different

protein kinases, D6 PROTEIN KINASE and PINOID. *eLife* **3**, e02860 (2014).

48.   Michniewicz, M. *et al.* Antagonistic Regulation of PIN Phosphorylation by PP2A and

PINOID Directs Auxin Flux. *Cell* **130**, 1044–1056 (2007).

49.   Titapiwatanakun, B. *et al.* ABCB19/PGP19 stabilises PIN1 in membrane microdomains in

Arabidopsis. *Plant J.* **57**, 27–44 (2009).

50.   Nagawa, S. *et al.* ROP GTPase-dependent actin microfilaments promote PIN1 polarization

by localized inhibition of clathrin-dependent endocytosis. *PLoS Biol.* **10**, e1001299 (2012).

51.   Heisler, M. G. *et al.* Alignment between PIN1 polarity and microtubule orientation in the

shoot apical meristem reveals a tight coupling between morphogenesis and auxin

transport. *PLoS Biol.* **8**, e1000516 (2010).

52.   Heisler, M. G. *et al.* Patterns of auxin transport and gene expression during primordium

development revealed by live imaging of the Arabidopsis inflorescence meristem. *Curr.*

Biol. **15**, 1899–1911 (2005).

53.   Gonçalves, B. *et al.* A conserved role for *CUP-SHAPED COTYLEDON* genes during ovule

development. *Plant J.* **83**, 732–742 (2015).

54.   Marin, E. *et al.* mir390, Arabidopsis TAS3 tasiRNAs, and their AUXIN RESPONSE

FACTOR targets define an autoregulatory network quantitatively regulating lateral root

growth. *Plant Cell* **22**, 1104–1117 (2010).

55.   Smyth, D. R., Bowman, J. L. & Meyerowitz, E. M. Early flower development in Arabidopsis.

Plant Cell **2**, 755–767 (1990).

56.   Strauss, S. *et al.* Using positional information to provide context for biological image

analysis with MorphoGraphX 2.0. *eLife* **11**, e72601 (2022).

57.   Schindelin, J. *et al.* Fiji: An open-source platform for biological-image analysis. *Nat.*
647   *Methods* **9**, 676–682 (2012).

648   58.   Schindelin, J., Rueden, C. T., Hiner, M. C. & Eliceiri, K. W. The ImageJ ecosystem: An
649   open platform for biomedical image analysis. *Mol. Reprod. Dev.* **82**, 518–529 (2015).

650   59.   Majda, M. *et al.* Elongation of wood fibers combines features of diffuse and tip growth. *New*
651   *Phytol.* **232**, 673–691 (2021).

652   60.   Mosca, G. *et al.* Modeling Plant Tissue Growth and Cell Division. in *Mathematical Modelling*
653   *in Plant Biology* 107–138 (2018). doi:10.1007/978-3-319-99070-5_7.

654   61.   Federl & Prusinkiewicz. Virtual laboratory: an interactive software environment for
655   computer graphics. in *1999 Proceedings Computer Graphics International* 93–100 (1999).
656   doi:10.1109/CGI.1999.777921.

657   62.   R Core Team (2021). R: A language and environment for statistical computing. R
658   Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

659   63.   Wickham, H. *ggplot2: Elegant Graphics for Data Analysis*. (Springer, 2009).
660   doi:10.1007/978-0-387-98141-3.

ACKNOWLEDGEMENTS

We thank Isabella Burda, Trevor Cross, Michelle Heeney, Henrik Jönsson, Byron Rusnak, and

Avilash Singh Yadav for comments and suggestions on the manuscript. We thank Elliot

Meyerowitz and Arnavaz Garda for the *DR5::3xVENUS-N7*/*PIN1::GFP* (Ler) seeds and the

pCUC1::3xVENUS-N7 construct. We thank Mitsuhiro Aida for the *pCUC1::CUC1-GFP* seeds. We

thank Alexis Maizel for the *DR5::ER-mRFP1.2* seeds. We thank Arabidopsis Biological Resource

Center for providing seeds used in this research. We thank Addgene for providing the *5mCUC1*

pGreenII0129 construct (#12067). Research reported in this publication was supported by the

National Institute of General Medical Sciences of the National Institutes of Health (NIH) under

award number R01GM134037 to A.H.K.R., and a Biotechnological and Biological Sciences

Research Council (BBSRC) Institute Strategic Program Grant (BB/X01102X/1) to R.S.S. The

content is solely the responsibility of the authors and does not necessarily represent the official

views of the National Institutes of Health.

AUTHOR CONTRIBUTIONS

Conceptualization and design of experiments were done by S.K., M.Z., and A.H.K.R. Experiments

were carried out by S.K. Data analysis was done by S.K. and D.P. The MorphoDynamX modeling

platform was developed by B.L. and R.S.S. The floral meristem model was developed by S.K.

based on scripts written by B.L. and R.S.S. Manuscript was prepared by S.K. and edited by all

the authors. Funds were provided by A.H.K.R. and R.S.

COMPETING INTERESTS

The authors declare no competing interests.

MATERIALS AND CORRESPONDENCE

Material requests and correspondence should be addressed to A.H.K.R.

FIGURES

Fig. 1. CUC1 is upregulated in *drmy1* with disrupted spatial pattern.

(a-b) Floral phenotype of the *drmy1* mutant. **(a)** Stage 12 buds of WT (left) and *drmy1* (right). In

drmy1, asterisk indicates a gap on the side due to uneven sepal positions; arrowhead indicates a

gap on the top caused by unequal sepal length. Scale bars, 250 µm. **(b)** Two stage 6 buds of WT

(left), and two *drmy1* buds with matching outer sepal size. Arrowheads point to sepal primordia.

Note that sepal primordia in *drmy1* are of variable size, number, and position, leaving regions in

the bud periphery with no sepal outgrowth (bracket). Scale bars, 25 µm.

(c) Expression of *CUC* genes in floral tissue of WT vs *drmy1* (in Ler *ap1 cal AP1-GR* background),

as determined by RNA-seq. n = 3 samples per genotype. Adjusted p-values from DESeq2: CUC1,

$3.710{\times}10^{-13}$; CUC2, 0.6650; CUC3, 0.2292. Complete dataset is in Kong et al. (2023)[17].

(d-e) Expression pattern of *CUC1* in WT **(d)** and *drmy1* **(e)**, live imaged through time. For 0 to 36

hours, top row shows *pCUC1::3xVENUS-N7*, and bottom row shows *pCUC1::3xVENUS-N7*

(yellow) merged with Chlorophyll (magenta; to show flower morphology). For 54 hours, the

chlorophyll channel and its longitudinal section (dashed lines) are shown. Note that in WT, four

boundaries with high *CUC1* expression form at robust positions between the incipient sepal

primordia and the center of the floral meristem at stage 2 (arrowheads), corresponding to the

robust initiation of four sepals at stage 3. In *drmy1*, *CUC1* expression expands to the bud

periphery (brackets). Although this peripheral expression later narrows to boundary domains

(arrowheads), it correlates with limited sepal outgrowth at stage 3 (asterisk). *CUC1* expression in

the inter-sepal regions are also expanded in *drmy1* (asterisks). See Supplementary Movie 1.

Images are representative of n = 4 WT buds and n = 5 *drmy1* buds. Note that plants are

heterozygous for *pCUC1::3xVENUS-N7*. Scale bars, 25 μm.

(f-g) Protein accumulation pattern of CUC1 in WT **(f)** and *drmy1* **(g)**, live imaged through time.

For 0 to 36 hours, top row shows *pCUC1::CUC1-GFP*, and bottom row shows *pCUC1::CUC1-*

GFP (yellow) merged with Chlorophyll (magenta). For 60 hours, the chlorophyll channel and its

longitudinal section (dashed lines) are shown. Note that in WT, four boundaries with high CUC1

protein accumulation form at robust positions between the incipient primordia and the center of

the floral meristem at stage 2 (arrowheads), corresponding to the robust initiation of four sepals

at stage 3. In *drmy1*, CUC1 protein accumulation is initially much higher and less boundary-

restricted, and often in the bud periphery (brackets). In some regions, CUC1 accumulation

narrows to boundary domains (arrowheads), while in other regions, it remains in the bud

periphery, correlated with limited sepal outgrowth at stage 3 (asterisk). See Supplementary Movie

2. Images are representative of n = 2 WT buds and n = 4 *drmy1* buds. Scale bars, 25 μm.

Fig. 2. *CUC1* **upregulation is sufficient and necessary for variable sepal initiation**

(a-e) *CUC1* upregulation causes variable sepal initiation. **(a-b)** Two representative WT **(a)** and

5mCUC1 **(b)** buds stained with PI. Arrowheads show sepal primordia. "p" shows petal primordia.

Scale bars, 25 μm. **(c)** Quantification of sepal primordium number in each bud, color-coded by

number and grouped by genotype. **(d)** Illustration of quantifying variability in sepal position. Buds

in which sepal primordia are evenly distributed around the bud have low CV values of angular

distanced between adjacent sepal primordia (left). Buds in which sepal primordia are irregularly

positioned have high CV values of angles between adjacent sepals (right). Reproduced from Kong

et al. (2023)[17]. **(e)** Quantification of positional variability in WT vs *5mCUC1*. Sample size: WT, n

= 51 buds; *5mCUC1*, n = 70 buds. Asterisks indicate statistically significant differences from WT

in a Fisher's contingency table test (p = 4.535×10^{-14}) **(c)** or Wilcoxon rank sum test (p = 2.878×10^{-16}) **(e)**.

(f-k) The *cuc1* mutation rescues variability in sepal number and position in *drmy1*. **(f-i)**

Representative buds of WT **(f)**, *cuc1* **(g)**, *drmy1* **(h)**, and *drmy1 cuc1* **(i)**. The *cuc1* bud was PI-

stained, and others carried the *35S::mCitrine-RCI2A* membrane marker. Arrowheads point to

sepal primordia. Scale bars, 25 μm. **(j-k)** Quantification of sepal primordium number **(j)** and

positional variability **(k)** as described above. Sample size: WT, n = 66 buds; *cuc1*, n = 52 buds;

drmy1, n = 87 buds; *drmy1 cuc1*, n = 63 buds. Asterisks indicate statistically significant differences

in a Fisher's contingency table test **(j)** or Wilcoxon rank sum test **(k)**, and ns means no significant

differences. P-values for **(j)**: WT vs. *cuc1*, p = 1; *drmy1* vs. *drmy1 cuc1*, p = 2.575×10^{-4}. P-values

for **(k)**: WT vs. *cuc1*, p = 2.665×10^{-6}; *drmy1* vs. *drmy1 cuc1*, p = 4.922×10^{-3}.

Fig. 3. CUC1 promotes rapid sepal initiation.

(a-c) The outer sepal initiates more rapidly in *5mCUC1* than WT. **(a,b)** show one bud of WT **(a)**

and one bud of *5mCUC1* **(b)** live imaged every 6 hours for 18 hours. Chlorophyll channel is shown

on the top, and Gaussian curvature of extracted surfaces is on the bottom. The buds were

matched for size, indicating matched developmental stage. Asterisks indicate sepal initiation

events. Note that outer sepal initiation was much earlier in *5mCUC1* than WT. **(c)** Quantification

of bud size (area in Z-projections; as a proxy for developmental stage) at the time of sepal

initiation. Shown are mean (bars) and individual buds (dots). Sample size, WT outer sepal, n =

13; *5mCUC1* outer sepal, n = 20; WT inner sepal, n = 8; *5mCUC1* inner sepal, n = 15. An ANOVA

model of Size ~ Genotype * Sepal was fit. Genotype, degree of freedom (df) = 1, F = 26.823, p =

3.67×10^{-6}. Sepal, df = 1, F = 53.946, p = 1.40×10^{-9}. Interaction, df = 1, F = 8.387, p = 5.52×10^{-3}.

(d-f) Sepals initiate more slowly in *cuc1* than WT. **(d,e)** show one bud of WT **(d)** and one bud of

cuc1 **(e)** live imaged every 6 hours for 24 hours. Chlorophyll is shown on the top, and Gaussian

curvature of extracted surfaces is on the bottom. The buds were matched for size, indicating

matched developmental stage. Asterisks indicate sepal initiation events. Note that overall sepal

initiation is delayed in *cuc1* compared to WT. **(f)** Quantification of bud size (area in Z-projections;

as a proxy for developmental stage) at the time of sepal initiation. Shown are mean (bars) and

individual buds (dots). Sample size, 26 buds for WT and 23 buds for *cuc1*. An ANOVA model of

Size ~ Genotype * Sepal was fit. Genotype, df = 1, F = 13.198, p = 3.91×10^{-4}. Sepal, df = 2, F =

60.760, p < 2×10^{-16}. Interaction, df = 2, F = 0.685, p = 0.5055. Scale bar in all images, 25 μm.

Fig. 4. CUC1 overexpression is necessary and sufficient for variability in auxin maxima

patterning

(a-d) Auxin patterning is disrupted in *5mCUC1* and *drmy1* but rescued in *drmy1 cuc1*. **(a)**

Representative images of *DR5::3xVENUS-N7* (cyan, top row) and DR5 merged with the

chlorophyll channel (magenta, bottom row) in stage 2 buds of WT, *5mCUC1*, *cuc1*, *drmy1*, and

drmy1 cuc1. **(b)** Quantification of total DR5 signal (mean ± SD). Asterisks indicate statistically

significant differences in two-tailed t-tests (WT vs. *5mCUC1*, p = 0.0266; WT vs. *cuc1*, p =

$2.09{\times}10^{-6}$; *drmy1* vs. *drmy1 cuc1*, p = $1.32{\times}10^{-6}$). **(c)** Illustration of circular histogram analysis in

(d) and **(j-n)**. Each bud was aligned so that the incipient outer sepal was on the top and the

inflorescence meristem on the bottom. DR5 signal was quantified in 4°-bins around the Z-axis

starting between the incipient lateral and outer sepal, so that the incipient outer sepal would be at

45°. **(d)** Circular histograms of DR5 in each genotype showing mean ± SD. Sample size: WT, n =

30 buds; *5mCUC1*, n = 29 buds; *cuc1*, n = 19 buds; *drmy1*, n = 24 buds; *drmy1 cuc1*, n = 32 buds.

(e-n) *CUC1* upregulation promotes auxin maxima formation, but can amplify sporadic auxin

patches. **(e-i)** Live imaging of stage 2 buds of WT, *5mCUC1*, *cuc1*, *drmy1*, and *drmy1 cuc1*

carrying *DR5::3xVENUS-N7*. Shown is DR5 (cyan) merged with the chlorophyll channel

(magenta). On the left shows time relative to the beginning of stage 2 (second row), an indicator

of developmental progression. **(j-n)** Kymographs showing DR5 signal through time, in the same

buds as in **(e-i)**. O, incipient outer sepal; I, incipient inner sepal; L, incipient lateral sepal; CB,

cryptic bract. Brackets indicate diffuse bands of auxin signaling that later form distinct, variably

positioned auxin maxima (arrowheads). Arrows indicate additional auxin maxima that form in the

space between existing ones. Asterisks indicate a sporadic auxin patch that gradually disappears

in *drmy1 cuc1*. Scale bar in all images, 25 μm.

Fig. 5. CUC1 amplifies sporadic auxin patches to form variably positioned auxin maxima and primordia.

(a-f) WT **(a,b)**, *cuc1* **(c,d)**, and *5mCUC1* **(e,f)** were treated with mock (DMSO) **(a,c,e)** or 1 µM

BAP **(b,d,f)** for 32 hours and then transferred onto non-treatment media (0 h) and live-imaged

every 24 hours for four time points. For 0 h, 24 h, and 48 h, top rows show *DR5::3xVENUS-N7*

(cyan) and bottom rows show DR5 merged with the Chlorophyll channel (magenta). For 72 h,

buds were PI-stained. Note that BAP treatment disrupts robust auxin patterning and causes

sporadic auxin patches. In *cuc1*, sporadic patches disappear (asterisk), whereas in WT and

5mCUC1, they are amplified to form variably positioned auxin maxima and sepal primordia

(arrowheads). Scale bars, 25 µm.

(g) Circular histograms of DR5 in late stage 2 and early stage 3 buds at 48 h post-treatment.

Shown is mean ± SD. Note that BAP treatment causes regions of great variability in WT (large

SD, black arrows), which is less pronounced in *cuc1* and more pronounced in *5mCUC1*. Sample

size: WT + mock, n = 11 buds; WT + BAP, n = 17 buds; *cuc1* + mock, n = 11 buds; *cuc1* + BAP,

n = 19 buds; *5mCUC1* + mock, n = 13 buds; *5mCUC1* + BAP, n = 20 buds.

(h-k) Quantification of variability in primordium number **(h,i)** and position **(j,k)** in stage 3 and 4

buds at 72 h post-treatment. **(h)** shows the number of primordia in each bud, grouped by

genotype. An ANOVA model of Number ~ Genotype * Treatment was fit. Genotype, df = 2, F =

10.04, p = 1.06×10^{-4}. Treatment, df = 1, F = 39.81, p = 7.51×10^{-9}. Interaction, df = 2, F = 14.01, p

= 4.27×10^{-6}. **(j)** shows a violin plot of the CV of angles between adjacent primordia in each bud.

An ANOVA model of CV ~ Genotype * Treatment was fit. Genotype, df = 2, F = 27.790, p =

2.53×10^{-10}. Treatment, df = 1, F = 63.585, p = 2.58×10^{-12}. Interaction, df = 2, F = 6.932, p =

1.52×10^{-3}. Note that both sepal and petal primordia were counted, as they were hard to distinguish

in many cases. **(i,k)** shows the respective interactions plots of mean primordium number **(i)** and

mean CV of position **(k)**. Note that *5mCUC1* further increases the variability caused by BAP

treatment, while *cuc1* is largely resistant to the effect of BAP. Sample size: WT + mock, n = 13

buds; WT + BAP, n = 18 buds; *cuc1* + mock, n = 15 buds; *cuc1* + BAP, n = 21 buds; *5mCUC1* +

mock, n = 17; *5mCUC1* + BAP, n = 23.

Fig. 6. The noise-amplifying effect of CUC1 can be fully explained by an increase in PIN repolarization.

(a) Diagram of auxin-CUC1 interaction implemented in the model. Auxin represses *CUC1* expression, whereas CUC1 increases PIN repolarization. PIN1 polarizes toward neighboring cells with highest auxin concentration and moves auxin out of the cell into its neighbors. **(b)** Modeling approach. The floral meristem is modeled with a 2D disk of cells, which starts patternless. Molecular interactions in **(a)** generate auxin distribution patterns. Auxin maxima within the shaded middle ring (1/3 < r < 2/3 of total radius) were extracted and quantified. **(c)** How genotypes were modeled. The *drmy1* mutation is modeled as having higher noise in auxin production. The *cuc1* mutation is modeled by removing CUC1 (which decreases PIN1 repolarization). **(d-i)** Removing CUC1 rescues the variability in auxin patterning of *drmy1*. Shown are simulations of WT **(d)**, *cuc1* **(e)**, *drmy1* **(f)**, and *drmy1 cuc1* **(g)**. Note that while a sporadic auxin patch in *drmy1* was amplified to form an auxin maximum (arrowhead), a similar patch in *drmy1 cuc1* dissipates (arrow). Growth rate, 0.8. **(h)** shows simulated auxin concentration averaged across all auxin maxima within each bud (mean ± SD). **(i)** shows the number of auxin maxima in each bud at t=100. Quantifications

were done across n=500 simulations of each genotype. Letters show multiple comparison using

Tukey's HSD. **(j)** A tradeoff exists between the speed and robustness of morphogenesis. Having

CUC1 promotes rapid morphogenesis under low noise conditions (top left), yet disrupts

robustness under high noise conditions such as *drmy1* or BAP treatment (bottom left). Removing

CUC1 reduces auxin maxima intensity and sepal initiation speed (top right), but allows additional

time for noise to dissipate and robust pattern to form under high noise conditions (bottom right).