Somatic Mosaicism in Amyotrophic Lateral Sclerosis and Frontotemporal Dementia Reveals Widespread Degeneration from Focal Mutations

Zinan Zhou[1,2,3,12], Junho Kim[1,2,3,4,12], August Yue Huang[1,2,3,12], Matthew Nolan[5], Junseok Park[1,2,3], Ryan Doan[1,3], Taehwan Shin[1,2,3], Michael B. Miller[1,6], Brian Chhouk[1,2,3], Katherine Morillo[1,2,3], Rebecca C. Yeh[1,2,3], Connor Kenny[1,2,3], Jennifer E. Neil[1,2,3,11], Chao-Zong Lee[5], Takuya Ohkubo[7,8], John Ravits[8], Olaf Ansorge[9], Lyle W. Ostrow[10], Clotilde Lagier-Tourenne[5,13], Eunjung Alice Lee[1,2,3,13] and Christopher A. Walsh[1,2,3,11,13]

Affiliations:
1. Division of Genetics and Genomics, Boston Children's Hospital, Boston, MA, USA.
2. Manton Center for Orphan Disease, Boston Children's Hospital, Boston, MA, USA.
3. Department of Pediatrics, Harvard Medical School, Boston, MA, USA .
4. Department of Biological Sciences, Sungkyunkwan University, Suwon, South Korea.
5. Department of Neurology, The Sean M. Healey and AMG Center for ALS at Mass General, Massachusetts General Hospital, Harvard Medical School, Boston, MA, USA.
6. Department of Pathology, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA.
7. Department of Neurology, Yokohama City Minato Red Cross Hospital, Yokohama, Kanagawa, Japan.
8. Department of Neurosciences, School of Medicine, University of California at San Diego, La Jolla, CA, USA.
9. Nuffield Department of Clinical Neurosciences, University of Oxford, Oxford, Oxfordshire, UK.
10. Department of Neurology, Lewis Katz School of Medicine at Temple University, Philadelphia, USA.
11. Howard Hughes Medical Institute, Boston Children's Hospital, Boston, MA, USA.
12. These authors contributed equally: Zinan Zhou, Junho Kim, August Yue Huang.
13. These authors jointly supervised this work: Clotilde Lagier-Tourenne, Eunjung Alice Lee, Christopher A. Walsh. Email: clagier-tourenne@mgh.harvard.edu; ealee@childrens.harvard.edu; christopher.walsh@childrens.harvard.edu.

Abstract

      Although mutations in dozens of genes have been implicated in familial forms of amyotrophic lateral sclerosis (fALS) and frontotemporal degeneration (fFTD), most cases of these conditions are sporadic (sALS and sFTD), with no family history, and their etiology remains obscure. We tested the hypothesis that somatic mosaic mutations, present in some but not all cells, might contribute in these cases, by performing ultra-deep, targeted sequencing of 88 genes associated with neurodegenerative diseases in postmortem brain and spinal cord samples from 404 individuals with sALS or sFTD and 144 controls. Known pathogenic germline mutations were found in 20.6% of ALS, and 26.5% of FTD cases. Predicted pathogenic somatic mutations in ALS/FTD genes were observed in 2.7% of sALS and sFTD cases that did not carry known pathogenic or novel germline mutations. Somatic mutations showed low variant allele fraction (typically <2%) and were often restricted to the region of initial discovery, preventing detection through genetic screening in peripheral tissues. Damaging somatic mutations were preferentially enriched in primary motor cortex of sALS and prefrontal cortex of sFTD, mirroring regions most severely affected in each disease. Somatic mutation analysis of bulk RNA-seq data from brain and spinal cord from an additional 143 sALS cases and 23 controls confirmed an overall enrichment of somatic mutations in sALS. Two adult sALS cases were identified bearing pathogenic somatic mutations in *DYNC1H1* and *LMNA,* two genes associated with pediatric motor neuron degeneration. Our study suggests that somatic mutations in fALS/fFTD genes, and in genes associated with more severe diseases in the germline state, contribute to sALS and sFTD, and that mosaic mutations in a small fraction of cells in focal regions of the nervous system can ultimately result in widespread degeneration.

Introduction

Amyotrophic lateral sclerosis (ALS), a disease in which premature loss of upper and lower motor neurons (UMNs and LMNs) leads to fatal paralysis, shows clinical, genetic, and pathological overlap with frontotemporal dementia (FTD), a neurodegenerative disorder characterized by behavioral, language, and memory dysfunction[1]. 5-22% of individuals with ALS develop FTD, and ≈ 15% of those with FTD eventually develop ALS[2]. ALS and FTD also share common pathology, with cytoplasmic inclusions of TAR DNA binding protein (TDP-43) found in almost all ALS brains and in half of FTD brains[3,4]. FTD brains lacking TDP-43 inclusions mainly show tau pathology. ALS typically begins focally and spreads regionally as the disease progresses[5,6], although whether degeneration begins in UMNs, LMNs, or both simultaneously, has remained controversial[7,8], with some studies suggesting that focality can manifest independently in UMNs and LMNs[5,9]. TDP-43 pathology also follows stereotypical patterns in ALS and FTD brains[9-11], thought to reflect focal onset and intercellular transmission of TDP-43 inclusions in a prion-like manner, as shown in cell and animal models[12-18].

Whereas over 30 genes are implicated in ALS and FTD[19], most causative genes are linked to familial ALS (fALS) and FTD (fFTD), while 90-95% of cases are sporadic ALS (sALS) and FTD (sFTD) without a family history[20]. Prospective studies of ALS revealed a higher number of cases stemming from a genetic basis, regardless of whether a family history is documented[21], with the underestimation of genetic cases probably reflecting multiple factors, including incomplete ascertainment, death from other causes before diagnosis, and incomplete disease penetrance. Therefore, genetic screening of ALS/FTD genes is needed to fully examine the contribution of germline mutations in sporadic cases.

The focal onset of ALS and FTD, their stereotypical spread, and the increased risk in smokers[22], have raised interest in potential roles of somatic mosaic mutations in the pathogenesis of ALS and FTD[23]. Somatic mutations are increasingly recognized as prevalent in normal-appearing tissues, but somatic mutations responsible for neurological conditions are often limited to the central nervous system (CNS)[24], and hence undetectable through DNA sequencing of non-CNS tissues. Recent studies have evaluated the contributions of somatic mutation to Alzheimer's and Parkinson's disease directly using postmortem brain tissues[25].

In this study, we assessed potential contributions of germline and somatic mutations — distinguished by their variant allele frequencies (VAFs) — to sALS and sFTD using ultra-deep sequencing of a panel of neurodegeneration-associated genes on postmortem tissues of various brain regions and spinal cords from >400 unique sALS and sFTD cases. Our study revealed that pathogenic germline mutations are more common than previously appreciated in sALS and sFTD cases, supporting the underestimation of ALS and FTD cases with underlying genetic causes. In addition, we identified novel predicted pathogenic somatic mutations in 2.7% of the sALS and sFTD cases without known or novel pathogenic germline mutations. Protein-altering (missense/nonsense/frameshift) somatic mutations showed significant enrichment in sALS and sFTD cases and in disease-affected brain regions, supporting roles in disease pathogenesis. Regional analysis revealed focality of predicted pathogenic somatic mutations in primary motor cortex and spinal cord, supporting independent disease initiation in UMNs and LMNs, but also strongly supporting models of ALS and FTD in which the disease spreads beyond a relatively confined region containing a somatic mutation.

Results
Ultra-deep targeted sequencing of neurodegenerative genes in sALS and sFTD brains

To directly detect somatic mutations in sALS and sFTD brains, we obtained post-mortem frozen tissues of several brain regions and spinal cords from individuals diagnosed with sALS or sFTD, as well as from age-matched controls through the Massachusetts Alzheimer's Disease Research Center, Oxford Brain Bank, and Target ALS Foundation (Fig. 1a and Supplementary Table 1). Additional brain tissues from ALS, FTD and control cases, without a record of family history but with an age of death above 45 years old, were also obtained from the NIH NeuroBioBank. We designed a molecular inversion probe (MIP) panel targeting the exons and

exon-intron junctions of 88 neurodegeneration-related genes[26], which included 34 ALS/FTD
genes, 10 Alzheimer's disease genes, 28 Parkinson's disease genes, and 16 genes associated with
other rare neurodegenerative disorders (Supplementary Table 2). We performed MIP panel
sequencing at ~1,800X average sequencing depth (Fig. 1b and Extended Data Fig. 1), with a
similar distribution of sequencing depth across batches, disease conditions, and tissue regions
(Extended Data Fig. 1). The variance of depth, along with the batch and sample information,
were considered as factors in the mutation burden test . A total of 937, 364, and 516 samples
from 291 ALS, 117 FTD, and 144 neurotypical control individuals respectively were sequenced
(Fig. 1a, 1c and Supplementary Table 1). Of the ALS and FTD cases, 8 were diagnosed with
both ALS and FTD, and were therefore counted for each condition.
116
Pathogenic germline mutations in sALS and sFTD cases
We first identified pathogenic germline single-nucleotide variants (SNVs) and short
insertions and deletions (indels) using GATK followed by multiple variant filters (Fig. 1d). The
functional impact and predicted pathogenicity of identified germline mutations were annotated
using ANNOVAR[27] and multiple clinical databases. In addition, the most common inherited
cause of ALS and FTD, a hexanucleotide repeat expansion in the *C9ORF72* gene[28,29], was
genotyped by a repeat-primed PCR assay (Extended Data Fig. 2). Overall, 20.6% (60/291) of
ALS, 26.5% (31/117) of FTD and 0.7% (1/144) of control cases showed *C9ORF72* repeat
expansions or pathogenic germline mutations in ALS and FTD genes that have been previously
reported (Fig. 2a, Supplementary Table 3, 4). Known and novel missense mutations in ALS/FTD
genes represented the most prevalent mutation type (Fig. 2b). *C9ORF72* repeat expansion was
the most frequently mutated gene followed by known and novel pathogenic germline mutations
in *SOD1* for ALS, and *GRN* and *MAPT* mutations for FTD cases (Fig. 2c and 2d). The overall
fractions of *C9ORF72* repeat expansion carriers — 10.6% for ALS-only cases and 12.0% for
FTD-only cases — slightly exceeded those reported in previous studies, yet they remained
notably lower than the rates observed in fALS and fFTD cases[30-32]. Three carriers of the
C9ORF72 repeat expansion also had known pathogenic mutations in other genes associated with
ALS/FTD (Fig. 2d and Supplementary Table 3), aligning with previous studies that have
demonstrated instances of oligogenic inheritance involving *C9ORF72* repeat expansions and
other pathogenic mutations in certain sALS and sFTD cases[33,34].
Our pathogenicity prediction found pathogenic germline mutations in dominant
ALS/FTD genes besides *C9ORF72* repeat expansions in 14.1% of ALS, 19.7% of FTD, and
5.6% of control cases (Fig. 2a, Supplementary Table 3, 4). The odds ratios for the presence of
pathogenic mutations in ALS and FTD cases, compared to control cases, were 2.78 (95% CI:
1.24-7.07, p=9.3×10^{-3}) and 4.14 (95% CI: 1.70-11.17, p=8.2×10^{-4}) respectively, suggesting
pathogenic mutations are enriched in both ALS and FTD cases. Not surprisingly, all previously
reported pathogenic mutations were predicted to be pathogenic. Most novel pathogenic
mutations were nonsynonymous SNVs that would require experimental validation to confirm
their functional impact. However, two novel *GRN* frameshift mutations (p.L46Rfs*18 and
p.D250Tfs*6) identified in FTD cases are probably disease-causing (Supplementary Table 3),
since loss-of-function *GRN* mutations are known to cause FTD in a dominant manner[35,36].
When we considered previously unreported but likely pathogenic germline mutations,
another 12 disease cases exhibited potential instances of oligogenic inheritance (Fig. 2d). Of
these, five individuals carried *C9ORF72* repeat expansions alongside novel pathogenic germline
mutations in other ALS/FTD genes, while another five cases had known pathogenic germline

mutations in *GRN*, *SOD1*, and *MAPT* genes, in combination with novel predicted pathogenic
germline mutations in other ALS/FTD genes. Two patients carried multiple novel pathogenic
germline mutations. These findings provide additional evidence for oligogenic inheritance of
ALS and FTD[33,34,37,38] (Fig. 2d). We also found 13 FTD cases to have germline mutations in
genes previously linked to ALS only (*NEK1*, *SETX*, *ATP13A2*, *ALS2*, *ANXA11*, *DCTN1*, *FIG4*
and VAPB) and one ALS case to have a predicted pathogenic missense mutation in the FTD-
associated *MAPT* gene (Fig. 2d). These crossover mutations between ALS and FTD reinforce the
overlap between both diseases from shared underlying mechanisms.
160
Identification of somatic SNVs and indels from MIP sequencing data
We developed a custom pipeline integrating RePlow[39], Mutect2[40], and Pisces[41] for calling
somatic SNVs and indels in our MIP sequencing data (Fig. 1d). We selected somatic mutations
identified by at least two of the three callers (double-called mutations) followed by multi-step
variant filters to remove false positive candidates. Unlike heterozygous germline mutations with
variant allele fractions (VAFs) around 50%, heterozygous somatic mutations have VAFs less
than 50%, and we only called somatic mutations with VAFs below 40%. To benchmark our
pipeline, we performed a spike-in experiment by mixing two human samples from the Genome
in a Bottle Consortium (GIAB) at variant allele fractions (VAFs) of 10%, 5%, 2.5%, 1%, and
0.5% (Extended Data Fig. 3a). Double-called mutations identified by Mutect2 and Pisces were
excluded from the final call set due to high false positive and low validation rates (Extended
Data Fig. 3b, c). High sensitivity and precision were achieved for the remaining Replow-based
double-called mutations (Replow-Mutect2 and Replow-Pisces) while maintaining a low false
positive rate across the low VAFs compared to the somatic mutations called by each caller. The
MIP sequencing and our custom pipeline together allowed us to confidently identify somatic
mutations with a low false positive rate at VAF as low as 0.5%. The observed VAFs of somatic
mutations were well in line with the target VAFs at all five VAF levels (Extended Data Fig. 3).
The custom pipeline identified 167 somatic SNVs and indels from our MIP sequencing
data (Supplementary Table 5). The VAF distribution of identified somatic mutations was similar
between disease and control cases at high VAF levels (>5%), but low-VAF mutations were more
common in disease cases (Extended Data Fig. 4). Forty-one somatic candidates were selected for
validation and 87.8% of them were confirmed by deep amplicon sequencing (Supplementary
Table 6). The VAFs of validated candidates in amplicon sequencing showed a strong correlation
with their original VAFs in the MIP sequencing data (Fig. 3a).
185
Somatic mutations in disease-relevant genes are enriched in ALS and FTD cases lacking
pathogenic germline mutations
To examine the burden and potential roles of somatic mutations in ALS and FTD, we
focused on cases that lacked known or novel pathogenic germline mutations (referred to as
germline-free cases). Ninety-five unique somatic mutations in neurodegeneration-related genes
were identified in 696, 243, and 516 samples from 216 ALS germline-free cases, 76 FTD
germline-free cases, and 144 neurotypical controls, respectively. Most somatic mutations (80%,
76 out of 95 unique mutations) were focal, identified only in one tissue region of an individual
(Fig. 3b), and at very low VAFs (Extended Data Fig. 4), suggesting that they likely arose after
gastrulation[42], and are likely to have been confined to nervous tissue. Mutational signature
analysis using Mutalisk[43] demonstrated that clock-like signatures (SBS5 and SBS1) were the
predominant mutational signatures (Extended Data Fig. 5). Recent work has identified their

presence in brain development[44,45], and SBS1 reflects deamination of methylated cytosine during
cell division and mitosis.
Our MIP panel contained not only ALS/FTD genes but also genes involved in other
dementia. We first focused on somatic mutations in all the neurodegenerative genes. For the
somatically mutated genes, there was a clear separation between the disease and control groups
(Fig. 3c). Indeed, just one protein-altering somatic mutation was observed among all controls,
while 15 and 7 were observed in ALS and FTD cases, respectively. These protein-altering
somatic mutations were significantly enriched in ALS and FTD cases (Fig. 4a; p=0.013 and
p=0.011) when tested using a linear mixed-effect regression model, which considers multiple
potential confounding factors, suggesting that some or all of them were potentially disease-
causing.
The enrichments of somatic mutations in neurodegenerative genes showed striking
topographic patterns, with exonic somatic mutations showing enrichment exclusively in disease-
affected tissue regions for both FTD and ALS germline-free cases. The prefrontal cortex showed
enrichment for somatic FTD mutations, and the primary motor cortex for ALS (Fig. 4b), while
the premotor cortex—located immediately in between these two regions—showed no enrichment
for either condition, as was the case for other tested cerebral cortical regions as well. The spinal
cord in ALS had only a modest increase in protein-altering somatic mutations, although this
analysis is limited by a small number of control spinal cord samples and resultant wide
confidence intervals (Fig. 4b). For the prefrontal cortex of FTD and the primary motor cortex of
ALS, enrichments of protein-altering somatic mutations in germline-free cases were even more
significant than the overall enrichments of exonic somatic mutations (Fig. 4b; p=0.043 and
p=9.1×10^{-3}, p=6.8×10^{-3} and p=2.4×10^{-3} for exonic and protein-altering mutations in ALS and
FTD germline-free cases, respectively; linear mixed model), further supporting the pathogenic
roles of the identified somatic mutations.
We further assessed somatic mutations in genes specifically related to ALS and FTD and
found that somatic mutations in each were enriched in genes relevant to that corresponding
condition. Exonic and protein-altering mutations were specifically enriched in ALS-related genes
in germline-free ALS samples (Fig. 4c; p=0.029 and p=0.017 for exonic and protein-altering
mutations, linear mixed model). Moderate enrichments were observed for exonic and protein-
altering mutations in FTD-related genes in germline-free FTD samples. In fact, less than half of
FTD cases have pathological TDP-43 protein aggregates, while the other half have Tau protein
aggregates[4]. We thus checked the contribution of Tau proteinopathy-related genes, including
genes associated with Alzheimer's disease (AD), together with FTD-related genes and found
nominally significant enrichment of exonic and protein-altering somatic mutations only in
germline-free FTD cases (Fig. 4c; p=0.046 for both exonic and protein-altering mutations, linear
mixed model). Our FTD cases could not be categorized into those related to TDP-43 or Tau
proteinopathies due to the lack of relevant pathological information, hindering our ability to
examine the potential enrichment of somatic mutations within these distinct categories. On the
other hand, no protein-altering mutation was observed in any of the ALS/FTD genes in control
samples (Fig. 3c). The exclusive and diagnosis-specific enrichments of functional somatic
mutations suggest that most or all somatic mutations contribute to the pathogenesis of sALS and
sFTD.
241
242

Pathogenic somatic mutations have restricted regional distributions and are enriched in hypodiploid cells

Pathogenicity prediction of somatic mutations resulted in 8 predicted pathogenic somatic SNVs in previously known ALS and FTD/Tau-proteinopathy genes (Supplementary Table 7), which account for 3.2% and 2.6% of germline-free ALS and FTD cases, respectively (2.7% for all the germline-free sALS and sFTD cases). All mutations in ALS cases were observed in primary motor cortex or spinal cord, the most severely affected regions in ALS, emphasizing the remarkable topographic specificity of the mutations. In addition, a predicted pathogenic somatic SNV in *APP* (p.R328Q) was identified in primary motor cortex of a sporadic case that showed both ALS and FTD. All somatic mutations occurred in disease genes with dominant inheritance when found in the germline setting, except for one sALS case with a somatic *ALS2* (p.T787R) mutation identified in spinal cord. *ALS2* is an autosomal recessive disease gene[46,47], and the same individual carried an *ALS2* (p.Q24R) germline mutation in addition to the identified somatic mutation. Both *ALS2* mutations were predicted to be pathogenic, suggesting that they initiate disease in a "second hit" autosomal recessive manner at the cellular level in a small proportion of cells in the spinal cord and again further supporting the likely contribution of somatic variants to pathogenesis.

We selected four predicted pathogenic somatic SNVs in ALS/FTD genes-- *TIA1* (p.H54R), *MATR3* (p.K594I), *ALS2* (p.T787R), and *TARDBP* (p.L248F), and the predicted pathogenic *APP* somatic SNV (p.R328Q)--to study in greater detail in terms of regional and cell-type distributions. Amplicon sequencing across multiple brain and spinal cord regions showed that three of the five somatic SNVs [*MATR3* (p.K594I), *APP* (p.R328Q), *TARDBP* (p.L248F)] were restricted to the primary motor cortex (Fig. 5a and Supplementary Table 8). The other two somatic SNVs [*TIA1* (p.H54R) and *ALS2* (p.T787R)] had the highest VAFs in the spinal cord [2.16% for *TIA1* (p. H54R) and 0.97% for *ALS2* (p.T787R)], where they were originally identified, and were also present in other brain regions at very low VAFs [0.15-1.05% for *TIA1* (p.H54R), 0.16% - 0.59% for *ALS2* (p.T787R)] (Fig. 5a and Supplementary Table 8). All five somatic SNVs were absent in cerebellum. The ultra-low levels and limited distribution of these somatic SNVs suggest that they probably arose late in development and were thus likely excluded from non-CNS tissues. Together with the enrichment of exonic and protein-altering somatic mutations in disease-affected tissue regions, these findings also support the focal onset of ALS at the genetic level in these somatic cases. Cells carrying damaging somatic mutations could form initial lesions, likely TDP-43 inclusions, in UMNs and LMNs, but these must have ultimately spread to other regions of the motor system that lacked or carried exceedingly low levels of the mutation, but which nonetheless showed robust pathology post mortem otherwise indistinguishable from germline cases.

We then determined the presence of these five somatic SNVs in different cell types by performing amplicon sequencing of DNA from neuronal (NeuN+), glial (NeuN-), diploid, polyploid, and hypodiploid nuclei isolated by fluorescence-activated nuclei sorting (FANS) from the tissue regions in which they were originally identified (Extended Data Fig. 6). Interestingly, *TIA1* (p.H54R), *MATR3* (p.K594I), and *ALS2* (p.T787R) mutations were enriched in hypodiploid nuclei (Fig. 5b), which likely represent apoptotic cells with DNA fragmentation and cell death[48,49]. Enrichment of these three mutations in hypodiploid cells indicates a possible role in the pathogenic process, suggesting that they might be involved in inducing cell death. Surprisingly, these three mutations were identified in all cell fractions, but were more enriched in non-neuronal cells compared to neurons (Fig. 5b). This finding also implies that neurons may

exhibit a cell-type specific vulnerability to damaging somatic mutations in ALS/FTD genes. In
contrast, the depletion of the *APP* mutation from hypodiploid cells, and its relative enrichment in
non-neuronal cells compared to neurons (Fig. 5b), align with models proposing important actions
of AD risk genes in non-neuronal cells including microglia and astrocytes, potentially leading to
secondary neuronal loss[50]. However, further research is needed to confirm and better understand
these potential associations and mechanisms. The *TARDBP* (p.L248F) mutation was found in a
primary motor cortex sample with a very low VAF ($\approx 0.5\%$ upon validation). However, when
isolated cell fractions were tested, the mutation was not detected in any of them. This suggests
that the mutation was only present in the specific area where it was initially discovered and did
not extend to nearby regions. This conclusion was confirmed by amplicon sequencing of a
second tissue sample from the primary motor cortex, where it was also absent.
300
301   **RNA-MosaicHunter identifies additional pathogenic somatic mutations in bulk RNA-seq**
302   **data of sALS cases**
303   To complement our targeted sequencing of neurodegenerative genes, which identified
pathogenic somatic mutations in a small proportion of sALS and sFTD cases in known genes, we
performed a transcriptome-wide screen for somatic mutations using RNA-seq data to explore
whether genes not normally associated with these conditions might cause them in the mosaic
state. We profiled pathogenic somatic mutations in all expressed genes in bulk RNA-seq data
generated from 789 postmortem brain and spinal cord tissue samples of 143 sALS cases and 23
age-matched controls by the New York Genome Center ALS Consortium (Supplementary Table
9; 81 and 11 of the sALS and control cases respectively were included in our MIP sequencing)
using RNA-MosaicHunter, a tool capable of calling clonal somatic mutations from bulk RNA-
seq data with a Bayesian probabilistic model. Because of the limited coverage of bulk RNA-seq
data, RNA-MosaicHunter only has sensitivity to detect somatic mutations VAFs >$\approx$5%, and
discards somatic mutations at ultra-low levels. We found significant increases in total somatic
mutations in sALS cases not carrying pathogenic germline mutations (Extended data Fig. 7;
p=0.007). Additionally, there was a higher burden of somatic mutations predicted to be
damaging in germline-free sALS cases; although this trend did not reach statistical significance
(Extended data Fig. 7; p=0.058). Overall, these findings further confirmed that somatic mutations
may contribute to the development of sALS.
320   Interestingly, somatic SNVs in *DYNC1H1* and *LMNA* were identified in multiple CNS
regions of two sALS cases that did not harbor other pathogenic germline or somatic mutations
(Fig. 6a and Supplementary Table 10, both cases were included in the MIP sequencing).
Heterozygously acting, generally de novo, mutations in *DYNC1H1* and *LMNA* have been found
in patients with phenotypes resembling spinal muscular atrophy (SMA)[51-54], a motor neuron
disease genetically distinct but sharing some pathological overlap with ALS, including loss of
lower motor neurons, denervation of neuromuscular junction, and muscle atrophy[55]. Analysis of
whole-genome sequence data of the two cases for germline mutations in *SMN1*, the most
commonly mutated genes in SMA, did not identify pathogenic germline mutations. Both
individuals carrying these somatic mutations had leg-onset ALS with TDP43 pathology
predominantly in spinal cord and to a lesser extent in motor cortex (Fig. 6a-c). We further
investigated their regional mutation distribution using amplicon sequencing. The *LMNA*
(p.H566Y) somatic mutation was detected in all the tested brain and spinal cord regions with
VAFs ranging from 5.3 to 12.3% (Fig. 6d and Supplementary Table 8). The *DYNC1H1*
(p.R1962C) somatic mutation was also detected in all the tested CNS regions with VAFs ranging

from 0.1% to 5.2%, but the VAFs of the mutation were extremely low in the cerebellum (0.1%),
thoracic spinal cord (0.8%) and lumbar spinal cord (0.8%) (Fig. 6d and Supplementary Table 8).
Notably, the *DYNC1H1* (p.R1962C) mutation was undetectable in cultured fibroblasts from the
patient (Supplementary Table 8), indicating that the mutation arose late in development and was
likely limited to the CNS. The broad distribution of these two somatic mutations aligns with our
previous finding that somatic mutations with more than 5% VAFs are typically detected
throughout the CNS[56], with the low levels of the mutation in lumbar spine potentially reflecting
death of the motor neurons carrying this mutation. The *DYNC1H1* p.R1962C mutation is known
to be highly pathogenic, as it completely abolishes the motor function of the dynein complex *in*
vitro[57], and germline *DYNC1H1* p.R1962C mutations have been found in patients with
malformations of cortical development and delayed psychomotor development[58,59]. Although the
LMNA (p.H566Y) mutation was not previously reported, *LMNA* mutations cause autosomal
dominant laminopathies including Hutchinson-Gilford progeria and congenital muscular
dystrophy, which are characterized by congenital defects and increased early lethality[60,61]. Thus,
both genes can cause lethal diseases with pediatric age of onset, which may ordinarily preclude
the appearance of ALS, but the mosaic state could allow for a normal early life and the onset of a
degenerative disorder later in life. These data suggest that further genome-wide exploration of
brain tissue for somatic mutations could reveal additional ALS genes that cause early lethality in
the germline state.
354
Discussion
Our data provide several important insights into sALS and sFTD. First, we found that
about 30% of both conditions show known or novel, likely pathogenic germline mutations in
ALS or FTD genes, which advocates for a shift from family history-based to genetic testing-
based classification of ALS and FTD cases. Second, we find that a small but important fraction
(~2.7%) of germline-free sporadic cases harbor predicted pathogenic somatic variants in known
ALS or FTD genes, with the distribution of these mutations being disease and brain region-
specific, providing proof of concept of a potentially important contribution of somatic mutations
to pathogenesis. Finally, we find examples of genes associated with severe pediatric degenerative
diseases that can be present in ALS in the somatic state, potentially broadening the spectrum of
causative genes for these conditions.
While the case-control enrichment of somatic variants suggests a role in pathogenesis,
these somatic variants are present at surprisingly low VAFs and with patterns of topographic
restriction that match disease onset. It is very likely that these pathogenic somatic mutations
arose at a late stage of development and were not shared by other tissue regions. In the most
extreme case, the *TARDBP* (p. L248F) somatic SNV was even undetectable in tissue adjacent to
the original sampling site. The nature of these focal somatic events would prevent them from
being identified through routine genetic testing with blood or other peripheral samples. The
focality of these mutations in the nervous system also suggests a mechanism by which
degeneration may spread from a site containing mutant cells to eventually cause loss of neurons
in regions that do not carry the mutation. This process is thought to involve the TDP-43
proteinopathy as supported by recent studies in cell and animal models[12-18]. Identification of
predicted pathogenic somatic mutations in the primary motor cortex and in spinal cord from
individuals with ALS suggests potential onset of disease in either UMNs or LMNs but eventual
involvement of both. Our cell-type analysis revealed that several predicted pathogenic somatic
mutations were more enriched in glia than neurons. However, the reduced abundance in neurons

might also reflect the loss of neurons carrying these somatic mutations. This was reinforced by
our observations that three out of the four tested somatic mutations were more prevalent in
hypodiploid cells, which likely represent apoptotic cells. The potential harm inflicted on neurons
by these mutations once again bolsters the concept of a focal onset of ALS. Neurons carrying
these mutations constitute the initial lesion and subsequently undergo cell death. The demise of
these neurons could further reduce their presence, leading to a reduction in the VAFs of the
mutations compared to their levels at the time of initial emergence.
Although only about 2.7% of germline-free ALS and FTD cases had predicted
pathogenic somatic mutations in our MIP sequencing data, this is likely greatly underestimated
because of the limited sensitivity of even our deep panel sequencing approach to detect somatic
mutations at ultra-low levels (Extended data Fig. 4). The detection of somatic mutations with low
VAFs remains a technical challenge[45], but improved duplex sequencing approaches promise the
ability in the future to systematically sample somatic mutations at virtually all allele frequency
levels. Given that somatic mutations at very low levels and in focal regions appear capable of
creating a spreading disease, it will require very deep analysis to determine the lower allele
frequency range of variants that is capable of initiating this process. Variant detection is also
limited by availability of samples from regions across the CNS.
Our identification of candidate somatic SNVs in *DYNC1H1* (p. R1962C) and *LMNA* (p.
H566Y) using RNA-seq analysis of sALS cases suggests that genes that predispose their carriers
to ALS and FTD by somatic mutations may include genes distinct from those discovered in
germline cases. Certain alleles in both *DYNC1H1* and *LMNA* are associated with motor neuron
degeneration in the form of SMA, so they are capable of predisposing to neuronal degeneration,
but also in both cases, other alleles (including the *DYNC1H1* p. R1962C allele[58,59]) cause severe
pediatric disease that would normally mask the possibility of late-life ALS. This result suggests
that a wider range of ALS genes and alleles could exist in the somatic state that cannot be
observable in the germline state due to their association with early-onset severe disease. This
raises an exciting prospect that future genome-wide approaches, such as deep whole-genome or
exome sequencing of a cohort of ALS cases, could shed light not only on additional somatic
genetic mechanisms and their contributions to ALS, but also on the topographic patterns of
spread of pathology from focal sites.
411

412
Fig: 1. Experimental and analysis strategies. (a) Overall scheme of the experiments. Genomic
DNA isolated from 1,817 postmortem tissue samples of multiple brain regions and spinal cords
of 144 control, 291 ALS, and 117 FTD cases were used for molecular inversion probe (MIP)
capture sequencing with ultra-high depth. (b, c) Mean sequencing depth and number of tissue
samples in different brain regions and spinal cords of control, ALS, and FTD cases. Control,
n=516; ALS, n=937; FTD, n=364. CB: cerebellum; PMC: primary motor cortex; PFC: prefrontal
cortex; PreMC: premotor cortex; SC: spinal cord; OC: occipital cortex; AC: anterior cingulate
cortex. Error bars, 95% CI (d) Methodological pipelines to identify germline and somatic
variants. Germline variants were called by GATK HaplotypeCaller. *C9ORF72* genotype of ALS
and FTD cases were determined by repeat-primed PCR. Somatic variants were called by
RePlow, MuTect2, and Pisces. Additional somatic variants were called from 789 bulk RNA-seq
profiles of multiple brain regions and spinal cords of ALS cases generated by the New York
Genome Center ALS Consortium using RNA-MosaicHunter.
426

Fig. 2: *C9ORF72* **repeat expansion and pathogenic germline variants in ALS/FTD genes are prevalent in ALS and FTD.** (a) Proportions of ALS and FTD cases with *C9ORF72* repeat expansion, known, and novel pathogenic germline variants of ALS/FTD genes. Cases with multiple pathogenic mutations are indicated with '+' sign. (b) Distribution of *C9ORF72* repeat expansion and known and novel pathogenic germline variants in ALS/FTD genes classified by mutation types. (c) Ranking of the top 10 mutated ALS/FTD genes.(d) Visualization of ALS and FTD cases (vertical columns) with known and novel pathogenic germline variants (horizontal rows) in ALS/FTD genes. Color codes indicate the types of mutations. Rectangular outline represents novel variants. Genes are grouped by their known involvement in the diseases. * indicates cases with multiple pathogenic mutations.

Fig. 3: Somatic variants in MIP sequencing data tend to be focal, protein-altering and are almost exclusively restricted to disease cases. (a) The observed VAFs of somatic variants in amplicon sequencing validation were consistent with the VAFs in original MIP sequencing. Forty somatic variants were validated and included in the plot. (b) Total somatic variant counts classified by the number of brain regions in which a given variant was identified. (c) Distribution of somatic variants in all neurodegenerative genes. Color codes indicate variant types. Note that somatic variants identified in controls are unlikely to alter function, with just one missense mutation (red) and the remaining being synonymous or noncoding substitutions.

Fig. 4: Somatic variants are enriched in ALS and FTD cases and disease-related tissue
regions. (a) Enrichment of somatic variants in different genomic regions of germline-free ALS
and FTD cases compared to normal controls. (b) Enrichment of somatic variants in different
brain regions of germline-free ALS and FTD cases compared to normal controls. Significance of
enrichment and 95% CI was estimated while controlling for potential confounding factors
including average read-depth, sequencing batch, sampled individual using a linear mixed model.
(c) Enrichment of exonic and protein-altering somatic variants in two different groups of disease-
related genes (ALS genes and FTD/Tau-proteinopathy genes) compared to normal controls
457

Fig. 5: Pathogenic somatic mutations have restricted regional distributions and are
enriched in hypodiploid cells. (a) Regional distribution of VAFs of somatic variants in
individual brains and spinal cords. Brain cortex is annotated by Brodmann areas. The color
spectrum indicates the VAFs of somatic variants in amplicon sequencing. Dots indicate
unavailable regions and white indicates regions without the somatic variants. Red highlight
indicates the region of initial detection by MIP sequencing. (b) VAFs of somatic variants in
FANS sorted cell types. Five hundred neuronal (NeuN+), non-neuronal (NeuN-), diploid
(DAPI), hyperdiploid (High DAPI) and hypodiploid (Low DAPI) cells were each sorted for
amplicon sequencing with four replicates. Error bars, 95% CI.
468

Fig. 6: Somatic variants in *DYNC1H1* and *LMNA* in sALS. (a) Two pathogenic somatic SNVs that were shared by multiple tissue regions of the ALS cases. (b) Sections of the lumbar spinal cord, motor cortex, and hippocampus of the two sALS cases stained with a phospho-TDP43 antibody. Scale bar = 40 um. Arrowheads indicate the cells shown in the insets, which are magnified to twice their original size. (c) Quantification of phospho-TDP43 staining of CNS tissue sections of the two sALS cases with *DYNC1H1* and *LMNA* somatic mutations. Error bars indicate SD (n = 5). PFC: prefrontal cortex. MC: primary motor cortex. HC: hippocampus. TC: temporal cortex. PC: parietal cortex. OC: Occipital cortex. SC: spinal cord. (d) Regional distribution of VAFs of somatic variants in individual brains and spinal cords. Brain cortex is annotated by Brodmann areas. The color spectrum indicates the VAFs of somatic variants in amplicon sequencing. Dots indicate unavailable regions and white indicates regions without the somatic variants.

483
References

1.   Ferrari, R., Kapogiannis, D., Huey, E.D. & Momeni, P. FTD and ALS: a tale of two
diseases. *Curr Alzheimer Res* **8**, 273-94 (2011).
2.   Saxon, J.A. *et al.* Examining the language and behavioural profile in FTD and ALS-FTD.
J Neurol Neurosurg Psychiatry **88**, 675-680 (2017).
3.   Lagier-Tourenne, C., Polymenidou, M. & Cleveland, D.W. TDP-43 and FUS/TLS:
emerging roles in RNA processing and neurodegeneration. *Hum Mol Genet* **19**, R46-64
(2010).
4.   Ling, S.C., Polymenidou, M. & Cleveland, D.W. Converging mechanisms in ALS and
FTD: disrupted RNA and protein homeostasis. *Neuron* **79**, 416-38 (2013).
5.   Ravits, J.M. & La Spada, A.R. ALS motor phenotype heterogeneity, focality, and spread:
deconstructing motor neuron degeneration. *Neurology* **73**, 805-11 (2009).
6.   Kanouchi, T., Ohkubo, T. & Yokota, T. Can regional spreading of amyotrophic lateral
sclerosis motor symptoms be explained by prion-like propagation? *J Neurol Neurosurg*
Psychiatry **83**, 739-45 (2012).
7.   Eisen, A., Kim, S. & Pant, B. Amyotrophic lateral sclerosis (ALS): a phylogenetic
disease of the corticomotoneuron? *Muscle Nerve* **15**, 219-24 (1992).
8.   Chou, S.M. & Norris, F.H. Amyotrophic lateral sclerosis: lower motor neuron disease
spreading to upper motor neurons. *Muscle Nerve* **16**, 864-9 (1993).
9.   Gromicho, M. *et al.* Spreading in ALS: The relative impact of upper and lower motor
neuron involvement. *Ann Clin Transl Neurol* **7**, 1181-1192 (2020).
10.   Brettschneider, J. *et al.* Stages of pTDP-43 pathology in amyotrophic lateral sclerosis.
Ann Neurol **74**, 20-38 (2013).
11.   Brettschneider, J. *et al.* Sequential distribution of pTDP-43 pathology in behavioral
variant frontotemporal dementia (bvFTD). *Acta Neuropathol* **127**, 423-439 (2014).
12.   Polymenidou, M. & Cleveland, D.W. Biological Spectrum of Amyotrophic Lateral
Sclerosis Prions. *Cold Spring Harb Perspect Med* **7**(2017).
13.   Porta, S. *et al.* Patient-derived frontotemporal lobar degeneration brain extracts induce
formation and spreading of TDP-43 pathology in vivo. *Nat Commun* **9**, 4220 (2018).
14.   Laferriere, F. *et al.* TDP-43 extracted from frontotemporal lobar degeneration subject
brains displays distinct aggregate assemblies and neurotoxic effects reflecting disease
progression rates. *Nat Neurosci* **22**, 65-77 (2019).
15.   Peng, C., Trojanowski, J.Q. & Lee, V.M. Protein transmission in neurodegenerative
disease. *Nat Rev Neurol* **16**, 199-212 (2020).
16.   De Rossi, P. *et al.* FTLD-TDP assemblies seed neoaggregates with subtype-specific
features via a prion-like cascade. *EMBO Rep* **22**, e53877 (2021).
17.   Tamaki, Y. *et al.* Spinal cord extracts of amyotrophic lateral sclerosis spread TDP-43
pathology in cerebral organoids. *PLoS Genet* **19**, e1010606 (2023).
18.   Kumar, S.T. *et al.* Seeding the aggregation of TDP-43 requires post-fibrillization
proteolytic cleavage. *Nat Neurosci* **26**, 983-996 (2023).
19.   Rosen, D.R. *et al.* Mutations in Cu/Zn superoxide dismutase gene are associated with
familial amyotrophic lateral sclerosis. *Nature* **362**, 59-62 (1993).
20.   Turner, M.R. *et al.* Controversies and priorities in amyotrophic lateral sclerosis. *Lancet*
Neurol **12**, 310-22 (2013).

21. Andersen, P.M. & Al-Chalabi, A. Clinical genetics of amyotrophic lateral sclerosis: what
do we really know? *Nat Rev Neurol* **7**, 603-15 (2011).
22. Wang, H. *et al.* Smoking and risk of amyotrophic lateral sclerosis: a pooled analysis of 5
prospective cohorts. *Arch Neurol* **68**, 207-13 (2011).
23. Armon, C. Acquired nucleic acid changes may trigger sporadic amyotrophic lateral
sclerosis. *Muscle Nerve* **32**, 373-7 (2005).
24. Jamuar, S.S. *et al.* Somatic mutations in cerebral cortical malformations. *N Engl J Med*
371, 733-43 (2014).
25. Proukakis, C. Somatic mutations in neurodegeneration: An update. *Neurobiol Dis* **144**,
105021 (2020).
26. Hardenbol, P. *et al.* Multiplexed genotyping with sequence-tagged molecular inversion
probes. *Nat Biotechnol* **21**, 673-8 (2003).
27. Wang, K., Li, M. & Hakonarson, H. ANNOVAR: functional annotation of genetic
variants from high-throughput sequencing data. *Nucleic Acids Res* **38**, e164 (2010).
28. Renton, A.E. *et al.* A hexanucleotide repeat expansion in C9ORF72 is the cause of
chromosome 9p21-linked ALS-FTD. *Neuron* **72**, 257-68 (2011).
29. DeJesus-Hernandez, M. *et al.* Expanded GGGGCC hexanucleotide repeat in noncoding
region of C9ORF72 causes chromosome 9p-linked FTD and ALS. *Neuron* **72**, 245-56
(2011).
30. Majounie, E. *et al.* Frequency of the C9orf72 hexanucleotide repeat expansion in patients
with amyotrophic lateral sclerosis and frontotemporal dementia: a cross-sectional study.
Lancet Neurol **11**, 323-30 (2012).
31. Byrne, S. *et al.* Cognitive and clinical characteristics of patients with amyotrophic lateral
sclerosis carrying a C9orf72 repeat expansion: a population-based cohort study. *Lancet*
Neurol **11**, 232-40 (2012).
32. Mahoney, C.J. *et al.* Frontotemporal dementia with the C9ORF72 hexanucleotide repeat
expansion: clinical, neuroanatomical and neuropathological features. *Brain* **135**, 736-50
(2012).
33. van Blitterswijk, M. *et al.* Evidence for an oligogenic basis of amyotrophic lateral
sclerosis. *Hum Mol Genet* **21**, 3776-84 (2012).
34. Testi, S., Tamburin, S., Zanette, G. & Fabrizi, G.M. Co-occurrence of the C9ORF72
expansion and a novel GRN mutation in a family with alternative expression of
frontotemporal dementia and amyotrophic lateral sclerosis. *J Alzheimers Dis* **44**, 49-56
(2015).
35. Baker, M. *et al.* Mutations in progranulin cause tau-negative frontotemporal dementia
linked to chromosome 17. *Nature* **442**, 916-9 (2006).
36. Cruts, M. *et al.* Null mutations in progranulin cause ubiquitin-positive frontotemporal
dementia linked to chromosome 17q21. *Nature* **442**, 920-4 (2006).
37. Kuuluvainen, L. *et al.* Oligogenic basis of sporadic ALS: The example of SOD1
p.Ala90Val mutation. *Neurol Genet* **5**, e335 (2019).
38. Goutman, S.A. *et al.* Emerging insights into the complex genetics and pathophysiology of
amyotrophic lateral sclerosis. *Lancet Neurol* **21**, 465-479 (2022).
39. Kim, J. *et al.* The use of technical replication for detection of low-level somatic
mutations in next-generation sequencing. *Nat Commun* **10**, 1047 (2019).
40. Benjamin, D. *et al.* Calling Somatic SNVs and Indels with Mutect2. *bioRxiv*, 861054
(2019).

41.   Dunn, T. *et al.* Pisces: an accurate and versatile variant caller for somatic and germline
575   next-generation sequencing data. *Bioinformatics* **35**, 1579-1581 (2019).
576   42.   Bizzotto, S. *et al.* Landmarks of human embryonic development inscribed in somatic
577   mutations. *Science* **371**, 1249-1253 (2021).
578   43.   Lee, J. *et al.* Mutalisk: a web-based somatic MUTation AnaLyIS toolKit for genomic,
579   transcriptional and epigenomic signatures. *Nucleic Acids Res* **46**, W102-W108 (2018).
580   44.   Chung, C. *et al.* Comprehensive multi-omic profiling of somatic mutations in
581   malformations of cortical development. *Nat Genet* **55**, 209-220 (2023).
582   45.   Huang, A.Y. & Lee, E.A. Identification of Somatic Mutations From Bulk and Single-Cell
583   Sequencing Data. *Front Aging* **2**, 800380 (2021).
584   46.   Hadano, S. *et al.* A gene encoding a putative GTPase regulator is mutated in familial
585   amyotrophic lateral sclerosis 2. *Nat Genet* **29**, 166-73 (2001).
586   47.   Yang, Y. *et al.* The gene encoding alsin, a protein with three guanine-nucleotide
587   exchange factor domains, is mutated in a form of recessive amyotrophic lateral sclerosis.
588   *Nat Genet* **29**, 160-5 (2001).
589   48.   Ferlini, C., Biselli, R., Scambia, G. & Fattorossi, A. Probing chromatin structure in the
590   early phases of apoptosis. *Cell Prolif* **29**, 427-36 (1996).
591   49.   Young, N.A. *et al.* Use of flow cytometry for high-throughput cell population estimates
592   in brain tissue. *Front Neuroanat* **6**, 27 (2012).
593   50.   Hansen, D.V., Hanson, J.E. & Sheng, M. Microglia in Alzheimer's disease. *J Cell Biol*
594   **217**, 459-472 (2018).
595   51.   Rudnik-Schoneborn, S. *et al.* Mutations of the LMNA gene can mimic autosomal
596   dominant proximal spinal muscular atrophy. *Neurogenetics* **8**, 137-42 (2007).
597   52.   Harms, M.B. *et al.* Mutations in the tail domain of DYNC1H1 cause dominant spinal
598   muscular atrophy. *Neurology* **78**, 1714-20 (2012).
599   53.   Tsurusaki, Y. *et al.* A DYNC1H1 mutation causes a dominant spinal muscular atrophy
600   with lower extremity predominance. *Neurogenetics* **13**, 327-32 (2012).
601   54.   Iwahara, N., Hisahara, S., Hayashi, T., Kawamata, J. & Shimohama, S. A novel lamin
602   A/C gene mutation causing spinal muscular atrophy phenotype with cardiac involvement:
603   report of one case. *BMC Neurol* **15**, 13 (2015).
604   55.   Bowerman, M. *et al.* Pathogenic commonalities between spinal muscular atrophy and
605   amyotrophic lateral sclerosis: Converging roads to therapeutic development. *Eur J Med*
606   *Genet* **61**, 685-698 (2018).
607   56.   Lodato, M.A. *et al.* Somatic mutation in single human neurons tracks developmental and
608   transcriptional history. *Science* **350**, 94-98 (2015).
609   57.   Hoang, H.T., Schlager, M.A., Carter, A.P. & Bullock, S.L. DYNC1H1 mutations
610   associated with neurological diseases compromise processivity of dynein-dynactin-cargo
611   adaptor complexes. *Proc Natl Acad Sci U S A* **114**, E1597-E1606 (2017).
612   58.   Poirier, K. *et al.* Mutations in TUBG1, DYNC1H1, KIF5C and KIF2A cause
613   malformations of cortical development and microcephaly. *Nat Genet* **45**, 639-47 (2013).
614   59.   Yang, H. *et al.* De Novo Variants in the DYNC1H1 Gene Associated With Infantile
615   Spasms. *Front Neurol* **12**, 733178 (2021).
616   60.   Eriksson, M. *et al.* Recurrent de novo point mutations in lamin A cause Hutchinson-
617   Gilford progeria syndrome. *Nature* **423**, 293-8 (2003).
618   61.   Quijano-Roy, S. *et al.* De novo LMNA mutations cause a new form of congenital
619   muscular dystrophy. *Ann Neurol* **64**, 177-86 (2008).

62. Boyle, E.A., O'Roak, B.J., Martin, B.K., Kumar, A. & Shendure, J. MIPgen: optimized
modeling and design of molecular inversion probes for targeted resequencing.
Bioinformatics **30**, 2670-2 (2014).
63. Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads.
2011 **17**, 3 (2011).
64. Li, H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
arXiv preprint arXiv:1303.3997 (2013).
65. Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *Gigascience* **10**(2021).
66. Quinlan, A.R. & Hall, I.M. BEDTools: a flexible suite of utilities for comparing genomic
features. *Bioinformatics* **26**, 841-2 (2010).
67. Au, C.H., Ho, D.N., Kwong, A., Chan, T.L. & Ma, E.S.K. BAMClipper: removing
primers from alignments to minimize false-negative mutations in amplicon next-
generation sequencing. *Sci Rep* **7**, 1567 (2017).
68. Smith, T., Heger, A. & Sudbery, I. UMI-tools: modeling sequencing errors in Unique
Molecular Identifiers to improve quantification accuracy. *Genome Res* **27**, 491-499
(2017).
69. Van der Auwera, G.A. *et al.* From FastQ data to high confidence variant calls: the
Genome Analysis Toolkit best practices pipeline. *Curr Protoc Bioinformatics* **43**, 11 10
1-11 10 33 (2013).
70. Sherry, S.T. *et al.* dbSNP: the NCBI database of genetic variation. *Nucleic Acids Res* **29**,
308-11 (2001).
71. Genomes Project, C. *et al.* A global reference for human genetic variation. *Nature* **526**,
68-74 (2015).
72. Karczewski, K.J. *et al.* The ExAC browser: displaying reference data information from
over 60 000 exomes. *Nucleic Acids Res* **45**, D840-D845 (2017).
73. Karczewski, K.J. *et al.* The mutational constraint spectrum quantified from variation in
141,456 humans. *Nature* **581**, 434-443 (2020).
74. Fu, W. *et al.* Analysis of 6,515 exomes reveals the recent origin of most human protein-
coding variants. *Nature* **493**, 216-20 (2013).
75. Scott, E.M. *et al.* Characterization of Greater Middle Eastern genetic variation for
enhanced disease gene discovery. *Nat Genet* **48**, 1071-6 (2016).
76. Glusman, G., Caballero, J., Mauldin, D.E., Hood, L. & Roach, J.C. Kaviar: an accessible
system for testing SNV novelty. *Bioinformatics* **27**, 3216-7 (2011).
77. Jaganathan, K. *et al.* Predicting Splicing from Primary Sequence with Deep Learning.
Cell **176**, 535-548 e24 (2019).
78. Landrum, M.J. *et al.* ClinVar: improving access to variant interpretations and supporting
evidence. *Nucleic Acids Res* **46**, D1062-D1067 (2018).
79. Stenson, P.D. *et al.* Human Gene Mutation Database (HGMD): 2003 update. *Hum Mutat*
21, 577-81 (2003).
80. Liu, X., Wu, C., Li, C. & Boerwinkle, E. dbNSFP v3.0: A One-Stop Database of
Functional Predictions and Annotations for Human Nonsynonymous and Splice-Site
SNVs. *Hum Mutat* **37**, 235-41 (2016).
81. Kumar, P., Henikoff, S. & Ng, P.C. Predicting the effects of coding non-synonymous
variants on protein function using the SIFT algorithm. *Nat Protoc* **4**, 1073-81 (2009).
82. Adzhubei, I.A. *et al.* A method and server for predicting damaging missense mutations.
Nat Methods **7**, 248-9 (2010).

83. Chun, S. & Fay, J.C. Identification of deleterious mutations within three human genomes. *Genome Res* **19**, 1553-61 (2009).

84. Schwarz, J.M., Rodelsperger, C., Schuelke, M. & Seelow, D. MutationTaster evaluates disease-causing potential of sequence alterations. *Nat Methods* **7**, 575-6 (2010).

85. Reva, B., Antipin, Y. & Sander, C. Predicting the functional impact of protein mutations: application to cancer genomics. *Nucleic Acids Res* **39**, e118 (2011).

86. Shihab, H.A. *et al.* Predicting the functional, molecular, and phenotypic consequences of amino acid substitutions using hidden Markov models. *Hum Mutat* **34**, 57-65 (2013).

87. Shihab, H.A. *et al.* An integrative approach to predicting the functional effects of non-coding and coding sequence variation. *Bioinformatics* **31**, 1536-43 (2015).

88. Choi, Y., Sims, G.E., Murphy, S., Miller, J.R. & Chan, A.P. Predicting the functional effect of amino acid substitutions and indels. *PLoS One* **7**, e46688 (2012).

89. Dong, C. *et al.* Comparison and integration of deleteriousness prediction methods for nonsynonymous SNVs in whole exome sequencing studies. *Hum Mol Genet* **24**, 2125-37 (2015).

90. Zook, J.M. *et al.* Integrating human sequence data sets provides a resource of benchmark SNP and indel genotype calls. *Nat Biotechnol* **32**, 246-51 (2014).

91. Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21 (2013).

92. DePristo, M.A. *et al.* A framework for variation discovery and genotyping using next-generation DNA sequencing data. *Nat Genet* **43**, 491-8 (2011).

93. Huang, A.Y. *et al.* MosaicHunter: accurate detection of postzygotic single-nucleotide mosaicism through next-generation sequencing of unpaired, trio, and paired samples. *Nucleic Acids Res* **45**, e76 (2017).

94. Genomes Project, C. *et al.* An integrated map of genetic variation from 1,092 human genomes. *Nature* **491**, 56-65 (2012).

95. Tennessen, J.A. *et al.* Evolution and functional impact of rare coding variation from deep sequencing of human exomes. *Science* **337**, 64-9 (2012).

96. Lek, M. *et al.* Analysis of protein-coding genetic variation in 60,706 humans. *Nature* **536**, 285-91 (2016).

97. Evrony, G.D. *et al.* Single-neuron sequencing analysis of L1 retrotransposition and somatic mutation in the human brain. *Cell* **151**, 483-96 (2012).

98. Nolan, M. *et al.* Quantitative patterns of motor cortex proteinopathy across ALS genotypes. *Acta Neuropathol Commun* **8**, 98 (2020).

Methods
Tissue sources and sample preparation
Fresh frozen postmortem human brain and spinal cord tissues were collected by the
Massachusetts Alzheimer's Disease Research Center, Oxford Brain Bank, Target ALS
Foundation, and NIH NeuroBioBank (Supplementary Table 1) according to their respective
institutional protocols, written authorization and informed consent; they were subsequently
obtained for this study with the approval of the Boston Children's Hospital Institutional Review
Board. Research on these deidentified specimens and data was performed at Boston Children's
Hospital with approval from the Committee on Clinical Investigation. Sporadic ALS and FTD
cases were selected based on available clinical records. ALS and FTD cases without clear
recording of family histories were also selected if the age of death was above 45 years old.
gDNA of these tissue samples was extracted using the EZ1 Advanced XL (Qiagen) system
followed by an additional purification using AMPure XP beads (Beckman Coulter).
715
MIP panel design
A double-stranded DNA MIP panel targeting 1.4Mb across exons and exon-intron junctions of
88 neurodegenerative genes was designed using custom scripts incorporating MIPgen[62] using the
human reference genome, hg19, with Mly1 restriction sites masked with 'N' using bedtools. The
final panel of 26,439 MIPs captures an average fragment length of 209bp, including the
extension and ligation arms to ensure overlapping of the forward and reverse sequencing read.
The panel successfully targets 92.7% of bases including flanking intronic regions, with >98% of
exonic bases covered with an average of at least 2 unique MIPs. All MIPs were designed to
include a custom backbone consisting of primer binding sites and dual 5nt unique molecular
indexes (UMI). MIPs were rebalanced in the pool based on the percent of GC content within the
regions. Common primer binding and Mly1 restriction enzyme sites were added to both ends of
the oligo sequences to enable blunt-end removal of the primer binding sites. The forward and
reverse compliment sequences were printed into a single ssDNA pool by CustomArray (Bothell,
WA). The resulting panel was amplified at a low cycle number (12X), digested with Mly1
enzyme for 12 hrs at 37C, and purified using Qiagen Nucleotide removal kit.
731
MIP capture and library construction
Two hundred fifty ng of gDNA was first hybridized in a 15 ul reaction with 1.5 ul of
Ampligase® 10X Reaction Buffer (VWR), 1.5 ul of the reverse blocking oligo (5'-
NNNNGAAGTCGAAGGGCTATAGGCTGCCATCACANNNN-3') and the MIP pool at 63
nM for
10 min at 95 °C and 24 hrs at 60 °C. Gap-fill/ligation was then performed by adding 1 unit of
Phusion™ High-Fidelity DNA Polymerase (Thermo Fisher), 4 units of Ampligase® DNA
Ligase (Epicentre), 0.2 ul of Ampligase® 10X Reaction Buffer, 0.6 ul of dNTPs (10 mM) and 1
ul of nuclease-free water to the MIP capture product and incubated at 60 °C for 1 hr. For
exonuclease digestion, 50 units of Exonuclease III (Thermo Fisher), 10 units of Exonuclease I
(Thermo Fisher), 0.2 ul of Ampligase® 10X Reaction Buffer (VWR), and 2.05 ul of nuclease-
free water was added to the Gap-fill/ligation product, which was incubated for 40 min at 37 °C
and 5 min at 95 °C. Ten ul of the captured library is amplified in a 50 ul final reaction by adding
1 unit of Phusion Hot Start II DNA Polymerase (Thermo Fisher), 10 ul of 5X HF buffer, 1 ul of
dNTPs (10mM), 1 ul of the universal MIP barcode forward primer (10 uM), 1 ul of the
individual barcode reserve primer (10 uM) and 26.5 ul of nuclease-free water. MIP library

amplification was then performed under the following conditions: 98 °C for 30 s; 16 cycles of
98 °C for 10 s, 60 °C for 30 s and 72 °C for 30 s; 72 °C for 2 min. MIP library was then purified
using 2X AMPure XP Beads (Beckman Coulter,) and quantified by the Quant-iT™ dsDNA
Assay HS Kit (Thermo Fisher). Ninety-six MIP libraries were pooled together and sequenced on
one lane of Illumina Hiseq X.
753
Pre-processing and read mapping of MIP sequencing data
MIP sequencing primers were removed first from the raw FASTQ files using Cutadapt[63] (v2.4,
5' adapter of the first read: CATACGAGATCCGTAATCGGGAAGCTGAAG, 3' adapter of the
first read: ACACTACCGTCGGATCGTGCGTGT, 5' adapter of the second read:
GCTAAGGGCCTAACTGGCCGCTTCACTG, 3' adapter of the second read:
CTTCAGCTTCCCGATTACGGATCTCGTATG). Trimmed reads were aligned to the human
reference genome (GRCh37) using BWA-mem[64] (v0.7.15) and sorting and indexing were
performed using samtools[65] (v1.3.1). From the aligned BAM file, off-target reads were removed
by checking the overlaps with the target regions using bedtools[66] (v2.26.0). MIP arm regions
were masked by soft-clipping for each read using BAMClipper[67] (v1.0.1). Unique molecular
identifier (UMI) information was extracted, and then mapped reads were deduplicated based on
the mapping coordinate and the shared UMI using UMI-tools[68] (v1.0.0). Base quality score
recalibration and local realignment were performed using the Genome Analysis Toolkit (GATK,
v3.7)[69], generating final analysis-ready BAMs.
768
Variant calling for pathogenic germline mutations
Initial candidates of germline SNVs and indels were identified using GATK HaplotypeCaller
with default parameter settings. Low-quality candidates were filtered out if any of the following
conditions is not satisfied: 1) ≥ 10 variant-supporting reads, 2) ≥ 20 total read-depth at the
variant site, 3) VAF ≥ 0.3, 4) GATK QUAL ≥ 50, and 5) identified in all brain regions from the
same individual except for the samples failed to cover the variant site (<10 reads). Possible
pathogenic germline variants were further selected by satisfying all the following conditions: 1)
present in less than 0.1% of the population in any ethnic group of public databases including
dbSNP[70], the 1000 Genomes Project[71], the Exome Aggregation Consortium (ExAC)[72], the
Genome Aggregation Database (gnomAD)[73], the NHLBI Exome Sequencing Project
(ESP6500)[74], the Greater Middle East variome project (GME)[75], and Kaviar database[76], 2)
candidates observed only in disease or control groups but not in both, 3) possible protein-altering
candidates (missense, nonsense, frame-shift, or splicing variants), and 4) affecting 30 ALS- and
FTD-related genes. Pathogenicity prediction module (see computational prediction of variant
pathogenicity section below) was then applied to the remaining candidates, and predicted
pathogenic variants were reported as final pathogenic germline mutations. ANNOVAR[27] was
used to annotate the genomic region, affected genes, population allele frequency, and exonic
variant functions. SpliceAI[77] was additionally utilized to identify more splice-altering variants.
Candidates with delta score > 0.5 were considered to be possible splicing variants.
788
C9ORF72 **repeat expansion genotyping**
Repeat-primed PCR (RP-PCR) of the *C9ORF72* repeat expansion was performed in a 30 ul PCR
reaction with 150 ng of gDNA, 15 ul of 2X FastStart™ PCR Master (Roche), 2 ul of DMSO, 5
ul of 5X Q-solution (Qiagen), 1 ul of 5 mM 7-deaza-dGTP (NEB), 1 ul of 25 mM MgCl$_2$
(Qiagen) and 1 ul of the primer mix (40 uM of the Forward primer: 5'-/56-

FAM/AGTCGCTAGAGGCGAAAGC-3', 20 uM of the Reverse primer: 5'-
TACGCATCCCAGTTTGAGACGGGGGGCCGGGGCCGGGGCCGGGG-3' and 40 uM of the
Anchor/tail primer: 5'-TACGCATCCCAGTTTGAGACG-3'. The reaction was performed with
touchdown PCR cycling conditions consisting of 15 minutes at 95°C, followed by cycles of 94°C
for 1 minute, annealing starting at 70°C for 1 minute, and extension at 72°C for 3 minutes,
ending with a final extension step of 10 minutes at 72°C. The annealing temperature was
decreased in 2°C steps as follows: 70°C for two cycles, 68°C for three cycles, 66°C for four
cycles, 64°C for five cycles, 62°C for six cycles, 60°C for seven cycles, 58°C for eight cycles,
and 56°C for five cycles. The RP-PCR products were separated by the SeqStudio Genetic
Analyzer (Thermo Fisher) with the GeneScan™ 600 LIZ™ Dye Size Standard (Thermo Fisher).
Results of fragment sizes were analyzed by Peak Scanner™ Software v1.0 (Thermo Fisher).
805
Somatic variant calling from MIP sequencing data
Three different callers RePlow (v1.1.0)[39], Mutect2 (v4.1.5)[40], and Pisces (v5.2.11)[41] were used to
generate initial candidate sets. Each sample was analyzed by all three callers with the single-
sample mode. Default parameter settings were used except for the adjustments for disabling the
coverage limit. Variants that passed all the filters from each caller were used to make three
different initial sets. Candidates identified by only one caller were discarded, and those called at
least two callers were retained as a double-call set. For indels, double-calls between Mutect2 and
Pisces were used as somatic indel candidates since RePlow does not support indel detection. For
SNVs, among double-calls Mutect2-Pisces pairs were additionally filtered out due to high false
positive rates and low validation rates in the benchmarking data set (Supplementary Fig. 3).
Remaining RePlow-based SNV double-calls and indel candidates were subject to multi-step
variant filters to further remove false positive candidates.
Unlike germline variant calling, somatic variant calling aims to reliably detect low-VAF
mutations up to ~0.5%, which requires enough supporting evidence to control the false positive
rate. Calling thresholds such as variant-supporting read count, read-depth at the variant site, and
average base-call quality were determined based on the benchmarking data. Somatic variants
were selected satisfying all the following conditions: 1) ≥ 50 total read-depth at the variant site,
2) ≥ 15 variant-supporting reads excluding the reads with the variant allele on their probe-arm
regions, 3) > 30 average base-call quality of variant allele, 4) ≥ 2 different types of variant-
supporting amplicons, 5) $0.001 \leq VAF \leq 0.4$, 6) ≤ 3 variant candidates within 20 bp window
from the same sample, 7) present in less than 0.1% of the population in any ethnic group of
public databases and 8) observed in < 5 different individuals.
We additionally found that low-level contamination of DNA from another sample occurred in a
few samples. Germline variants from the contaminant mimicked low-VAF somatic mutations
and generated false positive calls. We therefore implemented a module to identify low-level
contamination and filter out candidates that originated from the contaminant. By comparing a
somatic candidate set from a given sample with the germline call set of every individual, sample
contamination was determined if the given sample has ≥ 40 low-VAF somatic candidates that are
also observed in a specific individual as germline variants. In this case, germline variants of the
matched individual are considered to be possible sources of false positive calls and all somatic
candidates that are matched with these germline variants from the individual were filtered out.
The remaining candidates were reported as final somatic variants.
Pathogenic somatic variants were further annotated with similar criteria for selecting pathogenic
germline variants. Among final candidates, variants that are 1) observed only in disease or

control groups but not in both, 2) possible protein-altering variants, and 3) affecting ALS- and
FTD-related genes were selected and applied for the pathogenicity prediction module.
ANNOVAR and SpliceAI were utilized to annotate variants with various genomic information
and detect additional splice-altering variants, respectively.
844
Computational prediction of variant pathogenicity
Pathogenicity prediction module was applied to filtered germline and somatic variants to refine
the pathogenic candidate sets. Variants that were previously reported as benign/likely benign in
the clinical databases (ClinVar[78] and Human Gene Mutation Database[79]) were excluded from the
pathogenic candidate set. Nonsense, frameshift, and canonical splicing variants (±1-2 splice
sites) were assumed to disrupt gene function and were included in the pathogenic set. For
missense variants, the dbNSFP database[80] was utilized to adopt multiple computational
algorithms (SIFT[81], PolyPhen2[82], LRT[83], MutationTaster[84], MutationAssessor[85], FATHMM[86],
FATHMM-MKL[87], PROVEAN[88], MetaSVM[89], MetaLR[89]), considering damaging effects at
different levels such as biochemical property, protein structure, and evolutionary conservation.
Categorical prediction results of each algorithm were delivered by ANNOVAR. A missense
variant was selected to be pathogenic if at least three different algorithms predicted damaging
effects (deleterious for SIFT, LRT, FATHMM, PROVEAN, MetaSVM and MetaLR; probably
damaging for PolyPhen2; disease_causing for MutationTaster), while excluding possibly/likely
damaging predictions from the counts for more conservative selection. For ALS/FTD-related
genes, previously reported inheritance patterns (dominant/recessive) were carefully checked. For
recessive genes, two independent mutations in the same gene were required to determine whether
a given individual was affected by pathogenic mutations.
863
Benchmarking with spike-in datasets
Two Coriell cell lines (GM12878 and GM24695) were used to generate a spike-in data.
Extracted DNA were mixed at five different levels to mimic low-level somatic mutations,
targeting the VAFs of 0.5%, 1%, 2.5%, 5%, and 10%. Genomic DNA from GM12878 cells was
spiked into DNA from GM24695, therefore unique germline SNPs in GM12878 were served as
somatic mutations. Genomic position and genotype information for germline SNPs of Coriell
samples were obtained from NIST high-confidence call sets[90]. A total of 165 SNPs (57
homozygous and 108 heterozygous SNPs) covered by our designed MIP panel were used as the
benchmark variant set. RePlow, Mutect2, Pisces, and their combinations were tested. Detected
mutations not in the benchmark set were considered to be false positives, except for GM24695
germline SNPs.
875
Somatic variant calling from RNA-seq data
Raw bam files of RNA-seq and matched WGS data for sALS and control cases of the New York
Genome Center ALS Consortium were obtained from the New York Genome Center. RNA-seq
reads extracted from raw bam files were aligned to the GRCh38 human reference genome by
STAR (v2.5.0a)[91] in the two-pass mode with the reference gene annotation (Gencode version
39). The aligned bam files were processed by Picard (v1.138) to remove duplicates, and then by
GATK (v3.6)[92] for SplitNCigarReads, indel realignment, and base quality recalibration. We
further excluded reads that were improperly paired or with ambiguous alignment.
Somatic SNVs were called by RNA-MosaicHunter (v1.0) with default parameters
(https://gitlab.aleelab.net/august/rna-mosaichunter; manuscript in submission). Derived from

MosaicHunter[93], which was designed for somatic mutation calling in DNA sequencing, RNA-
MosaicHunter incorporates a Bayesian genotyper and a series of empirical filters to
systematically distinguish somatic mutations from technical artifacts and germline mutations,
with 59% sensitivity and 94% precision benchmarked using cancer datasets. Specifically,
germline mutations identified from the matched WGS data from the same individual were
excluded. We excluded A-to-G candidates because they are most likely led by the widespread A-
to-I(G) RNA editing events in the human genome. To remove recurrent artifacts, we only
considered exonic candidates that were called in one or two individuals. We further excluded
candidates present in human polymorphism databases including dbSNP[70], the 1000 Genomes
Project[94], the Exome Sequencing Project[95], and the Exome Aggregation Consortium[96].
896
Nuclei isolation and whole genome amplification
Isolation of total (DAPI+), neuronal (NeuN+), non-neuronal (NeuN-), and damaged (low DAPI)
nuclei were achieved by FANS together with nuclear staining of NeuN (Millipore, MAB377)
and DAPI following a previously published study[97]. Five hundred nuclei of each cell population
were sorted into wells of 96-well plates.
Sorted nuclei were subjected to genome amplification using the Primary Template-directed
Amplification kit (BioSkryb, 100136) following the manufacturer's protocol.
904
Amplicon sequencing
Primer sets targeting each identified somatic SNV were designed using BatchPrimer3
(Supplementary Table 11). Amplicon was amplified for 25 cycles in a 50 ul PCR reaction with
50 ng of gDNA, 1 unit of Phusion Hot Start II DNA Polymerase (Thermo Fisher), 10 ul of 5X
HF buffer, 1 ul of dNTPs (10mM) and 10 ul of each primer (10 uM). Amplicon PCR products
were then purified by a 0.65X + 1.05X double size selection with AMPure XP Beads (Beckman
Coulter, A63882). Purified amplicons were then pooled based on the concentrations measured by
the Quant-iT™ dsDNA Assay HS Kit (Thermo Fisher) and sequenced using Amplicon-EZ
(Genewiz).
914
Burden analysis of somatic mutations using linear mixed model
Linear mixed-effect regression model was used to compare somatic mutation burden between
clinical conditions while accounting for other covariates that may affect the burden. Clinical
conditions and covariates of interest (e.g. age, gender, sequencing depth) were modeled as fixed
effects and the batch and individual (donor) information were modeled as random effects,
considering the uncertainty caused by sample clusters from the same origin (donor or batch).
Somatic mutation count in each sample was normalized per megabase pair and modeled as a
dependent variable. A covariate with a p-value < 0.05 was considered to be significant, based on
a t-test using the Satterthwaite approximation of degrees of freedom. To test the burden of
somatic mutations in different genomic regions, a linear mixed model was fitted to the mutation
counts of specific type (e.g. exonic). To test the burden of somatic mutations in different brain
regions, samples were first divided by the sequenced region and then a linear mixed model was
fitted for each region group.
928
929 **Immunohistochemistry**
930 Immunohistochemistry was performed using DAB (3,3'-Diaminobenzidine) detection as
931 previously described[98]. Briefly, 7μm formalin-fixed, paraffin-embedded (FFPE) sections were

dewaxed using citrisolve, before being rehydrated through decreasing concentrations of ethanol. Antigen retrieval was performed using sodium citrate buffer pH 6.0 at 121°C for 15 mins. Endogenous peroxidases were blocked using 3% hydrogen peroxide solution, and non-specific binding was blocked using 10% normal goat serum. Sections were then incubated overnight at 4°C with primary antibody (pTDP-43 mouse monoclonal, CosmoBio CAC-TIP-PTD-P03, 1:10,000). After washing with TBS-Triton, sections were incubated with a Horseradish peroxidase (HRP)-conjugated Goat anti-mouse secondary (Dako) for one hour at room temperature. HRP signal was detected using DAB substrate (Dako) applied for 15 minutes. Counterstaining was performed using Coles hematoxylin for 1 minute. Sections were then dehydrated, cleared using citrisolve, and mounted using glass coverslips. All sections were viewed using a Leica upright light microscope and assessed for section quality prior to whole-slide digital scanning.

Quantification of p-TDP43 burden by immunohistochemistry

Stained sections were scanned using a NanoZoomer whole-slide digital imager at 40X magnification. Images were then visualized and quantified using QuPath image analysis software and algorithms described previously[98]. Briefly, for cortical/cerebellar sections 5 ROI measuring 3mm2 (1000 x 3000µm) were placed equidistantly around a single gyrus with the short end of the ROI placed at the pial surface. Pathology was then quantified using a positive pixel count within each ROI and measurements were averaged to provide an output of positive pixels/mm2. For spinal cord sections, square ROI (2.25mm^2) was placed on each side of the central canal within the anterior horn and measurements were averaged.

Data availability

The bulk RNA-seq data for the NYGC ALS Consortium samples can be obtained upon request through the NYGC. The MIP targeted gene panel sequencing data generated in this study will be deposited to dbGaP with controlled use conditions set by human privacy regulations. Germline and somatic mutations identified and validated in this study are listed in the supplementary tables.

Code availability

The source code and default configuration file of RNA-MosaicHunter are available at https://gitlab.aleelab.net/august/rna-mosaichunter.git. The implemented codes for preprocessing of MIP sequencing data, statistical test, and visualization will be available before publication.

Acknowledgements

We thank the Massachusetts Alzheimer's Disease Research Center, Oxford Brain Bank, Target ALS Foundation (Biobank Core Facility at St. Joseph's Hospital and Barrow Neurological Institute, Georgetown Brain Bank, Eleanor and Lou Gehrig ALS Center at Columbia University and UCSD ALS bank) and NIH NeuroBioBank (Harvard Brain Tissue Resource Center, Mount Sinai/JJ Peters VA Medical Center NIH Brain and Tissue Repository, Brain Endowment Bank of University of Miami, University of Pittsburgh Neuropathology Brain Bank, University of Maryland Brian and Tissue Bank and UCLA Human Brain and Spinal Fluid Resource Center) for providing fresh frozen human tissues. We thank the Target ALS Human Postmortem Tissue Core, New York Genome Center for Genomics of Neurodegenerative Disease, Amyotrophic Lateral Sclerosis Association and TOW Foundation for providing the bulk RNA-seq data. We thank the donors and families for their contributions, and J. E. Neil and J. Gonzalez for

assistance with tissue procurement. We thank the Research Computing group at Harvard Medical
School and Boston Children's Hospital. The brains in Fig. 5 and Fig. 6 were illustrated by A. Lai
with input from the authors. This work was supported by the PRMRP Discovery Award
W81XWH2010028 (Z.Z.); the Edward R. and Anne G. Lefler Center postdoctoral fellowship
(Z.Z.); the American Heart Association Career Development Award 23CDA1046074 (Z.Z.); the
National Research Foundation of Korea (NRF) 2022R1C1C1010430 (J.K.); the Alzheimer's
Association research fellowship (A.Y.H.); R56 AG079857 (A.Y.H., C.A.W. and E.A.L.); A
Cullen Education and Research Foundation Young Investigator Award from the Healey Center
(M.N.); a Holloway Postdoctoral Fellowship from the Association for Frontotemporal
Degeneration (M.N.); K08 AG065502 (M.B.M.); donors of the Alzheimer's Disease Research
program of the BrightFocus Foundation A20201292F (M.B.M.); the Doris Duke Charitable
Foundation Clinical Scientist Development Award 2021183 (M.B.M.); K01 AG051791 (E.A.L.);
the Suh Kyungbae Foundation (E.A.L.), DP2 AG072437 (E.A.L.); R01 NS032457 (C.A.W.);
R01 AG070921 (C.A.W. and E.A.L.); a Massachusetts Alzheimer's Disease Research Center
pilot grant (C.L.-T. and C.A.W.); and the Allen Discovery Center program, a Paul G. Allen
Frontiers Group advised program of the Paul G. Allen Family Foundation (C.A.W. and E.A.L.).
C.L.-T. is supported by the Araminta Broch-Healey Endowed Chair in ALS. C.A.W. is an
Investigator of the Howard Hughes Medical Institute. The funders had no role in the study
design, data collection and analysis, decision to publish or preparation of the manuscript.
998

Author contributions

Z.Z., J.K. and A.Y.H. conceived and designed the study. Z.Z. performed tissue processing, MIP
panel sequencing, cell sorting and amplicon sequencing. J.K. performed bioinformatic analysis
for MIP sequencing data and validation with assistance from R.D. and T.S.. A.Y.H. performed
bioinformatic analysis for bulk RNA-seq data with assistance from J.P.. M.N. optimized and
performed immunofluorescent imaging and quantification, and generated data shown in this
manuscript. Z.Z., M.M. and R.D. designed the MIP panel. B.C., K.M., and R.Y. helped with
tissue processing and amplicon sequencing. C.K. provided technical support for MIP sequencing.
J.E.N. contributed tissue procurement and ethics expertise. T.O. and J.R. provided
immunofluorescent images and interpretation of disease pathology. L.W.O and O.A. provided
fresh frozen human tissues and interpretation of disease pathology. C.A.W., E.A.L. and C.L.-T.
supervised the study. Z.Z., J.K., A.Y.H., C.A.W., E.A.L. and C.L.-T. wrote the manuscript.