Developmental remodeling repurposes larval neurons for sexual behaviors in adult *Drosophila*

Julia A. Diamandi, Julia C. Duckhorn, Kara E. Miller, Mason Weinstock, Sofia Leone, Micaela R. Murphy, and Troy R. Shirangi[1],[*]

Villanova University, Department of Biology, 800 East Lancaster Ave, Villanova, PA 19085 U.S.A.

[1] Lead Contact

[*] Correspondence: troy.shirangi@villanova.edu

SUMMARY

Most larval neurons in *Drosophila* are repurposed during metamorphosis for functions in adult life, but their contribution to the neural circuits for sexually dimorphic behaviors is unknown. Here, we identify two interneurons in the nerve cord of adult *Drosophila* females that control ovipositor extrusion, a courtship rejection behavior performed by mated females. We show that these two neurons are present in the nerve cord of larvae as mature, sexually monomorphic interneurons. During pupal development, they acquire the expression of the sexual differentiation gene, *doublesex*, undergo *doublesex*-dependent programmed cell death in males, and are remodeled in females for functions in female mating behavior. Our results demonstrate that the neural circuits for courtship in *Drosophila* are built in part using neurons that are sexually reprogrammed from former sex-shared activities in larval life.

INTRODUCTION

Many behaviors of adult animals develop during a period of maturation, *e.g.*, puberty, as juveniles transform into adults[1]. The central nervous system changes dramatically during this time, including the birth of new neurons that build adult circuits[2]. However, in some animals like insects[3] and worms[4], the adult nervous system is also assembled with reprogrammed neurons that were formerly active in the juvenile.

During *Drosophila* metamorphosis, most neurons of the larval central nervous system are recycled for use in adult circuits[5]. The moonwalker descending neuron, for instance, triggers backward locomotion in crawling larvae and is remodeled during pupal life to regulate backward walking in adult flies[6]. Additionally, several input and output neurons of the larval mushroom body trans-differentiate during metamorphosis and contribute to entirely different circuits in the adult brain[3]. Despite these cases, there are currently no examples of recycled larval neurons that contribute to sexually dimorphic behaviors of adult flies. Indeed, lineage analyses of adult neurons with sexual identity in *Drosophila* suggest that most are born post-embryonically and contribute exclusively to sexually dimorphic behaviors in adults[7,8].

We previously identified a small sexually dimorphic population of interneurons in the abdominal ganglion of adult flies, the DDAG neurons, that co-express the *tailless*-like orphan nuclear receptor, *dissatisfaction* (*dsf*), and the sex differentiation gene, *doublesex* (*dsx*)[9]. Here, we identify two female-specific DDAG neurons that influence the extrusion of the ovipositor performed by mated females to reject courting males. These two DDAG neurons are born during embryogenesis and exist as segmental homologs of a sexually monomorphic interneuron in the larval abdominal ganglion called A26g. During early pupal life, the A26g neuron at abdominal segments five and six acquire the expression of *dsx*, undergo *dsx*-dependent programmed cell death in males, and are remodeled in females for use in the circuitry for ovipositor extrusion. Our results demonstrate that the neural circuits for sexually dimorphic behaviors in *Drosophila* include sexually reprogrammed neurons with former activities in the juvenile larva of both sexes.

RESULTS

The DDAG neurons are anatomically diverse

Drosophila (*D.*) *melanogaster* females and males have eleven and three *dsf*- and *dsx*-co-expressing abdominal ganglion (DDAG) neurons, respectively, which contribute to several female- and male-specific mating behaviors (Figure 1A)[9]. As a first step toward associating specific courtship functions to specific

DDAG neurons, we employed a stochastic labeling method to determine the anatomy of individual

54   DDAG neurons in females and males.

The DDAG neurons are labeled by a genetic intersectional strategy whereby a Flp recombinase,

56   driven in *dsx*-expressing cells by *dsx*LexA::p65 [10], excises a transcriptional stop cassette from an upstream

activating sequence (UAS)-regulated *myr::gfp* transgene. Expression of *myr::gfp* is activated in *dsf*-co-

58   expressing cells by the *dsf*Gal4 allele[9]. To stochastically label individual DDAG neurons using a similar

intersectional strategy, we utilized *FlpSwitch*[11], a Flp recombinase fused to the ligand binding domain of

60   the human Progesterone Receptor (hPR). The activity of the FlpSwitch recombinase is dependent upon

the presence of the progesterone mimic, mifepristone. We constructed flies carrying the *dsf*Gal4 and

62   *dsx*LexA::p65 alleles, a UAS-regulated *FlpSwitch*, and a LexAop-regulated *myr::gfp* transgene containing a

transcriptional stop cassette that is conditionally excised by the FlpSwitch recombinase. When these flies

64   were fed food containing mifepristone for a relatively short period of time, single DDAG neurons were

often labeled in adult flies. We examined approximately 60 and 20 single cell clones in female and male

66   ventral nerve cords, respectively, and identified five anatomically distinct DDAG subtypes (Figure 1B).

One of these subtypes, DDAG_A, is present in both sexes, whereas the DDAG_B–E subtypes are specific

68   to females. All DDAG neuronal subtypes arborize extensively in the abdominal ganglion, whereas the

DDAG_C–E neurons also extend an anteriorly projecting branch that innervates thoracic neuropils. We

70   conclude that the DDAG neuronal population consists minimally of five anatomical subtypes. Additional

DDAG subtypes may exist that were not labeled in our experiments, and the number of neurons of each

72   subtype is unclear.

74   **DDAG_C and DDAG_D neurons contribute to ovipositor extrusion**

The five DDAG subtypes we identified in females may contribute to different reproductive

76   behaviors. To develop driver lines that target subsets of DDAG neurons, CRISPR/Cas9-mediated

homology-directed repair was used to create new alleles of *dsf* that express *LexA::p65*, or the Split-Gal4

78   hemi-drivers, *p65AD::Zp* and *Zp::GAL4DBD*, in *dsf*-expressing cells. In a search for Split-Gal4 drivers that

target subsets of DDAG neurons, we found that intersecting *dsf*p65AD::Zp with the enhancer line, *VT026005-*

80   *Zp::GDBD*, labeled four female-specific DDAG neurons per hemisphere of the adult ventral nerve cord

(Figure 2A, B). Two cell bodies are located at the posterior tip of the nerve cord and the other two are at

82   the dorsal side of the abdominal ganglion. Labeling of individual neurons by the multicolor Flp-out

technique[12] identified the two neurons at the tip as a local interneuron corresponding to the DDAG_B

84   subtype, whereas the two dorsally located cell bodies are the DDAG_C and DDAG_D neurons (Figure

2C). The DDAG_C and DDAG_D neurons are segmental homologs (see below) with several anatomical similarities (Video S1). Both neurons extend a primary branch off the cell body forming a "ventral arch" that projects across the midline and then anteriorly on the contralateral dorsal side. Unlike the DDAG_D neuron, the DDAG_C neuron extends a dorsally projecting medial branch off the ventral arch that gives rise to arbors within the abdominal ganglion. The DDAG_D neuron is located anterior to the DDAG_C neuron. We were unable to differentiate the two DDAG_B neurons, however they may exhibit subtle anatomical differences that were undetected in our analysis and may contribute to female behavior differently.

We asked how the neurons labeled by *dsf*[p65AD::Zp] ∩ *VT026005-Zp::GDBD* influence female behavior. During courtship, unmated *D. melanogaster* females signal their willingness to mate by opening their vaginal plates and partially exposing the tube-like ovipositor, a behavior called "vaginal plate opening" or VPO[13,14]. Mated females, however, reject courting males by fully extruding their ovipositor, which may block copulation or male courtship drive[14,15]. The length of the abdomen increases when unmated females open their vaginal plates and mated females extrude their ovipositor, but the change in abdominal length is greater during an ovipositor extrusion than during an opening of the vaginal plates (Figure 2D). We previously showed that transient photoactivation of all DDAG neurons using the mVenus-tagged, red-light-gated cation channelrhodopsin, *CsChrimson::mVenus*[16], caused unmated females to open their vaginal plates and mated females to extrude their ovipositor[9] (Figure 2D). Upon photoactivation, *dsf*[p65AD::Zp] ∩ *VT026005-Zp::GDBD* > *CsChrimson::mVenus* females fully extruded their ovipositor regardless of their mating status (Figure 2D, E; Video S2, S3). Extrusion of the ovipositor was penetrant and occurred largely during the photoactivation period (Figure 2F), and quantitatively similar behaviors were observed across a range of stimulus intensities (Figure 2G).

We next tested whether activity of the DDAG_B–D neurons was required for ovipositor extrusion in mated females. In addition to labeling a subset of female-specific DDAG neurons, the intersection of *dsf*[p65AD::Zp] and *VT026005-Zp::GDBD* labels neurons in the brain (Figure 2A). To target the DDAG_B–D neurons specifically, we intersected *dsf*[p65AD::Zp] ∩ *VT026005-Zp::GDBD* with *dsx*[LexA::p65] (Figure 3A) and used the three-way intersection to suppress the activity of the DDAG_B–D neurons by driving the expression of a GFP-tagged version of the inwardly rectifying K+ channel, *Kir2.1*[17] (Figure 3B). Unmated females expressing *Kir2.1::gfp* in the DDAG_B–D neurons copulated with males at a rate that was similar to control unmated females (Figure 3C) and opened their vaginal plates during courtship at a frequency comparable to control females (Figure 3D). However, mated *dsf*[p65AD::Zp] ∩ *VT026005-Zp::GDBD* ∩ *dsx*[LexA::p65] > *Kir2.1::gfp* females extruded their ovipositor during courtship with a modest reduction in frequency

(Figure 3E) and laid fewer eggs (Figure 3F) compared to controls. Thus, the activity of one or more DDAG

118   subtypes labeled by the *dsf*[p65AD::Zp] ∩ *VT026005-Zp::GDBD* intersection contribute to ovipositor extrusion

and egg laying in mated females. The modest effects on ovipositor extrusion and egg laying frequency

120   from silencing these neurons may suggest the involvement of additional neural circuit elements.

       To determine the specific DDAG neuronal subtype that influences ovipositor extrusion, we

122   randomly expressed *CsChrimson::mVenus* in one or more neurons labeled by the *dsf*[p65AD::Zp] ∩ *VT026005-*

Zp::GDBD intersection using a Flp-based approach. By stochastically expressing a Flp recombinase under

124   the control of a heat-inducible promoter, we randomly excised a transcriptional stop cassette from a UAS-

regulated *CsChrimson::mVenus* transgene whose expression was driven by *dsf*[p65AD::Zp] ∩ *VT026005-*

126   *Zp::GDBD*. Using this strategy, we generated a population of mosaic females (n=103) that randomly

expressed *CsChrimson::mVenus* in one or more of the DDAG neurons labeled by *dsf*[p65AD::Zp] ∩ *VT026005-*

128   *Zp::GDBD*, producing 18 distinct groups of mosaic females (Figure 2H). Most of these females expressed

CsChrimson::mVenus in one or two DDAG neurons (Figure 2H inset). Females were tested in an

130   optogenetic activation experiment before their ventral nerve cords were dissected and stained with

antibodies to GFP to reveal the identity of DDAG neurons that expressed the *CsChrimson::mVenus* in each

132   female.

       All mosaic females expressing *CsChrimson::mVenus* specifically in a bilateral pair of DDAG_C

134   (n=5) or DDAG_D (n=4) neurons extruded their ovipositor during bouts of photoactivation (Figure 2H, I;

Video S4, S5). Unilateral activation of the DDAG_C (n=15) or DDAG_D (n=13) neurons caused a change

136   in abdominal posture with no extrusion of the ovipositor (Figure 2H, Video S6, S7). It is unclear if the

postural change is associated with any displacement of the vaginal plates. Photoactivation of a unilateral

138   DDAG_B neuron (n=16) failed to evoke any obvious behavior (Figure 2H, Video S8). We did not obtain

mosaic females that expressed *CsChrimson::mVenus* specifically in a bilateral pair of DDAG_B neurons

140   (Figure 2H). However, bilateral activation of the DDAG_B neurons did not modify the postural change

induced by photoactivation of a single DDAG_C or DDAG_D neuron (Figure 2H, Video S9). Taken

142   together, these results demonstrate that the DDAG_C and DDAG_D neurons contribute to ovipositor

extrusion in mated females. A functional difference between the DDAG_C and DDAG_D neurons was

144   not detected in our experiments. The contribution of the DDAG_B neurons to female behavior is

currently unclear.

The DDAG neurons originate as embryonic-born neurons in the larval ventral nerve cord

In each hemisphere of the late third-instar larval abdominal ganglion, the *dsf*[Gal4] allele labels two segmentally repeating interneurons from A1 to A8, and four interneurons at the terminal segments

(Figure 4A). The number and gross anatomy of these twenty neurons is similar in female and male larvae, in newly hatched larvae, and in larvae aged 24 and 48 hours after hatching (Figure 4A), indicating that the

neurons are born during embryogenesis, and that the expression of the *dsf*[Gal4] allele is stable through larval life. We posited that a subset of these twenty *dsf*-expressing neurons in each hemisphere of the

larval abdominal ganglion become the DDAG neurons of adult females and males. We tested this by visualizing *dsf*[Gal4] and *dsx*[LexA::p65] expression in the abdominal ganglion of pupae staged at several times

during pupal development. From 0 to 18 hours after puparium formation (APF), the twenty *dsf*-expressing neurons we observed in the abdominal ganglion of larvae were identifiable in the ventral

nerve cord of female and male pupae (Figure 4B, C). By 18 hours APF, approximately eleven of these neurons had gained *dsx*[LexA::p65] expression in both sexes (Figure 4B, C). In neuromeres A3–A7, one of the

two *dsf*-expressing neurons in each hemisegment was labeled by *dsx*[LexA::p65], and all six *dsf*-expressing neurons in A8 and the terminal segments co-expressed *dsx*[LexA::p65] (Figure 4B). The gain of *dsx*[LexA::p65]

expression in *dsf*-expressing neurons occurred gradually and monomorphically in both sexes from 0 to 18 hours APF, but by 36 hours APF, approximately eight *dsf*[Gal4] and *dsx*[LexA::p65] co-expressing neurons were

absent in males (Figure 4C). We previously demonstrated that the difference in DDAG neuron number between adult females and males was due to *dsx*-dependent apoptosis in males[9], indicating that the loss

of *dsf*[Gal4] and *dsx*[LexA::p65] co-expressing neurons in male pupae aged 36 hours APF is due to cell death. At 48 hours APF, approximately eleven and three *dsf*[Gal4] and *dsx*[LexA::p65] co-expressing neurons were present in

the abdominal ganglion of females and males, respectively, corresponding to the number of DDAG neurons in adults (Figure 4C).

To further test that the DDAG neurons are derived from *dsf*-expressing neurons in the abdominal ganglion of larvae, we repeated the Flp-based genetic intersection between *dsf*[Gal4] and *dsx*[LexA::p65], but this

time, we used *UAS-FlpSwitch* to conditionally activate the recombinase in *dsf*[Gal4]-expressing cells only during larval life. Larvae were fed mifepristone, causing the FlpSwitch to excise a transcriptional stop

cassette from a LexAop-controlled *myr::gfp* transgene in larval cells labeled by the *dsf*[Gal4]. The *dsx*[LexA::p65] allele was used to drive the expression of *myr::gfp* in *dsx*-expressing neurons of adults. GFP expression

was observed in the DDAG neurons of adults of both sexes (Figure S1), consistent with the notion that the DDAG neurons originate as *dsf*-expressing neurons in the larval abdominal ganglion. We conclude that

the entire population of DDAG neurons are embryonic-born *dsf*-expressing neurons that are present in larvae of both sexes. During pupal life, approximately eleven *dsf*-expressing neurons in the abdominal

ganglion of both sexes gain *dsx* expression (Figure 4B, C), eight of which are subsequently lost in males

due to *dsx*-mediated apoptosis[9].

Two segmental homologs of the A26g neuron in larvae become the DDAG_C and DDAG_D neurons

184   **of adult females**

We next sought to identify the larval counterparts of specific DDAG neuronal subtypes. We

186   found that the intersection of *dsf*[p65AD::Zp] and *VT026005-Zp::GDBD* used above to target the DDAG_B–D

subtypes labeled a single, bilateral, and segmentally repeating *dsf*-expressing interneuron in segments

188   A4–A6 of the larval abdominal ganglion of both sexes (Figure 5A, B). Stochastic labeling of individual

neurons labeled by the Split-Gal4 demonstrated that these neurons correspond to the A26g interneuron

190   (Figure 5C; J. Truman, personal communication).

The A26g neuron of larvae and the DDAG_C and DDAG_D neurons of adult females display

192   several anatomical similarities (Video S1). All three neurons have a cell body on the dorsal side of the

abdominal ganglion, a ventral arch, and a dorsally-located contralateral branch that projects anteriorly.

194   We therefore hypothesized that two segmental homologs of the A26g neuron labeled in larvae by

dsf[p65AD::Zp] ∩ *VT026005-Zp::GDBD* metamorphose into the DDAG_C and DDAG_D neurons of adult

196   females. To test this, we first visualized the A26g neurons labeled by *dsf*[p65AD::Zp] ∩ *VT026005-Zp::GDBD*

over the course of pupal development in females and probed DSX-F protein expression. At the onset of

198   pupariation, the A26g neurons at A4–A6 were labeled by *dsf*[p65AD::Zp] ∩ *VT026005-Zp::GDBD* but none of

them co-expressed DSX-F (Figure 5D). By 24 hours APF, however, the A26g neuron at A5 and A6, but not

200   A4, had gained DSX-F expression (Figure 5E). Two additional neurons, one at A7 and one at A8, were

labeled by the Split-Gal4 and both were also marked by DSX-F (Figure 5E). By 48 hours APF, the gross

202   morphology of the neurons labeled by *dsf*[p65AD::Zp] ∩ *VT026005-Zp::GDBD* had transformed to the likeness

of the DDAG_B–D neurons of adult females (Figure 5F, G). The cell bodies of the DSX-F-expressing A5

204   and A6 neurons were positioned on the dorsal side of the abdominal ganglion where the soma of the

DDAG_D and DDAG_C neurons are normally located, and the A7 and A8 cell bodies were at the tip of

206   the nerve cord, where the DDAG_B neurons are found. By adulthood, the DSX-F-non-expressing A26g

neuron at A4 had disappeared (Figure 5G). The DDAG_D neuron is anterior to the DDAG_C neuron,

208   suggesting that the A26g neuron at segments A5 and A6 transform into the DDAG_D and DDAG_C

neurons, respectively. The DDAG_B neurons are likely derived from *dsf*-expressing neurons at A7 and A8

210   that become marked by *dsf*[p65AD::Zp] ∩ *VT026005-Zp::GDBD* during pupal development prior to 24 hours

APF. The larval counterparts of the DDAG_B neurons are currently not known.

212   To confirm that the A26g neurons in larvae become the DDAG_C and DDAG_D neurons, we repeated the FlpSwitch-based genetic intersection we described above but between *dsx*[LexA::p65] and

214   *dsf*[p65AD::Zp] ∩ *VT026005-Zp::GDBD.* The Split-Gal4 was used to drive the expression of *UAS-FlpSwitch* in the A26g neurons and larvae were fed mifepristone to activate the FlpSwitch during larval life. The FlpSwitch

216   then excised a transcriptional stop cassette from a LexAop-controlled *myr::gfp* transgene driven by *dsx*[LexA::p65]. The DDAG_C and DDAG_D neurons of adult females were labeled by GFP (Figure 5H),

218   further confirming that the DDAG_C and DDAG_D neurons are indeed derived from two segmental homologs of the A26g neurons in larvae.

DSF and DSX-M regulate the survival of the DDAG_C and DDAG_D neurons

We previously showed that DSF activity is required for the survival of a subset of DDAG neurons in females, whereas DSX-M promotes the cell death of female-specific DDAG neurons in males[9]. We

sought to determine how DSF and DSX contribute to the development of the DDAG_C and DDAG_D neurons specifically. The *dsf*[p65AD::Zp] ∩ *VT026005-Zp::GDBD* intersection was used to visualize the

DDAG_C and DDAG_D neurons while driving the expression of a validated UAS-regulated short hairpin/miRNA (ShmiR) targeting *dsf* or *dsx* transcripts. *Dsx* transcripts are sex-specifically spliced and

translated to produce female- and male-specific isoforms of DSX proteins[18]. Knock-down of *dsx* transcripts in females did not cause any obvious change in DDAG_C or DDAG_D anatomy (Figure S2A,

B), whereas depletion of male-specific *dsx* transcripts caused a gain of DDAG_C and DDAG_D neurons with an arborization pattern like that of wild-type females (Figure S2A′, B′). Female-like DDAG neurons

are resurrected in males with the ectopic expression of the cell death inhibitor, P35[9], suggesting that the gain of the DDAG_C and DDAG_D neurons in *dsf*[p65AD::Zp] ∩ *VT026005-Zp::GDBD > UAS-dsx_ShmiR* males

is due to loss of cell death. Optogenetic activation of the resurrected DDAG_C and DDAG_D neurons in *dsf*[p65AD::Zp] ∩ *VT026005-Zp::GDBD > UAS-dsx_ShmiR* males induced an extrusion of the male's terminalia

(Video S10), a behavior reminiscent of ovipositor extrusion in mated females. Consistent with the absence of an anatomical phenotype in their DDAG_C and DDAG_D neurons, photoactivation of *dsf*[p65AD::Zp] ∩

VT026005-Zp::GDBD > UAS-dsx_ShmiR females caused an extrusion of the ovipositor (Figure S2G, Video S11).

In contrast, the DDAG_C and DDAG_D neurons were lost in *dsf*[p65AD::Zp] ∩ *VT026005-Zp::GDBD > UAS-dsf_ShmiR* females (Figure S2A, C) and in females carrying loss-of-function mutations in *dsf* (Figure

S2D, E). When *dsf*[p65AD::Zp] ∩ *VT026005-Zp::GDBD* was used to drive the expression of *UAS-dsf_ShmiR* and *UAS-CsChrimson::mVenus,* females failed to extrude their ovipositor during bouts of photoactivation

(Figure S2G, Video S12). To confirm that the loss of the DDAG_C and DDAG_D neurons in *dsf* mutant

females was due to cell death, we used *dsf*[Gal4] to drive the expression of *UAS-P35*. Indeed, blockage of cell

246   death by ectopic expression of P35 in *dsf*-expressing neurons of *dsf* mutant females rescued the DDAG_C

and DDAG_D neurons (Figure S2D–F). The rescued DDAG_C and DDAG_D neurons appear to lack

248   contralateral ascending projections, suggesting that *dsf* may contribute to neuronal development beyond

acting as a pro-survival factor. Knock-down of *dsf* transcripts in *dsf*[p65AD::Zp] ∩ *VT026005-Zp::GDBD > UAS-*

250   *dsf_ShmiR* males had no effect on the survival of the DDAG_B–D neurons (Figure S2A', C'). Taken

together, these data suggest that DSF promotes the survival of the DDAG_C and DDAG_D neurons in

252   females, whereas DSX-M promotes their cell death in males. Knock-down of *dsf* transcripts in *dsf*[p65AD::Zp] ∩

VT026005-Zp::GDBD > UAS-dsf_ShmiR larvae had no obvious effect upon the development of the A26g

254   neurons (Figure S2H), suggesting that DSF regulates the survival of the DDAG_C and DDAG_D neurons

during pupal development.

DISCUSSION

258   Many neurons of the *Drosophila* larval nervous system persist through metamorphosis and

contribute to adult neural circuits, yet their contribution to sexually dimorphic adult behaviors is unclear.

260   In this paper, we address this gap with two key findings: First, we identify two interneurons in the

abdominal ganglion of adult females, the DDAG_C and DDAG_D neurons, that contribute to ovipositor

262   extrusion, a behavior performed primarily by mated and mature females to reject courting males. The

neural circuitry that mediates ovipositor extrusion in mated females has been partially delineated

264   recently[15]. In response to the male's song, *dsx*-expressing pC2l neurons activate a descending neuron

called DNp13 that triggers the motor circuits in the abdominal ganglion for ovipositor extrusion[15].

266   Interestingly, the ability of DNp13 to induce ovipositor extrusion in mated but not unmated females

depends upon mechanosensory input during ovulation[15]. The DDAG_C/D neurons may function

268   downstream of DNp13 and ovulation-sensing mechanosensory neurons, perhaps integrating their inputs,

but upstream of motor circuits for ovipositor extrusion.

270   Photoactivation of all DDAG neurons induces vaginal plate opening in unmated females and

ovipositor extrusion in mated females[9]. Activating the DDAG_B–D neurons, however, induces ovipositor

272   extrusion regardless of mating status (Figure 2D). This suggests the possibility that the DDAG_A or

DDAG_E subtypes or both integrate mating status to inhibit DDAG_C/D-driven ovipositor extrusion in

274   unmated females. Testing this will require the development of new genetic tools that provide access to

the DDAG_A and DDAG_E subtypes.

276   Second, we find that the DDAG_C/D neurons are present in the larval abdominal ganglion as

mature sexually monomorphic neurons corresponding to two segmental homologs of the A26g neuron.

278   The function of A26g in larvae is currently unknown. During metamorphosis, the A26g neurons at

segments A5 and A6 acquire *dsx* expression in both sexes and are then remodeled in females to become

280   the DDAG_D and DDAG_C neurons, respectively. In males, expression of DSX-M promotes

programmed cell death of the A26g neurons during pupal life[9], whereas DSF activity is necessary for the

282   survival the A26g neurons in females. The mechanism by which DSF and DSX-M regulate A26g apoptosis

is unknown. One possibility is that DSX-M antagonizes DSF function in males, thereby allowing the

284   neurons to die during pupal life.

Our results demonstrate that the neural circuits for courtship behavior in *Drosophila* are

286   constructed in part from sex-non-specific larval neurons that are sexually reprogrammed during

metamorphosis for functions in adults (Figure 6). Similar observations have been made in *C. elegans*

288   where sex-specific patterns of synaptic connectivity in adults develop from neurons in juvenile worms

with sexually monomorphic connections[19]. How much of the circuitry for dimorphic courtship behaviors

290   in flies is built from reprogrammed larval neurons? Lineage analyses of neurons with sexual identity in

the adult brain and thoracic ganglia have shown that most, if not all, are born during post-embryonic

292   neurogenesis and function only in adults[7,8]. However, this may differ in the abdominal nervous system.

The abdominal ganglion of adult flies is largely specialized for functions in reproduction, and indeed the

294   majority of *dsx*-expressing neurons are located in the abdominal ganglion[20–22]. Some of these neurons are

specific to adults and are born during larval and pupal life from two terminal abdominal neural stem

296   cells, *i.e.*, neuroblasts, with sex-specific patterns of neurogenesis[23–25]. But many others, like the DDAG

neurons, are likely to be derived from embryonic lineages. In contrast to the brain and thoracic nervous

298   system, the vast majority of neuroblasts in the abdominal neuromeres (*e.g.*, A2–A8) finish producing

neurons by the end of embryonic life and add very few adult-specific neurons[23]. The contribution of

300   remodeled larval neurons to the circuits for sexually dimorphic behaviors may thus be relatively greater

in the abdominal ganglion than in other regions of the fly nervous system.

302   Insect neurons exhibit impressive plasticity as the central nervous system metamorphoses from

its larval to adult form. Some larval neurons undergo programmed cell death, but most persist through

304   pupal life to contribute to adult circuits[5]. Larval neurons that persist remodel their axonal and dendritic

arbors to regulate similar processes in adults[3,6,26–30] or trans-differentiate to obtain altogether different

306   functions[3,31]. Courtship and neuronal sexual identity are specific to adults, suggesting that the A26g-to-

DDAG_C/D transformation may be a case of trans-differentiation. How the A26g neurons acquire sexual

308   identity and become repurposed during metamorphosis may provide a system to study the regulatory

mechanisms underlying developmental reprogramming of the nervous system.

ACKNOWLEDGEMENTS

312   We thank J. Truman (University of Washington) for identifying the A26g neuron and suggesting driver

lines to target them in larvae; J. Cande and D. Stern (HHMI) for contributing *dsf* alleles; Y. Ding

314   (University of Pennsylvania), J. Lillvis (HHMI), and E. Preger-Ben Noon (Technion) for helpful

discussions and comments on the manuscript; and A. McStravog for administrative assistance. T.R.S. is

316   supported by the National Science Foundation (IOS-1845673).

318   **AUTHOR CONTRIBUTIONS**

Conceptualization, T.R.S.; Investigation, J.A.D, J.C.D, K.E.M, M.W., S.L., M.R.M., T.R.S.; Writing –

320   Original Draft, T.R.S., J.A.D.

322   **DECLARATION OF INTERESTS**

The authors declare no competing interests.

MAIN TEXT FIGURE LEGENDS

326   **Figure 1. The DDAG neurons are anatomically diverse.** (A) Confocal images of ventral nerve cords

(VNCs) from $dsf^{Gal4} \cap dsx^{LexA::p65} > myr::gfp$ males and females. GFP-expressing neurons are labeled in black

328   and DNCad (neuropil) is shown in light gray. (B) Confocal images of individual DDAG neurons from

dsf^{Gal4}/*LexAop-frt.stop-myr::gfp*; $dsx^{LexA::p65}$/*UAS-hPR::Flp* males and females fed food containing

330   mifepristone for about two hours during adulthood. A total of one and five DDAG subtypes were

identified in males and females, respectively. Coronal and sagittal views of each neuron is shown.

Figure 2. The DDAG_C and DDAG_D neurons contribute to ovipositor extrusion in unmated and

334   **mated females.** (A and B) Confocal images of brains and VNCs from $dsf^{p65AD::Zp} \cap VT026005\text{-}ZP::GDBD >$

CsChrimson::mVenus females and males probed with antibodies to GFP. *CsChrimson::mVenus*-expressing

336   neurons are labeled in black and DNCad (neuropil) is shown in light gray. Four VNC neurons are labeled

per hemisphere. (C) Confocal images of individual DDAG_B, DDAG_C, and DDAG_D neurons from

338     $dsf^{p65AD::Zp} \cap VT026005\text{-}ZP::GDBD >$ multicolor flp-out females. Arrows point to the ventral arch of the

DDAG_C and DDAG_D neurons. The DDAG_C neuron has a medial branch off the ventral arch that is

340   absent in the DDAG_D neuron. (D) Artificial activation of the DDAG_B–D neurons induces ovipositor

extrusion in unmated and mated females. The change in abdominal length when an unmated (black

342   squares) or mated (magenta squares) *Canton S* female performs vaginal plate opening or ovipositor

extrusion, respectively, during courtship is shown. Photoactivation of the entire DDAG population in

344   unmated (black triangles) and mated (magenta triangles) females induces a change in abdominal length

like that observed when *Canton S* females open their vaginal plates or extrude their ovipositor,

346   respectively[9]. Photoactivation of the DDAG_B–D subtypes in unmated (black circles) and mated

(magenta circles) females induces a change in abdominal length like that observed when *Canton S* mated

348   females extrude their ovipositor. (E) Still-frame images of decapitated *dsf*[p65AD::Zp] ∩ *VT026005-ZP::GDBD* >

CsChrimson::mVenus unmated (top) and mated (bottom) females before (left) and during (right)

350   illumination with 10.5 μW/mm^2 of red light to activate DDAG_B–D neurons. In both cases,

photoactivation results in ovipositor extrusion. (F) Average fraction of time unmated and mated females

352   extrude their ovipositor during (red) and before or after (black) three sequential 15-second bouts of

photoactivation (*i.e.*, oe index) of the DDAG subtypes labeled by *dsf*[p65AD::Zp] ∩ *VT026005-ZP::GDBD*. Lights

354   are off for 45 seconds before each bout. Indices are above zero when lights are off because it takes a few

seconds for females to fully retract the ovipositor once the photoactivation bout ends. (G) The change in

356   abdominal length upon photoactivation of *dsf*[p65AD::Zp] ∩ *VT026005-ZP::GDBD* > *CsChrimson::mVenus* in

unmated (black) and mated (magenta) females is similar across increasing light intensities. (H) Histogram

358   showing the number of unmated mosaic females in which *CsChrimson::mVenus* was either not expressed

(first column) or expressed randomly in DDAG_B–D neurons uni- or bilaterally (columns 2–19) via

360   expression of a heat-inducible Flp recombinase. Most mosaic females expressed *CsChrimson::mVenus* not

at all or in one or two DDAG neurons (inset histogram). Upon photoactivation, females performed either

362   no behavior (black bars), an abdominal postural change (orange bars), or an ovipositor extrusion (blue

bars). All females within each expression category along the x-axis performed similar behaviors upon

364   photoactivation. (I) Artificial bilateral activation of the DDAG_C or DDAG_D neurons induces ovipositor

extrusion. No other neurons were activated in these females. (right) Still-frame images of mosaic

366   *dsf*[p65AD::Zp] ∩ *VT026005-ZP::GDBD* > *CsChrimson::mVenus* unmated females before (left) and during (right)

illumination with 10.5 μW/mm^2 of red light to activate the DDAG_C or DDAG_D neurons bilaterally and

368   specifically. (left) *CsChrimson::mVenus* expression in the posterior nerve cord of the females shown in the

still frames. (D, F, G) show individual points, mean, and SD. A one-way ANOVA Turkey-Kramer test for

370   multiple comparisons was used to measure significance ($P<0.05$). Same letter indicates no significant

difference ($P>0.05$).

Figure 3. Activity of neurons labeled by *dsf*[p65AD::Zp] ∩ *VT026005-ZP::GDBD* is necessary for ovipositor extrusion and egg laying in mated females. (A) Confocal images of the brain and VNC from *dsf*[p65AD::Zp] ∩ *VT026005-ZP::GDBD* ∩ *dsx*[LexA::p65] > *myr::gfp* females. GFP-expressing neurons are labeled in black and DNCad (neuropil) is shown in light gray. (B) Confocal image of a VNC from *dsf*[p65AD::Zp] ∩ *VT026005-ZP::GDBD* ∩ *dsx*[LexA::p65] > *kir2.1::gfp* females. Kir2.1::GFP-expressing neurons are labeled in black and DNCad (neuropil) is shown in light gray. (C, D) Unmated females with inhibited DDAG_B–D neurons using *kir 2.1::gfp* are similarly receptive to male courtship relative to two control genotypes. (C) Fraction of unmated females that mated with a naïve *Canton S* male over a 30-min period. Significance (*P*<0.05) was measured using a Logrank test. n.s. = not significant. (D) Number of times an unmated female opened her vaginal plates per minute of active courtship from a naïve *Canton S* male. (E, F) Ovipositor extrusion and egg laying are reduced in mated females with inhibited DDAG_B–D. (E) Number of times a mated female extruded her ovipositor per minute of active courtship from a naïve *Canton S* male. (F) Number of eggs laid by a mated female 24-hr after mating. (D–F) show individual points, mean, and SD. A one-way ANOVA Turkey-Kramer test for multiple comparisons was used to measure significance (*P*<0.05). Same letter indicates no significant difference (*P*>0.05).

Figure 4. The DDAG neurons originate as embryonic-born *dsf*-expressing interneurons in the abdominal ganglion of larvae. (A) Confocal images of central nervous systems (CNSs) from *dsf*[Gal4] > *myr::gfp* larvae at various hours post-hatching (hph). GFP-expressing neurons are labeled in black and DNCad (neuropil) is shown in light gray. *dsf*[Gal4] labels 20 interneurons per hemisphere in the abdominal ganglion of female and male larvae. These neurons are observed in newly hatched larvae and are thus born during embryogenesis. The sex of the larvae at 0–2 and 24–26 hph was not determined. (B) A subset of *dsf*-expressing neurons in the abdominal ganglion of larvae acquire *dsx* expression during pupal life. (left) Confocal images of the VNC from *dsf*[Gal4] > *myr::rfp*, *dsx*[LexA::p65] > *myr::gfp* female and male pupae at 18 hours after puparium formation (APF). RFP-expressing neurons are labeled in magenta, GFP-expressing neurons are in green, and DNCad (neuropil) is shown in blue. (right) *dsf*[Gal4]-expressing neurons at each abdominal neuromere are shown. Neurons that co-express *dsx*[LexA::p65] are labeled with an asterisk. From A3–A7, one of two *dsf*[Gal4]-expressing neurons are also positive for *dsx*[LexA::p65], whereas all *dsf*[Gal4]-expressing neurons posterior to A7 co-express *dsx*[LexA::p65]. An illustration summarizing the expression of *dsf*[Gal4] and *dsx*[LexA::p65] in the abdominal ganglion at 18 hours APF is shown to the right. Cells that co-express *dsf*[Gal4] and *dsx*[LexA::p65] are shown in magenta. (C) The number of *dsf*[Gal4]-expressing neurons (circles) and *dsf*[Gal4]-,

404     $dsx^{\text{LexA::p65}}$-co-expressing (*i.e.*, DDAG) neurons (diamonds) per hemisphere in the abdominal ganglion of

females and males is shown. From 0–18 hours APF, both sexes have ~20 dsf^{Gal4}-expressing neurons, and

406   the number of dsf^{Gal4}- and $dsx^{\text{LexA::p65}}$-co-expressing neurons gradually increases to ~11 neurons. Between

18–48 hours APF, ~8 DDAG neurons gradually disappear in males but not in females leaving a total of 11

408   and 3 DDAG neurons in females and males, respectively. Individual points, mean, and SD are shown.

410   **Figure 5. Two A26g neurons in the larval abdominal ganglion become the DDAG_C and DDAG_D**

neurons of adult females. (A, B) Confocal images of larval CNSs from $dsf^{\text{p65AD::Zp}} \cap VT026005\text{-}ZP\text{::}GDBD >$

412   *myr::gfp* (A) females and (B) males. GFP-expressing neurons are labeled in black and DNCad (neuropil) is

shown in light gray. The A26g neuron is labeled in segments A4–A6. (C) A confocal image of a single

414   A26g neuron at A4 or A5 using the multicolor Flp-out technique. Coronal and sagittal views are shown.

(D–G) Confocal images of the abdominal ganglion at various time points after puparium formation (APF)

416   from $dsf^{\text{p65AD::Zp}} \cap VT026005\text{-}ZP\text{::}GDBD > myr::gfp$ females. GFP-expressing neurons are labeled in green

and DSX-F-expressing neurons are labeled in magenta. (D) At 0 hours APF, the A26g neuron at A4–A6 is

418   labeled by GFP, but none of them express DSX-F. (E) At 24 hours APF, the A26g neuron at A5 and A6

express DSX-F. Two additional neurons at A7 and A8 are labeled by $dsf^{\text{p65AD::Zp}} \cap VT026005\text{-}ZP\text{::}GDBD$,

420   both of which express DSX-F. (F) By 48 hours APF, the anatomy of the neurons labeled by $dsf^{\text{p65AD::Zp}} \cap$

$VT026005\text{-}ZP\text{::}GDBD$ is similar to that of the DDAG_B–D neurons of adults. (G) By adulthood, only

422   neurons at A5 (DDAG_D), A6 (DDAG_C), A7, and A8 (DDAG_B) are labeled by GFP and DSX-F. (H)

Confocal images of posterior VNCs from $dsf^{\text{p65AD::Zp}} \cap VT026005\text{-}ZP\text{::}GDBD > UAS\text{-}hPR\text{::}Flp$, $dsx^{\text{LexAp65}} >$

424   *LexAop2-frt.stop-myr::gfp* adult females fed ethanol- or mifepristone-containing food during larval life.

GFP-expressing neurons are labeled in black and DNCad (neuropil) is shown in light gray. Expression of

426   GFP in the DDAG_C and DDAG_D neurons of adults indicates that the cells existed among the $dsf^{\text{p65AD::Zp}}$

$\cap VT026005\text{-}ZP\text{::}GDBD$-expressing neurons in the larval abdominal ganglion. Although the Split-Gal4

428   labels the DDAG_B neurons after larval life (*i.e.*, between 0–24 hours APF), the DDAG_B neurons were

also labeled in these experiments. This was likely due to the perdurance of residual mifepristone after

430   pupariation.

432   **Figure 6. A26g neurons in larvae become the DDAG_D and DDAG_C neurons in females but undergo**

***dsx*-dependent cell death in males.** A model summarizing the results described in this paper. See text for

434   details.

 STAR METHODS

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
rabbit anti-GFP	Invitrogen	Cat#A11122; RRID:AB_221569
rat anti-DN-cadherin	Developmental Studies Hybridoma Bank	Cat#DN-Ex#8; RIDD:AB_528121
rabbit anti-DSX-F	Peng et al.[32]	N/A
mouse anti-DSX-M	Peng et al.[32]	N/A
mouse anti-HA.11	BioLegend	Cat#MMS-101P; RRID:AB_291261
rat anti-FLAG	Novus Biologicals	Cat#NBP1-06712; RRID:1625981
Fluorescein (FITC)-conjugated donkey anti-rabbit	Jackson ImmunoResearch	Cat#711-095-152; RIDD:AB_2315776
goat anti-rat AlexaFluor 647	Invitrogen	Cat#A21247; RRID:AB_141778
donkey anti-rat AlexaFluor 568	Invitrogen	Cat#A78946; RRID:AB_2910653
donkey anti-mouse AlexaFluor 647	Invitrogen	Cat#A31571; RRID:AB_162542
goat anti-mouse AlexaFluor 488	Invitrogen	Cat#A32723; RRID:AB_2633275
Chemicals, peptides, and recombinant proteins		
DPX	Sigma-Aldrich	Cat#06522
all-*trans*-Retinal	Sigma-Aldrich	Cat#R2500
Mifepristone (RU-486)	Sigma-Aldrich	Cat#475838-50MG
Experimental models: Organisms/strains		
Canton S	Duckhorn et al.[9]	N/A
w[1118]	Duckhorn et al.[9]	N/A
dsx[LexA::p65]*/TM6B*	Zhou et al.[33]	N/A
dsf[Gal4]*/CyO*	Duckhorn et al.[9]	N/A
pJFRC29-10XUAS-IVS-myr::GFP-p10 (attP2)	Janelia Research Campus (JRC), HHMI	N/A
pJFRC12-10XUAS-IVS-myr::GFP (attP2)	JRC	
pJFRC79-8XLexAop-2-FlpL (attP40)	JRC	N/A
pJFRC41-10XUAS-FRT-STOP-FRT-myr::gfp (su(Hw)attP1)	JRC	N/A
pJFRC108-20XUAS-IVS-hPR-Flp-p10 (attP2)	JRC	N/A
pJFRC40-13XLexAop-FRT-STOP-FRT-myr::gfp (attP40)	JRC	N/A
pJFRC56-10XUAS-FRT-STOP-FRT-kir2.1::gfp (attP2)	JRC	N/A
pBPhsFlp2::PEST (attP3)	JRC	N/A
pJFRC201-10XUAS-FRT-STOP-FRT-myr::smGFP-HA (VK0005)	JRC	N/A
pJFRC240-10XUAS-FRT>STOP>FRTmyr::smGFP-V5-THS-10XUAS-FRT>STOP>FRT-myr::smGFP-FLAG (su(Hw)attP1)	JRC	N/A
20XUAS-FRT>STOP>FRT-CsChrimson::mVenus (VK5)	JRC	N/A

VT026005-Zp::GDBD (attP2)	JRC	N/A
UAS-P35[BH1]	BDSC	RRID:BDSC_5072
UAS-dsx_ShmiR (attP2)	Duckhorn et al.[9]	RRID:BDSC_35645
UAS-dsf_ShmiR (attP2)	Duckhorn et al.[9]	N/A
dsf[Del]	Duckhorn et al.[9]	N/A
dsf[p65AD::Zp]/CyO	This study	N/A
dsf[DBD::Zp]/CyO	This study	N/A
dsf[LexA::p65]/CyO	This study	N/A
Software and algorithms		
Fiji	NIH, USA	https://imagej.net/fiji
MATLAB	Mathworks	https://www.mathworks.com/products/matlab.html

RESOURCE AVAILABILITY

440 **Lead contact**

Further information and requests for resources and reagents should be directed to and will be fulfilled by

442 the lead contact, Troy Shirangi (troy.shirangi@villanova.edu).

444 **Materials availability**

Fly lines generated in this study are available from the lead contact.

Data and code availability

448 All data reported in this paper will be shared by the lead contact upon request. This paper does not report

original code. Any additional information required to reanalyze the data reported in this paper is

450 available from the lead contact upon request.

452 **EXPERIMENTAL MODEL AND SUBJECT DETAILS**

Drosophila melanogaster stocks were maintained on standard cornmeal and molasses food at 25°C and

454 ~50% humidity in a 12-hr light/dark cycle unless otherwise noted. Fly stocks used in this study are listed

in the Key Resources Table. *dsf[p65AD::Zp]*, *dsf[Zp::GDBD]*, and *dsf[LexA::p65]* alleles were generated using the same

456 strategy as that used to build the *dsf[Gal4]* allele[9] except the Gal4 sequence in the donor construct was

replaced with sequences encoding *p65AD::Zp*, *Zp::GDBD*, or *LexA::p65*.

METHOD DETAILS

460 **Immunohistochemistry**

Nervous systems were dissected in 1X phosphate-buffered saline (PBS), fixed in 4% paraformaldehyde in

462   PBS for 35 minutes, then rinsed and washed in PBT (PBS with 1% Triton X-100). If a blocking step was

performed, nervous systems were incubated in 5% normal goat serum or 5% normal donkey serum in

464   PBT for 30 minutes. Tissues were then incubated with primary antibodies diluted in PBT or PBT with

block overnight at 4°C. The next day, three washes were performed over the course of several hours

466   before nervous systems were incubated in secondary antibodies diluted in PBT or PBT with block

overnight at 4°C. Tissues were then washed three times over the course of several hours and placed on

468   cover slips coated in poly-lysine, dehydrated in an increasing ethanol concentration series, and cleared in

a xylene series. Nervous systems were mounted onto slides using DPX mounting medium and imaged on

470   a Leica TCS SP8 Confocal Microscope at 40X magnification. For MultiColor FlpOut experiments, vials

containing larvae aged between first and second instars (or 3–4-day old adults) were placed in a 37°C

472   water bath for 1–10 minutes then dissected 2 days later. To stochastically label the DDAG neurons, male

and female adults (deprived of food overnight) were placed in vials with food containing 100mM

474   mifepristone (RU-486; Sigma 475838-50MG) for 2 hours and kept in darkness. Flies were then transferred

back to vials containing untreated food for 3 days before their VNCs were dissected for staining. For

476   immortalization experiments using mifepristone, three days after crosses were set-up on normal food, 60

μL of 100mM RU-486 was added directly to the food and larvae were raised in darkness on RU-486-

478   treated food until pupariation. Pupae were collected and transferred to vials containing untreated food

before eclosure. Control groups were also kept in darkness but were treated with 60 μL of ethanol instead

480   of RU-486. Adult nervous systems were dissected in PBS. The following primary antibodies were used:

rabbit anti-GFP (Invitrogen #A11122; 1:1000), rat anti-DN-cadherin (DN-Ex#8, Developmental Studies

482   Hybridoma Bank; 1:50), anti-DSX-F (1:200)[32], mouse anti-HA.11 (BioLegend #MMS-101P; 1:250), and rat

anti-FLAG (Novus Biologicals #NBP1-06712; 1:200). The following secondary antibodies were used:

484   Fluorescein (FITC) conjugated donkey anti-rabbit (Jackson ImmunoResearch #711-095-152; 1:500), AF-647

goat anti-rat (Invitrogen #A21247; 1:500), AF-568 donkey anti-rat (Invitrogen #A78946; 1:500), AF-647

486   donkey anti-mouse (Invitrogen #A31571; 1:500), AF-488 goat anti-mouse (Invitrogen #A32723), AF-647

goat anti-rat (Invitrogen #A21247; 1:500).

Optogenetic assays

490   Unmated females used in optogenetic assays were raised in darkness and on food containing 0.2mM all-

trans-retinal (sigma-Aldrich #R2500) and were incubated at 25°C and ~50% humidity. Once collected,

492   unmated females were grouped in vials consisting of 15–20 flies for 8–12 days before testing. Flies were

anesthetized on ice for ~2 minutes, decapitated under low-intensity light, and were given 15–20 minutes

494   to recover before being transferred to individual behavioral chambers (diameter: 10 mm, height: 3 mm). A

FLIR Blackfly S USB3, BFS-U3-31S4M-C camera with a 800 nm long-pass filter (Thorlabs, FEL0800) was

496   used to record optogenetic videos in SpinView. Upon testing, chambers were placed on top of an LED

panel with continuous infrared (850 nm) light and recurring photoactivating red (635 nm) light using an

498   Arduino script. To measure change in abdominal length before and during vaginal plate opening or

ovipositor extrusion, a ruler (cm/mm) was included in the frame to set the scale. The change in abdominal

500   length was calculated as the difference in the abdominal length from the base of the scutellum to the tip

of the abdomen before and during photoactivation. Behavior indices were measured by calculating the

502   average fraction of time spent preforming the behavior during the first three 15-second lights-on periods

and the first three 45-second light-off periods. For stochastic optogenetic activation, unmated females

504   carrying *hs-Flp*, *UAS-frt.stop-CsChrimson::mVenus*, and the Split-Gal4 were reared on retinal-containing

food, grouped in vials consisting of 15–20 flies, and aged for ~3 days before being placed in a 37°C water

506   bath for periods ranging from 20–60 minutes. Females were then transferred to new vials containing

retinal food and aged for an additional 5 days before being tested in an optogenetic activation experiment

508   as described above. Following optogenetics, the VNC of each female was dissected and placed singly in

wells of a 60-well mini tray (Fischer Scientific #12-565-155) for staining. Each VNC was subsequently

510   mapped to the female in the optogenetics experiment from which the VNC was obtained.

512   **Behavioral assays**

Unmated females and males were collected were under CO_2 and aged for 7–10 days in a 12-hour

514   light/dark cycle and incubated at 25°C and ~50% humidity. Unmated females were group-housed in vials

consisting of 15–20 flies, and *Canton S* males were individually housed. Courtship assays were done

516   within the first two hours of the subjective day. Unmated females and *Canton S* males were transferred to

individual behavioral chambers (diameter: 10 mm, height: 3 mm) and recorded for 30 minutes using a

518   Sony Vixia HFR700 video camera at 25°C under white light. For experiments using mated females,

unmated females were housed with males for 24 hours, anesthetized on ice for ~2 minutes, and mated

520   females were collected into a new vial and given 30 minutes to recover. Mated females and *Canton S*

males were loaded to chambers and recorded as described above. Courtship index was measured as the

522   total time the male preformed courtship behaviors divided by the total recording time. Courtship index

was measured as the total time the male performed courtship behaviors divided by the observation time

524   which was usually about 5 minutes. Vaginal plate opening (vpo) and ovipositor extrusion (oe) frequency

was measured as the total number of times a female performed a vpo or oe in a 6-min period of active

526   male courtship. Egg laying was measured by allowing females to mate with males before transferring

them to individual vials to for 24 hours. The total number of eggs laid in 24 hours by each female was

528   then counted.

530   **QUANTIFICATION AND STATISTICAL ANALYSIS**

Data were analyzed using one-way ANOVA with Tukey-Kramer tests for multiple comparisons, Rank

532   Sum tests, or Logrank tests. All p-values were measured in MATLAB.

534   **Legends for Supplemental Videos**

Video S1. Anatomy of the DDAG_B, DDAG_C, DDAG_D, and A26g neurons, Related to Figure 2.

536   **Video S2. Optogenetic activation of the DDAG_B–D neurons in unmated females, Related to Figure 2.**

Video S3. Optogenetic activation of the DDAG_B–D neurons in mated females, Related to Figure 2.

Video S4. Bilateral optogenetic activation of the DDAG_C neurons in unmated females, Related to Figure 2.

Video S5. Bilateral optogenetic activation of the DDAG_D neurons in unmated females, Related to Figure 2.

Video S6. Unilateral optogenetic activation of a DDAG_C neuron in unmated females, Related to Figure 2.

Video S7. Unilateral optogenetic activation of a DDAG_D neuron in unmated females, Related to Figure 2.

Video S8. Unilateral optogenetic activation of a DDAG_B neuron in unmated females, Related to Figure 2.

Video S9. Bilateral optogenetic activation of the DDAG_B neurons in unmated females, Related to Figure 2.

Video S10. Optogenetic activation of a $dsf^{p65AD::Zp} \cap VT026005\text{-}Zp::GDBD > CsChrimson::mVenus$ male with depleted dsx transcripts, Related to Figure 5.

Video S11. Optogenetic activation of a $dsf^{p65AD::Zp} \cap VT026005\text{-}Zp::GDBD > CsChrimson::mVenus$ unmated female with depleted dsx transcripts, Related to Figure 5.

Video S12. Optogenetic activation of a $dsf^{p65AD::Zp} \cap VT026005\text{-}Zp::GDBD > CsChrimson::mVenus$ unmated female with depleted dsf transcripts, Related to Figure 5.

REFERENCES

1. Sisk, C.L., and Zehr, J.L. (2005). Pubertal hormones organize the adolescent brain and behavior. Front
   Neuroendocrinol *26*, 163–174. 10.1016/j.yfrne.2005.10.003.

2. Ahmed, E.I., Zehr, J.L., Schulz, K.M., Lorenz, B.H., DonCarlos, L.L., and Sisk, C.L. (2008). Pubertal
   hormones modulate the addition of new cells to sexually dimorphic brain regions. Nat Neurosci *11*,
   995–997. 10.1038/nn.2178.

3. Truman, J.W., Price, J., Miyares, R.L., and Lee, T. (2023). Metamorphosis of memory circuits in
   Drosophila reveals a strategy for evolving a larval brain. Elife *12*, e80594. 10.7554/eLife.80594.

4. Witvliet, D., Mulcahy, B., Mitchell, J.K., Meirovitch, Y., Berger, D.R., Wu, Y., Liu, Y., Koh, W.X.,
   Parvathala, R., Holmyard, D., et al. (2021). Connectomes across development reveal principles of brain
   maturation. Nature *596*, 257–261. 10.1038/s41586-021-03778-8.

5. Truman, J.W., and Riddiford, L.M. (2023). Drosophila postembryonic nervous system development: a
   model for the endocrine control of development. Genetics *223*, iyac184. 10.1093/genetics/iyac184.

6. Carreira-Rosario, A., Zarin, A.A., Clark, M.Q., Manning, L., Fetter, R.D., Cardona, A., and Doe, C.Q.
   (2018). MDN brain descending neurons coordinately activate backward and inhibit forward
   locomotion. Elife *7*, e38554. 10.7554/eLife.38554.

7. Cachero, S., Ostrovsky, A.D., Yu, J.Y., Dickson, B.J., and Jefferis, G.S.X.E. (2010). Sexual dimorphism in
   the fly brain. Curr Biol *20*, 1589–1601. 10.1016/j.cub.2010.07.045.

8. Ren, Q., Awasaki, T., Huang, Y.-F., Liu, Z., and Lee, T. (2016). Cell Class-Lineage Analysis Reveals
   Sexually Dimorphic Lineage Compositions in the Drosophila Brain. Curr Biol *26*, 2583–2593.
   10.1016/j.cub.2016.07.086.

9. Duckhorn, J.C., Cande, J., Metkus, M.C., Song, H., Altamirano, S., Stern, D.L., and Shirangi, T.R. (2022).
   Regulation of Drosophila courtship behavior by the Tlx/tailless-like nuclear receptor, dissatisfaction.
   Curr Biol *32*, 1703-1714.e3. 10.1016/j.cub.2022.02.031.

10. Pan, Y., Robinett, C.C., and Baker, B.S. (2011). Turning males on: activation of male courtship behavior
    in Drosophila melanogaster. PLoS One *6*, e21144. 10.1371/journal.pone.0021144.

11. Harris, R.M., Pfeiffer, B.D., Rubin, G.M., and Truman, J.W. (2015). Neuron hemilineages provide the
    functional ground plan for the Drosophila ventral nervous system. Elife *4*, e04493. 10.7554/eLife.04493.

12. Nern, A., Pfeiffer, B.D., and Rubin, G.M. (2015). Optimized tools for multicolor stochastic labeling
    reveal diverse stereotyped cell arrangements in the fly visual system. Proc Natl Acad Sci U S A *112*,
    E2967-2976. 10.1073/pnas.1506763112.

13. Wang, K., Wang, F., Forknall, N., Yang, T., Patrick, C., Parekh, R., and Dickson, B.J. (2021). Neural
    circuit mechanisms of sexual receptivity in Drosophila females. Nature *589*, 577–581. 10.1038/s41586-
    020-2972-7.

592 14. Mezzera, C., Brotas, M., Gaspar, M., Pavlou, H.J., Goodwin, S.F., and Vasconcelos, M.L. (2020). Ovipositor Extrusion Promotes the Transition from Courtship to Copulation and Signals Female
594 Acceptance in Drosophila melanogaster. Curr Biol *30*, 3736-3748.e5. 10.1016/j.cub.2020.06.071.

15. Wang, F., Wang, K., Forknall, N., Parekh, R., and Dickson, B.J. (2020). Circuit and Behavioral
596 Mechanisms of Sexual Rejection by Drosophila Females. Curr Biol *30*, 3749-3760.e3. 10.1016/j.cub.2020.07.083.

598 16. Klapoetke, N.C., Murata, Y., Kim, S.S., Pulver, S.R., Birdsey-Benson, A., Cho, Y.K., Morimoto, T.K., Chuong, A.S., Carpenter, E.J., Tian, Z., et al. (2014). Independent optical excitation of distinct neural
600 populations. Nat Methods *11*, 338–346. 10.1038/nmeth.2836.

17. Baines, R.A., Uhler, J.P., Thompson, A., Sweeney, S.T., and Bate, M. (2001). Altered electrical properties
602 in Drosophila neurons developing without synaptic transmission. J Neurosci *21*, 1523–1531. 10.1523/JNEUROSCI.21-05-01523.2001.

604 18. Burtis, K.C., and Baker, B.S. (1989). Drosophila doublesex gene controls somatic sexual differentiation by producing alternatively spliced mRNAs encoding related sex-specific polypeptides. Cell *56*, 997–
606 1010. 10.1016/0092-8674(89)90633-8.

19. Oren-Suissa, M., Bayer, E.A., and Hobert, O. (2016). Sex-specific pruning of neuronal synapses in
608 Caenorhabditis elegans. Nature *533*, 206–211. 10.1038/nature17977.

20. Lee, G., Hall, J.C., and Park, J.H. (2002). Doublesex gene expression in the central nervous system of
610 Drosophila melanogaster. J Neurogenet *16*, 229–248. 10.1080/01677060216292.

21. Robinett, C.C., Vaughan, A.G., Knapp, J.-M., and Baker, B.S. (2010). Sex and the single cell. II. There is a
612 time and place for sex. PLoS Biol *8*, e1000365. 10.1371/journal.pbio.1000365.

22. Rideout, E.J., Dornan, A.J., Neville, M.C., Eadie, S., and Goodwin, S.F. (2010). Control of sexual
614 differentiation and behavior by the doublesex gene in Drosophila melanogaster. Nat Neurosci *13*, 458–466. 10.1038/nn.2515.

616 23. Truman, J.W., and Bate, M. (1988). Spatial and temporal patterns of neurogenesis in the central nervous system of Drosophila melanogaster. Dev Biol *125*, 145–157. 10.1016/0012-1606(88)90067-x.

618 24. Taylor, B.J., and Truman, J.W. (1992). Commitment of abdominal neuroblasts in Drosophila to a male or female fate is dependent on genes of the sex-determining hierarchy. Development *114*, 625–642.
620 10.1242/dev.114.3.625.

25. Birkholz, O., Rickert, C., Berger, C., Urbach, R., and Technau, G.M. (2013). Neuroblast pattern and
identity in the Drosophila tail region and role of doublesex in the survival of sex-specific precursors. Development *140*, 1830–1842. 10.1242/dev.090043.

26. Truman, J.W., and Reiss, S.E. (1976). Dendritic reorganization of an identified motoneuron during metamorphosis of the tobacco hornworm moth. Science *192*, 477–479. 10.1126/science.1257782.

27. Technau, G., and Heisenberg, M. (1982). Neural reorganization during metamorphosis of the corpora pedunculata in Drosophila melanogaster. Nature *295*, 405–407. 10.1038/295405a0.

28. Williams, D.W., and Shepherd, D. (1999). Persistent larval sensory neurons in adult Drosophila melanogaster. J Neurobiol *39*, 275–286.

29. Lee, K., and Doe, C.Q. (2021). A locomotor neural circuit persists and functions similarly in larvae and adult Drosophila. Elife *10*, e69767. 10.7554/eLife.69767.

30. Consoulas, C., Restifo, L.L., and Levine, R.B. (2002). Dendritic remodeling and growth of motoneurons during metamorphosis of Drosophila melanogaster. J Neurosci *22*, 4906–4917. 10.1523/JNEUROSCI.22-
12-04906.2002.

    31. Sprecher, S.G., and Desplan, C. (2008). Switch of rhodopsin expression in terminally differentiated
636   Drosophila sensory neurons. Nature *454*, 533–537. 10.1038/nature07062.

    32. Peng, Q., Chen, J., Su, X., Wang, R., Han, C., and Pan, Y. (2022). The sex determination gene doublesex
638   is required during adulthood to maintain sexual orientation. J Genet Genomics *49*, 165–168. 10.1016/j.jgg.2021.08.006.

640   33. Zhou, C., Pan, Y., Robinett, C.C., Meissner, G.W., and Baker, B.S. (2014). Central brain neurons expressing doublesex regulate female receptivity in Drosophila. Neuron *83*, 149–163.
642   10.1016/j.neuron.2014.05.038.

A

male

coronal   sagittal

ant   ant
↑   vent↖

female

coronal   sagittal

DDAG_all

*dsf*GAL4 ∩ *dsx*LexAp65 > *myr::GFP*

B

coronal   sagittal

ant   ant
↑   vent↖

DDAG_A^male

coronal   sagittal   coronal   sagittal   coronal   sagittal   coronal   sagittal   coronal   sagittal

ant   ant
↑   vent↖

DDAG_A^female   DDAG_B   DDAG_C   DDAG_D   DDAG_E

*dsf*GAL4 > *hPR::Flp* ∩ *dsx*LexAp65 > *LexAop2-frt.stop-myr::gfp*

A

brain	ventral nerve cord

female

coronal   sagittal

ant ↑   ant ↑ vent ←

$dsf^{p65AD::Zp}$ ∩ $VT026005$-$Zp::GDBD$ ∩ $dsx^{LexAp65}$ > $myr::gfp$

B

kir2.1::gfp

C

(n≥40)   n.s.

fraction that mated
1.0
0.8
0.6
0.4
0.2
0
0   5   10   15   20   25   30
time (min)

D

α   α   α

vpo/min of courtship
4

E

α   α   β

oe/min of courtship
5

2.5

F

α   α   β

eggs/24 hrs
100

● X/X; $dsf^{p65AD::Zp}/+$; $VT026005$-$Zp::GDBD$, $dsx^{LexAp65}/+$ ● X/X; $+/LexAop2$-$FlpL$; $+/UAS$-$frt.stop$-$kir2.1::gfp$ ● X/X; $dsf^{p65AD::Zp}/LexAop2$-$FlpL$; $VT026005$-$Zp::GDBD$, $dsx^{LexAp65}/UAS$-$frt.stop$-$kir2.1::gfp$

A female

B male

C coronal sagittal

$dsf^{p65AD::Zp} \cap VT026005\text{-}Zp::GDBD > myr::gfp$

A26g_A4/5

0 hour APF   24 hour APF   48 hour APF   adult

GFP

DSX-F

merge

$dsf^{p65AD::Zp} \cap VT026005\text{-}Zp::GDBD > myr::gfp$

H larval ethanol   coronal sagittal   larval mifepristone   DDAG_B coronal sagittal   DDAG_C coronal sagittal   DDAG_D coronal sagittal

$dsf^{p65AD::Zp} \cap VT026005\text{-}Zp::GDBD > UAS\text{-}hPR::Flp, dsx^{LexAp65} > LexAop2\text{-}frt.stop\text{-}myr::gfp$