High-throughput experimental evolution reveals the mutation-driven landscape of high-level clinical resistance acquired under low-level environmental antibiotic stresses

Hanqing Wang[1], Hui Lu[2], Chao Jiang[3], Lizhong Zhu[1]*, Huijie Lu[1]*

1. College of Environmental and Resource Sciences, Zhejiang University, Hangzhou 310058, China
2. ZJU-Hangzhou Global Scientific and Technological Innovation Center, Zhejiang University, Hangzhou 311215, China
3. Life Sciences Institute, Zhejiang University, Hangzhou, 310058, China

*To whom correspondence should be addressed:
Huijie Lu (luhuijie@zju.edu.cn)
Lizhong Zhu (zlz@zju.edu.cn)

Abstract

Potential pathogens exposed to low-level environmental antibiotics could develop high-level clinically relevant antibiotic resistance detrimental to the health of the general population. However, the underlying evolutionary landscapes remain poorly understood. We conducted a high-throughput experimental evolution study by exposing an environmentally isolated pathogenic *Escherichia coli* strain to 96 typical antibiotics at 10 μg l^{-1} for 20 days. Antibiotic resistance phenotypic (IC$_{90}$ against 8 clinically used antibiotics) and genetic changes of the evolved populations were systematically investigated, revealing a universal increase in antibiotic resistance (up to 349-fold), and mutations in 2,432 genes. Transposon sequencing was further employed to verify genes potentially associated with resistance. A core set of mutant genes conferring high-level resistance was analyzed to elucidate their resistance mechanisms by analyzing the functions of interacted genes within the gene co-fitness network and performing gene knockout validations. We developed machine-learning models to predict antibiotic resistance phenotypes from antibiotic structures and genomic mutations, enabling the resistance predictions for another 569 antibiotics. Importantly, 14.6% of the 481 key mutations were observed in clinical and environmental *E. coli* isolates retrieved from the NCBI database, and several were over-represented in clinical isolates. Deciphering the evolutionary landscapes underlying resistance exposed to low-level environmental antibiotics is crucial for evaluating the emergence and risks of environment-originated clinical antibiotic resistance.

Introduction

The emergence of antimicrobial resistance (AMR) in the environment is a growing concern in the One Health paradigm[1–5]. Antibiotics are increasingly released into the environment via wastewater discharge and disposal of solid waste, where 40-90% are excreted as parent compounds[6]. Environmental bacteria exposed to these antibiotics at low-level, sub-minimal inhibitory concentrations (sub-MIC) can evolve antibiotic resistance by accumulating diverse low-frequency mutants, which are different from those evolved at clinical concentrations[7–11]. The environmental research community essentially needs to elucidate the contribution of mutation-driven resistance evolution in the environment by crafting analytical measures to pinpoint the responsible selective forces (i.e., antibiotics) driving AMR development[12]. Therefore, exploring the evolutionary landscape under low-level antibiotic exposures is increasingly important at the interface between environmental and clinical microbiology pertaining to antibiotic resistance[4,13].

Traditional low-throughput experimental evolution has provided an incomplete view of the AMR evolutionary landscape, as the number of stress conditions, mutants, and resistance-conferring genes examined were limited[14,15]. High-throughput experimental evolution enables the rapid generation and systematic investigation of population-level antibiotic resistance under a variety of stress conditions, allowing better characterization of the structure- or dose-response relationships among stressors and induced resistance[16,17]. In addition, antibiotic resistance genes (ARGs) were traditionally identified by sequencing and verified through phenotypic assays/knockout experiments, bearing the risks of overlooking critical insights from genetic interactions. Transposon sequencing (Tn-Seq) measures genome-wide gene fitness across large-scale transposon mutagenesis libraries. By combining Tn-Seq with deep resequencing, researchers can map the accumulated mutations onto complex and coordinated interaction networks. The

prevalence and clinical significance of identified mutations in broad environments

can be examined by comparative genomics[18]. The above novel approaches could

help address critical knowledge gaps in the field of environmental AMR, for

example: (1) What is the association between the antibiotic resistance of evolved

populations and the chemical structures of antibiotics under low-level exposures?

(2) What cellular functions are affected by mutations potentially via new antibiotic

resistance mechanisms? (3) Are mutations that arise under low-level antibiotic

exposures observed in clinically isolated resistant strains?

This study utilized a soil-isolated opportunistic pathogen *E. coli* strain to

perform high-throughput experimental evolution under low-level exposures to 96

antibiotic stressors (**Supplementary Fig. 1**). The antibiotic resistance profiles of

evolved populations and their core mutations were characterized. To explore the

resistance mechanisms conferred by these core mutant genes, a genome-wide

co-fitness interaction network was constructed based on Tn-seq. Chemical

fingerprint-based modeling was built to predict antibiotic resistance profiles

induced by another 569 antibiotics sharing similar structural features with the 96

stressors. The occurrence of core mutations in broader environmental and clinical

isolates was examined by comparative genomics. These findings underscore the

importance of AMR developed under low-level antibiotic exposures in

environmental settings and its potential spread to clinical contexts.

Results

Antibiotic resistance profiles of evolved *E. coli* QSHQ1 populations

   E. coli strain QSHQ1 isolated from soil evolved under exposure to 96 antibiotic stresses at an environmentally relevant concentration of 10 μg l^{-1} for 20 days, which corresponded to approximately 200 generations (**Supplementary Data 1**). These antibiotics can be classified into ten classes and mainly act through four mechanisms, including inhibition of DNA/RNA (DSI), protein (PSI), cell wall (CWSI), and membrane (MSI) synthesis (**Fig. 1a**). All populations survived and displayed varying degrees of growth inhibition compared to control populations without antibiotics (**Supplementary Fig. 2**). Evolved populations on day 20 were tested for antibiotic resistance (indicated by IC$_{90}$) to eight clinically relevant antibiotics, including azithromycin (AZI), ciprofloxacin (CIP), tetracycline (TET), sulfamethoxazole with trimethoprim (SMZ+TMP), tobramycin (TOB), ceftazidime (CAZ), meropenem (MER), and polymyxin B (PMB). These IC$_{90}$ values were compared with CLSI's breakpoint values for susceptible and resistant bacteria (**Fig. 1b** and **Supplementary Data 2**).

   The wild-type (WT) strain was initially susceptible or intermediate to the eight clinically relevant antibiotics. All 96 evolved populations conferred resistance (increased IC$_{90}$) to at least one of the eight antibiotics (Multiple unpaired *t*-tests, FDR < 0.1, **Fig. 1a**), where resistance to AZI (78.9%) and TOB (18.8%) were the most prevalent. A similar evolution of AZI-resistant strains was recently reported for *Comamonas testosteroni*[10]. Only two evolved populations stressed by marbofloxacin and ceftriaxone displayed significantly decreased IC$_{90}$ values to at least one antibiotic. Specifically, a ciprofloxacin-stressed population displayed the most significant increase in CIP resistance (IC$_{90}$ increased by 345-fold), and a cefazolin-stressed population displayed a 349-fold increase in MER resistance (**Fig. 1b**). Previous studies[19,20] reported 256-fold and >300-fold increase in the MIC

for ciprofloxacin of *E. coli* and *S. aureus* cells after < 20 passages, respectively. In fact, the WT strain was susceptible to CIP and MER. Exposure to quinolones was the primary driver of CIP resistance evolution (average IC_{90} fold-change: 24.6 ± 14.9), while exposure to aminoglycosides highly contributed to the evolution of TOB resistance (average IC_{90} fold-change: 2.19 ± 0.26) (**Fig. 1c**, **Supplementary Figs. 3** and **4**).

Populations evolved under each class of antibiotics displayed variations in resistance to the eight antibiotics tested, where 5/8 displayed either significantly positive or negative correlations induced by a specific class of antibiotics (**Fig. 1d** and **Supplementary Fig. 5**). Overall, positive correlations (Pearson's *r* > 0.6, *P* < 0.05) were much more abundant than negative ones, where the latter was only observed for macrolides-stressed populations and between their AZI and MER resistance (*r* = −0.85, *P* < 0.05). The 96 antibiotic stressors were further clustered based on the antibiotic resistance profiles they induced (**Fig. 1e** and **Supplementary Fig. 6**). Most aminoglycosides and macrolides (both acting via PSI) were clustered (VI), as well as sulfonamides and tetracyclines (acting via DSI and PSI, respectively, cluster II). Quinolones formed their distinct cluster (cluster III), separated from other antibiotic classes.

Fig. 1: Antibiotic resistance profiles of evolved populations. **a,** Antibiotic resistance profiles of the evolved populations. Antibiotic resistance changes were indicated by log2-transformed fold changes in IC90 values (relative to the control populations without antibiotic stress). The tree was constructed based on Euclidean distances. Antibiotics stressors: 96; Stressor classes: 10 (different colors of the inner circle); Stressor groups: 4 (DSI, CWSI, PSI, and MSI); Antibiotic resistance assay: 8 (heatmap circles); Biological replicates: 4. **b,** Distribution of antibiotic resistance changes in evolved populations. Grey dots: log2-transformed fold changes in IC90 values of all independently evolved populations; Curves: normal distribution approximation; Colored sticks: susceptible (blue) and resistant (red) breakpoints for the tested antibiotic resistance according to CLSI guidelines[21]. Box plot: center

line, median; small square, mean; left and right sides of the box, upper and lower quartiles; whiskers, 1.5× interquartile range. **c,** Averaged resistance changes ($\log_2$-transformed fold changes in IC_{90} values) of evolved populations stressed by the same class of antibiotics. **d,** Positive and negative correlations between different antibiotic classes according to the antibiotic resistance profiles (Pearson's correlation coefficient $|r| > 0.6$, $P < 0.05$). **e,** Uniform manifold approximation and projection (UMAP) embedding of evolved populations based on their resistance to eight antibiotics. **c**, **d**, and **e** are assigned the same color code as **a** for antibiotic classes.

Phenotypic convergence and divergence of antibiotic resistance evolution

Pairwise antibiotic stressors induced convergent (Spearman's correlation coefficient $\rho > 0.2$, 76%) and divergent ($\rho < -0.2$, 4.6%) antibiotic resistance phenotypes among the 96 antibiotics (**Figs. 2a and 2b**). Convergent phenotypes (a phenomenon whereby developing similar phenotypes) were predominant and occurred more frequently within the same class of stressors (85.3% vs. 73.9%, **Fig. 2c**). Divergent phenotypes (orthogonal), otherwise, were more found across different antibiotic classes than within the same class (5.1% vs. 2.3%). Overall, the frequency of convergence was much higher than the previously reported value (60%) for *P. aeruginosa* strains that evolved under high-level antibiotic exposures[22]. We hypothesize that the survival of more diverse resistant subpopulations under low-level antibiotic exposures likely contributes to widespread elevated population antibiotic resistance, as these subpopulations may possess similar off-target mutations[23]. However, the consistency of the mutation similarity and convergent phenotypes await confirmation via deep sequencing. Furthermore, stressors within aminoglycosides and lipopeptides (LPS) induced the most convergent resistance phenotypes, while quinolones induced the most divergent resistance patterns compared to other antibiotic classes (**Fig. 2d**). Convergent resistance phenotypes were widespread across classes. Divergent phenotypes were more frequently observed between aminoglycosides-quinolones and aminoglycosides-tetracyclines, e.g., kanamycin and sparfloxacin ($\rho = -0.81$; $P = 0.0149$) than other pairwise combinations (**Fig. 2e** and **Supplementary Fig. 7**). Our analyses of these stressor-stressor pairs corroborated the previous findings of resistance evolution leading to phenotypic convergence and identified new population-level convergent resistance evolution in sub-MIC antibiotic selections[22].

184 n = 3,666 pairs ■ Convergent evolution ■ Divergent evolution

Fig. 2: Convergent and divergent evolution relationships based on antibiotic resistance profiles induced by low-level antibiotic exposures. **a,** Total number of convergent (con.) and divergent (div.) evolution relationships. Strong convergent or divergent evolution is presented in dark red or blue, respectively (Spearman's rank correlation |ρ| > 0.6); light red or blue suggests weak convergent or divergent evolution (0.2< |ρ| <0.6). **b,** Convergent and divergent evolution relationships between stressors belonging to different (across) or same (within) antibiotic classes. Node size reflects the number of stressors within each antibiotic class. Edge thickness reflects the number of convergent or divergent evolution pairs (|ρ| > 0.2). **c,** Convergent and divergent resistance evolution occurred between stressors across and within classes. Convergent: red; divergent: blue. *P* value: chi-square

test significance. **d,** Frequency of convergent evolution (strong, $\rho > 0.6$, red) and divergent evolution ($\rho < -0.2$, blue) relationships among antibiotic classes. Dot size reflects the number of stressor pairs within (diagonal) or across (upper and lower triangles) any two antibiotic classes. **e,** Proportions of convergent and divergent evolution relationships 'across' and 'within' antibiotic classes. **e** is assigned the same color code as **b** for antibiotic classes.

Mutations in evolved populations

For each stress condition, the four evolved cultures were pooled for deep resequencing (coverage: 1,642 ± 466, n = 97, including one control group without antibiotic exposure). A total of 56,744 mutation events were detected by filtering the mutations in the control group, averaging 591 mutations per condition (**Supplementary Fig. 8** and **Supplementary Data 3**), causing 4,750 amino acid changes (AACs) involved in 2,432 protein-coding genes (**Fig. 3a**). These mutations were classified into single nucleotide polymorphisms (non-synonymous SNPs, 3,179), insertions (379), and deletions (1,428). Here, we confirmed the consistency of convergent phenotypes with similar mutational backgrounds (**Fig. 3b**). Additionally, all antibiotic stressors selected more non-synonymous than synonymous substitutions (dN/dS > 1, positive selection), and there were negative correlations between dN/dS and the number of point mutations (**Fig. 3c**). Overall, DSIs induced more point mutations than other stressors at the same dN/dS level, consistent with their mechanisms of action.

The mutation counts of the 2,432 genes exhibit a power-law distribution with a long tail, where the top ten mutant genes account for 6% of the total mutations (**Fig. 3d**). In order to find mutation hotspots along the chromosomal DNA, we performed mutation enrichment analysis in 5-kb sliding windows. The top three mutation hotspots were flagged for the seven classes of antibiotic stressors (**Fig. 3e**). Some sites with frequent mutations included, e.g., the acid stress response gene *evgS* and the multidrug efflux pump gene *emrY*. Interestingly, *kdpD* mutation frequency was relatively high in both macrolides and aminoglycosides-stressed populations. Recent studies report that KdpD promotes resistance to osmotic, oxidative and antimicrobial stresses[24]. *yeaY* with frequent mutation under β-lactam stressors encodes a putative outer membrane lipoprotein[25]. *rcsC* flagged under LPS stress is a known ARG that exhibited increasing expression in response to

polymyxin B. Polymyxin E can activate the RcsC/YojN/RcsB two-component

system, responding to alterations in the bacterial membrane[26].

Unsupervised clustering UMAP embedding was used to test the hypothesis

that antibiotics with the same mechanism of action would enrich mutations leading

to similar AACs (**Fig. 3f**). The hypothesis was largely verified with one exception-

tetracyclines (PSI), which enriched AACs similar to those enriched under

sulfonamides and quinolones stresses (DSI) rather than aminoglycosides and

macrolides (PSI). This aligned well with the observed antibiotic resistance

phenotypes stressed by these antibiotics. Tetracyclines could likely induce DNA

damage due to their strong binding affinity to DNA and their ability to generate free

radicals in close proximity to the DNA backbone[27,28]. Among the 2,718 highly

variable AACs, 991 were enriched by at least one of the four action mechanisms

of stressors (Fisher's exact test $P < 0.05$). DSI antibiotics contributed the most to

the overall enrichment (722 hits), and the majority of overlapping hits were

observed between PSI and CWSI antibiotics (**Fig. 3g**).

We next used Gene Ontology (GO) enrichment analysis to understand the co-

occurrence of mutant gene-associated biological processes for the seven classes

of antibiotics (**Fig. 3h**). 'DNA metabolic process' was represented among the top

hits by sulfonamides and quinolones (DSI), and 'outer membrane' was the most

targeted gene category by LPS stressors (MSI). Meanwhile, tetracyclines caused

the most mutant genes in the 'transporter' category, including but not limited to

those related to carbohydrate transport and metal ion transport. Methylation of the

16S rRNA is an emerging mechanism of high-level resistance to PSI antibiotics via

ribosomal methyltransferases[29,30]. Mutant genes associated with this process were

significantly more prevalent in the presence of macrolides and aminoglycosides

compared with other antibiotics.

Fig. 3: Mutations detected in evolved populations. a, Distribution of mutation events. SNPs are classified into non-synonymous (red) and synonymous (grey) ones. **b,** Mutation similarity in evolved populations exposed to stressors exhibiting convergent or divergent evolutionary relationships. Box plots: center line, median; '+', mean; limits, first and third quartiles; upper and lower whiskers, the highest and lowest values within 1.5× the interquartile range from the hinge. Mutation similarity levels are significantly different for evolved populations exhibiting convergent, other, and divergent relationships (F = 44.59, $P < 0.0001$; one-way ANOVA,

Tukey's honestly significant difference). **c,** Relationships between non-

synonymous to synonymous mutation ratio (dN/dS) and the number of point

mutations. Dashed curves represent non-linear regressions performed for the four

groups of stressors (R^2 = 0.59-0.99, all P < 0.05). **d,** The distribution of mutation

counts for a total of 2,432 mutant genes. The top ten mutant genes harboring the

most mutations are displayed. **e,** Mutation hotspots in the chromosomes of evolved

populations under different antibiotic stresses. The top three mutation hotspots are

highlighted for each antibiotic class. **f,** UMAP clustering of highly variable AACs (n

= 2,718) occurred under the stresses of seven antibiotic classes, which are further

grouped based on the action mechanisms of antibiotics (four groups, rectangle

frames). **g,** UMAP clustering of highly variable AACs among the four groups of

stressors. **h,** GO biological processes associated with mutant genes for different

classes of antibiotic stresses.

Mutant resistance-relevant genes and co-fitness interaction network

Recognizing that not all mutant genes directly contribute to high-level resistance, we employed Tn-Seq to determine gene essentiality and identify potentially resistance-relevant genes (PRGs) across the whole genome. Successful construction of the Tn library using the WT strain enabled high-confidence essentiality and fitness quantification in the presence and absence of eight antibiotics used in MIC assays (**Supplementary Fig. 9**). Overall, the WT strain genome contained 468 essential, 3,910 non-essential, and 367 uncertain genes (**Fig. 4a**). Genes with absolute fitness change $|\Delta W| > 0.15$ for at least one antibiotic were considered as PRGs (n = 2,830, **Fig. 4b** and **Supplementary Data 4**). Their proportions increased from 55% across the whole genome to approximately 73% among mutant genes (**Fig. 4a**). This highlighted that PRGs were indeed selected during experiment evolution under antibiotic stresses.

To map the co-fitness interactions of pairwise PRGs, a 2,830 × 2,830 correlation matrix was calculated based on their conditional fitness profiles. A total of 22,811 co-fitness interactions were identified for 2,152 PRGs (Pearson's $|r| >$ 0.75, corrected $P < 0.00002$), including 15,403 positive and 7,408 negative interactions that were used for constructing the high-confident co-fitness interaction network (**Fig. 4c**). Highly connected hub PRGs (Maximal Clique Centrality, MCC score > 1e20) were identified, forming two central regions in the interaction network. Spatial analysis of functional enrichment (SAFE) further identified 15 network regions spatially enriched for similar bioprocesses, encompassing 1,093 PRGs in total (high-enrichment PRGs, **Fig. 4d** and **Supplementary Data 5**). In addition to some well-studied resistance-associated bioprocesses such as DNA replication and repair, response to stimulus, adhesion, and metabolism, the SAFE analysis also uncovered the enrichment of bioprocesses previously unreported (e.g., carbon/nitrogen/sulfur utilization) or less

known (e.g., metabolism or cell motility[31,32]) to be associated with antibiotic resistance. How these cellular bioprocesses influence resistance evolution under low antibiotic stresses warrants further investigation.

Mutant PRGs ($n = 1,572$) that potentially contribute to the high-level antibiotic resistance of evolved populations, i.e., core mutant PRGs ($n = 131$, $P < 0.01$, Fisher's exact test), were identified by comparing those enriched in populations exhibiting the top 10 versus bottom 40 resistance increase (**Supplementary Fig. 10**). The majority of core mutant PRGs were linked to high-level resistance against TOB, CAZ, MER, and PMB (**Fig. 4e, 4f,** and **Supplementary Data 6**). A representative subset comprising 22 out of the 131 core mutant PRGs was selected for more detailed investigation on their mechanisms of resistance contribution, including 9 hub genes identified in **Fig. 4c**. Some of these representative core mutant PRGs participate in bioprocesses that were spatially enriched by SAFE analysis (high enrichment, **Fig. 4g**), such as response to stimulus (e.g., *ynfB*[33]), signaling (*focC*[34]), and carbohydrate transport (*ycjV*[35]). While other core mutant PRGs contribute to bioprocesses that were not spatially enriched (low-enrichment), such as transcriptional reprogramming genes (e.g., *rtcR* and *narL*), they may still play roles in the development of AMR. This is likely because compared with high-enrichment PRGs, the low-enrichment ones exhibit higher degrees of co-fitness interactions (**Supplementary Fig. 11**), indicating their potential to interact with broader PRGs and thereby facilitate AMR development.

Fig. 4: Gene co-fitness interaction network and spatial analysis of functional enrichment. **a,** Number of essential, non-essential, uncertain genes in the whole genome and mutant genes. ****: chi-square test $P < 1e–4$. **b,** Fitness differences (ΔW) of all Tn-Seq library genes for the 8 tested antibiotics. **c,** Gene interaction (GI) network of PRGs (n = 2,152) based on their co-fitness matrix. **d,** Enriched 15 bioprocesses by SAFE analysis on the PRG GI network in **c**. **e,** Core mutant PRGs conferring high-level resistance to the eight antibiotics (nodes, n = 131). Node size

reflects the co-occurrence frequency of core mutant PRGs. **f,** Hierarchical

clustering of the eight antibiotics based on their associated core mutant PRGs. **g,**

Mutant PRGs (blue) and core mutant PRGs (yellow) are highlighted on a

schematic representation of the GI network. The 22 labelled representative core

mutant PRGs are associated with SAFE high-enrichment bioprocesses (within the

circle, P < 0.05, corrected for multiple testing) or low-enrichment bioprocesses

(outside the circle).

Resistance mechanisms of representative core mutant PRGs

Among the 22 representative core mutant PRGs, 16 have reported links to specific resistance mechanisms, including biofilm formation (e.g., *focC* and *pdeR*), efflux pump (e.g., *ycjV*), drug inactivation (e.g., *MJ1187*), target bypass (e.g., *yafK*), etc. We are therefore interested in uncovering the contribution mechanisms of the remaining 6 representative core mutant PRGs with unreported links to resistance (i.e., *metH, ynfB, ulaG, ftsK, rtcR,* and *hflK*) by constructing their gene interaction modules (module I-VI, respectively, **Fig. 5a**). Within their modules, these PRGs interacted with genes encoding biofilm formation, metabolism, efflux pump, target alteration, membrane permeability, and DNA replication & repair-related proteins. On the one hand, these connections indicated that the 6 PRGs likely contribute to AMR development through similar mechanisms. On the other hand, analyzing the fitness differences (ΔW) of PRG-connected genes in response to antibiotic exposure further refined the proposed resistance mechanisms (**Fig. 5b**). For example, the fimbrial adhesin protein-coding gene *faeJ* and biofilm formation associated gene *vasD*, which interacted with *rtcR* (module V), exhibited significant fitness differences ($|\Delta W| = 0.30 \pm 0.05$ under SMZ+TMP stress). Similarly, the interactions of *hflK* (module VI) with the fimbrial adhesin protein-coding genes *yehB*, *afaA*, and biofilm formation associated gene *suhB* ($|\Delta W| = 0.35 \pm 0.19$ in response to TOB, MER, and PMB stresses) suggested that *hflK* may also contribute to AMR development through affecting biofilm formation.

The proposed resistance mechanisms of *rtcR* and *hflK* were further validated via gene knockout experiments. The two knockout strains exhibited increased susceptibility to their respective antibiotics (**Fig. 5c**), prolonged lag phases in growth ($P < 0.05$, Student's *t*-test; **Fig. 5d**), and decreased biofilm formation (approximately 72% lower than that of the WT strain, **Fig. 5e**). These results collectively suggest that $rtcR^M$ and $hflK^M$ may confer resistance by enhancing

371 biofilm formation, thereby limiting antibiotics diffusion into the population. The
372 resistance mechanisms of the remaining four representative core mutant PRGs
373 (*metH, ynfB*, *ulaG,* and *ftsK)* could be proposed and validated using similar
374 approaches in future studies.

Fig. 5: Gene co-fitness interaction modules of selected core mutant PRGs and
verification of their proposed resistance mechanism. **a,** Core mutant PRGs
displayed within modules extracted from the gene co-fitness interaction network

(in bold, with unknown resistance mechanism) or outside modules (with known

resistance mechanisms). **b,** Fitness differences of genes in modules V and VI

under specific antibiotic exposures. Grey regions represent abs(ΔW) < 0.15. **c,**

Dose–response curves for the three strains exposed to antibiotics at varied

concentrations. Error bars represent standard deviations from biological triplicate.

d, Decreased resistance of $\Delta rtcR$ and $\Delta hflK$ strains to SMZ+TMP and TOB,

respectively, compared with the WT strain. **e,** Biofilm formation potentials of the

WT, $\Delta rtcR$, and $\Delta hflK$ strains determined by the crystal violet assay. N = 6 for all

strains. ****: P < 0.0001.

Resistance evolution prediction by machine-learning models

A long-standing challenge in AMR research has been to predict the outcome of resistance evolution based on the chemical structure and/or dose of stress compounds. The 96 antibiotics used in this study are adequately representative of the structure diversity of commonly applied antibiotics. We developed machine learning models that can correlate the chemical structures of antibiotics, represented by extended-connectivity fingerprints (ECFPs), with the antibiotic resistance profiles of evolved strains. Automated generations by genetic algorithms were employed for feature selection and model optimization. The ensembled regression models demonstrated varied modeling performance on the training datasets, with coefficient of determination (R^2) values ranging from approximately 0.3 to 0.8 for the eight antibiotics (**Fig. 6a**). The modeling performance on CIP resistance (R^2 = 0.79 ± 0.01) was better than those for other antibiotics (F = 31.48, P < 1e–4, two-way ANOVA). We then used CIP resistance prediction results as an example to understand the contributions of monomer structures to antibiotic resistance. The top five antibiotics leading to high-level CIP resistance changes ($\log_2$ fold-change from 5.16 to 7.32) were annotated for their atomic groups' contributions to the CIP resistance (**Supplementary Fig. 12)**. The key contributors to CIP resistance changes were the fluorine atom introduced at the C-6 position and the cyclopropyl group introduced at the N-1 position of the parent nucleus.

To extend the applicability of the machine learning models, we established the applicability domain based on the Tanimoto similarity network. Among the additional 1,563 antibiotics selected from PubChem, 569 (36.4%) with a Tanimoto similarity >0.2 compared to the 96 antibiotics used in this study were considered applicable (**Fig. 6b** and **Supplementary Data 7**). The best-performing models were used to predict the antibiotic resistance profiles stressed by these applicable

antibiotics, resulting in resistance fold changes ($\log_2$ transformed) from −3.32 to 5.58 (**Fig. 6c** and **Supplementary Data 8**). To further evaluate the predictive accuracy of the models, we conducted experimental evolution using 16 external antibiotics (within the applicability domain and encompassing eight antibiotic classes, **Supplementary Table 1**). The predicted resistance levels to CIP and CAZ exhibited agreement with their actual values ($R^2_{test} > 0.75$, **Supplementary Fig. 13**). These chemical structure-based models demonstrate promising potential for predicting antibiotic evolution under low-level antibiotic stresses.

In addition to ECFP modeling, we also explored mutation-based antibiotic resistance prediction. The AAC model was constructed based on the random forest (RF) algorithm, which indeed improved the prediction performance (R^2_{AAC}: 0.73 ± 0.05 vs. R^2_{ECFP}: 0.51 ± 0.15 for the training dataset, **Fig. 6a** and **Supplementary Fig. 14**). Despite proof-of-concept demonstrations, the applications of AAC-based modeling in predicting resistance phenotypes remain to be validated due to the complexity in intrinsic and extrinsic factors influencing mutation outcomes, e.g., genetic background and nutrient availability.

Fig. 6: Performance of machine learning models and prevalence of key point mutations in clinical/environmental isolates. **a,** Performance of the machine learning models. The error bars represent the standard error of mean from the top three best-performing models. **b,** Tanimoto similarity network demonstrating the applicability domain (Tanimoto similarity ≥ 0.2) of ECFP-based models. **c,** Split violin plots showing the normalized distributions of predicted resistance levels for the training (n = 96) and predicting antibiotics (n = 569). **d,** Top 25 most important

SNPs (features) contributing to the accuracy of the AAC-based RF model for each antibiotic. Dot size: importance of features; Shared counts: occurrence of particular SNPs in the circle. **e,** Occurrence of key SNPs (feature-importance SNPs in **d** and SNPs in core mutant PRGs conferring high-level resistance, n = 481 in total) in NCBI-retrieved clinical and environmental isolates. **f,** Left: key SNPs associated with resistance to the eight antibiotics. Right: number of clinical and environmental isolates containing the top 40 key SNPs. Total isolates: 1,263 retrieved from the NCBI Pathogen Detection database. Fisher's exact test P values indicate the overrepresentation of SNPs in clinical isolates and are colored in varying degrees of purple, from highly to not significant (dark to light purple, respectively).

Clinical and environmental prevalence of resistance-relevant mutations

Assessing the prevalence of key mutations in a broader range of clinical or environmental isolates is crucial for evaluating their risks in the 'One Health' paradigm. We focus on two groups of point mutations, comprising a total of 481 key SNPs, i.e., feature-importance SNPs contributing to the accuracy of the AAC-based RF models (**Fig. 6d**) and SNPs conferring high-level antibiotic resistance (**Supplementary Fig. 10**). These key SNPs and 1,263 *E. coli* genomes (clinical: 577, environmental: 686) retrieved from the NCBI Pathogen Detection database were subjected to comparative genomic analysis (**Supplementary Data 9 and 10**). Overall, 70 of the 481 key SNPs (14.6%) were identified in at least one of the external genomes (**Fig. 6e**), with the highest occurrence in 487 genomes (*yddB* Arg458His, **Fig. 6f**). Their relative prevalence in clinical and environmental isolates was summarized in **Supplementary Fig. 15**. Importantly, 62.9% of the key SNPs were identified in clinical isolates, and three were significantly more prevalent in clinical than environmental isolates, e.g., TetR family of transcription regulator gene *rutR* Ala15Thr and lipoprotein-related gene *bor* Ala66Thr. Furthermore, 74.3% of the key SNPs were identified in environmental isolates, and three were significantly more prevalent, e.g., oxidative stress response gene *btuE* Ala54Val. Notably, co-occurring SNPs (i.e., detected in both clinical and environmental isolates) accounted for 37.1% of the total key SNPs. These results suggested that mutations that significantly contribute to antibiotic resistance are widespread in both clinical and environmental isolates, which may be acquired under sub-MIC antibiotic exposure conditions.

Discussion

Despite the growing attention that the development of AMR could be driven by the low-level selective pressure of antibiotics released into the environment, the relationship between AMR phenotypes and genotypes, and whether mutations conferring high-level resistance are prevalent remain key questions to be addressed. In this study, we constructed an unprecedented phenotype-genotype atlas of resistance evolution under 96 low-level antibiotic stress conditions. We used high-throughput experimental evolution and deep-resequencing approaches to determine, at the population level, how bacteria can evolve high-level resistance upon exposure to a wide range of antibiotics at low levels. Tn-Seq further facilitated the identification of PRGs and the elucidation of associated resistance mechanisms. ECFP- and AAC-based machine learning models have shown promise in areas such as predicting resistance phenotypes and identifying feature-importance antibiotic substructures and mutations. We leveraged an extensive collection of publicly available genomes to determine the prevalence of key mutations in a broader range of environmental and clinical settings. Some of the main findings extend beyond our understanding of resistance evolution under low-level antibiotic exposures, shedding insights into the origins and dissemination of antibiotic resistance in the environment.

From a phenotypic perspective, most stressors showed positive correlations in causing consistent antibiotic resistance changes in evolved populations. Cross-resistance was much more prevalent than collateral sensitivity[16], reinforcing and complementing the previous findings of cross-resistance and collateral sensitivity relationships identified at clinical concentrations[36,37]. Some of the newly identified collateral sensitivity relationships (e.g., kanamycin or gentamicin vs. sparfloxacin, and aztreonam vs. ciprofloxacin; $P < 0.05$) offer valuable insights for developing alternative clinical antibiotic dosing strategies. The study is notable for pioneering

the establishment of a link between stressor's chemical structures and resistance evolution. For instance, quinolone stressors elicited the highest CIP resistance compared with other antibiotics, and their particular atomic groups, e.g., the C-6 fluorine atom and the cyclopropyl group, were found likely to contribute to high-level resistance evolution.

From a genotypic perspective, the results provide a comprehensive understanding of how critical mutations contribute to clinically relevant resistance and the evolutionary landscape. Frequently encountered ARGs such as *evgS* and *ermY* were among the most common targets of mutations under antibiotic stresses. Tn-Seq further enabled the construction of the GI network in response to antibiotics. We, therefore, identified a set of core mutant PRGs (n = 131) contributing to high-level resistance evolution through the phenotype-genotype atlas. In addition to the well-characterized ARGs, e.g., *pdeR* involved in biofilm formation[38], *ycjV* involved in efflux pump[35,39], *MJ1187* involved in drug inactivation[40], *yafK* involved in target bypass[41,42], and yjiR involved in transcriptional reprogramming[43], others may confer high-level resistance via unreported mechanisms, e.g., *metH*, *ynfB*, *ulaG*, *ftsK*, *rtcR*, and *hflK*. By locating these understudied core mutant PRGs in the gene co-fitness interaction network and analyzing the functions of interacted genes, their resistance mechanisms were proposed. The resistance contribution mechanism of two core mutant PRGs, namely *rtcR* and *hflK*, through biofilm formation, was experimentally confirmed. Importantly, this demonstrated that the gene co-fitness network generated based on Tn-Seq data highlights valuable interactions that can be explored to uncover resistance mechanisms of potential ARGs. The SAFE analysis further suggests that genome-wide mutations affecting broad biological processes[44,45], such as metabolism and cell motility, can significantly contribute to the evolution of antibiotic resistance[31,32], extending beyond independent, target-gene-related mutations.

The ECFP- and AAC-based machine learning models exhibited promising levels of accuracy and scalability in elucidating the intricate relationships between resistance phenotype, genotype, and underlying drivers (i.e., antibiotics), leveraging the power of vast data that are unattainable through traditional low-throughput evolutionary experiments. Feature-importance antibiotic substructures identified may have important implications for the development of next-generation antibiotics and the targeted pollutant removal from contaminated environments[46]. The ECFP-based models exhibit broad applicability across a wide range of antibiotics, enabling relatively reliable prediction of antibiotic resistance in *E. coli* strains under low-level antibiotic exposures without the need for intensive, high-throughput experimental evolution. Moreover, the predicted resistance levels can be used as a reference, to some extent, for assessing the risks of environmentally acquired resistance.

Key SNPs selected at sub-MICs indeed exhibited both environmental and clinical relevance across a vast collection of *E. coli* isolates, and their environment-clinic co-occurrence rates were up to 37.1%. Some of these key SNPs were even enriched in clinical isolates, suggesting that the variants may already pose health risks. Consequently, antibiotic-contaminated aquatic and soil environments could serve as reservoirs of evolved resistant bacteria carrying clinically relevant, high-resistance-conferring mutations. A few studies have attempted to determine the genetic relationships between environmental and clinical isolates in public databases to infer transmission dynamics[47,48]. Our results made additional contributions to the identification of key mutations and ARGs of concern, as well as the quantification of their prevalence.

Disentangling the AMR risk is challenging and requires a holistic understanding of environment-clinic AMR dissemination, including but not limited to its evolution under low-level antibiotic exposures. Some considerations should

be taken into account when extending our results to other studies on environmental resistance. First, environmental pathogens are remarkably diverse, likely resulting in species-specific response and resistance evolution patterns that differ from those of *E. coli*. Second, environmental changes are far more drastic and dynamic in nature, and resistance evolution is an ongoing process. Efforts to predict resistance profiles, either based on chemical structure or mutations, are inevitably constrained by temporal limitations and complex interplay of influencing factors. Third, uncovering the resistance-conferring mechanisms of core mutant PRGs is based on a hypothesis-verification approach. However, even if the hypotheses are experimentally validated, they may not be the primary mechanisms conferring resistance due to the pleiotropy of genes. We are actively investigating the extent to which these lab-scale findings from experimental evolution mirror AMR development in real-world settings, particularly antibiotic-contaminated environments. Nonetheless, these efforts provide valuable scientific support for combating antibiotic resistance within the One Health framework.

Methods

Bacterial strains, culture conditions, and antibiotics

E. coli strain QSHQ1 (GenBank accession number: CP128989) was a plasmid-free opportunistic pathogen that was originally isolated from soil in Hangzhou, China (**Supplementary Fig. 16**). Cells were cultivated with LB medium (150 rpm, 37°C) in Abgene™ 96-well 2.2 ml polypropylene deep-well plates (Thermo Scientific, #AB0661) with breathable sealing film (Corning, #1223S04). All antibiotics used were purchased from Yuanye-Bio, Macklin, and Solarbio (details in **Supplementary Data 1**). Antibiotic stock solutions (dissolved in 10% sterile DMSO) were filter-sterilized and stored at −20°C before use.

Experimental evolution under 96 antibiotic stresses

Pre-adaptation: A single colony of *E. coli* strain QSHQ1 was picked from the LB-agar plate and grown in LB medium without antibiotics for 24 h. Cells were sub-cultured daily by 500-fold dilutions in fresh LB medium over a 10-day period, corresponding to approximately 9 generations per day.

Experimental evolution: 96 antibiotics were added to the LB-medium at a uniform concentration of 10 μg l^{-1}. Four independent cultures (biological quadruplicate) were grown under each antibiotic stress and were sub-cultured daily by 500-fold dilutions for 20 days (1 ml medium, 96-well plates, 150 rpm, 37°C). The antibiotic concentration was selected to reflect the upper range of environmentally relevant concentrations detected in, e.g., wastewater treatment plants[49]. Four control lineages were cultured under the same condition without antibiotics. Cell growth was monitored by measuring OD$_{600}$ using the Spark microplate reader (Tecan Inc.). After the 20-day experimental evolution, end-point cultures were frozen at −80°C in 30% glycerol for whole-genome sequencing. No growth was observed in the negative control wells throughout the experimental

599   evolution.

Automated high-throughput MIC assays of evolved populations

601   The automated high-throughput MIC assays were carried out on the iBioFoundry platform at ZJU-Hangzhou Global Scientific and Technological Innovation Center. The platform consisted of a MultiDrop Combi microplate dispenser (Thermo Scientific, #5840300), a Tecan Freedom Evo 200 Base laboratory automation workstation, a static incubator, and a CLARIOstar Plus microplate reader (BMG LABTECH Inc.), which were operated by two six-axis robotic arms (Thermo Scientific, TS-F7000).

608   Antibiotic resistance against azithromycin (AZI), ciprofloxacin (CIP), tetracycline (TET), sulfamethoxazole with trimethoprim (SMZ+TMP), tobramycin (TOB), ceftazidime (CAZ), meropenem (MER), and polymyxin B (PMB) were measured for evolved cultures at the end of the experimental evolution. The end-point glycerol-frozen cultures were revived in 1 ml LB medium for 16 h at 37°C with (evolved populations) or without (control) stressing antibiotics. The revived cultures were 2000-fold diluted in freshly prepared MIC plates to a final volume of 130 μl. The 2-fold serial dilutions were performed over 12 steps, starting from the highest concentrations of 400, 25, 100, 200, 100, 200, 25, and 400 mg l^{-1}, respectively, for the eight antibiotics. After 20 h static incubation, the OD_{600} values were measured. The dose-response curves generated from biological quadruplicates were fitted to the Hill equation, and the antibiotic concentration corresponding to an OD_{600} of 0.2 was defined as IC_{90}. In the event of curve overflow (exceeding 100% inhibition) or underfill (below 90% inhibition), the IC_{90} values were assigned as the maximum tested concentration or half of the minimum tested concentration, respectively.

DNA extraction from evolved populations

In the antibiotic resistance assays, iBioFoundry identified the first well with

detectable cell growth (OD_{600} > 0.2) in the presence of antibiotics, and pooled cells

from up to 32 of these wells together (8 antibiotics × 4 replicates, ~4 ml). Genomic

DNA (gDNA) was extracted from pooled cultures using the DNeasy Blood & Tissue

kit (Qiagen) in accordance with the manufacturer's instructions. DNA quantity and

quality were determined by the Colibri microvolume spectrometer (Titertek-

Berthold Inc., Germany). High-quality DNA (A_{260}/A_{280} = 1.8, > 6 μg) was used for

whole-genome sequencing.

Whole-genome sequencing and mutation analysis

Whole-genome sequencing was performed for the WT strain *E. coli* QSHQ1

as well as the 96 evolved populations, following the procedures described

previously[10]. Briefly, the WT strain was sequenced using PacBio RS combined with

Illumina sequencing (NovaSeq6000). The Illumina data were used to evaluate the

complexity of the genomes and correct errors in PacBio long reads. The high-

quality complete genome of the WT strain (5,084,285 bp) served as the reference

for mutation detection (**Supplementary Fig. 17**). Pooled evolved populations were

deep resequenced on the Illumina platform (150 bp×2, ~1,500x coverage,

Shanghai BIOZERON Co., Ltd). Mutations detected in the control lineages were

excluded from further analysis. Variants with a frequency greater than 0.2% in the

population were identified. The non-synonymous (d*N*) and synonymous (d*S*)

substitutions were obtained under each stress condition, and the proportion of

beneficial mutations was indicated by the *dN/dS* ratio, which was set to 1 for neutral

evolution (without antibiotic stress).

Transposon sequencing and gene fitness assays

A transposon mutant library for the WT strain was generated by

electroporation using the EZ-Tn5 <KAN-2> Tnp transposome kit (Lucigen Inc.).

The library consisted of approximately 40,000 insertion mutants, achieving a

genome-wide saturation of 97.5%. The library was divided into 9 aliquots, 500x

diluted in LB medium and grew for 24 h (4-8 generations) with no antibiotic or one

of the eight antibiotics at sub-MIC levels. The gDNA was extracted, and a specific

PCR forward primer (GATCCTCTAGAGTCGACCTGCAGG) was designed to

amplify the regions flanking the Tn5 insertion (TA) sites. TA sites were sequenced

on Illumina NovaSeq6000, and the sequencing data were analyzed using the

TRANSIT package v3.2.7[50]. Essential genes were identified based on the Gumbel

method by calling long consecutive sequences of TA sites without insertions. For

non-essential genes with valid TA sites > 5 and mean insertion counts (N) > 5,

roughly 3,500 in total, their gene fitness W was calculated as follows:

670 $$W_i = ln(N_i(t_2)/N_i(t_1))$$

where $N_i(t_1)$ and $N_i(t_2)$ were the mean insertion counts of non-essential gene i

detected in a particular library before and after antibiotic selection, respectively.

The differences in the gene fitness were therefore calculated as $\Delta W = W_{ABX} -$

W_{noABX}. Genes with absolute fitness difference values $|\Delta W|$ larger than 0.15 under

at least one antibiotic stress condition were identified as potentially resistance-

relevant genes (PRGs).

Gene co-fitness interaction network and spatial analysis of functional

enrichment (SAFE)

A co-fitness interaction network for PRGs was constructed as described

previously[51]. In brief, a gene fitness matrix was built for a total of 2,830 PRGs,

which was further used to calculate Pearson's correlation coefficients (r) for all

pairwise genes. A high-confidence co-fitness interaction network was established

using Cytoscape, comprising 2,152 PRGs (nodes) and 22,811 interactions (edges)

with $|r|$ > 0.75 and corrected P < 0.00002 (threshold: 0.05, corrected by the number

of genes). The plugin CytoHubba was used to calculate the degree and maximal clique centrality (MCC) of all nodes in the network. The plugin ClusterViz was employed to identify distinct modules based on network topology using the FAG-EC algorithm. PRGs were annotated using the Swiss-Prot, GO, CARD, VFDB, and BacMet databases. SAFE highlighted regions of the gene interaction network enriched in specific functional attributes.

Knockout of selected PRGs and biofilm formation potential measurement

Two core mutant PRGs, *rtcR* and *hflK* were subjected to knockout experiments to verify their hypothesized involvement in biofilm formation. The single guide RNAs (sgRNAs) were designed using the online design tool (https://en.rc-crispr.com/). The pair of oligos for the two target genes were annealed and ligated to the CB-001 vector (Ubigene Biosciences Co., Ltd., Guangzhou, China). The CB-001-*rtcR* (or *hflK*) [gRNA] plasmids containing each target sgRNA sequence and donor were transformed into the WT strain by electroporation. Cells were spread onto LB agar and incubated overnight. Transformants were identified by colony PCR and Sanger sequencing. The sequences of sgRNAs and primers for CRISPR were provided in **Supplementary Tables 2** and **3**, respectively. Growth curves of the WT and knockout strains were obtained without antibiotics and with two low-level antibiotics (SMZ+TMP: 1 µg/ml, TOB: 5 µg/ml). The biofilm formation potentials of WT and knockout strains were determined based on the microtiter dish assay. In brief, strains were overnight grown in LB medium at 150 rpm and 37°C, and then diluted to OD_{600} = 0.05. The 96-well plates were statically incubated at 37°C for 24 h to allow biofilm formation, and the ultimate OD_{600} was determined. For staining, 200 µl of 0.1% crystal violet (CV) was added to each well and incubated for 15 min at 37°C. The wells were then washed three times with water and dried overnight. The stained biofilm was

disrupted by adding 200 µl of 95% ethanol solution (95% EtOH, 4.95% distilled

H_2O, 0.05% Triton X-100), and OD_{550} was measured afterward. Biofilm formation

potential was calculated as OD_{550}/OD_{600}.

Machine learning methods

To elucidate the relationships between the chemical structures of antibiotics

and the antibiotic resistance profiles they induced, quantitative structure-property

relationship (QSPR) models were constructed based on a published study[52]. In

brief, molecular fingerprints, including MACCS 166 and ECFPs and about 6,000

descriptors, were computed for the 96 stress-inducing antibiotics, which can be

fine-tuned with a set of features. The antibiotic resistance profiles of evolved

strains were used to train a number of ensemble regression models, including

linear regression (OLS), partial least squares (PLS), k-nearest neighbors (KNN),

and support vector machines (SVM). Automatic model generation (n = 10,000

iterations) by genetic algorithms was performed to find the best combination of

features. The performance of a model was evaluated by using statistical

parameters, i.e., in-sample coefficients of determination R^2, RMSE, and out-of-

sample random 10-fold cross-validation R^2. The visualization strategy for tracing

the atomic contribution was previously described[53,54]. To assess the applicability

domain of the best-performing models, the structural data of 1,563 antibiotics,

excluding the 96 used in experimental evolution, were downloaded from PubChem

(https://pubchem.ncbi.nlm.nih.gov/#query=antibiotic).   A   pairwise   Tanimoto

similarity matrix was computed for the 1,659 antibiotics in total. An antibiotic was

considered as inside domain if its Tanimoto similarity with at least one of the 96

antibiotics exceeded 0.2. The best-performing models were employed to predict

antibiotic resistance development under stress conditions by the in-domain

antibiotics. 16 out of the 569 in-domain antibiotics were randomly selected to

evaluate model prediction performance, following the same procedures used in experimental evolution.

In addition to molecular fingerprint-based modeling, the relationships between 4,750 AACs identified in evolved strains and the resulting antibiotic resistance profiles were also explored using RF models. In brief, the Boole matrix was constructed based on the relative fold-change values and AACs for the 96 antibiotic stressors. The RF models employed 100 individual decision trees, each with a variable range of splits from 10 to 2,000 and a minimum split size of 5 per tree. Bootstrapping was employed to mitigate the inherent overfitting of models, where a 10-fold cross-validation (Shuffle Split) method was used for each machine-learning algorithm.

Prevalence of point mutations based on comparative genomics

Comparative genomic analysis was performed using a well-established pipeline as described previously[31]. Briefly, all *E. coli* genomes with L50 < 20 were downloaded from the NCBI database (https://www.ncbi.nlm.nih.gov/pathogens/) on 3/24/2023, including 'clinical' (n = 3,376) and 'environmental' (n = 2,545) isolates. Genomes were searched for all key SNPs (n = 481) identified in this study. For each of the 5,921 genomes, Blast was used to identify the genomes hit for the QSHQ1 strain. If the BLAST e-value was 1e–10 or less, and the query genome had a coverage of at least 85%, the CDS sequences from hit clinical (n = 577) and environmental (n = 686) genomes were extracted, and multiple alignment was performed using Muscle. Mutated positions in all genomes in the database that aligned with the QSHQ1 SNPs were tabulated. Mutations that did not appear at least once in the entire pathogen database were removed from consideration; P values for over-representation in clinical (or environmental) isolates were calculated using Fisher's exact test for all remaining mutations.

Data availability

The raw sequence data of Reseq and Tn-Seq analyses are available in the NCBI Sequence Read Archive under the accession number PRJNA987173. All relevant data and models in this study are available from the corresponding authors upon reasonable request. Source data are provided with this paper.

References

1. Programme, U. N. E. Frontiers 2017: Emerging issues of environmental concern. Preprint at (2017).

2. Larsson, D. G. J. & Flach, C.-F. Antibiotic resistance in the environment. *Nat. Rev. Microbiol.* **20**, 1–13 (2021).

3. Martínez, J. L. Antibiotics and antibiotic resistance genes in natural environments. *Science* **321**, 365–367 (2008).

4. Allen, H. K. *et al.* Call of the wild: antibiotic resistance genes in natural environments. *Nat. Rev. Microbiol.* **8**, 251–259 (2010).

5. FAO, UNEP, WHO & OIE), the W. O. for A. H. (WOAH founded as. *One Health Joint Plan of Action (2022–2026): Working together for the health of humans, animals, plants and the environment*. 86 https://www.who.int/publications/i/item/9789240059139 (2022).

6. Polianciuc, S. I., Gurzău, A. E., Kiss, B., Ştefan, M. G. & Loghin, F. Antibiotics in the environment: causes and consequences. *Med. Pharm. Rep.* **93**, 231–240 (2020).

7. Wistrand-Yuen, E. *et al.* Evolution of high-level resistance during low-level antibiotic exposure. *Nat. Commun.* **9**, 1599 (2018).

8. Liang, H., Zhang, J., Hu, J., Li, X. & Li, B. Fluoroquinolone residues in the environment rapidly induce heritable fluoroquinolone resistance in *Escherichia coli*. *Environ. Sci. Technol.* **57**, 4784–4795 (2023).

9. Xing, Y., Wu, S. & Men, Y. Exposure to environmental levels of pesticides stimulates
and diversifies evolution in *Escherichia coli* toward higher antibiotic resistance. *Environ.*
Sci. Technol. **54**, 8770–8778 (2020).

10. Wang, H., Feng, Y. & Lu, H. Low-level cefepime exposure induces high-level
resistance in environmental bacteria: molecular mechanism and evolutionary dynamics.
Environ. Sci. Technol. **56**, 15074–15083 (2022).

11. Jørgensen, K. M. *et al.* Sublethal ciprofloxacin treatment leads to rapid development
of high-level ciprofloxacin resistance during long-term experimental evolution of
Pseudomonas aeruginosa. Antimicrob. Agents Ch. **57**, 4215–21 (2013).

12. Jin, L. *et al.* Integrating environmental dimensions of "One Health" to combat
antimicrobial resistance: essential research needs. *Environ. Sci. Technol.* (2022)
doi:10.1021/acs.est.2c01651.

13. Theuretzbacher, U. *et al.* Critical analysis of antibacterial agents in clinical
development. *Nat. Rev. Microbiol.* **18**, 286–298 (2020).

14. Fait, A. *et al.* Adaptive laboratory evolution and independent component analysis
disentangle complex vancomycin adaptation trajectories. *Proc. National. Acad. Sci.* **119**,
e2118262119 (2022).

15. Remigi, P., Masson-Boivin, C. & Rocha, E. P. C. Experimental evolution as a tool to
investigate natural processes and molecular functions. *Trends Microbiol.* **27**, 623–634
(2019).

16. Maeda, T. *et al.* High-throughput laboratory evolution reveals evolutionary
constraints in *Escherichia coli. Nat. Commun.* **11**, 5970 (2020).

17. Lukačišinová, M., Fernando, B. & Bollenbach, T. Highly parallel lab evolution
reveals that epistasis can curb the evolution of antibiotic resistance. *Nat. Commun.* **11**,
3105 (2020).

18. Palomino, A. *et al.* Metabolic genes on conjugative plasmids are highly prevalent in
Escherichia coli and can protect against antibiotic treatment. *ISME J.* **17**, 151–162
(2023).

19. D'Lima, L. *et al.* No decrease in susceptibility to NVC-422 in multiple-passage
studies with methicillin-resistant *Staphylococcus aureus*, *S. aureus*, *Pseudomonas*
aeruginosa, and *Escherichia coli. Antimicrob. Agents Chemother.* **56**, 2753–2755 (2012).

20. Pollard, J. E. *et al.* In vitro evaluation of the potential for resistance development to
ceragenin CSA-13. *J. Antimicrob. Chemother.* **67**, 2665–2672 (2012).

21. CLSI. Performance standards for antimicrobial susceptibility testing. 33rd ed.
Preprint at (2023).

22. Imamovic, L. *et al.* Drug-driven phenotypic convergence supports rational treatment
strategies of chronic infections. *Cell* **172**, 121-134.e14 (2018).

23. Stracy, M. *et al.* Minimizing treatment-induced emergence of antibiotic resistance in
bacterial infections. *Science* **375**, 889–894 (2022).

24. Alegado, R. A., Chin, C., Monack, D. M. & Tan, M. The two-component sensor
kinase KdpD is required for *Salmonella typhimurium* colonization of *Caenorhabditis*
elegans and survival in macrophages. *Cell. Microbiol.* **13**, 1618–1637 (2011).

25. Rezuchova, B., Miticka, H., Homerova, D., Roberts, M. & Kormanec, J. New
members of the *Escherichia coli* σE regulon identified by a two-plasmid system. *FEMS*
Microbiol. Lett. **225**, 1–7 (2003).

26. Groisman, E. A. & Mouslim, C. Sensing by bacterial regulatory systems in host and
non-host environments. *Nat. Rev. Microbiol.* **4**, 705–709 (2006).

27. Khan, M. A., Mustafa, J. & Musarrat, J. Mechanism of DNA strand breakage induced
by photosensitized tetracycline–Cu(II) complex. *Mutat. Res.Fundam. Mol. Mech.*
Mutagen. **525**, 109–119 (2003).

28. Pushparajan, A. R., Ramachandran, R., Reji, J. G. & Kumar, R. A. *Mycobacterium*
tuberculosis TetR family transcriptional regulator Rv1019 is a negative regulator of the
mfd-mazG operon encoding DNA repair proteins. *FEBS Lett.* **594**, 2867–2880 (2020).

29. Doi, Y., Wachino, J. & Arakawa, Y. Aminoglycoside resistance the emergence of
acquired 16S ribosomal RNA methyltransferases. *Infect. Dis. Clin. North Am.* **30**, 523–
537 (2016).

30. Darby, E. M. *et al.* Molecular mechanisms of antibiotic resistance revisited. *Nat. Rev.*
Microbiol. 1–16 (2022) doi:10.1038/s41579-022-00820-y.

31. Lopatkin, A. J. *et al.* Clinically relevant mutations in core metabolic genes confer
antibiotic resistance. *Science* **371**, eaba0862 (2021).

32. Piskovsky, V. & Oliveira, N. M. Bacterial motility can govern the dynamics of
antibiotic resistance evolution. *Nat. Commun.* **14**, 5584 (2023).

33. Seo, S. W., Kim, D., O'Brien, E. J., Szubin, R. & Palsson, B. O. Decoding genome-
wide GadEWX-transcriptional regulatory networks reveals multifaceted cellular
responses to acid stress in *Escherichia coli*. *Nat. Commun.* **6**, 7970 (2015).

34. Nüesch-Inderbinen, M., Treier, A., Stevens, M. J. A. & Stephan, R. Whole genome
sequence-based characterisation of Shiga toxin-producing *Escherichia coli* isolated from
game meat originating from several European countries. *Sci. Rep.* **13**, 3247 (2023).

35. Moussatova, A., Kandt, C., O'Mara, M. L. & Tieleman, D. P. ATP-binding cassette
transporters in *Escherichia coli*. *Biochimica. Et. Biophysica. Acta. Bba. – Biomembr.*
1778, 1757–1771 (2008).

36. Liu, D. Y. *et al.* Collateral sensitivity profiling in drug-resistant *Escherichia coli*
identifies natural products suppressing cephalosporin resistance. *Nat. Commun.* **14**, 1976
(2023).

37. Imamovic, L. & Sommer, M. O. A. Use of collateral sensitivity networks to design
drug cycling protocols that avoid resistance development. *Sci. Transl. Med.* **5**, 204ra132
(2013).

38. Wang, N., Gao, J., Xiao, S. & Zhuang, G. Overexpression of pdeR promotes biofilm
formation of *Paracoccus denitrificans* by promoting ATP production and iron
acquisition. *Front. Microbiol.* **13**, 966976 (2022).

39. Bevers, L. E., Hagedoorn, P.-L., Krijger, G. C. & Hagen, W. R. Tungsten transport
protein A (WtpA) in *Pyrococcus furiosus* : the first member of a new class of tungstate
and molybdate transporters. *J. Bacteriol.* **188**, 6498–6505 (2006).

40. Bult, C. J. *et al.* Complete genome sequence of the methanogenic archaeon,
Methanococcus jannaschii. *Science* **273**, 1058–1073 (1996).

41. Kuru, E. *et al.* Fluorescent D-amino-acids reveal bi-cellular cell wall modifications
important for *Bdellovibrio bacteriovorus* predation. *Nat. Microbiol.* **2**, 1648–1657 (2017).

42. Voedts, H., Kennedy, S. P., Sezonov, G., Arthur, M. & Hugonnet, J.-E. Genome-wide
identification of genes required for alternative peptidoglycan cross-linking in *Escherichia*
coli revealed unexpected impacts of β-lactams. *Nat. Commun.* **13**, 7962 (2022).

43. Rodionova, I. A. *et al.* A systems approach discovers the role and characteristics of
seven LysR type transcription factors in *Escherichia coli*. *Sci. Rep.* **12**, 7274 (2022).

44. Baryshnikova, A. Systematic functional annotation and visualization of biological
networks. *Cell Syst.* **2**, 412–421 (2016).

45. Costanzo, M. *et al.* A global genetic interaction network maps a wiring diagram of
cellular function. *Science* **353**, aaf1420–aaf1420 (2016).

46. Zhang, L., Liu, H., Dai, J., Xu, P. & Tang, H. Unveiling degradation mechanism of
PAHs by a *Sphingobium* strain from a microbial consortium. *mLife* **1**, 287–302 (2022).

47. Wyres, K. L. & Holt, K. E. *Klebsiella pneumoniae* as a key trafficker of drug
resistance genes from environmental to clinically important bacteria. *Curr. Opin.*
Microbiol. **45**, 131–139 (2018).

48. Pornsukarom, S., Vliet, A. H. M. van & Thakur, S. Whole genome sequencing
analysis of multiple *Salmonella serovars* provides insights into phylogenetic relatedness,
antimicrobial resistance, and virulence markers across humans, food animals and
agriculture environmental sources. *BMC Genom.* **19**, 801 (2018).

49. Geng, J., Liu, X., Wang, J. & Li, S. Accumulation and risk assessment of antibiotics
in edible plants grown in contaminated farmlands: A review. *Sci. Total Environ.* 158616
(2022) doi:10.1016/j.scitotenv.2022.158616.

50. DeJesus, M. A., Ambadipudi, C., Baker, R., Sassetti, C. & Ioerger, T. R. TRANSIT -
a software tool for Himar1 TnSeq analysis. *Plos Comput. Biol.* **11**, e1004401 (2015).

51. Leshchiner, D. *et al.* A genome-wide atlas of antibiotic susceptibility targets and
pathways to tolerance. *Nat. Commun.* **13**, 3165 (2022).

52. Mauri, A. & Bertola, M. Alvascience: A new software suite for the QSAR workflow
applied to the blood–brain barrier permeability. *Int. J. Mol. Sci.* **23**, 12882 (2022).

53. Polishchuk, P. G., Kuz'min, V. E., Artemenko, A. G. & Muratov, E. N. Universal
approach for structural interpretation of QSAR/QSPR models. *Mol. Inform.* **32**, 843–853
(2013).

54. Riniker, S. & Landrum, G. A. Similarity maps - a visualization strategy for molecular
fingerprints and machine-learning methods. *J. Cheminformatics* **5**, 43 (2013).

Acknowledgements

This work was supported by the 'Major Program of National Natural Science Foundation of China (Grant No. 22193061 to Lizhong Zhu)', the 'Zhejiang Provincial Natural Science Foundation of China (Grant No. LR23B070002 to Huijie Lu)', the 'General Program of National Natural Science Foundation of China (Grant No. 22276165 to Huijie Lu)', the 'National Key Research and Development Program (Grant No. 2022YFC3704700 to Huijie Lu)', and ZJU Foundation of Institute of Intelligent Biological and Chemical Manufacturing. The authors thank Dr. Haoxuan Liu from Zhejiang University Centre for Evolutionary & Organismal Biology and Dr. Lei Huang from ZJU-Hangzhou Global Scientific and Technological Innovation Center for valuable discussions.

Author contributions

Conceptualization, H.W. and Huijie.L.; Software, H.W. and Hui.L.; Investigation, H.W. and Huijie.L.; Writing-original draft, H.W.; Writing-review & editing, H.W., Huijie.L., C.J. and L.Z.; Supervisions, Huijie.L. and L.Z. All authors approved the final paper.

Competing interests

The authors declare no competing interests.

Materials & Correspondence

Correspondence and material requests should be addressed to Huijie Lu.