Accounting for observation biases associated with counts of young when estimating fecundity: case study on the arboreal-nesting red kite (*Milvus milvus*)

Sollmann Rahel*[1], Adenot Nathalie[2], Spakovszky Péter[3], Windt Jendrik[2], Mattsson Brady J.[2]

[1] Department of Ecological Dynamics, Leibniz Institute for Zoo and Wildlife Research – Berlin, Germany
[2] Institute of Wildlife Biology and Game Management, Department of Integrative Biology and Biodiversity Research, University of Natural Resources and Life Sciences – Vienna, Austria
[3] TB Raab GmbH – Deutsch-Wagram, Austria

*Corresponding author
Correspondence: sollmann@izw-berlin.de

ABSTRACT

Counting the number of young in a brood from a distance is common practice, for example in tree-nesting birds. These counts can, however, suffer from over and undercounting, which can lead to biased estimates of fecundity (average number of nestlings per brood). Statistical model development to account for observation bias has focused on false negatives (undercounts), yet it has been shown that these models are sensitive to the presence of false positives (overcounts) when they are not accounted for. Here, we develop a model that estimates fecundity while accounting for both false positives and false negatives in brood counts. Its parameters can be estimated using a calibration approach that combines uncertain counts with certain ones, which can be obtained by accessing the brood, for example during ringing. The model uses multinomial distributions to estimate the probabilities of observing y young conditional on the true state of a brood z (i.e., true number of young) from paired uncertain and certain counts. These classification probabilities are then used to estimate the true state of broods for which only uncertain counts are available. We use a simulation study to investigate bias and precision of the model and parameterize the simulation with empirical data from 26 red kite nests visited with ground and nest-based counts during 2021 and 2022 in central Europe. In these data, bias in counts was at most 1 in either direction, more common in larger broods, and undercounting was more common than overcounting. This led to an overall 5% negative bias in fecundity in uncertain counts. The model produced essentially unbiased estimates (relative bias < 2%) of fecundity across a range of sample sizes. This held true whether or not fecundity was the same in the paired data and in the data with uncertain counts only. But the model could not estimate parameters when true states were missing from the paired data, which happened frequently in small sample sizes (n = 10 or 25).

Further, we projected populations 50 years into the future using fecundity estimates
corrected for observation biases from the multinomial model, and based on "raw" uncertain
observations. We found that ignoring observation bias led to strong negative bias in projected
population size for growing populations, but only minor negative bias in declining
populations. Accounting for apparently minor biases associated with ground counts is
important for ensuring accurate estimates of abundance and population dynamics especially
for increasing populations. This could be particularly important for informing conservation
decisions in projects aimed at recovering depleted populations.
51
52 ***Keywords:*** false positives, false negatives, multinomial model, classification, calibration, population
53 dynamics
54

Introduction

Understanding spatiotemporal mechanisms for changes in animal abundance rests on information about survival, movement, and fecundity (Sample et al. 2018, Sen and Akçakaya 2022). Challenges when estimating these parameters due to observation bias are notorious (Kéry and Royle 2015), and correcting for these observation biases requires the use of statistical methods such as capture-recapture models when estimating survival, recruitment and/or abundance (Williams et al. 2002). Capture-recapture models also serve as a core building block of integrated population models (Zipkin and Saunders 2018, Zhao 2020). Several quantities have been proposed to measure fecundity (Etterson et al. 2011). Existing methods for estimating fecundity, however, typically do not address inaccurate counts of offspring (but see Sen and Akçakaya 2022). Failing to account for such bias may undermine inferences about population dynamics and their drivers at this critical part of the annual life cycle, and in turn compromise efforts to conserve species.

A particular challenge when estimating fecundity is ascertaining the number of young produced per litter or per nest (henceforth, brood). The number of young can be determined accurately by accessing the brood close to fledging in altricial species, as is often done in bird ringing programs (e.g., Katzenberger et al. 2019). Depending on nest location, doing so may require specialized training and equipment (e.g., climbing or using a lifting platform for tree-nesting birds), is time-intensive (Fuller and Mosher 1981) and can stress and disturb the adults (Andersen 1990), which could reduce productivity (Ibáñez-Alamo et al. 2012). In some cases, accessing a nest may not be possible at all if the structure supporting the nest is too fragile (e.g. a thin branch that cannot support human weight). To minimize disturbance, costs, risk to the observer, and simplify field logistics, counts of young are therefore often obtained at a distance from the ground, for example in tree-nesting raptors (Brown et al. 2013). Nest monitoring can also be done from the air using helicopters (White and Sherrod 1973) or more recently drones (Gallego and Sarasola 2021). In this context, counts of young may not reflect true brood size, and obtaining an accurate count likely depends on several factors, including observation distance, time spent observing, level of camouflage, activity level of the young as well as their position in the nest (i.e., central or close to the edge).

Determining brood size, when done via observation from a distance, potentially suffers from two types of observation error, false negatives (when young are missed; e.g., Goszczyński 1997) and false positives (when young are double-counted, as occurred in our surveys; see below). False negatives have long been acknowledged as pervasive in wildlife surveys, and there is a wealth of statistical models dealing with false negatives: capture-recapture models (Otis et al. 1978), N-mixture (Royle 2004), or distance sampling models (Buckland et al. 2015) estimate abundance when the probability of detecting any individual is < 1; and occupancy models (MacKenzie et al. 2002, Tyre et al. 2003) estimate occurrence when species detection is < 1. Even though several studies have shown that false positives are also common in wildlife surveys (Clement et al. 2014, Brack et al. 2018, McKibben et al. 2023) and that models accounting only for false negatives are very sensitive to the presence of false positives (Miller et al. 2011, Lahoz-Monfort et al. 2016, Link et al. 2018, Nakashima 2020), the development of models accounting for false positives has lagged behind (Dénes et al. 2015). Existing efforts to formally account for false positives have largely focused on occupancy models (Royle and Link 2006, Miller et al. 2011, Chambert et al. 2015), which require some form of additional information to be uniquely identifiable, including constraints on parameters (Royle and Link 2006) or detections with and without false positives (Miller et al. 2011, Chambert et al. 2015). In the context of abundance estimation, the Poisson-Poisson version of the N-mixture model (Kéry and Royle 2015) allows for counting an individual more than once but requires repeated visits. In contrast, here, we develop and test a method that accounts for both over and undercounting based on a single visit to each nest. We are unaware of any studies that have accounted for false positive bias when estimating brood size.

To do so, we treat brood size as a state that is imperfectly observed when the young are counted from a distance. We then expand the approach of Royle and Link (2006) to formulate observation state probabilities (here, observed number of young) as conditional on true states (here, true number of young) to >2 states. Without additional information, parameters in this model are not uniquely identifiable, and we adopt a modified version of the calibration approach by Chambert et al. (2015), in which for some sites the true state is confirmed (here, nests that are climbed and young counted without error) to inform the imperfect observation process. We evaluate the performance of this model using a case study on the red

107 kite (*Milvus milvus*) in central Europe. Finally, we use population projections to investigate the implications
108 of not accounting for observation error in brood size counts for population trajectories.

Methods

Empirical data collection
We collected data on red kite brood size during June of 2021 and 2022 during the fledging period. From
the middle of June onwards and approximately three weeks after hatching, young reach a development
phase that is sufficient for tagging and ringing (Cramp and Simmons 1980). Nests were identified from the
ground at the beginning of the breeding season by members and partners of the LIFE EUROKITE (LIFE18
NAT/AT/000048) project, and checked again at the end of June to confirm occupancy. In 2021, we sampled
8 nests in Germany. In 2022, 14 of the 18 sampled nests were located in stands of deciduous forest within
Lower Austria, along the March River that forms the border with Slovakia. Remaining nests that year (i.e.,
3 in Upper Austria and 1 in Slovakia) were located in mixed deciduous-coniferous forests.
Each nest was visited once by two observers. The ground observer aimed to find the best vantage point
between 50 m and 100 m from the nest, and used a spotting scope or binoculars to observe nestlings.
Counting from the ground proceeded before and during climbing of the tree by the second observer, a
person certified for tree-climbing along with tagging and ringing of the young red kites. The ground
observation started as soon as the best vantage point to observe the nest was found and stopped when
the climber reached the nest. The climber did not reveal the number of young in the nest until after the
ground-count was conducted.

Model development
Royle and Link (2006) developed an occupancy model allowing for false positive and false negative
detections by formulating probabilities of observed states y (0 or 1) conditional on true states z (0 or 1).
For example, when a site is not occupied ($z = 0$), we can either obtain a true negative observation ($y = 0$)
with probability p, or we can obtain a false positive observation ($y = 1$) with probability $1 - p$. They
conceptually expanded this model to more than 2 states; for each true state, the probabilities of all possible
observed states sum to 1, and observations y for a given state z are multinomial random variables.
Here, we adapt this approach and consider brood size as the true state with Z possible categories
corresponding to possible numbers of nestlings. Because only nests with at least 1 observed nestling are
included in nestling count data sets, and we do not expect empty nests to be classified as occupied, $z = 0$
is not a possible state, but the model could easily be expanded to allow for true and/or observed empty
nests. Z is given by the biological upper limit of how many offspring the focal species can raise:

(1) $z \sim Multinomial(\boldsymbol{\pi}, 1)$

Here, $\boldsymbol{\pi}$ is the vector of probabilities that a nest falls into a given state category (henceforth, nest
probabilities). The expected number of nestlings in any given nest is $\lambda = \sum_{i=1}^{Z} i \times \pi_i$. We refer to λ as
fecundity.
Observed states, y, can fall into one of K categories (where K may be equal to Z), and the probability of
observing a particular category k, $P(y = k)$, depends on the true state z.

(2) $y|z \sim Multinomial(\mathbf{p}_z, 1)$

Thus, $\mathbf{p}$ is a $Z \times K$ classification matrix, and we refer to values in $\mathbf{p}$ as classification probabilities.
Royle and Link (2006) showed that this model is not uniquely identifiable for the 2-state and K-state
case; further development by Miller et al. (2011) and Chambert et al. (2015) allowed for the incorporation
of additional information to make parameters identifiable. Here, we adopt a modified version of the
'calibration' approach by Chambert et al. (2015), in which at some sites, the true state is confirmed so that
direct information on the classification process is available. Specifically, we propose to combine imperfect
or undertain counts y (e.g., ground counts of nestlings), which provide imperfect representations of the
true state, with confirmed or certain counts (e.g., climb counts for raptors) which provide z without error.
Collecting both types of data, y and z, at a set of nests allows estimation of the parameters in the

classification matrix **p**. We can then use estimates of **p** to obtain estimates of the expected number of
nestlings in a data set consisting only of imperfect counts, *y.new*. This differs from the calibration approach
of Chambert et al. (2015) in that certain observations of *z* also inform the state process (ie, estimates of **π**
and λ). We consider two possible sampling scenarios:
(1) The paired counts (i.e., imperfect ground counts, *y*, with perfect climb counts, *z*; sample size *n*) and
the imperfect-only counts, *y.new* (sample size *n.new*) come from the same population, i.e., they share **π**
and λ. In this situation, one could use only measures of *z* (i.e., only the climb counts from the paired data
set) to determine **π** and λ. But adding *y.new* increases sample size and thus may yield more precise
estimates of these parameters. Note that in the set of formulas below, *z.new* is not observed, whereas for
the paired data, *z* is observed.

(3) $z.new \sim Multinomial(\boldsymbol{\pi}, 1)$
(4) $y.new | z.new \sim Multinomial(\mathbf{p}_{z.new}, 1)$

(2) The paired *y* and *z* versus *y.new* come from different populations that may differ in fecundity (i.e.,
in one population, birds may produce more/fewer nestlings, on average). In this case, we can still use *y* and
z to estimate **p** – which we assume is shared between both populations – but have to estimate a separate
π.new.

(5) $z.new \sim Multinomial(\boldsymbol{\pi}.\mathbf{new}, 1)$
(6) $y.new | z.new \sim Multinomial(\mathbf{p}_{z.new}, 1)$

This is the more interesting scenario, as in contrast to **π, π.new** cannot be estimated from the paired
data alone. Accuracy of **π.new** will depend on how accurately **p** is estimated. Depending on *Z* and *K*, the
number of parameters in **p** can quickly become unwieldy (Royle and Link 2006) and require unrealistically
large sample sizes to be estimated well. We address this issue in our simulation by constraining the mis-
identification process such that *y* can differ from *z* by no more than 1. In the Discussion, we further present
alternative model formulations that simplify the detection/classification process and reduce the number
of parameters, along with some preliminary exploration of model behavior. A full implementation and
presentation of these alternative formulations is beyond the scope of the current study.

Simulation – parameter estimation
We used a simulation approach to determine how well parameters of the above described model can
be estimated. We refer to data sets in which measures of both *z* and *y* from the same nests are available
as 'paired data sets'; and data sets with paired data plus additional uncertain counts *y.new* as 'combined
data sets'.
We parameterized our simulation based on the empirical data on the 26 paired (uncertain) ground and
(certain) climb counts described above and a literature review of reproductive values in red kites. Reported
values of clutch size based on a global literature review ranged from 1.9 - 3.2, and number fledged per
successful pair from 1.4 to 2.3 (Sergio et al. 2019). As nestlings have not yet fledged when they are counted,
we used reported clutch size as a guideline to set three levels of reproduction. Our baseline $\boldsymbol{\pi} = $
$\{0.1, 0.45, 0.4, 0.05\}$, resulting in fecundity $\lambda = 2.4$. We also created two alternative vectors of **π**, one with
a higher fecundity of $\lambda_{high} = 2.8$ and a high rate of 3 nestlings: $\boldsymbol{\pi}_{high} = \{0.05, 0.2, 0.65, 0.1\}$; one with a
lower fecundity of $\lambda_{low} = 2$ and a more even distribution of nestling probabilities: $\boldsymbol{\pi}_{low} = $
$\{0.3, 0.45, 0.2, 0.05\}$.
Our parameterization of **p** was strongly informed by the empirical data, as to our knowledge no other
published data sets for paired nestling counts are available. We assumed *K=Z* (implying that no observer
would report more than 4 nestlings, given the knowledge of the species' breeding habits, and the dataset
would not contain nests in which observes recorded 0 nestlings) and constructed **p** using the following
rules: (1) the probability of observing the correct number of nestlings is higher than the probability of
observing the wrong number of nestlings but decreases as true number of nestlings increases; (2) when
observing the wrong number of nestlings, missing a nestling (false negative) is more common than
overcounting (false positive); and (3) at most 1 nestling is missed/invented. The last rule was especially
important to limit the number of cells in **p** that need to be estimated (reducing it from 16 to 10); it may be

relaxed in theory but that will increase the necessary sample size to estimate all values in **p**; moreover, not
fixing probabilities of highly unlikely combinations of z and y (e.g., observing 4 nestlings in a nest that has
1) may cause numerical problems. Based on this parameterization, we attempted to answer three
questions:

*Question 1: How does the sample size of a paired data set affect estimates of **p**?*
In paired data sets, we have direct measures of z, and we therefore expect unbiased estimates of **π**
and λ, with precision increasing with sample size. We expect a much stronger impact of sample size on
estimates of **p**, because sample size per element of **p** is much lower than for **π**. Also, how often certain
nestling numbers are observed is affected by how common they are in the population, as well as the
classification process; thus, for several cells in **p**, we may have 0 observations even at intermediate sample
size. We therefore expect small sample bias in **p**, particularly in cells referring to rare states, that will decline
with increasing n.
We generated 100 paired data sets for n = 10, 25, 50, 100 and 250 nests, mirroring a range of sample
sizes in red kite monitoring programs (Sergio et al. 2019). We did so for each of the three alternative levels
of fecundity. We estimated **π,** λ and **p** from the resulting data sets using our nestling count model and
compared estimates to truth. We focus on absolute bias in estimates of **p** and relative bias and precision
(CV = estimate/standard error) in estimates of λ. We report absolute bias for **p** and abstain from reporting
a CV because several true values of **p** are fairly small, leading to extreme relative bias/CV values.

Question 2: How much precision do we gain in estimating fecundity when combining a small paired data
set with uncertain counts, when both come from the same population (i.e., share fecundity)?
Climb counts are logistically and financially challenging to implement at large spatial scales;
representative sampling of a population may require large sample sizes that are unfeasible with climb
counts. It may therefore be beneficial to combine a fairly small set of paired counts with a larger data set
of uncertain ground counts only. In this scenario, again, because direct information on z is available from
the paired data, we expect **π** and λ to be unbiased, but we expect precision to increase as we add uncertain
count only data sets of increasing size.
We created combined data sets by combining paired data sets with n = 25 with uncertain count only
data (generated from the same values of **π** as the paired data) of size $n.new$ = 10, 25, 50, 100 or 250 (100
combined data sets for each value of $n.new$). We estimated **π,** λ and **p** from the resulting data sets and
compared estimates to truth, as well as to estimates from the corresponding scenario from Question 1
(i.e., with n = 25 paired counts and no additional ground counts). We did so for all three levels of fecundity.

Question 3: How accurately can we estimate fecundity for a set of uncertain counts when we combine it
with paired data sets of different sample sizes from a population with different fecundity?
This is likely the most important aspect of this model, as it would allow using paired counts from one
population to estimate **π** and λ in a different population (with different fecundity) assuming the
classification process is the same.
We created combined data sets by combining paired data (generated from baseline **π**) of size n = 10,
25, 50, 100 or 250 (100 combined data sets for each value of n) with uncertain count only data of size $n.new$
= 50 generated from **π. new**. We did so separately for **π. new** $= \textbf{π}_{high}$ and **π. new** $= \textbf{π}_{low}$. We estimated
π. new, $\lambda.new$ and **p** from the resulting data sets and compared estimates to truth.
For all scenarios/questions, we further compared bias in model-estimated λ to bias if we calculated λ
directly from uncertain counts (λ_y), i.e., ignored any observation bias.

Model implementation
We implemented the model in a Bayesian framework, for the ease with which this framework handles
latent state variables. Specifically, we used the package nimble v.0.12.2 (de Valpine et al. 2017, 2022) in R
v.4.2.1 (R Core Team 2022). To improve convergence of MCMC chains, we used the log-linear formulation
of the multinomial model for both **π** and **p**, which estimates $log(\gamma)$, then rescales to multinomial cell
probabilities using $\frac{log(\gamma)}{\sum log(\gamma)}$. We wrote a custom Nimble function to fix some values for **p** to 0. We used
moderately informative Normal priors ($\mu = 0, \sigma = 2$) on $log(\gamma)$ to ensure chain convergence. Note that

when estimating only $\boldsymbol{\pi}$ from climb counts, convergence was not an issue and informative priors were not
necessary, but extending the model to estimate $\mathbf{p}$, even only from paired data, led to convergence issues
in the absence of these moderately informative priors. For all models under Question 1 and 2, we ran 3
parallel chains with 5000 iterations per chain, discarding the first 2500 as burn-in; for Question 3, we ran
chains for 10000 iterations, discarding the first 5000. We checked convergence by calculating the Gelman-
Rubin statistic (Brooks and Gelman 1998) and treated chains as converged when $\hat{r} < 1.2$ (this is a little less
conservative than the typical cutoff of 1.1; for time constraints we wanted to avoid running a large number
of models for longer, but for applications of the model to real data we would urge users to run the model
long enough to obtain $\hat{r} < 1.1$). We also visually inspected trace plots of every 20th simulation iteration
per scenario. We excluded non-converged simulation iterations from summary calculations and plots. The
numbers of excluded simulation iterations are provided in Appendix 1: Tables S1-3.

Problematic datasets

Traceplot inspections showed that for datasets in which some true state category was not represented
at all in the paired counts (henceforth, datasets with missing truths), some values for $\mathbf{p}$ were not estimable
(posteriors were flat over the 0-1 interval). This is intuitive: the model cannot estimate classification
probabilities for a state not represented in the data. We therefore removed such data sets, leading to a
lower number of simulation iterations: for n = 10, 70-75% of data sets had to be excluded; for n = 25, 20-
45%; for $n > 25$, <10% of data sets had to be excluded. Because not all state categories are equally likely to
be missing, however, this led to a biased sample of retained datasets, particularly for n = 10. For baseline
λ and λ_{low}, z = 4 was the rarest category and retained datasets (which had at least one nest with z = 4)
therefore had an average number of nestlings higher than λ or λ_{low}. For λ_{high}, z = 1 was the rarest category
and retained datasets (with at least one nest with z = 1) had a lower average number of nestlings than
λ_{high}. Only for n = 10 and 25 did this bias in retained data exceed 1% (Appendix 1: Table S4). To assess bias
in model estimates of λ fairly for n = 10 and 25, we compared estimates against the retained sample mean
$\bar{\lambda}$, instead of the original true input value of λ.
Because dataset removal due to missing truths was applied only to paired data, there is no systematic
bias in additional uncertain count data, which were generated independently to address Question 2 and 3.
For Question 2, however, the model assumes λ to be the same in the paired and the uncertain count only
data, which is not the case when datasets with missing truths are excluded from the paired data. To avoid
this mismatch in λ between paired and uncertain count data in Question 2, we simulated these additional
uncertain counts using the sample average values of $\bar{\boldsymbol{\pi}}$ of retained datasets and compared estimates of λ
from combined paired and uncertain counts against the retained sample mean $\bar{\lambda}$. For simplicity, we still
refer to scenarios in Question 1 and 2 by the original data-generating input values of λ (i.e., 2.4, 2 and 2.8).

Population projections

To investigate how bias in fecundity affects population projections, we implemented a second
simulation, projecting an age-structured population 50 years into the future, considering females only.
Input parameter values for starting age distribution, age-specific proportion of the population that breeds,
and age-specific survival were taken/adjusted from Sergio et al. (2021) and are summarized in Appendix 1:
Table S5. We used two sets of age-specific survival probabilities, one that led to a growing population and
one that led to a declining population. We combined each with values of $\boldsymbol{\pi}$ from the parameter estimation
simulation to generate the total number of nestlings. Assuming a 50:50 sex ratio at nestling stage, we then
randomly assigned each nestling a sex and only considered females going forward in time, to keep with the
female-only projection. We projected the population with the true values of $\boldsymbol{\pi}$ (high and low), the
corresponding expected observed values of $\boldsymbol{\pi}$ if we used ground counts only and ignored observation bias
($\boldsymbol{\pi}_y$), and the average (across simulation iterations) model-estimated values of $\widehat{\boldsymbol{\pi}.\mathbf{new}}$ from Question 3,
combining 50 uncertain counts generated from $\boldsymbol{\pi}.\mathbf{new}$ with 50 paired counts generated from baseline $\boldsymbol{\pi}$.
Using the average over simulation iterations ignores the fact that estimates at each individual iteration
were likely more different from truth than the average. But our interest here is to see the general impact
on population trajectories when ignoring observation error vs when accounting for it; thus, we ignored
variability in estimates in our projection. We used a starting population size of 100 females and we
projected the population 1000 times over 50 years. We calculated average trajectories and 95% confidence

intervals over the 1000 iterations, as well as average difference in population size in year 50 between the
three projections using true, observed uncorrected, and estimated fecundity.

Figure 1 - Absolute error (boxplots) and bias (median line) of estimates of classification probabilities
p of observing *y* young when the true number of young is *z* from paired certain and uncertain counts
of different sample sizes (*n*) from a model accounting for over- and undercounting of young. Red
dashed line indicates 0 error/bias. Values of **p** for missing combinations of *y* and *z* were fixed to 0.

 Results

Empirical data

312   The average number of young observed in climb counts at the 26 surveyed nests was 2.15 (range: 1 to
313 3); the average number based on ground counts for the same nests was 2.04 (range: 1 to 4), thus showing
314 approximately 5% negative bias. At 19 nests, ground counts yielded the same values as climb counts; climb
315 counts were biased low by one nestling in 5 and biased high by one nestling in 2 nests. All combinations of
316 ground and climb counts are available along with the code to implement simulations.

Simulation - parameter estimation

318   Input parameter settings of nest probabilities (**π**) and classification probabilities (**p**) led to 7% negative
319 bias in fecundity when ignoring observation bias for λ and λ_{high}, and 6% negative bias for λ_{low}.

320   For Questions 1-3, median bias across all kept simulation iterations, CV of λ, as well the RMSE of all
321 parameters for all scenarios can be found in Appendix 2.

Question 1

As expected, when analyzing paired counts only, estimates of λ were essentially unbiased, with relative
bias < 2% for all scenarios (compared to retained sample mean for *n* = 10 and 25, compared to true input
value for all other *n*; Appendix1: Figure S1A) and the CV of λ declined with increasing sample size, but was
already low at *n* = 10 (ranging from 8.9% for λ_{high} to 11.4% for λ_{low}; Appendix1: Figure S1B). Estimates of
π showed mild bias at *n* = 10 and 25, but were essentially unbiased at larger sample sizes (Appendix1:
Figure S2). As expected, estimates of **p** showed considerable bias for small sample sizes (*n* = 10 and 25) and
were only largely unbiased at *n* = 250, with some bias remaining in some of the probabilities (Figure 1 for
λ, Appendix1: Figure S3 for λ_{low} and λ_{high}).

Question 2

Adding uncertain ground counts *y.new* to 25 paired counts, with both datasets having the same
underlying fecundity, bias in λ was reduced with 10 additional ground counts; adding more ground counts
slightly increased bias, though it remained very low overall (< 1.6%; Figure 2A). Bias in λ was generally
negative. Adding ground counts only marginally improved the CV of λ, e.g. by 2-3% comparing 250 to 0
additional ground counts (Figure 2B). Overall, patterns of bias in **π** were similar to those in λ, with 10
additional ground counts improving on estimates, but more ground counts increasing bias again slightly
(Appendix1: Figure S4). Similarly, bias in **p** (e.g., Appendix1: Figure S5) remained very similar to that
observed with no additional ground counts, as expected, since *y.new* does not contribute information to
estimating **p**. Interestingly, the more ground counts were added, the higher the rate of non-convergence
(Appendix 1: Table S2); this was not the case in Question 3.

Question 3

When combining paired counts of different sample sizes *n* from a population with nest probabilities **π**
with 50 uncertain counts from a population with **π.new**, estimates of $\lambda.new$ were largely unbiased across
all sample sizes (Figure 3). There was a tendency for bias to increase with increasing *n* for $\lambda.new = \lambda_{low}$,
however, even at *n* = 250, bias was only 2%. Whereas in Question 1 all values of **π** became essentially
unbiased at *n* = 50 or 100, here, some bias in **π.new** remained at all sample sizes, with no obvious trends
(Appendix1: Figure S6). Bias in estimates of **p** decreased with increasing paired sample size; estimates of **p**
also became increasingly similar to those under Question 1 with increasing sample size (Appendix 1: Figure
S7).

Figure 2 - Relative error (boxplots) and bias (median) (A) and coefficient of variation (CV; B) of estimates of fecundity λ (average number of young per nest) from a model accounting for over- and undercounting of young, fit to 25 paired certain and uncertain counts plus varying numbers of additional uncertain ground counts (x axis), when both data sets come from the same population (with the same fecundity; Question 2). Red dashed line and blue lines in A show 0 bias/error and relative bias if fecundity was calculated based on uncertain counts only, ignoring observation bias, respectively.

Figure 3 - Relative error (boxplots) and bias (median) of estimates of fecundity λ (average number of young per nest) from a model accounting for over- and undercounting of young, fit to 50 uncertain counts from a population with $\lambda = 2$ or $\lambda = 2.8$ plus varying numbers of paired certain and uncertain counts (x axis) from a populations with $\lambda = 2.4$. Red dashed line shows 0 bias/error, and blue lines show bias if fecundity was calculated based on uncertain counts only, ignoring observation bias.

Population projections

Using nest probabilities based on imperfect ground counts only, ignoring observation bias, led to negative bias in projected population trajectories. That bias was stronger in growing populations and at higher fecundity (π_{high}), because of the exponential growth underlying the projections. In agreement with the largely unbiased estimates of fecundity from the parameter estimation simulations, using average model-estimated nest probabilities led to population trajectories that were largely identical to those using true input values (Figure 4). Consequently, relative bias in population size in year 50 was negligible when using estimated nest probabilities (< 1% to 5% negative bias), but considerable when not accounting for observation bias (25% to 42% negative bias; Table 1).

Table 1 - Percent bias in projected population size at time T=50 when using the observed uncertain counts of nestlings to determine fecundity (observed) and when using estimated fecundity accounting for observation error in uncertain counts (estimated). Median (2.5th and 97.5th percentile) across 1000 simulated population trajectories, relative to the trajectory using true fecundity for increasing and decreasing populations with either high (input π_{high}) or low (input π_{low}) fecundity.

Population	Observed vs true	Estimated vs true
High, increasing	-41.73 (-61.79, -18.11)	-4.83 (-36.79, 35.41)
High, decreasing	-36.20 (-77.38, 30.38)	-2.26 (-59.30, 90.97)
Low, increasing	-31.28 (-61.30, 12.15)	5.59 (-38.45, 64.11)
Low, decreasing	-25.46 (-90.06, 103.74)	-0.62 (-85.09, 148.58)

Discussion

In count based surveys, counting individuals more than once (i.e., a form of false positive error) is a common problem (e.g., in camera trapping - McKibben et al. 2023; unmanned aerial surveys - Brack et al.

2018). Yet, few modeling approaches explicitly account for such false positives (but see Kéry and Royle
2015). In our classification model, we treat the true state - the number of nestlings - as categorical. This
allowed us to adopt the strategies developed for multi-state occupancy models that account for false
positives and state uncertainty (Royle and Link 2005, 2006, Miller et al. 2011). In particular, we used a
calibration approach that combines uncertain with certain counts to learn about the probabilities of
observing a certain number of nestlings conditional on the true number of nestlings, **p**, and then using
those probabilities to correct uncertain counts in a separate dataset (Chambert et al. 2015).

We found that our approach produced largely unbiased estimates of fecundity for nestling counts
suffering from both false positive and false negative observation error. Unbiased estimates persisted
regardless of whether the separate dataset had the same data-generating value of fecundity (i.e., came
from the same biological population) or a different value (i.e., came from a different population). Ignoring
observation errors in the number of nestlings led to underestimating forecasted abundances, particularly
in large or growing populations. Using estimates that corrected for both false negatives and false positives,
however, led to population trajectories that closely resembled true trajectories.

Figure 4 - Population projections for growing (A) and declining (B) populations with two levels of
fecundity (λ). Solid lines show median trajectory and colored polygons 2.5th and 97.5th percentiles of
1000 projections using true input value of λ. Dashed lines show median trajectory (across 1000
projections) when calculating λ based on uncertain counts of young only, and dotted line shows
median trajectory (across 1000 projections) when estimating λ with a model accounting for over- and
undercounting of young.

The most crucial variable determining the performance of our classification model was the sample size
of the paired counts (i.e., the number of nests for which both certain and uncertain observations were
available). Specifically, some classification probabilities **p** were not estimable when a dataset of true

nestlings was missing a nestling category (e.g., no nest in the dataset had 4 nestlings), as in those cases the
dataset contains no information on the classification probabilities related to the missing true state. Missing
true states were, of course, more common in small data sets; in our simulation, a paired $n = 50$ was
generally sufficient to avoid this problem. The necessary sample size, however, depends on the distribution
of true states probabilities, $\boldsymbol{\pi}$. In particular, the rarer one or some categories, the larger the sample has to
be to capture all categories.

A truncated Poisson distribution is a potential alternative for modeling nest probabilities, $\boldsymbol{\pi}$. With a
single parameter (as opposed to Z-1 parameters in the multinomial model), it is more parsimonious and its
specific properties can inform nest probabilities for categories with no observations. We found, however,
that a truncated Poisson did not fit the empirical data well and led to biased estimates of $\boldsymbol{\pi}$. Limited trials
suggested that this poor fit increased bias in estimates of fecundity when combining paired data with
uncertain counts from populations with different fecundities. A more thorough evaluation would be
necessary to determine the extent of this bias as a function of the degree of lack of fit.

As an alternative, unobserved or very rare state categories could be excluded when estimating
fecundity; that has the added advantage of reducing the size of the classification matrix $\mathbf{p}$. We did not
explore any potential effect of excluding non-represented nestling categories on bias and precision in
estimates of fecundity. Even in datasets without missing true categories, many classification probabilities
showed large bias; in individual simulation iterations, some probabilities were often barely estimable, with
large uncertainty, and we had to use mildly informative priors on the log-probabilities of the classification
model to ensure chain convergence. All of these issues are likely due to small sample sizes for individual
classification probabilities, even though we already reduced their number by limiting the difference
between true states and observations to 1. These limitations highlight that with an increasing number of
true states, the size of the classification matrix becomes unwieldy and parameters comprising this matrix
become challenging to estimate (Royle and Link 2006). Other constraints, such as imposing that true
classification probabilities are higher than false classification probabilities (Royle and Link 2005, 2006,
Miller et al. 2011), and that undercounting is more likely than overcounting are justified based on our
empirical data and could help further improve the performance of the model we developed. Still, based on
our simulation results we believe this classification model is of limited use for species with a wider possible
range of brood sizes than what we considered here.

There are ways in which the classification model could be simplified. For example, the imperfect
observations could be regarded as Poisson random variables with mean $z \times \varrho$, where ϱ is a detection rate
correcting for net bias in nestling counts (for example, in our empirical data, undercounts were more
frequent than overcounts, thus, net bias was negative and ϱ would be < 1). This is akin to the observation
process in the Poisson-Poisson N-mixture model (Kéry and Royle 2015), which allows for both over and
undercounting. Preliminary trials with our simulated data suggest that this model will lead to bias in
estimates of fecundity when paired data are combined with uncertain-only data, even though the paired
data provide direct information on ϱ. N-mixture models rely on temporally replicated sampling, whereas
our data has a single observation per nest. When it is easier to obtain repeated uncertain counts from nests
than to climb them to ascertain the true number of nestlings, a PP N-mixture model may be a viable
alternative to our classification model. The standard PP N-mixture model has been shown to provide largely
unbiased estimates of abundance (Nakashima 2020). How well a modified version with a multinomial (or
truncated Poisson) state process would perform for estimating fecundity remains to be evaluated.

Our empirical data suggested that both over and undercounting misses the true number of nestlings
by at most 1. If that is the case, the classification process can be broken down into two Bernoulli processes:
whether the number of nestlings is counted correctly (with probability p_c), and if not, then whether the
number is overcounted (with p_{over}; as opposed to undercounted). The classification probabilities can then
be written as a function of these Bernoulli probabilities. If either or both probabilities can be assumed to
be the same regardless of the true state, this will reduce the number of observation parameters that need
to be estimated. Initial trials, using our original simulated data and allowing p_c to vary by true state but
fixing p_{over} suggested that this simplified model may yield similarly unbiased estimates of fecundity as the
more complex data generating model (though in our case, the two model versions differed only by one
free parameter).

Adding uncertain counts from the same biological population (i.e., with the same fecundity) to 25
paired counts (Question 2) only marginally improved the precision in estimates of λ and led to more

convergence problems. Thus, in a study focusing on a single population with constant fecundity, there is
little to no benefit from investing effort in additional ground counts. All effort in such a case should be
dedicated to collecting unbiased brood size information at as many locations as possible. In contrast,
adding even a small number of paired counts to a dataset of uncertain counts from a population with
different fecundity greatly reduced bias in estimates of λ_{new} compared to ignoring observation bias
altogether. As long as the observation process (i.e., classification probabilities) can be assumed to be the
same in different populations, information on these probabilities from one population can be leveraged to
obtain unbiased estimates of reproduction in another.

The validity of detection parameters beyond the spatial and temporal scope of sampling used to
estimate them needs to be evaluated carefully (e.g., Williams et al. 2002). An important future step is
therefore to identify factors affecting classification probabilities and to integrate them into the
classification model. For example, nest visibility (and hence, the probability of correctly identifying the
number of nestlings) may be higher in deciduous than in mixed or coniferous forests. Identifying important
predictors of classification probabilities would make estimates of **p** more transferable to other populations
as sources of variation in the classification parameters could then be taken into account.

Alternatively to count-based methods to estimate brood size or fecundity, capture-mark-recapture
methods can be used to estimate age-specific population sizes and derive fecundity as number of juveniles
per adult (e.g., Sen and Akçakaya 2022). Capture and recapture of individuals, however, requires significant
expenditures of time and money by investigators (Lieury et al. 2017). Such repeated sampling may thus not
be feasible in many situations.

Recommendations for data collection

There are several important considerations when obtaining data necessary for applying our modeling
framework. Unless researchers adapt this framework to incorporate covariates for the classification
probabilities, potential biases should be minimized by keeping the observation conditions as consistent as
possible. These include weather, time of day, nestling age, duration of observation, method of observation
(e.g., binocular, spotting scope), and distance from the ground observer's position to the young. Fuller et
al. (1995) recommended observing the nest for 30 to 60 minutes from the ground to accurately count the
young. Observing young for > 1 hour likely increases disturbance and may increase the risk of overcounting
the number of nestlings.

The paired ground and climb counts should be implemented during a single visit when at least one
observer is present at all times, to ensure no young are lost (e.g., due to predation) between the two
observations. A possible exception is when ground counts are paired with perfect counts obtained via video
documentation of the nest. Installing a camera would require climbing to the nest at an earlier stage,
disturbing both parents and nestlings when they are more vulnerable (Götmark 1992, Ibáñez-Alamo et al.
2012). Cameras could be installed during the non-breeding season at previously used nests but without
guarantee that these will be occupied again.

Conclusion

Fecundity is a key parameter affecting population dynamics (Sibly and Hone 2002), and we showed that
even apparently low bias in fecundity estimates can lead to considerable bias in population projections
under certain conditions. This can have implications for species conservation, for example, when
conducting a population viability analysis (Boyce 1992). This highlights the need for statistical models that
can produce unbiased estimates of fecundity taking into account both false positive and false negative
observation bias. We believe the model presented here represents a useful approach to dealing with both
kinds of observation bias while taking into account the specific characteristics of nestling count data,
particularly, the limited number of true and observed states. The approach hinges on the assumption that
the classification process is the same for data with and without certain observations. To improve
transferability of classification probabilities between points in space or time, we believe the most
important next step is to examine potential predictors of these probabilities (e.g., time spent observing),
and to extend the model to incorporate them. Including such covariates would be easier if the number of
classification parameters could be reduced and we sketch out several possibilities for such simplified
models, the behavior of which under different scenarios remains to be explored. More broadly, our
approach contributes to the field of models accounting for biases induced by false-positive observations,

which remains underrepresented in the literature even though its importance has been highlighted
repeatedly.
515

Appendices

517   Appendix 1: Additional information and results from estimation simulation and population projections
518   Appendix 2: Median bias, precision and accuracy of parameter estimates
519   Appendices can be found at doi.org/10.5281/zenodo.10245398 (Sollmann et al. 2023).

Acknowledgements

521   We thank R. Raab for allowing the student assistants to join the telemetry team in visiting the nests for
522 data collection. F. Knufinke, M. Kolbe, and E. Schöll provided useful input in the development of the
523 manuscript. We thank M. Malvezin for assistance with refining the field sampling protocol and data
524 collection.

Data, scripts, code, and supplementary information availability

526   Data, scripts and code can be found at doi.org/10.5281/zenodo.10245803 (Sollmann 2023).

Conflict of interest disclosure

528   The authors declare that they comply with the PCI rule of having no financial conflicts of interest in
529 relation to the content of the article.

Funding

531   The Institute of Wildlife and Game Management at the University of Natural Resources and Life
532 Sciences provided travel funding for the field work. No other funding to declare.

References

Andersen, D. E. 1990. Nest-defense behavior of red-tailed hawks. Condor 92:991-997.
https://doi.org/10.2307/1368735
Boyce, M. S. 1992. Population viability analysis. Annual Review of Ecology and Systematics 23:481-506.
https://doi.org/10.1146/annurev.es.23.110192.002405
Brack, I. V., A. Kindel, and L. F. B. Oliveira. 2018. Detection errors in wildlife abundance estimates from
Unmanned Aerial Systems (UAS) surveys: Synthesis, solutions, and challenges. Methods in Ecology and
Evolution 9:1864-1873. https://doi.org/10.1111/2041-210X.13026
Brooks, S. P., and A. Gelman. 1998. General methods for monitoring convergence of iterative simulations.
Journal of Computational and Graphical Statistics 7:434-455. https://doi.org/10.2307/1390675
Brown, J. L., K. Steenhof, M. N. Kochert, and L. Bond. 2013. Estimating raptor nesting success: Old and new
approaches. Journal of Wildlife Management 77:1067-1074. https://doi.org/10.1002/jwmg.566
Buckland, S. T., E. A. Rexstad, T. A. Marques, and C. S. Oedekoven. 2015. Distance sampling: methods and
applications. Springer International Publishing, Cham, Switzerland.
Chambert, T., D. A. W. Miller, and J. D. Nichols. 2015. Modeling false positive detections in species
occurrence data under different study designs. Ecology 96:332-339. DOI: https://doi.org/10.1890/14-
1507.1
Clement, M. J., T. J. Rodhouse, P. C. Ormsbee, J. M. Szewczak, and J. D. Nichols. 2014. Accounting for false-
positive acoustic detections of bats using occupancy models. Journal of Applied Ecology 51:1460-1467.
https://doi.org/10.1111/1365-2664.12303

Cramp, S., and K. Simmons. 1980. Handbook of the Birds of Europe the Middle East and North Africa: The
Birds of the Western Palearctic, vol. 2: Oxford University Press.
de Valpine, P., C. Paciorek, D. Turek, N. Michaud, C. Anderson-Bergman, F. Obermeyer, C. Wehrhahn
Cortes, A. Rodrìguez, L. D. Temple, and S. Paganin. 2022. NIMBLE: MCMC, particle filtering, and
programmable hierarchical modeling. https://doi.org/10.5281/zenodo.1211190, R package version
0.12.2, https://cran.r-project.org/package=nimble.
de Valpine, P., D. Turek, C. J. Paciorek, C. Anderson-Bergman, D. T. Lang, and R. Bodik. 2017. Programming
with models: writing statistical algorithms for general model structures with NIMBLE. Journal of
Computational and Graphical Statistics 26:403-413. https://doi.org/10.1080/10618600.2016.1172487
Dénes, F. V., L. F. Silveira, and S. R. Beissinger. 2015. Estimating abundance of unmarked animal
populations: accounting for imperfect detection and other sources of zero inflation. Methods in Ecology
and Evolution 6:543-556. https://doi.org/10.1111/2041-210X.12333
Etterson, M. A., S. N. Ellis-Felege, D. Evers, G. Gauthier, J. A. Grzybowski, B. J. Mattsson, L. R. Nagy, B. J.
Olsen, C. M. Pease, M. Post van der Burg, and A. Potvien. 2011. Modeling fecundity in birds: Conceptual
overview, current models, and considerations for future developments. Ecological Modelling 222:2178-
2190. https://doi.org/10.1016/j.ecolmodel.2010.10.013
Fuller, M. R., J. S. Hatfield, and E. L. Lindquist. 1995. Assessing ground-based counts of nestling bald eagles
in northeastern Minnesota. Wildlife Society Bulletin 23:169-174.
Fuller, M. R., and J. A. Mosher. 1981. Methods of detecting and counting raptors: A review. Studies in Avian
Biology 6:235-246.
Gallego, D., and J. H. Sarasola. 2021. Using drones to reduce human disturbance while monitoring breeding
status of an endangered raptor. Remote Sensing in Ecology and Conservation 7:550-561.
https://doi.org/10.1002/rse2.206
Goszczyński, J. 1997. Density and productivity of Common Buzzard Buteo buteo and Goshawk Accipiter
gentilis populations in Rogów, Central Poland. Acta Ornithologica 32:149–155.
Götmark, F. 1992. The effects of investigator disturbance on nesting birds. Current Ornithology 9:63-104.
Ibáñez-Alamo, J. D., O. Sanllorente, and M. Soler. 2012. The impact of researcher disturbance on nest
predation rates: a meta-analysis. Ibis 154:5-14. https://doi.org/10.1111/j.1474-919X.2011.01186.x
Katzenberger, J., E. Gottschalk, N. Balkenhol, and M. Waltert. 2019. Long-term decline of juvenile survival
in German red kites. Journal of Ornithology 160:337-349. https://doi.org/10.1007/s10336-018-1619-z
Kéry, M., and J. A. Royle. 2015. Applied hierarchical modeling in ecology: Analysis of distribution,
abundance and species richness in R and BUGS. Academic Press, London, UK.
Lahoz-Monfort, J. J., G. Guillera-Arroita, and R. Tingley. 2016. Statistical approaches to account for false-
positive errors in environmental DNA samples. Molecular Ecology Resources 16:673-685.
https://doi.org/10.1111/1755-0998.12486
Lieury, N., S. Devillard, A. Besnard, O. Gimenez, O. Hameau, C. Ponchon, and A. Millon. 2017. Designing
cost-effective capture-recapture surveys for improving the monitoring of survival in bird populations.
Biological Conservation 214:233-241. https://doi.org/10.1016/j.biocon.2017.08.011
Link, W. A., M. R. Schofield, R. J. Barker, and J. R. Sauer. 2018. On the robustness of N-mixture models.
Ecology 99:1547-1551. https://doi.org/10.1002/ecy.2362
MacKenzie, D. I., J. D. Nichols, G. B. Lachman, S. Droege, J. A. Royle, and C. A. Langtimm. 2002. Estimating
site occupancy rates when detection probabilities are less than one. Ecology 83:2248-2255.
https://doi.org/10.1890/0012-9658(2002)083[2248:ESORWD]2.0.CO;2
McKibben, F. E., F. Abadi, and J. K. Frey. 2023. To model or not to model: false positive detection error in
camera surveys. Journal of Wildlife Management 87:e22365. https://doi.org/10.1002/jwmg.22365
Miller, D. A., J. D. Nichols, B. T. McClintock, E. H. C. Grant, L. L. Bailey, and L. A. Weir. 2011. Improving
occupancy estimation when two types of observational error occur: non-detection and species
misidentification. Ecology 92:1422-1428. https://doi.org/10.1890/10-1396.1
Nakashima, Y. 2020. Potentiality and limitations of N-mixture and Royle-Nichols models to estimate animal
abundance based on noninstantaneous point surveys. Population Ecology 62:151-157.
https://doi.org/10.1002/1438-390X.12028
Otis, D. L., K. P. Burnham, G. C. White, and D. R. Anderson. 1978. Statistical-inference from capture data on
closed animal populations. Wildlife Monographs:1-135.

R Core Team. 2022. R: A language and environment for statistical computing. R Foundation for Statistical
Computing, Vienna, Austria. Online: https://www.R-project.org

Royle, J. A. 2004. N-mixture models for estimating population size from spatially replicated counts.
Biometrics 60:108-115. https://doi.org/10.1111/j.0006-341X.2004.00142.x

Royle, J. A., and W. A. Link. 2005. A general class of multinomial mixture models for anuran calling survey
data. Ecology 86:2505-2512. https://doi.org/10.1890/04-1802

Royle, J. A., and W. A. Link. 2006. Generalized site occupancy models allowing for false positive and false
negative   errors.   Ecology   87:835-841.   https://doi.org/10.1890/0012-
9658(2006)87[835:GSOMAF]2.0.CO;2

Sample, C., J. M. Fryxell, J. A. Bieri, P. Federico, J. E. Earl, R. Wiederholt, B. J. Mattsson, D. T. T. Flockhart, S.
Nicol, J. E. Diffendorfer, W. E. Thogmartin, R. A. Erickson, and D. R. Norris. 2018. A general modeling
framework for describing spatially structured population dynamics. Ecol Evol 8:493-508.
https://doi.org/10.1002/ece3.3685

Sen, B., and H. R. Akçakaya. 2022. Fecundity and density dependence can be estimated from mark-
recapture data for making population projections. Ornithological Applications 124, duab064.
https://doi.org/10.1093/ornithapp/duab064

Sergio, F., A. Tanferna, J. Chicano, J. Blas, G. Tavecchia, and F. Hiraldo. 2019. Protected areas under
pressure: decline, redistribution, local eradication and projected extinction of a threatened predator,
the red kite, in Doñana National Park, Spain. Endangered Species Research 38:189-204.
https://doi.org/10.3354/esr00946

Sergio, F., G. Tavecchia, J. Blas, A. Tanferna, and F. Hiraldo. 2021. Demographic modeling to fine-tune
conservation targets: importance of pre-adults for the decline of an endangered raptor. Ecological
Applications:e2266. https://doi.org/10.1002/eap.2266

Sibly, R. M., and J. Hone. 2002. Population growth rate and its determinants: an overview. Philosophical
Transactions   of   the   Royal   Society   B-Biological   Sciences   357:1153-1170.
https://doi.org/10.1098/rstb.2002.1117

Sollmann, R. 2023. rasrage/Sollmann_et_al_Nestling_counts_PCJ: Version 1 (Version v1). Zenodo.
doi.org/10.5281/zenodo.10245803

Sollmann, R., N. Adenot, P. Spakovszky, J. Windt, and B. J. Mattsson. 2023. Appendix 1+2 for 'Accounting
for observation biases associated with counts of young when estimating fecundity: case study on the
arboreal-nesting red kite (Milvus milvus)'. Zenodo. doi.org/10.5281/zenodo.10245398

Tyre, A. J., B. Tenhumberg, S. A. Field, D. Niejalke, K. Parris, and H. P. Possingham. 2003. Improving precision
and reducing bias in biological surveys: Estimating false-negative error rates. Ecological Applications
13:1790-1801. https://doi.org/10.1890/02-5078

White, C., and S. Sherrod. 1973. Advantages and disadvantages of the use of rotor-winged aircraft in raptor
surveys. Raptor Research 7:97-104.

Williams, B. K., J. D. Nichols, and M. J. Conroy. 2002. Analysis and management of animal populations.
Academic press, San Diego, California, USA.

Zhao, Q. 2020. On the sampling design of spatially explicit integrated population models. Methods in
Ecology and Evolution 11:1207-1220. https://doi.org/10.1111/2041-210X.13457

Zipkin, E. F., and S. P. Saunders. 2018. Synthesizing multiple data types for biological conservation using
integrated   population   models.   Biological   Conservation   217:240-250.
https://doi.org/10.1016/j.biocon.2017.10.017