Spatial segregation and aging of metabolic processes underlie phenotypic heterogeneity

in mycobacteria

Authors: Celena M. Gwin[a,*], Kuldeepkumar R. Gupta[a,*], Yao Lu[a], Lin Shao[a], E. Hesper Rego[a,1]

Affiliations

[a] Department of Microbial Pathogenesis, Yale University School of Medicine, New Haven, CT

06519

* equal contribution

[1]To whom correspondence should be addressed: hesper.rego@yale.edu

Individual cells within clonal populations of mycobacteria vary in size, growth rate, and antibiotic

susceptibility. Heterogeneity is, in part, determined by LamA, a protein found exclusively in

mycobacteria. LamA localizes to sites of new cell wall synthesis where it recruits proteins

important for polar growth and establishing asymmetry. Here, we report that in addition to this

function, LamA interacts with complexes involved in oxidative phosphorylation (OXPHOS) at a

subcellular location distinct from cell wall synthesis. Importantly, heterogeneity depends on a

unique extension of the mycobacterial ATP synthase, and LamA mediates the coupling between

ATP production and cell growth in single cells. Strikingly, as single cells age, concentrations of

proteins important for oxidative phosphorylation become less abundant, and older cells rely less

on oxidative phosphorylation for growth. Together, our data reveal that central metabolism is

spatially organized within a single mycobacterium and varies within a genetically identical

population of mycobacteria. Designing therapeutic regimens to account for this heterogeneity may

help to treat mycobacterial infections faster and more completely.

Introduction

For a bacterial infection to linger after antibiotic treatment, only a few bacteria need to remain.

What is different about surviving bacteria, and how do these differences arise? Despite extensive

research, no single model has emerged explaining this phenomenon, and it is likely that the

answers to these questions will vary depending on the bacterial species. In the case of model

bacteria, like *Escherichia coli*, much of the focus has been on stochastic mechanisms that underlie

cells switching into rare drug-tolerant states (*1, 2*). However, stochasticity is just one way of

generating diversity, and other mechanisms of heterogeneity are more deterministic in nature.

For example, clonal populations of mycobacteria, a genus that includes the human pathogen

Mycobacterium tuberculosis, exhibit more variability than model bacterial species (*3*). At least

some of the heterogeneity is created every time a mycobacterium divides when it produces two

cells with different sizes, growth rates, and susceptibilities to antibiotics (*4*). Importantly,

heterogeneity is hard-coded in the genome, as deletion of a single gene unique to mycobacteria

– *lamA* – collapses morphological heterogeneity and leads to fast and uniform killing by several

antibiotics (*5*). LamA localizes to sites of new cell wall synthesis and recruits proteins important

for polar growth (*6*). However, many bacterial species divide asymmetrically but do not exhibit as

much heterogeneity as mycobacteria, suggesting that asymmetric division is only one factor

responsible for creating a heterogeneous population (*7*). Here, we sought to understand the

mechanism by which LamA creates heterogeneity in growth.

Results

A conserved tyrosine regulates the localization and function of LamA

The predicted structure of LamA includes two regions of high disorder separated by a single

transmembrane domain (**Fig. 1A**). The carboxy terminus of the protein is predicted to encode an

MmpS domain. Proteins with these domains are often encoded by operons that include *mmpL*

genes, but LamA's operon partner is a gene of unknown function. The amino terminus of LamA

Figure 1. A conserved phosphorylated tyrosine is important for LamA function and localization.
(A) Schematic of the predicted structure of LamA. **(B)** A multiple sequence alignment of the N-terminal sequence of LamA. Stars indicate residues mutated in panel C. **(C)** Cell size of WT, $\Delta lamA$, or cells expressing the indicated *lamA* allele in $\Delta lamA$. (dark black lines indicate medians; n=100-125 cells for all strains; ****p<0.0001 by one-way ANOVA comparing means to *lamA*_{WT} corrected for multiple comparisons). **(D)** Asymmetry in the daughter cell size at the time of division, measured by the old pole daughter as a fraction of the total size of both daughters. The dotted line marks symmetric division. (n=195, 236, 223 daughter cell pairs for WT, $\Delta lamA$, and *lamA*_{Y50A}, respectively). **(E)** The elongation rate, λ, of the indicated strains. (CV = coefficient of variation, n=239, 226, 267 single cells for WT, $\Delta lamA$, and *lamA*_{Y50A}, respectively). **(F)** Phase contrast and fluorescence images of either msfGFP-LamA or msfGFP-LamA_{Y50A} expressed in single copy from the native promoter. **(G)** Fractional dwell time of strains in panel F calculated by the amount of time a focus spent at the pole before dissipating, normalized to the total time of a fast-interval timelapse (n=15 cells for each strain). **(H)** Intensity profiles were measured across several cells, normalized to cell length, and averaged (n =140-165 cells for each strain). In panels D & E, LamA_{Y50A} refers to the strain mutated on the chromosome in an otherwise wild type background. In all others, *lamA*_{Y50A} is expressed in single copy in $\Delta lamA$.

is highly acidic, with nearly 20% of the residues in this region being either aspartic or glutamic

acid. Additionally, several whole-cell proteomic studies have mapped phosphorylation events at

serine, threonine, and tyrosine residues in this region (*8-13*). We performed a multiple sequence

alignment of LamA proteins found throughout the mycobacterial genus and observed that, despite

the predicted disorder of this region, several tyrosine and serine residues were highly conserved

(**Fig. 1B**). To test the function of these residues, we created phage-integrating plasmids carrying

alanine mutations at each of these sites, transformed these into $\Delta lamA$ *M. smegmatis,* and

imaged the resulting strains by phase contrast microscopy. To quantify the morphology of single

cells, we trained the U-Net machine-learning software package to detect and segment single

mycobacterial cells by phase contrast (*14, 15*). Using this approach, we found that one mutant, a

tyrosine mutated to an alanine at position 50, resulted in cells that were slightly larger than the

other mutants and wild type (**Fig. 1C**). Increased cell size was due to an increase in cell length

rather than width (**Fig. S1**). Importantly, tyrosine 50 is found to be phosphorylated in several

whole-cell proteomic studies (*9-11*). To confirm that this mutation is important at the native locus,

we used marker-less single-strained recombineering (*16*) to recode this tyrosine to an alanine at

the native locus and performed timelapse microscopy. Compared to wild type cells, a

subpopulation of cells expressing $LamA_{Y50A}$ was indeed larger, driving the increased size of cells

across the population (**Fig. S2**). This also led to more heterogeneity in cell size at the time of

division; however, it did not change the asymmetry at division (**Fig. 1D, Fig S2**). Both phenotypes

suggested that $LamA_{Y50A}$ functions similarly to $LamA_{WT}$ or may be a gain-of-function mutant.

However, this mutant was deficient in other functions performed by LamA. Specifically,

LamA also affects the heterogeneity in single-cell growth rate. Biochemical fluctuations within

metabolic pathways result in individual cells growing at slightly different rates centered around a

population average (*17*). To compare growth rate distributions between strains, we relied on the

observation that single *M. smegmatis* cells grow exponentially and computed an average growth

rate, $\lambda = \ln(S_d/S_b)/\Delta T$, where S_d is the size of the cell at division, S_b is the size of the cell at birth,

and ΔT is the time between birth and division (*18*). Wild type *M. smegmatis* displayed 16%

variation in λ, in agreement with prior studies that computed λ by measuring length instead of

area (*18, 19*). In contrast, both $\Delta lamA$ and $LamA_{Y50A}$ cells were less variable than wild type (CV_λ

= 12% and 11%, respectively) (**Fig. 1E**). Computing growth by other metrics resulted in the same

conclusion (**Fig. S3A**). Importantly, expression of $LamA_{WT}$ from an integrative plasmid restored

heterogeneity in $\Delta lamA$ (**Fig. S3B**). Thus, LamA has at least two functions. One establishes

asymmetry in cell size at the time of division; the other is dependent on tyrosine 50 and mediates

heterogeneity in the growth rate of single cells.

87   To understand what was different about LamA$_{Y50A}$, we fused msfGFP to either the wild

type or mutant variant of LamA at their N-termini and expressed these from the native promoter

at a phage integration site in Δ*lamA*. We had previously determined that LamA localized to the

septum, but were unable to visualize membrane localization, possibly due to over-expression of

the fusion construct (*5*). By expressing msfGFP fusions at native levels, we observed that, in

addition to localizing to the septum, msfGFP-LamA$_{WT}$ also localizes along the sides and is

occasionally found at the poles (**Fig. 1E**). Timelapse imaging at 5-minute intervals confirmed that

msfGFP-LamA$_{WT}$ is highly dynamic between the pole and sides of the cell, at a timescale that is

too fast to represent new synthesis and maturation of the fluorescence protein fusion construct

(**Fig. 1F-H**). We next localized msfGFP-LamA$_{Y50A}$ and found its localization to be much less

dynamic, with the protein primarily localized to the poles (**Fig. 1F-H**). Interestingly, the human

pathogen *M. tuberculosis* divides less asymmetrically than *M. smegmatis* but, like *M. smegmatis*,

exhibits a wide variation in single-cell growth rates (*20*). As we have connected LamA to both

asymmetric division and growth rate heterogeneity, we wondered if the *M. tuberculosis* LamA

variant localized differently than *M. smegmatis* LamA. Indeed, we find the *M. tuberculosis* variant

of LamA expressed in *M. smegmatis* is mainly found along the side walls, with less polar

localization than we observe with the *M. smegmatis* variant (**Fig. S4**).

Taken together, these data reveal that the functions of LamA are performed at different

subcellular sites and that LamA dynamically localizes between these sites, possibly in a

phosphorylation-dependent manner. Specifically, we find that a mutant of LamA that cannot be

phosphorylated at a conserved residue is locked at the poles and grows more uniformly. This

suggests that the mechanisms used to create heterogeneity in growth are performed along the

sides of the bacterium. Furthermore, these data suggest that LamA functions at the pole to

establish asymmetric polar growth. In fact, we have recently shown that LamA is important for

recruiting certain proteins to the poles, which are required to establish asymmetry (*6*).

LamA precipitates with proteins involved in oxidative phosphorylation, which are excluded

from the poles and septa

As LamA is performing different functions at distinct subcellular sites, we hypothesized

that it would interact with different proteins depending on its localization. To examine this, we

performed a series of immunoprecipitations to find potential protein-protein interaction partners.

We created a strain in which the sole copy of LamA was fused to the 3X-FLAG epitope. We fixed

these cells with a chemical crosslinker to capture potentially transient interactions, incubated

lysates with α-FLAG beads, and identified co-precipitating peptides by mass spectrometry.

Precipitated peptides largely fell into two categories of proteins – those associated with cell

elongation (MmpL3, PgfA, MurA, and PknA), and those associated with cellular respiration and

oxidative phosphorylation (components of the electron transport chain and ATP synthase

complex) (**Table S1**). To verify these results, we conducted a reciprocal co-immunoprecipitation

with AtpG, the gamma subunit of ATP synthase, since it was highly enriched in multiple biological

replicates. For this, we created strains expressing both AtpG-3XFLAG and LamA-strep and found

that we could precipitate LamA-strep with α-FLAG beads only when AtpG-3XFLAG was present

(**Fig. 2A**).

Perhaps shedding light on these interactions, the operon containing *lamA* is arranged in

the middle of two operons containing many of the genes that encode proteins in the cytochrome

III/IV supercomplex, which performs the last step in the electron transport chain (ETC). This

genomic synteny is conserved across mycobacterial species (**Fig. 2B,C**). To further investigate

the connection between LamA and the ETC, we tracked the localization of QcrB, a subunit of the

III/IV supercomplex, by making an in-frame fusion to msfGFP at the chromosomal locus. We also

Figure 2. LamA interacts with proteins involved in OXPHOS, which are spatially separated from those used to grow the cell.
(A) Strains expressing the indicated epitope-tagged proteins were incubated with α-FLAG beads, and the precipitates were probed with an α-strep antibody via Western blot. **(B)** The genomic region surrounding *lamA* in the indicated organisms. **(C)** The proteins in the ETC that correspond to the genes in panel B. **(D)** Live cells expressing the indicated fusions were imaged by 3D-SIM. A maximum projection is shown. **(E)** Cells expressing AtpG-msfGFP were followed over time by 3D-SIM at 15-minute increments. At each time point single slices from the 3D stacks are shown. Arrows indicate AtpG dynamically infiltrating the tips of cells. **(F)** Cells expressing fusions to QcrB and Wag31 were imaged by live-cell 3D-SIM. A maximum projection is shown. **(G)** A model summarizing the localization of LamA and the complexes involved in polar growth and OXPHOS.

created fusions to AtpG and AtpA, the gamma and alpha subunits of the ATP synthase complex,

respectively. As expected, by conventional fluorescence microscopy, the proteins were localized

to the plasma membrane (**Fig. S5A**). To visualize the localization of these proteins more closely,

we loaded cells into a microfluidic device and performed live-cell three-dimensional structured-

illumination microscopy (3D-SIM), a super-resolution technique. As SIM requires collecting

several images to reconstruct one super-resolved image, we focused on AtpG and QcrB fusions

because these were the brightest. We collected images at single time points and observed that

both proteins were excluded from the tips of the cells as well as the septa of dividing cells (**Fig.**

2D). By timelapse 3D-SIM at 15-minute increments, we occasionally saw AtpG-msfGFP infiltrate

the tip of the cell but primarily remain localized to the sides of the cell (**Fig. 2E**). To visualize the

localization in the context of proteins that direct polar growth, we created a strain that encoded

key polar growth scaffolding protein, Wag31, fused to mKate2 in the background of cells

expressing QcrB-msfGFP and performed two-color live-cell 3D-SIM (**Fig. 2F,G**). We observed

little to no colocalization between these two proteins, showing that the complexes that perform

the last steps of oxidative phosphorylation are spatially distinct from those involved in growing the

bacterium (**Fig. 2F, Fig. S5B**). Moreover, as LamA is localized dynamically between the

poles/septa and sides, these data also suggest that the interaction between LamA and the

electron transport chain is transient and largely confined to the sides, rather than poles, of the

bacterium (**Fig. 2G**).

LamA inhibits oxidative phosphorylation, and reliance on oxidative phosphorylation is

associated with uniformity.

If LamA is working with the respiratory chain, we reasoned that changes to the membrane

potential and cellular ATP levels might reveal if it has a negative or positive regulatory role. We

measured membrane potential by quantifying the accumulation of a charged fluorescence

molecule, TMRM, by flow cytometry (*21, 22*). As a control, we treated cells with a known

protonophore, CCCP, at greater-than-MIC concentrations to depolarize the membrane and

observed the expected decrease in signal (**Fig. 3A**). Compared to wild type, Δ*lamA* was slightly

depolarized (**Fig. 3A**), supporting the idea that LamA is functioning with members of the electron

transport chain. Next, we measured ATP levels using a standard luminescence-based assay and

found that ATP levels were slightly higher in Δ*lamA* cells (**Fig. S6A**). Together, these data suggest

that LamA may be inhibiting the production of ATP via oxidative phosphorylation.

Oxidative phosphorylation is essential

for mycobacteria, but mycobacteria can also

produce ATP in other ways, like substrate-level

phosphorylation when grown on sugars. We

wondered how generalizable the connection

between metabolism and heterogeneity was.

Specifically, we asked if cells grown in

conditions promoting the exclusive use of

oxidative phosphorylation would phenocopy

Δ*lamA*. To test this, we grew cells in defined

minimal media and supplied succinate as the

sole carbon source. Consistent with our

hypothesis, wild type *M. smegmatis* grew

uniformly in minimal succinate medium, and

deletion of *lamA* had no additional effect on

heterogeneity (**Fig. 3B**). *M. smegmatis* cells

cultured in acetate also grow uniformly (*19*),

suggesting this phenomenon is not specific to

succinate-grown cells. To understand if this

was also true in slow-growing mycobacteria,

we followed single *Mycobacterium bovis* BCG

cells for approximately four doublings (~4

days) in both our normal medium (7H9) and

minimal medium supplied with succinate.

Consistent with our findings in *M. smegmatis*,

BCG cells cultured in 7H9 grew more

Figure 3. LamA-mediated heterogeneity is tightly linked to ATP production. (A) Accumulation of TMRM is measured by flow cytometry. As a control, cells were treated for 15 minutes with 500μM of CCCP. The bars represent the medians of three experiments (each with 3 biological replicates (triangles)) assayed on different days. Error bars are 95% confidence intervals. **p<0.002 by one-way ANOVA. **(B)** The indicated strains were grown in either 7H9 or minimal media (HdB for *Msm* or MMAT for *BCG*) supplied with succinate as the sole carbon source and imaged by phase contrast timelapse microscopy to compute λ for single cells. **(C)** Mycobacteria encode an unusual domain in the alpha subunit of the ATP synthase that prevents hydrolysis. We hypothesize that this positively charged extension interacts with the negatively charged cytoplasmic tail of LamA. Cells missing this extension were imaged over time to measure λ. (n = 152, 172, and 149 cells for WT, Δ*lamA,* and *atpAΔC* cells, respectively). **(D)** WT or Δ*lamA* cells expressing the QUEEN-2m ATP biosensor were imaged over time, and the ratio of the two QUEEN excitation wavelengths was used to obtain relative ATP levels. For each cell, the variation within a cell cycle (as measured by the standard deviation of the QUEEN-2m signal divided by its mean) was compared to the exponential growth rate (n = 67 and 59 complete cell cycles for WT and Δ*lamA*, respectively; lines represent linear regressions; *r* is Pearson's correlation coefficient; and, *p* is the *p*-value).

heterogeneously than those fed only succinate (CV_λ = 17% and 13%, respectively) (**Fig. 3B**).

Together, these data show that cells undergoing more oxidative phosphorylation, either through

deletion of *lamA*, or through carbon source availability, grow more uniformly.

An unusual extension on the ATP synthase alpha subunit mediates heterogeneity.

We hypothesized that the interaction between LamA and OXPHOS proteins is important

for creating heterogeneity in the growth rate of single cells. The complexes that perform oxidative

phosphorylation in mycobacteria are comprised of highly conserved proteins, but there are

several unique characteristics specific to mycobacteria and related species (*23*). For instance, in

many organisms, ATP synthase can both synthesize and hydrolyze ATP depending on cellular

conditions. However, in some actinobacteria, including mycobacteria, the alpha subunit of ATP

synthase has a disordered extension (residues V519 – A548) that interacts with the gamma

subunit to prevent the enzyme from hydrolyzing ATP (*24*). The ATP synthase operon is duplicated

in *M. smegmatis (25)*; therefore, we created a strain in which we deleted the extension in both

copies of AtpA (*atpA*$_{\Delta C}$). Consistent with the known function of this extension to block ATP

hydrolysis, *atpA*$_{\Delta C}$ cells have slightly lower ATP levels (**Fig. S6B**).

Because the extension on AtpA is highly positively charged, and the N-terminal extension

of LamA is negatively charged both by amino acid residues and by multiple phosphorylated

residues, we hypothesized that these two proteins might be interacting electrostatically (**Fig. 3C**).

To that end, we reasoned that if LamA interacts with this extension to inhibit ATP synthase, a

condition we have shown is associated with more heterogeneity, then we should expect that

AtpA$_{\Delta C}$ cells would phenocopy $\Delta lamA$ cells with regards to both membrane potential and single-

cell growth heterogeneity. To test membrane potential, we used the same TMRM assay as before

and found that the depolarization in *atpA*$_{\Delta C}$ cells was similar in magnitude and direction as that of

$\Delta lamA$ cells (**Fig. 3A**). Next, we analyzed single-cell growth by phase contrast timelapse

microscopy, and consistent with our hypothesis, *atpA*$_{\Delta C}$ cells exhibit less variability in growth than

wild type cells (**Fig. 3C**), but not division asymmetry **(Fig. S7).** Further experiments will be needed

to determine if LamA and ATP synthase interact directly, a challenging directive as our data

suggests any interaction is likely transient. Nevertheless, we conclude that both LamA and an

actinobacteria-specific feature of the ATP synthase are needed to create phenotypic

heterogeneity within a genetically identical mycobacterial population.

LamA mediates the coupling between ATP fluctuations and growth in single cells.

Cellular ATP levels are often assumed to be uniformly distributed across a population and stable

over the course of a cell cycle. However, fluorescent biosensors that dynamically report on ATP

concentrations in single bacterial cells have revealed that ATP levels vary from cell to cell and

fluctuate dynamically within a cell cycle (*26, 27*). For single cells within a clonal population, large

ATP fluctuations during a cell cycle are associated with slower growth (*28*). While the biological

basis for this phenomenon remains unknown, it offers a potential explanation for phenotypic

heterogeneity in growth rate. Since we show that LamA is connected to ATP generation through

oxidative phosphorylation, we hypothesized that ATP fluctuations would be altered in Δ*lamA* cells.

To assay this, we expressed a codon-optimized version of the QUEEN-2m biosensor in *M.*

smegmatis cells with and without *lamA (26)*. ATP concentration within bacterial cells is linearly

proportional to the ratio of fluorescence excited at 405nm versus 488nm detected by the same

emission bandpass (*26*). Thus, we computed pixel-by-pixel ratios of the fluorescence values

collected at these wavelengths. As in *E. coli (26)*, single-cell measurements revealed a negative

correlation between the amplitude of fluctuation of QUEEN-2m signal (*i.e.* [ATP]) and growth rate

in wild type *M. smegmatis* cells. On average, Δ*lamA* cells displayed a similar magnitude of ATP

fluctuation, but these fluctuations were not correlated with growth rate (**Fig. 3D**). Together, these

data show that LamA mediates the coupling between ATP production and growth rate across a

clonal population of *M. smegmatis* cells.

245 **Mycobacterial single-cell aging is associated with metabolic heterogeneity.**

246   After division, rod-shaped cells inherit a new pole formed from the most recent division

247 event and an old pole that was formed during a prior division event. For mycobacteria, this means

248 that cells inherit growing poles of various ages (**Fig. 4A**). Cells with the oldest poles are born

249 larger but grow more slowly and with more variability (*19, 29-31*). As we have connected growth

250 variability to the production of ATP, we wondered how the abundance of OXPHOS components

251 differed as single cells aged. By timelapse microscopy, we observed that the fluorescence

252 intensity of QcrB-msfGFP (a proxy for the cellular concentration of QcrB) was inherited unevenly

253 at division. On average, "old pole" cells inherited a lower concentration of QcrB than their "new

Figure 4. Metabolic aging in single mycobacterial cells.
(A) Schematic of aging cells and their associated pole ages. **(B)** QcrB-msfGFP fluorescence in the new pole daughter is compared to the fluorescence in the old pole daughter at the time of division in the indicated strains. The dotted line represents equal inheritance. **** p<0.0005 by one-way ANOVA. (n = 187, 191, and 159 sister pairs for WT, ΔlamA, ΔlamA +lamA, respectively). **(C)** QcrB-msfGFP fluorescence was followed over time in old pole daughters as they age. Two representative lineages are shown. **(D)** QcrB-msfGFP fluorescence over multiple generations, coloring corresponds to percent signal normalized to the maximum (n = 10 lineages). The initial mother cell is of unknown age as indicated by the * in panel A. **(E)** Cells grown in a microfluidic device and treated with 50μM CCCP. Size was measured pre- and post-treatment. (From left to right: n = 43, 40, 64, 47 over two different experiments, indicated by the triangles). P<0.05 by one-way ANOVA comparing the means of the triangles.

pole" siblings. This difference was reduced in ∆*lamA,* with a more uniform inheritance of subunits

between sisters (**Fig. 4A**).

We next asked how these differences propagated through the generations in a wild type

population. Thus, we identified the youngest cells in our timelapse data and followed them for at

least two generations. For many old pole daughters, we observed that fluorescence decreased at

time of division and declined steadily over multiple divisions (**Fig. 4C, 4D**). Analyzing multiple

lineages showed that QcrB concentration decreased in the old pole daughter cells by

approximately 25% after the third division (**Fig. S8A,B**). Reconstructing complete lineages

revealed that this phenomenon occurred to a greater or lesser degree depending on the identity

of the mother cell and was most pronounced in cells with the oldest mothers (**Fig. 4D**). We

repeated these measurements with cells expressing AtpG-msfGFP and observed similar trends

(**Fig. S8C,D**). Together, these data suggested that new pole progeny may be performing more

oxidative phosphorylation than older cells. To test this, we inhibited the proton motive force with

the protonophore CCCP and watched the response at the single-cell level. Consistent with our

hypothesis, the growth of new pole cells was affected by CCCP, while old pole cells were largely

unaffected (**Fig. 4E)**. These results phenocopy LamA overexpression (*5*), further supporting the

notion that LamA inhibits oxidative phosphorylation. Together, these data suggest a model

whereby mycobacteria aging at the single-cell level is associated with less reliance on oxidative

phosphorylation. Consequently, in an asynchronous population of mycobacterial cells, flux

through central metabolism varies from cell to cell.

Conclusion

Despite their small size, bacteria encode diverse mechanisms to spatially structure their internal

biochemical processes. For example, many bacteria rely on concentration gradients along their

long axis to spatially organize the macromolecular machines that perform division (*32-35*).

Additionally, at division, the concentration of secondary messengers like cyclic-di-GMP, can be

asymmetrically distributed, an event needed

for the pathogenic lifecycle of *P. aeruginosa*

(36). Our work shows that central metabolic

processes like those that produce energy

can be subcellularly orchestrated in bacteria.

Further work will be needed to understand

the source of the connection between

fluctuations in ATP levels and growth of

single cells, which in mycobacteria is

mediated by LamA. We speculate that the

subcellular utilization of the proton motive

force is an important component of this

correlation. The proton motive force is a key

resource for a bacterial cell – it powers the

molecular machine needed to synthesize

ATP and is used by numerous integral

membrane proteins to transport various

Figure 5. Model for spatial segregation and aging of metabolic processes in mycobacteria.
(A) In mycobacteria, the complexes that perform oxidative phosphorylation are physically separated from those used to grow the bacterium. Both complexes rely on the proton motive force to transport molecules across the plasma membrane. LamA dynamically interacts with both complexes and is important for coupling ATP production to growth rate in single cells. **(B)** As single mycobacterial cells age, they rely less on oxidative phosphorylation for growth. Thus, in a genetically identical population of mycobacteria cells, the flux through central metabolic pathways differs from cell to cell.

substrates, including molecules that make up the cell envelope, across the plasma membrane. In

bacteria that grow by incorporating new envelope along their sides, complexes that use the proton

motive force to grow and make ATP spatially intermingle. In contrast, we show that these two

processes are spatially distinct in mycobacteria, with the proteins needed for oxidative

phosphorylation found along the sides of the bacterium and certain pumps like MmpL3 and MurJ

mainly localized at the poles *(37-40)* (**Fig. 5A**). Furthermore, our data suggests that LamA tra-

nsiently and stochastically interacts with both pathways. Supporting this observation are the

localizations of the likely kinase and phosphatase involved. PknA and PknB, essential kinases

involved in cell growth and division, are predicted to phosphorylate LamA and localize to the poles

(*41, 42*); in contrast, the only known protein phosphatase in mycobacteria, PstP, localizes along

the side walls and to the septum (*43, 44*). These data suggest that LamA could be constantly in

motion to generate asymmetry and heterogeneity in growth and division.

Mycobacteria can simultaneously catabolize different carbon sources, a feature that is

important for the survival of pathogenic mycobacteria which rely on a mixture of fatty acids,

cholesterol, and carbohydrates at different points during infection (*45-47*). Our data show that the

metabolism of mycobacteria is even more unusual than previously recognized, as a rapidly

growing population can diversify its metabolism at the single-cell level (**Fig. 5B**). While *M.*

tuberculosis is thought to mainly rely on cholesterol early in infection, multiple lines of evidence

suggest that this species uses glycolysis at later stages of infection when the bacterial burden is

high (*48, 49*). Our data suggest that this may also lead to metabolic and morphological

heterogeneity that may be critical for the lifecycle of the pathogen. Specifically, our model

suggests that inhibition of multiple metabolic subpopulations may help kill mycobacterial

populations faster and more completely. Indeed, simultaneous inhibition of oxidative

phosphorylation and glycolysis leads to rapid and complete sterilization of *M. tuberculosis (50)*.

Thus, our data add to the growing body of evidence that understanding - and accounting for - the

complexity of mycobacterial physiology at the single-cell level may be the key to improving TB

therapy (*51-55*).

References

1. J. N. Carey *et al.*, Regulated Stochasticity in a Bacterial Signaling Network Permits
Tolerance to a Rapid Environmental Change. *Cell* **173**, 196-207.e114 (2018).
2. N. Q. Balaban, J. Merrin, R. Chait, L. Kowalik, S. Leibler, Bacterial Persistence as a
Phenotypic Switch. *Science* **305**, 1622-1625 (2004).
3. E. S. Chung, W. C. Johnson, B. B. Aldridge, Types and functions of heterogeneity in
mycobacteria. *Nature Reviews Microbiology* **20**, 529-541 (2022).
4. B. B. Aldridge *et al.*, Asymmetry and Aging of Mycobacterial Cells Lead to Variable
Growth and Antibiotic Susceptibility. *Science* **335**, 100-104 (2012).
5. E. H. Rego, R. E. Audette, E. J. Rubin, Deletion of a mycobacterial divisome factor
collapses single-cell phenotypic heterogeneity. *Nature* **546**, 153-157 (2017).

6.   K. R. Gupta *et al.*, An essential periplasmic protein coordinates lipid trafficking and is required for asymmetric polar growth in mycobacteria. *eLife* **11**, e80395 (2022).

7.   M. Campos *et al.*, A constant size extension drives bacterial cell size homeostasis. *Cell* **159**, 1433-1446 (2014).

8.   J. Zeng *et al.*, Protein kinases PknA and PknB independently and coordinately regulate essential Mycobacterium tuberculosis physiologies and antimicrobial susceptibility. *PLoS Pathog* **16**, e1008452 (2020).

9.   S. Fortuin *et al.*, Phosphoproteomics analysis of a clinical Mycobacterium tuberculosis Beijing isolate: expanding the mycobacterial phosphoproteome catalog. *Front Microbiol* **6**, 6 (2015).

10.   K. C. Nakedi *et al.*, Identification of Novel Physiological Substrates of Mycobacterium bovis BCG Protein Kinase G (PknG) by Label-free Quantitative Phosphoproteomics. *Mol Cell Proteomics* **17**, 1365-1377 (2018).

11.   S. Prisic *et al.*, Extensive phosphorylation with overlapping specificity by Mycobacterium tuberculosis serine/threonine protein kinases. *Proceedings of the National Academy of Sciences* **107**, 7521-7526 (2010).

12.   R. Verma *et al.*, Quantitative Proteomic and Phosphoproteomic Analysis of H37Ra and H37Rv Strains of Mycobacterium tuberculosis. *Journal of Proteome Research* **16**, 1632-1645 (2017).

13.   U. Kusebauch *et al.*, *Mycobacterium tuberculosis* supports protein tyrosine phosphorylation. *Proceedings of the National Academy of Sciences* **111**, 9265-9270 (2014).

14.   T. Falk *et al.*, U-Net: deep learning for cell counting, detection, and morphometry. *Nature Methods* **16**, 67-70 (2019).

15.   S. Berg *et al.*, ilastik: interactive machine learning for (bio)image analysis. *Nature Methods* **16**, 1226-1232 (2019).

16.   K. C. Murphy. (Springer US, 2021), pp. 301-321.

17.   D. J. Kiviet *et al.*, Stochasticity of metabolism and growth at the single-cell level. *Nature* **514**, 376-379 (2014).

18.   M. M. Logsdon *et al.*, A Parallel Adder Coordinates Mycobacterial Cell-Cycle Progression and Cell-Size Homeostasis in the Context of Asymmetric Growth and Organization. *Current biology : CB* **27**, 3367-3374.e3367 (2017).

19.   M. Priestman, P. Thomas, B. D. Robertson, V. Shahrezaei, Mycobacteria Modify Their Cell Size Control under Sub-Optimal Carbon Sources. *Front Cell Dev Biol* **5**, 64 (2017).

20.   E. S. Chung, P. Kar, M. Kamkaew, A. Amir, B. B. Aldridge, Mycobacterium tuberculosis grows linearly at the single-cell level with larger variability than model organisms. *bioRxiv*, (2023).

21.   J. M. Benarroch, M. Asally, The Microbiologist's Guide to Membrane Potential Dynamics. *Trends in Microbiology* **28**, 304-314 (2020).

22.   J. M. Kralj, D. R. Hochbaum, A. D. Douglass, A. E. Cohen, Electrical Spiking in Escherichia coli Probed with a Fluorescent Voltage-Indicating Protein. *Science* **333**, 345-348 (2011).

23.   H. Guo *et al.*, Structure of mycobacterial ATP synthase bound to the tuberculosis drug bedaquiline. *Nature* **589**, 143-147 (2021).

24.   C. F. Wong *et al.*, Structural Elements Involved in ATP Hydrolysis Inhibition and ATP Synthesis of Tuberculosis and Nontuberculous Mycobacterial F-ATP Synthase Decipher New Targets for Inhibitors. *Antimicrobial Agents and Chemotherapy* **66**, e01056-01022 (2022).

25.   M. G. Montgomery, J. Petri, T. E. Spikes, J. E. Walker, Structure of the ATP synthase from Mycobacterium smegmatis provides targets for treating tuberculosis. *Proceedings of the National Academy of Sciences* **118**, e2111899118 (2021).

26. H. Yaginuma *et al.*, Diversity in ATP concentrations in a single bacterial cell population
revealed by quantitative single-cell imaging. *Scientific Reports* **4**, 6522 (2014).
27. S. Manuse *et al.*, Bacterial persisters are a stochastically formed subpopulation of low-
energy cells. *PLOS Biology* **19**, e3001194 (2021).
28. W.-H. Lin, C. Jacobs-Wagner, Connecting single-cell ATP dynamics to overflow
metabolism, cell growth, and the cell cycle in Escherichia coli. *Current Biology* **32**, 3911-
3924.e3914 (2022).
29. K. Ginda *et al.*, The studies of ParA and ParB dynamics reveal asymmetry of
chromosome segregation in mycobacteria. *Molecular Microbiology* **105**, 453-468 (2017).
30. M. M. Logsdon *et al.*, A Parallel Adder Coordinates Mycobacterial Cell-Cycle
Progression and Cell-Size Homeostasis in the Context of Asymmetric Growth and
Organization. *Current Biology* **27**, 3367-3374.e3367 (2017).
31. M. T. M. Hannebelle *et al.*, A biphasic growth model for cell pole elongation in
mycobacteria. *Nature Communications* **11**, (2020).
32. Y. E. Chen *et al.*, Spatial gradient of protein phosphorylation underlies replicative
asymmetry in a bacterium. *Proceedings of the National Academy of Sciences* **108**, 1052-
1057 (2011).
33. M. Thanbichler, L. Shapiro, MipZ, a Spatial Regulator Coordinating Chromosome
Segregation with Cell Division in Caulobacter. *Cell* **126**, 147-162 (2006).
34. S. Kretschmer, P. Schwille, Pattern formation on membranes and its role in bacterial cell
division. *Current Opinion in Cell Biology* **38**, 52-59 (2016).
35. D. Kiekebusch, M. Thanbichler, Spatiotemporal organization of microbial cells by protein
concentration gradients. *Trends in Microbiology* **22**, 65-73 (2014).
36. M. Christen *et al.*, Asymmetrical Distribution of the Second Messenger c-di-GMP upon
Bacterial Cell Division. *Science* **328**, 1295-1297 (2010).
37. C. Carel *et al.*, Mycobacterium tuberculosis Proteins Involved in Mycolic Acid Synthesis
and Transport Localize Dynamically to the Old Growing Pole and Septum. *PLoS ONE* **9**,
e97148 (2014).
38. J. Zhu *et al.*, Spatiotemporal localization of proteins in mycobacteria. *Cell Reports* **37**,
110154 (2021).
39. C. L. Gee *et al.*, A Phosphorylated Pseudokinase Complex Controls Cell Wall Synthesis
in Mycobacteria. *Science Signaling* **5**, ra7-ra7 (2012).
40. L. Thouvenel, J. Rech, C. Guilhot, J.-Y. Bouet, C. Chalut, In vivo imaging of MmpL
transporters reveals distinct subcellular locations for export of mycolic acids and non-
essential trehalose polyphleates in the mycobacterial outer membrane. *Scientific
Reports* **13**, (2023).
41. M. Mir *et al.*, The extracytoplasmic domain of the Mycobacterium tuberculosis Ser/Thr
kinase PknB binds specific muropeptides and is required for PknB localization. *PLoS
Pathog* **7**, e1002182 (2011).
42. S. N. Nagarajan *et al.*, Protein kinase A (PknA) of Mycobacterium tuberculosis is
independently activated and is critical for growth in vitro and survival of the pathogen in
the host. *J Biol Chem* **290**, 9626-9645 (2015).
43. Iswahyudi *et al.*, Mycobacterial phosphatase PstP regulates global serine threonine
phosphorylation and cell division. *Scientific Reports* **9**, (2019).
44. F. Shamma, E. H. Rego, C. C. Boutte, Mycobacterial serine/threonine phosphatase
<scp>PstP</scp> is phosphoregulated and localized to mediate control of cell wall
metabolism. *Molecular Microbiology* **118**, 47-60 (2022).
45. A. D. Baughn, K. Y. Rhee, Metabolomics of Central Carbon Metabolism in
Mycobacterium tuberculosis. *Microbiology Spectrum* **2**,
10.1128/microbiolspec.mgm1122-0026-2013 (2014).

46.   L. P. de Carvalho *et al.*, Metabolomics of Mycobacterium tuberculosis reveals
compartmentalized co-catabolism of carbon substrates. *Chem Biol* **17**, 1122-1131
(2010).
47.   Dany J. V. Beste *et al.*, 13C-Flux Spectral Analysis of Host-Pathogen Metabolism
Reveals a Mixed Diet for Intracellular Mycobacterium tuberculosis. *Chemistry & Biology*
20, 1012-1021 (2013).
48.   J. Marrero, C. Trujillo, K. Y. Rhee, S. Ehrt, Glucose phosphorylation is required for
Mycobacterium tuberculosis persistence in mice. *PLoS Pathog* **9**, e1003116 (2013).
49.   W. Y. Phong *et al.*, Characterization of phosphofructokinase activity in Mycobacterium
tuberculosis reveals that a functional glycolytic carbon flow is necessary to limit the
accumulation of toxic metabolic intermediates under hypoxia. *PLoS One* **8**, e56037
(2013).
50.   J. S. Mackenzie *et al.*, Bedaquiline reprograms central metabolism to reveal glycolytic
vulnerability in Mycobacterium tuberculosis. *Nat Commun* **11**, 6092 (2020).
51.   C. Toniolo, O. Rutschmann, J. D. McKinney, Do chance encounters between
heterogeneous cells shape the outcome of tuberculosis infections? *Current Opinion in*
Microbiology **59**, 72-78 (2021).
52.   A. M. Cadena, S. M. Fortune, J. L. Flynn, Heterogeneity in tuberculosis. *Nature*
reviews.Immunology, (2017).
53.   P. L. Lin *et al.*, Sterilization of granulomas is common in active and latent tuberculosis
despite within-host variability in bacterial killing. *Nature medicine* **20**, 75-79 (2014).
54.   Q. Liu *et al.*, Tuberculosis treatment failure associated with evolution of antibiotic
resilience. *Science* **378**, 1111-1118 (2022).
55.   M. Mistretta, N. Gangneux, G. Manina, Microfluidic dose-response platform to track the
dynamics of drug response in single mycobacterial cells. *Sci Rep* **12**, 19578 (2022).
462
463

Acknowledgments. We thank members of the Rego lab for thoughtful suggestions and

Dr. Andy Goodman for critical reading of the manuscript.

Funding: This work was supported by the Burroughs Wellcome Fund, the Searle

Foundation, the Pew Foundation, and National Institute of Health Grant R01AI148255 to

E.H.R. We also thank Yale Flow Cytometry for their assistance with BD Symphony

analyzer service. The Core is supported in part by an NCI Cancer Center Support Grant

NIH P30 CA016359. The BD Symphony was funded by shared instrument grant # NIH

S10 OD026996. We also thank the MS & Proteomics Resource at Yale University for

providing the necessary mass spectrometers and the accompanying biotechnology tools

funded in part by the Yale School of Medicine and by the Office of The Director, National

Institutes of Health (S10OD02365101A1, S10OD019967, and S10OD018034). The

funders had no role in study design, data collection and analysis, decision to publish, or

preparation of the manuscript.

Author contributions: Conceptualization: C.M.G., K.R.G, and E.H.R. Investigation:

C.M.G., K.R.G, and E.H.R. Resources. C.M.G., K.R.G, Y.L., L.S., and E.H.R. Software:

L.S. Supervision: E.H.R. Funding acquisition: E.H.R. Writing - original draft: C.M.G.,

K.R.G, and E.H.R. Writing – review and editing: C.M.G., K.R.G, Y.L., L.S., and E.H.R

Competing interests: The authors declare no competing interests.

Data and materials availability: Strains developed during the study will be made

available from E.H.R.

486