Strain-specific evolution and host-specific regulation of transposable elements in the model plant symbiont *Rhizophagus irregularis*

Jordana Inácio Nascimento Oliveira[1], Nicolas Corradi[1*]

[1]Department of Biology, Faculty of Sciences, University of Ottawa, Ottawa, ON, Canada

[*]Corresponding author ncorradi@uottawa.ca

Abstract

Transposable elements (TE) are repetitive DNA that can create variability in genome structure and regulation. The genome of *Rhizophagus irregularis*, a widely studied arbuscular mycorrhizal fungus (AMF), comprises approximately 50% repetitive sequences that include transposable elements (TE). Despite their abundance, two-thirds of TE remain unclassified, and their regulation among AMF life-stages remains unknown. Here, we aimed to improve our understanding of TE diversity and regulation in this model species by curating repeat datasets obtained from chromosome-level assemblies and by investigating their expression across multiple conditions. Our analyses uncovered new TE superfamilies and families in this model symbiont and revealed significant differences in how these sequences evolve both within and between *R. irregularis* strains. With this curated TE annotation, we detected that the number of upregulated TE families in colonized roots is four times higher than extraradical mycelium, and their overall expression differs depending on the plant host. This work provides a fine-scale view of TE diversity and evolution in model plant symbionts and highlights their transcriptional dynamism and specificity during host-microbe interactions. We also provide Hidden Markov Model profiles of TE domains for future manual curation of uncharacterized sequences (https://github.com/jordana-olive/TE-manual-curation/tree/main).

Key words

Introduction

Arbuscular mycorrhizal fungi (AMF) are an ancient group of plant symbionts capable of colonizing thousands of different species. The AMF provides vitamins and minerals from the soil to the plants and receives carbohydrates and lipids in return [1]. This relationship may have been established since plants first conquered the land [2], and there is evidence showing that AMF improve plant growth and ecosystem productivity [3]. More recently, it was also reported that these symbionts can retain atmospheric carbon, indicating these play a significant role in the global climate balance [4,5]. Nonetheless, despite their relevance for plant fitness and long-term evolution, AMF show low morphological variability with no apparent plant specificity [6].

In addition to being ecologically and agriculturally important, AMF harbours peculiar cellular features. Their spores and hyphae always carry thousands nuclei, and to date, no stage where one or two nuclei co-exist in one cell has ever been observed [7]. Furthermore, despite their longevity, no sexual reproduction has been formally observed in these organisms. However, evidence has now shown that AMF strains can either carry one or two nuclear genotypes in their cells, a genetic characteristic found in sexually multinucleated strains of ascomycetes and basidiomycetes, suggesting that sexual reproduction does exist in these prominent symbionts [8,9].

The low morphological diversity of AMF masks remarkable differences in gene content and structure in this group. For example, species vary significantly in genome size and gene counts [10], and the variation in genome size is significantly correlated with the abundance of transposable elements (TE) [11]. Recent studies based on chromosome-level assemblies of *Rhizophagus irregularis,* also showed this model species carries two different (A/B) compartments, which are reminiscent of the "two-speed genome" structure previously reported in plant pathogens [9,12]. In AMF, the A-compartment carries most core genes and shows significantly higher gene expression, while the B-compartment has a higher density in repetitive DNA, as well as in secreted proteins and candidate effectors involved in the molecular dialogues between the partners of the mycorrhizal symbiosis [12].

TE are sparse repetitive DNA classified into retrotransposons (Class I), which use RNA molecules as intermediates for "copy-and-paste", and transposons (Class II), which

spread through "cutting-and-pasting" the DNA [13]. Each class is divided into orders and

superfamilies based on pathways of transposition and phylogenetic relationships [14,15], and

thus classifying TE is important to describe the evolution of the genome and infer its

impact on the biology of any organism [16,17]. For example, by modifying chromatin status

and attracting transcript factors, these elements can also promote regulation of the gene

expression [18].

In the model AMF *R. irregularis,* over 50% of the genome is composed by TE

[9,12,19]. Their higher abundance in the B-compartment is linked to higher rates of

rearrangements [9], and it was proposed that elevated TE expression in germinating spores

may lead to new expansions of TE in this AMF species [20]. Despite recent findings, key

questions regarding the diversity and evolution of TE, as well as their role in mycorrhizal

interactions, remain unanswered. For example, approximately two thirds of TE remain

unclassified, making it difficult to infer their function in AMF genome biology and

evolution [20,21]. Similarly, because analyses of TE expression have so far centred on

germinating spores, it is unknown how these elements are controlled during host

colonisation, and whether some show host-specific regulation. The present study

addresses these questions by providing an improved, curated classification of TE families

in all *R. irregularis* strains with chromosome-level assemblies and by investigating their

expression among multiple hosts.

Material and methods

Curation and classification of TEs families

We used chromosome level assemblies of five homokaryotic (4401, A1, B3, C2 and

DAOM197198) [12] and four heterokaryotic strains (A4, A5, G1 and SL1) [9]

Supplementary material S1) as a source to build repeat libraries. The manual curation

for nonmodel species followed the most recommended guides [16,22]. Firstly, the repeat

libraries were generated using RepeatModeler2.0.3 [22] with -LTRstruct mode for detecting

Long Terminal Repeats sequences implemented by LTRharvest and LTR_retriever. The

libraries from all strains were merged to create a single reference for the manual curation.

The unique library was submitted to TEclass which separates the sequences into

the order level: nonLTR, LTR or DNA [23]. This step helped us to distinguish orders with

similar protein domains, such as DIRS (nonLTR) and Crypton (DNA). Tirvish, from genome-tools tools (http://genometools.org/), was used to detect Terminal Inverted Repeats (TIR) in DNA transposons elements (for order DNA/TIR) with the parameter -mintirlength 8. Hidden Markov Models (hmm) of specific TE superfamilies or orders domains were generated from a combination of conserved regions described in **Supplementary material S2**. The sequences for each domain were first aligned using mafft [24], converted to stockholm format using esl-reformat and finally submitted to hmmbuild (version 3.1b2) to generate the hmm profiles [25].

Lastly, we provided hmm profiles for detecting elements with reverse transcriptase (LINE, DIRS, PLE, LTR, Bel, Copia, Gypsy), specific transposase superfamilies (Academ, CMC, Ginger, KDZ, Kolobok, MULE, Merlin, Novsib, P, PIF-Harbinger, PiggyBac, Plavaka, Sola-1, Sola-2, Sola-3, Tc-Mariner, Transib, Zator and hAT) and tyrosine recombinase (DIRS and Crypton) (https://github.com/jordana-olive/TE-manual-curation/tree/main/TE-domains). The open reading frames (ORF) from the reference library were generated using the getorf with 200 amino acids as minimum size (https://www.bioinformatics.nl/cgi-bin/emboss/help/getorf). The hmmrsearch (version 3.1b2) was used to find the sequences with TE domains from above step. We removed sequences with expanded genes domains (Sel1, BTB, Kelch, Protein Kinase, TPR), the models are available at https://github.com/jordana-olive/TE-manual-curation/tree/main/expanded-genes.

A transposon (DNA/TIRS) was kept in the final library when detected as DNA by TEclass, presents a transposon domain and TIR sequence, with a size between 1kb to 17kb. Sequences with TIRS, ranging between 50 to 1kb and lacking transposase domains were classified as MITEs. Due to the similarity and closest relationship of certain elements, after the hmmrsearch, we constructed a phylogenetic tree with the putative LINE, DIRS, PLE, LTR, Ginger, Crypton and Maverick sequences. The protein sequences of these above-mentioned orders were aligned using mafft [24], then submitted to RAxML [26] for generating the phylogenetic tree, using PROTGAMMA model with 1000 of bootstraps. The best tree resolution was visualized using ggtree in R [27]. The final classification of these sequences was based on the relationships (**Supplementary Figure S1**).

In summary, a known sequence (with classification from RepeatModeler) was
kept if correctly detected by TEclass and presence of a respective transposition domain.
A new sequence was classified according to TEclass order, presence of a transposition
domain and its relationship to a known sequence according to the phylogenetic tree. The
final library, and models are available on https://github.com/jordana-olive/TE-manual-
curation/tree/main and can be reproduced in any other reference to custom TE
characterization.

Repeat landscapes of genomes and compartments

We ran the RepeatMasker (version 4.1.2-p1) [28] for all strains, with the parameters -a -s -
lib using the curated library. The repeat landscapes were generated from modified
createlandscape.pl and calculedivergence.pl scripts, provided in RepeatMasker files. The
landscapes also were generated to available A and B-compartment [9,12]. We applied a
paired t-test on R of the TE percentages along Kimura divergence points in A and B-
compartments annotation. A variation between the landscape was considered significative
when $p < 0.05$.

Transposable element and gene expression analysis

We evaluate the expression of genes and TEs using available RNAseq from different
tissues (germinating spores [20], intraradical mycelium, arbuscules [29] and extraradical
mycelium [30]), and mycorrhized roots from different plant hosts colonized by
DAOM197198 (*Allium schoenoprasum*, *Medicago truncatula*, *Nicotiana benthamiana* [29]
and *Brachypodium distachyon* [31]). The accesses to the data are described in
Supplementary material S3. The reads were filtered using Trimmomatic [32] and aligned
to DAOM197198 reference genome using Bowtie2 [33]. The read count was accessed by
TEtranscripts [34] guided by the TE annotation performed in this study and gene annotation
executed by [12]. Using DESeq2 [35], for each host condition and tissue, the differential
expression was generated comparing germinated spores as control. A transcript was
considered differentially expressed when padj (adjusted p-value) is equal or lower than
0.05.

Transposable element nearby gene correlation

For local copy expression, we used available RNAseq generated through Oxford
Nanopore Technology (ONT) sequencing [19]. The long ONT-RNAseq reads were filtered
and trimmed using pychopper (https://github.com/epi2me-labs/pychopper) (7.98% did
not pass the quality parameters and were discarded). The nucleotide correction was
performed based on self-clustering using isONTcorrect [36]. The filtered and corrected
reads were aligned to DAOM197198 genome using hisat2 and annotated using stringtie
[37] guided by TE annotation performed in this study and gene annotation executed by [12].
In the same way, stringtie generated the counts of the transcripts, used in the expression
analysis. For detecting TEs upstream of genes, we extracted the genomic regions up to -
1000 to the transcript start position and then intersected with TE annotations using
bedtools [38]. The correlation function based on the Pearson method in R was used to
determine the association between genes and their TE upstream pairs.

Results

A curated database reveals new TE families in <u>Rhizophagus irregularis</u>

Using RepeatModeler and RepeatMasker, recent analyses of *R. irregularis* chromosome-
level datasets indicated that strains of this species carry an average of 50% of TE [9,12].
However, only one third of their repeat content could be classified, and thus on average
30% of all available genomes are composed by unclassified TE sequences [9,12,19,21]. To
address this, we used five chromosome-assembly level from homokaryon and four
heterokaryon strains to generate curated repeat libraries. When all genomes are
considered, out of total of 9,257 TE sequences identified by RepeatModeler, only 2,369
(approx. 25%) can be considered well defined, *bona-fide* non-redundant consensus
sequences harboring transposition domains. The notable reduction in TE numbers is due
to non-curated datasets containing highly degenerated TE without domains (relics), and
repeats that cannot be classified with current knowledge of TE evolution.

In our final curated library, 1,458 sequences belong to families previously
classified by RepeatModeler (K, **Figure 1a**), while 636 sequences represent newly
identified families (N, **Figure 1a**) of the following orders; SINE, LINE, LTR,
DNA/TIRS, DNA/Crypton, Maverick and RC/Helitron orders (**Figure 1a**).

Using the curated library to annotate TE in the DAOM-197198 genome,
sequences belonging to these orders increased in number by respectively 4-fold on
average, and similar results were obtained for all investigated *R. irregularis* strains
(**Figure 1b**). We also assessed the influence of the manually curated library on the repeat
landscape of the DAOM197198 genome (see **Figures 1c-e**). Following TE annotation,
the percentage of the DAOM197198 genome is represented by classified TE increased
from 12% (**Figure 1c-d**) to 36% (**Figure 1e**).

Figure 1 – TE curated library description. **a** Proportion of known (detected by RepearModeler, in light color, K) and novel (in bold colors, N) curated sequences. **b** RepeatModeler and curated library annotation comparison across the strains. The *y-axis* represents the *R. irregularis* strains, while the *x-axis* shows the relative abundance of the TE in the genome from the homology-based annotation (RepeatModeler, left) and the current annotation (manual curation, right).

The newly curated *R. irregularis* TE library now includes novel families of non-
LTR retrotransposons: SINE, LINE/CR1-Zenon, LINE/L1-Tx1, LINE/L1-R1, LINE/R2-
Hero, LINE/RTE-BovB, LINE/I, which were detected based on phylogenetic analyses
(**Supplementary Figure S1**). A novel superfamily of DNA transposons – e.g., Transib, -

and novel families belonging to known AMF TE superfamilies were also detected,

including Crypton, CMC, hAT-19, MULE, Sola-2, Tc1-Mariner and Plavaka (**Figure 2**).

The Plavaka family, which is part of CACTA/CMC/EnSpm superfamily, is

particularly prevalent in DAOM197198, A1, B3, C2, A5 and G1. This family has already

been identified in fungi [39,40], but is not deposited in publicly available repeat databases,

which is why the RepeatModeler could not detect this family.

Figure 2 – Annotation of TEs families in *R. irregularis* strains. The heatmaps show the percentage families in each strain considering the RepeatMasker (**a**) and the curated library of known and novel TE families (**b**). Blank areas mean absence of certain family in that annotation, while the black rectangles highlight the new families found during the curation.

In the *R. irregularis* isolate DAOM197198, TE are differently mobilized and retained
across the genome [20], but we found these patterns can differ among *R. irregularis* strains
with chromosome-level assemblies. Specifically, in most of strains TE expansions are
recent, as highlighted by the high number of elements with Kimura substitutions ranging
from 0 to 5, supporting the findings that new TE expansion bursts exceed older TE
insertions in DAOM197198 [20] (**Figure 3**). However, other strains can have different TE
distributions. For example, the strains A4 and SL1 carry a larger proportion of TE with
older Kimura substitution levels, indicating strain-specific patterns of TE emergence,
evolution, and retention. Remarkably, the genome of C2 has a much larger proportion of
younger, and thus likely more active TE, and it is plausible that this feature is linked to
the larger genome size of this strain compared to relatives - i.e., the C2 genome is 162Mb
compared to an average of 147Mb for other *R. irregularis* strains (**Figure 3**).

Figure 3 – Repeat landscape of *R. irregularis* strains. **a-h** Each plot shows the sequence divergence from its consensus (*x-axis*) in relation to the number of copies in the genome (*y-axis*). Newer insertions are shown in the left peaks, while the older and more degenerate insertions are on the right side of the graph. The strains with recent burst of TE expansions are indicated with the green arrow, while the ones representing early TE degeneration are highlighted with red arrow.

The distribution of TE based on Kimura substitution levels also differ
significantly between A and B-compartment in all strains (**Figure 4**). Specifically, the
proportion of TE with Kimura substitutions between 10 and 20 – i.e., older TE - is always
higher in the B-compartment (p<0.05), indicating that these elements are maintained over
time at higher levels in this portion of the genome. In contrast, old and degenerated TE
appear to be more rapidly eliminated in the A-compartment (**Figure 4**).

TE and gene expression is positively correlated within the B-compartment

The localization of TE nearby genes can impact their expression [18,41]. We tested if the
presence of TE upstream of genes is linked with gene expression in the compartments by
identifying TE located 1000 nucleotides upstream of genes using available long reads
RNA data [19]. We found no correlation between the expression of TE and coding genes
located in A-compartment (r=0.21, p<0.0001) (**Figures 4f**), however, a positive and
significant correlation exists for genes and TE in the B-compartment (r=0.84, p<0.0001)
(**Figures 4g**). This finding suggests that TE and genes in B-compartment are being co-
expressed.

Figure 4 – **a-e** A and B-compartments repeat landscapes in *R. irregularis*. The arrows point to the
divergence between the compartments. **f-g** Coding genes and upstream TE correlation of expression. Each
point represents a coding gene-upstream TE pair, where the *y-axis* is the TE expression (logTPM) and the
x-axis is the coding gene expression (logTPM). The green line represents the regression, followed by the
confidence interval in light grey.

Transposable elements are significantly more expressed in colonized roots compared to
extraradical mycelium

To obtain additional insights into the biology of TE, we investigated their regulation
during host colonization using *R. irregularis* DAOM197198 RNA-seq data from multiple
tissues, including micro dissected cells of arbuscules (ARB) and intraradical mycelium
(IRM), and extraradical mycelium (ERM) (**Figure 5a**) in symbiosis with *Medicago*
truncatula roots (**Figures 5b-d**).

Figure 5 – Expression of TE in *R. irregularis* through life-stages of colonization. **a** Schematic view of the different life stages compared to spores. ERM, extraradical mycelium; IRM, intraradical mycelium, ARB; arbuscules. **b-d** Volcano plots of expressed TE in each colonization stage, evidencing the number of down and upregulated sequences, in blue and red, respectively. The ones expressed but not significative different from spores are shown in black as NO. **f** AGO protein gene expression under different conditions. Colonization stages collected by laser dissection of ARB and IRM; ERM and mycorrhized roots. The t-test was performed comparing all the conditions with spores, where: * $p < 0.05$; ** $p < 0.001$ and ns is no significance. The box limits represent the range of 50% of the data, the central line marks the median, and the vertical lines out of the boxes are the upper and lower values. normTPM – normalized transcripts per million.

246   The number of upregulated TE is more than four times higher in IRM and ARB
247   than in extraradical mycelium in the same host (**Figure 5b-d**). TE significantly
248   upregulated in colonized tissue (ARB and IRM) include DNA/TIRS, LTR, LINE and
249   SINES orders, among them, LTR/Gypsy, LINE, CMC-EnSpm, Plavaka, hAT-Tag1,

hAT-Ac and Helitron are the superfamilies with more expressed families (**Supplementary Material S4**).

One mechanism to control TE mobilization is through RNAi and AGO proteins [42]. The *R. irregularis* DAOM197198 genome contains 25 AGO domain proteins [43], and we found that ERM samples express significantly more (p < 0.05) AGO genes compared to the other conditions (**Figure 5b** and **f**). In stark contrast, arbuscules and intraradical mycelium laser dissected cells have significantly reduced AGO expression, which again differs from the significantly higher expression of TEs in these conditions (p < 0.05) (**Figure 5b** and **f**).

TE regulation changes with different hosts and correlates with host genome size

Analyses of RNA-seq data roots from multiple hosts colonized by DAOM197198 (*Allium schoenoprasum*, *Medicago truncatula*, *Nicotiana benthamiana* and *Brachypodium distachyon*), also reveals that TE expression differs significantly among hosts, with some families being expressed only in one of four hosts investigated (**Figure 6** and **Supplementary Material S4**). For example, a total of 404 families are upregulated during colonization with *A. schoenoprasum* roots (**Figure 6a**), including both families from Class II elements (e.g., Sola-1, hATm, Academ-1, DIRS) and Class I elements (e.g. CR1-Zenon and DIRS) that are only upregulated with this host. In *N. benthamiana* and *M. truncatula* roots, the symbiont upregulates a smaller number of TE families compared to *A. schoenoprasum*, 251 and 197 respectively, while *B. distachyon* was the condition with lowest TE upregulation overall (116 families) (**Figure 6d**).

The differences in TE expression among hosts are not linked to variability in expression of AMF AGO genes – i.e., hosts with highest TE expression do not always have low AGO expression and vice-versa. However, a significant and positive correlation ($r = 0.91$, $p = 0.001$) exist between the repeat-content of the host genome and number of overexpressed families in the DAOM197198 symbiont. For example, *A. schoenoprasum* is the host with most TE and his colonized roots have the highest AMF TE upregulation. In contrast, *B. distachyon* (272Mb) has the lowest TE content and has significantly lower TE upregulation in the symbiont.

Figure 6 – Differential expression of *TE* in *R. irregularis* during symbiosis with different plant hosts *compared to germinated spores*. **a-d** Volcano plots showing the down and upregulated TE in relation to spores, in blue and red respectively, considering padj <0.05. In black are the ones with no significant difference in expression in relation to the control. **e** Number of overexpressed TE families in each host. **f** AGO protein gene expression under colonization with different hosts. The t-test was performed comparing all the conditions with spores, where: * p < 0.05, * p < 0.001 and ns is no significance. The box limits represent the range of 50% of the data, the central line marks the median, and the vertical lines out of the boxes are the upper and lower values. normTPM – normalized transcripts per million. **g** Correlation between the repeated content in host genome (in base pairs) and the number of families overexpressed by the AMF during the symbiosis. Brachy – *Brachypodium.*

Discussion

An improved view of TE family diversity and evolution in a model plant symbiont

Through manual curation of *R. irregularis* repeat libraries, we first improved the proportion of annotated families in these model plant symbionts and uncovered novel sequences representing the largest proportion of their repetitive sequences. Our work also revealed how *R. irregularis* strains differ in how they maintain or remove TE over time within two genome compartments. For example, our findings indicate that strains with largest genome sizes (C2) derive from a combination of a higher rate of TE emergence and retention of these elements. We also uncovered notable differences in how TE evolve

within each strain, with some (A1, B3) having have much higher proportions of very young TE compared to others (SL1, A4) that carry similar levels of Kimura substitution rates indicative of early repeat degeneration and fewer recent expansion.

TE retention rates are different between compartments

By investigating the degree at which TE accumulate mutations over time, a significant distinction emerged between A/B compartments. Specifically, all A-compartment landscapes show a continuing invasion of novel/young TE insertions, and the high methylation present in this compartment [12], and/or purifying selection, might be needed for their control and rapid removal; as evidenced by the lower TE density in this compartment. In contrast, these insertions accumulate in the B-compartment, leading to a notable inflation of these elements over time, as shown by the stable TE density along the axis that defines the Kimura substitution rates. The accumulation of TE in the B-compartment might be linked with their domestication [44,45] and/or the emergence of new functions, a view supported by the significant positive correlations we observed in expression of TE and genes and similar findings from plant pathogens [46,47].

TE regulation and evolution further underpin similarities between AMF and known fungal pathogens

Obvious similarities between the genomes of AMF and those of plant pathogens have been known for some time [48,49]. These include enrichments in transposable elements, and genomes subdividing into highly diverging regions dense in effector genes and TE, and more conserved ones composed of core genes and low repeat density.

In the plant pathogen *Verticilium dahiliae,* TE often locate in proximity to highly expressed pathogenicity-related genes within fast evolving adaptive genomic regions [50]. These regions are reminiscent of *R. irregularis* B-compartments [12,49], which are also enriched in TE and secreted proteins that promote symbiosis with different hosts [51]. As such, the significant correlation we observed between TE and genes specific to the B-compartment may mirror identical processes in AMF and plant pathogens.

The increased upregulation of genes in close proximity to TE during the colonization stages in the B-compartment could also indicates a TE control for de-

repressing those regions of the genome [52]. Indeed, it has been proposed that TE-effectors

regulation is well-timed; i.e. both are expressed during the infection and repressed in the

absence of the host [47]. In filamentous fungi, the variability of TE can also allow for

escaping mechanisms of recognition by the plant immunity system [46,47], and it is thus

possible that in AMF the expression of transposable elements could aid plant-symbiont

communication during colonization.

What drives TE upregulation and host-specificity during colonization?

TE expression is active in germinating spores [20], but our study shows their expression in

colonized roots is much higher still. Within this context, we found that the AGO genes,

which are known TE regulators [53], are significantly more expressed in extraradical

mycelium compared to colonized roots [43], suggesting RNAi is one of the key factors

implicated in the regulation of TE across stages of the mycorrhizal symbiosis -i.e.

downregulation in extraradical mycelium, and upregulation *in planta*.

Notably, AMF AGO expression did not vary significantly among hosts, and thus

other factors could be responsible for the host-specific variation in the TE expression we

observed. One intriguing possibility is that TE expression in the hosts directly or

indirectly influences TE expression in the AMF, as seen in multiple plant-microbe

interactions in cross-talk regulations [54–57]. In support of this view, we found a positive

correlation between TE abundance in the host and TE expression in AMF. With mounting

evidence of molecular cross-communication between the mycorrhizal partners, including

RNA from the fungi interacting with mRNA from the hosts [43,58,59], it is likely that

molecular dialogues between hosts also result in the increased TE expression we observed

in the fungal symbiont.

Acknowledgements

Our research is funded by the Discovery program of the Natural Sciences and Engineering

Research Council (RGPIN2020-05643) and a Discovery Accelerator Supplements

Program (RGPAS-2020-00033). NC is a University of Ottawa Research Chair in

Microbial Genomics. JINO is founded by Mitacs Accelerate Program (IT16902) and a

Discovery Accelerator Supplements Program (RGPAS-2020-00033).

Competing interests

The authors declare no competing interests.

Author contributions

JINO and NC designed the study. JINO performed the experiments and analysis of the

data. JINO and NC wrote the paper. NC mentored and supervised all the processes.

ORCID

Jordana Inácio Nascimento Oliveira https://orcid.org/0000-0003-2511-1746

Nicolas Corradi https://orcid.org/0000-0002-7932-7932

Data availability

The genomes and RNAseq used in this study are described in **Supplementary material**

S1 and **S3**. The ONT RNAseq can be accessed at SRR21968700.

Supporting information

Supplementary file S1 – Chromosome level assemblies of *R. irregularis* strains used in

this study.

Supplementary file S2 – Domains used to construct the hmm models.

Supplementary file S3 – RNAseq data and alignments stats used in this study.

Supplementary file S4 – Number of upregulated TE in each condition.

Supplementary figure S1 – Phylogenetic tree of TE from different orders with similar

domains. Figure in high resolution can be downloaded here https://github.com/jordana-

olive/TE-manual-curation/blob/main/Supplementary-figure-S1.png.

References

1.   Martin, F. M., Uroz, S. & Barker, D. G. Ancestral alliances: Plant mutualistic

symbioses with fungi and bacteria. *Science* **356**, eaad4501 (2017).

2.   Delaux, P.-M. & Schornack, S. Plant evolution driven by interactions with symbiotic
and pathogenic microbes. *Science* **371**, (2021).

3.   Bonfante, P. & Anca, I.-A. Plants, mycorrhizal fungi, and bacteria: a network of
interactions. *Annu. Rev. Microbiol.* **63**, 363–383 (2009).

4.   Horsch, C. C. A., Antunes, P. M., Fahey, C., Grandy, A. S. & Kallenbach, C. M.
Trait-based assembly of arbuscular mycorrhizal fungal communities determines soil
carbon formation and retention. *New Phytol.* **239**, 311–324 (2023).

5.   Hawkins, H.-J. *et al.* Mycorrhizal mycelium as a global carbon pool. *Curr. Biol.* **33**,
R560–R573 (2023).

6.   Corradi, N. & Brachmann, A. Fungal Mating in the Most Widespread Plant
Symbionts? *Trends Plant Sci.* **22**, 175–183 (2017).

7.   Kokkoris, V., Stefani, F., Dalpé, Y., Dettman, J. & Corradi, N. Nuclear Dynamics in
the Arbuscular Mycorrhizal Fungi. *Trends Plant Sci.* **25**, 765–778 (2020).

8.   Ropars, J. *et al.* Evidence for the sexual origin of heterokaryosis in arbuscular
mycorrhizal fungi. *Nature Microbiology* **1**, (2016).

9.   Sperschneider, J. *et al.* Arbuscular mycorrhizal fungi heterokaryons have two nuclear
populations with distinct roles in host-plant interactions. *Nat. Microbiol.* (2023)
doi:10.1038/s41564-023-01495-8.

10. Morin, E. *et al.* Comparative genomics of Rhizophagus irregularis, R. cerebriforme,
R. diaphanus and Gigaspora rosea highlights specific genetic features in
Glomeromycotina. *New Phytol.* **222**, 1584–1598 (2019).

11. Mathu, M. C. *et al.* The early branching arbuscular mycorrhizal fungus Paraglomus
occultum carries a small and repeat poor genome compared to relatives in the
Glomeromycotina. *Microbial Genomics* (2022) doi:10.1099/mgen.0.000810.

12. Yildirir, G. *et al.* Long reads and Hi-C sequencing illuminate the two-compartment
genome of the model arbuscular mycorrhizal symbiont Rhizophagus irregularis. *New*
Phytol. **233**, 1097–1107 (2022).

13. Wicker, T. *et al.* A unified classification system for eukaryotic transposable
elements. *Nat. Rev. Genet.* **8**, 973–982 (2007).

14. Seberg, O. & Petersen, G. A unified classification system for eukaryotic transposable
elements should reflect their phylogeny. *Nature reviews. Genetics* vol. 10 276
(2009).

15. Wicker, T. *et al.* Reply: A unified classification system for eukaryotic transposable
elements should reflect their phylogeny. *Nat. Rev. Genet.* **9**, 411–412 (2008).

16. Wells, J. N. & Feschotte, C. A Field Guide to Eukaryotic Transposable Elements.
Annu. Rev. Genet. **54**, 539–561 (2020).

17. Rech, G. E. *et al.* Population-scale long-read sequencing uncovers transposable
elements associated with gene expression variation and adaptive signatures in
Drosophila. *Nat. Commun.* **13**, 1948 (2022).

18. Bourque, G. *et al.* Ten things you should know about transposable elements. *Genome
Biol.* **19**, 199 (2018).

19. Manley, B. F. *et al.* A highly contiguous genome assembly reveals sources of
genomic novelty in the symbiotic fungus Rhizophagus irregularis. *G3 (Bethesda)* **13**,
(2023).

20. Dallaire, A. *et al.* Transcriptional activity and epigenetic regulation of transposable
elements in the symbiotic fungus Rhizophagus irregularis. *Genome Res.* **31**, 2290–
2302 (2021).

21. Miyauchi, S. *et al.* Large-scale genome sequencing of mycorrhizal fungi provides
insights into the early evolution of symbiotic traits. *Nat. Commun.* **11**, 5125 (2020).

22. Flynn, J. M. *et al.* RepeatModeler2 for automated genomic discovery of transposable
element families. *Proceedings of the National Academy of Sciences* **117**, 9451–9457
(2020).

23. Abrusán, G., Grundmann, N., DeMester, L. & Makalowski, W. TEclass—a tool for
automated classification of unknown eukaryotic transposable elements.
Bioinformatics **25**, 1329–1330 (2009).

24. Nakamura, T., Yamada, K. D., Tomii, K. & Katoh, K. Parallelization of MAFFT for
large-scale multiple sequence alignments. *Bioinformatics* **34**, 2490–2492 (2018).

25. Eddy, S. HMMER user's guide. *Department of Genetics, Washington University*
School of Medicine **2**, 13 (1992).

26. Stamatakis, A. RAxML version 8: A tool for phylogenetic analysis and post-analysis
of large phylogenies. *Bioinformatics* **30**, 28–36 (2014).

27. Yu, G., Smith, D. K., Zhu, H., Guan, Y. & Lam, T. T.-Y. Ggtree : An r package for
visualization and annotation of phylogenetic trees with their covariates and other
associated data. *Methods Ecol. Evol.* **8**, 28–36 (2017).

28. Smit, A. F. A., Hubley, R. & Green, P. RepeatMasker.
https://www.repeatmasker.org/ (1996).

29. Zeng, T. *et al.* Host- and stage-dependent secretome of the arbuscular mycorrhizal
fungus Rhizophagus irregularis. *Plant J.* **94**, 411–425 (2018).

30. Tsuzuki, S., Handa, Y., Takeda, N. & Kawaguchi, M. Strigolactone-Induced Putative
Secreted Protein 1 Is Required for the Establishment of Symbiosis by the Arbuscular
Mycorrhizal Fungus Rhizophagus irregularis. *Mol. Plant. Microbe. Interact.* **29**,
277–286 (2016).

31. Kamel, L. *et al.* The Comparison of Expressed Candidate Secreted Proteins from
Two Arbuscular Mycorrhizal Fungi Unravels Common and Specific Molecular
Tools to Invade Different Host Plants. *Front. Plant Sci.* **8**, 124 (2017).

32. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

33. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat.*
Methods **9**, 357–359 (2012).

34. Jin, Y., Tam, O. H., Paniagua, E. & Hammell, M. TEtranscripts: a package for
including transposable elements in differential expression analysis of RNA-seq
datasets. *Bioinformatics* **31**, 3593–3599 (2015).

35. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

36. Sahlin, K. & Medvedev, P. Error correction enables use of Oxford Nanopore
technology for reference-free transcriptome analysis. *Nat. Commun.* **12**, 2 (2021).

37. Pertea, M. *et al.* StringTie enables improved reconstruction of a transcriptome from
RNA-seq reads. *Nat. Biotechnol.* **33**, 290–295 (2015).

38. Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing
genomic features. *Bioinformatics* **26**, 841–842 (2010).

39. Iyer, L. M. *et al.* Lineage-specific expansions of TET/JBP genes and a new class of
DNA transposons shape fungal genomic and epigenetic landscapes. *Proc. Natl.*
Acad. Sci. U. S. A. **111**, 1676–1683 (2014).

40. Muszewska, A., Steczkiewicz, K., Stepniewska-Dziubinska, M. & Ginalski, K.
Transposable elements contribute to fungal genes and impact fungal lifestyle. *Sci.*
Rep. **9**, 4307 (2019).

41. Drongitis, D., Aniello, F., Fucci, L. & Donizetti, A. Roles of Transposable Elements
in the Different Layers of Gene Expression Regulation. *Int. J. Mol. Sci.* **20**, (2019).

42. Kelleher, E. S., Azevedo, R. B. R. & Zheng, Y. The Evolution of Small-RNA-
Mediated Silencing of an Invading Transposable Element. *Genome Biol. Evol.* **10**,
3038–3057 (2018).

43. Silvestri, A. *et al.* In silico analysis of fungal small RNA accumulation reveals
putative plant mRNA targets in the symbiosis between an arbuscular mycorrhizal
fungus and its host plant. *BMC Genomics* **20**, 169 (2019).

44. Zhang, Y. *et al.* Transposon molecular domestication and the evolution of the RAG
recombinase. *Nature* **569**, 79–84 (2019).

45. Modzelewski, A. J., Gan Chong, J., Wang, T. & He, L. Mammalian genome
innovation through transposon domestication. *Nat. Cell Biol.* **24**, 1332–1340 (2022).

46. Fouché, S. *et al.* Stress-Driven Transposable Element De-repression Dynamics and
Virulence Evolution in a Fungal Pathogen. *Mol. Biol. Evol.* **37**, 221–239 (2019).

47. Fouché, S., Oggenfuss, U., Chanclud, E. & Croll, D. A devil's bargain with
transposable elements in plant pathogens. *Trends Genet.* **38**, 222–230 (2022).

48. Mathieu, S., Cusant, L., Roux, C. & Corradi, N. Arbuscular mycorrhizal fungi:
intraspecific diversity and pangenomes. *New Phytol.* **220**, 1129–1134 (2018).

49. Reinhardt, D., Roux, C., Corradi, N. & Di Pietro, A. Lineage-Specific Genes and
Cryptic Sex: Parallels and Differences between Arbuscular Mycorrhizal Fungi and
Fungal Pathogens. *Trends Plant Sci.* **26**, 111–123 (2021).

50. Torres, D. E., Thomma, B. P. H. J. & Seidl, M. F. Transposable Elements Contribute
to Genome Dynamics and Gene Expression Variation in the Fungal Plant Pathogen
Verticillium dahliae. *Genome Biol. Evol.* **13**, evab135 (2021).

51. Teulet, A. *et al.* A pathogen effector FOLD diversified in symbiotic fungi. *bioRxiv*
(2022) doi:10.1101/2022.12.16.520752.

52. Oggenfuss, U. & Croll, D. Recent transposable element bursts are associated with
the proximity to genes in a fungal plant pathogen. *PLoS Pathog.* **19**, e1011130
(2023).

53. Thomson, D. W. *et al.* Assessing the gene regulatory properties of Argonaute-bound
small RNAs of diverse genomic origin. *Nucleic Acids Res.* **43**, 470–481 (2015).

54. Melayah, D., Bonnivard, E., Chalhoub, B., Audeon, C. & Grandbastien, M. A. The
mobility of the tobacco Tnt1 retrotransposon correlates with its transcriptional
activation by fungal factors. *Plant J.* **28**, 159–168 (2001).

55. Weiberg, A. *et al.* Fungal small RNAs suppress plant immunity by hijacking host
RNA interference pathways. *Science* **342**, 118–123 (2013).

56. Ren, B., Wang, X., Duan, J. & Ma, J. Rhizobial tRNA-derived small RNAs are signal
molecules regulating plant nodulation. *Science* **365**, 919–922 (2019).

57. Wong-Bajracharya, J. *et al.* The ectomycorrhizal fungus Pisolithus microcarpus
encodes a microRNA involved in cross-kingdom gene silencing during symbiosis.
Proc. Natl. Acad. Sci. U. S. A. **119**, (2022).

58. Ledford, W. C. *et al.* A journey into the world of small RNAs in the arbuscular
mycorrhizal symbiosis. *New Phytol.* (2023) doi:10.1111/nph.19394.

59. Silvestri, A. *et al.* Different Genetic Sources Contribute to the Small RNA Population
in the Arbuscular Mycorrhizal Fungus Gigaspora margarita. *Front. Microbiol.* **11**,
(2020).