Getting there in one piece: The Rac pathway prevents cell fragmentation in a nonprotrusively migrating leader cell during organogenesis

Noor Singh[1], Karen Jian Li[1], Kacy Lynn Gordon[1,2]

[1]Department of Biology, The University of North Carolina at Chapel Hill, Chapel Hill, North Carolina 27599
[2]To whom correspondence should be addressed, kacy.gordon@unc.edu

Keywords: Rac, GTPase, gonad, leader cell, migration, fragmentation, germline, stem cell niche

SUMMARY

The *C. elegans* hermaphrodite distal tip cell (DTC) leads gonadogenesis. Loss-of-function mutations in a *C. elegans* ortholog of the Rac1 GTPase (*ced-10*) and its GEF complex (*ced-5*/DOCK180, *ced-2*/CrkII, *ced-12*/ELMO) cause gonad migration defects related to directional sensing; we discovered an additional defect class of gonad bifurcation in these mutants. Using genetic approaches, tissue-specific and whole-body RNAi, and *in vivo* imaging of endogenously tagged proteins and marked cells, we find that loss of Rac1 or its regulators causes the DTC to fragment as it migrates. Both products of fragmentation—the now-smaller DTC and the membranous patch of cellular material—localize important stem cell niche signaling (LAG-2/DSL ligand) and migration (INA-1/integrin subunit alpha) factors to their membranes, but only one retains the DTC nucleus and therefore the ability to maintain gene expression over time. The enucleate patch can lead a bifurcating branch off the gonad arm that grows through germ cell proliferation. Germ cells in this branch differentiate as the patch loses LAG-2 expression. While the nucleus is surprisingly dispensable for aspects of leader cell function, it is required for stem cell niche activity long-term. Prior work found that *Rac1[-/-];Rac2[-/-]* mouse erythrocytes fragment; in this context, our new findings support the conclusion that maintaining a cohesive but deformable cell is a conserved function of this important cytoskeletal regulator.

INTRODUCTION

The formation of each of the two arms of the *C. elegans* hermaphrodite gonad is led by a post-embryonic migration of a somatic gonad leader cell, the distal tip cell (DTC)[1–3]. DTCs begin migrating–nonprotrusively[4]–in opposite directions from the site of the future vulva along the ventral midline at the L2 larval stage. In the L3 larval stage, each DTC executes two 90-degree turns that establish the gonad bend. DTC migration continues along the dorsal body wall during the L3 and L4 larval stages until the two DTCs both reach the dorsal midbody, where they remain throughout adulthood[2,5,6]. Proliferating germ cells provide a propulsive force to the DTC, which interacts with the extracellular matrix to steer the direction of migration[4].

The distal-most end of the adult germline—capped by the DTC—is composed of germ stem cells followed proximally by their still-mitotic descendants, together comprising the proliferative zone[7]. The DTC acts as the germ stem cell niche by expressing the ligand LAG-2[8], which is transduced by the GLP-1/Notch receptor in germ cells. Active Notch signaling in the germline leads to post-transcriptional repression of meiotic entry factors to prevent differentiation[9]. Proximal to the proliferative zone, germ cells lose active Notch signaling, lose the repression of meiotic entry factors, enter meiosis, and eventually differentiate as gametes[10–12]. Hermaphrodites make sperm cells in the L4 larval stage and oocytes as adults[13,14]. An essential feature of the adult hermaphrodite germline is that some meiotic cells do not form gametes, but instead take on a nurse-cell-like role in which they donate their cytoplasmic contents to oocytes and undergo apoptosis, becoming engulfed by the somatic gonadal sheath cells that wrap around the germline[15,16].

Apoptosis and engulfment of dead cells are regulated by highly conserved factors, first identified
in a *C. elegans* genetic screen for animals with abnormal cell death (*ced* mutants)[17–25].
Engulfing cells activate two partially redundant *ced* pathways[26], one of which has also been
shown to also be required for DTC migration[20–23]. This pathway is composed of CED-
10/Rac1[22]–a regulator of the cytoskeleton[27] conserved among eukaryotes[28]–and factors that
have since been discovered to comprise its upstream activating guanine nucleotide exchange
factor (GEF) complex: CED-5/DOCK180[29–31], CED-2/CrkII[22], CED-12/ELMO[20,21,30,31]. (CED-12
has also recently been discovered to have GAP function in the context of F-actin nucleation in
C. elegans embryonic morphogenesis[32]). Rac GTPases, members of the Rho family of
GTPases, are required for cytoskeletal remodeling in protrusions at the leading edge of many
migratory cells[33,34]. However, the migratory DTC has a smooth leading edge[4,35,36], both
distinguishing it from the classic model of leader cell migration[37–39] and raising the question of
how the Rac-pathway *ced* module–a canonical regulator of lamellipodia–regulates migration in
the non-protrusive cell leading gonadogenesis. The DTC may serve as a useful model for the
molecular regulation of tubular organs that form with migration of non-protrusive leader cells.
One way these Rac pathway *ced* genes regulate DTC migration is as downstream effectors of
noncanonical Wnt signaling[40,41], culminating in repression of the netrin receptor UNC-5 during
DTC dorsal migration[40].
70
Genetic control of gonadogenesis is well-studied[1,42,43], and many mutants with polarity
defects[40,41,44] and other gonad malformations have been identified[22,45–51]. Genetic studies of the
three *C. elegans* Rac-family GTPase-encoding genes *ced-10*[22], *rac-2*[52], and *mig-2*[53] have
revealed that two play a role in DTC migration[22,52,54]. *mig-2*[47,54] and *ced-10*[22] cause DTC
migration defects when mutated. *ced-10* and *rac-2* are nearly identical at the sequence level[52]
and act in different subsets of cell types (*rac-2* is not expressed in the DTC[55]). How DTC-
migration genes act at a cellular level to coordinate the cell movements of organogenesis is an
area of increasing interest. Recent work using live-imaging techniques reveals the key role of
interactions between the DTC and extracellular matrix, mediated, for example, by integrin[4]
(which in *C. elegans* is composed of INA-1/PAT-3 and PAT-2/PAT-3 dimers[56–58]). We set out to
investigate a previously unreported class of gonad defect—bifurcation—in genetic mutants that
have long been known to have DTC migration defects.
83
Here we report that loss-of-function mutations in the Rac-pathway *ced* genes lead to the
fragmentation of the DTC as it migrates. Both products of fragmentation—the distal tip cell
(reduced in size) and the membranous patch of cellular material—localize important migration
(an integrin alpha subunit, INA-1) and signaling (the Notch ligand LAG-2) factors to their plasma
membrane. However, only one structure retains the DTC nucleus and therefore the ability to
maintain gene expression and synthesize new protein throughout adulthood. The patch can
promote local germ cell proliferation for a time, leading to a bifurcating branch off the gonad arm
that grows through germ cell proliferation until the patch loses LAG-2 signal due to its lack of
nucleus. Cell fragmentation and changes in deformability have been documented in mammalian
cell types after Rac1 loss-of-function[59–61]; we conclude that Rac1 is required to keep the DTC in
one piece during migration, and that the Rac pathway plays a fundamental and broadly-
conserved function in maintaining cellular integrity in a number of Eukaryotic cell types.
96
RESULTS
Loss of function of Rac-pathway *ced* genes causes gonad bifurcation
Mutations of the cell death pathway genes *ced-10/*Rac1, *ced-5/*DOCK180, *ced-2/*CrkII, and *ced-*
*12/*ELMO have long been observed to cause gonad migration defects[20–22,29,52]. We examined
ced-10(n1993)[22]*, ced-5(n1812)*[29]*, ced-2(n1994)*[22]*,* and *ced-12(n3261)*[21] mutants, all of which are
putative null with the exception of *ced-10(n1993),* a hypomorph[22,52].

Typically, each gonad arm forms a single U-shaped, blind tube[1,6]. In all four mutants, we
observed gonad bifurcation (Figure 1) which was not identified in the original or subsequent
reports of gonad defects caused by mutations in *ced* genes (Gumienny et al., 2001, wrote of a
different *ced-12(oz167)* allele: "*ced-12* mutants also occasionally show …short branching or
distensions"). At the L4 (Figure 1B) and young adult (Figure 1C) stages, we consistently
observe bifurcation in 10-16% of animals (Figure 1D). While misdirected DTC migration or
turning defects have been investigated in mutants for these[40] and other genes[35], gonad
bifurcation is strikingly rare[62–64]. We decided to focus on this most severe phenotype to elucidate
additional roles of Rac-pathway *ced* genes in regulating DTC migration.
112
The Rac pathway acts cell-autonomously to regulate DTC leader cell activity, and loss of
function causes somatic gonad cell defects as well as gonad anatomy defects
To determine whether gonad bifurcation is caused by cell-autonomous loss of the Rac pathway
ced genes, we used a strain with DTC-specific sensitivity to RNAi and a distal tip cell membrane
marker[65] (see Methods and Figure 2A) to knock down *ced-10, ced-5, ced-2,* and *ced-12* in the
DTC. Indeed, *ced* gene knockdown in the DTC causes cell-autonomous defects that result in
bifurcation (Figure 2B). We next examined the DTC itself in RNAi-treated animals. We made
two striking observations. First, when the gonad bifurcates, both tips have a focus of
mNeonGreen (mNG) signal (Figure 2B), in contrast to the single focus of mNG signal in the
control RNAi-treated animals (Figure 2A). Second, even when the gonad does not bifurcate,
there is a second patch of mNG signal near the bend of the gonad in 17-29% of L4 larval
animals (Figure 2B'). Similar defects were also observed after RNAi of *mig-2*/RhoG, another
small Rac-family GTPase, but were not observed after knockdown of the parallel engulfment
pathway members *ced-1* and *ced-6* (Figure S1), in agreement with previously observed
specificity of the Rac pathway *ced* genes, but not other *ced* genes, for DTC migration[29,52].
128
We next examined the DTC in the mutant animals. A cell membrane marker that is specific for
the DTC (*cpIs122[lag-2p::mNeonGreen::PLC$^{\delta PH}$]*[65] shows a single locus of expression in each
gonad arm in otherwise wild-type animals (Figure 2A'). In c*ed-10(n1993), ced-5(n1812), ced-*
2(n1994), and *ced-12(n3261)* mutants, this marker reveals the presence of a patch of mNG
fluorescence on both tips of bifurcated gonads (Figure 2C) and near the bend in gonads that do
not bifurcate (Figure 2C'). The second site of mNG expression appears with high penetrance
across mutants (57-80% of mutant animals, N=107), and is always present when we see gonad
bifurcation (N=14/14 bifurcated mutant gonads with the DTC marker). Put another way, every
bifurcated gonad we observed has a second mNG+ patch, but not every gonad with an ectopic
mNG+ patch bifurcates. On the other hand, we did not observe a patch every time we observed
DTC pathfinding defects, suggesting these two defects reflect different aspects of Rac pathway
function in the DTC. We therefore conclude that Rac-pathway loss-of-function causes the
formation of this mNG+ patch, and hypothesize that the patch sometimes causes gonad
bifurcation.
143
Somatic gonad defects appear during or just after DTC turning
We next sought the origin of the ectopic mNG+ patch. We examined c*ed-10(n1993), ced-*
5(n1812), ced-2(n1994), and *ced-12(n3261)* mutants expressing the *cpIs122[lag-*
2p::mNeonGreen::PLC$^{\delta PH}$][65] cell membrane marker during larval development. Early in gonad
migration (Phase I: ventral migration), a single site of mNG is visible at the growing tip of each
gonad arm in wild-type and mutant gonads (Figure 3). As the DTC makes its turns or just after
(Phase II: turning), a second mNG+ patch appears in the mutant gonads, attached to the DTC
by a filament. Because the patch is connected to the DTC at its first appearance, we
hypothesize that the DTC is the source of the patch. Patches are observed in all four mutants by
Phase III: dorsal migration.

The DTC fragments after Rac-pathway *ced* loss of function

We tested several hypotheses about how the DTC might give rise to the patch. A bifurcated gonad phenotype has been observed after RNAi treatments that interfere with the cell cycle[63,64], which cause the normally postmitotic DTC to divide aberrantly, or another somatic gonad cell to be mis-specified as an extra DTC[62,66]. If the second focus of mNG is caused by aberrant cell division or misspecification of another gonad cell, the "patch" would be a second cell with DTC identity. We also considered that overexpression of membrane fluorescent transgene-encoded proteins could cause abnormal cell growth[67]; in that case, strains without such a transgene should not form a patch. Finally, we considered that the DTC breaks apart without dividing, and the patch is a membranous fragment of cellular material.

We first tested whether the patch was or was not a second distal tip cell. We examined a strain expressing an allele encoding endogenously tagged GFP::HLH-2 protein[68], a tagged transcription factor that localizes to the nucleus of the DTC (among other cells) in control gonads, with some dim cytoplasmic DTC fluorescence signal also observed (Figure 4A). We exposed the strain bearing this allele to *ced-5* and *ced-12* RNAi (as these were the most efficient RNAi treatments for causing gonad bifurcation) and observed GFP::HLH-2 expression in bifurcated gonads. If we detected a bright focus of nuclear signal of this tagged protein in both bifurcated gonad tips, we would conclude that the DTC divided or a second somatic gonad cell became mis-specified as a DTC. However, in 19/19 animals with bifurcated gonads, we observe nuclear localization in only one of the two structures at the tips of the bifurcated gonads and dim cytoplasmic expression in the other (Figure 4B-C). Of note, the tip containing the nucleus of the DTC is not always the tip closest to the correct anatomical position of the DTC (Figure S2). Because GFP::HLH-2 is not encoded by an overexpressed transgene, nor does it encode a membrane-localized fluorescent protein, this experiment also rules out the possibility that driving an overexpressed fluorescent membrane protein in the DTC is responsible for the formation of the structure at the second tip.

We observed the same pattern in a strain coexpressing a *lag-2(bmd202[lag-2::P2A::H2B::mT2])* histone tag[69] and *lag-2p::mNG* membrane marker (*lag-2p::mNeonGreen::PLC$^{\delta PH}$*). In an otherwise wild-type genetic background, the DTC membrane and nucleus are the only sites of expression of these markers in the distal gonad (Figure 4D). After crossing these markers into a *ced-5(n1812)* background, nuclear expression is visible in only one of the two membranous bodies (Figure 4E).

We used this same experiment to quantify the sizes of the DTC and enucleate patch appearing in *ced-5(n1812)* mutant animals coexpressing the *lag-2::P2A::H2B::mT2* histone tag and the *lag-2p::mNG* membrane marker vs. that of the otherwise wild-type DTC in L4 animals. After fragmentation, *ced-5* mutant DTCs are smaller in their projected area than control DTCs, with the patches smaller still (Figure 4F). However, the summed sizes of the *ced-5* mutant DTC and patch are not significantly different from the sizes of wild-type L4 DTCs, suggesting that we are seeing two pieces of a fragmented DTC (Figure 4F). We conclude that the ectopic patch is not a daughter of the DTC nor a mis-specified second DTC in the affected gonad arm—it is a cell fragment. This led us to ask how a fragment of cellular material can be competent to lead gonad outgrowth.

Both fragments of the DTC localize integrin, required for DTC migration, throughout larval development

We next sought to characterize the molecular repertoire of the structures that result from DTC fragmentation. We hypothesized that the cell fragment must localize key regulators of both DTC leader cell and stem cell niche functions in order to lead to the bifurcation of the gonad. Integrin,

a receptor that mediates cell interactions with extracellular matrix, is required during DTC
migration[4,56,58]. We examined a strain with an endogenously tagged INA-1::mNG allele[70], which
is localized to the surfaces of many cells, including the DTC (Figure 5A). We asked if the patch
was decorated by INA-1::mNG protein. We treated the strain with whole-body *ced-5* RNAi to
induce DTC fragmentation and imaged gonads with a second focus of INA-1::mNG expression
at or near the gonad bend. In the L4 stage, the smaller DTC and the ectopic patch both continue
to localize endogenously tagged INA-1::mNG on their surfaces (Figure 5B and 5C). This is true
whether the DTC is the fragment that migrates in the correct (Figure 5B) or incorrect (Figure 5C)
direction. The ectopic patch therefore contains at least some of the cellular machinery that is
known to be required for DTC migration.
215
Both DTC fragments localize the stem cell niche stemness signal in larvae but only one
maintains expression in adults
LAG-2 is a ligand that activates the GLP-1/Notch receptor on germ cells, acting as the stemness
cue, and is necessary for germ cell mitosis throughout gonad migration and adulthood[8,71,72].
Germ cell mitosis provides the pushing force necessary for gonad elongation[4], so we
hypothesized that in order for a gonad to bifurcate, both branches must support germ cell
proliferation with a source of LAG-2, at least for a time. We observed L4 animals of a strain
expressing endogenously tagged LAG-2::mNG and *lag-2p::myr::TdTomato*[73] to mark the DTC
membrane (Figure 6A, top). We treated this strain with whole-body *ced-5* RNAi to induce DTC
fragmentation and investigated LAG-2::mNG localization (Figure 6B). Indeed, LAG-2::mNG
signal was apparent on both structures that were positive for the *lag-2p::myr::TdTomato*
membrane marker upon RNAi knockdown of *ced-5* (Figure 6C), however the patch has less
total LAG-2::mNG than the DTC (Figure 6F). While the "patch" is not a cell with DTC identity, it
localizes a protein that can maintain stemness in nearby germ cells.
230
The ectopic patch does not maintain niche signaling protein long-term, but both DTC
fragments persist in adulthood
We interpreted the membrane-localized protein signal on the ectopic patch to come from
proteins present at the time of DTC fragmentation or produced soon thereafter. We
hypothesized that the enucleate patch has a minimal ability to synthesize new proteins because
it lacks a genome. We tested whether the patch can sustain stem cell niche function by
examining LAG-2::mNG expression in adult animals. In control young adults at the onset of
oogenesis, the LAG-2::mNG protein localizes to the DTC body and its processes (Figure 6D). In
young adult *ced-5* RNAi-treated animals, we first observed the *lag-2p::myrTdTomato*
membrane-marking signal decorating bifurcated gonad tips and patches perdures (Figure 6E,
6E" and 6C). The membranous cell fragment is not eliminated. However, LAG-2::mNG protein
signal is no longer detected in this structure in more than half of adult animals (Figure 6C), and
overall LAG-2::mNG expression is significantly decreased in adult patches relative to L4
patches (Figure 6F, magenta bar). The *ced-5* RNAi-treated animals maintain LAG-2::mNG in
the DTC itself at levels indistinguishable from control DTCs (Figure 6E' and 6C), from which we
draw two conclusions. First, the loss of LAG-2::mNG signal from the DTC fragment is not due to
a global failure of *ced-5* RNAi-treated animals to produce LAG-2::mNG. Second, the DTC that
fragmented in the larval stage persists and is both morphologically and molecularly normal
hours to days after it breaks apart, suggesting a surprising capacity for cellular recovery in the
DTC (Figure 6E', 6F).
251
Bifurcated germlines are mispatterned, with differentiating germ cells at the second tip
Since LAG-2::mNG signal is reduced on the patch relative to the DTC in L4 and is reduced on
the surface of the patch in adults (Figure 6F), we predicted that one branch of bifurcated gonads
would fail to support stem-like germ cells in later adulthood. We studied germline patterning in

branched gonads compared to controls using a germline-specific promoter-driven fluorescent
histone *(naSi2(mex-5p::H2B::mCherry))*[74] to visualize the nuclear morphology of germ cells in
otherwise wild-type and *ced-5(n1812)* mutant animals.
259
A wild-type adult hermaphrodite germline has a distinctive pattern of germ cell nuclear
morphology (Figure 7A) extending through the formation of oocytes (Figure 7B). Distal germ
cells have small, uniform nuclei (Figure 7C) and visible mitotic figures; these are the cells of the
proliferative zone. Proximal to these cells are cells of the transition zone, which have distinctive
crescent-shaped nuclear fluorescence (Figure 7D); these are at the leptotene/zygotene stage of
meiosis I[14]. Proximal to these cells are meiotic germ cells with "bowl of spaghetti" nuclear
morphology of meiotic pachytene[14] (Figure 7E). As germ cells differentiate into oocytes, the
fluorescence signal reveals condensed bivalent chromosomes of diakinesis, and in DIC we
observe the cells becoming larger and more cytoplasm-rich (Figure 7B-B").
269
In branched gonads of 2-day adults, imaging is complicated by the displacement deep into the
animal of one of the two gonad branches. However, in specimens in which both tips are visible,
we observe one normal looking branch with the expected distal-to-proximal pattern of
undifferentiated-to-differentiated germ cell nuclear morphology (Figure 7F-H). In the second tip,
we consistently observed bulbous masses and disorganized H2B::mCherry signal (Figure 7F, I,
I'). This includes H2B::mCherry signal consistent with the condensing chromosomes of
oogenesis (compare Figure 7I' to 7B'). A normal distal tip germline stem cell niche is not
maintained at the second tip, and germ cells appear to be differentiating in this tip. The
persistence of the branch despite the collapse of the ectopic niche suggests that the germ cells
do not flow from the branch into the "assembly line" of germline output, but instead differentiate
in place.
281
Aberrant patterning is also visible in younger branched gonads. In control L4 larvae (Figure
S3A, S3C-D), most of the distal germline is proliferative, with meiotic transition zone nuclei
visible only near the bend. In the proximal germline, chromatin condenses for
spermatogenesis[14,75]. In branched gonads of *ced-5(n1812)* L4 larvae undergoing
spermatogenesis (Figure S3B), we observe two tubes of germ cells, one of which has the
expected patterning with undifferentiated mitotic cells distal (Figure S3E). The second branch
tip, however, has germ cells with the crescent nuclear morphology typical of the transition zone
of meiotic prophase (Figure S3F). While anatomically it is difficult to identify which is the distal
tip and which is the second branch at this stage (see Figure 1), germ cell nuclear morphology
reveals that only one branch maintains a stem-like population of germ cells.
292
As branched gonads lose LAG-2::mNG signal on the patch membrane in adulthood, the germ
cells in the second branch become disorganized, lose stemness, and appear to differentiate
(taking on crescent-shaped nuclear morphology or showing condensed bivalent chromosomes,
and making larger, blocky cells that resemble oocytes). We conclude that, while both DTC
fragments resulting from Rac-pathway *ced* loss-of-function are initially competent to lead gonad
growth via germ cell proliferation, the enucleate patch cannot sustain adult stem cell niche
activity.
300
DISCUSSION
During larval development, the *C. elegans* hermaphrodite DTC is both a leader cell and a stem
cell niche[1] and must coordinate cell-biological processes like cell signaling, basement
membrane secretion, and adhesion[4] to form a gonad arm populated by proliferating germ cells.
As a leader cell, it is non-protrusive, yet it has long been known that its proper migration
requires the canonical "leading edge" Rac pathway members CED-10/Rac1, CED-5/DOCK180,

CED-2/CrkII, and CED-12/ELMO. These factors play a role in polarity sensing[40], however
defects in DTC migration polarity have not been reported to cause gonad bifurcation. We found
that without Rac pathway function, the DTC fragments during migration into two semi-equivalent
structures capable of leading gonad elongation and supporting germ cell proliferation,
sometimes driving gonad bifurcation. However, one structure lacks a nucleus and thus lacks the
ability to continue to produce LAG-2 protein and thereby act as a stem cell niche through
adulthood.
314
Cell migration depends on major cytoskeletal rearrangements regulated by a complex network
of signal transduction pathways involving the Rho-family GTPases, including the highly
conserved (in eukaryotes) regulator of lamellipodial cell migration, Rac1[76–78]. Migrating cells
often form F-actin-rich protrusions at their leading edge, such as lamellipodia, filopodia, or
blebs[38,79]. The DTC curiously does not form protrusions during migration[4,35]. Nevertheless, DTC
migration defects have long been observed after loss of function of *ced-10*/Rac1[20–22,29].
Migration events required for branching morphogenesis in the vertebrate lung, kidney,
mammary, and salivary glands are also driven by leader cells with a smooth leading edge[80–83].
The mechanisms behind leader cell migration in these processes is largely unknown. Therefore,
elucidating regulators of DTC migration could provide a model for vertebrate organogenesis and
its misregulation, which occurs in certain cancers[84].
326
Rac-pathway factors, including *ced-10*, play many important roles in *C. elegans* development[85].
Rac pathway activity is essential for axon pathfinding[52,86–88] and dendrite regeneration[89] of
neurons. Rac pathway members have recently been discovered to interact with a CLIC chloride
channel protein during the formation of the excretory canal (another tubular organ led by–and in
that case composed entirely of–a single cell)[90]. Embryogenesis also requires *ced-10*/Rac in
several events, including epidermal cell intercalation[91,92], gastrulation[93], mitotic spindle
orientation[41], and of course apoptotic cell corpse clearance[41]. It is not known whether loss-of-
function of *ced-10*/Rac or its GEF complex in these other contexts causes cell fragmentation.
335
Other *C. elegans* cells shed portions of themselves at other stages of development that are
cleared by entosis[94]. For example, membranous lobes are either trimmed away or left behind
after an entosis event in the primordial germ cells (in a *ced-10*-dependent manner)[95] and the
linker cell (it is not known if this occurs in a Rho-GTPase-dependent manner)[96]. However, these
cellular remodeling events are transient, and the residual bodies are eliminated by neighboring
cells, whereas the DTC patch is long-lasting and signal-active, at least for a time.
342
The dividing of a cell without DNA replication into functionally equivalent but genetically distinct
daughters (meiosis notwithstanding) is not unprecedented in animals. The superficial epithelial
cells of the larval zebrafish skin were recently discovered to divide after terminal differentiation
and produce hypoploid descendants[97]. Unlike DTC fragments, however, these descendants
each have reduced nuclear content but none lack a nucleus entirely. An example of mouse cells
fragmenting has recently been reported in the early embryo as a consequence of ectopic
activation of the polar body extrusion pathway in blastomeres[98]. The behavior of the DTC after
loss of Rac pathway activity is one of rather few documented instances of cell fragmentation
without cell death. How loss of Rac-pathway activity causes the DTC to fragment (and why it
only sometimes drives gonad bifurcation) are questions for future study.
353
The fragmented DTC more closely resembles mutant phenotypes observed in mouse
erythrocytes after loss of Rac pathway function. *Rac1*[-/-]*; Rac2*[-/-] erythrocytes are fragmented and
misshapen, and these cells are experimentally determined to be less deformable than cells with
Rac1 expression[59] (the best *C. elegans* BLAST hit of both mouse *Rac1* and *Rac2* is *ced-10*).

Rac1 also functions in human carcinoma cells to downregulate contractile myosin and make
cells more deformable; Rac1 dominant-active cells outcompete Rac1-deficient mutant cells via
entosis while its depletion causes cells to be stiffer and lose out to more deformable cells[60]. On
the other hand, in fibroblasts, Rac1 depletion leads to increased deformability[61], so Rac1
function has a significant but not strictly directional influence on cell deformability. Taken
together, the existing literature and our results suggest that Rac1 is important for cells to
dynamically deform without breaking apart. We conclude that, although the DTC does not
undergo lamellipodial-driven migration, it nevertheless relies on CED-10/Rac1 to retain its
polarity, shape, and integrity at the tip of the growing gonad. Without Rac1 function when
cellular cohesion is lost and the cell breaks apart, each fragment can retain leader cell function
and support gonad elongation, suggesting that the nucleus is surprisingly dispensable for
completion of gonad growth, but is required for long-term maintenance of the stem cell niche.
370
371 **METHODS**
372 Sections of this text are adapted from Li et al., 2022[99], as they describe our standard laboratory
373 practices.
374
Strains
Some strains were provided by the CGC, which is funded by NIH Office of Research
Infrastructure Programs (P40 OD010440). In strain descriptions, we designate linkage to a
promoter with a *p* following the gene name and designate promoter fusions and in-frame fusions
with a double semicolon (::). Some integrated strains (xxIs designation) may still contain for
example the unc-119 (ed4) mutation and/or the unc-119 rescue transgene in their genetic
background, but these are not listed in the strain description for the sake of concision, nor are
most transgene 3' UTR sequences. Complete strain list in Key Resources Table.
383
384 **Worm rearing**
385 *C. elegans* strains were kept at 20°C unless otherwise described on standard NGM media and
386 fed *E. coli* OP50. All animals assessed were hermaphrodites, as males have nonmigratory
387 DTCs. Worm populations were synchronized at L1 arrest for developmental staging by standard
388 egg preps[100].
389
Confocal imaging
All images were acquired at room temperature on a Leica DMI8 with an xLIGHT V3 confocal
spinning disk head (89 North) with a 63× Plan-Apochromat (1.4 NA) objective and an
ORCAFusion GenIII sCMOS camera (Hamamatsu Photonics) controlled by microManager[101].
RFPs were excited with a 555 nm laser, GFPs were excited with a 488 nm laser, and CFP was
excited by a 445 nm laser. Z-stacks through the gonad were acquired with a step-size of 1μm
unless otherwise noted. Worms were mounted on agar pads with 0.01 M sodium azide.
397
RNAi
E. coli HT115(DE3) containing the L4440 plasmid with or without a dsRNA trigger insert from
the Ahringer[102] or Vidal[103] RNAi libraries was grown as a single-colony overnight culture at 37°C,
expression induced with IPTG (Apex BioResearch Products, cat# 20-109) for one hour at 37°C,
and was plated and allowed to grow at least overnight on NGM plates. Note, Ahringer *ced-10*
clone number C09G12.8A has <30 nt homology to the *ced-10* 3' UTR and yielded inconsistent
knockdown; Vidal *ced-10* clone AAA28140 was used preferentially. Worm populations were
synchronized by bleaching according to a standard egg prep protocol[100], plated on NGM plates
seeded with RNAi-expressing bacteria as arrested L1 larvae, and kept on RNAi until the time of
imaging. Whole-body RNAi treatment was conducted at 20°C, and DTC-specific RNAi treatment
was conducted at 16°C due to a temperature-sensitive *rrf-3* mutation in the DTC-specific RNAi

strain. Whole-body *ced-12* RNAi on the *hlh-2(ar623[gfp::hlh-2])* strain included specimens
imaged after two generations on RNAi.
411
412 DTC-specific RNAi was performed using strain NK2115[65], a genetic background carrying an
413 *rde-1(ne219)* loss of function that prevents RNAi activity globally, with RNAi function restored in
414 the DTC (in an operon along with a coding sequence for membrane-tethered mNeonGreen) by
415 a transgene *lag-2p::mNG::PLC^δPH::F2A::rde-1* and a *rrf-3(pk1426)* mutation that enhances
416 RNAi.
417

Image analysis

419 Images were processed in FIJI[104] (Version: 2.14.1/1.54f), and images that required stitching
420 used the ImageJ Stitching Plugin[105]. Detailed descriptions of image analysis for different
421 experiments are provided below.
422

Measurement of DTC vs patch area

To compare the area of the WT DTC with the mutant DTC and enucleate patch, WT and mutant
gonads coexpressing the *lag-2p::mNeonGreen::PLC^δPH* membrane marker and *lag-*
2::P2A::H2B::mT2 nuclear marker were imaged with a 1.0 µm Z-step size. Only late L4 animals
were imaged for both WT and mutants and only mutants with a patch or bifurcation (both
marked by a second focus of mNG expression) were imaged. Late L4 animals were defined as
between the L4.6-L4.9 larval stage, based on vulval morphology[106]. Using FIJI[104], maximum
intensity z-projections were made through the depth of the entire DTC and patch structures. The
structures were outlined by hand (based on membranous mNG signal in the GFP channel)
using the freehand selections tool to define ROIs that corresponded to the boundary of the
structure. The area of these ROIs was then measured in µm^2.
434

Fluorescence intensity of LAG-2::mNeonGreen

For quantitative comparisons of fluorescence intensity shown in FIGURES, gonads were
imaged with uniform laser power and exposure times with 0.3 µm Z-steps. For the L4 stage,
only L4.6-L4.9 (late L4) animals were imaged, staged based on vulval morphology[106]. Young
Adults were identified as ~24h post the L4 stage and by the presence of some fertilized
embryos. Using FIJI[104], sum-intensity z-projections were made through the depth of the focal
DTC and patch structures as visualized with the TdTomato signal in the RFP channel, excluding
deeper Z-slices with appearance of large, bright gut granules in the footprint of the DTC. The
structures were outlined with the Threshold function to make ROIs, and Mean Gray Value (and
Raw Integrated Density) were measured inside the ROI in arbitrary units. The same ROIs were
shifted onto worm gonad background and GFP fluorescence intensity was measured again.
This background was subtracted from the signal for the final value of LAG-2::mNG detected in
the structure. In the *ced-5* RNAi group, only RNAi-treated specimens with patches or
bifurcations were analyzed; escapers were discarded. Because different regions of the gonads
in adults are often in different focal planes, not every sample had both a visible DTC and a
visible patch. Along those lines, we measure lower LAG-2::mNG signal in adult control and *ced-*
5 RNAi DTCs compared to L4 DTCs. We attribute this to imaging adult DTCs at a greater tissue
depth, which reduces signal capture, and with greater background autofluorescence in adults
relative to L4s. Patches in the adult, on the other hand, were usually on the surface of the gonad
and were never under the gut. We never observed adult DTCs (either control or *ced-5* RNAi-
treated) that lacked LAG-2::mNG signal, while the patches of *ced-5* RNAi-treated adults lacked
noticeable signal in more than half of all samples.
457
458
459

460   **Statistical analyses**
461   Sample sizes, tests, test statistics, and p values are given for each analysis in the
462   accompanying figure legends. All statistical testing was carried out using Prism (GraphPad
463   Prism Version 10.10 (264) for macOS), GraphPad Software, San Diego, CA. Significance bars
464   in each graph were produced by Dunnett's T3 Test for multiple comparisons of selected groups.
465
466   **ACKNOWLEDGEMENTS**

467   We thank Taylor Medwig-Kinney, Pu Zhang, Rob Dowen and his lab members, Brian Kinney,
468   Fred Koitz, and other Gordon Lab members for feedback on the manuscript and sharing
469   resources. Funded by NIGMS Grant 1R35GM147704 to KLG.

470
FIGURE LEGENDS
Figure 1. Rac pathway *ced* mutants have gonad bifurcation with moderate penetrance.
(A) Cartoon illustrating wild-type and bifurcated gonads in *C. elegans* hermaphrodites. (**B**)
Micrographs made with DIC imaging of *C. elegans* hermaphrodites in late larval L4 stage for
wild-type N2 and four mutant strains. Images are Z-projections through thickness of the gonad
required to capture both tips. Visible gonad outlined in black dashed line. Yellow asterisks show
the anatomical tips of gonad branches of the gonad on the right-hand side of the image. White
asterisk, visible tip of other gonad arm. (**C**) Same as B, but in the young adult stage. Scale bar
20 µm. **(D)** Graph showing percentage of N2 controls and mutants of each genotype (L4s and
young adults) with gonad bifurcation (yellow), other gonad defects (blue), or no defect (black).
N2 WT control N=51, *ced-5(n1812)* N=56, *ced-10(n1993)* N=46, *ced-12(n3261)* N=70, *ced-*
2(n1994) N=45.

483
Figure 2. Bifurcation is a DTC cell-autonomous defect caused by Rac-pathway loss of
function, and a DTC-membrane marker reveals associated somatic gonad cell
morphology defects. (A) Strain expressing a combined *lag-2* promoter-driven membrane
marker and *rde-1* rescue transgene *lag-2p::mNG::PLC^{δPH}::F2A::rde-1* in a genetic background
that is *rde-1(ne219)* loss of function and *rrf-3(pk1426)* hypersensitive to RNAi[65]. On control
RNAi (empty vector L4440), one site of strong mNG expression is visible in each gonad arm in
late L4 hermaphrodites–the distal tip cell. **(A')** Otherwise wild-type N2 strain bearing
cpIs122[lag-2p::mNeonGreen::PLC^{δPH}][65] marker of the DTC. **(B)** DTC-specific RNAi knockdown
of Rac-pathway *ced* genes causes a range of cellular and anatomical gonad defects: gonad
bifurcation, in which mNG expression is always observed on both tips, and the formation of a
second "patch" of mNG expression near the bend of an otherwise anatomically normal gonad
(B'). (C) Mutants of the Rac-pathway *ced* genes with the *cpIs122* marker of the DTC manifest
the same defects as animals treated with DTC-specific RNAi to those genes: gonad bifurcation,
in which mNG expression is always observed on both tips, and the formation of a second
"patch" of mNG expression near the bend of an otherwise anatomically normal gonad **(C')**.
Maximum projection of GFP fluorescence channel through Z-slices with mNG signal merged
with maximum projection of DIC image through slices capturing the gonad tip(s). Imaged at late
L4 stage. Visible gonad outlined in black dashed line. Yellow asterisks mark gonad tips of focal
gonads, white asterisk marks tip of other gonad arm if visible. Yellow arrowhead marks the
patch, and white arrowhead marks patch of other gonad arm if visible. Autofluorescence of the
gut is visible as green punctae; this is unrelated to expression of the fluorescent protein. Scale
bars 20 µm.
506
507
508
509

Figure 3. Somatic gonad cell morphology defects manifest at or just after DTC turning.
(A) Developmental time series of otherwise wild-type N2 animals bearing *cpIs122[lag-*
2p::mNeonGreen::PLC$^{\delta PH}$][65] marker of the DTC during ventral DTC migration (left), turning
(center), and dorsal migration just after turning (right). The DTC has a gumdrop shape with only
very thin trailing filaments. **(B)** Rac-pathway *ced* mutants expressing the same *cpIs122* marker
of DTC at the same stages. Patches appear at or after the turning stage. Maximum projection of
GFP fluorescence channel through all Z-slices with focal mNG signal merged with maximum
projection of DIC image through slices capturing the gonad tip(s). Visible gonad outlined in black
dashed line. Yellow asterisks mark gonad tips of focal gonads, white asterisk marks tip of other
gonad arm if visible, yellow arrowheads mark the patch. Scale bars 20 µm.
520
Figure 4. The second tip is not led by a cell, but by a DTC fragment (A-C) A strain
expressing an endogenously tagged *hlh-2(ar623[gfp::hlh-1])* allele encoding GFP::HLH-2
transcription factor[68] on control and experimental whole-body RNAi. Left, fluorescence images
merged with DIC projection through gonad tip(s). Center, inset of gonad tip. Right, GFP::HLH-2
alone. Yellow asterisks mark DTC; yellow carats mark proximal gonad cells expressing
GFP::HLH-2. **(A)** GFP::HLH-2 strain on control RNAi (empty vector L4440) has concentrated
fluorescence in the DTC nucleus and dimmer, diffuse cytoplasmic GFP in the DTC. **(B)** RNAi
knockdown of *ced-5* in the strain bearing the *gfp::hlh-2* allele. Upper left, maximum projection of
GFP fluorescence channel through the more superficial tip of a bifurcated gonad in GFP merged
with DIC. Inset right, the more superficial tip has diffuse cytoplasmic GFP::HLH-2 signal. Below,
maximum projection of GFP fluorescence channel through the deeper tip of same bifurcated
gonad merged with DIC. Inset right, the deeper tip has nuclear GFP::HLH-2 signal. Only
animals with a second site of HLH-2::GFP expression were imaged of which N=9 bifurcated
gonads. **(C)** RNAi knockdown of *ced-12* in the strain bearing the *gfp::hlh-2* allele. Upper left,
maximum projection of GFP fluorescence channel through the more superficial tip of a
bifurcated gonad in GFP merged with DIC. Inset right, the more superficial tip has nuclear
GFP::HLH-2 signal. Below, maximum projection of GFP fluorescence channel through the
deeper tip of same bifurcated gonad merged with DIC. Inset right, the deeper tip has diffuse
cytoplasmic GFP::HLH-2 signal. N=10 bifurcated gonads. **(D-E)** A strain bearing a transgene
that marks the membrane of the DTC, *cpIs122[lag-2p::mNeonGreen::PLC$^{\delta PH}$]*, and a nuclear
marker inserted at the endogenous *lag-2* locus (*lag-2::P2A::H2B::mT2)*[69] in otherwise (D)
wildtype and (E) *ced-5(n1812)* genetic backgrounds. Left, maximum projections through all Z-
slices with focal mNG (green) and/or mT2 (magenta) signal merged with DIC projection through
gonad tip(s). Center, inset of tip, *lag-2p::mNG::PLC$^{\delta PH}$* only. Right, inset of tip, *H2B::mT2* only.
(D) Otherwise wild-type strain with DTC membrane and nuclear markers. **(E)** Mutant with
bifurcated gonad *ced-5(n1812)* with DTC membrane and nuclear markers. Inset, top, the DTC
membrane (center) and nuclear (right) signal. Inset, bottom, the membranous fragment (center)
lacks histone signal (right). Visible gonad outlined in black or white dashed line. Yellow asterisks
mark gonad tip with DTC nucleus, yellow arrowhead marks the second tip. Yellow boxes show
position of insets in larger image. Autofluorescence of the gut is visible as punctae; this is
unrelated to expression of the fluorescent proteins. Imaged at L4 stage (L4.6-L4.9), based on
vulval morphology[106]. Scale bars 20 µm. **(F)** Graph showing that both the *ced-5(n1812)* DTC
and patch are smaller than wild-type DTCs, but the sum of their projected area is not different
from that of wild-type DTCs. WT DTC N=10, mutant DTC N=9, mutant patch N=11, mutant DTC
+ patch N=9. Graphed data is presented with median and interquartile range. One-way ANOVA
testing the effect of structure type on size. $F_{3.00, 32.26}= 70.17$, $p<0.0001$. Dunnett's T3 multiple
comparisons test found that the mean value of structure area was significantly different between
WT DTC vs. mutant DTC (p= 0.0126, 95% CI = [9.587, 85.10]) and mutant DTC vs. mutant
patch (p < 0.0001, 95% CI=[90.45, 153.6]). However, the structure area was not significantly
different between the WT DTC vs. mutant DTC + patch (p=0.9990 95% CI=[-39.54, 43.41]).

Figure 5. The DTC fragment retains INA-1, a key regulator of gonadogenesis. A strain
expressing an endogenously tagged allele of *ina-1(qy23[ina-1::mNG])* has mNG signal on the
surface of many cells, including the DTC. Left, merged images of maximum projection through
Z-slices with mNG signal and DIC image showing gonad tip(s). Right, GFP channel only. Insets,
gonad tip(s). **(A)** INA-1::mNG-expressing strain on control RNAi (empty vector L4440). **(B)**
Specimen of INA-1::mNG-expressing strain on *ced-5* RNAi. Note that the DTC is closer to its
correct anatomical position than the patch. **(C)** Another specimen of INA-1::mNG-expressing
strain on *ced-5* RNAi. Note that the DTC is migrating in the wrong direction with the patch closer
to the correct anatomical DTC position; the DTC is deeper in the specimen and is therefore
dimmer. Visible gonad outlined in black dashed line. Yellow asterisks mark gonad tip with DTC
nucleus, yellow arrowhead marks the enucleate patch. Yellow boxes show positions of insets in
larger images. Autofluorescence of the gut is visible as punctae; this is unrelated to expression
of the fluorescent protein. Imaged at the L4 stage. Scale bars 20 µm.
574
Figure 6. The DTC fragment retains the germline stem cell niche stemness cue in larvae
but not adults. (A) Top, larval gonad of an animal from a strain expressing an endogenously
tagged allele of *lag-2(cp193[lag-2:: mNeonGreen^3xFlag])* as well as a transgene to mark the
DTC, *qIs154(lag-2p:: myr::tdTomato)* on control RNAi (empty vector L4440) shows LAG-2::mNG
protein (magenta) in the DTC (membrane in green). Left, merged images of maximum
projection through Z-slices with mNG and/or TdTomato signal and DIC image showing gonad
tip(s). Center, inset of DTC, merged fluorescence channels only. Right, LAG-2::mNG channel
alone. **(B)** Larval gonad of an animal from the strain expressing endogenously tagged LAG-2
(magenta) and a DTC membrane marker (green) on *ced-5* RNAi. Note that both structures
expressing the DTC membrane marker also have LAG-2::mNG protein on their surfaces. **(B')**
Inset showing merged fluorescence in the DTC (left) and LAG-2::mNG (right). **(B")** Inset
showing merged fluorescence and LAG-2::mNG expression in the second tip. **(C)** Graph
showing incidence of LAG-2::mNG signal in the DTC or patch/second tip at L4 stage and in
Adults in control and RNAi-treated animals. L4 control DTC, N = 20; L4 RNAi DTC, N = 14; L4
RNAi patch = 16; Adult control DTC, N = 19; Adult RNAi DTC, N = 9; Adult RNAi patch = 17.
(D-E) The same strain and treatments imaged as Day 1 Adults (staging described in Methods).
(D) Control RNAi (empty vector L4440). All channels merged. Center, inset of DTC, merged
fluorescence channels. Right, LAG-2::mNG channel alone. **(E)** Same strain on *ced-5* RNAi. All
channels merged. **(E')** Center, inset of DTC, merged fluorescence channels. Right, LAG-
2::mNG channel. **(E")** Center, inset of second tip, merged fluorescence channels. Right, LAG-
2::mNG channel. Yellow asterisks mark DTC, yellow arrowheads mark the second tip. Yellow
boxes show positions of insets in larger images. Autofluorescence of the gut is visible as
punctae; this is unrelated to expression of the fluorescent proteins. Scale bars 20 µm. **(F)** Graph
showing quantification of LAG-2::mNG fluorescence Mean Gray Value (arbitrary units) in control
and *ced-5* RNAi animals in the DTC or patch/second tip. L4 control DTC, N = 20; L4 RNAi-
treated DTC = 14; L4 RNAi-treated patch = 16; Adult control DTC, N = 19; Adult RNAi-treated
DTC, N= 9; Adult RNAi-treated patch = 17. Graphed data presented with median and
interquartile range. One-way ANOVA testing the effect of condition on LAG-2::mNG expression.
$F_{5.000, 28.09} = 30.14$, p <0.0001. Dunnett's T3 multiple comparisons test found that mean value of
LAG-2::mNG expression was significantly different between L4 RNAi DTC vs. L4 RNAi patch
(p= 0.0002, 95% CI = [1415, 4243]), L4 RNAi DTC vs. Adult RNAi DTC (p=0.0007, 95% CI =
[1002, 3862]), L4 Control DTC vs. Adult Control DTC (p = 0.0024, 95% CI = [307.3, 1823]),
Adult RNAi DTC vs. Adult RNAi patch (p = 0.0019, 95% CI = [321.5, 1172]) and highlighted in
magenta, L4 RNAi patch vs. Adult RNAi patch (p = 0.0024, 95% CI = [112.3, 586.3]). However,
LAG-2::mNG was not significantly different between the L4 Control DTC vs. L4 RNAi DTC
(p=0.3631, 95% CI = [-2469, 522.9]) or Adult control DTC vs. Adult RNAi DTC (p=0.2368, 95%
CI = [-136.2, 924.0]).

Figure 7. Bifurcated germlines are mispatterned and have differentiating germ cells at the distal end. Micrographs comparing features of adult germlines between control (A-E) and *ced-5(n1812)* (F-I) animals. Germ cell nuclei visualized with the *naSi2(mex-5p::H2B::mCherry)* transgene[74], magenta in merge. **(A)** Z-projection through DIC image of two-day adult (48 hours post L4) control gonad merged with germ cell histone fluorescence. Boxes show positions of insets that follow. **(B)** Enlargement of cellularizing oocyte showing increased cytoplasm in DIC and, single slice through plane of nucleolus **(B')** condensed bivalent chromosomes at diakinesis. **(C)** Enlargement of distal tip fluorescence image showing undifferentiated germ cell nuclei. **(D)** Enlargement of the fluorescence image of the meiotic transition zone showing its distinctive crescent-shaped nuclear morphology. **(E)** Enlargement of meiotic pachytene nuclei. **(F)** Z-projection through DIC image of 2-day adult *ced-5(n1812)* gonad merged with germ cell histone fluorescence. Yellow boxes show positions of insets in larger images. **(G)** Enlargement of distal tip fluorescence image showing undifferentiated germ cell nuclei. **(H)** Enlargement of meiotic pachytene nuclei (transition zone is obscured in this sample by the second gonad branch). **(I)** Enlargement of single plane through nucleoli of DIC image of second gonad tip of sample shown in (F). Note large, irregular cells inside. **(I')** Enlargement of Z-projection through fluorescence image of second gonad tip shown in (I). Note abnormal fluorescent bodies including notable condensed, bivalent-like structures (yellow arrows). Gonads outlined in black or white dashed lines. Autofluorescence of the gut is visible as punctae; this is unrelated to expression of the fluorescent proteins. Scale bars: 20 μm.

REFERENCES

1. Kimble, J. & White, J. G. On the Control of Germ Cell Development in Caenorhabditis Elegans. doi:10.1016/0012-1606(81)90284-0.

2. Lehmann, R. Cell migration in invertebrates: clues from border and distal tip cells.

3. Hubbard, E. J. A. & Greenstein, D. The Caenorhabditis elegans gonad: A test tube for cell and developmental biology. Dev. Dyn. 218, 2–22 (2000).

4. Agarwal, P., Shemesh, T. & Zaidel-Bar, R. Directed cell invasion and asymmetric adhesion drive tissue elongation and turning in C. elegans gonad morphogenesis. Dev. Cell 57, 2111-2126.e6 (2022).

5. Genetics of cell and axon migrations in Caenorhabditis elegans | Development | The Company of Biologists.

6. Kimble, J. & HIrsh, D. The Postembryonic Cell Lineages of the Hermaphrodite and Male Gonads in Caenorhabditis elegans. doi:10.1016/0012-1606(79)90035-6.

7. Hubbard, E. J. A. & Schedl, T. Biology of the Caenorhabditis elegans Germline Stem Cell System. Genetics 213, 1145–1188 (2019).

8. Gao, D. & Kimble, J. APX-1 can substitute for its homolog LAG-2 to direct cell interactions throughout Caenorhabditis elegans development. Proc. Natl. Acad. Sci. U. S. A. 92, 9839–9842 (1995).

9. Gordon, K. L., Zussman, J. W., Li, X., Miller, C. & Sherwood, D. R. Stem cell niche exit in C. elegans via orientation and segregation of daughter cells by a cryptic cell outside the niche. eLife 9, e56383 (2020).

10. Kimble, J. & Crittenden, S. L. Controls of Germline Stem Cells, Entry into Meiosis, and the Sperm/Oocyte Decision in Caenorhabditis elegans. Annu. Rev. Cell Dev. Biol. 23, 405–433 (2007).

11. Austin & Kimble, J. glp-1 Is Required in the Germ Line for Regulation of the Decision between Mitosis and Meiosis in C. elegans. doi:10.1016/0092-8674(87)90128-0.

12. Rosu, S. & Cohen-Fix, O. Live-imaging analysis of germ cell proliferation in the C. elegans adult supports a stochastic model for stem cell proliferation. Dev. Biol. 423, 93–100 (2017).

13. Zanetti, S. et al. The sperm-oocyte switch in the C. elegans hermaphrodite is controlled through steady-state levels of the fem-3 mRNA. RNA (2012) doi:10.1261/rna.031237.111.

14. Pazdernik, N. & Schedl, T. Introduction to Germ Cell Development in Caenorhabditis elegans. in Germ Cell Development in C. elegans (ed. Schedl, T.) 1–16 (Springer, 2013). doi:10.1007/978-1-4614-4015-4_1.

15. Gumienny, T. L., Lambie, E., Hartwieg, E., Horvitz, H. R. & Hengartner, M. O. Genetic control of programmed cell death in the Caenorhabditis elegans hermaphrodite germline. Dev. Camb. Engl. 126, 1011–1022 (1999).

16. Reddien, P. W. & Horvitz, H. R. The Engulfment Process of Programmed Cell Death in Caenorhabditis Elegans. Annu. Rev. Cell Dev. Biol. 20, 193–221 (2004).

17. Ellis, H. M. & Horvitz, H. R. Genetic control of programmed cell death in the nematode C. elegans. Cell 44, 817–829 (1986).

18. Hedgecock, E. M., Sulston, J. E. & Thomson, J. N. Mutations Affecting Programmed Cell Deaths in the Nematode Caenorhabditis eleganss. Science 220, 1277–1279 (1983).

19. Ellis, R. E., Jacobson, D. M. & Horvitz, H. R. Genes required for the engulfment of cell
corpses during programmed cell death in Caenorhabditis elegans. Genetics 129, 79–94
(1991).

20. Gumienny, T. L. et al. CED-12/ELMO, a Novel Member of the CrkII/Dock180/Rac Pathway,
Is Required for Phagocytosis and Cell Migration. Cell 107, 27–41 (2001).

21. Zhou, Z., Caron, E., Hartwieg, E., Hall, A. & Horvitz, H. R. The C. elegans PH Domain
Protein CED-12 Regulates Cytoskeletal Reorganization via a Rho/Rac GTPase Signaling
Pathway. Dev. Cell 1, 477–489 (2001).

22. Reddien, P. W. & Horvitz, H. R. CED-2/CrkII and CED-10/Rac control phagocytosis and
cell migration in Caenorhabditis elegans. Nat. Cell Biol. 2, 131–136 (2000).

23. Wu, Y.-C., Tsai, M.-C., Cheng, L.-C., Chou, C.-J. & Weng, N.-Y. C. elegans CED-12 Acts in
the Conserved CrkII/DOCK180/Rac Pathway to Control Cell Migration and Cell Corpse
Engulfment. Dev. Cell 1, 491–502 (2001).

24. Hengartner, M. O., Ellis, R. E. & Horvitz, H. R. Caenorhabditis elegans gene ced-9 protects
cells from programmed cell death. Nature 356, 494–499 (1992).

25. Conradt, B., Wu, Y.-C. & Xue, D. Programmed Cell Death During Caenorhabditis elegans
Development. Genetics 203, 1533–1562 (2016).

26. Mangahas, P. M. & Zhou, Z. Clearance of apoptotic cells in Caenorhabditis elegans.
Semin. Cell Dev. Biol. 16, 295–306 (2005).

27. Shakir, M. A. et al. The Arp2/3 activators WAVE and WASP have distinct genetic
interactions with Rac GTPases in Caenorhabditis elegans axon guidance. Genetics 179,
1957–1971 (2008).

28. Cotteret, S. & Chernoff, J. The evolutionary history of effectors downstream of Cdc42 and
Rac. Genome Biol. 3, reviews0002.1-reviews0002.8 (2002).

29. Wu, Y.-C. & Horvitz, H. R. C. elegans phagocytosis and cell-migration protein CED-5 is
similar to human DOCK180. Nature 392, 501–504 (1998).

30. Brugnera, E. et al. Unconventional Rac-GEF activity is mediated through the Dock180-
ELMO complex. Nat. Cell Biol. 4, 574–582 (2002).

31. Wang, X. et al. Cell Corpse Engulfment Mediated by C. elegans Phosphatidylserine
Receptor Through CED-5 and CED-12. Science 302, 1563–1566 (2003).

32. Venkatachalam, T., Mannimala, S. & Soto, M. C. CED-5/CED-12 (DOCK/ELMO) can
promote and inhibit F-actin formation via distinct motifs that target different GTPases.
BioRxiv Prepr. Serv. Biol. 2023.10.04.560868 (2023) doi:10.1101/2023.10.04.560868.

33. Hanna, S. & El-Sibai, M. Signaling networks of Rho GTPases in cell motility. Cell. Signal.
25, 1955–1961 (2013).

34. Nobes, C. D. & Hall, A. Rho, rac, and cdc42 GTPases regulate the assembly of
multimolecular focal complexes associated with actin stress fibers, lamellipodia, and
filopodia. Cell 81, 53–62 (1995).

35. Sherwood, D. R. & Plastino, J. Invading, Leading and Navigating Cells in Caenorhabditis
elegans : Insights into Cell Movement in Vivo. Genetics 208, 53–78 (2018).

36. Kim, H.-S. et al. VAB-10 spectraplakin acts in cell and nuclear migration in Caenorhabditis
elegans. Dev. Camb. Engl. 138, 4013–4023 (2011).

37. Gov, N. S. Collective cell migration patterns: Follow the leader. Proc. Natl. Acad. Sci. 104,
15970–15971 (2007).

38. Friedl, P. & Gilmour, D. Collective cell migration in morphogenesis, regeneration and
cancer. Nat. Rev. Mol. Cell Biol. 10, 445–457 (2009).
39. Qin, L., Yang, D., Yi, W., Cao, H. & Xiao, G. Roles of leader and follower cells in collective
cell migration. Mol. Biol. Cell 32, 1267–1272 (2021).
40. Levy-Strumpf, N., Krizus, M., Zheng, H., Brown, L. & Culotti, J. G. The Wnt Frizzled
Receptor MOM-5 Regulates the UNC-5 Netrin Receptor through Small GTPase-Dependent
Signaling to Determine the Polarity of Migrating Cells. PLoS Genet. 11, e1005446 (2015).
41. Cabello, J. et al. The Wnt Pathway Controls Cell Death Engulfment, Spindle Orientation,
and Migration through CED-10/Rac. PLoS Biol. 8, e1000297 (2010).
42. Kimble, J. Germline proliferation and its control. WormBook (2005)
doi:10.1895/wormbook.1.13.1.
43. Gordon, K. Recent Advances in the Genetic, Anatomical, and Environmental Regulation of
the C. elegans Germ Line Progenitor Zone. J. Dev. Biol. 8, 14 (2020).
44. Levy-Strumpf, N. & Culotti, J. G. Netrins and Wnts function redundantly to regulate antero-
posterior and dorso-ventral guidance in C. elegans. PLoS Genet. 10, e1004381 (2014).
45. Blelloch, R. et al. The gon-1 gene is required for gonadal morphogenesis in Caenorhabditis
elegans. Dev. Biol. 216, 382–393 (1999).
46. Cram, E. J., Clark, S. G. & Schwarzbauer, J. E. Talin loss-of-function uncovers roles in cell
contractility and migration in C. elegans. J. Cell Sci. 116, 3871–3878 (2003).
47. Tannoury, H. et al. CACN-1/Cactin interacts genetically with MIG-2 GTPase signaling to
control distal tip cell migration in C. elegans. Dev. Biol. 341, 176–185 (2010).
48. Kubota, Y., Nagata, K., Sugimoto, A. & Nishiwaki, K. Tissue architecture in the
Caenorhabditis elegans gonad depends on interactions among fibulin-1, type IV collagen
and the ADAMTS extracellular protease. Genetics 190, 1379–1388 (2012).
49. Meighan, C. M. & Schwarzbauer, J. E. Control of C. elegans hermaphrodite gonad size and
shape by vab-3/Pax6-mediated regulation of integrin receptors. Genes Dev. 21, 1615–1620
(2007).
50. B, I. et al. SRC-1, a non-receptor type of protein tyrosine kinase, controls the direction of
cell and growth cone migration in C. elegans. Dev. Camb. Engl. 132, (2005).
51. Su, M. et al. Regulation of the UNC-5 netrin receptor initiates the first reorientation of
migrating distal tip cells in Caenorhabditis elegans. Dev. Camb. Engl. 127, 585–594 (2000).
52. Lundquist, E. A., Reddien, P. W., Hartwieg, E., Horvitz, H. R. & Bargmann, C. I. Three C.
elegans Rac proteins and several alternative Rac regulators control axon guidance, cell
migration and apoptotic cell phagocytosis. Development 128, 4475–4488 (2001).
53. Zipkin, I. D., Kindt, R. M. & Kenyon, C. J. Role of a New Rho Family Member in Cell
Migration and Axon Guidance in C. elegans. Cell 90, 883–894 (1997).
54. Xu, X., Rongali, S. C., Miles, J. P., Lee, K. D. & Lee, M. pat-4/ILK and unc-112/Mig-2 are
required for gonad function in Caenorhabditis elegans. Exp. Cell Res. 312, 1475–1483
(2006).
55. Hutter, H., Ng, M.-P. & Chen, N. GExplore: a web server for integrated queries of protein
domains, gene expression and mutant phenotypes. BMC Genomics 10, 529 (2009).
56. Baum, P. D. & Garriga, G. Neuronal migrations and axon fasciculation are disrupted in ina-
1 integrin mutants. Neuron 19, 51–62 (1997).

57. Lee, M., Cram, E. J., Shen, B. & Schwarzbauer, J. E. Roles for βpat-3 Integrins in
Development and Function ofCaenorhabditis elegans Muscles and Gonads *. J. Biol.
Chem. 276, 36404–36410 (2001).

58. Meighan, C. M. & Schwarzbauer, J. E. Temporal and spatial regulation of integrins during
development. Curr. Opin. Cell Biol. 20, 520–524 (2008).

59. Kalfa, T. A. et al. Rac GTPases regulate the morphology and deformability of the
erythrocyte cytoskeleton. Blood 108, 3637–3645 (2006).

60. Sun, Q. et al. Competition between human cells by entosis. Cell Res. 24, 1299–1310
(2014).

61. Kunschmann, T. et al. The Small GTPase Rac1 Increases Cell Surface Stiffness and
Enhances 3D Migration Into Extracellular Matrices. Sci. Rep. 9, 7675 (2019).

62. Hong, Y., Roy, R. & Ambros, V. Developmental regulation of a cyclin-dependent kinase
inhibitor controls postembryonic cell cycle progression in Caenorhabditis elegans. Dev.
Camb. Engl. 125, 3585–3597 (1998).

63. Fay, D. S., Keenan, S. & Han, M. fzr-1 and lin-35/Rb function redundantly to control cell
proliferation in C. elegans as revealed by a nonbiased synthetic screen. Genes Dev. 16,
503–517 (2002).

64. Fujita, M., Takeshita, H. & Sawa, H. Cyclin E and CDK2 Repress the Terminal
Differentiation of Quiescent Cells after Asymmetric Division in C. elegans. PLOS ONE 2,
e407 (2007).

65. Linden, L. M. et al. Identification of regulators of germ stem cell enwrapment by its niche in
C. elegans. Dev. Biol. 429, 271–284 (2017).

66. Kostic, I., Li, S. & Roy, R. cki-1 links cell division and cell fate acquisition in the C. elegans
somatic gonad. doi:10.1016/j.ydbio.2003.07.001.

67. Kintaka, R., Makanae, K. & Moriya, H. Cellular growth defects triggered by an overload of
protein localization processes. Sci. Rep. 6, 31774 (2016).

68. Attner, M. A., Keil, W., Benavidez, J. M. & Greenwald, I. HLH-2/E2A Expression Links
Stochastic and Deterministic Elements of a Cell Fate Decision during C. elegans
Gonadogenesis. Curr. Biol. CB 29, 3094-3100.e4 (2019).

69. Medwig-Kinney, T. N., Sirota, S. S., Gibney, T. V., Pani, A. M. & Matus, D. Q. An in vivo
toolkit to visualize endogenous LAG-2/Delta and LIN-12/Notch signaling in C. elegans.
Open Access.

70. Jayadev, R. et al. α-Integrins dictate distinct modes of type IV collagen recruitment to
basement membranes. J. Cell Biol. 218, 3098–3116 (2019).

71. Henderson, S. T., Gao, D., Lambie, E. J. & Kimble, J. lag-2 may encode a signaling ligand
for the GLP-1 and LIN-12 receptors of C. elegans. Dev. Camb. Engl. 120, 2913–2924
(1994).

72. Nadarajan, S., Govindan, J. A., McGovern, M., Hubbard, E. J. A. & Greenstein, D. MSP
and GLP-1/Notch signaling coordinately regulate actomyosin-dependent cytoplasmic
streaming and oocyte growth in C. elegans. Dev. Camb. Engl. 136, 2223–2234 (2009).

73. Gordon, K. L. et al. Ectopic germ cells can induce niche-like enwrapment by neighboring
body wall muscle. Curr. Biol. CB 29, 823-833.e5 (2019).

74. Roy, D., Kahler, D. J., Yun, C. & Hubbard, E. J. A. Functional Interactions Between rsks-
1/S6K, glp-1/Notch, and Regulators of Caenorhabditis elegans Fertility and Germline Stem
Cell Maintenance. G3 Bethesda Md 8, 3293–3309 (2018).
75. Chu, D. S. et al. Sperm chromatin proteomics identifies evolutionarily conserved fertility
factors. Nature 443, 101–105 (2006).
76. Ridley, A. J., Paterson, H. F., Johnston, C. L., Diekmann, D. & Hall, A. The small GTP-
binding protein rac regulates growth factor-induced membrane ruffling. Cell 70, 401–410
(1992).
77. Wu, Y. I. et al. A genetically encoded photoactivatable Rac controls the motility of living
cells. Nature 461, 104–108 (2009).
78. Mehidi, A. et al. Transient Activations of Rac1 at the Lamellipodium Tip Trigger Membrane
Protrusion. Curr. Biol. 29, 2852-2866.e5 (2019).
79. Hall, A. Rho GTPases and the actin cytoskeleton. Science 279, 509–514 (1998).
80. Ewald, A. J., Brenot, A., Duong, M., Chan, B. S. & Werb, Z. Collective Epithelial Migration
and Cell Rearrangements Drive Mammary Branching Morphogenesis. Dev. Cell 14, 570–
581 (2008).
81. Harunaga, J. S., Doyle, A. D. & Yamada, K. M. Local and global dynamics of the basement
membrane during branching morphogenesis require protease activity and actomyosin
contractility. Dev. Biol. 394, 197–205 (2014).
82. Andrew, D. J. & Ewald, A. J. Morphogenesis of epithelial tubes: Insights into tube
formation, elongation, and elaboration. Dev. Biol. 341, 34–55 (2010).
83. Williams, J. M. & Daniel, C. W. Mammary ductal elongation: differentiation of
myoepithelium and basal lamina during branching morphogenesis. Dev. Biol. 97, 274–290
(1983).
84. Gray, R. S., Cheung, K. J. & Ewald, A. J. Cellular mechanisms regulating epithelial
morphogenesis and cancer invasion. Curr. Opin. Cell Biol. 22, 640–650 (2010).
85. Sáenz-Narciso, B., Gómez-Orte, E., Zheleva, A., Gastaca, I. & Cabello, J. Control of
developmental networks by Rac/Rho small GTPases: How cytoskeletal changes during
embryogenesis are orchestrated. Bioessays 38, 1246–1254 (2016).
86. Nørgaard, S., Deng, S., Cao, W. & Pocock, R. Distinct CED-10/Rac1 domains confer
context-specific functions in development. PLOS Genet. 14, e1007670 (2018).
87. Quinn, C. C., Pfeil, D. S. & Wadsworth, W. G. CED-10/Rac1 mediates axon guidance by
regulating the asymmetric distribution of MIG-10/lamellipodin. Curr. Biol. CB 18, 808–813
(2008).
88. Nørgaard, S. & Pocock, R. Rac GTPases: domain-specific functions in neuronal
development. Neural Regen. Res. 14, 1367–1368 (2019).
89. Brar, H. K. et al. Dendrite regeneration in C. elegans is controlled by the RAC GTPase
CED-10 and the RhoGEF TIAM-1. PLoS Genet. 18, e1010127 (2022).
90. Af, A., J, E. & Dd, S. A metazoan-specific C-terminal motif in EXC-4 and Gα-Rho/Rac
signaling regulate cell outgrowth during tubulogenesis in C. elegans. Dev. Camb. Engl.
149, (2022).
91. Walck-Shannon, E., Reiner, D. & Hardin, J. Polarized Rac-dependent protrusions drive
epithelial intercalation in the embryonic epidermis of C. elegans. Dev. Camb. Engl. 142,
3549–3560 (2015).

92. Zhu, Y. & Hardin, J. TIAM-1 regulates polarized protrusions during dorsal intercalation in
the C. elegans embryo through both its GEF and N-terminal domains. BioRxiv Prepr. Serv.
Biol. 2023.07.24.550374 (2023) doi:10.1101/2023.07.24.550374.
93. Soto, M. C. et al. The GEX-2 and GEX-3 proteins are required for tissue morphogenesis
and cell migrations in C. elegans. Genes Dev. 16, 620–632 (2002).
94. Overholtzer, M. et al. A nonapoptotic cell death process, entosis, that occurs by cell-in-cell
invasion. Cell 131, 966–979 (2007).
95. Abdu, Y., Maniscalco, C., Heddleston, J. M., Chew, T.-L. & Nance, J. Developmentally
programmed germ cell remodelling by endodermal cell cannibalism. Nat. Cell Biol. 18,
1302–1310 (2016).
96. Lee, Y. et al. Entosis Controls a Developmental Cell Clearance in C. elegans. Cell Rep. 26,
3212-3220.e4 (2019).
97. Chan, K. Y. et al. Skin cells undergo asynthetic fission to expand body surfaces in
zebrafish. Nature 605, 119–125 (2022).
98. Pelzer, D. et al. Ectopic activation of the polar body extrusion pathway triggers cell
fragmentation in preimplantation embryos. doi:10.1101/2022.12.22.521568.
99. Li, X. et al. The C. elegans gonadal sheath Sh1 cells extend asymmetrically over a
differentiating germ cell population in the proliferative zone. eLife 11, e75497 (2022).
100. Stiernagle, T. Maintenance of C. elegans. WormBook Online Rev. C Elegans Biol. 1–11
(2006) doi:10.1895/wormbook.1.101.1.
101. Edelstein, A., Amodaj, N., Hoover, K., Vale, R. & Stuurman, N. Computer control of
microscopes using µManager. Curr. Protoc. Mol. Biol. Chapter 14, Unit14.20 (2010).
102. Kamath, R. S. & Ahringer, J. Genome-wide RNAi screening in Caenorhabditis elegans.
Methods San Diego Calif 30, 313–321 (2003).
103. Rual, J.-F. et al. Toward Improving Caenorhabditis elegans Phenome Mapping With an
ORFeome-Based RNAi Library. Genome Res. 14, 2162–2168 (2004).
104. Schindelin, J. et al. Fiji: an open-source platform for biological-image analysis. Nat.
Methods 9, 676–682 (2012).
105. Preibisch, S., Saalfeld, S. & Tomancak, P. Globally optimal stitching of tiled 3D microscopic
image acquisitions. Bioinformatics 25, 1463–1465 (2009).
106. Mok, D. Z. L., Sternberg, P. W. & Inoue, T. Morphologically defined sub-stages of C.
elegans vulval development in the fourth larval stage. BMC Dev. Biol. 15, 26 (2015).
882
883

Figure 1. Rac pathway *ced* mutants have gonad bifurcation with moderate penetrance.

Figure 2. Bifurcation is a DTC cell-autonomous defect, and a DTC membrane marker reveals somatic gonad cell morphology defects.

Figure 4. The second tip is not led by a cell, but by a DTC fragment

A **RNAi control** *hlh-2(ar623[gfp::hlh-2])* merge tip GFP::HLH-2

B *ced-5* RNAi superficial z-projection deep z-projection

C *ced-12* RNAi superficial z-projection deep z-projection

D *lag-2p::mNG::PH; lag-2(bmd202[lag-2::P2A::H2B::mT2])* DTC membrane Nuclear signal
control

E *ced-5(n1812)*

F WT DTC mutant DTC mutant patch mutant DTC + patch
Area (μM²)

Patch membrane No nuclear signal

Figure 5. The DTC fragment retains INA-1, a key regulator of gonadogenesis.

A *ina-1(qy23[ina-1::mNG])*, DIC merge   INA-1::mNG

RNAi control

B

ced-5 RNAi

C

ced-5 RNAi

Figure 6. The DTC fragment retains the germline stem cell niche
stemness cue LAG-2 in larvae but not adults

Figure 7. Bifurcated germlines are mispatterned and have differentiating germ cells at the distal end

A *mex-5p::H2B::mCherry*

B oocyte, DIC

C Distal tip

D Transition zone

E Meoitic nuclei

B' oocyte histone

F *ced-5(n1812); mex-5p::H2B::mCherry*

G Distal tip

H Meiotic nuclei

I Second tip, DIC

I' Second tip histone

Figure S1. DTC-specific knockdown of *mig-2* Rac1 GTPase but not parallel cell-engulfment pathway *ced* genes causes DTC cell morphology and migration/bifurcation defects.

Figure S1. DTC-specific knockdown of *mig-2/*RhoG GTPase, but not parallel cell-engulfment pathway *ced* genes causes DTC cell morphology and migration/bifurcation defects. Strain expressing a combined *lag-2* promoter-driven membrane marker and *rde-1* rescue transgene *lag-2p::mNG::PLC^{δPH}::F2A::rde-1* in a genetic background that is *rde-1(ne219)* loss of function and *rrf-3(pk1426)* hypersensitive to RNAi. **(A)** On control RNAi (empty vector L4440), one site of strong mNG expression is visible in each gonad arm in late L4 hermaphrodites–the distal tip cell. No expression is observed near the bend of the gonad (inset). **(B)** DTC-specific RNAi knockdown of the *mig-2/*RhoG gene causes a range of cellular and anatomical gonad defects: gonad bifurcation (left), in which mNG expression is always observed on both tips, and the formation of a second "patch" of mNG expression near the bend of an otherwise anatomically normal gonad (right). N=34 total RNAi-treated animals scored, 7/34 had second patch only and an additional 2/34 had a bifurcated gonad. *mig-2* acts upstream of the Rac pathway *ced* genes[40]. **(C)** DTC-specific knockdown of *ced-1* (N=33) and *ced-6* (N=29), members of a parallel engulfment pathway to *ced-10, ced-5, ced-2,* and *ced-12.* No DTC or gonad morphology defects were observed, though in 3/62 samples a delay in dorsal elongation was noted in a single gonad arm. Maximum projection of GFP fluorescence channel through all Z-slices with mNG signal merged with maximum projection of DIC image through slices capturing the gonad tip(s). Imaged at late L4 stage. Visible gonad outlined in black dashed line. Yellow asterisks mark gonad tips of focal gonads, white asterisk marks tip of other gonad arm, yellow arrowhead marks the patch. Autofluorescence of the gut is visible as green punctae; this is unrelated to expression of the fluorescent protein. Scale bars 20 µm.

Figure S2. The nuclear signal is not always found in the distal-most tip.

Figure S2. The nuclear DTC signal is not always found in the distal-most tip. (A) RNAi knockdown of *ced-5* in the strain bearing the *gfp*::*hlh-2* allele. Upper left, maximum projection of GFP fluorescence channel through the more superficial tip of a bifurcated gonad in GFP merged with DIC. Inset right, the more superficial tip has diffuse cytoplasmic GFP::HLH-2 signal, despite falling in the anatomically correct position of a gonad distal tip. Below, maximum projection of GFP fluorescence channel through the deeper tip of same bifurcated gonad merged with DIC. Inset right, the deeper tip has nuclear GFP::HLH-2 signal. N = 1/9 total bifurcated gonads imaged **(B)** RNAi knockdown of *ced-12* in the strain bearing the *gfp*::*hlh-2* allele. Upper left, maximum projection of GFP fluorescence channel through the more superficial tip of a bifurcated gonad in GFP merged with DIC. Inset right, the more superficial tip has diffuse cytoplasmic GFP::HLH-2 signal. Below, maximum projection of GFP fluorescence channel through the deeper tip of same bifurcated gonad merged with DIC. Inset right, the deeper tip has nuclear GFP::HLH-2 signal, despite having made an extra turn towards the viewer. N = 1/10 total bifurcated gonads imaged. Visible gonad outlined in black or white dashed line. Yellow asterisks mark gonad tip with DTC nucleus, yellow arrowhead marks the enucleate patch. Yellow boxes show position of insets in larger image. Autofluorescence of the gut is visible as punctae; this is unrelated to expression of the fluorescent proteins. Imaged at L4 stage. Scale bars 20 μm.

Figure S3. Bifurcated germlines begin differentiating at the second tip in L4 larva

Figure S3. Bifurcated germlines are mispatterned and have differentiating germ cells at the distal end in the L4 stage. Micrographs comparing features of adult germlines between control (A, C, D) and *ced-5(n1812)* (B, E, F) animals. Germ cell nuclei visualized with the *naSi2(mex-5p::H2B::mCherry)* transgene[74], magenta in merge. **(A)** Z-projection through DIC image of L4 control gonad merged with germ cell histone fluorescence. Boxes show positions of insets that follow. **(B)** Z-projection through DIC image of L4 *ced-5(n1812)* gonad merged with germ cell histone fluorescence. Boxes show positions of insets that follow. **(C)** Enlargement of control distal tip fluorescence image showing undifferentiated germ cell nuclei. **(D)** Enlargement of the fluorescence image of the control meiotic transition zone showing its distinctive crescent-shaped nuclear morphology. **(E)** Enlargement of distal tip of *ced-5(n1812)* fluorescence image of gonad distal tip showing undifferentiated germ cell nuclei. **(F)** Enlargement of second tip of *ced-5(n1812)* mutant goand fluorescence image showing crescent-shaped germ cell nuclei characteristic of cells that have entered meiosis. Gonads outlined in black or white dashed lines. Autofluorescence of the gut is visible as punctae; this is unrelated to expression of the fluorescent proteins. Scale bars: 20 µm.