Proteomic and Physiological Signatures of Altitude Adaptation in a *Myrsine coriacea*

Population under Common Garden Conditions

Short title: **Altitude Adaptation in a *Myrsine coriacea* Population**

Roberta Pena da Paschoa[1,2], Vitor Batista Pinto[5], Jéssica Priscilla Pereira[3], Paulo Cezar

Cavatte[3], Mário Luís Garbin[3], Tiago de Oliveira Godinho[4], Lucas Rodrigues Xavier[1,2],

Tatiana Tavares Carrijo[3*], Vanildo Silveira[1,2*]

[1] Laboratório de Biotecnologia, Centro de Biociências e Biotecnologia (CBB), Universidade

Estadual do Norte Fluminense Darcy Ribeiro (UENF), Av. Alberto Lamego, 2000, Campos

dos Goytacazes, RJ, 28013-602, Brazil.

[2] Unidade de Biologia Integrativa, Setor de Genômica e Proteômica, UENF, Brazil.

[3] Universidade Federal do Espírito Santo, Centro de Ciências Exatas, Naturais e da Saúde,

Depto. Biologia, Lab. Botânica, Alto Universitário, Guararema, Alegre, ES, Brazil.

[4] Reserva Natural Vale, Rodovia BR 101, km 122 s/n Zona Rural, Linhares - ES, 29900-111,

Brazil.

[5] Laboratório de Biologia Celular e Tecidual, CBB, UENF, Brazil.

* Correspondence: tcarrijo@gmail.com and vanildo@uenf.br

E-mail addresses:

robertappaschoa@gmail.com

vitorbp@uenf.br

jessicapri.pereira@yahoo.com.br

cavattepc@hotmail.com

mlgarbin@gmail.com

tiago.godinho@vale.com

rxlucas@outlook.com

tcarrijo@gmail.com

vanildo@uenf.br

Date of submission:

Number of tables: 1

Number of Figures: 6

Word count: 2511

Supplementary data: 3 Figures and 1 table

Highlight

This study unveils proteomic and physiological adaptations in a high-altitude *M. coriacea*
population with reduced carbon fixation and energy metabolism.

Abstract

Plants exhibit phenotypic plasticity in response to environmental variations, which can
lead to stable genetic and physiological adaptations if exposure to specific conditions is
prolonged. *Myrsine coriacea* demonstrates this through its ability to thrive in diverse
environments. The objective of the article is to investigate the adaptive responses of *M. coriacea*
by cultivating plants from seeds collected at four different altitudes in a common garden
experiment. Through integrated physiological and proteomic analyses, we identified 170
differentially accumulated proteins and observed significant physiological differences among
the populations. The high-altitude population (POP1) exhibited a unique proteomic profile with
significant down-regulation of proteins involved in carbon fixation and energy metabolism,
suggesting a potential reduction in photosynthetic efficiency. Physiological analyses showed
lower leaf nitrogen content, net CO_2 assimilation rate, specific leaf area, and relative growth
rate in stem height for POP1, alongside higher leaf carbon isotopic composition ($\delta 13C$) and leaf

carbon (*C*) content. These findings provide insight into the complex interplay between

proteomic and physiological adaptations in *M. coriace*a and underscore the importance of local

adaptations. This study enhances our understanding of how altitude-specific selection pressures

can shape plant molecular biology and physiology, offering valuable perspectives for predicting

plant responses to global environmental changes.

Keywords Altitude, Altitudinal gradient, Climate change, Comparative proteomics, *Myrsine*

coriacea, Natural population, Physiological parameters, Primulaceae.

Abbreviation list

A - Net assimilation rate of CO_2

C - Leaf carbon

Ci - Internal CO_2 concentration

CIT - Total chlorophyll

DAPs - differentially accumulated proteins

FASP - filter-aided sample preparation

GO - Gene ontology

gs - Stomatal conductance

KEGG - Kyoto Encyclopedia of Genes and Genomes

Kleaf - Leaf hydraulic conductivity

N - Leaf nitrogen

PCA - principal component analysis

POP1 - Population originating from Dores do Rio Preto, at an altitude of 2019 meters

POP2 - Population originating from Domingos Martins at an altitude of 1229 meters

POP3 - Population originating from Mimoso do Sul at an altitude of 930 meters

POP4 - Population originating from Muqui at an altitude of 657 meters

PPIs - protein-protein interaction

RGR_H - Relative growth rate in height

SLA - Specific leaf area

TCA - trichloroacetic acid

UPGMA - Unweighted Pair Group Method using Arithmetic averages

VD - Vein density

$\delta13C$ - Leaf carbon isotopic composition

Introduction

In a rapidly changing climate, plants face distinct environmental challenges compared to

those experienced by their ancestors in recent evolutionary history. One way in which plants

respond to these changes is through environmentally induced changes in phenotype (Mendonça

et al., 2019; Pérez-Ramos *et al.*, 2019). Plant species can adjust to these new conditions through

phenotypic plasticity, adapt through natural selection, or disperse and establish in areas with

conditions similar to those where they are adapted, and these options are not mutually exclusive

(Crispo, 2008; Malhi *et al.*, 2010).

Understanding how plants respond to environmental changes is essential for

understanding the processes of plant adaptation and speciation. Altitude gradients are

particularly intriguing due to their pronounced changes in various physical environmental

characteristics, such as temperature, atmospheric pressure, humidity, sunlight hours, ultraviolet

radiation, wind, season duration, and geology (Korner, 2007; Chen *et al.*, 2014; Ahmad *et al.*,

2016; Rahman *et al.*, 2019). Altitude gradients present challenges for successful plant

adaptation. As altitude increases, it is observed that certain plant species are replaced by others,

giving rise to different vegetation types. Some species are restricted to a specific area, while

others grow across a wide range of altitudes, adapting to particular conditions through

phenotypic plasticity and genetic modification (Clausen et al., 1940; Hovenden and Vander

Schoor, 2004; Byars et al., 2007).

Plant populations occurring along an altitudinal gradient can vary in their morpho-

anatomical and physiological characteristics (Ahmad *et al.*, 2016; Rahman *et al.*, 2020). While

physiological variations occurring in plants with increasing altitude are well described (Ahmad *et al.*, 2016; Hniličková *et al.*, 2016; Ibañez *et al.*, 2017; Liu *et al.*, 2020; Lim and Burns, 2022), the underlying processes leading to these phenotypic variations are not clearly understood. The continuous challenge lies in identifying the molecular pathways and genes of adaptive significance in this context. Our understanding remains incomplete regarding the specific genes and proteins intricately involved in the local adaptation of diverse populations within tropical species. Consequently, the translation of genetic variations into phenotypic adaptations across varied tropical environments still contains gaps in information. Common garden experiments help in understanding local adaptation by enabling the assessment of the relative importance of plastic and molecular responses within the same species (de Villemereuil *et al.*, 2016)

Proteomics can significantly contribute to the elucidation of the molecular mechanisms involved in species adaptation to diverse environments (Diz *et al.*, 2012; Li *et al.*, 2014; Ma *et al.*, 2015; Voelckel *et al.*, 2017). A comprehensive understanding of how proteins are regulated in species with a wide distribution range holds the potential to provide valuable insights into the impact of environmental changes on organisms, the evolutionary processes shaping proteins over time, and the strategies employed by organisms to adapt to their specific environmental conditions.

Myrsine coriacea is an angiosperm with a broad capacity to inhabit different environments (Sánchez-Tapia *et al.*, 2018), exhibiting substantial phenotypic variation (Pipoly, 1991; Ricketson and Pipoly III, 1997; Carrijo *et al.*, 2017). In the Atlantic Forest, the species can be found from coastal ecosystems to high-altitude Campos (Carrijo *et al.*, 2017). The genetic (Paschoa *et al.*, 2018), morphological, and physiological (Pereira *et al.*, 2022) diversity in populations distributed across an altitudinal gradient in the Atlantic Forest seems to indicate high phenotypic plasticity, considering the demonstrated capacity of these plants to acclimate to diverse environmental conditions. A better understanding of the influence of the relative roles of genetic and environmental factors on phenotypic variation in *M. coriacea* helps to understand the potentially adaptive phenotypic variation in plants in nature and the adaptive and survival strategies of populations.

In this study, we subjected plants germinated from *M. coriacea* seeds originating from natural populations distributed across an altitudinal gradient to common garden conditions. The objective of the article is to investigate the adaptive responses of *M. coriacea* by cultivating plants from seeds collected at four different altitudes in a common garden experiment. Our

findings provide new insights regarding the molecular mechanisms involved in the local
adaptation of different populations of *M. coriacea*.

Materials and Methods

Seed collection and common garden establishment

The seeds were collected from natural populations located in four municipalities in
Espírito Santo State, Brazil, along a 1,517 m altitude gradient (Table 1). Ripe fruits were
collected in November 2017 from four branches of ten selected adult trees in each population.
After collection, the fruits were subjected to mechanical scarification using a steel sieve and
dried in the shade. Then, the seeds were immediately taken to a greenhouse located in Jerônimo
Monteiro-ES (20º47'45''/41º24'18''). In the nursery, the seeds were planted in polypropylene
trays with a capacity of 19.5 cm³ containing washed sand substrate. In March 2018, the
seedlings were transplanted into polypropylene tubes with a capacity of 280 cm³ filled with a
commercial substrate Tropstrato, HT Hortaliças (Vida Verde®, SP, Brazil). The seedlings were
kept for 12 months under shade cloth with irradiance levels reduced by 50% and automated
micro-sprinkler irrigation activated three times during the daytime period. In March 2019, the
seedlings were transferred to three-liter pots containing a substrate composed of a 3:1 ratio of
the same commercial substrate and sand, where they remained until the setup of the field
experiments.

A common garden was established in December 2019 in the municipality of Marechal
Floriano (Espírito Santo, Brazil) (20°25'38.41"/40°51'29.08") located at approximately 914 m
altitude. The planting was conducted in a randomized block design, with three blocks
containing ten plants from each population, totaling 40 plants per block. A total of 87 mixed
seedlings were planted in the borders, spaced 2 meters within rows and 3 meters between rows,
in a total area of 1.482 m^2.

Each block consisted of five rows containing eight individuals (two from each
population). The seedlings were planted with a spacing of two meters within the rows and three
meters between rows. Planting was carried out in pits with dimensions of 30x30x30 cm, where
200 g of superphosphate (19% P) was added 15 days before planting. To ensure the
establishment of the seedlings, regular irrigations were performed using manual watering cans.
Weed control was carried out through manual weeding and chemical targeted application of the

herbicide Roundup Original® (Bayer CropScience, SP, Brazil) Whenever necessary, ant control

was carried out using the ant insecticide (Mirex-SD®, SP, Brazil)

Proteomic analysis

Protein extraction and digestion

To trace the proteomic profile of the populations, expanded leaves located throughout the

canopy were collected from three individuals per population in each block seven months after

the common garden was established. The samples collected in the field were stored in sealed

paper bags and covered with silica gel. The leaf collection time started at 12:00 PM and finished

at 2:00 PM, with an average temperature of 25 °C, relative humidity of 57%, and wind speed

of 10 km/h. Subsequently, the samples were wrapped in aluminum foil and stored at -80 °C

until proteomic analysis.

For protein extraction, the three individuals collected in each population within each

block were pooled and macerated under liquid nitrogen to comprise each biological replicate.

protein extraction. Proteins were extracted from 300 mg of powered fresh matter (FM) aliquots

using the trichloroacetic acid (TCA)/acetone precipitation method described by Damerval et

al., (1986), with modifications as described by Passamani *et al.,*(2017). The protein

concentration was determined using the 2-D Quant Kit (Cytiva, Marlborough, MA, USA).

Before the trypsin digestion step, aliquots of 100 µg of proteins were precipitated using

the methanol/chloroform methodology to remove any interferents from the samples (Nanjo *et*

al., 2012). After precipitation, the samples were resuspended in a 7 M urea/2 M thiourea buffer,

and tryptic digestion of proteins (V5111; Promega, Madison, WI, USA; final enzyme-to-protein

ratio of 1:100) was performed using the filter-aided sample preparation (FASP) method as

described by Wiśniewski et al., (2009) with modifications by Reis *et al*., (2021). The resulting

peptides were quantified using the protein and peptide method at 205 nm using a NanoDrop

2000c spectrophotometer (Thermo Fisher Scientific, Waltham, USA).

Mass spectrometry analysis

Mass spectrometry was performed using a nanoACQUITY ultra-performance liquid

chromatograph (UPLC) coupled to a Q-TOF SYNAPT G2-Si instrument (Waters, Manchester,

UK) as described by Reis *et al.,* (2021). The runs consisted of three biological replicates of 2.0

µg of peptides. During separation, the samples were loaded onto a nanoAcquity UPLC M-Class

Symmetry C18 5 µm trap column (100 Å, 5 µm, 180 µm × 20 mm, 2D; Waters) at a flow rate

of 5 µL/min for 3 min and then onto an analytical reversed-phase nanoAcquity M-Class HSS

T3 1.8 µm column (100 Å, 1.8 µm, 75 µm × 150 mm; Waters) at a flow rate of 400 nL/min.

The column temperature was set to 45 °C. A binary gradient was used for peptide elution, where

mobile phase A consisted of water (Tedia, Fairfield, Ohio, USA) with 0.1% formic acid (Sigma-

Aldrich), and mobile phase B consisted of acetonitrile (Sigma-Aldrich) with 0.1% formic acid.

The gradient elution was performed as follows: 5% B for 3 min, increasing from 5 to 41% B

over 92.00 min, increasing from 41 to 97% B over 96.00 min, holding at 97% B for 100.00 min,

and decreasing to 5% B by 102.00 min.

Mass spectrometry was performed in positive mode and resolution mode (mode V), with

35,000 full widths at half maximum (FWHM) and ion mobility, and in data-independent

acquisition (DIA) mode. The ion mobility separation (IMS) used an IMS wave velocity of 800

m s^{-1} (HDMS^E); the transfer collision energy increased from 19 to 55 V in high-energy mode;

the cone and capillary voltages were 30 V and 3000 V, respectively; and the source of

temperature was 100 °C. For time-of-flight (TOF) parameters, the scan time was set to 0.5 s in

continuum mode, and the mass range was 50–2000 Da. Human [Glu1] fibrinopeptide B at 100

fmol µL−1 was used as an external calibrant, and lock mass acquisition was performed every

30 s. Mass spectrum acquisition was performed by MassLynx software (version 4.1, Waters).

224   Proteomic data analysis

Spectra processing and database search conditions were performed using ProteinLynx

Global SERVER (PLGS) software (version 3.02, Waters). The HDMS^E analysis followed the

following parameters: Apex3D of 150 counts for low-energy threshold; 50 counts for elevated-

energy threshold; 750 counts for intensity threshold; one missed cleavage; minimum fragment

ions per peptide equal to three; minimum fragment ions per protein equal to seven; minimum

peptides per protein equal to two; fixed modifications of carbamidomethyl (C) and variable

modifications of oxidation (M) and phosphoryl (STY); default false discovery rate (FDR) of

1%; automatic peptide and fragment tolerance.

To select the proteome of the species phylogenetically closest to *M. coriacea*, a

phylogenetic tree was constructed using the PhyloT website (https://phylot.biobyte.de/) based

on the National Center for Biotechnology Information (NCBI) taxonomy. The proteome of the species *Actinidia chinensis* (ID: UP000241394), available on UniProtKB (https://www.uniprot.org/), was used, as it is the species phylogenetically closest to *M. coriacea*.

Label-free quantification analysis was performed using ISOQuant software v.1.8 (Distler *et al.*, 2014, 2016). The parameters used were as follows: peptide and protein FDR 1%; sequence length of at least six amino acid residues; and minimum peptide score equal to six. Samples were normalized by a multidimensional normalization process, which corrects peak intensities based on the intensity and retention time domains. The software performed relative protein quantification based on the TOP3 method. Based on the relative abundances of uniquely assigned peptides, the abundances of shared peptides were redistributed to the respective source proteins followed by TOP3-based quantification (Distler *et al.*, 2014).

For the comparative analysis, only the proteins present or absent (for unique proteins) in the three biological replicates were accepted for differential abundance analysis. Comparative analyses were performed on POP1 relative to the others. The following comparisons were performed: POP1/POP2, POP1/POP3 and POP1/POP4. Data were analyzed by Student's t-test (two-tailed). Proteins with a p-value < 0.05 were deemed up-accumulated if the Log_2 of the fold change (FC) > 0.6 and down-accumulated if the Log_2 of the FC was less than < -0.6. The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE (Perez-Riverol *et al*., 2022) partner repository with the dataset identifier PXD047424. Functional enrichment analysis, including Gene Ontology (GO) analysis and Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway analysis, was conducted on differentially accumulated proteins (DAPs) using OmicsBox software 3.0.29 (https://www.biobam.com/omicsbox).

The interaction networks of DAPs used the first level of interaction retrieved by STRING version 11.5 (http://string-db.org) search. To generate a protein-protein interaction (PPIs) network, Actinidia chinensis was considered the reference plant species using a minimum required interaction score of 0.7.

To visualize the relationships among the populations, a heatmap plot was generated using the gplots package (Warnes *et al.*, 2022) and a principal component analysis (PCA) was performed using the prcomp function in package stats in R (R Core Team, 2023).

Physiological analysis

Physiological analysis of the populations was performed 13 months after the common garden was established using two plants per population in each block, resulting in 6 biological replicates per population.

For growth analysis, plant height was measured using a graduated ruler (precision of 0.1 cm). Height was determined from ground level to the apical bud of the stem. Based on this measurement, the relative growth rate of stem height (RGR_H, mm.mm^{-1}dia^{-1}) was calculated with a three-month interval between the two measurements.

To determine the chlorophyll index, four fully expanded leaves located throughout the canopy were selected, and the analysis was performed using ClorofiLOG (CFL 1030, Falker, Rio Grande do Sul, BR). The average value for each plant was used for analysis. Leaf hydraulic conductance ($Kleaf$, mmol m^{-2}s^{-1}. MPa^{-1}) was determined using one leaf per plant following the method described by Martins *et al.,*(2019): $Kleaf$ = E* / (Ψw pd - Ψw pm-dark), where E is the transpirational flux and Ψw is the leaf water potential. Leaf vein density (VD; mm mm^{-2}) was measured using two leaves per plant, following the protocol outlined by Pereira *et al.,*(2022).

Gas exchange and chlorophyll fluorescence analyses were performed using an infrared gas analyzer (IRGA, Li 6400XT, Li-Cor, Lincoln, USA) equipped with a 2 cm^2 leaf chamber (Li 6400-40 LCF, Li-Cor, Lincoln, EUA). The measurements of net CO_2 assimilation (*A,* μmol CO_2 m^{-2} s^{-1}), stomatal conductance (*gs,* mol H_2O m^{-2} s^{-1}) and internal CO_2 concentration CO_2 (*Ci*, μmol CO_2 m^{-1}) were conducted between 8:30-10:30 under saturating photosynthetically active radiation of 1.000 µmol m^{-2} s^{-1}, a partial pressure of CO_2 of 400 ppm, and a block temperature of 28 °C.

Ten leaves were collected for specific leaf area (*SLA*; cm^2 g^{-1}) determination and carbon isotopic composition (*δ13C*; ‰) according to Cavatte *et al.,*(2011). Total leaf carbon (*C*) and leaf nitrogen (*N*) concentrations were determined using an elemental analyzer (Carlo Erba, Milan, Italy), while isotopic compositions were measured using a stable isotope ratio mass spectrometer (IRMS Delta Plus, Finnigan Mat, San Jose, USA) at the Center for Nuclear Energy in Agriculture, University of São Paulo. The equations proposed by Farquhar *et al.,*(1989) were used to calculate *δ13C* using the Vienna Pee Dee Belemnite as the standard.

The physiological data were analyzed using a linear model with a source of variation (POP 1 to 4) and one variable at a time (for both physiological variables and isotope

measurements). Subsequently, the estimated means of the populations were compared by
overlaying confidence intervals (95% confidence intervals), where the lack of overlap indicated
significant differences between the means. Finally, the results were presented graphically with
bar charts and their respective confidence intervals. All analyses were performed using R (R
core Team, 2023), using the function emmeans from the package emmeans (Russell V. Lenth,
2023).

Results

Proteomic analysis

The comparative proteomic analysis allowed the identification of 590 proteins in the *M.*
coriacea samples from the four populations included in the experimental planting. Comparative
analysis between POP1 and the other populations revealed that 170 DAPs (Supplementary
Table S1). Heatmap analysis, using the unweighted pair group method using arithmetic
averages (UPGMA) algorithm based on the total ion count of proteins in the biological
replicates of POP1, POP2, POP3, and POP4, allowed visualization of the separation into two
major groups, one comprising POP1 and the other consisting of the remaining populations. This
analysis revealed that POP1 exhibited a distinct protein profile, with most proteins displaying
lower total ion count values (Fig. 1A).

Additionally, POP1 distinguished itself from the other studied populations in the PCA
analysis (Fig. 1B). Each biological replicate for a specific population was evaluated, and their
relationship is indicated by an enclosed circle. The first two eigenvalues of PCA explained
54.9% of the total variation, with 37.2% explained by component 1 and 17.7% by component
2.

In Gene Ontology (GO) analysis, the most representative biological processes in all
populations were the "cellular process," "biosynthetic process," "response to cadmium ion," and
"generation of precursor metabolites and energy." Many proteins were found to be down-
accumulated in POP1 (Supplementary Fig. S1).

The KEGG pathway analysis revealed that the most representative pathway in all three
comparisons was "Carbon fixation in photosynthetic organisms," followed by
"glycolysis/gluconeogenesis" for the POP1/POP2 and POP1/POP3 comparisons and "cysteine
and methionine metabolism" for the POP1/POP4 comparison (Supplementary Fig. S2).

The predicted analysis of protein-protein interactions (PPIs) resulted in interactions

among potential key proteins involved in the differential regulation of POP1/POP2 (Fig. 2),

POP1/POP3 (Fig. 3), and POP1/POP4 (Fig. 4) comparisons.

PPIs among Triosephosphate isomerase (A0A2R6RWY2), Glucose-6-phosphate

isomerase (A0A2R6R2W5), Fructose-bisphosphate aldolase (A0A2R6PR43), and

Phosphoglycerate kinase (A0A2R6PR64) were visualized in the comparisons between

populations (Fig. 2, 3, and 4; highlighted arrows). These proteins are involved in the pathways

of "Carbon fixation in photosynthetic organisms" and "Glycolysis/gluconeogenesis" and are

down-accumulated in POP1 compared to the other populations. There was the exception of the

POP1/POP2 comparison, which was unchanged, but also showed a lower accumulation in

POP1 in this comparison. Chlorophyll a-b binding proteins (A0A2R6Q6C6, A0A2R6RTT1,

A0A2R6RG56) that were also highlighted in the interaction network, participate in the

"photosynthesis-antenna proteins" pathway. These proteins are up-accumulated in POP1

compared to the other populations (Fig. 2, 3, and 4; highlighted arrows).

Physiological parameters

The analysis of physiological parameters revealed significant differences among the

populations (Fig. 5, 6 and Supplementary Fig. S3). $\delta 13C$ (Fig. 5A) and C (Fig. 5B) were

significantly different in POP1 compared to the other populations, with POP1 showing higher

values. POP1 exhibited the lowest values of N (Fig. 5C) and RGR_H (Fig. 5D) compared to POP2

and POP4. In $Kleaf$ (Fig. 6A), A (Fig. 6B), and SLA (Fig. 6C), there was also a significant

difference between POP1 and POP2, with POP1 showing lower values. There were no

significant differences among all four populations for gs, CIT, VD, and Ci (Supplementary Fig.

S3). POP2, POP3, and POP4 were statistically similar in all physiological analyses.

Discussion

Proteomic analysis revealed differential regulation of proteins among the four populations

originating from the altitudinal gradient and subjected to a common garden condition. Based

on the cluster analysis with a heatmap and PCA (Fig. 1) of the identified proteins, it was possible

to separate the natural populations into two main groups, one containing the highest altitude

population (POP1) and the other comprising the remaining populations at lower altitudes. Our

results also revealed a differential pattern for POP1 based on physiological analyses (Fig. 5 and

6). These results indicate that the high-altitude population (POP1) (Table 1), exhibits distinct

molecular and physiological adaptations when compared to the other populations. This

distinction suggests a specific local adaptation of POP1 to the environmental conditions unique

to its original high-altitude habitat, which becomes particularly evident when these populations

are subjected to common garden conditions at 914 m altitude.

POP1 was separated from the other populations, likely due to the edaphoclimatic

differences to which the population is exposed in its original environment. According to the

latest update of the Köppen and Geiger climate classification by Alvares *et al.,*(2013), the

region of POP1 is characterized as a tropical highland climate, situated at approximately 2019

m altitude, where temperatures can drop to -4 °C in winter. In contrast, the regions of origin of

the other populations, POP2 and POP3, are classified as having a warm temperate climate,

while POP4 originates from a rainy tropical climate. These latter regions, in comparison to

POP1, are marked by lower altitudes and higher temperatures year-round.

Our comparative proteomic analysis revealed that 'carbon fixation in photosynthetic

organisms' and 'glycolysis/gluconeogenesis' were the most representative pathways in POP1,

in comparison to the other populations. This was evidenced by a notable down-accumulation

of many related proteins (Supplementary Fig. S2; Supplementary Table S1). These results

suggest that POP1 exhibits differential protein regulation, particularly in photosynthesis

pathways, when subjected to common garden conditions. Natural populations of *M. coriacea*

in their native environment showed differences in morphological and stomatal characteristics

due to altitude, affecting the photosynthetic capacity of the species at higher altitudes (Pereira

et al., 2022). Individuals from the high-altitude population (POP1; 2019 m) had leaves with

higher vein density (*VD*), trichome density, and leaf carbon isotopic composition (*δ13C*) but

lower guard cell length, maximum stomatal conductance, and specific leaf area (*SLA*) compared

to the other populations. In our study, *δ13C* also exhibited higher values, and *SLA* showed lower

values in POP1 than in the other populations under common garden conditions. However, *VD*

did not differ among populations. These results suggest that the species *M. coriacea* is capable

of local adaptation in response to specific environmental conditions, such as altitude, through

changes in morphological and stomatal characteristics. The high-altitude population (POP1)

may have developed specific adaptations, such as increased trichome density, higher *δ13C*

levels, and lower *SLA*, to cope with the conditions of high altitude. These adaptations could

affect the species' photosynthetic capacity of the at higher altitudes.

393   Additionally, alongside the higher values of $\delta13C$ and C content, POP1 exhibited lower

394 *Kleaf* and *A*), suggesting that plants from this population may have reduced their stomatal

395 opening to minimize water loss. $\delta13C$ is an important indicator of the environmental conditions

396 in which the plant has grown. Soil water availability and climatic conditions are factors that

397 affect this composition, with drought leading to a higher proportion of heavy carbon isotopes

398 in the leaf (Vantyghem *et al.*, 2022). This is because the plant tends to reduce stomatal opening

399 to avoid excessive water loss, which can result in lower leaf hydraulic conductance (*Kleaf*)

400 (Jákli *et al.*, 2016). Additionally, drought can lead to a reduction in the net assimilation rate of

401 CO_2 (*A*) (Fresneau *et al.*, 2007) and an increased internal concentration of CO_2 (*Ci*) in the leaf

402 (Nunes *et al.*, 1968), indicating lower photosynthetic efficiency in terms of carbon fixation.

403   The down-accumulation of proteins related to energy metabolism might negatively

404 impact the photosynthetic process in the high-altitude population POP1, when compared to the

405 other populations. In a controlled greenhouse experiment with *Quercus ilex* from different

406 altitude ranges, a differential proteomic profile among populations was revealed, with the high-

407 altitude population exhibiting a down-regulation of proteins related to energy metabolism (San-

408 Eufrasio *et al.*, 2021). In our study, we identified down-accumulated proteins in POP1, when

409 compared to the other populations, that were involved in energy metabolism, such as

410 Triosephosphate isomerase (A0A2R6RWY2), Glucose-6-phosphate isomerase

411 (A0A2R6R2W5), Fructose-bisphosphate aldolase (A0A2R6PR43), and Phosphoglycerate

412 kinase (A0A2R6PR64). These proteins were found to participate in predicted protein-protein

413 interactions in the performed comparisons (Fig. 2, 3, and 4).

414   The decrease in proteins involved in energy metabolism suggests that POP1 has a

415 different energy yield compared to the other populations, consequently reducing its

416 photosynthetic capacity. This hypothesis is consistent with our physiological analysis results in

417 the studied populations. Physiological analyses revealed that POP1 exhibited higher values of

418 *C* and $\delta13CI$, and lower values of *A* and relative stem height growth rate (*RGRH*). Increasing

419 temperature leads to an increase in the carboxylation rate and a decrease in the specificity of

420 RUBISCO (ribulose-1,5-bisphosphate carboxylase/oxygenase) for CO_2 (Brooks and Farquhar,

421 1985). It has been demonstrated that leaves with higher $\delta13C$ values exhibit lower rates of

422 photosynthesis and stomatal conductance than those with lower values (Vitoria *et al.*, 2016).

423 Thus, photosynthesis and growth are affected by temperature, which limits biochemical

424 processes (Cooper and Taiton, 1968).

425   Nitrogen assimilation is a highly energy-demanding process (Bloom *et al.*, 1992). Once

426 absorbed, nitrogen can be assimilated in the root itself or transported to the leaves, where

427 assimilation takes place (Lee *et al.*, 1992). Since POP1 exhibited down-regulation of proteins

428 involved in energy metabolism, indicative of different energy yields compared to the other

429 populations and a reduced photosynthetic capacity, it may also interfere with nitrogen

430 assimilation, as evidenced by a lower leaf nitrogen concentration (N) in POP1 (Fig. 5C).

431   Energy deficit is a primary symptom of plants under stress (Baena-González *et al.*, 2007).

432 In this study, an up-regulation of Chlorophyll a-b binding proteins (A0A2R6RG56,

433 A0A2R6RTT1, A0A2R6Q6C6) was identified in POP1, when compared to the other

434 populations. The expression of Chlorophyll a-b binding protein encoding genes is affected by

435 light intensity, low temperature, high salinity, drought, and disease (Humbeck and Krupinska,

436 2003; Caffarri *et al.*, 2005; Umate, 2010; Xia *et al.*, 2012; Wang *et al.*, 2017). Regulating the

437 levels of Chlorophyll a-b binding proteins allows chloroplasts to respond flexibly and rapidly

438 to environmental stresses (Li *et al.*, 2020). The composition of pigments is altered to adapt to

439 specific environmental conditions and when a plant is exposed to lower irradiation, it produces

440 more chlorophyll to absorb light and capture more energy (Arena *et al.*, 2020). The up-

441 regulation of these Chlorophyll a-b binding proteins in POP1, may be related to the plants' need

442 to respond to environmental stress, such as lower irradiation and higher temperature compared

443 to their original location.

Conclusion

446   The populations studied displayed distinct proteomic and physiological profiles, and the

447 high-altitude population (POP1) demonstrating particularly unique characteristics. In POP1, a

448 notable decrease in proteins involved in energy metabolism was observed, including

449 Triosephosphate isomerase (A0A2R6RWY2), Glucose-6-phosphate isomerase

450 (A0A2R6R2W5), Fructose-bisphosphate aldolase (A0A2R6PR43), and Phosphoglycerate

451 kinase (A0A2R6PR64). This population also showed lower concentrations of N, A, SLA and

452 RGR_H, coupled with higher concentrations of $\delta 13C$ and C, indicating a reduced photosynthetic

453 capacity in POP1 under common garden conditions.

454   Furthermore, the up-regulation of Chlorophyll a-b binding proteins (A0A2R6RG56,

455 A0A2R6RTT1, A0A2R6Q6C6) in POP1 suggests an adaptation to environmental stress, which

456 is significant in the acclimation process of high-altitude populations of *M. coriacea*. The distinct

local adaptations in traits such as N, A, SLA, RGR_H, $\delta13C$, and C, and the observed differentially accumulated proteins related to energy metabolism underscore the role of abiotic selection pressures in the high-altitude environment in driving the proteomic and physiological changes observed in POP1, even under common garden conditions.

These findings highlight the importance of understanding local adaptations in response to abiotic stresses to better predict how *M. coriacea* populations might respond to climate change. Such knowledge is crucial for predicting their survival and genetic diversity in changing environmental conditions.

Supplementary data

Supplementary Fig. S1. Functional classification based on Gene Ontology (GO) of DAPs during comparative proteomic analysis between natural populations of *M. coriacea*.

Supplementary Fig. S2. KEGG pathway analysis of DAPs during comparative proteomic analysis between natural populations of *M. coriacea*.

Supplementary Fig. S31. Physiological parameters among the natural populations of *M. Coriacea*.

Supplementary Table S1. Complete list of proteins identified in the comparison of four populations of *M. coriacea*.

Acknowledgements

RPP is grateful for the scholarship funding provided by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), and RLX acknowledges the scholarship funding from the Fundação de Amparo à Pesquisa do Estado do Rio de Janeiro (FAPERJ). We extend our sincere thanks to the Laboratório de Botânica at the Universidade Federal do Espírito Santo for their invaluable assistance with seed collection, physiological analysis support, and the establishment of the common garden.

Author contributions

RPP, TCC, PCC, and VS conceived and coordinated the research. RPP performed data collection, analysis, and interpretation and wrote the draft of the manuscript. VBP performed data collection and analysis. JPP performed data collection and analysis. PCC, MLG, TOG, and

TTC set up the field experimental design. LRX assisted in data analysis and interpretation. All

authors revised and edited the final version of the manuscript.

Conflict of interest No conflicts of interest.

Funding

This research was supported by Fundação de Amparo à Pesquisa do Estado do Rio de Janeiro

(FAPERJ) (Proc E-26/211.310/2021; Proc E26/200.999/2021) and Conselho Nacional de

Desenvolvimento Científico e Tecnológico (CNPq) (307755/2019-3), awarded to VS;

Fundação Estadual de Amparo à Pesquisa do Estado do Espírito Santo (FAPES/VALE

#525/2016, PROCAP #08/2017, PROFIX #10), VALE (525/2016), and CNPq (311551/2022-

0), awarded to TTC; and CNPq (311522/2022-0) awarded to MLG. Coordenação de

Aperfeiçoamento de Pessoal de Nível Superior, Brazil (CAPES), Finance Code 001.

Data availability

The mass spectrometry proteomics data have been deposited to the ProteomeXchange

Consortium via the PRIDE partner repository with the dataset identifier PXD047424. The list

of all identified proteins is available in the supplementary material.

References

Ahmad KS, Hameed M, Fatima S, Ashraf M, Ahmad F, Naseer M, Akhtar N. 2016. Morpho-anatomical and physiological adaptations to high altitude in some Aveneae grasses from Neelum Valley, Western Himalayan Kashmir. Acta physiologiae plantarum 38, 1–14.

Alvares CA, Stape JL, Sentelhas PC, Gonçalves JL de M, Sparovek G. 2013. Köppen's climate classification map for Brazil. Meteorologische Zeitschrift 22, 711–728.

Arena ME, Pastur GM, Lencinas MV, Soler R, Bustamante G. 2020. Changes in the leaf nutrient and pigment contents of *Berberis microphylla* G. Forst. in relation to irradiance and fertilization. Heliyon 6, e03264.

Baena-González E, Rolland F, Thevelein JM, Sheen J. 2007. A central integrator of transcription networks in plant stress and energy signalling. Nature 448, 938–942.

Bloom AJ, Sukrapanna SS, Warner RL. 1992. Root respiration associated with ammonium and nitrate absorption and assimilation by barley. Plant Physiology 99, 1294–1301.

Brooks A, Farquhar GD. 1985. Effect of temperature on the CO_2/O_2 specificity of ribulose-1, 5-bisphosphate carboxylase/oxygenase and the rate of respiration in the light. Planta 165, 397–406.

Byars SG, Papst W, Hoffmann AA. 2007. Local adaptation and cogradient selection in the alpine plant, *Poa hiemata*, along a narrow altitudinal gradient. Evolution: International Journal of Organic Evolution 61, 2925–2941.

Caffarri S, Frigerio S, Olivieri E, Righetti PG, Bassi R. 2005. Differential accumulation of Lhcb gene products in thylakoid membranes of *Zea mays* plants grown under contrasting light and temperature conditions. Proteomics 5, 758–768.

Carrijo TT, Tuler AC, Luber J, Costa RG da, Santos MC dos, Paschoa RP da, Freitas M de F. 2017. Flora do Espírito Santo: Primulaceae. Rodriguésia 68, 1829–1856.

Cavatte PC, Oliveira ÁAG, Morais LE, Martins SC V, Sanglard LMVP, DaMatta FM. 2011. Could shading reduce the negative impacts of drought on coffee? A morphophysiological analysis. Physiologia plantarum 144, 111–122.

Chen B, Zhang X, Tao J, Wu J, Wang J, Shi P, Zhang Y, Yu C. 2014. The impact of climate change and anthropogenic activities on alpine grassland over the Qinghai-Tibet Plateau. Agricultural and Forest Meteorology 189, 11–18.

Clausen J, Keck DD, Hiesey WM. 1940. Experimental studies on the nature of species. 1, Effect of varied environments on western North American plants. *in press*.

Cooper JP, Taiton NM. 1968. Light and temperature requirements for the growth of tropical and temperate grasses.

Crispo E. 2008. Modifying effects of phenotypic plasticity on interactions among natural selection, adaptation and gene flow. Journal of evolutionary biology 21, 1460–1469.

Damerval C, de Vienne D, Zivy M, Thiellement H. 1986. Technical improvements in two-dimensional electrophoresis increase the level of genetic variation detected in wheat-seedling proteins. Electrophoresis 7, 52–54.

de Villemereuil P, Gaggiotti OE, Mouterde M, Till-Bottraud I. 2016. Common garden experiments in the genomic era: new perspectives and opportunities. Heredity 116, 249–254.

Distler U, Kuharev J, Navarro P, Levin Y, Schild H, Tenzer S. 2014. Drift time-specific collision energies enable deep-coverage data-independent acquisition proteomics. Nature methods 11, 167–170.

Distler U, Kuharev J, Navarro P, Tenzer S. 2016. Label-free quantification in ion mobility–enhanced data-independent acquisition proteomics. Nature protocols 11, 795–812.

Diz AP, Martínez-Fernández M, Rolán-Alvarez E. 2012. Proteomics in evolutionary ecology: linking the genotype with the phenotype. Molecular ecology 21, 1060–1080.

Farquhar GD, Ehleringer JR, Hubick KT. 1989. Carbon isotope discrimination and photosynthesis. Annual review of plant biology 40, 503–537.

Fresneau C, Ghashghaie J, Cornic G. 2007. Drought effect on nitrate reductase and sucrose-phosphate synthase activities in wheat (Triticum durum L.): role of leaf internal CO_2. Journal of Experimental Botany 58, 2983–2992.

Hniličková H, Kuklová M, Hnilička F, Kukla J. 2016. Effect of altitude and age of stands on physiological response of three dominant plants in forests of the Western Carpathians. Plant, Soil and Environment 62, 341–347.

Hovenden MJ, vander Schoor JK. 2004. Nature vs nurture in the leaf morphology of Southern beech, *Nothofagus cunninghamii* (Nothofagaceae). New Phytologist 161, 585–594.

Humbeck K, Krupinska K. 2003. The abundance of minor chlorophyll a/b-binding proteins CP29 and LHCI of barley (*Hordeum vulgare* L.) during leaf senescence is controlled by light. Journal of Experimental Botany 54, 375–383.

Ibañez VN, Berli FJ, Masuelli RW, Bottini RA, Marfil CF. 2017. Influence of altitude and enhanced ultraviolet-B radiation on tuber production, seed viability, leaf pigments and morphology in the wild potato species *Solanum kurtzianum* Bitter & Wittm collected from an elevational gradient. Plant Science 261, 60–68.

Jákli B, Tränkner M, Senbayram M, Dittert K. 2016. Adequate supply of potassium improves plant water-use efficiency but not leaf water-use efficiency of spring wheat. Journal of Plant Nutrition and Soil Science 179, 733–745.

Körner C. 2007. The use of 'altitude' in ecological research. Trends in ecology & evolution 22, 569–574.

Lee RB, Purves J v, Ratcliffe RG, Saker LR. 1992. Nitrogen assimilation and the control of ammonium and nitrate absorption by maize roots. Journal of Experimental Botany 43, 1385–1396.

Li X, Yang Y, Ma L, Sun X, Yang S, Kong X, Hu X, Yang Y. 2014. Comparative proteomics analyses of *Kobresia pygmaea* adaptation to environment along an elevational gradient on the central Tibetan Plateau. PloS one 9, e98410.

Li X-W, Zhu Y-L, Chen C-Y, Geng Z-J, Li X-Y, Ye T-T, Mao X-N, Du F. 2020. Cloning and characterization of two chlorophyll A/B binding protein genes and analysis of their gene family in *Camellia sinensis*. Scientific reports 10, 1–9.

Lim G, Burns KC. 2022. Adaptive fruit colour variation along an elevational gradient in a colour-polymorphic plant. doi: https://doi.org/10.21203/rs.3.rs-1709462/v1. [Preprint].

Liu W, Zheng L, Qi D. 2020. Variation in leaf traits at different altitudes reflects the adaptive strategy of plants to environmental changes. Ecology and Evolution 10, 8166–8175.

Ma L, Sun X, Kong X, Galvan JV, Li X, Yang S, Yang Y, Yang Y, Hu X. 2015. Physiological, biochemical and proteomics analysis reveals the adaptation strategies of the alpine plant *Potentilla saundersiana* at altitude gradient of the Northwestern Tibetan Plateau. Journal of proteomics 112, 63–82.

Malhi Y, Silman M, Salinas N, Bush M, Meir P, Saatchi S. 2010. Introduction: elevation gradients in the tropics: laboratories for ecosystem ecology and global change research. Global Change Biology 16, 3171–3175.

Martins SCV, Sanglard ML, Morais LE, Menezes-Silva PE, Mauri R, Avila RT, Vital CE, Cardoso AA, DaMatta FM. 2019. How do coffee trees deal with severe natural droughts? An analysis of hydraulic, diffusive and biochemical components at the leaf level. Trees 33, 1679–1693.

Mendonça AM das C, Lira JMS, Melo NC de, Rodrigues M, Barbosa JPRAD. 2019. Phenotypic Plasticity of *Pimenta pseudocaryophyllus* (Gomes) Landrum Under Different Light Conditions. Floresta e Ambiente 26.

Nanjo Y, Skultety L, Uváčková L, Klubicová K, Hajduch M, Komatsu S. 2012. Mass spectrometry-based analysis of proteomic changes in the root tips of flooded soybean seedlings. Journal of proteome research 11, 372–385.

Nunes MA, Bierhuizen JF, Ploegman C. 1968. Studies on productivity of coffee: I. Effect of light, temperature and CO_2 concentration on photosynthesis of *Coffea arabica*. Acta Botanica Neerlandica 17, 93–102.

Paschoa RP da, Christ JA, Valente CS, Ferreira MF da S, Miranda FD de, Garbin ML, Carrijo TT. 2018. Genetic diversity of populations of the dioecious *Myrsine coriacea* (Primulaceae) in the Atlantic Forest. Acta Botanica Brasilica 32, 376–385.

Passamani LZ, Barbosa RR, Reis RS, Heringer AS, Rangel PL, Santa-Catarina C, Grativol C, Veiga CFM, Souza-Filho GA, Silveira V. 2017. Salt stress induces changes in the proteomic profile of micropropagated sugarcane shoots. PLoS One 12, e0176076.

Pereira JP, Garbin ML, Carrijo TT, da Silva JA, Bourguignon TP, Cavatte PC. 2022. Lack of coordination between stomatal and vein traits provides functional benefits to the dioecious tropical tree Myrsine coriacea. Physiologia Plantarum, 174, e13719.

Pérez-Ramos IM, Matías L, Gómez-Aparicio L, Godoy Ó. 2019. Functional traits and phenotypic plasticity modulate species coexistence across contrasting climatic conditions. Nature communications 10, 1–11.

Perez-Riverol Y, Bai J, Bandla C, Hewapathirana S, García-Seisdedos D, Kamatchinathan S, Kundu D, Prakash A, Frericks-Zipper A, Eisenacher M, Walzer M, Wang S, Brazma A, Vizcaíno JA. 2022. The PRIDE database resources in 2022: A Hub for mass spectrometry-based proteomics evidences. Nucleic Acids Res 50, D543-D552.

Pipoly JJ. 1991. Systematic studies in the genus *Myrsine* L.(Myrsinaceae) in Guayana. Novon 1, 204–210.

R Core Team. 2023. R: A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing; https://www.R-project.org/.

Rahman IU, Afzal A, Iqbal Z, Hart R, Abd_Allah EF, Alqarawi AA, Alsubeie MS, Calixto ES, Ijaz F, Ali N. 2020. Response of plant physiological attributes to altitudinal gradient: Plant adaptation to temperature variation in the Himalayan region. Science of The Total Environment 706, 135714.

Rahman IU, Hart R, Afzal A, Iqbal Z, Abdallah EF, Alqarawi AA, Ijaz F, Ali N, Kausar R, Muzammil S. 2019. Phenological plasticity in *Berberis lycium* Royle along temporal and altitudinal gradients. Applied Ecology and Environmental Research 17, 331–341.

Reis RS, Vale EM, Sousa KR, Santa-Catarina C, Silveira V. 2021. Pretreatment free of 2,4-dichlorophenoxyacetic acid improves the differentiation of sugarcane somatic embryos by affecting the hormonal balance and the accumulation of reserves. Plant Cell, Tissue and Organ Culture (PCTOC) 145, 101–115.

Ricketson JONM, Pipoly III JJ. 1997. Nomenclatural notes and a synopsis of the genus *Myrsine* (Myrsinaceae) in Mesoamerica. SIDA, Contributions to Botany 17, 579–589.

Russell V. Lenth BBPBIG-VMHMJJLFMHRHS. 2023. Emmeans: estimated marginal means, aka least-squares means. R package version 1.

Sánchez-Tapia A, Garbin ML, Siqueira MF, Guidoni-Martins KG, Scarano FR, Carrijo TT. 2018. Environmental and geographical space partitioning between core and peripheral *Myrsine* species (Primulaceae) of the Brazilian Atlantic Forest. Botanical Journal of the Linnean Society 187, 633–652.

San-Eufrasio B, Castillejo MÁ, Labella-Ortega M, Ruiz-Gómez FJ, Navarro-Cerrillo RM, Tienda-Parrilla M, Jorrín-Novo J v, Rey M-D. 2021. Effect and response *of Quercus ilex* subsp. *ballota* [Desf.] Samp. seedlings from three contrasting Andalusian populations to individual and combined *Phytophthora cinnamomi* and drought stresses. Frontiers in Plant Science 12.

Umate P. 2010. Genome-wide analysis of the family of light-harvesting chlorophyll a/b-binding proteins in *Arabidopsis* and rice. Plant signaling & behavior 5, 1537–1542.

Vantyghem M, Merckx R, Stevens B, Hood-Nowotny R, Swennen R, Dercon G. 2022. The potential of stable carbon isotope ratios and leaf temperature as proxies for drought stress in banana under field conditions. Agricultural Water Management 260, 107247.

Vitoria AP, Vieira TO, Camargo PB, Santiago LS. 2016. Using leaf δ13C and photosynthetic parameters to understand acclimation to irradiance and leaf age effects during tropical forest regeneration. Forest Ecology and Management 379, 50–60.

Voelckel C, Gruenheit N, Lockhart P. 2017. Evolutionary transcriptomics and proteomics: insight into plant adaptation. Trends in plant science 22, 462–471.

Wang Y, Fan K, Wang J, Ding Z, Wang H, Bi C, Zhang Y, Sun H. 2017. Proteomic analysis of *Camellia sinensis* (L.) reveals a synergistic network in the response to drought stress and recovery. Journal of plant physiology 219, 91–99.

Warnes GR, Bolker B, Bonebakker L, *et al.* 2022. gplots: Various R Programming Tools for Plotting Data. https://github.com/talgalili/gplots.

Wiśniewski JR, Zougman A, Nagaraj N, Mann M. 2009. Universal sample preparation method for proteome analysis. Nature methods 6, 359–362.

Xia Y, Ning Z, Bai G, Li R, Yan G, Siddique KHM, Baum M, Guo P. 2012. Allelic variations of a light harvesting chlorophyll a/b-binding protein gene (Lhcb1) associated with agronomic traits in barley. PLoS One **7**, e37573.

Tables

Table 1. Location of the natural populations of *M. coriacea*.

Population code	Municipality	Locality	Latitude	Longitude	Altitude (M.S.L)
POP1	Dores do Rio Preto	Macieira in Caparaó National Park	20º27'27''S	41°48'38"W	2019
POP2	Domingos Martins	Parque Estadual Pedra Azul	20°23'38"S	41°01'36"W	1229
POP3	Mimoso do Sul	Pedra dos Pontões	20°59'35"S	41°21'08"W	930
POP4	Muqui	Private farm property	20º53'40S	41°20'12"W	657

Figures legends

Fig. 1. Proteomic analysis relationships between natural populations of *M. coriacea* based on the protein identification and quantitation results. A) Heatmap using the Unweighted Pair Group Method using Arithmetic averages (UPGMA); B) PCA.

Fig. 2. Protein-protein interaction (PPIs) network among the DAPs in the comparative proteomic analysis of POP1/POP2. The red nodes represent proteins involved in "carbon fixation in photosynthetic organisms," the blue nodes are related to "glyoxylate and dicarboxylate metabolism," the light green nodes are involved in "glycolysis/gluconeogenesis," the maroon nodes are related to "pyruvate metabolism," the dark green nodes represent proteins involved in "cysteine and methionine metabolism," the pink nodes are involved in "amino sugar and nucleotide sugar metabolism," and the light blue nodes are involved in "photosynthesis." The colored lines indicate various types of interaction evidence (dark blue: co-occurrence; black: co-expression; light blue: database evidence; green: neighboring genes; pink: experimental evidence; red: fusion evidence; yellow: text mining evidence; purple: protein homology). Arrows indicate the proteins highlighted for discussion.

Fig. 3. Protein-protein interaction (PPIs) network among the DAPs in the comparative proteomic analysis of POP1/POP3. The red nodes represent proteins involved in "carbon fixation in photosynthetic organisms," the blue nodes are related to "glyoxylate and dicarboxylate metabolism," the light green nodes are involved in "glycolysis/gluconeogenesis," the maroon nodes are related to "pyruvate metabolism," the dark green nodes represent proteins involved in "cysteine and methionine metabolism," the pink nodes are involved in "amino sugar and nucleotide sugar metabolism," and the light blue nodes are involved in "photosynthesis." The colored lines indicate various types of interaction evidence (dark blue: co-occurrence; black: co-expression; light blue: database evidence; green: neighboring genes; pink: experimental evidence; red: fusion evidence; yellow: text mining evidence; purple: protein homology). Arrows indicate the proteins highlighted for discussion.

Fig. 4. Protein-protein interaction (PPIs) network among the DAPs in the comparative proteomic analysis of POP1/POP4. The red nodes represent proteins involved in "carbon fixation in photosynthetic organisms," the blue nodes are related to "glyoxylate and dicarboxylate metabolism," the light green nodes are involved in "glycolysis/gluconeogenesis," the dark green nodes represent proteins involved in "cysteine and methionine metabolism," the pink nodes are involved in "amino sugar and nucleotide sugar metabolism," and the light blue

nodes are involved in "photosynthesis." The colored lines indicate various types of interaction evidence (dark blue: co-occurrence; black: co-expression; light blue: database evidence; green: neighboring genes; pink: experimental evidence; red: fusion evidence; yellow: text mining evidence; purple: protein homology). Arrows indicate the proteins highlighted for discussion.

Fig. 5. Physiological parameters among the natural populations of *M. Coriacea*. A) Leaf carbon isotopic composition (*δ13C*); B) leaf carbon (*C*); C) leaf nitrogen (*N*); and D) relative growth rate in height (*RGR_H*). Data are presented as the mean ± CI (95% confidence interval) of six independent replicates The error bars represent confidence intervals.

Fig. 6. Physiological parameters among the natural populations of *M. Coriacea*. A) Leaf hydraulic conductivity (*Kleaf*); B) net assimilation rate of CO_2 (*A*); and C) specific leaf area (*SLA*). Data are presented as the mean ± CI (95% confidence interval) of six independent replicates. The error bars represent confidence intervals.