Androgen deprivation triggers a cytokine signaling switch to induce immune suppression and prostate cancer recurrence

Kai Sha[1,6,8], Renyuan Zhang[1,6,9], Aerken Maolake[1], Shalini Singh[1,10],

Gurkamal Chatta[4], Kevin H. Eng[1,2,11], Kent L. Nastiuk[1,3,5,7] and John J. Krolewski[1,5,7]

Departments of [1]Cancer Genetics and Genomics, [2]Biostatistics and Bioinformatics, [3]Urology and [4]Medicine, Roswell Park Comprehensive Cancer Center, Buffalo, NY 14263

[5]Department of Biology and Program in Data Science and Analytics, Buffalo State University/ SUNY, Buffalo, NY 14222

[6,7]These authors contributed equally to this report

[8]Current address: LabCorp-Monogram Biosciences, South San Francisco, CA 95080
[9]Current address: Thermo-Fisher Scientific, Ann Arbor, MI 48104
[10]Current address: BiologyWorks, Los Angeles, CA 90004
[11]Current address: Eikon Therapeutics, Jersey City, NJ 07302

CORRESPONDENCE:

Dr. J.J. Krolewski, Buffalo State University/ SUNY, Department of Biology; 1300 Elmwood Avenue; Buffalo, NY 14222; krolewjj@buffalostate.edu; 949-872-9107

Dr. K.L. Nastiuk, Roswell Park Comprehensive Cancer Center, Department of Cancer Genetics and Genomics, Elm & High Streets, Buffalo, NY 14263; kent.nastiuk@roswellpark.org; 716-845-5771

KEY WORDS: androgen deprivation, tumor stem cell, CCL2, immunosuppression, resistance, TNF
CONFLICT OF INTEREST: The authors declare that there are no potential conflicts of interest
FINANCIAL SUPPORT: This work was supported by the Department of Defense Prostate Cancer Research Program No. W81XWH-19-1-0378 (to J.J. Krolewski and K.L. Nastiuk), institutional funds from the Roswell Park Alliance Foundation and the National Cancer Institute Cancer Center Support Grant P30-CA016056 to Roswell Park Comprehensive Cancer Center.

ABSTRACT

Androgen deprivation therapy (ADT) is an effective but not curative treatment for advanced and recurrent prostate cancer (PC). We investigated the mechanisms controlling the response to androgen-deprivation by surgical castration in genetically-engineered mouse models (GEMMs) of PC, using high frequency ultrasound imaging to rigorously measure tumor volume. Castration initially causes almost all tumors to shrink in volume, but many tumors subsequently recur within 5-10 weeks. Blockade of tumor necrosis factor (TNF) signaling a few days in advance of castration surgery, using a TNFR2 ligand trap, prevents regression in a PTEN-deficient GEMM. Following tumor regression, a stem cell-like population within the tumor increases along with TNF protein levels. Tumor cell lines in culture recapitulate these *in vivo* observations, suggesting the possibility that stem cells are the source of TNF. When TNF signaling blockade is administered immediately prior to castration, tumors regress but recurrence is prevented. This implies that a late wave of TNF secretion within the tumor – which coincides with the expression of NFκB regulated genes – drives recurrence. The inhibition of signaling downstream of an NFκB-regulated protein – chemokine C-C motif ligand 2 (CCL2) – prevents post-castration tumor recurrence, phenocopying post-castration (late) TNF signaling blockade. CCL2 was originally identified as a macrophage chemoattractant and indeed at late times after castration gene sets related to chemotaxis and migration are up-regulated. Importantly, enhanced CCL2 signaling during the tumor recurrence phase coincides with an increase in pro-tumorigenic macrophages and a decrease in CD8 T cells, suggesting that recurrence is driven at least in part by tumor immunosuppression. In summary, we demonstrate that a therapy-induced switch in TNF signaling induces an immunosuppressive tumor microenvironment and concomitant tumor recurrence.

INTRODUCTION

Androgen deprivation therapy (ADT) via surgical castration (1) and currently via medical therapy employing gonadotropin releasing hormone analogues (2), has been an effective treatment for advanced primary and recurrent prostate cancer (PC) for many decades. Within the past several years, the addition of either the microtubule toxin docetaxel or the androgen biosynthesis inhibitor abiraterone to ADT has improved the medical therapy (3). However, these ADT-combination therapies have only marginally extended overall survival and the fundamental issue for patients with advanced primary and recurrent PC remains the same: ADT (or an ADT-combination) rarely cures PC and resistant tumors are eventually lethal. Thus, there is a need to develop additional therapies for PC, preferably by identifying molecular targets that are mechanistically linked to the development of ADT resistance. ADT is effective because the growth of almost every prostate cancer is dependent on a program of gene expression mediated by the androgen receptor and a set of critical co-regulators. The androgen receptor (AR) gene is activated by genetic alteration in only a small fraction of primary PCs (4). Instead, one or two alterations of a small group of other genes (5) is sufficient to re-program the AR cistrome to transcribe a set of genes that promote immortalization and tumorigenesis (6). Additional alterations in a second class of genes driving proliferation and growth – such as the PTEN tumor suppressor or the c-MYC oncogene – completes the process of transformation to prostate cancer (5).

Tissue reconstitution studies (7) have shown that androgen blockade can act indirectly via soluble factors derived from the tumor micro-environment (TME). Consistent with this paradigm, in which proteins are secreted into the TME and act on the cell of origin (autocrine) or on other cells within the TME (paracrine), we have previously investigated the response of prostate cancers to androgen deprivation (8-13), using cell culture and murine models of human prostate cancer. A recurrent theme of this research has been the role of tumor necrosis factor (TNF) as a pleotropic signaling mediator within the PC TME. TNF binds and trimerizes its cognate receptor and signals via a complex network of intercellular protein-protein interactions to promote either an anti-tumor or pro-tumor fate, depending on the cellular context. In this report, we examine the role of TNF and TNF-regulated genes in the response

to castration in a genetically engineered mouse model (GEMM) of PC driven by deletion of the PTEN

gene (14). Castration induces regression and subsequent recurrence in this GEMM, resembling the

onset of castration resistant prostate cancer (CRPC). Both responses are dependent on TNF signaling

and can be separately blocked by differentially timing the administration of a TNF ligand trap molecule.

At late times post-castration, when the tumor is beginning to recur in a castration resistant form, TNF

induces NFκB-mediated gene expression. Moreover, we demonstrate that the NFκB-induced gene

CCL2 – which encodes a protein that is chemotactic for immune suppressive cells such as tumor

associated macrophages (TAMs) – is necessary for recurrence in the PTEN-loss model as well as in a

second GEMM of PC that is driven by MYC over-expression. Thus, a TNF signaling switch induces

immunosuppression in the tumor microenvironment to promote castration-resistant tumor recurrence.

MATERIALS AND METHODS

Animals and tumor imaging

All animal studies were approved by the Roswell Park Comprehensive Cancer Center (RPCCC)

institutional animal care and use committee. Male Pb-Cre4; PTEN$^{fl/fl}$ mice (14) and Hi-MYC (15) were

breed in the RPCCC laboratory animal shared resource. Mice were housed in environmentally-controlled

conditions on a 12-hour light/dark cycle with water and food. For both genetically engineered mouse

models (GEMMs) we studied, male pups were genotyped and tumor volumes were monitored by three-

dimensional high frequency ultrasound (HFUS) imaging, as previously described (9,16). In brief, mice

were anesthetized, the abdomen was depilated and B-mode image scans were acquired using a Vevo

2100 micro-ultrasound imaging system (Vevo LAZR; VisualSonics Inc., Toronto). Images were imported

into Amira software (Visualization Sciences Group, Burlington, MA) for 3D tumor reconstruction and

volume determination. Tumors developed in ~100% of male animals for both GEMMs. The histology,

lobular location and usual age to reach 300-500 mm^3 volume are provided for the two models in

Supplementary Table S1. Mice with tumor volumes of 300-500 mm^3 were randomly enrolled into

experimental groups, and surgically castrated (or sham-operated) via scrotal incision, under isoflurane

anesthesia (17). To systematically define the response of the GEMMs to castration, we applied a murine

approximation of the human clinical RECIST (response evaluation criteria in solid tumors) criteria (18).

Thus, we define regression as a volume decline of 16% or more post-castration and recurrence (of tumors

that had met the regression standard) as occurring when we observe volume increases of 11% from the

volume nadir, in at least two temporally distinct imaging sessions. Etanercept (sTNFR2-Fc; Amgen,

Thousand Oaks, CA, USA) was injected intra-peritoneally (4mg/kg) twice per week for up to 5 weeks

following castration, based on our previous studies (8,11). The CCR2 antagonist (CCR2a) BMS-741672

(Torcis Bioscience, Bristol, UK) was injected intraperitoneally (2 mg/kg) three times per week as

described (19), beginning one day prior to castration.

Tumor tissue collection and processing

After mice were sacrificed, individual prostate tumors were identified grossly, excised and cut in half.

Typically, a portion was flash frozen in liquid nitrogen and stored in a liquid nitrogen freezer while a

second portion was fixed in paraformaldehyde at room temperature overnight, washed in PBS, stored at

4°C and embedded in paraffin as required for the preparation of tissue sections. Mouse prostate tumor

and normal tissues were dissociated for single cell RNA sequencing using the Tumor Dissociation kits

from Miltenyi Biotec (cat. no. 130-096-730). Briefly, tissue was excised, washed with chilled 1x PBS,

placed in a petri dish on ice, cut into 2-4 mm^3 pieces, transferred to a gentleMACS C tube (cat. no. 130-

096-334) containing DMEM/enzyme mix and then attached to gentleMACS Octo Dissociator (cat. no.

130-096-427) with heating running the '37C_m_TDK_1' (mouse) program. The suspension was filtered

through a 40 µm cell strainer (Falcon). The cells were harvested by centrifugation at 400g for 10 min and

resuspended in chilled 1X Red Blood Cell Removal Solution (Miltenyi Biotec, cat. no.130-094-183).

Debris was then removed using the Debris Removal Solution (Miltenyi Biotec, cat. no. 130-109-398)

following the manufacturer's instructions.

Cell culture

The C4-2 cell line, from Jer-Tsong Hsieh, UT Southwestern Medical Center, Dallas, TX, was used to

model castration-resistant prostate cancer (20). C4-2 was periodically monitored by polymerase chain

reaction (PCR) for mycoplasma contamination (21) and DNA fingerprinting was employed to authenticate

this cell lines (22). C4-2 was cultured in RPMI-1640 media supplemented with 10% FCS and 1%

penicillin/streptomycin. Enzalutamide was from Selleck Chemicals (Houston, TX).

Enzyme-linked immunosorbent assays (ELISA)

In the case of prostate tumors, tissue powder was homogenized in NP-40 buffer using Sonic

Dismembrator (Fisher Scientific, Hampton, NH) at power 7 for 30 sec. The supernatant was collected by

centrifugation and stored at −80°C for subsequent analysis. In the case of cell culture, cells were seeded

at a density of $0.8\text{-}1\times10^5$/ml as mono-cultures and media stored at 4°C was used for subsequent analysis.

TNF and CCL2 were quantitated by ELISA (eBioscience, San Diego, CA), according to the

manufacturer's instructions. Protein concentration was determined using bicinchoninic acid reagent

(Pierce, ThermoFisher Scientific), according to the manufacturer's instructions.

Fluorescent activating cell sorting

Cells in culture were detached from cell culture plasticware using StemPro accutase (Gibco,

Gaithersburg, MD), washed with PBS and incubated with 1% BSA in 1x PBS for 10 min at 4°C. Human

cell lines were incubated with antibodies against CD166-FITC (3A6; Ancell, Stillwater, MN; 1:1250) and

CD49f-Alexa Fluor® 647 (GoH3; Biolegend, San Diego, CA; 1:200) for 30 min at 4°C. Mouse prostates

were disaggregated using 1mg/ml type I collagenase and 1mg/ml Dispase (Invitrogen, Carlsbad, CA) in

RPMI medium and filtered through 40 μm cell strainer. Cells were suspended in DMEM with 10% FBS

and 1% L-glutamine, and then incubated with fluor-labeled antibodies for 30 min at 4°C. Mouse cells

were incubated with antibodies against Sca-1-APC (D7; 1:500), CD49f-PE (GoH3; 1:333), and the

'lineage' markers (Ter119-FITC (TER-119; 1:250), CD31-FITC (390; 1:250) and CD45-FITC (30-F11;

1:250)). LSC FACS, first gating on lineage-negative (Lin$^-$) cells and then CD49f and SCA-1 was

performed as described by Witte and colleagues (23). All anti-mouse antibodies were purchased from

eBioscience (San Diego, CA). 7-AAD (Invitrogen, Carlsbad, CA; 1:1000) was used to determine dead

cells. FACS analysis was performed using the BD LSR-II (BD Bioscience, San Jose, CA) and analyzed

by Winlist 3D (Version 8.0; Verity Software House, Topsham, ME, USA). Cell sorting was done by BD

FACSAria-II (BD Bioscience, San Jose, CA).

Colony formation assay

Cells were seeded in 6-well plates at densities of 0.5×10^3, 1×10^3 and 1.5×10^3 cells per well and

cultured for 21 days. The media was removed, wells were washed once with PBS, stained with 2%

toluidine blue for 30 min and scanned with Epson Perfection 1240U (Suwa, Nagano Prefecture, Japan).

Colonies were imaged and then enumerated using ImageJ.

Immunohistochemistry

Immunohistochemical (IHC) staining was performed on 5 μm sections of formalin fixed paraffin-

embedded (FFPE) mouse prostate tumor tissue. Sections were incubated with monoclonal antibodies

against F4/80 (BM8; Biolegend, San Diego, CA, USA; 1:600), CD8alpha (D4W2Z; Cell Signaling

Technology, Danvers, MA, USA; 1:400), or CCL2 (2H5; Biolegend, San Diego, CA, USA; 1:100),

appropriate horse-radish peroxide secondary antibodies and reacted with 3,3-diaminobenzidine using an

automated tissue processor. IHC and H&E images were captured at 20x magnification using Olympus

BX45 microscope and cellSens software (Olympus, Shinjuku-ku, Tokyo, Japan). Ten random high

powered (200X) fields were captured for each section and were blinded prior to quantitation of stained

cells (F4/80$^+$ or CCL2$^+$ macrophages or CD8a$^+$ lymphocytes) using ImageJ.

RNA extraction, sequencing and analysis

RNA was extracted from liquid nitrogen frozen tumor samples by pulverizing tissue into a powder using

a pre-cooled mortar and pestle. TRIzol (Invitrogen, Carlsbad, CA) was then added by a ratio of 1 mL of

TRIzol per 100 mg of tumor tissue, and RNA extraction completed as per manufacturer's instructions

(24). Bulk RNA-seq data processing and analysis was performed as follows. FastQC v0.11.5 was used

for quality control; STAR v2.6.0a (25) was employed to align the reads to the murine reference genome

(GRCm38 M16) and gene expression was then quantified using RSEM (26). FastQC, RSEM, STAR

were all performed in a local Linux/Unix computing environment. Subsequent analyses were performed

using the following R-software packages from the Bioconductor website. Estimated gene counts from

RSEM were filtered and the upper quartile was normalized using edgeR (27). Differential gene

expression (DGE) was performed using Limma after Voom transformation followed by linear regression

(28,29). The significance threshold was set to false discovery rate (FDR) / adjusted p-value of less than

0.05 with a fold change greater than 2 (FDR < 0.05 and |log2 fold change| > 1). Sets of genes with

FDR<0.05 and |log2FC| from the DGE analysis were subjected to Gene Ontology analysis of biological

processes as described (30), using the website tool (http://pantherdb.org/index.jsp). Heatmaps were

generated using the package Complexheatmap (31). Single cell gene expression libraries were

generated using the 10X Genomics Chromium platform and sequenced on a NovaSeq 6000 (Illumina).

Processing of scRNA-seq data was performed as follows. For QC, normalization, clustering and

differential gene expression, the Cellranger v6.0.0 software (https://support.10xgenomics.com/single-

cell-geneexpression/software/) was used to align FASTQ sequencing reads to the mm10 (mouse)

reference transcriptome, generating single cell feature counts for each sample. Count data from all cells

were included in a 'SingleCellExperiment' (version 1.8.0) object (32). The 'scater' package (version

1.14.5) was used to filter cells and normalize counts per million (CPM) (33). Transcripts from a given cell

were removed from the analysis if the corresponding transcriptome met any of the following criteria: 20%

or more reads aligned to mitochondrial genes, 1000 total counts per cell or less, or 300 genes per cell or

less. The dimensionality of this transcriptomic data was reduced using the t-distributed stochastic

neighbor embedding (t-SNE) algorithm (34,35). The 'limma' package (29) was used to identify

differentially expressed genes to refine the assignment of clusters generated by the t-SNE algorithm.

Statistics

Data are presented as the mean ± SEM. Statistical analyses were conducted using JMP-Pro12 (SAS,

Cary, NC, USA). Differences between two means was assessed by Student's unpaired t-test and

differences among multiple means by one-way or two-way ANOVA followed by Tukey-Kramer HSD post-

hoc testing if the F-statistic was significant. For the small n, categorical data sets related to the studies

using GEM prostate cancer models shown in Figures 4A, 5C and 5D, we employed the exact binomial

test to determine the Clopper-Pearson confidence interval for the proportion and Fisher's exact test to

determine the *p*-values.   Supplementary Table S3 summarizes these statistical analyses.

RESULTS

Tumor recurrence following castration-induced regression in PTEN-deficient PC

Mice engineered to delete the PTEN gene in the prostate epithelium at puberty uniformly develop an

intraductal form of prostate adenocarcinoma (14) (Table S1).   To investigate the response of prostate

cancer to androgen deprivation therapies, we used high frequency ultrasound (HFUS) and three-

dimension reconstruction software (9,16) to serially determine the volume of tumors in live animals (Fig.

1A).   As seen in Figure 1B, the kinetics of tumor volume following castration is complex.   Immediately

following castration, tumor volume continues to increase similar to the non-castrated control.   We recently

demonstrated that this delay in tumor regression – which is not observed in the normal murine prostate

– is driven by glucocorticoid receptor signaling transiently substituting for tumor-specific re-programmed

AR signaling (9).   Following this delay, the tumors regressed as expected but within ~5 weeks most

began to increase in volume, implying tumor recurrence and mimicking the development of CRPC in

men.   Regression and recurrence events were defined by adapting standard human oncology imaging

protocols to mice, as described in the MATERIALS AND METHODS section and in the legend to

Supplementary Table S2.   Serial HFUS imaging of a large cohort castrated PTEN-deficient mice

(including those in Fig. 1) is summarized in Supplementary Table S2, demonstrating 40-80% recurrence

dependent on the period of post-castration observation.

We measured TNF protein levels in tumor lysates, and found an increase at 5 weeks post-castration,

when tumor recurrence is beginning (Fig. 1C).   Previously, we had observed an increase in *Tnf* mRNA

at 8 hours post-castration in the stromal compartment of the normal prostate, suggesting a relatively

direct effect of androgen-withdrawal on TNF expression (11).   In the case of PTEN-deficient tumors, the

increase in TNF protein at a late time post-castration suggests a more complex transcriptional and/or

translational regulatory mechanism in response to androgen withdrawal.   Alternately, the increase in TNF

at late times post-castration might be due to the expansion of a population of TNF-expressing cells as

the tumor evolves in response to androgen deprivation. We queried gene expression profiles in public

databases from two previously published studies of men who had either hormone-naïve PC or CRPC

(36,37), to gain insight into this question. Increased *TNF* mRNA expression in CRPC, relative to

castration-naive primary PC, correlated with elevated expression of stem cell-related genes in both

datasets (Fig. S1).

240 Androgen-deprivation induced TNF secretion is due to increased basal cell stemness

To determine if we could detect a change in the stem cell-like tumor epithelial cell populations as the

tumors began to increase in volume following castration-induced regression, we examined single cell

preparations from tumors for cell surface expression of stem cell markers, using fluorescent-activated

cell sorting (FACS). We observed an increase in Lin$^-$/Sca1$^+$/CD49f^{hi} (LSC^{hi}) cells at 25 days post-

castration, relative to non-castrated tumors (Fig 1D). These cells correspond to the cytokeratin-5$^+$/p63$^+$

basal stem cell population (38), a multipotent stem cell population that is required for the post-natal

development of the prostate (39). Basal stem cells are distinct from Lin$^-$/Sca1$^+$/CD49f^{med} (LSC^{med}) luminal

progenitor cells (identified in the lower right population in Fig. 1D) that are the likely tumor initiating cell

populations for most human prostate cancers (39,40).

The data in Figures 1 and S1 suggest that the TNF secreted during the post-castration recurrence

phase in the PTEN-deficient tumors might be derived from a cell population with increased levels of

'stemness'. To test this more thoroughly, we performed experiments using the C4-2 and LNCaP cell

culture models of PC. First, we observed that long-term culture in the anti-androgen enzalutamide (41)

significantly reduced the growth rate of C4-2 prostate cancer cells (Fig. 2A), as expected. Reduced

growth was accompanied by an increase in the fraction of cells that resembled basal stem cells (Fig. 2B-

C) and a coordinate increased in TNF levels (Fig. 2C). In the analyses for Figure 2, we employed CD166

as a basal stemness biomarker (42). We also performed the enzalutamide selection experiment on the

LNCaP cell line, an isogenic parental line for C4-2 (20), which is more androgen-responsive (Fig S2A).

We observed a similar correlation between stemness and TNF secretion (Fig. S2B-C). Finally, we

replicated enzalutamide selection for both C4-2 and LNCaP cell lines, using immunofluorescence

microscopy to identify CD49f^{hi} and CD166^{hi} cells (Fig. S3). Again, an increase in the fraction of cells

scoring for these stemness markers correlated with increased TNF secretion. Stem cell marker

enrichment >10% was typically required to detect an increase in TNF in the media of enzalutamide-

treated cultures. Next, we treated C4-2 cells with enzalutamide for 20 days and used preparative FACS

to partially purified cells into CD166^{lo} and CD166^{hi} fractions (Fig. 3A). TNF levels were higher in

conditioned media from the CD166^{hi} fraction relative to the CD166^{lo} fraction (Fig. 3B). We also

demonstrated that the efficiency of colony formation – a functional marker of stemness – correlated with

the level of expression of CD166 (Fig. 3C).

Differentially-timed blockade of TNF signaling

We previously employed cell culture and mouse models of prostate cancer to demonstrate that TNF

signaling mediates two distinct responses to ADT – an anti-tumorigenic apoptotic/regression phenotype

(8,43) and a pro-tumorigenic cell migration phenotype (44). Given the observations in Figure 1, we

hypothesized that TNF mediates both the regression and recurrence phases that occur after castration

(Fig. 1B). To test this, we treated tumor-bearing mice with etanercept – a sTNFR2-Fc fusion protein,

which competitively binds TNF (45) and prevents receptor binding – beginning at three distinct time

points: three days prior, one day prior, and three days following castration. When etanercept is

administered three days before castration (Fig. 4A, -3d), castration-induced regression is blocked, and

tumor growth is similar to the sham group (Fig. 1B). Etanercept given one day before castration does

not affect regression but blocks recurrence (Fig. 4A, -1d). Finally, when etanercept was administered

three days after castration there is no detectable effect on regression or recurrence (Fig. 4A, +3d).

Instead, the result resembles castration alone (Fig. 1B).

To gain insight into the nature of the different molecular processes occurring during regression

and recurrence, we performed gene set enrichment analysis (GSEA) on RNA sequence data from bulk

tumor mRNA. Specifically, we compared samples from sham castrated mice with those from either mice

six days after castration (Fig. 4B, upper panel) or mice 35 days after castration (Fig. 4B, lower panel).

Multiple gene sets related to TNF-induced transcription, most NFkB-related, were uniformly down-

regulated at six days post-castration but up-regulated 5 weeks after castration, relative to the sham

conditions. The switch in TNF-related gene sets from down-regulation to up-regulation is consistent with

a switch from apoptosis signaling (low NFkB levels) to signaling that promotes transcription of NFkB-

regulated genes (high NFkB levels).

CCL2 signaling is required for recurrence in murine prostate cancer models

CCL2 is a chemotactic cytokine (chemokine) (46) linked to prostate cancer pathogenesis (47). The

CCL2-CCR2 signaling axis has multiple pro-tumorigenic functions, most notably the recruitment and

polarization of tumor associated macrophages (TAMs) (48). Previously we demonstrated autocrine

CCL2 secretion following enzalutamide induction of TNF in mono-cultures of C4-2 cells (10), consistent

with CCL2 gene transcription via TNF induction of NFκB (49). We returned to the enzalutamide selection

experiments shown in Figure 2C and Supplementary Figure S2 and found that the secretion of CCL2

protein increased over time, in a manner dependent on secreted TNF, in enzalutamide treated C4-2 and

LNCaP cells (Fig. S4A-B)To assess paracrine CCL2 production in response to androgen deprivation, we

replicated these experiments using co-cultures of LNCaP or C4-2 plus the myofibroblast-like WPMY-1

cell line (50), grown in enzalutamide-containing media. Again, we observed coordinate secretion of TNF

and CCL2 (Fig. S4C). Co-cultures produced 5- to 20-fold more CCL2 than the prostate cancer cell line

mono-cultures. We also found that enzalutamide-treated co-cultures of C4-2 plus WPMY-1 produced

conditioned media that induced migration of THP-1, a CCR2$^+$ macrophage-like leukemia cell line that is

a model for TAMs (Fig. S4D).

These observations suggested that paracrine TNF induction of CCL2 in the prostate cancer TME

might be responsible for tumor recurrence at late times post-castration, possibly by recruiting

immunosuppressive TAMs (51). Indeed, we found that in the samples from Figure 1C, CCL2 increased

at 5 weeks (Fig. 5A), coincident with the rise in TNF that accompanies tumor recurrence. Tissue sections

corresponding to tumors 5 weeks post-castration showed an increase in stroma-localized macrophage-

like cells expressing CCL2 (Fig. 5B). Moreover, CCL2 expression was reduced in tumor tissue sections

from castrated mice that received etanercept, consistent with CCL2 regulation by TNF (Fig. 5B). To

determine if CCL2 signaling was necessary for post-castration tumor recurrence in prostate cancer

mouse models, mice bearing PTEN-deficient prostate cancers received an antagonist of CCR2 prior to

castration. Serial imaging revealed that none of the tumors met the criteria for recurrence (Fig. 5C),

indicating that CCL2-CCR2 signaling is indeed required for castration resistant recurrence in this model.

Importantly, the tumor kinetics produced by CCR2 blockade phenocopies the effect of etanercept

administered one-day prior to castration (Fig. 4A, -1d), consistent with TNF regulation of CCL2 gene

transcription via NFkB. To confirm the role of CCL2 signaling we replicated the CCR2 inhibition studies

in the Hi-MYC prostate cancer GEMM, which has a similar response to castration (regression delay,

regression then recurrence) as the PTEN-deficient GEMM (Fig. 5D, blue symbols; see also ref. (9)). We

observed that CCR2a treatment prior to castration was comparably effective at preventing post-castration

recurrence in Hi-MYC mice (Fig. 5D, red symbols; also see Fig. S5).

We used a transcriptomic approach to further investigate CCL2 signaling, immune cell migration,

and TAMs in the development of castration-resistant prostate cancer in the PTEN-deficient GEMM.

Differential gene expression (DGE) analysis was performed on bulk RNA-sequencing data from tumor

tissue and genes with FDR<0.05 and |log2FC| (see Fig. S6 and details in the legend) were subjected to

Gene Ontology (GO) enrichment analysis (Fig. 6). These analyses revealed that a very similar category

of gene sets – mediating immune cell chemotaxis and migration – are up-regulated in the bulk tumor

mRNA from mice 35 days post-castration (Fig. 6A) but are down-regulated in the bulk tumor mRNA from

mice 35 days post-castration which were treated with CCR2a (Fig. 6B). We also assembled a TAM-

related gene set and found that these genes are similarly regulated: 'up' in the 35 days post-castration

group and 'down' in the 35 days post-castrated group treated with CCR2a (Fig. 6C).

An immunosuppressive cell profile is induced in the TME during recurrence

Taken together, the analyses in Figures 5 and 6 indicate that CCL2 might be functioning to recruit TAMs

and drive the progression to, or recurrence as, castration resistant prostate cancer (52). Thus, to

determine how TNF-CCL2 signaling effects the tumor immune microenvironment in the PTEN-deficient

murine prostate cancer model, we used immunohistochemistry (IHC) to quantitate F4/80- and CD8-

stained cells in tissue sections from the tumors. As seen in Figure 7A, F4/80-stained cells (presumptive

TAMs) increase, most notably at 35 days post-castration, just as tumor re-growth in the recurrence phase

is beginning and CCL2 levels are peaking (Fig. 5A). In addition, etanercept partially blocks this increase

in TAMs and coordinately inhibits tumor recurrence. CD8 T cells (presumptive cytolytic T-lymphocytes;

CTLs) decreased over time following castration and this was also partially reversed by etanercept

administration to the mice bearing PTEN-deficient tumors. Figure 7B demonstrates that blocking CCL2

signaling with the CCR2 antagonist has the expected effect – it reduces F4/80-stained cells and

increased CD8-stained cells, reversing the castration-induced immunosuppressive cell profile and

coordinately preventing prostate cancer recurrence (Fig. 5C). Integrating the data on F4/80-staining,

CD8-staining and CCL2 ELISAs for tumors treated and harvested at various times post-castration reveals

a demonstrates a high degree of correlation of the key immune components we examined. Specifically,

there is positive correlation of macrophages and CCL2 secretion and a negative correlation of T cells

with macrophage as well as with CCL2 section (Fig. S7). We also observed a more pronounced CD45+

cell infiltrate in the recurrent tumors, relative to non-recurrent tumors (Fig. S7B).

Transcriptomic CD8/CD68 ratio predicts castration outcome

Next, we carried out transcriptomic profiling to confirm the IHC studies in Figure 7, which correlated an

immunosuppressive immune cell profile (elevated CD68, reduced CD8) with recurrence of the PTEN-

deficient tumor. Specifically, three mice bearing PTEN-deficient prostate cancers were castrated and

tumor volume was HFUS-monitored for 42 days (Fig. 8A). One mouse failed to regress and was labeled

'stable'. Two mice regressed, one of which showed recurrence 21 days after castration. We recovered

the tumors, and performed single cell RNA sequencing. Transcript count data was pooled and tSNE

dimensionality reduction was used, along with subjective cell annotation, to identify six cell populations

(Fig. 8B). Among CD45-positive immune cells, TAMs were identified as those cells expressing both

CD68 and CD80, while cytolytic T cells (CTL) were defined as those expressing CD3 and CD8 (see

details in the legend to Fig. 8). In Figure 8C, we plotted the ratio of CTL to TAM, and observed that

compared with the regressing tumor, the recurrent tumor had a lower CTL/TAM ratio. This is consistent

with the IHC analysis in Figure 7A and suggests that immune suppression drives tumor re-growth.

DISCUSSION

TNF signaling switch

Employing a HFUS-based imaging protocol to measure tumor volume in live mice, we are able to resolve

the response of genetically-engineered murine prostate cancers to castration into multiple,

mechanistically-distinct phases, as illustrated for the PTEN-deficient prostate cancer model in Figure 1

(the kinetics for the castration response in the Hi-MYC GEMM are similar).   Immediately following

castration, there is an increase in tumor volume in both prostate cancer GEMMs.   We recently

demonstrated that this seemingly paradoxical effect is dependent on glucocorticoid signaling transiently

substituting for tumor-specific androgen receptor signals that are blocked by the androgen ablative effects

of castration (9).   After 4-7 days of gradually increasing volume, the tumors begin to regress due to

apoptotic cell death of the epithelial tumor cells in the cancerous glands, consistent with the classical

morphological description of apoptosis (53).   In addition to tumor epithelial cell apoptosis, there is a loss

of vascular integrity within the tumor that contributes to the regression phase (54,55).   Some tumors

continue to regress for more than two months, but for the majority – up to 75% of individual tumors in the

prostate cancer GEMMs we studied – recurrence begins 4-6 weeks after castration (Fig. 1A, Fig 4D and

Supplementary Table S2b).

We sought to decipher the intra- and inter-cellular signaling events that occur within the tumor

and the surrounding microenvironment in response to androgen deprivation (castration).   Previously we

had shown that TNF – but not other common death receptor ligands (FasL and TRAIL) – is necessary

for apoptosis-driven regression of the normal murine prostate (11).   In addition, we also found that TNF

is required for so-called vascular regression of murine prostate cancers, probably by triggering

endothelial cell apoptosis (8). The loss of vascular integrity leads to tumor hypoxia within a day following

castration.   While these events occur multiple days prior to the onset of a measurable reduction in

prostate cancer volume, hypoxia may indirectly drive regression by inducing tumor epithelial cell

apoptosis via TP53 (8). In the current studies, when we administered etanercept to PTEN-deficient mice

three days prior to castration, regression was effectively inhibited (Fig. 4A and Supplementary Table

S2a). This observation is consistent with TNF-driven apoptosis of tumor epithelial and endothelial cells.

The increase in TNF protein 5 weeks post-castration (Fig. 1C) – at a point when most tumors

have regressed – suggested a second role for TNF signaling in the response to androgen deprivation.

Indeed, blockade of TNF signaling beginning one day prior to castration results in most tumors

regressing, but no recurrence (Fig. 4A and Supplementary Table S2a). Thus, TNF signaling switches

from pro-apoptotic signaling that activates caspase-8 and drives regression to pro-tumorigenic signaling

that activates the NFκB and/or PI3K/Akt pathways, promoting tumor cell survival and proliferation and

tumor recurrence. The results of transcriptional analysis (Fig. 4B) support the switch to a pro-tumorigenic

phenotype: we observed increased expression of NFκB-related gene sets in bulk mRNA recovered from

tumor samples at 35 days, versus 6 days, post-castration. A key consequence of NFκB signaling during

the recurrence phase is the induction of CCL2 protein levels 35 days post-castration (Fig. 5A). Moreover,

the administration of a CCR2 antagonist to tumor bearing mice prevented castration-induced recurrence

(Fig. 5C), phenocopying the effects of -1d TNF blockade by etanercept treatment (Fig. 4A, -1d panel).

This observation is consistent with CCL2 expression downstream of TNF-NFκB signaling. Similarly, the

number of CCL2-staining cells in the TME was reduced in tumors from mice receiving etanercept to block

TNF signaling (Fig. 5B).

Increased tumor cell stemness following androgen deprivation drives pro-tumor TNF signaling

While the pleotropic signaling effects of TNF are well-established, an important mechanistic question in

the context of this report is whether changes in the tumor and surrounding TME are responsible for the

post-castration switch in TNF signaling. Two potential determinants are the spatial and temporal pattern

of TNF production/ secretion and the context of the cells that TNF signaling targets. Previously, we

showed that in the normal rat prostate TNF mRNA and protein increased as early as 8 hours post-

castration, with the most significant increase occurring within the stromal compartment of the prostate

(11). However, in PTEN-deficient murine tumors there is no discernable increase in TNF protein in the

first 24 hours post-castration (Fig. 1B). Alternatively, death receptor driven apoptosis can be triggered

by down-regulation of the CASP8 inhibitor CFLAR (56). Indeed, we previously found castration reduced

CFLAR expression within 12 hours post-castration in the normal rat prostate (12,13). Moreover, CFLAR

protein is up-regulated in CRPC patient samples (57). To investigate the mechanism of the late (5 weeks

post-castration) increase in TNF, we blocked AR action with enzalutamide in prostate cancer cell culture

models (10). After 3 weeks of culture in enzalutamide, TNF secretion and stemness were increased in

both C42 and LNCaP cell lines (Figs. 2 and Supplementary Figs. S2-S3). Higher levels of TNF secretion

occurred in cultures enriched for expression of one of the stem cell-like markers (Fig. 3). The increased

expression of stem cell-like genes following androgen deprivation may reflect the death of luminal-like

cells (low stemness) and selection for, or expansion of, basal-like cells (higher stemness) that have

increased levels of NFκB and therefore secrete more TNF protein. We observed an increase in basal

stem-like cells at 25 days post-castration in the PTEN-deficient tumor model (Fig. 1D), suggesting a

similar phenomenon is occurring *in vivo*. It is also possible that the increase in stemness that we observe

post-castration is due to lineage plasticity of pre-existing cells, rather than selection and expansion of

bone fide tumor stem cells (58). Regardless, it appears most likely that the increase in TNF secretion

observed 5 weeks post-castration is a consequence of changes in cellular context rather than a

promoter/enhancer-driven transcriptional response.

Immune suppression is a key consequence of increased tumor cell stemness

By investigating the switch in TNF signaling, we identified CCL2 secretion as a key TNF-NFκB-induced

event required for the onset of tumor recurrence in the PTEN-deficient GEMM (Fig. 5). Moreover, CCL2

is also required for recurrence in the genetically distinct Hi-MYC GEMM, suggesting CCL2 may be a

common driver for recurrence post-castration, at least in murine models of prostate cancer. Although

CCL2 has multiple pro-tumorigenic immune-modulating functions in prostate and other cancers (59), the

transcriptomic studies described in Figure 6 are consistent with TAM recruitment. Specifically, Gene

Ontology analysis (Fig. 6A-B) indicates that CCL2 induces immune cell chemotaxis and migration at the

time of recurrence, which is the behavior expected of TAMs as well as other myeloid immune suppressive

cells, such as myeloid-derived suppressor cells (MDSC). The heat map in Figure 6C showed up-

regulation of TAM-related genes, while IHC analysis in Figure 7 (and Supplementary Fig. S7)

demonstrated a coordinate increase in F4/80+ macrophages and CCL2 at the time of recurrence. The

latter two observations support a critical role for TAMs, but do not rule out a role for MDSC. Identifying

the precise tumor immune cell populations within the TME that modulate the response of cancers to

therapy is the focus of considerable ongoing research (60). TAMs are a heterogenous set of immune

cells which are predominantly pro-tumorigenic (61) and have been implicated in human and murine

prostate cancers (62,63). Although an oversimplification of the immune response to cancer, a balance

between anti-tumor cytolytic T cells and immunosuppressive myeloid cell populations may predict the

success of emerging immunotherapies. Indeed, variants of the CTL/TAM ratio – similar to our

observations in Figures 7 and 8 – are potential prognostic biomarkers in patients with melanoma (64),

breast cancer (65) and hepatocellular carcinoma (66). **Prospective: a model for the development of**

CRPC

The models we employed are anatomically similar to non-metastatic CRPC (nmCRPC) in that the tumor

is localized to the prostate, whereas in metastatic CRPC (mCRPC) the TME is derived from the metastatic

site, such as bone. Most CRPC is discovered clinically as mCRPC and indeed nmCRPC is likely a

transient state that progresses to mCRPC (67). Castration of mice bearing PTEN-deficient or Hi-MYC

prostate cancers induces regression and then recurrence – a pattern very similar to that seen in the

development of human CRPC. Importantly, HFUS imaging provides real-time volume data to ensure that

the complex kinetics of tumor volume changes can be used to accurately categorize the response of

each tumor to androgen deprivation. Supplementary Figure S8 is a mechanistic model consistent with

our findings. It proposes that TNF secretion into the TME first drives regression and subsequently

induces an immunosuppressive state that leads to recurrence. The specific signal may vary depending

on the TME context. For example, in some PTEN-deficient GEMMs, IL1ß triggers myeloid immune

suppression (68). Most of our analyses were performed on bulk tumor tissue – this represents a

significant limitation of the studies in this report. Future studies employing multi-modal single cell

analyses will be critical to elucidating the molecular interactions between tumor cells and the

myofibroblasts, endothelial cells and immune cells that constitute the TME. These studies are underway

in our laboratories. Although the CCL2 antagonist was effective at preventing recurrence in our GEMMs,

a monoclonal antibody which binds CCL2 failed in a phase 2 patient trial in mCRPC (69). This may have

been due to technical issues with the drug (69), or possibly heterogeneity among human CRPCs. Indeed,

Giancotti and colleagues identified three subtypes of CRPC (70) and proposed that one of these –

mesenchymal stem-like prostate cancer – might use CCL2 signaling to promote myeloid immune

suppression. Thus, in the future patients with this CRPC subtype may be candidates for anti-CCL2

therapies.

ACKNOWLEDGEMENTS

We acknowledge: Chawnshang Chang and Shu-Yuan Yeh (University of Rochester) for encouraging us

to investigate the pro-tumorigenic role of CCL2 in prostate cancer; Yvonne Saenger (Albert Einstein

College of Medicine) for helpful discussions on the ratio of CTLs and TAMs in tumor immune suppression,

Bo Xu (Roswell Park Comprehensive Cancer Center) for discussions on the histopathology of murine

prostate cancers and Michalis Mastri (Thermo-Fisher/ GIBCO) for assistance with computational analysis

of RNA sequence data.

REFERENCES

1. Huggins C, Hodges CV (1941) Studies on prostatic cancer. I. The effect of castration, of estrogen and of androgen injection on serum phosphatases in metastatic carcinoma of the prostate. Cancer Res 1:293-297.

2. Labrie F, Belanger A, Luu-The V, Labrie C, Simard J, Cusan L, Gomez J, Candas B (2005) Gonadotropin-releasing hormone agonists in the treatment of prostate cancer. Endocr Rev 26:361-379.

3. Sartor O, de Bono JS (2018) Metastatic Prostate Cancer. N Engl J Med 378:645-657.

4. Cancer Genome Atlas Research Network. Electronic address scmo, Cancer Genome Atlas Research N (2015) The Molecular Taxonomy of Primary Prostate Cancer. Cell 163:1011-1025.

5. Copeland BT, Pal SK, Bolton EC, Jones JO (2018) The androgen receptor malignancy shift in prostate cancer. Prostate 78:521-531.

6. Grbesa I, Augello MA, Liu D, McNally DR, Gaffney CD, Huang D, Lin K, Ivenitsky D, Goueli R, Robinson BD, Khani F, Deonarine LD, Blattner M, Elemento O, Davicioni E, Sboner A, Barbieri CE (2021) Reshaping of the androgen-driven chromatin landscape in normal prostate cells by early cancer drivers and effect on therapeutic sensitivity. Cell Reports 36:109625.

7. Cunha GR, Hayward SW, Wang YZ, Ricke WA (2003) Role of the stromal microenvironment in carcinogenesis of the prostate. Int J Cancer 107:1-10.

8. Krolewski JJ, Singh S, Sha K, Jaiswal N, Turowski SG, Pan C, Rich LJ, Seshadri M, Nastiuk KL (2022) TNF signaling is required for castration-induced vascular damage preceding prostate cancer regression. Cancers 14:6020.

9. Maolake A, Zhang R, Sha K, Singh S, Pan C, Xu B, Chatta G, Eng KH, Nastiuk KL, Krolewski JJ (2021) Glucocorticoid signaling delays castration-induced regression in murine models of prostate cancer. bioRxiv doi.org/10.1101/2021.10.11.463722.

10. Sha K, Yeh S, Chang C, Nastiuk K, Krolewski JJ (2015) TNF signaling mediates an enzalutamide-induced metastatic phenotype of prostate cancer and microenvironment cell co-cultures. Oncotarget 6:25726-25740.

11. Davis JS, Nastiuk KL, Krolewski JJ (2011) TNF is necessary for castration-induced prostate regression, whereas TRAIL and FasL are dispensable. Mol Endocrinol 25:611-620.

12. Nastiuk KL, Kim JW, Mann M, Krolewski JJ (2003) Androgen regulation of FLICE-like inhibitory protein gene expression in the rat prostate. J Cell Physiol 196:386-393.

13. Cornforth AN, Davis JS, Khanifar E, Nastiuk KL, Krolewski JJ (2008) FOXO3a mediates the androgen-dependent regulation of FLIP and contributes to TRAIL-induced apoptosis of LNCaP cells. Oncogene 27:4422-4433.

14. Wang S, Gao J, Lei Q, Rozengurt N, Pritchard C, Jiao J, Thomas GV, Li G, Roy-Burman P, Nelson PS, Liu X, Wu H (2003) Prostate-specific deletion of the murine Pten tumor suppressor gene leads to metastatic prostate cancer. Cancer Cell 4:209-221.

15. Ellwood-Yen K, Wongvipat J, Sawyers C (2006) Transgenic mouse model for rapid pharmacodynamic evaluation of antiandrogens. Cancer Res 66:10513-10516.

16. Singh S, Pan C, Wood R, Yeh C-R, Yeh S, Sha K, Krolewski JJ, Nastiuk KL (2015) Quantitative volumetric imaging of normal, neoplastic and hyperplastic mouse prostate using ultrasound. BMC Urol 15:97.

17. Pan C, Jaiswal Agrawal N, Zulia Y, Singh S, Sha K, Mohler JL, Eng KH, Chakkalakal JV, Krolewski JJ, Nastiuk KL (2020) Prostate tumor-derived GDF11 accelerates androgen deprivation therapy-induced sarcopenia. JCI Insight 5:e127018.

18. Schwartz LH, Litiere S, de Vries E, Ford R, Gwyther S, Mandrekar S, Shankar L, Bogaerts J, Chen A, Dancey J, Hayes W, Hodi FS, Hoekstra OS, Huang EP, Lin N, Liu Y, Therasse P, Wolchok JD, Seymour L (2016) RECIST 1.1-Update and clarification: From the RECIST committee. Eur J Cancer 62:132-137.

19. Macanas-Pirard P, Quezada T, Navarrete L, Broekhuizen R, Leisewitz A, Nervi B, Ramirez PA (2017) The CCL2/CCR2 Axis Affects Transmigration and Proliferation but Not Resistance to Chemotherapy of Acute Myeloid Leukemia Cells. PLoS One 12:e0168888.

20. Thalmann GN, Anezinis PE, Chang SM, Zhau HE, Kim EE, Hopwood VL, Pathak S, von Eschenbach AC, Chung LW (1994) Androgen-independent cancer progression and bone metastasis in the LNCaP model of human prostate cancer. Cancer Res 54:2577-2581.

21. Pruckler JM, Pruckler JM, Ades E (1995) Detection by polymerase chain reaction of all common *mycoplasma* in a cell culture facility. Pathobiol 63:9-11.

22. Dirks WG, Drexler HG (2005) Authentication of scientific human cell lines: easy-to-use DNA fingerprinting. Methods Mol Biol 290:35-50.

23. Lukacs RU, Goldstein AS, Lawson DA, Cheng D, Witte ON (2010) Isolation, cultivation and characterization of adult murine prostate stem cells. Nat Protoc 5:702-713.

24. Chougoni KK, Grossman SR (2020) Extraction of high-quality RNA from mouse pancreatic tumors. MethodsX 7:101163.

25. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR (2013) STAR: ultrafast universal RNA-seq aligner. Bioinformatics 29:15-21.

26. Li B, Dewey CN (2011) RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome. BMC Bioinformatics 12:323.

27. Robinson MD, McCarthy DJ, Smyth GK (2010) edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. Bioinformatics 26:139-140.

28. Law CW, Chen Y, Shi W, Smyth GK (2014) voom: Precision weights unlock linear model analysis tools for RNA-seq read counts. Genome Biol 15:R29.

29. Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, Smyth GK (2015) limma powers differential expression analyses for RNA-sequencing and microarray studies. Nucleic Acids Res 43:e47.

30. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski K, Dwight SS, Eppig JT, Harris MA, Hill DP, Issel-Tarver L, Kasarskis A, Lewis S, Matese JC, Richardson JE, Ringwald M, Rubin GM, Sherlock G (2000) Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. Nat Genet 25:25-29.

31. Gu Z, Eils R, Schlesner M (2016) Complex heatmaps reveal patterns and correlations in multidimensional genomic data. Bioinformatics 32:2847-2849.

32. Amezquita RA, Lun ATL, Becht E, Carey VJ, Carpp LN, Geistlinger L, Marini F, Rue-Albrecht K, Risso D, Soneson C, Waldron L, Pages H, Smith ML, Huber W, Morgan M, Gottardo R, Hicks SC (2020) Orchestrating single-cell analysis with Bioconductor. Nat Methods 17:137-145.

33. McCarthy DJ, Campbell KR, Lun AT, Wills QF (2017) Scater: pre-processing, quality control, normalization and visualization of single-cell RNA-seq data in R. Bioinformatics 33:1179-1186.

34. Kobak D, Berens P (2019) The art of using t-SNE for single-cell transcriptomics. Nat Commun 10:5416.

35. Zhou B, Jin W (2020) Visualization of Single Cell RNA-Seq Data Using t-SNE in R. Methods Mol Biol 2117:159-167.

36. Yu YP, Landsittel D, Jing L, Nelson J, Ren B, Liu L, McDonald C, Thomas R, Dhir R, Finkelstein S, Michalopoulos G, Becich M, Luo JH (2004) Gene expression alterations in prostate cancer predicting tumor aggression and preceding development of malignancy. J Clin Oncol 22:2790-2799.

37. Ross-Adams H, Lamb AD, Dunning MJ, Halim S, Lindberg J, Massie CM, Egevad LA, Russell R, Ramos-Montoya A, Vowler SL, Sharma NL, Kay J, Whitaker H, Clark J, Hurst R, Gnanapragasam VJ, Shah NC, Warren AY, Cooper CS, Lynch AG, Stark R, Mills IG, Gronberg H, Neal DE, CamCa PSG (2015) Integration of copy number and transcriptomics provides risk stratification in prostate cancer: A discovery and validation cohort study. EBioMedicine 2:1133-1144.

38. Lawson DA, Xin L, Lukacs RU, Cheng D, Witte ON (2007) Isolation and functional characterization of murine prostate stem cells. Proc Natl Acad Sci U S A 104:181-186.

39. Sackmann Sala L, Boutillon F, Menara G, De Goyon-Pelard A, Leprevost M, Codzamanian J, Lister N, Pencik J, Clark A, Cagnard N, Bole-Feysot C, Moriggl R, Risbridger GP, Taylor RA, Kenner L, Guidotti JE, Goffin V (2017) A rare castration-resistant progenitor cell population is highly enriched in Pten-null prostate tumours. J Pathol 243:51-64.

40. Wang ZA, Toivanen R, Bergren SK, Chambon P, Shen MM (2014) Luminal Cells Are Favored as the Cell of Origin for Prostate Cancer. Cell Reports 8:1339-1346.

41. Schalken J, Fitzpatrick JM (2016) Enzalutamide: targeting the androgen signalling pathway in metastatic castration-resistant prostate cancer. BJU Int 117:215-225.

42. Mulholland DJ, Kobayashi N, Ruscetti M, Zhi A, Tran LM, Huang J, Gleave M, Wu H (2012) Pten loss and RAS/MAPK activation cooperate to promote EMT and metastasis initiated from prostate cancer stem/progenitor cells. Cancer Res 72:1878-1889.

43. Davis JS, Nastiuk KL, Krolewski JJ (2011) TNF is necessary for castration-induced prostate regression, whereas TRAIL and FasL are dispensable. Mol Endocrinol 25:611-620.

44. Sha K, Yeh S, Chang C, Nastiuk KL, Krolewski JJ (2015) TNF signaling mediates an enzalutamide-induced metastatic phenotype of prostate cancer and microenvironment cell co-cultures. Oncotarget 6:25726-25740.

45. Kaymakcalan Z, Sakorafas P, Bose S, Scesney S, Xiong L, Hanzatian DK, Salfeld J, Sasso EH (2009) Comparisons of affinities, avidities, and complement activation of adalimumab, infliximab, and etanercept in binding to soluble and membrane tumor necrosis factor. Clin Immunol 131:308-316.

46. Matsushima K, Larsen CG, DuBois GC, Oppenheim JJ (1989) Purification and characterization of a novel monocyte chemotactic and activating factor produced by a human myelomonocytic cell line. J Exp Med 169:1485-1490.

47. Zhang J, Patel L, Pienta KJ (2010) CC chemokine ligand 2 (CCL2) promotes prostate cancer tumorigenesis and metastasis. Cytokine Growth Factor Rev 21:41-48.

48. Fei L, Ren X, Yu H, Zhan Y (2021) Targeting the CCL2/CCR2 Axis in Cancer Immunotherapy: One Stone, Three Birds? Front Immunol 12:771210.

49. Jin F, Li Y, Dixon JR, Selvaraj S, Ye Z, Lee AY, Yen CA, Schmitt AD, Espinoza CA, Ren B (2013) A high-resolution map of the three-dimensional chromatin interactome in human cells. Nature 503:290-294.

50. Webber MM, Trakul N, Thraves PS, Bello-DeOcampo D, Chu WW, Storto PD, Huard TK, Rhim JS, Williams DE (1999) A human prostatic stromal myofibroblast cell line WPMY-1: a model for stromal-epithelial interactions in prostatic neoplasia. Carcinogenesis 20:1185-1192.

51. Larionova I, Tuguzbaeva G, Ponomaryova A, Stakheyeva M, Cherdyntseva N, Pavlov V, Choinzonov E, Kzhyshkowska J (2020) Tumor-associated macrophages in human breast, colorectal, lung, ovarian and prostate cancers. Frontiers Oncology 10:566511.

52. Loberg RD, Ying C, Craig M, Yan L, Snyder LA, Pienta KJ (2007) CCL2 as an important mediator of prostate cancer growth in vivo through the regulation of macrophage infiltration. Neoplasia 9:556-562.

53. Kerr JF, Wyllie AH, Currie AR (1972) Apoptosis: a basic biological phenomenon with wide-ranging implications in tissue kinetics. Br J Cancer 26:239-257.

54. Godoy A, Montecinos VP, Gray DR, Sotomayor P, Yau JM, Vethanayagam RR, Singh S, Mohler JL, Smith GJ (2011) Androgen deprivation induces rapid involution and recovery of human prostate vasculature. Am J Physiol Endocrinol Metab 300:E263-275.

55. Buttyan R, Ghafar MA, Shabsigh A (2000) The effects of androgen deprivation on the prostate gland: cell death mediated by vascular regression. Curr Opin Urol 10:415-420.

56. Irmler M, Thome M, Hahne M, Schneider P, Hofmann K, Steiner V, Bodmer J-L, Schröter M, Burns K, Mattmann C, Rimoldi D, French LE, Tschopp J (1997) Inhibition of death receptor signals by cellular FLIP. Nature 388:190-195.

57. McCourt C, Maxwell P, Mazzucchelli R, Montironi R, Scarpelli M, Salto-Tellez M, O'Sullivan JM, Longley DB, Waugh DJ (2012) Elevation of c-FLIP in castrate-resistant prostate cancer antagonizes therapeutic response to androgen receptor-targeted therapy. Clin Cancer Res 18:3822-3833.

58. Sandhu HS, Portman KL, Zhou X, Zhao J, Rialdi A, Sfakianos JP, Guccione E, Kyprianou N, Zhang B, Mulholland DJ (2022) Dynamic plasticity of prostate cancer intermediate cells during androgen receptor-targeted therapy. Cell Reports 40:111123.

59. Hao Q, Vadgama JV, Wang P (2020) CCL2/CCR2 signaling in cancer pathogenesis. Cell communication and signaling 18:82.

60. Mellman I, Chen DS, Powles T, Turley SJ (2023) The cancer-immunity cycle: Indication, genotype, and immunotype. Immunity 56:2188-2205.

61. Kloosterman DJ, Akkari L (2023) Macrophages at the interface of the co-evolving cancer ecosystem. Cell 186:1627-1651.

62. Gollapudi K, Galet C, Grogan T, Zhang H, Said JW, Huang J, Elashoff D, Freedland SJ, Rettig M, Aronson WJ (2013) Association between tumor-associated macrophage infiltration, high grade prostate cancer, and biochemical recurrence after radical prostatectomy. Am J Cancer Res 3:523-529.

63. Escamilla J, Schokrpur S, Liu C, Priceman SJ, Moughon D, Jiang Z, Pouliot F, Magyar C, Sung JL, Xu J, Deng G, West BL, Bollag G, Fradet Y, Lacombe L, Jung ME, Huang J, Wu L (2015) CSF1 receptor targeting in prostate cancer reverses macrophage-mediated resistance to androgen blockade therapy. Cancer Res 75:950-962.

64. Gartrell-Corrado RD, Chen AX, Rizk EM, Marks DK, Bogardus MH, Hart TD, Silverman AM, Bayan CY, Finkel GG, Barker LW, Komatsubara KM, Carvajal RD, Horst BA, Chang R, Monod A, Rabadan R, Saenger YM (2020) Linking Transcriptomic and Imaging Data Defines Features of a Favorable Tumor Immune Microenvironment and Identifies a Combination Biomarker for Primary Melanoma. Cancer Res 80:1078-1087.

65. DeNardo DG, Brennan DJ, Rexhepaj E, Ruffell B, Shiao SL, Madden SF, Gallagher WM, Wadhwani N, Keil SD, Junaid SA, Rugo HS, Hwang ES, Jirstrom K, West BL, Coussens LM (2011) Leukocyte complexity predicts breast cancer survival and functionally regulates response to chemotherapy. Cancer Discov 1:54-67.

66. Xin H, Liang D, Zhang M, Ren D, Chen H, Zhang H, Li S, Ding G, Zhang C, Ding Z, Wu L, Han W, Zhou W, Chen Y, Luo H, Wang Y, Zhang H, Liu S, Li N (2021) The CD68+ macrophages to CD8+ T-cell ratio is associated with clinical outcomes in hepatitis B virus (HBV)-related hepatocellular carcinoma. HPB 23:1061-1071.

67. Figueiredo A, Costa L, Mauricio MJ, Figueira L, Ramos R, Martins-da-Silva C (2022) Nonmetastatic Castration-Resistant Prostate Cancer: Current Challenges and Trends. Clin Drug Investig 42:631-642.

68. Wang D, Cheng C, Chen X, Wang J, Liu K, Jing N, Xu P, Xi X, Sun Y, Ji Z, Zhao H, He Y, Zhang K, Du X, Dong B, Fang Y, Zhang P, Qian X, Xue W, Gao WQ, Zhu HH (2023) IL-1beta Is an Androgen-Responsive Target in Macrophages for Immunotherapy of Prostate Cancer. Adv Sci (Weinh) 10:e2206889.

69. Pienta KJ, Machiels JP, Schrijvers D, Alekseev B, Shkolnik M, Crabb SJ, Li S, Seetharam S, Puchalski TA, Takimoto C, Elsayed Y, Dawkins F, de Bono JS (2013) Phase 2 study of carlumab (CNTO 888), a human monoclonal antibody against CC-chemokine ligand 2 (CCL2), in metastatic castration-resistant prostate cancer. Investigational new drugs Invest New Drugs 31:760-768.

70. Su W, Han HH, Wang Y, Zhang B, Zhou B, Cheng Y, Rumandla A, Gurrapu S, Chakraborty G, Su J, Yang G, Liang X, Wang G, Rosen N, Scher HI, Ouerfelli O, Giancotti FG (2019) The Polycomb Repressor Complex 1 Drives Double-Negative Prostate Cancer Metastasis by Coordinating Stemness and Immune Suppression. Cancer Cell 36:139-155 e110.

71. Horoszewicz JS, Leong SS, Kawinski E, Karr JP, Rosenthal H, Chu TM, Mirand EA, Murphy GP (1983) LNCaP model of human prostatic carcinoma. Cancer Res 43:1809-1818.

72. Tsuchiya S, Yamabe M, Yamaguchi Y, Kobayashi Y, Konno T, Tada K (1980) Establishment and characterization of a human acute monocytic leukemia cell line (THP-1). Int J Cancer 26:171-176.

73. Arefieva TI, Kukhtina NB, Antonova OA, Krasnikova TL (2005) MCP-1-stimulated chemotaxis of monocytic and endothelial cells is dependent on activation of different signaling cascades. Cytokine 31:439-446.

74. Franklin RA, Liao W, Sarkar A, Kim MV, Bivona MR, Liu K, Pamer EG, Li MO (2014) The cellular and molecular origin of tumor-associated macrophages. Science 344:921-925.

FIGURE 1

Figure 1. Castration induces tumor recurrence, and an increase in TNF and stemness in prostate tumors from PTEN-deficient mice. Mice bearing PTEN-deficient tumors were either castrated (Cx1-4) or sham-operated (Sh1-3) and serially imaged using HFUS. In some cases, tumor tissue was recovered for further analysis. **A**. 3D reconstruction of tumor-bearing ventral lobes (pseudo-colored green) performed as described in Materials & Methods. **B**. Tumor volume kinetics post-castration. Tumor volume was determined from HFUS images using Amira software. Mean tumor volume at the indicated times post-castration (n=4, red) or post-sham operation (n=3, blue) is shown as percent pre-castration volume (mean [%] ± SEM). **C**. TNF levels in tumors tissue extracts were measured by ELISA and normalized to total protein in the tissue sample, at the indicated times post castration. One-way ANOVA followed by Tukey-Kramer HSD test revealed a difference in TNF levels between 0 and 5 weeks (p<0.01). **D**. LSC FACS analysis of tumor cells from sham (left) and castrated (Cx; right) PTEN-deficient mice. Tumors were disaggregated and cells were processed for Lin⁻/Sca-1/CD49f FACS analysis (23). Relative fluorescence intensities are shown as contour plots. The LSC^{hi} population is labeled 'Basal/stem'. The population in the lower right quadrant (LSC^{med}) corresponds to luminal progenitors. A representative analysis is shown. Sham LSC FACS was replicated 5 times (mean[%]±SEM = 27.0±4.0); 25d post-castration LSC FACS was replicated 4 times (mean[%]±SEM = 38.2±1.7). Student's t-test revealed a difference between Sham and Cx in the LSC^{hi} population (p<0.05).

FIGURE 2

A

B

day 0   day 21

CD49f-Ax647

CD166-FITC

C

C4-2 cells were grown in media plus 10% serum and treated with vehicle (blue) or 10 µM enzalutamide (Enz; red) for the indicated time. **A.** C4-2 cell growth curve in the presence of enzalutamide. Trypan-blue excluding live cells were counted microscopically at the indicated times. Data are shown as mean ± s.e.m. (n=3). Two-way ANOVA followed by Tukey-Kramer HSD test revealed a difference in cell number for the Enz treated culture at 28d versus 0d ($p<0.01$). **B.** FACS analysis for basal cell stemness markers in enzalutamide treated C4-2 cells. Cells grown in enzalutamide for the indicated time, were incubated with the fluor-labeled antibodies, analyzed by FACS and relative fluorescence intensity represented as dot plots. The fraction of the cells that correspond to the CD49f^{hi}/CD166^{hi} population (upper, right quadrant) is indicated. Additional time points are in Supplementary Figure S2. **C.** Fraction of cells scoring CD49f^{hi}/CD166^{hi} and corresponding TNF secretion, at the indicated time of culture in media plus enzalutamide. Data from FACS analyses (see Fig. S2) is plotted on the left and TNF levels in the media determined by ELISA is plotted on the right. Values are given as mean ± s.e.m. (*n* = 2). ******, *p* < 0.01 ********p* < 0.001 (two-way ANOVA followed by Tukey-Kramer HSD test).

Figure 2.   Extended enzalutamide treatment selects for basal stemness and TNF secretion.

FIGURE 3

A

B

C

Figure 3. **Enzalutamide treatment selects for a TNF-expressing basal stem cell population.** C4-2 cells were grown for 20 days in media plus 10% serum containing 10 µM enzalutamide, enriched for CD166hi expression. ELISA was used to measure TNF and colony formation. **A.** CD166-enrichment was performed via FACS to sort CD166hi and CD166lo C4-2 cells treated as indicated. The purity of the unsorted and sorted CD166hi and CD166lo populations are shown. **B.** TNF secretion of the indicated cultures (left to right): unsorted, vehicle-treated (veh); unsorted, enzalutamide-treated (enz); and sorted, enzalutamide-treated (CD166lo, CD166hi). TNF measured by ELISA, and plotted as mean ± s.e.m. (n=3). **p<0.01, ***p<0.001 (One-way ANOVA followed by Tukey-Kramer HSD test). **C.** Colony formation assay of enriched CD166hi and CD166lo cell populations sorted as in **A**. Seeding densities were 500, 1000, and 1500 cells/well, respectively. Colony counts per well are plotted. *p<0.05 (Student's unpaired t-test).

FIGURE 4

A

Etanercept	Regression	Recurrence
None	4/4	3/4
-3d	0/5	NA
-1d	4/5	0/4
+3d	3/3	2/3

B

Sham vs CX_6d

Genesets	SIZE	NES	p.val	FDR
PHONG_TNF_RESPONSE_VIA_P38_PARTIAL	144	-2.98	<0.01	5.16E-05
PHONG_TNF_RESPONSE_VIA_P38_COMPLETE	208	-2.94	<0.01	5.65E-05
RUAN_RESPONSE_TO_TNF_TROGLITAZONE_DN	39	-2.70	<0.01	2.97E-04
PHONG_TNF_TARGETS_UP	60	-2.65	<0.01	3.72E-04
RUAN_RESPONSE_TO_TNF_DN	80	-2.62	<0.01	4.81E-04
ZHOU_TNF_SIGNALING_30MIN	51	-1.97	<0.01	2.35E-02
ZHANG_RESPONSE_TO_IKK_INHIBITOR_AND_TNF_UP	190	-1.88	0.01	3.52E-02
ZHOU_TNF_SIGNALING_4HR	49	-1.85	0.01	4.09E-02
WANG_TNF_TARGETS	18	-1.77	0.02	5.82E-02
RUAN_RESPONSE_TO_TNF_TROGLITAZONE_UP	15	-1.68	0.03	8.09E-02

Sham vs CX_35d

Genesets	SIZE	NES	p.val	FDR
ZHANG_RESPONSE_TO_IKK_INHIBITOR_AND_TNF_UP	193	3.96	<0.01	9.35E-06
PHONG_TNF_TARGETS_UP	61	3.96	<0.01	9.35E-06
PHONG_TNF_RESPONSE_NOT_VIA_P38	307	3.72	<0.01	9.35E-06
WANG_TNF_TARGETS	19	3.5032	<0.01	9.35E-06
TIAN_TNF_SIGNALING_VIA_NFKB	27	3.2884	<0.01	9.35E-06
TIAN_TNF_SIGNALING_NOT_VIA_NFKB	21	2.7396	<0.01	1.64E-04
SANA_TNF_SIGNALING_UP	65	2.7088	<0.01	2.21E-04
PHONG_TNF_RESPONSE_VIA_P38_PARTIAL	149	2.3381	<0.01	2.27E-03
SANA_TNF_SIGNALING_DN	82	2.1602	<0.01	6.18E-03
ZHOU_TNF_SIGNALING_30MIN	51	1.8286	<0.01	3.32E-02
ZHOU_TNF_SIGNALING_4HR	49	1.7752	0.02	4.19E-02

Figure 4. Differentially-timed TNF signaling blockade has distinct effects on tumor growth. Mice with PTEN-deficient tumors were castrated and etanercept (sTNFR2-Fc) was administered at the indicated times: three days before (-3d), one day before (-1d), or three days after (+3d) surgical castration. **A.** Tumor volume kinetics. Tumor volumes were determined by serial HFUS imaging and 3D reconstruction and plotted after normalizing to the initial tumor volume for each mouse. Different colors/symbols represent individual mice. The table summarizes regression and recurrence incidence for each experiment. Growth curves for castrated mice that did not receive etanercept (None) are not shown. The criteria for regression and recurrence are described in the Materials and Methods. Statistical analysis is provided in Supplementary Table S3A. **B.** Gene set enrichment analysis. RNA was extracted from PTEN-deficient tumors from the following mice: sham-operated, 6d post-castration and 35d post-castration. GSEA was performed on bulk RNA-seq data using gene sets corresponding to TNF-related pathways.   Normalized enrichment scores (NES) from RNA from tumors at 6d (down-regulated, blue) and 35d (up-regulated, red) post-castration were compared to the sham group. Gene sets common to the two analyses are highlighted in yellow.

FIGURE 5

A

B

C

PTEN$^{-/-}$ GEMM
CX+CCR2a:
Recurrence = 0/6

D

hi-MYC GEMM
CX+CCR2a:
Recurrence = 1/5

CX+vehicle:
Recurrence = 4/5

Figure 5. CCL2-CCR2 signaling is required for castration-resistant recurrence. Mice bearing PTEN-deficient (**A-C**) or hi-MYC (**D**) tumors were sham-operated or castrated at the indicated times. Volume was monitored by HFUS. Some mice received etanercept (E) or the CCR2a, as indicated. **A**. Tumor CCL2 levels following castration. CCL2 levels were measured by ELISA on the same tumor extracts prepared for Fig. 1C, normalized to total protein and plotted as open circles. The bar marks the mean. Two-way ANOVA followed by Tukey-Kramer HSD test revealed a difference in CCL2 levels between 0 and 5 weeks *(p < 0.0001).(* **B.** Representative images of IHC-stained CCL2$^+$ cells (red) in tumors from the indicated treatment groups. CCL2+ cells were counted and plotted. *******p* < 0.01, **p* < 0.05 (Two-way ANOVA followed by Tukey-Kramer HSD test). **C.** Treatment with a CCR2 antagonist phenocopies -1d etanercept treatment (Fig. 4A) in PTEN-deficient tumor-bearing mice. Tumor volumes were determined by serial HFUS imaging and 3D reconstruction and plotted after normalizing to the initial tumor volume for each mouse (left). Different colors/symbols represent individual mice. The recurrence incidence is summarized (right). **D.** CCR2 antagonist suppresses castration-induced regression in Hi-MYC prostate cancer-bearing mice. Blue boxes: mice were castrated and received vehicle. Red circles: mice were castrated and received CCR2a prior to castration. N=5 for each group. The recurrence incidence is summarized (right). Tumor volume kinetics for individual mice are in Supplementary Figure S5. Statistical analysis for **C-D** is provided in Supplementary Table S3B.

FIGURE 6

Figure 6. Transcriptomic changes in PTEN-deficient prostate cancers following CCL2 signaling blockade. RNA was extracted from PTEN-deficient tumors from the following mice: sham-operated (Sham), 35d post-castration (35d post-CX) and 35d post-castration after receiving CCR2a (35dp-CX+CCR2a). DGE analysis was performed on bulk RNAseq data (Fig. S6) followed by Gene Ontology analysis of biological processes comparing RNAseq transcript sets derived from tumors in mice that were sham-operated vs mice 35d post-castration (**A**) or from tumors in mice 35d post-castration vs mice treated with CCR2a 35d post-castration (**B**). Red-colored bars represent up-regulation and blue-colored bars represent down-regulation. Fold-enrichment >5. **C.** TAM-related transcripts regulated by CCL2 signaling. Heatmap representation of mRNA levels of select TAM-associated genes for the same two comparisons shown in **A** (left) and **B** (right).

FIGURE 7

Figure 7. TNF and CCR2 signaling blockade suppress castration-induced immunosuppression at late times following castration. Mice bearing PTEN-deficient tumors were sham-operator or castrated (Cx) and tumors harvested at the indicated times post-castration. Some mice received etanercept (Etan) or the CCR antagonist BMS IHC staining for stromal F4/80+ macrophages and CD8+ T cells was performed and the number of stained cells per unit area were plotted. **A**. TNF signaling blockade. Images of representative IHC-stained tissue sections for the surface marker protein noted on the left side. Mice were treated as shown across top and the cell densities plotted for each of the treatments, below. **B**. CCL2 signaling blockade. Similar to **A** with both surface marker protein and corresponding mouse treatments shown along the top. A complete set of images are provided in Supplementary Figure S7C. Staining and counting of the Cx-35d samples in **A** and **B** were performed separately. Data are given as mean ± s.e.m. (n=4). *p<0.05, **p<0.01, ***p<0.001 (One-way ANOVA followed by Tukey-Kramer HSD test). *P*-value for the differences for CD8+ T cells and F4/80 macrophages due to CCR2a treatment was >0.05 (not statistically significant).

FIGURE 8

Figure 8. Tumor recurrence following castration correlates with an immunosuppressive tumor microenvironment. **A.** Post-castration tumor volume kinetics for three representative PTEN-deficient mouse tumors. Tumor volumes were determined by serial HFUS imaging and 3D reconstruction and plotted after normalizing to the initial tumor volume for each mouse. The average starting volume for these tumors was ~960 mm^3. **B.** t-SNE plots for single cell RNAseq analysis. At 42d post-castration, mice were sacrificed, tumor tissue was disaggregated into single cell preparations, RNA sequenced and analyzed as described in the Methods. Dimensionality reduction of cell transcriptomes from all three mice was performed with tSNE to produce the plot in far left panel (Aggregate). Cells are assigned to 6 subset populations as coded in the key in the lower right corner. Color-coded cells from each of the three mice corresponding to the response patterns in **A** are shown in the remaining three panels of **B**. Cells which were in the aggregate population, but not identified in individual mice are grey colored. TAMs (defined in panel C) are within the large, heterogeneous myeloid population (green). **C.** CTL:TAM ratios for regressing, stable or recurrent mouse tumors. Transcriptomic data from **B** was used to define TAMs (CD45 and CD68and CD80; lighter grey) and CTLs (CD45 and [CD3e or Cd3d or CD3g] and [CD8a or CD8b1]; darker grey). See text for discussion.