Title: **Evolutionary change is remarkably time-independent across scales**

Authors: Stephen P. De Lisle[1]*, Erik I. Svensson[1,2]

Affiliations:
[1] Department of Environmental and Life Science, Karlstad University, Karlstad, Sweden

[2] Department of Biology, Lund University, Lund, Sweden

*Corresponding author. Email: stephen.de.lisle@kau.se

Classification: Biological Sciences, Evolution

Keywords: Evolutionary rate, microevolution, macroevolution, stasis paradox

Abstract Apparent time-scaling of evolutionary rates presents an evolutionary dilemma. Rates of
molecular, phenotypic, and lineage diversification typically scale negatively with time interval,
raising questions about the processes driving this seemingly general pattern. Here we show that
much apparent temporal scaling is an inevitable outcome of plotting a ratio against its
denominator. Highly unlikely relationships between timescale and accumulated evolutionary
change are required to produce anything other than negative relationships between rate and time.
Rate-time scaling relationships reflect unscaled magnitudes of evolution that are largely
unrelated to time, consistent with uniformitarianism and reframing the widely-held view of
disparate pace of micro- and macroevolution. Time-independence of evolutionary change raises
new questions about the factors that generate temporal consistency in evolution.

Significance Statement A longstanding finding across molecular, phenotypic, phylogenetic and
fossil data is that evolutionary rates depend strongly on the timescale over which evolution is
measured. This seemingly general pattern has stimulated many hypotheses of some shared
general features of the underlying processes that shape evolutionary rates. Here, we show that
evolutionary rates will inevitably correlate negatively with time interval, even if evolutionary
processes are time-independent. While the statistical effects behind this pattern are known, our
new analyses demonstrates that evolutionary rate-time relationships are nearly inevitable.
Empirical data suggest variable evolutionary rates across time, but no time-dependency or
systematic difference between micro- and macroevolutionary scales. Thus, evolutionary change
is largely uniform and independent of timescale.

Main Text: Evolutionary rates vary with the time interval over which they are measured. A seemingly general pattern with faster evolutionary rate over short timescales, and slower rates over longer timescales has been recognized for decades (1-5). Such apparent time-dependence has subsequently been documented consistently for nearly every possible measure of evolutionary rate (6), including rates of morphological evolution, rates of molecular evolution (7), and rates of lineage formation on phylogenies and in the fossil record (Figure 1). Negative rate-time scaling relationships were initially proposed as an artifact that needed to be taken in to account, because evolutionary rates over different time intervals were not directly comparable (5). Yet these negative relationships between evolutionary rate and time continue to generate discussions and stimulate new hypotheses about the possible shared features of the underlying evolutionary processes that could explain such general time-scaling (6, 8). Recent technical and statistical explanations to these apparent negative rate-time scaling relationships focus on sampling biases (9), such as a the "push of the past" (6) model mis-specification (2), and measurement error (10). This puzzling and general phenomenon of apparent widespread negative rate-time scaling relationships has been recently highlighted as a major unsolved challenge in evolutionary biology (2, 6, 8). The renewed interest in such apparent time-dependence of evolutionary rates has re-kindled long-standing ideas and hypotheses about fundamental disparities between micro- and macroevolution (11-15).

Here we show that much of this apparent paradox of strong time-dependency of evolutionary rate is readily explained by the inevitable negative correlation generated by plotting a ratio against its own denominator (16). While this is a well-known statistical issue across a range of scientific fields (5, 17-20), recent discussions of rate-time scaling patterns in evolutionary biology have either have either downplayed the importance of the mathematical dependence of rate and time (e.g., Box 1 in 2, 6, 8), or have focused on explanations arising from

specific statistical models used to estimate rates in a single context (9). By revisiting the

functional dependence of rates and time, analyzing large existing datasets and using new

simulations, we show that a negative relationship between evolutionary rate and time is nearly

inevitable, regardless of the underlying evolutionary process. We show that the key feature

underlying negative rate-time relationships in empirical data is in fact a weak, null, or variable

relationship between magnitude of evolution and time. Uninformed focus on rate-time

relationships therefore provide little insight into underlying evolutionary processes even under

highly idealized conditions. Instead, the empirical evidence suggest remarkable time-

independence and uniformity of evolutionary change across timescales and datasets.

The negative rate-time association is nearly inevitable

Estimates of evolutionary rate can be modelled as a function of evolutionary change scaled by

time:

$$y = d/t \tag{1}$$

where y is the rate-measure, d is some measure of evolutionary change (e.g., change in

population mean phenotype, base pair substitutions, log change in number of species in a clade)

and t is some measure of time interval over which change was observed or inferred (e.g.

generations, or years). Some classical estimates of evolutionary rate are calculated directly by

dividing phenotypic change with time, the latter typically measured in generations. For example,

Darwins (21) and Haldanes (4) have been used to describe morphological rate, and other types

of rate estimators measure per-capita origination rate of fossil lineages (22) and phylogenetic net

diversification rate (23). Other estimates are obtained from more sophisticated statistical models

of evolution (24-26). Regardless, all evolutionary rate measures are, per definition, scaled by

time (e.g., waiting times between observed branching events on a phylogeny are a direct

component of the likelihood in in maximum likelihood models of birth-death diversification

(24)).

The inevitable mathematical dependence between rate and time is an obvious challenge

in interpreting the time-dependence of evolutionary rate (5). Even a random association between

evolutionary change and time will result in a strong negative scaling relationship between rate

and time (27). The strength of this negative relationship will inevitably increase when a greater

range of timescales is sampled (16). The problem of interpreting such inevitable negative

relationships between a rate and time interval have been discussed in several fields outside

evolutionary biology, including geology (28) and paleontology (29). The problem has also been

raised in evolutionary biology, both in the context of rates of morphological evolution (3, 5, 18,

30-32) and lineage diversification (33-35).   Here, we revisit and elaborate on the issues with

interpreting rates as a function of time.

It is useful to ask, under what scenarios do we expect null or positive relationships

between rate and time? Consider rate as a function of time,

$$y = f(t) \tag{2}$$

106   Where y is evolutionary rate (as in eqn 1). We also consider the amount, or magnitude, of

107   accumulated evolution (rate numerator) as a function of time,

$$d = g(t) \tag{3}$$

where d is the accumulated evolutionary change . This could be the change in trait means

between allochronic or synchronic populations, difference in log number of lineages between

two sampling points in time (e.g., the stem age of a clade and present), or any other measure of

evolutionary change. Rates, $y = d/t$, will be constant over time when d scales linearly with t,

i.e. $g'(t)$ is a positive constant over the timescale in question. Rates will increase with time, i.e.

$f'(t) > 0$ when $g''(t) > 0$; that is, rates increase over time when the amount of evolutionary

change is an accelerating function of time.

Noteworthy is that under a null relationship between d and t, such that $g'(t) = 0$, the

logarithmic regression of rate versus time ($\log(y) \sim \mu + \beta*\log(t)$) will yield a negative slope

estimate of $\beta = -1$ (5). This can be seen most easily by expressing equation 3 as a power

function, e.g., $d = t^0$, in which case we can describe the regression of $\log(y) \sim \beta*\log(t)$ as

$\log(t^0/t) \sim \beta*\log(t)$; solving for $\beta = \log(t^{-1})/\log(t) = -1$.

This negative slope estimate provides one cutoff of interest in assessing rate-time scaling

relationships: log-log regressions of rate vs time that are shallower than or not significantly

different from -1 are consistent with either a null relationship evolutionary change and time (time

independence) or an increasing relationship between change and time, respectively.

The relationships between equations 1-3 are illustrated in Figure 2, which shows the

relationship between rate and time. Figure 2A illustrate differing relationships between rate

numerator and time illustrated in Figure 2B for a range of power functions. There are few

theoretical models that predict the type of accelerating relationship between magnitude of

evolution and time that are needed to produce a positive relationship between evolutionary rate

and time. For example, neutral and adaptive models of phenotypic and molecular evolution

predict either no relationship with time or linear accumulation (constant change) with time (36).

Similarly, birth-death models of lineage accumulation also predict either a linear or a

decelerating accumulation of log lineage number through time under most conditions. Even

models of time-dependent lineage diversification make no prediction about accelerating log

lineage number versus interval time across a sample of clades. Moreover, lineage-through-time

(LLT) plots for single clades diversifying under such a process can produce complex lineage

through time plots (37) that are consistent with several different underlying diversification

processes (38).

Some biological and ecological scenarios could potentially lead to accelerating change

with time, and thus generate positive relationships between evolutionary rates and time (Figure

2). Such scenarios include "diversity begets diversity" models of lineage diversification (24, 39),

evolution by positive frequency-dependent selection (40), runaway sexual selection driven by

female choice (41-43), "tipping points" in the evolution of reproductive isolation (44) and

populations crossing critical thresholds followed by phase transitions between alternative stable

states (45). How common such ecological and evolutionary scenarios are in nature is an open

empirical question. However, given the transient nature of such rapid and self-reinforcing

evolutionary processes driven by positive feedbacks (42, 43), we suspect that positive time-rate

scenarios are unlikely to be sustained over longer macroevolutionary time scales and should

quickly reach stable equilibria and largely be transient. Thus, such rapid and self-reinforcing

evolutionary processes are unlikely to proceed unchecked over millennia (45), as would be

required to generate positive rate-time relationships in empirical data such as that shown in

Figure 1.

Data and simulation reveal persistence of inevitable scaling relationships

Reanalysis of empirical datasets of evolutionary rate-time associations reveals little

consistent patterns in the relationship between amount of evolutionary change and time, despite a

pervasive negative relationship between rate and time. This holds for body size evolution (Figure

1A, E), where past work (18, 30, 31) has shown that body size changes are largely unrelated to

time, and in time-series of extant populations found no detectable change (46). Although a log-

log regression of rate vs time yields a slope that is nominally steeper than -1 (β = -1.019, 95%CI

-1.031, -1.007) we recover a breakpoint in the relationship between morphological change and

time ($F_{2, 5388}$ = 207.2, P < 2.2 x 10^{-6}, breakpoint = -2.63 log MY, SE = 0.147). The result is an

estimated weak negative relationship between time and evolutionary change, followed by an

increasing slope after the breakpoint (Fig. 1A, E), identical to that found by (30), but no overall

slowdown in accumulated evolutionary change across the entire time-series

Similar results are seen in patterns of lineage accumulation in the fossil record (Figure

1B, F). We first found evidence for a strong negative rate-time relationship reflecting weak and

variable relationship between evolutionary change and time (breakpoint regression $F_{2, 86}$ = 3.199,

P = 0.046, breakpoint = -109 MY, SE = 33.13). However, this regression model was not a

significantly better fit than a model with only an intercept; $F_{3, 86}$ = 2.6, P = 0.057. Consistent

with this finding, a log-log regression between rate and time yields a slope that is significantly

shallower than -1 (β = -0.24, 95%CI -0.36, -0.11)

On phylogenetic trees (Figure 1C, D, G, H), we also find a consistent striking negative

relationships between rate and time (8) that in fact corresponds to underlying variable association

between diversity and time (33, but see 47). We recover a weak positive linear relationship

between diversity and time in one dataset ($F_{1, 112}$ = 14.17, P = 0.000267; Figure 1H)

corresponding to a log-log regression that is significantly shallower than -1 (β = -0.53, 95%CI -

0.68, -.39), and a piecewise linear relationship in another dataset (breakpoint regression $F_{2, 430}$ =

12.67, P = 4.474 x10^{-6}, breakpoint = 4.6 log MY, SE = 0.108) corresponding to a log-log

regression of rate and time with a slope that is not significantly different from -1 (β = -1.04,

95%CI -1.1, -0.98).

Interestingly, we even recover a nonlinear negative relationship between evolutionary

rate and time when examining phenotypic change that occurred within a timeframe less than the

average generation time of the study organism (in this case, primarily long-lived fish; Figure 1E).

This within-generation dataset had a log-log regression of rate and time with a slope that was not

significantly different from -1 (β = -0.83, 95%CI -1.08, -0.58). This final example, again

demonstrates the appearance of a rate-time scaling relationship for a case that obviously defies

evolutionary explanation based on between-generation change. This conclusion is further

strengthened by the fact that the slope estimate does not significantly differ from the expectation

under a random relationship between rate numerator and time. That is, scaling phenotypic

change by time in order to calculate rate generates a non-linear negative relationship with time

even for this within-generation change (Figure 1E, J), that is likely to reflect phenotypic

plasticity, culling, or other non-evolutionary within-generational environmental effects.

In summary: the striking negative rate-time relationship shared by four independent

evolutionary datasets and one non-evolutionary (within generation) dataset reflects, primarily, a

lack of strong or consistent association between the amount of evolutionary change (rate

numerator) and the timescale over which this change took place (Figure 1 F-J). Although some

of these case-specific results have been pointed out before (e.g., (30), whose analysis we have in

part replicated here), we have shown that a decoupling of amount of evolution and timescale of

measurement is likely a general feature across a range of evolutionary datasets, from

morphological change to lineage diversification, that together describe the pace of life. In only

one case did we find any evidence of a significant negative association between evolutionary

change and time that was less than -1 on a log scale. Yet, even in this case, that of vertebrate

body size evolution, breakpoint regression indicates that the true relationship between time and

evolutionary change involves a complex combination of decreasing and increasing relationships

between time and evolutionary change through the history of vertebrates (30). We further note

that widespread time-independence of evolutionary change does not imply that magnitudes of

evolution are low or "slow". Rather, the vertical axes in Figure 1F-J often correspond to massive

changes in the number of lineages or morphology. The crucial point is that these large

evolutionary changes are typically decoupled from the time interval over which they occurred

and there is no evidence for any systematic time-dependency of accumulated evolutionary

change.

Simulations of classical constant-rate evolutionary models reveal that negative rate-time

associations readily emerge from these existing models. Phenotypic evolution by random

genetic drift results in a Brownian motion process wherein phenotypic variance across lineages

increases linearly with time (Figure 3A), and evolutionary rate is time-independent when

calculated for an entire clade (3E inset panel). Even this simple constant rate model results in a

negative scaling relationship between lineage-specific estimates of evolutionary rate and time

(Figure 3E). The problem is even further exacerbated an under adaptive model of natural

selection towards an optimum (an Ornstein-Uhlenbeck process), where a null relationship

between time and divergence (Figure 3B) generates a strong negative relationship between per-

lineage rate and time (Figure 3F), reminiscent of empirical data from vertebrate body size

evolution (Figure 1).

Finally, the simplest model of lineage accumulation, a constant-rate pure-birth process, also

leads to these types of associations. The log number of lineages accumulates approximately

linearly through time in a pure-birth process, as seen when sampling pure-birth phylogenies

differing in age (Figure 3C, 3D). Yet, pure-birth estimators of diversification rate from these

phylogenies (Kendall-Moran Yule rates, Figure 3G, S1) as well as maximum likelihood

estimates of net diversification from a birth-death diversification model (Figure S2) nonetheless

show a negative scaling relationship with time. In principle Yule rates do not inherently show

time dependence when estimates of zero rates, which rarely result in phylogenies of more than a

single taxon, are included. For example this can be shown by the crown-age estimates using

clade age – time data (Magellon-Sanderson rate estimates, Figure 3H). However, even in this

case non-zero rate estimates show negative time scaling (Figure 3H). Similarly, sampling

directly from the theoretical distribution of pure-birth richness and age yields zero-rate estimates

that can cancel out non-zero rates for young clades (10), we note that these zero rates are largely

unobservable in actual data. Our simulations and others (10) show that non-zero rates show

negative time-scaling even under a constant-rate pure-birth diversification process. This negative

scaling relationships is driven largely by the inclusion of high evolutionary rates estimated over

short timescales (Figure 3G), yet the scaling pattern can sometimes remain even when focusing

only on larger and older clades (Figure S1). This illustrates the inherent problems of interpreting

evolutionary change scaled by time. Moreover, we recover the same general pattern when

simulating phylogenies of varying age under an accelerating time-dependent diversification

process (Figure S3).   These simulation results are not in any way meant as evaluations of these

models, rather a demonstration that constant rate processes readily and almost inevitably will

produce negative rate-time scaling relationships at least in some circumstances.

Discussion

250   Here we have demonstrated three main results with direct implications for interpreting

evolutionary rate-time relationships and their bearing on the pace of life. First, by simply

formalizing the development of the expected relationship between rate and time, we have

demonstrated that there will only exist a limited range of relationships between amount of

evolution and time that generate anything other than negative rate-time relationships. Second,

our analysis of several independent datasets of evolutionary rates demonstrate that existing,

widely-documented negative rate-time scaling relationships clearly suggest weak and

inconsistent relationships between the amount of evolution and timescale of measurement, with

no evidence for a systematic slowdown in accumulated evolutionary change or time-dependency.

Third, using simulation of some of the most common process-based models of constant-rate

evolutionary change, we demonstrate that all readily generate negative rate-time scaling

relationships. Together, these results unequivocally demonstrate the challenge in interpreting

widespread (2, 5, 6, 8) negative rate-time scaling relationships across varying temporal scales.

That evolutionary rates scale negatively with the timeframe over which they are

measured is unsurprising given the inevitable statistical dependence between rate and time.

Negative scaling relationships are therefore expected under a wide range of relationships

between evolutionary change and time, in spite of fundamentally different underlying

evolutionary processes behind the patterns (Figure 2). The fact that the scaling pattern is

observed consistently across disparate types of data from different sources, for rates estimated in

very different ways, and even within generations, strongly suggest that these negative

relationships are unlikely to reflect a causal biological process (2). As Gingerich (5)

emphasized, the negative scaling relationship that is generated by plotting a rate against time

interval must be accounted for before rate variation across time can be meaningfully compared.

Although the likely-artifactual nature of evolutionary rate-time scaling relationships has been

acknowledged since (2, 3, 5, 11), the pervasive negative relationship between evolutionary rate

and time interval continues to inspire evolutionary explanation (2, 6-9, 48).

This renewed interest in rate-time relationships is unfortunate in light of alternative

approaches to the analysis of time-dependence of the evolutionary process. For example,

relationships between the magnitude of evolutionary change (e.g., phenotypic change or the log

number of new lineages arising) and time can be assessed directly, as we have done here and

which has also been done by others(30, 49). For example, piece-wise or segmented regression (3,

28, 30, 50) can identify significant breakpoints in evolutionary change across time, an approach

that avoids the problem of statistical confounders between rate and time interval that was

highlighted by Gingerich (5). These and related statistical approaches (30) can be used to

identify temporal shifts in the evolutionary process over time. It is also likely that some

estimators of evolutionary rate do not show the types of inevitable time scaling relationships we

have reported; for example, clade-level estimates of Brownian motion rate appear to be less

affected. A main conclusion from our analysis, however, is that we should generally be

surprised when evolutionary rates *do not* show negative time scaling.

Our analyses of rate numerators versus time reveal lack of consistent time-dependency

across datasets (Figure 1). We note that these diverse datasets contain information from many

thousands of species of extant and extinct plants, animals, and prokaryotes, range in time interval

from less than one generation to hundreds of millions of years, and together span the entire

history of life. A conclusion from these results, and the finding of log-log rate-time regression

slopes that are shallower than null expectations, is that the persistent rate-time relationship do not

provide evidence of shared features of evolution across these disparate data, *contra* some recent

suggestions (2, 6, 8). In many ways our reanalysis refutes the view that high microevolutionary

rates measured over short time intervals and low macroevolutionary rates measured over longer

time scales indicate fundamental disparity between micro- and macroevolution. By extension,

these results call into question the common view (2, 6, 8) that microevolutionary rates

documented in natural populations of extant organisms (51, 52) are exceptionally high in

comparison to macroevolutionary change over longer time scales (Fig. 1F-J; (see also 30, 53));

our results indicate that such a finding may be a nearly-inevitable outcome of comparing lineage-

specific rates of evolution measured over vastly different time intervals.

Existing data (Fig. 1F-J) suggest that evolutionary change (rate numerator) is remarkably

independent of time, across multiple temporal scales. When significant relationships between

time and amount of evolution do exist, they are weak and variable across datasets.   Such general

and widespread time-independence of evolution becomes a new challenge to explain. Time-

independence of phenotypic evolution be explained by relatively-stable or slowly moving

adaptive zones, within which lineages may evolve substantially (30, 54). For lineage

diversification, disparity in ecological opportunity among diversifying lineages may decouple

clade age and species richness across taxa (34). We emphasize that while we find little

consistent relationships between amount of evolution and interval time, evolutionary events can

still be substantial, as shown in Figure 1. We also note that timing of key events in the history of

life has, of course, played a major role shaping the origins of diversity (55). These include shifts

in body size evolution (30) or rapid jumps in bill evolution in birds (56). Our results are not

inconsistent with such important historical events, but indicate that there is no evidence of a

pervasive relationship between amount of evolution and interval time. There is also evidence of

substantial variation in evolutionary change in response to different selective agents, including

human disturbances and harvesting (46), but such change in response to different environmental

agents does not appear to be systematically linked to time interval over which evolution was

recorded, as we have shown here.

Our results and other recent findings (10, 57) call into question the value of rate-time

scaling patterns as empirical evidence underlying the apparent "paradox of stasis", which is an

observation of far less evolutionary change observed over long timescales than what is expected

based on microevolution (58). For example, consider the widely-appreciated fact that

microevolutionary quantitative genetic models of random genetic drift are inconsistent with the

amount of evolutionary change in deep time (13, 59); we often find far less evolutionary change

in deep time than expected via drift alone, based on patterns of standing extant genetic variance.

However, it is noteworthy that even if phenotypic divergence in deep time was generally

consistent with drift under observed levels of standing variance, we would still expect to see a

negative scaling relationship between lineage specific rates of evolution and time interval (Figure

3). Although it is often claimed that stasis is widespread and a predominant mode of evolution

(12, 13), the results in the present study and others (57, 60, 61) reframe the view that pure stasis

is predominant. Given that the stasis paradox has partly been stimulated by apparent rate

disparities through time, which have been interpreted as indicating a fundamental qualitative

difference between micro- and macroevolution (6, 62, 63), our results instead indicate that the

paradox of stasis could be reformulated as the "paradox of time-independence" of evolutionary

change. Such a reformulation is also consistent with more recent interpretations of the paradox

(30, 57). Our analysis suggests that the decoupling of amount of evolution and time applies not

only to morphological change, but also to lineage accumulation.

Time-invariant evolutionary change still poses other challenges for reconciling micro-

and macroevolution (30). Constancy of evolutionary rates, and thus linear accumulation of

change with time, across temporal scales is expected and often observed at the molecular level

(36, 64). It is more difficult to explain relative time-independence in phenotypic evolution, where

trait change is expected to accumulate with time given strong and variable natural selection (65,

66), or under neutral evolution (67, 68). However, new empirical findings indicate that

phenotypic evolution over a range of timescales is proportional to standing phenotypic and

genetic variation (69). Such variation-limited evolution, if general across taxa, could be one

possible explanation for time-constancy in evolutionary change, although it raises additional

questions on how and why variational properties could be conserved through time. Time-

dependence, or rather lack thereof, therefore remains a major open question in evolutionary

biology.

References

1.   P. D. Gingerich, Rates of evolution on the time scale of the evolutionary process. *Genetica* **112**, 127-144 (2001).

2.   L. J. Harmon *et al.*, Causes and Consequences of Apparent Timescaling Across All Estimated Evolutionary Rates. *Annual Review of Ecology and Systematics* **52**, 587-609 (2021).

3.   A. P. Hendry, M. Kinnison, Perspective: the pace of modern life: measuring rates of contemporary microevolution. *Evolution* **53**, 1637-1653 (1999).

4.   P. D. Gingerich, Quantification and comparison of evolutionary rates. *American Journal of Science* **293**, 453-478 (1993).

5.   P. D. Gingerich, Rates of evolution: Effects of time and temporal scaling. *Science* **222**, 159-161 (1983).

6.   J. Rolland *et al.*, Conceptual and empirical bridges between micro- and macroevolution. *Nature Ecology and Evolution* **7**, 1181-1193 (2023).

7.   S. Y. W. Ho *et al.*, Time dependency of molecular rate estimates and systematic overestimation of recent divergence times. *Molecular Ecology* **20**, 3087-3101 (2011).

8.   L. F. Henao Diaz, L. J. Harmon, M. T. C. Sugawara, M. W. Pennell, Macroevolutionary diversification rates show time dependency. *Proceedings of the National Academy of Sciences* **116**, 7403-7408 (2019).

9.   S. Louca, L. Francisco Henao-Diaz, M. W. Pennell, The scaling of diversification rates with age is likely explained by sampling bias. *Evolution* **76**, 1625-1637 (2022).

10.   B. C. O'Meara, J. M. Beaulieu, Error causes a perceived pull to the present in evolutionary rates. *BioRxiv* **https://doi.org/10.1101/2024.02.29.582777** (2024).

11.   S. J. Gould, Smooth curve of evolutionary rate: A psychological and mathematical artifact. *Science* **226**, 994-996 (1984).

12.   N. Eldredge, S. J. Gould, "Punctuated equilibria: an alternative to phyletic gradualism" in Models in Paleobiology*, T. J. M. Schopf, Ed. (Freeman Cooper, Sanfrancisco, 1972).

13.   T. F. Hansen, D. Houle, "Evolvability, stabilizing selection, and the problem of stasis" in Phenotypic Integration: Studying the Ecology and Evolution of Complex Phenotypes*, M. Pigliucci, K. Preston, Eds. (Oxford University Press, 2004).

14.   S. J. Gould, N. Eldredge, Punctuated equilibria: the tempo and mode of evolution reconsidered. *Paleobiology* **3**, 115-151 (1977).

15.   S. J. Gould, N. Eldredge, Punctuated equilibrium comes of age. *Nature* **366**, 223-227 (1993).

16. K. Pearson, Mathematical contributions to the theory of evolution.—On a form of
spurious correlation which may arise when indices are used in the measurement of
organs. *Proceedings of the Royal Society of London* **60**, 489-498 (1897).

17. Y. T. Prarie, D. F. Bird, Some misconceptionsabout the spurious correlation problem in
the ecological literature. *Oecologia* **81**, 285-288 (1989).

18. S. J. Arnold, Phenotypic evolution: the ongoing synthesis. *The American Naturalist* **183**,
729-746 (2014).

19. W. R. Atchley, C. T. Gaskins, D. Anderson, Statistical Properties of Ratios. I. Empirical
Results. *Systematic Zoology* **25**, 137-138 (1976).

20. D. A. Jackson, K. M. Somers, The spectre of 'spurous' correlations. *Oecologia* **86**, 147-
151 (1991).

21. J. B. S. Haldane, Suggestions as to Quantitative Measurement of Rates of Evolution.
Evolution **3**, 51-56 (1949).

22. M. Foote, Origination and extinction components of taxonomic diversity:General
problems. *Paleobiology* **26**, 74-102 (2000).

23. S. Magallon, M. J. Sanderson, Absolute diversification rates in Angiosperm clades.
Evolution **55**, 1762-1780 (2001).

24. S. Nee, R. M. May, P. H. Harvey, The reconstructed evolutionary process. *Philosophical*
Transactions of the Royal Society of London Series B-Biological Sciences **344**, 305-311
(1994).

25. D. L. Rabosky *et al.*, BAMMtools: an R package for the analysis of evolutionary
dynamics on phylogenetic trees. *Methods in Ecology and Evolution* **5**, 701-707 (2014).

26. R. G. FitzJohn, Diversitree: comparative phylogenetic analyses of diversification in R.
Methods in Ecology and Evolution **3**, 1084-1092 (2012).

27. B. C. Kenney, Beware of Spurious Self-Correlations! *Water Resources Research* **18**,
1041-1048 (1982).

28. W. Schlager, D. Marsal, P. A. G. van der Geest, A. Sprenger, Sedimentaion rates,
obersvation span, and the problem of spurious correlation. *Mathematical Geology* **30**,
547-556 (1998).

29. H. D. Sheets, C. E. Mitchell, Uncorrelated change produces the apparent dependence of
evolutionary rate on interval. *Paleobiology* **27**, 429-445 (2001).

30. J. C. Uyeda, T. F. Hansen, S. J. Arnold, J. Pienaar, The million year wait for
macroevolutionary bursts. *Proceedings of the National Academy of Sciences of the*
United States of America **108**, 15908-15913 (2011).

31. S. Estes, S. J. Arnold, Resolving the paradox of stasis: models with stabalizing selection
explain evolutionary divergence on all timescales. *American Naturalist* **169**, 227-244
(2007).

32. P. D. Roopnarine, Analysis of rates of morphologic evolution. *Annual Review of Ecology*
and Systematics **34**, 605-632 (2003).

33. D. L. Rabosky, G. J. Slater, M. E. Alfaro, Clade age and species richness are decoupled
across the Eukaryotic tree of life. *PLoS Biology* **10**, e1001381 (2012).

34. D. L. Rabosky, Ecological limits and diversification rate: alternative paradigms to explain
the variation in species richness among clades and regions. *Ecology Letters* **12**, 735-743
(2009).

35. J. J. Wiens, The causes of species richness patterns across space, time, and clades and the
role of "ecological limits.". *Quarterly Review of Biology* **86**, 75-96 (2011).

36. G. J. Morgan, Emile Zuckerlandl, Linus Pauling, and the Molecular Evolutionary Clock,
1959-1965. *Journal of the History of Biology* **31**, 155-178 (1998).

37. E. Paradis, Random phylogenies and the distribution of branching times. *Journal of Theoretical Biology* **387**, 39-45 (2015).

38. S. Louca, M. W. Pennell, Extant timetrees are consistent with a myriad of diversification histories. *Nature* **580**, 502-505 (2020).

39. B. C. Emerson, N. Kolm, Species diversity can drive speciation. *Nature* **434**, 1015-1017 (2005).

40. M. Chouteau, M. Arias, M. Jorun, Warning signals are under postive frequency-dependent selection in nature. *Proceedings of the National Academy of Sciences* **113**, 2164-2169 (2016).

41. R. A. Fisher, *The genetical theory of natural selection* (The Clarendon press, Oxford, 1930), pp. xiv, 272 p.

42. R. Lande, Models of speciation by sexual selection on polygenic traits. *Proceedings of the National Academy of Sciences of the United States of America-Biological Sciences* **78**, 3721-3725 (1981).

43. M. Kirkpatrick, Sexual selection and the evolution of female choice. *Evolution* **36**, 1-12 (1982).

44. P. Nosil, J. L. Feder, S. M. Flazman, Z. Gompert, Tipping points in the dynamics of speciation. *Nature Ecology and Evolution* **1**, 0001 (2017).

45. M. Scheffer, *Critical Transitions in Nature and Society*, Princeton Series in Complexity (Princeton University Press, Princeton, N.J., 2009).

46. S. Sanderson *et al.*, The pace of modern life, revisisted. *Molecular Ecology* **31**, 1028-1043 (2021).

47. J. P. Scholl, J. J. Wiens, Diversification rates and species richness across the tree of life. *Proceedings B* **283**, 20161334 (2016).

48. M. T. Kinnison, A. P. Hendry, The pace of modern life II: from rates of contemporary microevolution to pattern and process. *Genetica* **112**, 145-164 (2001).

49. E. I. Svensson, Eco-evolutionary dynamics of sexual selection and sexual conflict. *Functional Ecology* **33**, 60-72 (2019).

50. M. E. Clapham, J. A. Karr, Environmental and biotic controls on the evolutionary history of insect body size. *Proceedings of the National Academy of Sciences* **109**, 10927-10930 (2012).

51. J. B. Losos, K. I. Warheitt, T. W. Schoener, Adaptive differentiation following experimental island colonization in Anolis lizards. *Nature* **387**, 70-73 (1997).

52. D. Reznick, F. H. Shaw, F. H. Rodd, R. G. Shaw, Evaluation of the Rate of Evolution in Natural Populations of Guppies (Poecilia reticulata). *Science* **275**, 1934-1937 (1997).

53. E. I. Svensson, The speed of life-history evolution. *Trends in Ecology and Evolution* **12**, 380-381 (1997).

54. G. G. Simpson, *The Major Features of Evolution* (Columbia University Press, New York, 1953).

55. S. J. Gould, *Wonderful Life: The Burgess Shale and the Nature of History* (W.W. Norton & Co., 1989).

56. C. R. Cooney *et al.*, Mega-evolutionary dynamics of the adaptive radiation of birds. *Nature* **542**, 344-347 (2017).

57. K. L. Voje, Tempo does not correlate with mode in the fossil record. *Evolution* **70**, 2678-2689 (2016).

58. T. F. Hansen, D. Houle, "Evolvability, stabilizing selection, and the problem of stasis" in The Evolutionary Biology of Complex Phenotypes*, M. Pigliucci, K. Preston, Eds. (Oxford University Press, 2004).

485   59.   D. Houle, G. H. Bolstad, K. van der Linde, T. F. Hansen, Mutation predicts 40 million
486   years of fly wing evolution. *Nature* **548**, 447-450 (2017).
487   60.   K. L. Voje, J. Starrfelt, L. H. Liow, Model adequacy and microevolutionary explanations
488   for stasis in the fossil record. *The American Naturalist* **191**, 509-523 (2018).
489   61.   G. Hunt, The relative importance of directional change, random walks, and stasis in the
490   evolution of fossil lineages. *Proceedings of the National Academy of Sciences of the*
491   *United States of America* **104**, 18404-18408 (2007).
492   62.   N. Eldredge *et al.*, The dynamics of evolutionary stasis. *Paleobiology* **31**, 36-55 (2005).
493   63.   D. J. Futuyma, Evolutionary constraint and ecological consequences. *Evolution* **64**, 1865-
494   1884 (2010).
495   64.   L. Bromham, D. Penny, The modern molecular clock. *Nature Reviews Genetics* **4**, 216-
496   224 (2003).
497   65.   J. G. Kingsolver *et al.*, The strength of phenotypic selection in natural populations.
498   *American Naturalist* **157**, 245-261 (2001).
499   66.   J. Felsenstein, Phylogenies and quantitative characters. *Annual Review of Ecology and*
500   *Systematics* **19**, 445-471 (1988).
501   67.   M. Lynch, The rate of morphological evolution in mammals from the standpoint of the
502   neutral expectation. *The American Naturalist* **136**, 727-741 (1990).
503   68.   J. Felsenstein, Phylogenies and the comparative method. *American Naturalist* **125**, 1-15
504   (1985).
505   69.   T. F. Hansen, C. Pélabon, Evolvability: A quantitative genetics perspective. *Annual*
506   *Review of Ecology and Systematics* **52**, 153-175 (2021).
507   70.   R. Lande, Natural selection and random genetic drift in phenotypic evolution. *Evolution*
508   **30**, 314-334 (1976).
509   71.   S. Nee, Inferring speciation rates from phylogenies. *Evolution* **55**, 661-668 (2001).
510   72.   B. G. Baldwin, M. J. Sanderson, Age and rate of diversification of the Hawaiian
511   silversword alliance (Compositae). *Proceedings of the National Academy of Sciences* **95**,
512   9402-9406 (1998).
513   73.   L. J. Revell, Phytools: an R package for phylogenetic comparative biology (and other
514   things). *Methods in Ecology and Evolution* **3**, 217-223 (2012).
515   74.   S. Hoehna, Fast simulation of reconstructed phylogenies under global time-dependent
516   birth-death processes. *Biorinformatics* **29**, 1367-1374 (2013).

Fig. 1. Evolutionary rate-time relationships reflect null or variable underlying relationships between evolutionary change and time. Panels **A-E** show evolutionary rate plotted against the timescale of measurement (6), with dashed red lines showing the least squares fit of the function $rate = a*time^b$. Panels **F-J** plot the amount of evolutionary change (i.e., a rate numerator) against time frame for the same datasets, with dashed red lines showing the linear piecewise regression, if statistically significant. Panels **A** and **F** show vertebrate body size evolution (30), Panels **B** and **G** show diversification of genera from fossil lineages (plants, mammals, and marine animals)(8), Panels **C** and **H** show diversification data estimated from time-calibrated molecular phylogenies (8), while Panels **D** and **I** show a larger database of phylogenetic diversification rates (47) where rates were inferred directly from age and richness data (23). Panels **E** and **J** show within-generation morphological changes (mainly from fish harvesting). All nonlinear negative relationships plotted in **A-E** were statistically significant, whereas relationships in **F-J** were variable, with statistically significant breakpoints found for morphology, fossil lineage accumulation, and one estimator of phylogenetic diversification, and a positive linear relationship for another estimator of phylogenetic diversification. No significant relationship was found for the within-generation changes in panel **J**. Sample sizes for each column: **A** N = 5392

records of evolutionary change, **B** N = 90 lineages, **C** N = 114 clades, **D** N = 434 Orders, **E** N = 302 records of within-generation change.

Fig. 2. Extraordinary relationships between the accumulated amount of evolution and time are required to produce positive rate-time relationships. Panel **A** shows the relationship between rate and time for various functions of the rate numerator and time, which are plotted in the same color in Panel **B**. The black line is a reference point, where there is a null relationship between rate and time. Thus, each line in Panel **A** corresponds to a specific relationship between rate numerator and time, plotted in Panel **B**; only cases of an accelerating relationship between amount of evolution and time are expected to produce positive rate-time scaling relationships. We assume for this example a form of the relationship, *amount of evolution = time^b*, where the function was plotted for b ranging from -2 (blue) to 2 (red). Similar general patterns are observed for other functions of change and time. Note that because we are considering these relationships in light of large amalgamations of data from disparate groups (e.g., Figure 1), as opposed to within-lineage time series, negative relationships between amount of evolution and interval time are in principle possible, even if unlikely.

Fig. 3. Simulations of constant-rate evolutionary models reveal persistence of spurious negative rate-time relationships. Panels **A** and **E** show a simulation of phenotypic evolution under random genetic drift in the absence of natural selection, which produces a Brownian motion process (66) where variance among lineages accumulates linearly with time (**A**). Although the rate of drift is constant (**E** inset), we will nevertheless obtain a negative scaling relationship between the lineage-specific per-generation rate of evolution (**E**). Panels **B** and **F** show a simulation of phenotypic evolution under drift and stabilizing natural selection, an Ornstein Uhlenbeck process (70). This process generates no relationship between time and divergence (**B**), but again a strong negative scaling relationship with per-lineage per-generation evolutionary rate and time and hence a similar negative relationship (**F**). Panels **C** and **G** show a constant-rate pure-birth (i.e., no extinction) lineage diversification process (Yule process), where diversity accumulates approximately linearly with time (**C**). This scenario also produces a similar negative scaling relationship between the Kendall-Moran branch-length based rate estimator and time (**G**). Calculating Yule diversification rate for a different set of trees (**D**) based on the crown-age Magellon and Sanderson estimator yields a similar pattern (**H**), but in this case the inclusion of zero rate estimates (which are unlikely to be sampled in actual trees or in the simulated data in panel C) offsets (green line) the negative relationship between non-zero

observable rates and time (red line). The last 15 million years of the simulated data illustrates there is no relationship between rate and time for this time interval (**H** inset).

Methods

Data and code availability: All data is already publicly available, and we provide our

complete R script as a supplemental file.

Data. We analyze existing data on rates of vertebrate body size evolution (30), which consists of

data on log-fold change in body size as well as time interval over which this change occurred.

For phylogenetic lineage diversification, we analyze two datasets, one in which diversification

rates were estimated from the branching pattern of species-level phylogenies (8), and another

larger dataset (47) where diversification rates were estimated from clade diversity and age

information. For fossil data, we analyze an existing dataset (8) of emergence rates of new

genera. For each dataset, we plot the standard measure of rate against the timeframe over which

that rate was measured, fitting exponential linear models by nonlinear least squares (6). We also

plot a measure of the rate numerator against time for each of these datasets, and fit piecewise

linear regressions to identify significant linear relationships between amount of evolution and

time, as well as any potential shifts in the relationship between evolution and time. For each

dataset, we fit nonlinear models of the form $rate = a*time^b$ using nonlinear least squares (nls())

in R, as well as piecewise regression using the segmented package. We compared models fits of

an ordinary linear regression to a model with a single breakpoint, and also when appropriate to a

null model containing only an intercept, using the anova() function in R. Data presented in

Figure 1 were obtained from ref 11 (available at

https://github.com/fhenaod/evol_time_scaling/tree/main/data), ref 29 (Dryad

doi:10.5061/dryad.7d580), ref 46, and ref 47 (database available at:

https://proceeddatabase.weebly.com/).

<u>Simulation</u> We simulated three different constant rate evolutionary process. The aim of our

simulations was not to evaluate the performance of statistical models used to estimate rates.

Rather, our aim was simple to examine scaling patterns observed in the simplest cases of

simulated constant rate evolution. For morphological evolution, we simulated random genetic

drift for 1-1000 generations assuming a heritability of 0.50 and a population size of 100. We also

simulated evolution under genetic drift and stabilizing selection, with the same demographic and

genetic assumptions but assuming weak selection ($\omega^2 = 15$) towards an optimum. Each

simulation was started with the population at its optimum. For morphological evolution, we

plotted rates in units of Haldanes (4), which was natural as we simulated evolution on the scale

of phenotypic standard deviations.   For lineage diversification, we simulated pure-birth

phylogenies with a constant speciation rate of 0.15, corresponding to a expected number of 2.3

taxa at timepoint 1 under a continuous time pure birth process. We simulated pure-birth

diversification because, while biologically unrealistic, it excludes explanations for rate-time

scaling patterns that are based on sampling biases generated by extinction. We also note that

although many challenges exist in sampling all possible trees in a simulation, these are identical

to challenges observed in actual data; namely, inclusion of trees with zero diversification rate.

Given the aims of our simulations noted above, we proceeded by first simulating trees

conditioned on number of tips as opposed to clade age, focusing first on trees with at least 3 tip

taxa (conclusions remain unchanged by focusing instead on trees with 20 or more tips; Figure

S3). We then calculated the Kendall-Moran estimator (71, 72) for the Yule process for these

trees using the diversitree package (26). We also estimated maximum likelihood birth-death

diversification rates for the same set of trees constraining diversification rates to be positive.

Finally, we also generated trees with ages ranging from 1-35 million years, and plot the

maximum likelihood estimator of the crown-age Yule diversification rate, $[\log(N)\text{-}\log(2)]/t$ (23).

Our simulations of phenotypic evolution were performed using nested for() loops in R and

parameterized as described above. Our simulations of constant rate pure birth phylogenetic

diversification was performed using the pbtree() function in the phytools package (73). Our

simulations of time-dependent pure birth diversification (Figure S3) were performed using the

TESS package (74). We provide our complete R script as supplemental material.

Acknowledgments: We thank Masahito Tsuboi and Josef Uyeda for critical feedback that

improved the manuscript, as well as Chris Cooney, Charlie Cornwallis, Kjetil Voje, and

Miguel Gomez for comments and discussion of these issues. **Funding:** This work was

funded by grants from the Swedish Research Council (VR: grants no. 2019-03706 to SPD

and no.r 2020-03123 to EIS), The Crafoord Foundation (20220602), Formas (grant 2021-

01096), to SPD

Author contributions: SPD and EIS conceived of the problem and designed the study, SPD

performed the analysis and drafted the manuscript. Both authors wrote the manuscript.

Competing Interest Declaration: The Authors have no competing interests to declare

Fig. S1. Negative time-scaling of ML lineage diversification rates for trees greater than or equal to 20 tips. Shown are the scaling relationships for time and diversification estimators for pure-birth phylogenies of size equal to or greater than N = 20. Panel A shows the net diversification rate from a ML birth-death model, panel B shows the Yule estimator for the same set of trees. Other simulation conditions were as in Figure 3C, G.

Fig S2. Birth-death estimator diversification rate for trees plotted in Figure 3D, H. Shown

is the ML estimate of the birth-death diversification rate λ for the same set of trees shown in

Figure 3D, H. Red dashed line shows the least squares fit of the model *rate = a*time^b*.

Fig. S3. Accelerating time-dependent diversification produces negative rate scaling

patterns. Panels show simulations of time-dependent pure birth diversification, repeated for

different timeframes from 1-35 million years under the same time-dependent diversification

model. In each simulation birth rates were assumed to be linearly related to time, $\lambda \sim 0.008*time$,

extinction rate $\mu = 0$. Panel **A** shows log of total diversity plotted against time for each

simulation run. Panel **B** shows the estimate of net speciation rate $\log(N)/t$ as a function of time.

730   Qualitative conclusions remain unchanged under different parameter values, with the one salient

731   feature being that the expected positive rate-time relationship is only recovered in deep time.