Title: **Revisiting evolutionary rate-time relationships**

Abstract

Rates of molecular, phenotypic, and lineage diversification typically scale negatively with time interval of measurement, raising longstanding questions about time-dependency of evolutionary processes. These patterns and their potential meaning have recently re-entered evolutionary discussions. In this Perspective we revisit the general challenges in interpreting rate-time relationships. Much apparent temporal scaling of evolutionary rate is an inescapable outcome of plotting a ratio against its denominator, either directly or indirectly. Highly unlikely relationships between timescale and accumulated evolutionary change are required to produce anything other than negative rate-time relationships. Simulations reveal that constant rate evolutionary processes readily generate negative rate-time scaling relationships under many conditions, and that a range of rate-time scaling exponents can be generated by similar evolutionary processes. Reanalysis of six empirical datasets reveals unscaled magnitudes of evolution that are either unrelated to time and/or vary in their relationship with time, with over 99% of variation in rate-time relationships across six datasets explained by time variation alone. We further evaluated a recent hypothesis that evolutionary rate-time scaling reflects three modes of change, from micro- to macroevolutionary time scales using break-point regression, but we found no strong support for this hypothesis. Taken together, negative rate-time relationships are largely inevitable and difficult to interpret. In contrast, it is more straightforward to assess how evolutionary change accumulates with time.

Main Text: Estimates of evolutionary rate vary with the time interval over which they are measured. A general pattern with faster evolutionary rate over short timescales, and slower rates over longer timescales has been recognized for decades (Gingerich 1983, Gingerich 1993, Hendry and Kinnison 1999, Gingerich 2001, Harmon, Pennell et al. 2021). Such time-dependency has been documented for nearly every possible measure of evolutionary rate (Rolland, Francisco Henao-Diaz et al. 2023), including rates of morphological and molecular evolution (Alba and Castresana 2005, Ho, Shapiro et al. 2007, Ho, Lanfear et al. 2011), and rates of lineage formation on phylogenies (Ricklefs 2006, Scholl and Wiens 2016) and in the fossil record (Foote 1994, Foote 2000, Henao Diaz, Harmon et al. 2019) (Figure 1). This puzzling and general phenomenon of widespread negative rate-time scaling relationships was recently highlighted as a major unsolved challenge in evolutionary biology, with potential explanation based on shared time-dependent features of evolution (Henao Diaz, Harmon et al. 2019, Harmon, Pennell et al. 2021, Rolland, Francisco Henao-Diaz et al. 2023). Recent technical and statistical explanations for negative rate-time scaling relationships move beyond simple arguments of artefact and focus on sampling biases (Emerson 2007, Louca, Francisco Henao-Diaz et al. 2022, Reijenga and Close 2025), such as a the "push of the past" (Rolland, Francisco Henao-Diaz et al. 2023), model mis-specification (Harmon, Pennell et al. 2021), and measurement error (O'Meara and Beaulieu 2024). The renewed interest in time-dependency of evolutionary rates has rekindled long-standing and controversial ideas about fundamental disparities between micro- and macroevolution (Eldredge and Gould 1972, Gould and Eldredge 1977, Charlesworth, Lande et al. 1982, Gould 1984, Gould and Eldredge 1993, Hansen and Houle 2004).

Interpreting the causes and meaning of rate-time scaling relationships is difficult in light of the inevitable negative correlation generated by plotting a ratio against its own denominator (Pearson 1897, Atchley, Gaskins et al. 1976, Gingerich 1983, Prarie and Bird 1989, Jackson and

Somers 1991, Arnold 2014).   Despite widespread appreciation of this statistical issue, (e.g., see

Box 1 in Gingerich 1983, Gould 1984, Harmon, Pennell et al. 2021) a revisit to the interpretative

challenges of general rate-time relationships is timely and necessary in light of renewed interest

in negative rate-time scaling. For example, in a high-profile review on bridging micro and

macroevolution, Rolland et al (Rolland, Francisco Henao-Diaz et al. 2023) presented five

negative rate-time scaling relationships as evidence of a divide between evolution over short

versus deep time.

In this Perspective, we analyze a simple model of rate-time relationship that is purposefully

agnostic to any specific underlying evolutionary process. We illustrate how evolutionary change,

time, and their (co)variation contribute to generating nearly inevitable rate-time relationships.

We also re-analyzed six classical datasets that have been used in past work to demonstrate the

universality of rate-time scaling. A central conclusion of this Perspective is that pervasive

negative rate-time relationships say little on their own about the correspondence of evolutionary

processes across disparate timescales.

I.   Background: Negative rate-time relationships are nearly inevitable

Evolutionary rate can be conceptualized as a function of evolutionary change scaled by time

interval:

$$y = d/t \qquad (1)$$

where y is the rate-measure, d is some measure of evolutionary change (e.g., change in

population mean phenotype, base pair substitutions, log change in number of species in a clade)

and t is some measure of time interval over which change was observed or inferred (e.g.

generations, or years) and noting that rates are undefined at $t = 0$. Note that we use "time" and

"time interval" relatively interchangeably, although it is important to realize many archetypical

datasets of rate estimates contain time intervals sampled from different lineages (see below).

Some classical estimates of evolutionary rate for morphological traits are calculated directly by

dividing phenotypic change with time, the latter typically measured in generations or years. For

example, darwins (Haldane 1949) $\left(\frac{\ln{(\bar{z}_1/\bar{z}_2)}}{\Delta t * 10^6}\right)$, where t is units of years, and haldanes (Gingerich

1993) $\left(\frac{(\bar{z}_1 - \bar{z}_2)}{\Delta t}\right)$, where z is units of pooled within-population standard deviation and t is units of

generations, have been used to describe morphological rate, and other types of rate estimators

measure per-lineage origination rate of fossil lineages (Foote 2000) and phylogenetic net

diversification rate (Magallon and Sanderson 2001). Other rate estimates are obtained from more

sophisticated statistical models of evolution (Nee, May et al. 1994, Nee 2001, FitzJohn 2012,

Rabosky, Grundler et al. 2014). Regardless, all evolutionary rate measures are, per definition,

scaled by time (e.g., waiting times between observed branching events on a phylogeny are a

direct component of the likelihood in maximum likelihood models of birth-death diversification

(Nee, May et al. 1994)). Thus, the conceptualization of evolutionary rate in equation 1 is general,

if not precise.

The inevitable mathematical dependence between rate and time depicted in equation 1 is

an obvious challenge in interpreting the time-dependency of evolutionary rate, and has been

widely appreciated in evolutionary biology, both for rates of morphological and molecular

evolution (Gingerich 1983, Hendry and Kinnison 1999, Roopnarine 2003, Estes and Arnold

2007, Uyeda, Hansen et al. 2011, Arnold 2014) and lineage diversification (Rabosky 2009,

Rabosky 2010, Wiens 2011, Rabosky, Slater et al. 2012, Rabosky 2016, Rabosky and Benson

2021). There has also been considerable attention given to related issues outside of or tangential

to evolutionary biology, including geology (Sadler 1981, Schlager, Marsal et al. 1998),

116   paleontology (Foote 1994, Sheets and Mitchell 2001) ecology (Kenney 1982) and conservation

117   (Wiens and Saban 2025).

118   When considering the role for ratio effects in contributing to our expectation for a

119   preponderance of negative rate-time relationships, it is useful to ask: under what scenarios do we

120   expect null or positive relationships between rate and time? Consider rate as a function of time,

$$y = f(t) \tag{2}$$

122   Where y is evolutionary rate (as in eqn 1). We also consider the amount, or magnitude, of

123   accumulated evolution (rate numerator) as a function of time,

$$d = g(t) \tag{3}$$

where d is the accumulated evolutionary change. Note that in most analyses that assess rate-time

relationships, point estimates of rates from different lineages spanning different time intervals

are combined in a single analysis, and in this case the function $g(t)$ need not be strictly

increasing even for absolute values of morphological change. Rates, $y = d/t$, will be constant

over time when d scales linearly with t, i.e. $g'(t)$ is a positive constant over the timescale in

question, with prime denoting the first derivative. Rates will increase with time, i.e. $f'(t) > 0$

when $g''(t) > 0$; that is, rates increase over time when the amount of evolutionary change is an

accelerating function of time. Noteworthy is that under a null relationship between d and t, such

that $g'(t) = 0$, the logarithmic regression of rate versus time ($\log(y) \sim \mu + \beta*\log(t)$) will yield a

negative slope estimate of $\beta = -1$ (Gingerich 1983). This can be seen most easily by expressing

equation 3 as a power function, e.g., $d = t^0$, in which case we can describe the regression of

$\log(y) \sim \beta*\log(t)$ as $\log(t^0/t) \sim \beta*\log(t)$; solving for $\beta = \log(t^{-1})/\log(t) = -1$. This negative slope

estimate provides one cutoff of interest in assessing rate-time scaling relationships: log-log

regressions of rate vs time that are shallower than or not significantly different from -1 are

consistent with either a null relationship between evolutionary change and time (time

independence) or an increasing relationship between accumulated change and time, respectively.

The relationships between equations 1-3 are illustrated in Figure 2, which shows the

relationship between rate and time. Figure 2A illustrate differing relationships between rate

numerator and time illustrated in Figure 2B for a range of power functions. There are few

theoretical models that predict the type of accelerating relationship between magnitude of

evolution and time that are needed to produce a positive relationship between evolutionary rate

and time. Neutral and adaptive models of phenotypic and molecular evolution predict either no

relationship with time or linear accumulation (constant change) with time (Morgan 1998).

Models based on adaptive landscape theory predict a slowdown in evolutionary rates for

morphological traits, but a recent study of thousands and extinct and extant species did not find

any empirical support for such a slowdown. i. e. no evidence for time-dependent evolutionary

rates (Quintero 2025). Similarly, birth-death models of lineage accumulation also predict either

a linear or a decelerating accumulation of log lineage number through time under most

conditions. Even models of time-dependent lineage diversification make no prediction about

accelerating log lineage number versus interval time across a sample of clades. Moreover,

lineage-through-time (LLT) plots for single clades diversifying under such a process can produce

complex patterns (Paradis 2015) that are consistent with several different underlying

diversification processes (Louca and Pennell 2020).

Some scenarios in nature could potentially lead to accelerating change with time and thus

generate positive relationships between evolutionary rates and time (Figure 2). Such scenarios

include "diversity begets diversity" models of lineage diversification (Nee, May et al. 1994,

Emerson and Kolm 2005), evolution by positive frequency-dependent selection (Chouteau,

Arias et al. 2016), runaway sexual selection driven by female choice (Fisher 1930, Lande 1981,

Kirkpatrick 1982), "tipping points" in the evolution of reproductive isolation (Nosil, Feder et al.

2017) and populations crossing critical thresholds followed by phase transitions between

alternative stable states (Scheffer 2009). Tracking of a constant-rate moving optimum can

generate short-term positive rate-time scaling until a population reaches its stable expected lag

(Lynch and Lande 1993, Bürger and Lynch 1995) (elaborated below). How common such

scenarios are in nature is an open empirical question. However, given the transient nature of self-

reinforcing evolutionary processes driven by positive feedbacks (Lande 1981, Kirkpatrick 1982),

we suspect that positive rate-time relationships are unlikely to be sustained over longer

macroevolutionary time scales and should quickly reach stable equilibria and thus be largely

transient. Rapid and self-reinforcing evolutionary processes are unlikely to proceed unchecked

over millennia (Scheffer 2009), as would be required to generate positive rate-time relationships

in empirical data such as that shown in Figure 1.

We can decompose the covariance between rate and time, focusing on the log-scaled

case, where

$$Cov\left[log\left(\frac{d}{t}\right), \log\left(t\right)\right] = Cov[log(d) - \log\left(t\right), \log\left(t\right)]$$

$$= Cov[log(d), \log\left(t\right)] - Var[\log(t)] \tag{4}$$

Which shows that the relationship between rate and time is governed both by the variance in time

interval as well as the relationship between accumulated change (*d*) and time. Importantly, the

variance partitioning in equation 4 makes it possible to evaluate the contribution of change-time

relationships in explaining rate-time relationships. If rate-time relationships generally carry

meaningful information about how evolutionary change accumulates with time, we expect

$Cov[log(d), \log\left(t\right)]$ to be an important contributor to explaining variation in

$Cov\left[log\left(\frac{d}{t}\right), \log\left(t\right)\right]$. We can transform the covariance to a regression coefficient by

multiplication by $Var[\log(t)]^{-1}$, which results in $\beta_{y,t} = \beta_{d,t} - 1$, and thus further illustrating

that if there is no relationship between accumulated change d and time (such that $\beta_{d,t} = 0$) then

the expected log-log regression of rate and time, $\beta_{y,t}$, is equal to -1.

Simulations

Simulations of classical constant-rate evolutionary models reveal that negative rate-time

associations readily emerge from these existing models. Phenotypic evolution by random

genetic drift results in a Brownian motion process wherein phenotypic variance across lineages

increases linearly with time (Figure 3A), and estimated evolutionary rate is time-independent

when calculated for an entire clade (3E inset panel). Even this simple constant rate model results

in a negative scaling relationship between lineage-specific estimates of evolutionary rate and

time (Figure 3E), resulting in a scaling exponent of – ½ in a log-log regression of rate versus

time (Hansen 2024).

Lineage specific rates are even more strongly associated with time under an adaptive

model of natural selection towards a stationary optimum (an Ornstein-Uhlenbeck, or OU,

process), where a null relationship between time and divergence (Figure 3B) generates a strong

negative relationship between per-lineage rate and time (Figure 3F) with a log-log slope of -1

over long timescales (Hansen 2024), reminiscent of empirical data from vertebrate body size

evolution (Figure 1). However, we note that a stationary OU model readily produces negative

rate-time scaling exponents that deviate from -1 when sampling lineages evolving under different

values of genetic variance and strength of stabilizing selection (Figure 4). Adding any

movement of the optimum in an OU process produces dramatic shifts in the form of rate time

scaling yet also indicates that a range of models incorporating stabilizing selection readily

produce scaling relationships similar or identical to that expected under pure genetic drift (Figure 4).

Well established constant-rate models of lineage accumulation can also lead to the appearance of negative associations between rate and time when sampling is incomplete, due to ratio effects. In principle Yule rates do not inherently show time dependence when estimates of zero rates, which rarely result in phylogenies of more than a single taxon, are included, yet such species-depauperate clades are rarely included in actual data, and non-zero rates show negative time-scaling even under a constant-rate pure-birth diversification process (O'Meara and Beaulieu 2024). Of course, the Yule model is not a realistic description of biological diversification, since extinction has clearly played a major role in the history of life. Yet such a combination of sampling bias and ratio effects also generate apparent negative rate-time relationships under birth-death processes, which remain to some degree even after conditioning the likelihood of the birth-death model on known sampling biases (Louca, Francisco Henao-Diaz et al. 2022).

II.   **Empirical examples reveal variable patterns of accumulated change despite pervasive negative rate-time scaling**

We revisited and reanalyzed six published empirical datasets of evolutionary rate estimates that have featured centrally in recent discussions of rate-time scaling relationships (Scholl and Wiens 2016, Svensson 2019, Sanderson, Beausoleil et al. 2021, Rolland, Francisco Henao-Diaz et al. 2023). We first perform regression analyses to calculate rate-time and rate numerator-time scaling relationships, the former largely recapitulating the analyses of the original studies, the latter of which is more rarely presented. We then performed the novel

variance decomposition presented in equation 4, which essentially represents a metanalysis of

these rate estimators.

Vertebrate body sizes: evaluating Hansen's (2024) "three modes" of evolution

For our first example, vertebrate body size evolution (Figure 1A, G), past work (Estes

and Arnold 2007, Uyeda, Hansen et al. 2011, Arnold 2014) has shown that body size changes are

largely unrelated to time, and in time-series of extant populations there was no detectable change

(Sanderson, Beausoleil et al. 2021). Although a log-log regression of rate vs time yields a slope

that is nominally steeper than -1 (β = -1.019, 95% CI -1.031, -1.007), which suggests no

accumulation of change with interval time, there is a significant breakpoint in the relationship

between morphological change and time ($F_{2, 5388}$ = 207.2, $P < 2.2$ x 10^{-6}, breakpoint = -2.63 log

MY (0.072 MY), SE = 0.147). The result is first a weak negative relationship between time and

evolutionary change, followed by an increasing slope after the breakpoint (Fig. 1A, G; or

shallower negative slope on the log scale, Figure 1A inset panel), identical to that found by

Uyeda et al (2011). Thus, for body size evolution, amount of change is largely unrelated to time

interval except over the deepest periods of evolutionary time (Uyeda, Hansen et al. 2011).

Quintero (2025) reached a similar conclusion in a recent compilation of rates of morphological

evolution among thousands of extinct and extinct species and no evidence for time-dependency.

Next, we applied breakpoint regression models to the log-log relationship between body

size evolutionary rate and time to test Hansen's (Hansen 2024) recently proposed "three modes"

of evolution. Hansen (2024) argued that rate-time scaling may be near null over short

(microevolutionary) time intervals, approach -1 over intermediate time intervals, and relax to

around -1/2 over deep macroevolutionary time intervals. To evaluate the evidence for this

hypothesis we fit progressively more complex breakpoint regression models of log absolute

darwins versus log time interval (Fig. 5). There was statistical support (significantly reduced

AIC) for increasing complexity up to seven breakpoints (Figure 5, Table S1), with ΔAIC of over

1200 between the best fit model and one with only two breakpoints (i.e. three modes). Further,

examination of the scaling slopes from a model with only two breakpoints reveals scaling slopes

that all differ significantly from the values proposed by Hansen (2024); $\beta_1 = $ -0.72 (95% CI -

0.78, -0.67), $\beta_2 = $ -1.19 (95% CI -1.22, -1.17), $\beta_3 = $ -0.23 (95% CI -0.32, -0.13).

Secondary sexual traits

Svensson (2019) presents a compilation of contemporary rates of evolution of secondary

sexual traits in the wild. These traits include morphology, color, and behavior traits in fish,

birds, mammals, and insects that have been shown to be a target of sexual selection and span a

time range of 3-26 generations. We find a negative relationship between rates and generation

time interval (Figure 1B); a log-log regression of absolute haldanes versus time reveals a slope

that does not differ from -1 (β = -1.306, 95% CI -1.825, -0.786), corresponding to a null

relationship between haldane numerator and time (Figure 1H). This finding of a -1 scaling

relationship over timespans spanning a handful of generations also contradicts the proposal for

null scaling relationships over such short microevolutionary intervals (Hansen 2024).

Fossil lineages

Similar results are seen in patterns of lineage accumulation in the fossil record (Figure

1C, I), where a strong negative rate-time relationship (found by (Henao Diaz, Harmon et al.

2019) and replotted here) reflects a weak and variable relationship between evolutionary change

and time (breakpoint regression $F_{2, 86} = 3.199$, $P = 0.046$, breakpoint = -109 MY, SE = 33.13).

However, this breakpoint regression model was not a significantly better fit than a model with

only an intercept; $F_{3, 86} = 2.6$, $P = 0.057$. Consistent with this finding, a log-log regression

between rate and time yields a slope that is significantly shallower than -1 ($\beta = -0.24$, 95%CI -

0.36, -0.11) and no significant evidence for a breakpoint ($F_{2, 75} = 1.71$, $P = 0.18$), indicating

accumulation of diversity with interval time.

Phylogenetic trees

On phylogenetic trees (Figure 1D, E, J, K), there is also a consistent striking negative

relationships between rate and time (as found by the original studies; Scholl and Wiens 2016,

Henao Diaz, Harmon et al. 2019) that corresponds to underlying variable association between

diversity and time (Rabosky, Slater et al. 2012, but see Scholl and Wiens 2016). We recover a

weak positive linear relationship between diversity and time in one dataset ($F_{1, 112} = 14.17$, $P =$

0.000267; Figure 1J) corresponding to a log-log regression that is significantly shallower than -1

($\beta = -0.53$, 95%CI -0.68, -.39), indicating some accumulation of diversity with time, and a

breakpoint linear relationship in another dataset (Figure 1K; breakpoint regression $F_{2, 430} = 12.67$,

$P = 4.474 \times 10^{-6}$, breakpoint = 4.6 log MY, SE = 0.108) corresponding to a log-log regression of

rate and time with a slope that is not significantly different from -1 ($\beta = -1.04$, 95%CI -1.1, -

0.98), indicating the accumulated diversity and interval time are decoupled.

Within-generation change

Interestingly, we even recover a nonlinear negative relationship between evolutionary

rates and time when examining phenotypic change that occurred within a timeframe less than the

average generation time of the study organism (here primarily long-lived fish; Figure 1F). This

within-generation dataset had a log-log regression of rate and time with a slope that was not

significantly different from -1 (β = -0.83, 95%CI -1.08, -0.58), indicating that there is no

accumulation of change with interval time for intervals less than one generation, as we may

expect since such change is likely an outcome of non-accruing environmental effects. This final

example demonstrates the appearance of a rate-time scaling relationship for a case that obviously

defies evolutionary explanation based on between-generation change. This conclusion is further

strengthened by the fact that the slope estimate does not significantly differ from the expectation

under a random relationship between rate numerator and time. That is, scaling phenotypic

change by time to calculate rate generates a non-linear negative relationship with time even for

this within-generation change (Figure 1F, L), that is likely to reflect phenotypic plasticity,

culling, or other non-evolutionary within-generational environmental effects.

Decomposing the covariance between rate and time across datasets

We can then decompose the covariance between rate and time for each of these datasets

into its components associated with variation in time and covariance between accumulated

change and time, as in equation 4. A caveat in this exercise is that not all of the rate estimates

are calculated directly as in equation 1, for example phylogenetic lineage diversification rates in

(Henao Diaz, Harmon et al. 2019), and so we do not expect all of the variance in rate-time

relationships to be explained by these two components. Nonetheless, this analysis reveals that

99.5% (39% with morphology data dropped) of the variance in rate-time relationships can be

explained by variance in time alone ($F_{1,4}$ = 844.5, P = 8.3x10^{-6}, Figure 6A), while change-time

covariance alone explains only 10.6 % (29% with morphology data dropped) of the variance in

rate-time relationships ($F_{1,4}$ = 0.47, P = 0.53, Figure 6B). Placing both predictors in a multiple

regression reveals support for variance in time (t = -28.4, P = 9.5x10^{-5}, df = 3), but not

covariance between change and time (t = 1.13, P = 0.34, df = 3), as a significant predictor of

rate-time relationships. While all covariance relationships will depend strongly on variance in

their constituent variables, this analysis reveals the noteworthy contrast in lack of meaningful

contribution of covariance between accumulated change and time.

Discussion

In this perspective we have explored the problem of why all estimated evolutionary rates

appear to scale negatively with time, and what information may be gleaned from these rate-time

relationships. First, it is readily apparent there will only exist a limited range of relationships

between the amount of evolution and time that generate anything other than negative rate-time

relationships. Second, revisiting several independent datasets of evolutionary rates illustrates

that existing, widely documented negative rate-time scaling relationships clearly suggest weak

and inconsistent relationships between the amount of evolution and timescale of measurement,

with no evidence of consistent time-dependency in evolutionary change across datasets. Our

novel analysis of these datasets reveals that variation in rate-time scaling relationships are

primarily driven by variation in timespans rather than patterns of accumulated evolutionary

change. Given that most common process-based models of constant-rate evolutionary change

can readily generate a range of negative rate-time scaling relationships, the ubiquitous negative

rate-time relationships in empirical data is hardly surprising. Although some of these points

have been discussed before, our revisit to the problem unequivocally illustrates the challenge in

interpreting negative rate-time scaling relationships, timely in light of their re-emergence in

recent literature (Henao Diaz, Harmon et al. 2019, Harmon, Pennell et al. 2021, Rolland,

Francisco Henao-Diaz et al. 2023).

How meaningful are rate-time scaling relationships?

Despite clear interpretive challenges inherent in evolutionary rate-time relationships,

evolutionary processes must of course play a driving role in generating them (Kinnison and

Hendry 2001, Ho, Lanfear et al. 2011, Henao Diaz, Harmon et al. 2019, Harmon, Pennell et al.

2021, Louca, Francisco Henao-Diaz et al. 2022, Rolland, Francisco Henao-Diaz et al. 2023), and

so there is justified interest in understanding the drivers of rate-time scaling relationships beyond

the inevitable qualitative expectation of a negative relationship (Harmon, Pennell et al. 2021).

Analyses of rate-time relationships can perhaps provide new insights when coupled with explicit

null hypothesis tests against scaling parameters expected under various evolutionary process

models (as can be done for the case of morphological evolution; Hansen 2024, Holstad, Voje et

al. 2024).

Here, we performed such a test and found no strong evidence for the three modes of

morphological evolutionary rate-time scaling proposed by Hansen (Hansen 2024) neither in our

reanalysis of an existing database of vertebrate body size evolutionary rate (Figure 5) nor in our

finding of a -1 relationship between microevolutionary rate and time (Figure 1b). Moreover, it

is challenging to construct meaningful null expectations for other types of rate-time scaling

patterns, such as lineage diversification. For example, in their assessment of negative scaling of

diversification rates, Henao Diaz et al. (2019) constructed a null expectation by simulating birth-

death trees for the same age ranges as their empirical set of trees, under the mean values of birth

and death rate observed for the oldest groups in the study. While such an approach is useful in

identifying cases of negative rate-time scaling beyond what may be an inevitable outcome from

our imperfect statistical models (but see Reijenga and Close 2025), for many types of rate

estimate we lack quantitative rate scaling predictions. One point of our paper is that, without

some robust quantitative expectation for a rate scaling relationship, the existence of a negative

relationship between empirical rates and time is not very informative.

Nonetheless, it will often be difficult to reconcile quantitative rate-time scaling patterns

with specific evolutionary process models, especially in large datasets of evolutionary rate

estimates spanning disparate lineages. For example, although a stationary OU model can

generate a -1 scaling relationship of rate with time (Hansen 2024), but this is only strictly true

over moderate timescales when selection is strong and genetic variance is moderate to high.

Similarly, evolution by Brownian motion produces a scaling relationship of -1/2, but so can a

random walk of the optimum under stabilizing selection, a process which itself is not strictly

Brownian (in terms of trait evolution) unless all phenotypic variance has an additive genetic

basis. Thus, while dramatic shifts in rate-time scaling relationships do imply shifts in the mode

of evolution (Hansen 2024), many similar rate-time scaling relationships may be generated by

many disparate processes.   Similarly, widespread decoupling of clade size and clade age has led

to proposals that rate-time relationships and measures of net diversification rate carry little

conceptual value (Rabosky 2009, Rabosky 2010, Rabosky, Slater et al. 2012, Rabosky and

Hurlburt 2015, Rabosky and Benson 2021), although counterpoints have been made (Wiens

2011, Harmon and Harrison 2015, Kozak and Wiens 2016). Similar arguments could be made

for morphological evolution: if evolutionary change rarely accumulates in a consistent manner

with time, what is the value of scaling change by time in the first place?

Parameter vs summary rates: A distinction without difference?

In general, we should be surprised when estimated evolutionary rates *do not* show negative time scaling. However, some estimators of evolutionary rate do not show the types of inevitable time scaling relationships we have discussed; for example, clade-level estimates of Brownian motion rate appear to be less affected. Moreover, constancy of evolutionary rates, and thus linear accumulation of change with time is sometimes observed at the molecular level (Morgan 1998, Bromham and Penny 2003). Indeed, in their review, Harmon et al. (see also Hunt 2012, 2021) draw a distinction between "summary rates" and "parameter rates"; the former being descriptive summary statistics, such as haldanes, while the latter are inferences of instantaneous rate parameters estimated from a process-based model of evolutionary change, with Brownian motion rate or birth-death lineage diversification rate being two well known examples. In this distinction, and barring inadequate models or data, it has been proposed that parameter rates from constant rate models should be free from inevitable time-scaling (Hunt 2012, Harmon, Pennell et al. 2021). This need not be strictly true, since an estimator of a population parameter can be the maximum likelihood estimator yet still be biased in certain contexts (a classic example is Pearson's correlation coefficient, which is the MLE of the correlation for a bivariate normal population, but demonstrates small-sample bias; Fisher 1915). Many maximum likelihood estimators of parameter rates are ultimately simple functions of amount of evolution and time, and noteworthy is that even most summary rates could in fact be considered parameter rates under appropriate evolutionary models (Harmon, Pennell et al. 2021). The haldane, for example is a parameter rate under one very simple model of directional trait change (constant directional selection) but shows positive and negative time scaling in different process-based manifestations of directional evolution models (see Fig. 4C). Given that we can

never know the true process model producing an evolutionary dataset all our models will always

be wrong to some degree (Box 1976). It remains to be seen if there is a general distinction

between these categories of rate estimate in regard to prevalence or interpretability of time

scaling.

Analysis of rate numerators

Relationships between the magnitude of evolutionary change (e.g., phenotypic change or

the log number of new lineages arising) and time can be assessed directly, as we and others have

done (Hendry and Kinnison 1999, Uyeda, Hansen et al. 2011, Svensson 2019). Breakpoint

regression (Schlager, Marsal et al. 1998, Hendry and Kinnison 1999, Uyeda, Hansen et al. 2011,

Clapham and Karr 2012) can be used to identify significant breakpoints in patterns of

accumulated evolutionary change across time. Such an approach avoids the problem of statistical

confounders between rate and time interval, at least when applied to analysis of change-time

relationships, and also allow for the possibility of identifying shifts in the form of rate-time

scaling relationships (Hansen 2024). Although analysis of rate-time and change-time plots

present the same total information, uninformed comparisons of rate-time relationships will

readily generate the appearance of shared features of the evolutionary process. Our analysis

revealed that variation in evolutionary rate-time relationships across datasets revealed little

information about variation in how evolutionary change accumulated with time.

Conclusions

A negative rate-time relationship is nearly inevitable, and the meaning and importance of

negative rate-time relationships for the correspondence of microevolution and macroevolution is

still unclear. We end with a final example that illustrates this point. Consider the widely-

appreciated fact that microevolutionary quantitative genetic models of random genetic drift are

inconsistent with the amount of evolutionary change in deep time (Lynch 1990, Hansen and

Houle 2004, Houle, Bolstad et al. 2017). We often find far less evolutionary change in deep time

than expected via drift alone, based on patterns of standing extant genetic variance. Yet even if

phenotypic divergence in deep time is consistent with drift under observed levels of standing

variance, we would still expect to see a negative scaling relationship between lineage specific

rates of evolution and time interval (Figure 3). This fact alone is underappreciated and highlights

that great caution is needed in interpreting negative rate-time relationships.

References

Alba, M. and J. Castresana (2005). "Inverse relationship between evolutionary rate and age of
mammalian genes." Molecular Biology and Evolution **22**: 3598-3606.
Arnold, S. J. (2014). "Phenotypic evolution: the ongoing synthesis." The American Naturalist
183: 729-746.
Atchley, W. R., C. T. Gaskins and D. Anderson (1976). "Statistical Properties of Ratios. I.
Empirical Results." Systematic Zoology **25**: 137-138.
Box, G. E. P. (1976). "Science and statistics." Journal of the American Statistical Assosciation
71: 791-799.
Bromham, L. and D. Penny (2003). "The modern molecular clock." Nature Reviews Genetics **4**:
216-224.
Bürger, R. and M. Lynch (1995). "Evolution and extinction in a changing environment: a
quantitative-genetic analysis." Evolution **49**: 151-163.
Charlesworth, B., R. Lande and M. Slatkin (1982). "A neo-darwininan commentary on
macroevolution." Evolution **36**(3): 474-498.
Chouteau, M., M. Arias and M. Jorun (2016). "Warning signals are under postive frequency-
dependent selection in nature." Proceedings of the National Academy of Sciences **113**: 2164-
2169.
Clapham, M. E. and J. A. Karr (2012). "Environmental and biotic controls on the evolutionary
history of insect body size." Proceedings of the National Academy of Sciences **109**: 10927-
10930.

Eldredge, N. and S. J. Gould (1972). Punctuated equilibria: an alternative to phyletic gradualism. Models in Paleobiology. T. J. M. Schopf. Sanfrancisco, Freeman Cooper.

Emerson, B. C. (2007). "Alarm bells for the molecular clock? No support for Ho et al.'s model of time-dependent molecular rate estimates." Systematic Biology **56**: 337-345.

Emerson, B. C. and N. Kolm (2005). "Species diversity can drive speciation." Nature **434**: 1015-1017.

Estes, S. and S. J. Arnold (2007). "Resolving the paradox of stasis: models with stabalizing selection explain evolutionary divergence on all timescales." American Naturalist **169**: 227-244.

Felsenstein, J. (1988). "Phylogenies and quantitative characters." Annual Review of Ecology and Systematics **19**: 445-471.

Fisher, R. (1915). "Frequency distribution of the values of the correlation coefficient in samples from an indefinitely large population." Biometrika **10**: 507-521.

Fisher, R. A. (1930). The genetical theory of natural selection. Oxford, The Clarendon press.

FitzJohn, R. G. (2012). "Diversitree: comparative phylogenetic analyses of diversification in R." Methods in Ecology and Evolution **3**: 1084-1092.

Foote, M. (1994). "Temporal variation in extinction risk and temporal scaling of extinction metrics." Paleobiology **20**: 424-444.

Foote, M. (2000). "Origination and extinction components of taxonomic diversity:General problems." Paleobiology **26**: 74-102.

Gingerich, P. D. (1983). "Rates of evolution: Effects of time and temporal scaling." Science **222**: 159-161.

Gingerich, P. D. (1993). "Quantification and comparison of evolutionary rates." American Journal of Science **293**: 453-478.

Gingerich, P. D. (2001). "Rates of evolution on the time scale of the evolutionary process." Genetica **112**: 127-144.

Gould, S. J. (1984). "Smooth curve of evolutionary rate: A psychological and mathematical artifact." Science **226**: 994-996.

Gould, S. J. and N. Eldredge (1977). "Punctuated equilibria: the tempo and mode of evolution reconsidered." Paleobiology **3**: 115-151.

Gould, S. J. and N. Eldredge (1993). "Punctuated equilibrium comes of age." Nature **366**: 223-227.

Haldane, J. B. S. (1949). "Suggestions as to Quantitative Measurement of Rates of Evolution." Evolution **3**: 51-56.

Hansen, T. F. (2024). "Three modes of evolution? Remarks on rates of evolution and time scaling." Journal of Evolutionary Biology **in press**.

Hansen, T. F. and D. Houle (2004). Evolvability, stabilizing selection, and the problem of stasis. Phenotypic Integration: Studying the Ecology and Evolution of Complex Phenotypes, Oxford University Press.

Harmon, L. J. and S. Harrison (2015). "Species diversity is dynamiv and unbounded at local and continental scales." The American Naturalist **185**: 5584-5593.

Harmon, L. J., M. W. Pennell, L. Francisco Henao-Diaz, J. Rolland, B. N. Sipley and J. C. Uyeda (2021). "Causes and Consequences of Apparent Timescaling Across All Estimated Evolutionary Rates." Annual Review of Ecology and Systematics **52**: 587-609.

Henao Diaz, L. F., L. J. Harmon, M. T. C. Sugawara and M. W. Pennell (2019). "Macroevolutionary diversification rates show time dependency." Proceedings of the National Academy of Sciences **116**: 7403-7408.

Hendry, A. P. and M. Kinnison (1999). "Perspective: the pace of modern life: measuring rates of contemporary microevolution." Evolution **53**: 1637-1653.

Ho, S. Y. W., R. Lanfear, L. Bromham, M. J. Phillips, J. Soubrier, A. G. Rodrigo and A. Cooper (2011). "Time dependency of molecular rate estimates and systematic overestimation of recent divergence times." Molecular Ecology **20**: 3087-3101.

Ho, S. Y. W., B. Shapiro, M. J. Phillips, A. Cooper and A. J. Drummond (2007). "Evidence for time dependency of molecular rate estimates." Systematic Biology **2007**: 515-522.

Holstad, A., K. L. Voje, O. H. Opedal, G. H. Bolstad, S. Bourg, T. F. Hansen and C. Pélabon (2024). "Evolvability predicts macroevolution under fluctuating selection." Science **383**: 688-693.

Houle, D., G. H. Bolstad, K. van der Linde and T. F. Hansen (2017). "Mutation predicts 40 million years of fly wing evolution." Nature **548**: 447-450.

Hunt, G. (2012). "Measuring rates of phenotypic evolution and the inseperability of tempo and mode." Paleobiology **38**: 351-373.

Jackson, D. A. and K. M. Somers (1991). "The spectre of 'spurous' correlations." Oecologia **86**: 147-151.

Kenney, B. C. (1982). "Beware of Spurious Self-Correlations!" Water Resources Research **18**: 1041-1048.

Kinnison, M. T. and A. P. Hendry (2001). "The pace of modern life II: from rates of contemporary microevolution to pattern and process." Genetica **112**: 145-164.

Kirkpatrick, M. (1982). "Sexual selection and the evolution of female choice." Evolution **36**: 1-12.

Kozak, K. H. and J. J. Wiens (2016). "Testing the relationships between divsersification, species richness, and trait evolution." Systematic Biology **65**: 975-988.

Lande, R. (1976). "Natural selection and random genetic drift in phenotypic evolution." Evolution **30**: 314-334.

Lande, R. (1981). "Models of speciation by sexual selection on polygenic traits." Proceedings of the National Academy of Sciences of the United States of America-Biological Sciences **78**(6): 3721-3725.

Louca, S., L. Francisco Henao-Diaz and M. W. Pennell (2022). "The scaling of diversification rates with age is likely explained by sampling bias." Evolution **76**: 1625-1637.

Louca, S. and M. W. Pennell (2020). "Extant timetrees are consistent with a myriad of diversification histories." Nature **580**: 502-505.

Lynch, M. (1990). "The rate of morphological evolution in mammals from the standpoint of the neutral expectation." The American Naturalist **136**: 727-741.

Lynch, M. and R. Lande (1993). Evolution and extinction in response to environmental change. Biotic Interactions and Global Change. P. Kareiva, J. G. Kingsolver and R. Huey. Sunderland, MA, Sinauer Assosciates.

Magallon, S. and M. J. Sanderson (2001). "Absolute diversification rates in Angiosperm clades." Evolution **55**(9): 1762-1780.

Morgan, G. J. (1998). "Emile Zuckerlandl, Linus Pauling, and the Molecular Evolutionary Clock, 1959-1965." Journal of the History of Biology **31**: 155-178.

Nee, S. (2001). "Inferring speciation rates from phylogenies." Evolution **55**: 661-668.

Nee, S., R. M. May and P. H. Harvey (1994). "The reconstructed evolutionary process." Philosophical Transactions of the Royal Society of London Series B-Biological Sciences **344**: 305-311.

Nosil, P., J. L. Feder, S. M. Flazman and Z. Gompert (2017). "Tipping points in the dynamics of speciation." Nature Ecology and Evolution **1**: 0001.

O'Meara, B. C. and J. M. Beaulieu (2024). "Noise leads to the perceived increase in evolutionary rates over short time scales." PLOS Computational Biology **20**: e1012458.

Paradis, E. (2015). "Random phylogenies and the distribution of branching times." Journal of Theoretical Biology **387**: 39-45.

Pearson, K. (1897). "Mathematical contributions to the theory of evolution.—On a form of spurious correlation which may arise when indices are used in the measurement of organs." Proceedings of the Royal Society of London **60**: 489-498.

Prarie, Y. T. and D. F. Bird (1989). "Some misconceptionsabout the spurious correlation problem in the ecological literature." Oecologia **81**: 285-288.

Quintero, I. (2025). "The diffused evolutionary dynamics of morphological novelty." Proceedings of the National Academy of Sciences **122**: e2425573122.

Rabosky, D. L. (2009). "Ecological limits and diversification rate: alternative paradigms to explain the variation in species richness among clades and regions." Ecology Letters **12**: 735-743.

Rabosky, D. L. (2010). "Primary controls on species richness in higher taxa." Systematic Biology **59**: 634-645.

Rabosky, D. L. (2016). "Challenges in the estimation of extinctio from molecular phylogenies: A response to Beaulieu and O'meara." Evolution **70**: 218-228.

Rabosky, D. L. and R. B. J. Benson (2021). "Ecological and biogeographic drivers of biodiversity cannot be resolved using clade age-richness data." Nature Communications **12**: 2945.

Rabosky, D. L., M. Grundler, C. Anderson, P. Title, J. J. Shi, J. W. Brown, H. Huang and J. G. Larson (2014). "BAMMtools: an R package for the analysis of evolutionary dynamics on phylogenetic trees." Methods in Ecology and Evolution **5**: 701-707.

Rabosky, D. L. and A. H. Hurlburt (2015). "Species richness at continental scales is dominated by ecological limits." The American Naturalist(185): 572-583.

Rabosky, D. L., G. J. Slater and M. E. Alfaro (2012). "Clade age and species richness are decoupled across the Eukaryotic tree of life." PLoS Biology **10**: e1001381.

Reijenga, B. R. and R. A. Close (2025). "Apparent timescaling of fossil diversfication rates is caused by sampling bias." Current Biology **35**: 905-910.e903.

Ricklefs, R. E. (2006). "Global variation in the diversification rate of passerine birds." Ecology **87**(2468-2478).

Rolland, J., L. Francisco Henao-Diaz, M. Doebeli, R. M. Germain, L. J. Harmon, L. L. Knowles, L. H. Liow, J. E. Mank, A. Machac, S. P. Otto, M. W. Pennell, N. Salamin, D. Silvestro, M. Sugawara, J. C. Uyeda, C. E. Wagner and D. Schluter (2023). "Conceptual and empirical bridges between micro- and macroevolution." Nature Ecology and Evolution **7**: 1181-1193.

Roopnarine, P. D. (2003). "Analysis of rates of morphologic evolution." Annual Review of Ecology and Systematics **34**: 605-632.

Sadler, P. M. (1981). "Sediment accumulation rates and the completeness of stratigraphic sections." The Journal of Geology **89**: 569-584.

Sanderson, S., M.-O. Beausoleil, R. E. O'Dea, Z. T. Wood, C. Correa, V. Frankel, L. D. Gorné, G. E. Haines, M. T. Kinnison, K. B. Oke, F. Pelletier, Pérez-Jvostov, W. D. Reyes-Corral, Y. Ritchot, F. Sorbara, K. M. Gotanda and A. P. Hendry (2021). "The pace of modern life, revisisted." Molecular Ecology **31**: 1028-1043.

Scheffer, M. (2009). Critical Transitions in Nature and Society. Princeton, N.J., Princeton University Press.

Schlager, W., D. Marsal, P. A. G. van der Geest and A. Sprenger (1998). "Sedimentaion rates, obersvation span, and the problem of spurious correlation." Mathematical Geology **30**: 547-556.

Scholl, J. P. and J. J. Wiens (2016). "Diversification rates and species richness across the tree of life." Proceedings B **283**: 20161334.

Sheets, H. D. and C. E. Mitchell (2001). "Uncorrelated change produces the apparent dependence of evolutionary rate on interval." Paleobiology **27**: 429-445.

Svensson, E. I. (2019). "Eco-evolutionary dynamics of sexual selection and sexual conflict." Functional Ecology **33**: 60-72.

Uyeda, J. C., T. F. Hansen, S. J. Arnold and J. Pienaar (2011). "The million year wait for macroevolutionary bursts." Proceedings of the National Academy of Sciences of the United States of America **108**: 15908-15913.

Wiens, J. J. (2011). "The causes of species richness patterns across space, time, and clades and the role of "ecological limits."." Quarterly Review of Biology **86**: 75-96.

Wiens, J. J. and K. E. Saban (2025). "Questioning the sixth mass extinction." Trends in Ecology and Evolution **40**: 375-384.

Fig. 1. Evolutionary rate-time relationships reflect null or variable underlying relationships between evolutionary change and time. Panels **A**-**F** show evolutionary rate plotted against the timescale of measurement (Rolland, Francisco Henao-Diaz et al. 2023), with dashed red lines showing the least squares fit of the function *rate = a**time^b and inset panels showing the linear relationship on a log-log scale. Panels **G**-**L** plot the amount of evolutionary change (i.e., a rate numerator) against time frame for the same datasets, with dashed red lines showing the linear breakpoint regression, if statistically significant. Panels **A** and **G** show vertebrate body size evolution (Uyeda, Hansen et al. 2011), Panels **B** and H show microevolution of secondary sex traits (Svensson 2019), **C** and **I** show diversification of genera from fossil lineages (plants,

mammals, and marine animals)(Henao Diaz, Harmon et al. 2019), Panels **D** and **J** show diversification data estimated from time-calibrated molecular phylogenies (Henao Diaz, Harmon et al. 2019), while Panels **E** and **K** show a larger database of phylogenetic diversification rates (Scholl and Wiens 2016) where rates were inferred directly from age and richness data (Magallon and Sanderson 2001). Panels **F** and **L** show within-generation morphological changes (mainly from fish harvesting). All nonlinear negative relationships plotted in **A**-**F** were statistically significant, whereas relationships in **G-L** were variable, with statistically significant breakpoints found for morphology, fossil lineage accumulation, and one estimator of phylogenetic diversification, and a positive linear relationship for another estimator of phylogenetic diversification. No significant relationship was found for microevolution of sex traits (**H**) or the within-generation changes in panel **L**. Sample sizes for each column: **A** N = 5392 records of evolutionary change, **B** N = 23 records of evolutionary change, **C** N = 90 lineages, **D** N = 114 clades, **E** N = 434 Orders, **F** N = 302 records of within-generation change. Note that each of these datasets is an amalgamation of evolutionary events from many different lineages, and does not represent single anagenetic time series; thus decreasing numerator-time relationships are indeed possible.

A   **B**

Fig. 2. Extraordinary relationships between the accumulated amount of evolution and time are required to produce positive rate-time relationships. Panel **A** shows the relationship between rate and time for various functions of the rate numerator and time, which are plotted in the same color in Panel **B**. The black line is a reference point, where there is a null relationship between rate and time. Thus, each line in Panel **A** corresponds to a specific relationship between rate numerator and time, plotted in Panel **B**; only cases of an accelerating relationship between amount of evolution and time are expected to produce positive rate-time scaling relationships. We assume for this example a form of the relationship, *amount of evolution = time^{b}*, where the function was plotted for b ranging from -2 (blue) to 2 (red). Similar general patterns are observed for other functions of change and time. Note that because we are considering these relationships in light of large amalgamations of data from disparate groups (e.g., Figure 1), as opposed to within-lineage time series, negative relationships between amount of evolution and interval time are in principle possible, even if unlikely.

Fig. 3. Simulations of constant-rate evolutionary models reveal persistence of negative rate-time relationships in resulting data. Panels **A** and **B** show a simulation of phenotypic evolution under random genetic drift in the absence of natural selection, which produces a Brownian motion process (Felsenstein 1988) where variance among lineages accumulates linearly with time (**A**). Although the rate of drift is constant (**B** inset), we will nevertheless obtain a negative scaling relationship between the lineage-specific per-generation rate of evolution (**B**). Panels **C** and **D** show a simulation of phenotypic evolution under drift and stabilizing natural selection, an Ornstein Uhlenbeck process (Lande 1976). This process generates no relationship between time and divergence (**C**), but again a strong negative scaling relationship with per-lineage per-generation evolutionary rate and time and hence a similar negative relationship (**D**).

Fig 4. Heterogeneity in rate-time scaling exponents under different forms of morphological
evolution. Panels **A** and **B** show simulated evolution under drift and stabilizing selection; an OU
process. When there is zero variance in the optimum, this is a stationary OU process generating

a scaling relationship of -1 over this timespan (9000 generations) only when genetic variation is
relatively high (Panel **A**), and selection is strong (a low value of ω and thus a strong curvature of
the fitness surface, (Panel **B**). Any variance in the optimum leads to a dramatic shift in the
scaling relationship. Noteworthy is that a wide range of parameter space, corresponding to
models of pure drift as well as models of pure fluctuating selection, can all produce a range of
similar scaling relationships particularly in the range of 0 to -1/2. For each parameter value in **A**
and **B**, the average scaling slope is shown from 10 simulation runs of 36 lineages each spanning
a time range of 1-9000 generations. Panel **C** shows evolutionary rate versus time for three
forms of directional evolution. In black is the rate estimate versus time for evolutionary change
under the breeders equation, $\Delta\bar{z} = G\beta$. Blue shows the rate estimate for evolution towards an
optimum that shifted 5 phenotypic standard deviations at generation 0, assuming the same form
of stabilizing selection as in Figure 4B, which is high over short timescales but decelerates as the
population adapts. Red shows the relationship for evolution towards an optimum that moves at
contant rate of .1 standard deviations per generation, starting at generation 0 with a population on
the optimum. In this model the accelerating rate-time relationship is due to the lag between
evolutionary rate and rate of peak movement that equilibrates when the population reaches its
equilibrium lag behind the optimum(Lynch and Lande 1993, Bürger and Lynch 1995). All of
these models produce directional evolutionary change, yet only for one specific model, $\Delta\bar{z} =$
$G\beta$, which is often thought to capture the main features of the two other models of directional
change, do we see a lack of time scaling.

Figure 5. No evidence for three distinct modes of evolution revealed in vertebrate body size evolutionary rate-time scaling. Shown is the log rate-time scaling relationship for vertebrate body size data (Uyeda, Hansen et al. 2011). The black line shows the breakpoint regression with two breakpoints, corresponding to a model of three "modes" of rate-time scaling (Hansen 2024), while the red line shows the best-fit breakpoint regression (see Table S1).

758
759
Fig. 6. Time interval determines rate-time relationships across datasets. The covariance between rate and time
in the six empirical datasets shown in Figure 1 is almost entirely explained by variance in time (Panel **A**).
Covariance between change and time, also expected to be a component of the covariance between rate and time, is
not significantly associated (Panel **B**). Lines and 95% confidence intervals are from linear models fit separately to
each; a multiple regression with both var(time) and cov(change, time) as predictors yielded equivalent conclusions
(see text).
766
767

Methods Details

Data. We analyze existing data on rates of vertebrate body size evolution (Uyeda, Hansen et al. 2011), which consists of data on log-fold change in body size as well as time interval over which this change occurred. For phylogenetic lineage diversification, we analyze two datasets, one in which diversification rates were estimated from the branching pattern of species-level phylogenies (Henao Diaz, Harmon et al. 2019), and another larger dataset (Scholl and Wiens 2016) where diversification rates were estimated from clade diversity and age information. For fossil data, we analyze an existing dataset (Henao Diaz, Harmon et al. 2019) of emergence rates of new genera. For each dataset, we plot the standard measure of rate against the timeframe over which that rate was measured, fitting exponential linear models by nonlinear least squares (Rolland, Francisco Henao-Diaz et al. 2023). We also plot a measure of the rate numerator against time for each of these datasets, and fit breakpoint linear regressions to identify significant linear relationships between amount of evolution and time, as well as any potential shifts in the relationship between evolution and time. For each dataset, we fit nonlinear models of the form $rate = a*time^b$ using nonlinear least squares (nls()) in R, as well as breakpoint regression using the segmented package. We compared models fits of an ordinary linear regression to a model with a single breakpoint, and also when appropriate to a null model containing only an intercept, using the anova() function in R. Data presented in Figure 1 were obtained from (Henao Diaz, Harmon et al. 2019) (available at https://github.com/fhenaod/evol_time_scaling/tree/main/data), (Uyeda, Hansen et al. 2011)(Dryad doi:10.5061/dryad.7d580), (Scholl and Wiens 2016), and (Sanderson, Beausoleil et al. 2021) (database available at: https://proceeddatabase.weebly.com/).

Simulation We simulated three different constant rate evolutionary processes. The aim of our simulations was not to evaluate the performance of statistical models used to estimate rates. Rather, our aim was simple to examine scaling patterns observed in the simplest cases of

simulated constant rate evolution. For morphological evolution, we simulated random genetic

drift for 1-1000 generations assuming a heritability of 0.50 and a population size of 100. We also

simulated evolution under genetic drift and stabilizing selection, with the same demographic and

genetic assumptions but assuming weak selection ($\omega^2 = 15$) towards an optimum. Each

simulation was started with the population at its optimum. For morphological evolution, we

plotted rates in units of Haldanes (Gingerich 1993), which was natural as we simulated evolution

on the scale of phenotypic standard deviations. We then repeated our simulations of OU

evolution but exploring a range of parameter values for genetic variance, the strength of selection

(curvature of the fitness surface) and the stability of the optimum. For the later, we sampled

changes in the optimum from a normal distribution centered at zero and exploring different

values of the variance, producing a random walk of the optimum.

Table S1. Model fits for testing Hansen's "three modes" hypothesis.
Note that models with more than 7 breaks failed to fit

parameters	breakpoints	AIC
2	0	4687.959
4	1	4287.43
6	2	4065.896
8	3	3075.987
10	4	2956.282
12	5	2865.407
14	6	2821.511
16	7	2778.154