Title: **Evolutionary change is remarkably constant across timescales**

Authors: Stephen P. De Lisle[1]*, Erik I. Svensson[1,2]

Affiliations:
[1] Department of Environmental and Life Science, Karlstad University, Karlstad, Sweden
[2] Department of Biology, Lund University, Lund, Sweden
*Corresponding author. Email: stephen.de.lisle@kau.se

Apparent time-scaling of evolutionary rates presents an evolutionary dilemma; rates of
molecular, phenotypic, and lineage diversification scale strongly negatively with timeframe
of measurement, raising questions about the processes driving this general pattern. Here
we show that much of this apparent temporal scaling is an inevitable outcome of plotting a
ratio against its denominator. Highly unlikely relationships between timescale and amount
of evolutionary change are required to produce anything other than artefactual negative
relationships between rate and time. Unscaled magnitudes of evolution are largely
unrelated to time, are easily explained by existing evolutionary models, and reframe the
widely-held view of disparate pace of micro- and macroevolution. Time-independence of
evolutionary change raises new questions about the factors that generate such temporal
consistency in evolution.

Main Text: Evolutionary rates vary with the time interval over which they are measured. A seemingly general pattern with faster evolutionary rate over short timescales, and slower rates over longer timescales has been recognized for decades [1-5]. Such apparent time-dependence has subsequently been documented consistently for nearly every possible measure of evolutionary rate [6], including rates of morphological evolution, rates of molecular evolution [7], and rates of lineage formation on phylogenies and in the fossil record (Figure 1). Negative rate-time scaling relationships were initially proposed as an artifact that needed to be taken in to account, in that evolutionary rates over different time intervals were not directly comparable [5]. Yet these negative relationships between evolutionary rate and time continue to generate discussions and stimulate biological hypotheses for shared features of the underlying evolutionary processes that could explain such general time-scaling [6,8]. Recent technical and statistical explanations to these apparent negative rate-time scaling relationships focus on sampling biases [9], such as a the "push of the past" [6] and model mis-specification [2]. This puzzling and general phenomenon of apparent widespread negative rate-time scaling relationships has been recently highlighted as a major unsolved challenge in evolutionary biology [2,6,8]. The renewed interest in such apparent time-dependence of evolutionary rates has re-kindled long-standing ideas and hypotheses about fundamental disparities between micro- and macroevolution [10-14].

Here we show that much of this apparent paradox of strong time-dependency of evolutionary rate is readily explained by the inevitable negative correlation generated by plotting a ratio against its own denominator [15]. While this is a well-known statistical issue across a range of scientific fields [5,16-19], recent discussions of rate-time scaling patterns in evolutionary biology have either have either downplayed the importance of the mathematical dependence of rate and time [2,6,8], or have focused on explanations arising from specific statistical models used to estimate rates in a single context [9]. By revisiting the functional dependence of rates and time,

analyzing large existing datasets and using new simulations, we show that a negative relationship

between evolutionary rate and time is nearly inevitable, regardless of the underlying evolutionary

process. We show that the key feature underlying negative rate-time relationships in empirical

data is in fact a weak, null, or variable relationship between magnitude of evolution and time.

Uninformed focus on rate-time relationships therefore provide little insight into underlying

evolutionary processes even under highly idealized conditions, and the empirical evidence

instead suggest remarkable time-constancy of evolutionary change across timescales.

The negative rate-time association is nearly inevitable

All measures of evolutionary rate are a function of evolutionary change scaled by time

$$y = d/t \tag{1}$$

where y is some measure of rate, d is some measure of evolutionary change (e.g., change in

population mean phenotype, base pair substitutions, log change in number of species in a clade)

and t is some measure of time interval over which change was observed or inferred (e.g.

generations, or years). Some classical estimates of evolutionary rate are calculated directly by

dividing phenotypic change with time, the latter typically measured in generations. For example,

Darwins [20] and Haldanes [4] have been used to describe morphological rate, and other types of

rate estimators measure per-capita origination rate of fossil lineages [21] and phylogenetic net

diversification rate [22]. Other estimates are obtained from more sophisticated statistical models of

evolution [23-25]. Regardless, all of these measures of evolutionary rates are, by definition, scaled

by time.

The mathematical dependence between rate and time is an obvious challenge in

interpreting the time-dependence of evolutionary rate [5]. Moreover, even a random association

between evolutionary change and time will result in a strong negative scaling relationship

between rate and time [26]. The strength of this relationship will only increase with a greater range

of timescales sampled [15]. This problem of interpreting relationships between a rate and the

timescale over which rates have been measured have been discussed in other scientific fields

outside evolutionary biology, including geology [27] and paleontology [28]. This problem has also

raised in evolutionary biology, in the context of rates of morphological evolution [3,5,17,29-31] and

lineage diversification [32-34]. Here, we revisit and elaborate on the issues with interpreting rates

as a function of time.

It is useful to ask, under what scenarios do we expect null or positive relationships

between rate and time? Consider rate as a function of time,

$$y = f(t) \tag{2}$$

Where y is evolutionary rate (as in eqn 1). We also consider the amount, or magnitude, of

evolution (rate numerator) as a function of time,

$$d = g(t) \tag{3}$$

where d is evolutionary divergence, the magnitude of evolutionary change. This could be change

in trait means between allochronic or synchronic populations, difference in log number of

lineages between two sampling points in time (e.g., the stem age of a clade and present), or any

other measure of evolutionary change. Rates, $y = d/t$, will be constant over time when d scales

linearly with t, i.e. $g'(t)$ is a positive constant over the timescale in question. Rates will increase

with time, i.e. $f'(t) > 0$ when $g''(t) > 0$; that is, rates increase over time when the amount of

evolutionary change is an accelerating function of time.

Noteworthy is that under a null relationship between d and t, such that $g'(t) = 0$, the

logarithmic regression of rate versus time ($\log(y) \sim \mu + \beta*\log(t)$) will yield a slope estimate of β

= -1 [5]. Thus, this slope estimate provides one cutoff of interest in assessing rate-time scaling

relationships: log-log regressions of rate vs time that are shallower than or not significantly

different from -1 are consistent with either a null relationship evolutionary change and time (time

independence) or an increasing relationship between change and time, respectively. Thus, if

there would exist a true biological, non-artefactual relationship between evolutionary rates and

time driven by accelerating change with time, we would expect log-log regressions with slopes

that are steeper than -1.

The relationships between equations 1-3 are illustrated in Figure 2, which shows the

relationship between rate and time and rate numerator and time illustrated for a range of power

functions. There are few existing models of evolution that predict the type of accelerating

relationship between magnitude of evolution and time needed to produce a positive relationship

between evolutionary rate and time. For example, neutral and adaptive models of phenotypic and

molecular evolution predict either no relationship with time or linear accumulation (constant

change) with time [35]. Similarly, birth-death models of lineage accumulation also predict either

linear or a decelerating accumulation of log lineage number through time under most conditions.

Even models of time-dependent lineage diversification, including models of accelerating

diversification rate with time before or towards present during the diversification of a single

clade [23], make no prediction about accelerating log lineage number versus interval time across a

sample of clades. Moreover, lineage-through-time (LLT) plots for single clades diversifying

under such a process can produce complex lineage through time plots [36] that are consistent with a

range of diversification processes [37].

Some specific biological and ecological scenarios could lead to accelerating change with

time, and thus potentially generate positive relationships between evolutionary rates and time

(Figure 2). Such scenarios include "diversity begets diversity" models of lineage diversification

[23,38], evolution by positive frequency-dependent selection [39], runaway sexual selection driven by

female choice [40-42], "tipping points" in the evolution of reproductive isolation [43] and populations

crossing critical thresholds followed by phase transitions between alternative stable states [44].

How common such ecological and evolutionary scenarios are in nature is an open empirical

question. However, given the transient nature of such rapid and self-reinforcing evolutionary

processes driven by positive feedbacks [41,42], we suspect that positive time-rate scenarios are

unlikely to be sustained over longer macroevolutionary time scales but should instead rather

quickly reach stable equilibria. Thus, such rapid and self-reinforcing evolutionary processes

seem unlikely to proceed unchecked over millennia [44], as would be required to generate positive

rate-time relationships in empirical data such as that shown in Figure 1.

Data and simulation reveal persistence of inevitable scaling relationships

Reanalysis of well-known examples of evolutionary rate-time associations reveals little

consistent patterns in the relationship between amount of evolutionary change and time, despite a

pervasive negative relationship between rate and time. This holds for body size evolution (Figure

1A, E), where past work [17,29,30] has shown that body size changes are largely unrelated to time,

and in time-series of extant populations found no detectable change [45]. Although a log-log

regression of rate vs time yields a slope that is significantly steeper than -1 (β = -1.019, 95%CI -

1.031, -1.007) we recover a breakpoint in the relationship between morphological change and

time ($F_{2, 5388}$ = 207.2, $P < 2.2 \times 10^{-6}$, breakpoint = -2.63 log MY, SE = 0.147). The result is an

estimated weak negative relationship between time and evolutionary change, followed by an

increasing slope after the breakpoint (Fig. 1A, E).

Similar results are seen in patterns of lineage accumulation in the fossil record (Figure

1B, F), where we found evidence for a strong negative rate-time relationship reflecting weak and

variable relationship between evolutionary change and time (breakpoint regression $F_{2, 86} = 3.199$,

$P = 0.046$, breakpoint = -109 MY, SE = 33.13). However, this regression model was not a

significantly better fit than a model with only an intercept; $F_{3, 86} = 2.6$, $P = 0.057$. Consistent

with this finding, a log-log regression between rate and time yields a slope that is significantly

shallower than -1 ($\beta = -0.24$, 95%CI -0.36, -0.11)

On phylogenetic trees (Figure 1C, D, G, H), we also find a consistent striking negative

relationships between rate and time [8] that in fact corresponds to underlying variable association

between diversity and time [32,]but see [46]. We recover a weak positive linear relationship between

diversity and time in one dataset ($F_{1, 112} = 14.17$, $P = 0.000267$; Figure 1H) corresponding to a

log-log regression that is significantly shallower than -1 ($\beta = -0.53$, 95%CI -0.68, -.39), and a

piecewise linear relationship in another dataset (breakpoint regression $F_{2, 430} = 12.67$, $P = 4.474$

x10^{-6}, breakpoint = 4.6 log MY, SE = 0.108) corresponding to a log-log regression of rate and

time with a slope that is not significantly different from -1 ($\beta = -1.04$, 95%CI -1.1, -0.98).

Interestingly, we even recover a nonlinear negative relationship between evolutionary

rate and time when examining phenotypic change that occurred within a timeframe less than the

average generation time of the study organism (in this case, primarily fish; Figure 1E). This

within-generation dataset had a log-log regression of rate and time with a slope that was not

significantly different from -1 ($\beta = -0.83$, 95%CI -1.08, -0.58). This demonstrates the appearance

of a rate-time scaling relationship for a case that obviously defies evolutionary explanation based

on between-generation change. This conclusion is further strengthened by the fact that the slope

estimate does not significantly differ from the expectation under a random relationship between

rate numerator and time. That is, scaling phenotypic change by time in order to calculate rate

generates a non-linear negative relationship with time even for this within-generation change

(Figure 1E, J), that is likely to reflect phenotypic plasticity, culling, or other non-evolutionary

within-generational environmental effects.

In summary: the striking negative rate-time relationship shared by a diverse array of

evolutionary datasets reflects, primarily, a lack of strong or consistent association between the

amount of evolutionary change and the timescale over which this change took place (Figure 1 F-

J). In only one case did we find evidence of a significant negative association between

evolutionary change and time that was less than -1 on a log scale. Yet, even in this case, that of

vertebrate body size evolution, breakpoint regression indicates that the true relationship between

time and evolutionary change involves a complex combination of decreasing and increasing

relationships between time and evolutionary change through the history of vertebrates [29]. We

further note that widespread time-independence of evolutionary change does not imply that

magnitudes of evolution are low or "slow". Rather, the vertical axes in Figure 1F-J often

correspond to massive changes in the number of lineages or morphology, yet these evolutionary

changes are largely decoupled from the time interval over which they occurred.

Simulations of classical constant-rate evolutionary models reveal that negative rate-time

associations readily emerge as statistical artifacts. Phenotypic evolution by random genetic drift

results in a Brownian motion process wherein phenotypic variance across lineages increases

linearly with time (Figure 3A), and evolutionary rate is time-independent when calculated for an

entire clade (3D inset panel). Even this simple constant rate model results in a negative scaling

relationship between lineage-specific estimates of evolutionary rate and time (Figure 3D). The

problem is even further exacerbated an under adaptive model of natural selection towards an

optimum (an Ornstein-Uhlenbeck process), where a null relationship between time and

divergence (Figure 3B) generates a strong negative relationship between per-lineage rate and

time (Figure 3E), reminiscent of empirical data from vertebrate body size evolution (Figure 1).

Finally, the simplest model of lineage accumulation, a constant-rate pure-birth process, also

leads to these types of associations. The log number of lineages accumulates approximately

linearly through time in a pure-birth process, as seen when sampling pure-birth phylogenies

differing in age (Figure 3C). Yet, pure-birth estimates of diversification rate nonetheless show a

negative scaling relationship with time (Figure 3F) that cannot be fully explained by the "push of

the past" [8], as we recover this effect in the absence of extinction and in model parameterizations

that yield high expected values of diversity early in time, minimizing the potential of only

sampling clades with stochastically high rates early in their history (see Materials and Methods).

This latter negative scaling relationships is driven entirely by the inclusion of high evolutionary

rates estimated over short timescales (Figure 3F). This illustrates the inherent problems of

interpreting evolutionary change scaled by time even when the evolutionary model is not mis-

specified. Moreover, we recover the same general pattern when simulating phylogenies of

varying age under an accelerating time-dependent diversification process (Figure S1).

Discussion

That evolutionary rates scale negatively with the timeframe over which they are

measured is unsurprising given the inevitable statistical dependence between rate and time.

Negative scaling relationships are therefore expected under a wide range of relationships

between evolutionary change and time, in spite of fundamentally different underlying

evolutionary processes behind the patterns (Figure 2). The fact that the scaling pattern is

observed consistently across disparate types of data from different sources, for rates estimated in

very different ways, and even within generations, strongly suggest that these negative

relationships are most likely statistical artifacts [2]. As Gingerich [5] emphasized, the negative

scaling relationship that is generated by plotting a rate against time interval must be accounted

for before rate variation across time can be meaningfully compared. Although the likely-

artifactual nature of evolutionary rate-time scaling relationships has been acknowledged since

[2,3,5,10], the pervasive negative relationship between evolutionary rate and time interval continues

to inspire evolutionary explanation [2,6-9,47].

This renewed interest in artefactual rate-time relationships is unfortunate in light of

alternative approaches to the analysis of time-dependence of the evolutionary process. For

example, relationships between the magnitude of evolutionary change (e.g., phenotypic change

or the log number of new lineages arising) and time can be assessed directly, as we have done

here and which has also been done by others [29,48]. For example, piece-wise or segmented

regression [3,27,29,49] can identify significant breakpoints in evolutionary change across time, an

approach that avoids the problem of statistical confounders between rate and time interval that

was highlighted by Gingerich [5]. These and related statistical approaches [29] can be used to identify

temporal shifts in the evolutionary process over time.

Our analyses of rate numerators versus time reveal lack of consistent time-dependency

across datasets (Figure 1). A conclusion from these results, and the finding of log-log rate-time

regression slopes that are shallower than null expectations, is that the persistent rate-time

relationship do not provide evidence of shared features of evolution across these disparate data,

contra recent suggestions [2,6,8]. Our reanalysis refutes the view that high microevolutionary rates

measured over short time intervals and low macroevolutionary rates measured over longer time

scales indicates fundamental disparity between micro- and macroevolution. By extension, the

common view [2,6,8] that microevolutionary rates documented in natural populations of extant

organisms [50,51] are exceptionally high in comparison to macroevolutionary change over longer time scales is not empirically supported, neither in this study nor by other empirical evidence (Fig. 1F-J; see also [29,52]).

Existing data (Fig. 1F-J) instead suggest that evolutionary change is remarkably independent of time, across multiple temporal scales. Such time-constancy becomes a new challenge to explain. Time-independence of phenotypic evolution be explained by relatively-stable or slowly moving adaptive zones, within which lineages may evolve substantially [29,53]. For lineage diversification, disparity in ecological opportunity among diversifying lineages may decouple clade age and species richness across taxa [33]. We emphasize that while we find little consistent relationships between amount of evolution and interval time, evolutionary events can still be substantial, as shown in Figure 1. We also note that timing of key events in the history of life has, of course, played a major role shaping the origins of diversity [54]. These include shifts in body size evolution [29] or rapid jumps in bill evolution in birds [55]. Our results are not inconsistent with such important historical events, but indicate that there is no evidence of a pervasive relationship between amount of evolution and interval time. There is also evidence of substantial variation in evolutionary change in response to different selective agents, including human disturbances and harvesting [45], but such change in response to different environmental agents does not appear to be systematically linked to time interval over which evolution was recorded, as we have shown here.

Our results and other recent findings [56] also call into question the empirical evidence underlying the apparent "paradox of stasis", which is an observation of far less evolutionary change observed over long timescales than what is expected based on microevolutionary rates [57]. Although it is often claimed that stasis is widespread and a predominant mode of evolution [11,12], the results in the present study and others [56,58,59] cast doubt on the view that stasis is predominant.

Given that the stasis paradox has partly been stimulated by apparent rate disparities through time, which have been interpreted as indicating a fundamental qualitative difference between micro- and macroevolution [6,57,60,61], our results instead indicate that the paradox of stasis could be reformulated as the "paradox of time-independence" of evolutionary change. Such a reformulation is also consistent with more recent interpretations of the paradox [29,56].

Time-invariant evolutionary change still poses other challenges for reconciling micro- and macroevolution [29]. Constancy of evolutionary rates, and thus linear accumulation of change with time, across temporal scales is expected and often observed at the molecular level [35]. It is more difficult to explain relative time-independence in phenotypic evolution, where trait change is expected to accumulate with time given strong and variable natural selection [62,63], or under neutral evolution [64,65]. However, new empirical findings indicate that phenotypic evolution over a range of timescales is proportional to standing phenotypic and genetic variation [66]. Such variation-limited evolution, if general across taxa, could be one possible explanation for time-constancy in evolutionary change, although it raises additional questions on how and why variational properties could be conserved through time. Time-dependence, or rather lack thereof, of the evolutionary process therefore remains a major open question in evolutionary biology.

References

1   Gingerich, P. D. Rates of evolution on the time scale of the evolutionary process. *Genetica* **112**, 127-144 (2001).
2   Harmon, L. J. *et al.* Causes and Consequences of Apparent Timescaling Across All Estimated Evolutionary Rates. *Annual Review of Ecology and Systematics* **52**, 587-609 (2021).
3   Hendry, A. P. & Kinnison, M. Perspective: the pace of modern life: measuring rates of contemporary microevolution. *Evolution* **53**, 1637-1653 (1999).
4   Gingerich, P. D. Quantification and comparison of evolutionary rates. *American Journal of Science* **293**, 453-478 (1993).

5   Gingerich, P. D. Rates of evolution: Effects of time and temporal scaling. *Science* **222**,
159-161 (1983).

6   Rolland, J. *et al.* Conceptual and empirical bridges between micro- and macroevolution.
Nature Ecology and Evolution **7**, 1181-1193 (2023).

7   Ho, S. Y. W. *et al.* Time dependency of molecular rate estimates and systematic
overestimation of recent divergence times. *Molecular Ecology* **20**, 3087-3101 (2011).

8   Henao Diaz, L. F., Harmon, L. J., Sugawara, M. T. C. & Pennell, M. W.
Macroevolutionary diversification rates show time dependency. *Proceedings of the*
National Academy of Sciences **116**, 7403-7408 (2019).

9   Louca, S., Francisco Henao-Diaz, L. & Pennell, M. W. The scaling of diversification
rates with age is likely explained by sampling bias. *Evolution* **76**, 1625-1637 (2022).

10   Gould, S. J. Smooth curve of evolutionary rate: A psychological and mathematical
artifact. *Science* **226**, 994-996 (1984).

11   Eldredge, N. & Gould, S. J. in *Models in Paleobiology* (ed T.J.M. Schopf) (Freeman
Cooper, 1972).

12   Hansen, T. F. & Houle, D. in *Phenotypic Integration: Studying the Ecology and*
Evolution of Complex Phenotypes (Oxford University Press, 2004).

13   Gould, S. J. & Eldredge, N. Punctuated equilibria: the tempo and mode of evolution
reconsidered. *Paleobiology* **3**, 115-151 (1977).

14   Gould, S. J. & Eldredge, N. Punctuated equilibrium comes of age. *Nature* **366**, 223-227
(1993).

15   Pearson, K. Mathematical contributions to the theory of evolution.—On a form of
spurious correlation which may arise when indices are used in the measurement of
organs. *Proceedings of the Royal Society of London* **60**, 489-498 (1897).

16   Prarie, Y. T. & Bird, D. F. Some misconceptionsabout the spurious correlation problem
in the ecological literature. *Oecologia* **81**, 285-288 (1989).

17   Arnold, S. J. Phenotypic evolution: the ongoing synthesis. *The American Naturalist* **183**,
729-746 (2014).

18   Atchley, W. R., Gaskins, C. T. & Anderson, D. Statistical Properties of Ratios. I.
Empirical Results. *Systematic Zoology* **25**, 137-138 (1976).

19   Jackson, D. A. & Somers, K. M. The spectre of 'spurous' correlations. *Oecologia* **86**,
147-151 (1991).

20   Haldane, J. B. S. Suggestions as to Quantitative Measurement of Rates of Evolution.
Evolution **3**, 51-56 (1949).

21   Foote, M. Origination and extinction components of taxonomic diversity:General
problems. *Paleobiology* **26**, 74-102 (2000).

22   Magallon, S. & Sanderson, M. J. Absolute diversification rates in Angiosperm clades.
Evolution **55**, 1762-1780 (2001).

23   Nee, S., May, R. M. & Harvey, P. H. The reconstructed evolutionary process.
Philosophical Transactions of the Royal Society of London Series B-Biological Sciences
344, 305-311 (1994).

24   Rabosky, D. L. *et al.* BAMMtools: an R package for the analysis of evolutionary
dynamics on phylogenetic trees. *Methods in Ecology and Evolution* **5**, 701-707 (2014).

25   FitzJohn, R. G. Diversitree: comparative phylogenetic analyses of diversification in R.
Methods in Ecology and Evolution **3**, 1084-1092 (2012).

26   Kenney, B. C. Beware of Spurious Self-Correlations! *Water Resources Research* **18**,
1041-1048 (1982).

Schlager, W., Marsal, D., van der Geest, P. A. G. & Sprenger, A. Sedimentaion rates, obersvation span, and the problem of spurious correlation. *Mathematical Geology* **30**, 547-556 (1998).

Sheets, H. D. & Mitchell, C. E. Uncorrelated change produces the apparent dependence of evolutionary rate on interval. *Paleobiology* **27**, 429-445 (2001).

Uyeda, J. C., Hansen, T. F., Arnold, S. J. & Pienaar, J. The million year wait for macroevolutionary bursts. *Proceedings of the National Academy of Sciences of the United States of America* **108**, 15908-15913 (2011).

Estes, S. & Arnold, S. J. Resolving the paradox of stasis: models with stabalizing selection explain evolutionary divergence on all timescales. *American Naturalist* **169**, 227-244 (2007).

Roopnarine, P. D. Analysis of rates of morphologic evolution. *Annual Review of Ecology and Systematics* **34**, 605-632 (2003).

Rabosky, D. L., Slater, G. J. & Alfaro, M. E. Clade age and species richness are decoupled across the Eukaryotic tree of life. *PLoS Biology* **10**, e1001381 (2012).

Rabosky, D. L. Ecological limits and diversification rate: alternative paradigms to explain the variation in species richness among clades and regions. *Ecology Letters* **12**, 735-743 (2009).

Wiens, J. J. The causes of species richness patterns across space, time, and clades and the role of "ecological limits.". *Quarterly Review of Biology* **86**, 75-96 (2011).

Morgan, G. J. Emile Zuckerlandl, Linus Pauling, and the Molecular Evolutionary Clock, 1959-1965. *Journal of the History of Biology* **31**, 155-178 (1998).

Paradis, E. Random phylogenies and the distribution of branching times. *Journal of Theoretical Biology* **387**, 39-45 (2015).

Louca, S. & Pennell, M. W. Extant timetrees are consistent with a myriad of diversification histories. *Nature* **580**, 502-505 (2020).

Emerson, B. C. & Kolm, N. Species diversity can drive speciation. *Nature* **434**, 1015-1017 (2005).

Chouteau, M., Arias, M. & Jorun, M. Warning signals are under postive frequency-dependent selection in nature. *Proceedings of the National Academy of Sciences* **113**, 2164-2169 (2016).

Fisher, R. A. *The genetical theory of natural selection*. (The Clarendon press, 1930).

Lande, R. Models of speciation by sexual selection on polygenic traits. *Proceedings of the National Academy of Sciences of the United States of America-Biological Sciences* **78**, 3721-3725 (1981).

Kirkpatrick, M. Sexual selection and the evolution of female choice. *Evolution* **36**, 1-12 (1982).

Nosil, P., Feder, J. L., Flazman, S. M. & Gompert, Z. Tipping points in the dynamics of speciation. *Nature Ecology and Evolution* **1**, 0001 (2017).

Scheffer, M. *Critical Transitions in Nature and Society*. (Princeton University Press, 2009).

Sanderson, S. *et al.* The pace of modern life, revisisted. *Molecular Ecology* **31**, 1028-1043 (2021).

Scholl, J. P. & Wiens, J. J. Diversification rates and species richness across the tree of life. *Proceedings B* **283**, 20161334 (2016).

Kinnison, M. T. & Hendry, A. P. The pace of modern life II: from rates of contemporary microevolution to pattern and process. *Genetica* **112**, 145-164 (2001).

48 Svensson, E. I. Eco-evolutionary dynamics of sexual selection and sexual conflict.
Functional Ecology **33**, 60-72 (2019).
49 Clapham, M. E. & Karr, J. A. Environmental and biotic controls on the evolutionary
history of insect body size. *Proceedings of the National Academy of Sciences* **109**, 10927-
10930 (2012).
50 Losos, J. B., Warheitt, K. I. & Schoener, T. W. Adaptive differentiation following
experimental island colonization in Anolis lizards. *Nature* **387**, 70-73 (1997).
51 Reznick, D., Shaw, F. H., Rodd, F. H. & Shaw, R. G. Evaluation of the Rate of Evolution
in Natural Populations of Guppies (Poecilia reticulata). *Science* **275**, 1934-1937 (1997).
52 Svensson, E. I. The speed of life-history evolution. *Trends in Ecology and Evolution* **12**,
380-381 (1997).
53 Simpson, G. G. *The Major Features of Evolution*. (Columbia University Press, 1953).
54 Gould, S. J. *Wonderful Life: The Burgess Shale and the Nature of History*. (W.W.
Norton & Co., 1989).
55 Cooney, C. R. *et al.* Mega-evolutionary dynamics of the adaptive radiation of birds.
Nature **542**, 344-347 (2017).
56 Voje, K. L. Tempo does not correlate with mode in the fossil record. *Evolution* **70**, 2678-
2689 (2016).
57 Hansen, T. F. & Houle, D. in *The Evolutionary Biology of Complex Phenotypes* (eds M.
Pigliucci & K. Preston) (Oxford University Press, 2004).
58 Voje, K. L., Starrfelt, J. & Liow, L. H. Model adequacy and microevolutionary
explanations for stasis in the fossil record. *The American Naturalist* **191**, 509-523 (2018).
59 Hunt, G. The relative importance of directional change, random walks, and stasis in the
evolution of fossil lineages. *Proceedings of the National Academy of Sciences of the*
United States of America **104**, 18404-18408 (2007).
60 Eldredge, N. *et al.* The dynamics of evolutionary stasis. *Paleobiology* **31**, 36-55 (2005).
61 Futuyma, D. J. Evolutionary constraint and ecological consequences. *Evolution* **64**, 1865-
1884 (2010).
62 Kingsolver, J. G. *et al.* The strength of phenotypic selection in natural populations.
American Naturalist **157**, 245-261 (2001).
63 Felsenstein, J. Phylogenies and quantitative characters. *Annual Review of Ecology and*
Systematics **19**, 445-471 (1988).
64 Lynch, M. The rate of morphological evolution in mammals from the standpoint of the
neutral expectation. *The American Naturalist* **136**, 727-741 (1990).
65 Felsenstein, J. Phylogenies and the comparative method. *American Naturalist* **125**, 1-15
(1985).
66 Hansen, T. F. & Pélabon, C. Evolvability: A quantitative genetics perspective. *Annual*
Review of Ecology and Systematics **52**, 153-175 (2021).
67 Lande, R. Natural selection and random genetic drift in phenotypic evolution. *Evolution*
30, 314-334 (1976).
68 Revell, L. J. Phytools: an R package for phylogenetic comparative biology (and other
things). *Methods in Ecology and Evolution* **3**, 217-223 (2012).
69 Hoehna, S. Fast simulation of reconstructed phylogenies under global time-dependent
birth-death processes. *Biorinformatics* **29**, 1367-1374 (2013).

423
424

Fig. 1. Evolutionary rate-time relationships reflect null or variable underlying relationships between evolutionary change and time. Panels **A-E** show evolutionary rate plotted against the timescale of measurement [6], with dashed red lines showing the least squares fit of the function *rate = a*time*[b]. Panels **F-J** plot the amount of evolutionary change (i.e., a rate numerator) against time frame for the same datasets, with dashed red lines showing the linear piecewise regression, if statistically significant. Panels **A** and **F** show vertebrate body size evolution [29], Panels **B** and **G** show diversification of genera from fossil lineages (plants, mammals, and marine animals)[8], Panels **C** and **H** show diversification data estimated from time-calibrated molecular phylogenies [8], while Panels **D** and **I** show a larger database of phylogenetic diversification rates [46] where rates were inferred directly from age and richness data [22]. Panels **E** and **J** show within-generation morphological changes (mainly from fish harvesting). All nonlinear negative relationships plotted in **A-E** were statistically significant, whereas relationships in **F-J** were variable, with statistically significant breakpoints found for morphology, fossil lineage accumulation, and one estimator of phylogenetic diversification, and a positive linear relationship for another estimator

439   of phylogenetic diversification. No significant relationship was found for the within-generation
440   changes in panel **J**.

456
457
Fig. 2. Extraordinary relationships between the amount of evolution and time are required
to produce positive rate-time relationships. Panel **A** shows the relationship between rate and
time for various functions of the rate numerator and time, which are plotted in the same color in
Panel **B**. The black line is a reference point, where there is a null relationship between rate and
time. We assume for this example a form of the relationship, *amount of evolution = time^b*, where
the function was plotted for b ranging from -2 (blue) to 2 (red). Similar general patterns are
observed for other functions of change and time.
465
466

Fig. 3. Simulations of constant-rate evolutionary models reveal persistence of spurious negative rate-time relationships. Panels **A** and **D** show a simulation of phenotypic evolution under random genetic drift in the absence of natural selection, which produces a Brownian motion process [63] where variance among lineages accumulates linearly with time (**A**). Although the rate of drift is constant (**D** inset), we will nevertheless obtain a negative scaling relationship between the lineage-specific per-generation rate of evolution (**D**).   Panels **B** and **E** show a simulation of phenotypic evolution under drift and stabilizing natural selection, an Ornstein Uhlenbeck process [67]. This process generates no relationship between time and divergence (**B**), but again a strong negative scaling relationship with per-lineage per-generation evolutionary rate and time and hence a similar spurious negative relationship(**E**). Panels **C** and **F** show a constant-rate pure-birth (i.e., no extinction) lineage diversification process, where diversity accumulates approximately linearly with time (**C**). This scenario also produces a similar spurious a negative scaling relationship between the rate estimator [22] and time (**F**). That this relationship is a

482 statistical artifact with no need for any evolutionary explanation can be seen by fitting a log-
483 linear model to the last 15 million years of the simulated data, which shows there is no
484 relationship between rate and time for this time interval (**F** inset).

Methods

Data and code availability: All data is already publicly available, and we provide our

complete R script as a supplemental file.

<u>Data.</u> We analyze existing data on rates of vertebrate body size evolution [29], which consists of

data on log-fold change in body size as well as time interval over which this change occurred.

For phylogenetic lineage diversification, we analyze two datasets, one in which diversification

rates were estimated from the branching pattern of species-level phylogenies [8], and another

larger dataset [46] where diversification rates were estimated from clade diversity and age

information. For fossil data, we analyze an existing dataset [8] of emergence rates of new genera.

For each dataset, we plot the standard measure of rate against the timeframe over which that rate

was measured, fitting exponential linear models by nonlinear least squares [6]. We also plot a

measure of the rate numerator against time for each of these datasets, and fit piecewise linear

regressions to identify significant linear relationships between amount of evolution and time, as

well as any potential shifts in the relationship between evolution and time. For each dataset, we

fit nonlinear models of the form $rate = a*time^b$ using nonlinear least squares (nls()) in R, as well

as piecewise regression using the segmented package. We compared models fits of an ordinary

linear regression to a model with a single breakpoint, and also when appropriate to a null model

containing only an intercept, using the anova() function in R. Data presented in Figure 1 were

obtained from ref 6 (available at https://github.com/fhenaod/evol_time_scaling/tree/main/data),

ref 24 (Dryad doi:10.5061/dryad.7d580), ref 40, and ref 39 (database available at:

https://proceeddatabase.weebly.com/).

<u>Simulation</u> We simulated three different constant rate evolutionary process. For morphological

evolution, we simulated random genetic drift for 1-1000 generations assuming a heritability of

0.50 and a population size of 100. We also simulated evolution under genetic drift and stabilizing

selection, with the same demographic and genetic assumptions but assuming weak selection (ω^2

= 15) towards an optimum. Each simulation was started with the population at its optimum. For

morphological evolution, we plotted rates in units of Haldanes [4], which was natural as we

simulated evolution on the scale of phenotypic standard deviations. For lineage diversification,

we simulated pure-birth phylogenies with a constant speciation rate of 0.15, corresponding to a

expected number of 2.3 taxa at timepoint 1 under a continuous time pure birth process. We

generated trees with ages ranging from 1-35 million years, and plot the maximum likelihood

estimator of the pure-birth diversification rate, $\log(N)/t$ [22]. Our simulations of phenotypic

evolution were performed using nested for() loops in R and parameterized as described above.

Our simulations of constant rate pure birth phylogenetic diversification was performed using the

pbtree() function in the phytools package [68]. Our simulations of time-dependent pure birth

diversification (Figure S1) were performed using the TESS package [69].

553 **Acknowledgments:** We thank Masahito Tsuboi, Chris Cooney, Kjetil Voje, and Miguel Gomez

554   for comments on the manuscript, and Charlie Conrwallis for discussion of these issues.

555   **Funding:** This work was funded by grants from the Swedish Research Council (VR:

556   grants no. 2019-03706 to SPD and no.r 2020-03123 to EIS), The Crafoord Foundation

557   (20220602), Formas (grant 2021-01096), to SPD

558 **Author contributions:** SPD and EIS conceived of the problem and designed the study, SPD

559   performed the analysis and drafted the manuscript. Both authors wrote the manuscript.

560 **Competing Interest Declaration:** The Authors have no competing interests to declare

Fig. S1. Accelerating time-dependent diversification produces negative rate scaling

patterns. Panels show simulations of time-dependent pure birth diversification, repeated for

different timeframes from 1-35 million years under the same time-dependent diversification

model. In each simulation birth rates were assumed to be linearly related to time, $\lambda \sim 0.008*time$,

extinction rate $\mu = 0$. Panel **A** shows log of total diversity plotted against time for each

simulation run. Panel **B** shows the estimate of net speciation rate $\log(N)/t$ as a function of time.

570   Qualitative conclusions remain unchanged under different parameter values, with the one salient

571   feature being that the expected positive rate-time relationship is only recovered in deep time.