Structural basis of the mechanism and inhibition of a human ceramide synthase

Tomas C. Pascoa[1,†], Ashley C.W. Pike[1], Christofer S. Tautermann[2], Gamma Chi[1], Michael

Traub[2], Andrew Quigley[1,4], Rod Chalk[1], Saša Štefanic[3,5], Sven Thamm[2], Alexander Pautsch[2],

Elisabeth P. Carpenter[1,†], Gisela Schnapp[2,†], David B. Sauer[1,†]

[1] Centre for Medicines Discovery, Nuffield Department of Medicine, University of Oxford,

Oxford, UK

[2] Boehringer Ingelheim Pharma GmbH & Co KG, Birkendorfer Straße 65, 88397, Biberach,

Germany

[3] Institute of Parasitology, Vetsuisse and Medical Faculty, University of Zürich, Zürich,

Switzerland

[4] Present address: Membrane Protein Laboratory, Research Complex at Harwell, Diamond

Light Source Ltd, Harwell Science and Innovation Campus, Didcot, UK

[5] Present address: Nanobody Service Facility, University of Zurich, AgroVet-Strickhof, 8315,

Lindau, Switzerland

[†] Correspondence to: T.C.P. (tomas.pascoa@cmd.ox.ac.uk), E.P.C. (lizcarpen1@gmail.com),

G.S. (gisela.schnapp@boehringer-ingelheim.com) or D.B.S. (david.sauer@cmd.ox.ac.uk)

Abstract

Ceramides are bioactive sphingolipids that play pivotal roles in regulating cellular

metabolism. Ceramides and dihydroceramides are synthesized by a family of six ceramide

synthase enzymes (CerS), each with distinct specificity for the acyl-CoA substrate.

Importantly, the acyl chain length plays a key role in determining the physiological function

of ceramides, as well as their role in metabolic disease. Ceramide with an acyl chain length of

16 carbons (C16 ceramide) has been implicated in obesity, insulin resistance and liver

disease, and the C16 ceramide-synthesizing CerS6 is regarded as an attractive drug target for

obesity-associated disease. Despite their importance, the molecular mechanism underlying

ceramide synthesis by CerS enzymes remains poorly understood. Here, we report cryo-

electron microscopy structures of human CerS6, capturing covalent intermediate and product-

bound states. These structures, together with biochemical characterization using intact protein

and small molecule mass spectrometry, reveal that CerS catalysis proceeds via a ping-pong

reaction mechanism involving a covalent acyl-enzyme intermediate. Notably, the product-

bound structure was obtained upon reaction with the mycotoxin fumonisin B_1, providing new

insights into its inhibition of CerS. These results provide a framework for understanding the

mechanisms of CerS function, selectivity, and inhibition, and open new directions for future

drug discovery targeting the ceramide and sphingolipid pathways.

Introduction

Ceramides are the precursors for the synthesis of complex sphingolipids and are bioactive

lipids with crucial roles in signaling. In particular, ceramides have been proposed to act as

key metabolic sensors to promote fatty acid utilization and storage during excessive fatty acid

availability[1]. However, abnormal ceramide accumulation is associated with metabolic

dysfunction, and elevated levels of ceramides have been observed in obesity-related

metabolic disorders such as diabetes, non-alcoholic fatty liver disease (NAFLD) and non-

alcoholic steatohepatitis (NASH)[2-4].

Ceramides are composed of a sphingosine (d18:1) long-chain base with an N-linked acyl

chain, the length of which is critical to the lipids' biological functions and roles in

pathophysiology[5]. For example, C16:0 ceramide is the most common ceramide in adipose

tissue and its levels are elevated in this tissue of obese humans[2]. In addition, insulin

resistance is correlated with plasma C16:0 and C18:0 ceramides, subcutaneous adipose tissue

C16:0 ceramides, and hepatic C16:0 and C18:0 ceramides[6-8]. Moreover, the levels of total

ceramides and C16:0, C22:0 and C24:1 dihydroceramides were found to be elevated in the

liver of insulin-resistant patients with NASH[3].

In mammals, *de novo* synthesis of ceramides is preceded by synthesis of dihydroceramides

through the N-acylation of the sphingoid long-chain base sphinganine (dihydrosphingosine;

d18:0) by one of six ceramide synthases (CerS1-6)[9]. The resulting dihydroceramides are

subsequently converted into ceramides by one of two dihydroceramide Δ4-desaturases

(DES1-2)[10]. Alternatively, CerS can directly re-acylate recycled sphingosine, in the salvage

pathway[11]. Of these enzymes, recent observations from human and mouse studies have

highlighted CerS6 as an attractive target for treating obesity-associated metabolic disease,

including type 2 diabetes and NASH[2,5,12-14]. Deletion of *CerS6* in mice granted protection

against diet-induced obesity, steatohepatitis, and insulin resistance[2,14], and liver-specific

deletion improved glucose tolerance and mitochondrial morphology[14]. Strikingly, this effect

was shown to only occur in CerS6$^{\Delta/\Delta}$ mice but not in CerS5$^{\Delta/\Delta}$ mice, though both show a

preference for C16-CoA and therefore primarily produce C16 ceramides[15] and hepatic C16:0

ceramide levels were reduced in both knock-outs[14]. This surprising effect was found to be a

result of the specific interaction of CerS6-derived C16:0 sphingolipids in mitochondria with

the mitochondrial fission factor (Mff)[14]. This revealed that the subcellular localization of

ceramide production can lead to drastically different physiological outcomes[14]. In further

support of the therapeutic potential of inhibiting CerS6, ablation of the protein's expression in

an obese insulin resistance mouse model led to reduced body fat and improved oral glucose

tolerance and insulin sensitivity[12].

Central to its enzymatic function, CerS contain a TRAM-Lag1-CLN8 (TLC) homology

domain[16], which includes a 52-residue Lag1p motif containing conserved histidines and

aspartates required for activity[17,18]. In addition, CerS2-6 contain a non-essential Hox-like

domain between TM1-2[19], flanked by two essential and conserved positively charged

residues[20]. The activity of each CerS is subject to complex regulation, which includes

phosphorylation[21], glycosylation[22], dimerisation[23] and other protein-protein interactions[24,25].

The six mammalian CerS also have different acyl chain-length specificities and tissue

expression patterns, further influencing the tissue-specific distribution of the different

ceramides[5,9,26]. However, the molecular mechanisms underlying synthesis and regulation of

ceramides and dihydroceramides remain unclear.

Despite the interest in pharmacologically reducing CerS6 activity, at present, no specific

inhibitors have been described for this family member. The best characterized CerS inhibitor

is fumonisin B_1 (FB$_1$), the most prevalent member of the fumonisin family of mycotoxins

produced by *Fusarium* species. FB$_1$ is a potential carcinogen and teratogen and therefore of

significant concern as it is a common contaminant in maize, rice, other cereals, and cereal-

based food stocks[27-29]. FB$_1$ is a potent non-selective inhibitor of all human CerS, and is

competitive towards both sphinganine and acyl-CoA substrates[30]. In addition, FTY720

(fingolimod, Gilenya), a prodrug administered to functionally antagonize the sphingosine-1-

phosphate receptor 1 in the treatment of multiple sclerosis, is also a non-selective CerS

inhibitor[31,32]. Further, a non-phosphorylatable analogue of FTY720, P053, has recently been

shown to potently and selectively inhibit CerS1[33], showcasing the possibility of achieving

isoform-specific CerS inhibition.

Here, to understand the molecular basis of ceramide and dihydroceramide synthesis by the

ceramide synthases, we employed a combination of cryogenic electron microscopy (cryo-

EM), intact protein mass spectrometry, and small-molecule high-resolution and tandem mass

spectrometry to probe the catalytic mechanism of human CerS6.

Results

Cryo-EM structure of human CerS6

We recombinantly expressed the human CerS6 protein, truncated at Asp350 to remove the predicted disordered C-terminus, in suspension-adapted HEK293 cells using the BacMam system. The protein was purified in glyco-diosgenin (GDN), yielding a mixture of monomeric and dimeric species on size-exclusion chromatography (Extended Data Fig. 1a-d). While we selected the dimer species for structural studies to maximize particle size and possible symmetry for single-particle cryo-EM, its relatively low molecular weight (84 kDa) is still challenging for current methods. We therefore raised camelid nanobodies against the purified protein and solved the structure of a CerS6 dimer with one nanobody bound per monomer at a nominal resolution of 3.2 Å resolution (Fig. 1a,b; Extended Data Table 1; Extended Data Fig. 2). The CerS6-Nb22 complex was well resolved throughout the membrane-spanning region, but the Hox-like domain was less well-defined. This allowed for unambiguous tracing of CerS6 and Nb22 with the exception of the Hox-like domains, where residues 72-119 were fitted into the envelope of the cryo-EM map as rigid bodies (see Methods) (Extended Data Fig. 2e; Extended Data Fig. 3a).

Figure 1. Cryo-EM structure of human CerS6. a, Cryo-EM map of the CerS6 dimer (blue) with one copy of Nb22 (gray) bound to each CerS6 monomer. **b,** Overall cartoon cylinder representation of the CerS6 dimer structure. One of the monomers is rainbow-coloured from purple (N-terminus) to red (C-terminus). **c,** schematic representation of the CerS6 7-TM helix topology. **d,** Cartoon representation of the transmembrane domain of a CerS6 monomer. The covalent acyl-imidazole species is shown in stick representation (acyl chain: pink carbon atoms), and the Coulombic potential map for this covalent species is shown as a pink transparent surface. The Hox-like domain was omitted for clarity. **e,** Denaturing intact protein MS analysis of purified CerS6 protein, revealing the presence of a covalent modification (+238.45 Da) which matches the expected mass shift corresponding to covalent attachment of a palmitoyl (Palm; C16:0) chain. This adduct peak was present in all purifications tested (n=6).

The cryo-EM structure reveals a CerS6 homodimer with one nanobody bound per monomer on the luminal face. Each monomer has 7 transmembrane helices (Fig. 1b,c), with the dimer interface formed by TM1a, the C-terminal end of TM3, and the TM3-TM4 loop. The TLC

domain (TM2-7) forms a barrel containing two 3-TM units arranged as inverted repeats

(Extended Data Fig. 4a), while TM1 is attached to the outside of the barrel through contacts

with TM2/3. The TM2-7 barrel is assembled around a central cavity, which is open to the

cytoplasm and closed to the ER lumen by a loop between TM6-7. This central cavity has a

large cytoplasmic opening (approximately 23 x 12 Å), creating a vestibule that funnels

towards the midpoint of the membrane, where it forms a ~5 Å wide tunnel that stretches

towards the ER side.

Interestingly, the fold of the TM2-7 barrel of CerS6 resembles that of the 6-TM barrels found

in very-long-chain fatty acid elongase ELOVL7[34] and TMEM120A/B[35,36], as identified by

DALI[37]. Additionally, a structurally homologous 6-TM barrel is predicted in the AlphaFold2

models of 3-hydroxyacyl-CoA dehydratases (HACD1-4) (Extended Data Fig. 4b), which

participate in the same four-enzyme acyl-CoA elongation cycle as ELOVLs[38]. The

observation that different acyl-CoA-modifying enzyme families share a similar 6-TM barrel

structure is intriguing, suggesting that this architecture is likely optimized for the specific

recognition and reaction with acyl-CoA substrates.

CerS6 co-purifies with a covalently bound C16:0 acyl chain

Surprisingly, the cryo-EM density map revealed the presence of a long density extending

from near the membrane mid-point to the occluded ER end of the central cavity, spanning the

entire length of the narrow tunnel (Fig. 1d; Fig. 2a). This density was continuous between

protein and the unknown molecule, suggesting a covalent modification by a lipid. To probe

the presence of covalent adducts, we performed denaturing intact protein mass spectrometry

(MS) analysis of purified protein samples, and identified a +238.45 Da mass shift (Fig. 1e

and Extended Data Fig. 1f,g). Notably, this adduct mass agrees with a covalently attached

C16:0 acyl chain (theoretical: +238.41 Da). Strikingly, the shape and length of the density in

the cryo-EM map is also consistent with a bound C16 acyl chain. A covalent bond is seen

linking the acyl chain to the imidazole Nε of His211 (TM4) (Fig. 2b; Extended Data Fig. 3b),

an absolutely conserved histidine that is required for CerS activity[17,18]. Thus, based on the

extended density in the cryo-EM map and the observed mass adduct, we have modelled a

palmitoyl (C16:0) chain covalently attached to His211.

Figure 2. CerS6 contains an acyl chain-binding tunnel buried deep in the membrane. a, Cutaway

molecular surface representation of the CerS6 transmembrane region, revealing the presence of a long

central cavity spanning the entire length of the protein. The palmitoyl chain covalently attached to

His211 is bound in a narrow tunnel on the ER luminal half of the central cavity. **b**, Coulombic

potential map in the region around the covalent linkage of the acyl chain to His211. **c-e**, CerS6's acyl-

binding tunnel, viewed from the **(c,d)** membrane plane or **(e)** from the ER face. Side-chains lining and

capping the tunnel are shown as sticks. Hydrogen bonds in the active site are shown as dashed lines.

The acyl chain is surrounded in the narrow tunnel primarily by hydrophobic side chains from

TM4, TM5 and TM7 (Fig. 2c-e). The tunnel is sealed at the ER end by the TM6-7 loop (Fig.

2e), revealing how the chain-length preference of CerS6 for C16-CoA is determined by

limiting the number of carbons that can fit in the acyl-binding tunnel. This observation is in

agreement with earlier reports that this loop determines CerS acyl chain specificity[22].

Catalysis by CerS6 proceeds via a ping-pong mechanism

Acyltransferases catalyze two-substrate, two-product reactions that can proceed via either

ternary-complex or double-displacement (ping-pong) mechanisms[39]. In a ternary-complex

mechanism, both the acyl donor and acceptor bind simultaneously and the enzyme catalyzes

the direct acyl transfer from one substrate to the other. In contrast, a ping-pong mechanism

involves two independent steps. Initially, reaction with the first substrate (the acyl donor)

results in the transfer of the acyl chain to a nucleophilic residue, forming a covalent acyl-

enzyme intermediate. This acyl chain is then transferred to the second substrate (the acyl

acceptor) in the second step of the reaction.

The CerS6 structure reveals that there is not enough space for both substrates to bind in the

central cavity and access the active site at the same time, thereby indicating a ternary-

complex mechanism is unlikely. Furthermore, the C16:0 chain covalently attached to the

conserved His211 fits with CerS6's selectivity for palmitoyl-CoA and suggests that this

species could correspond to the acyl-enzyme intermediate of a ping-pong type reaction

mechanism. To investigate whether the acylated enzyme species corresponds to a real

catalytic intermediate, we incubated the purified protein with the enzyme's second substrate,

the long-chain base sphinganine (Fig. 3a). Using denaturing intact protein MS, we observed

the complete loss of the C16:0-modified protein after sphinganine addition (Fig. 3b and

Extended Data Fig. 5a). Furthermore, high-resolution mass spectrometry (HRMS) revealed

that incubation of the acylated protein with sphinganine led to the production of C16:0

dihydroceramide (observed *m/z*: 540.5380; theoretical *m/z*: 540.5350; mass error: +5.55 ppm)

(Fig. 3c). Overall, these results clearly demonstrate that sphinganine reacts with the acyl-

enzyme intermediate to de-acylate the enzyme and generate the expected reaction product of

dihydroceramide, thus supporting a ping-pong reaction mechanism for CerS6. Taken together

with the structural data, this implies that His211 acts as the nucleophile in the first step of the

reaction.

Figure 3. Unraveling the catalytic mechanism of CerS6 using mass spectrometry. a, chemical

structures of the natural long-chain base substrate sphinganine (dihydrosphingosine) and the drug

FTY720 (fingolimod, Gilenya). The homologous primary amines are highlighted in salmon. **b,** Intact

mass analysis of protein samples after incubation in the absence of substrates (black) or in the

presence of sphinganine (green) (n = 3 biological replicates; see Extended Data Fig. 5 for replicate

traces). Deconvoluted mass peaks are indicated as follows: unmodified enzyme, orange icon; covalent

acyl-enzyme species, orange and pink icon; background species present in all traces, gray circle. **c,**

LC-HRMS detection of the reaction products. Extracted ion chromatograms (EICs) for the expected

[M+H]$^+$ ions of each of the reaction products after incubation of the acyl-enzyme intermediate with

sphinganine (C16:0 dihydroceramide, green) or FTY720 (N-palmitoyl FTY720, orange). EICs

obtained after incubation in the absence of substrates are shown in black. **d,** CerS6's active site,

viewed from the plane of the membrane. The central cavity is shown as a transparent gray surface,

highlighting the presence of a side pocket (highlighted in green) adjacent to the acyl carbonyl.

Hydrogen bonds are shown as dashed lines. **e,** Mutational analysis of the CerS6 active site by

comparison of the C16:0 dihydroceramide synthase activity of wild-type (WT) and active site

mutants. n=3 independent biological replicates. Gray bars correspond to the mean of the biological

replicates. Data points represent each biological replicate, corresponding to the mean of four technical

replicates. Error bars show the SEM.

Notably, the CerS substrate sphinganine and FTY720 are chemically similar (Fig. 3a), with

potential clinical consequences to drug pharmacokinetics as 15% of FTY720 becomes N-

acylated in human subjects[40]. Although the major N-acyl FTY720 metabolites identified were

N-stearoyl and N-2-hydroxystearoyl FTY720, there is evidence suggesting that a small

amount of N-palmitoyl (C16:0) FTY720 is formed[40]. Which enzyme catalyzes the N-

acylation of FTY720 has not been determined, but a putative role for CerS enzymes or the

reverse reaction of ceramidases has been proposed[40]. Incubation of acyl-CerS6 with FTY720

results in a small increase in the protein's thermostability, suggesting it binds to the purified

enzyme (Extended Data Fig. 1e). To investigate whether purified CerS6 can N-acylate

FTY720 *in vitro*, we incubated the enzyme with this compound, and subsequently identified

C16:0 FTY720 in the reaction mixture (observed *m/z*: 546.4869; theoretical *m/z*: 546.4881;

mass error: -2.20 ppm) (Fig. 3c), the expected product of a reaction between FTY720 and the

CerS6 bound acyl chain. The proposed structure of this C16:0 FTY720 reaction product was

further validated based on the product ion MS/MS spectrum of its [M+H]$^+$ ion (Extended

Data Fig. 6). However, no obvious loss of the acyl-enzyme mass peak was observed when the

reaction mixture was analyzed by denaturing intact protein MS (Extended Data Fig. 5b),

suggesting that FTY720 is a poor acyl acceptor for acylated CerS6.

Highly conserved residues line the catalytic site

CerS6's Lag1p motif (Arg202-Tyr253) (TM4-5) contains the highly conserved

His211/His212 (TM4) and Asp239/Asp242 (L5a-b/TM5), which have been previously

suggested to be located in the catalytic site and have been shown to be essential for function

in mammalian CerS[17] and *S. cerevisiae* Lag1p[18].

Examining the CerS6 structure, we noted these residues line the central cavity near the mid-

point of the membrane (Fig. 3d). Notably, the co-purified covalently linked C16:0 chain is

covalently attached to the first histidine of the Lag1p motif (His211) (Fig. 2a-d), which

appears to act as the nucleophile in the first step of the reaction and is required for CerS6

function (Fig. 3e). Although the observation that the mechanism of CerS6 involves a

nucleophilic histidine was surprising - as cysteines or serines are more commonly employed

as nucleophiles - there are multiple well-documented examples of other enzymes using

nucleophilic histidines in their catalytic mechanisms[34,41-45].

Histidine nucleophiles are typically oriented and activated by hydrogen-bonding between a

proton at Nδ of the imidazole ring and a hydrogen bond acceptor, ensuring that Nε remains

unprotonated[34,42,46]. Examining our structure, we noted the proximity of His212 to the

acylated His211 in CerS6, which immediately suggested this residue is the necessary

hydrogen bond acceptor. While the Nδ of His212 is located 3.84 Å away from the Nδ of

His211, it is plausible that the two side chains may interact in the acyl-CoA-bound state.

Indeed, we found that mutation of His212 to alanine results in the complete loss of activity

(Fig. 3e). Residues equivalent to His212 of CerS6 are also essential in the homologous mouse

and human CerS1[17,47] and yeast Lag1p[18]. This demonstrates the second histidine plays a key

role in catalysis, and explains the pathogenic effect of mutating the equivalent His of CerS1

to a Gln in patients with progressive myoclonic epilepsy and dementia[47]. Moreover, we have

recently demonstrated that the reaction of human ELOVL7 with acyl-CoA substrates also

involves a double histidine motif, where the first histidine acts as a nucleophile and is

activated by hydrogen-bonding to the second[34]. Strikingly, structural alignment of CerS6 and

ELOVL7 places the histidine pairs in equivalent positions (Extended Data Fig. 4c),

suggesting that these residues may play similar roles in the first step of their respective

reactions.

Formation of the acyl-enzyme intermediate proceeds via an oxyanion transition state, which

is usually stabilized by hydrogen bonding. Within the acyl-CerS6 structure, the carbonyl

oxygen of the covalently attached C16:0 chain is within hydrogen-bonding distance (2.8 Å)

of the Nδ of Asn315 (TM7). Based on the proximity of these moieties, we hypothesized that

this interaction could stabilize the formation of the oxyanion transition state during catalysis.

Consistent with this notion, an Asn is present at the equivalent position in CerS1, and CerS2-

5, Lag1p and Lac1p have a His at this position (Extended Data Fig. 7) which may likewise

act as a hydrogen-bond donor. Indeed, we found that an Asn315His mutant preserves activity.

Surprisingly, however, mutation of Asn315 to either an Ala or an Asp did not result in loss of

activity (Fig. 3e). It is unclear if the hydrogen bond in the wild-type structure is not required

for function, or if these mutations allow a water molecule to enter the reaction site and act as

a substitute hydrogen bond acceptor. Asn315 is involved in a hydrogen-bonding network

involving the conserved Arg275 (L6a-b) and Asp242 (TM5b). We found that an Arg275Lys

mutant retained only approximately 20 % of wild-type activity (Fig. 3e), suggesting that the

interactions established by the guanidino group of Arg275 to align neighbouring residues in

the active site are important for function.

There is a third histidine on TM5a which is conserved in mammalian CerS but is a

methionine in yeast Lag1p and Lac1p, and is therefore of ambiguous importance to the

enzymes' reaction cycle. In the CerS6 structure this residue, His238, is located above the

acyl-imidazole carbonyl and lays against the hydrocarbon chain (Fig. 3d). We found that

mutation of this His238 to alanine reduced activity by over 50 % (Fig. 3e), and therefore

suggest this residue may participate in aligning the acyl chain.

Finally, we noticed the presence of a side pocket in the active site of CerS6, adjacent to

His211 (Fig. 3d) and lined by Tyr142 (TM2), Tyr189 (TM3), His212 (TM4), Asp239 (L5a-

b), Asp242 (TM5b) and Arg275 (L6a-b). The position and chemistry of this pocket relative to

the carbonyl of the acyl-enzyme intermediate suggests this is likely to be the site where the

amino alcohol moiety of the second (long-chain base) substrate binds. Supporting this notion,

the residues of this pocket are highly conserved (Extended Data Fig. 7). This pocket therefore

likely ensures the primary amine of the long-chain base is ideally located for a nucleophilic

attack on the carbonyl carbon of the acyl-imidazole intermediate in the second step of the

reaction.

Structure of N-acyl fumonisin B_1-bound CerS6

Having captured the intermediate state of the CerS6 enzyme, we next set out to examine the structural basis of CerS inhibition by FB_1. To this end, we incubated purified acyl-CerS6 with fumonisin B_1, and determined the complex's structure to a nominal resolution of 2.95 Å (Fig. 4a,b and Extended Data Fig. 8). In comparison with the covalent intermediate state, the overall structure of each CerS6 monomer did not change (transmembrane region backbone RMSD = 0.39 Å), although the flexible Hox-like domains rotated by approximately 10°.

The cryo-EM map of the putative FB_1-bound complex contains a long, continuous non-protein density starting within the ER luminal end of the narrow tunnel and extending to the cytoplasmic entrance to the central cavity (Fig. 4a and Extended Data Fig. 3c,d). The observed density is too long to be FB_1 alone. Notably, incubation of palmitoyl-CerS6 with FB_1 completely removed the acyl group from the enzyme (Fig. 4c and Extended Data Fig. 5a and generated C16:0-fumonisin B_1 (observed m/z: 960.6260; theoretical m/z: 960.6254; mass error: +0.62 ppm) (Fig. 4d). Noting the covalent bond between the enzyme and the palmitoyl group has been replaced by a bond to the toxin, we modeled this density as C16:0-FB_1 (Fig. 5a and Extended Data Fig. 3c). These results indicate FB_1 has reacted with the covalent acyl-enzyme intermediate, releasing His211 and generating the N-acyl FB_1 product. This product remains bound to the enzyme, thus acting as an inhibitor. This finding is consistent with earlier reports that CerS enzymes can N-acylate FB_1 in cultured cells and in rats[48,49].

Figure 4. Cryo-EM structure of CerS6 in complex with N-palmitoyl fumonisin B_1. a, Cartoon representation of CerS6 with bound N-palmitoyl fumonisin B_1 (shown as sticks; cyan carbon atoms). The cryo-EM density of the bound product is shown as a transparent cyan surface. **b**, Cutaway molecular surface representation, revealing that the N-palmitoyl fumonisin B_1 species occupies the entire length of the central cavity. The Hox-like domain has been omitted for clarity. **c**, Intact mass analysis of protein samples after incubation in the absence of substrates (black) or in the presence of the mycotoxin fumonisin B_1 (blue) (n = 3 biological replicates; see Extended Data Fig. 5 for replicate traces). **d**, LC-HRMS detection of N-palmitoyl fumonisin B_1. The EIC for its expected $[M+H]^+$ ion is shown after incubation of the acyl-enzyme intermediate with fumonisin B_1 (blue) or in the absence of the toxin (black). *Inset:* chemical structure of fumonisin B_1. Its primary amine (salmon circle) and tricarballylic acid groups (gray circles) are highlighted.

The N-acyl FB_1 species spans the entire length of the central cavity, with the acyl chain remaining bound in the narrow tunnel at the occluded ER end of the cavity, and the FB_1 portion sitting in the wider cytoplasmic entrance (Fig. 4b and Fig. 5b-c). The newly formed amide bond sits in the active site (Fig. 5c), with the carbonyl oxygen forming a hydrogen bond to Tyr189 (TM3), and the amide NH hydrogen bonding to the Nε of His211 (TM4). The

3-hydroxyl group of FB$_1$, which is also present in sphingoid base substrates, forms a

hydrogen bond with the strictly conserved and catalytically essential Asp242 (TM5b)[17,18].

Such an interaction with the 3-hydroxyl group of sphingoid base substrates would optimally

place the sphingoid primary amine to attack the carbonyl carbon. However, further work is

required to investigate this hypothesis.

Below the active site, the cytoplasmic vestibule of the central cavity has both polar and non-

polar surfaces, where we noted that the hydrocarbon chain of FB$_1$ interacts with the long

hydrophobic face formed by TM5-7 (Fig. 5b,d). This hydrocarbon chain lies against the

conserved Trp318 of TM7 just beneath the active site. Mutation of the corresponding residue

in CerS5 to an alanine has been previously shown to abolish enzymatic activity[50]. In contrast,

introducing a phenylalanine at this position by a W318F mutation in CerS6 retained

approximately wild-type levels of activity (Fig. 3e). From these results, we hypothesize the

conserved Trp orients the sphingoid base substrate to place its amino alcohol moiety in the

side pocket within the active site for the second step of the reaction.

The face of the central cavity opposite the FB$_1$ hydrocarbon binding surface displays a polar

character, and is lined by conserved charged residues that may participate in CoA binding

(Fig. 5b,d). Accordingly, the positive charges provided by Lys125 and Arg131, located at the

cytoplasmic entrance to the cavity, adjacent to the Hox-like domain, have been previously

demonstrated to be important for the function of CerS5 and CerS6[20]. Furthermore, mutation

of Lys131 in CerS3, equivalent to CerS6's Arg131, to alanine has been implicated in

autosomal recessive congenital ichthyosis (ARCI)[51], a disorder of epidermal keratinization

which can be caused by loss of $\geq$ C26 ceramides due to mutations in *CERS3*[52,53].

The two tricarballylic acid (TCA) groups of FB$_1$ span the width of the cytoplasmic end of the

central cavity, where one of the groups interacts with Arg131 (TM2) and the other with

Lys328 (TM7) (Fig. 5b). Thus, through these charge-charge interactions, FB_1 links the first

helix of the first 3-TM bundle and the last helix of the second. Supporting the notion that FB_1

acts to restrain the structure, atomistic molecular dynamics (MD) simulations of covalent

intermediate and N-acyl FB_1-bound CerS6 monomers indicate that a bound N-palmitoyl FB_1

reduces the overall flexibility of the protein, including TM6, TM7, and the Hox-like domain

(Extended Data Fig. 9a-c). Moreover, while Lys125 is not resolved in the cryo-EM maps, in

our simulations Lys125 forms a salt bridge with the same TCA group that contacts Arg131

(Extended Data Fig. 9d). Further, the simulations show that this TCA also contacts Lys249

(TM5b), Tyr253 (TM5b), as well as Arg118 and Arg121 of the last helix of the Hox-like

domain. Therefore, FB_1 appears to mimic a number of CoA interactions through the binding

of one of its TCA moieties at the polar face of the cytoplasmic cavity. Finally, on the

opposite side of the cavity, the other TCA moiety can form salt bridges with Lys328 or

Arg332 (TM7) (Extended Data Fig. 9d). Overall, our cryo-EM structure and simulations

reveal that FB_1 is anchored by polar interactions at opposite sides of the 6TM barrel, which

likely hinder product release, arresting the enzyme in an inhibitory product-bound state.

Figure 5. Binding mode of N-palmitoyl fumonisin B$_1$. a, Close-up view of the CerS6 active site in
the covalent acyl-enzyme intermediate and N-palmitoyl FB$_1$-bound states, showing the transfer of
the palmitoyl chain from His211 to the toxin. **b**, Cytoplasmic portion of the central cavity, viewed
from the membrane plane. Residues lining the cavity are shown as sticks. Polar interactions between
the carboxylates of the tricarballylic acid groups of fumonisin B$_1$ and positively charged residues on
TM2 and TM7 are shown as dashed lines. **c**, Active site, viewed from the plane of the membrane. **d**,
Polar and non-polar surfaces on the cytoplasmic half of the central cavity. Cutaway molecular
surface view, showing that the hydrocarbon chain of fumonisin B$_1$ interacts with the large
hydrophobic face formed by TM5-7.

Bound lipid

In examining our CerS6 cryo-EM maps, we noted both conditions clearly contain an ordered

non-protein density near the dimer interface, adjacent to the LH-TM1 loop and TM4 and

likely corresponding to a bound lipid. There, we modelled a phosphatidylcholine molecule, as

its head group is consistent with the shape, size and environment of the observed density

(Extended Data Fig. 3e). The choline head group is nestled between conserved Trp9 and

Trp21 in a classic aromatic box for binding quaternary amines[54]. Adjacent to the choline-

binding site is Trp14, which is strictly conserved in CerS2-6. Interestingly, mutation of the

corresponding tryptophan residue in CerS3 to an arginine has been found in patients with

ARCI, and the mutant protein was found to be catalytically inactive[52]. In the CerS6 structure,

Trp14 does not directly contact the choline head group, but it could play a structural role in

forming the lipid-binding site.

Discussion

Ceramide synthases carry out an essential step in sphingolipid biosynthesis and are promising

drug targets. Yet, our limited understanding of their structures and biochemical mechanism

has hindered their therapeutic targeting. Here, we provide structural snapshots of CerS6 at

two stages of its reaction cycle, revealing its ping-pong (double-displacement) reaction

mechanism which uses a histidine nucleophile that attacks the acyl-CoA thioester to form a

stable acyl-imidazole intermediate. This intermediate subsequently reacts with the sphingoid

long-chain base substrate to yield the final ceramide or dihydroceramide reaction product

(Fig. 6).

Figure 6. Proposed double-displacement (ping-pong) mechanism of ceramide synthases.
Initially, the acyl-CoA substrate binds with the acyl chain buried deep within the central tunnel and
the CoA moiety sitting near the cytoplasmic entrance to the central cavity. In the first step, the
nucleophilic attack of His211 on the acyl-CoA thioester carbonyl results in thioester cleavage,
covalent acyl-imidazole intermediate formation, and release of CoA. Subsequently, the long-chain
sphingoid base substrate binds, with its hydrocarbon chain interacting with the hydrophobic face of
the central cavity and its amino alcohol moiety sitting in the side pocket in the active site. In the
second step of the reaction, the primary amine of the long-chain base attacks the acyl-imidazole
intermediate, leading to covalent intermediate breakdown and formation of the final N-acyl
sphingoid base (ceramide) product.

Although CoA and the hydrocarbon chain of long-chain bases likely interact at distinct faces
of the cytoplasmic vestibule, there is not enough space for the acyl-CoA and long-chain base
substrates to bind at the same time. Therefore, upon thioester cleavage and covalent
intermediate formation, the CoA product needs to exit the central cavity before the long-chain
base substrate can bind. We hypothesize that the amino alcohol group of the long-chain base
would bind in a discrete side pocket lined by polar residues conserved across the entire CerS
family. This long-chain base is further oriented for the second step of the reaction by
hydrophobic residues located at the entrance to this pocket, particularly by the conserved
Trp318 (TM7). The primary amine of the long-chain base then attacks the acyl-enzyme
intermediate to form the final reaction product. Release of this product into the membrane

likely occurs between two transmembrane helices, and the two halves of the 6TM barrel

immediately suggest an egress route between TM4 and TM7. Indeed, we found that the TCA

groups of N-palmitoyl FB_1 attach to both halves of the barrel and restrict the movement of

TM7, suggesting these interactions likely prevent that product's release. Indeed, this

proposed role for the TCA groups of FB_1 is consistent with previous reports that the

hydrolysed form of FB_1, lacking the TCA moieties, is a weaker CerS inhibitor[55]. However,

FB_1 is competitive towards both substrates[30]. This suggests FB_1's mode of inhibition of CerS

is likely to be multifaceted, sterically blocking substrate binding to the apo state, and locking

the protein in a product bound state as shown here.

Recently, Zelnik et al. postulated that CerS2, and ceramide synthases more generally, use a

ternary complex reaction mechanism[50]. In this model, the substrates bind orthogonally, and

the long-chain base would access the active site through a side entrance located near the

membrane mid-point, where it would bind parallel to the membrane plane. This proposed

substrate-binding arrangement is similar to that of small-molecule membrane-bound O-

acyltransferases (MBOATs) and led to the suggestion that CerS may catalyze the direct

transfer of the acyl chain from acyl-CoA to the long-chain base in a single step involving the

two active site histidines[50]. In addition, while this manuscript was under review, the cryo-EM

structure of yeast Lac1p bound to an acyl-CoA was reported. That structure revealed a small

lateral opening in Lac1p adjacent to the active site[56]. This observation led the authors to

hypothesize that it could potentially correspond to the entry route for sphingoid base

substrates[56]. However, this lateral opening is not present in our cryo-EM structures of CerS6

or in our molecular dynamics simulations. Instead, our experimental structures and

biochemical studies of CerS6 present several pieces of evidence which support a ping-pong

type mechanism, and challenge a ternary complex reaction. First, CerS possess a completely

different fold to that of the MBOATs. Indeed, CerS6's structure is strikingly similar to the

fatty acid elongase ELOVL7, forming a 6-TM barrel that encloses a narrow tunnel with an

essential and structurally superimposable active site histidine pair, which we have also

recently shown to catalyze acyl chain transfer via a ping-pong reaction mechanism[34]. Further,

purification of a stable acyl-CerS6 adduct species is incompatible with a ternary complex

reaction mechanism. Rather, by incubating acyl-CerS6 with sphinganine, FB_1 or FTY720, we

show that this species is a *bona fide* reaction intermediate which results in the expected

product upon exposure to the second substrate. Additionally, we captured the product-bound

state in the N-acyl FB_1-bound structure, where the toxin's hydrocarbon chain binds

perpendicularly to the membrane plane and interacts with the hydrophobic face of the central

cavity formed by TM5-7. This binding mode in the N-acyl FB_1-bound structure conflicts with

a MBOAT-style substrate coordination, but is still consistent with the action of CerS on the

fluorescently labelled NBD-sphinganine[57] as the bulky NBD moiety would sit near the wide

cytoplasmic entrance of the central cavity. Crucially, FB_1 is a structural analog of

sphinganine which undergoes N-acylation similarly to sphingoid base substrates, and its

mode of inhibition is competitive towards sphinganine[30], implying that FB_1 and sphinganine

likely bind to CerS in a similar manner. In aggregate, our data do not support a ternary

complex reaction mechanism. Rather, an ELOVL-type ping-pong model is consistent with

both our newly determined structures and biochemical results, and previously reported CerS

mutants[17,18,47].

Our observation that CerS6 can transfer a C16:0 acyl chain to the primary amine of FTY720

has important consequences for using compounds that mimic sphingoid bases in a clinical

setting. The action of CerS partially explains the N-acylated FTY720 species found during

the FTY720 drug trials[40], and by extension this enzyme family may play a role in the

pharmacokinetics of other sphingoid analogs. Further, our FB_1 results reveal how this toxin

can arrest the protein in a product-bound state (Extended Data Fig. 10), supporting the notion

that unintended substrates of this enzyme family may result in the formation of inhibitory

products. Taking advantage of this inhibition, ceramide or N-acyl FB_1 mimetics could

provide novel chemical scaffolds for new CerS inhibitors.

Methods

CerS6 cloning and expression

The *Homo sapiens CERS6* gene, encoding the CerS6 protein (Uniprot ID Q6ZMG9) was cloned into the baculovirus transfer vector pHTBV1.1-CT10H-SIII-LIC (adapted from the BacMam vector pHTBV1.1, provided by Frederick Boyce, Massachusetts General Hospital, Cambridge, MA) upstream of C-terminal TEV-cleavable 10x His and Twin-Strep tags. The construct used for structural determination lacked the final 42 amino acids, which are predicted to be disordered. For *in vitro* biotinylation, the avi-tagged CerS6 construct contained a GGGS linker between CerS6 and the Avi tag (GLNDIFEAQKIE), located upstream of the TEV cleavage site. Site-directed mutagenesis was performed using the Q5 site-directed mutagenesis kit (NEB).

Baculoviral DNA was generated by transposition of DH10Bac with the baculovirus transfer vector. Baculovirus was produced by transfecting *Sf*9 cells (Thermo Fisher Scientific) with the baculoviral DNA using the Insect GeneJuice transfection reagent (Merck Milipore). The virus was amplified by infecting *Sf*9 cells in the presence of 2 % fetal bovine serum and incubating for 72h at 27 $^{\circ}$C. For large-scale protein expression in Expi293F GnTI- cells (Thermo Fisher Scientific) in Freestyle 293 expression medium (Thermo Fisher Scientific), cells were transduced with baculovirus and supplemented with 5 mM sodium butyrate. Cells were incubated in a humidity-controlled orbital shaker at 37 $^{\circ}$C with 8 % CO_2 for 48h, and subsequently harvested by centrifugation at 900 x g for 15 min. Pelleted cells were washed with phosphate buffered saline (PBS), pelleted again by centrifugation, flash-frozen in liquid N_2 and stored at -80 $^{\circ}$C. To compare mutant activity, wild-type and mutant proteins were expressed in Expi293F cells (Thermo Fisher Scientific) using the same protocol.

Large-scale CerS6 purification

CerS6-overexpressing cells were re-suspended in Buffer A (50 mM HEPES pH 7.5, 200 mM

NaCl, 5 % (v/v) glycerol) and solubilized with 1 % (w/v) lauryl maltose neopentyl glycol

(LMNG) (Anatrace) and 0.1 % (w/v) cholesteryl hemisuccinate (CHS) (Sigma-Aldrich).

Insoluble material was removed by centrifugation at 35,000 x g, and the protein was purified

from the supernatant by batch binding to Strep-Tactin XT Superflow resin (IBA

Lifesciences). The resin was washed initially with Buffer A containing 0.05 % (w/v) GDN,

and subsequently with Buffer A supplemented with 1 mM ATP (Sigma-Aldrich), 10 mM

MgCl$_2$, and 0.05 % (w/v) GDN. CerS6 was eluted from the resin using Buffer A

supplemented with 100 mM D-biotin (Fluorochem) and 0.02 % (w/v) GDN, and the C-

terminal tag was cleaved by TEV protease digestion overnight. The His-tagged TEV protease

was removed by binding to Co^{2+}-charged TALON resin (Clontech) and the flow-through

concentrated using a 100 kDa molecular weight cut off concentrator (Corning). The protein

was finally purified by size-exclusion chromatography on a Superdex 200 Increase 10/300

GL column (GE Healthcare), pre-equilibrated in SEC buffer (20 mM HEPES pH 7.5, 200

mM NaCl, and 0.01 % (w/v) GDN).

For nanobody generation and selection, GDN was replaced in the purification buffers with

0.003 % (w/v) LMNG and 0.0003 % (w/v) CHS. For *in vitro* biotinylation of C-terminally

avi-tagged CerS6, during the TEV cleavage step, the protein sample was further

supplemented with 15 mM MgCl$_2$, 15 mM ATP, 50 mM bicine pH 8.3, and BirA at a 1:15

(w/w) BirA:CerS ratio. Biotinylation efficiency was routinely monitored by denaturing intact

protein mass spectrometry.

For structural determination, CerS6 was mixed with a 1.5 x molar excess of nanobody Nb22

and incubated for 1h on ice. The CerS6-Nb22 complex was purified by size-exclusion

chromatography, concentrated to 5 mg mL^{-1} using a 100 kDa molecular weight cut off

concentrator (Sartorius), and processed immediately for cryo-electron microscopy.

Nanobody library generation and selection

Purified CerS6 was reconstituted at a 1:20 (protein:lipid (w/w) ratio into liposomes composed

of POPC:POPG:DMPC:DMPG (7:3:7:3 (w/w)) as previously described[58].

To obtain anti-CerS6 nanobodies, alpacas were immunized and the nanobody library was

generated as previously described[59], with the exception that 200 µg purified CerS6 in

proteoliposomes were used for each immunization. All the procedures concerning alpaca

immunization were approved by the Cantonal Veterinary Office of Zurich, Switzerland

(License No. ZH 198/17). The resulting nanobody library was screened by biopanning

against CerS6. Subsequently, 190 single clones from the enriched nanobody library were

analyzed by ELISA for binding to CerS6. 96 ELISA-positive clones were Sanger sequenced

and grouped in families according to their CDR length and sequence diversity[60]. Of these, 42

unique nanobodies were identified as belonging to 26 nanobody families and were taken for

further validation.

Nanobodies were expressed in 50 mL scale in WK6 cells and purified from periplasmic

extracts using Ni^{2+}-NTA resin, as previously described[60]. Unique nanobodies were screened

by biolayer interferometry (BLI) in an Octet Red 384 system (Sartorius) using Streptavidin

SA biosensors (Sartorius) loaded with 40 µg mL^{-1} biotinylated CerS6 in SEC buffer

containing 0.003 % (w/v) LMNG, 0.0003 % (w/v) CHS. Nanobodies with slow off rates were

identified, and nanobody-22 was prioritized for structural studies.

Thermal stability measurements

Thermal unfolding experiments were carried out on a Prometheus NT.48 instrument (NanoTemper Technologies) using 5 µM CerS6 solubilized in 0.003% LMNG and 0.0003% CHS. The protein was incubated for 1h at 4 °C in the presence or absence of 20 or 100 µM FTY720 (Sigma-Aldrich) or fumonisin B_1 (Sigma-Aldrich). Proteins were heated from 20 °C to 95 °C, at a rate of 1 °C min^{-1}, and unfolding was monitored by the ratio of fluorescence emission at 350 nm and 330 nm. The melting temperature was determined from the inflection point of the transition using the PR.ThermControl software (NanoTemper Technologies).

Cryo-EM sample preparation and data acquisition

The purified CerS6-Nb22 complex (5 mg mL^{-1}) was applied to freshly glow-discharged Quantifoil 200 mesh Au R1.2/1.3 grids. Plunge freezing in liquid ethane was carried out using a Vitrobot Mark IV (Thermo Fisher Scientific) set to 4 °C and 100 % humidity. For the CerS6-Nb22-FB$_1$ complex, the SEC-purified CerS6-Nb22 complex was incubated with 120 µM fumonisin B_1 (Sigma-Aldrich) for 10 minutes on ice prior to protein concentration and subsequently for an additional 2h on ice prior to grid preparation. EM grids were screened on a Glacios (OPIC, Oxford) and high resolution data were collected on a Titan Krios G3 microscope (LISCB, University of Leicester) operating at 300 kV and equipped with a Bioquantum Energy Filter (Gatan) (operated at 20 eV slit width) and a K3 direct electron detector (Gatan) at 130,000 x nominal magnification, in super-resolution mode (2x binning; physical pixel size: 0.656 Å px^{-1}), using a defocus range between -0.8 µm and -2.4 µm. The total exposure dose was 56.3 e$^-$ Å^{-2}, fractionated over 50 frames. In total, 14,309 and 18,386 movies were collected for the CerS6-Nb22 and CerS6-Nb22-FB$_1$ datasets, respectively.

Cryo-EM data processing and model building

Movies were motion-corrected using MotionCor2[61], and CTF estimation was carried out in cryoSPARC[62] (version 3.3.1). Micrographs with bad CTF fitting (>5 Å) were excluded, yielding 14,196 and 16,647 micrographs for further analysis for the CerS6-Nb22 and for the CerS6-Nb22-FB$_1$ datasets, respectively.

For the covalent acyl-enzyme intermediate dataset, particles were initially blob picked and extracted from a subset of micrographs, and representative well-resolved 2D classes were used for template-based picking of the entire dataset. 3,400,118 template-picked particles were extracted in a 360 px box Fourier cropped to 120 px corresponding to a pixel size of 1.968 Å px^{-1} and classified in two rounds of reference-free 2D classification yielding 1,261,928 good particles. Four a*b-initio* models were generated and used as reference models in heterogeneous refinement. Particles belonging to the well-resolved 2:2 CerS6-Nb dimer complex class (53 % of input particles) were selected and subjected to a further round of heterogeneous refinement. The best resolved set of particles (497,504) were then subjected to non-uniform refinement[63] with C2 symmetry imposed, yielding a 3.21 Å reconstruction (FSC = 0.143). These aligned particles were imported into RELION using the csparc2star.py script from the UCSF pyem package[64] and re-extracted in a 360 px box Fourier cropped to 270 px, corresponding to a pixel size of 0.875 Å px^{-1}. In RELION, CTF parameters were refined, followed by Bayesian particle polishing[65]. The polished particles were subjected to 3D classification without image alignment (k =10, T=12). Certain classes displayed worse side-chain density and local deviations from C2 symmetry. Therefore, to improve map quality, the 93,680 particles belonging to the two highest-resolution C2-symmetric classes were re-imported into cryoSPARC, and subjected to a final non-uniform refinement with C2 symmetry, yielding a 3.22 Å reconstruction (FSC = 0.143). Importantly, even though CTF refinement, Bayesian polishing and 3D classification in RELION did not improve the

nominal resolution, the final reconstruction reveals better defined features, including

improved side-chain density in the CerS6 active site (Extended Data Fig. 2f,g).

The N-acyl FB_1 dataset was processed using the same workflow as the covalent acyl-enzyme

intermediate dataset, with the following changes. 3,998,935 template-picked particles were

classified in two rounds of reference-free 2D classification, yielding 1,353,329 good

particles. Four a*b-initio* models were generated and used as reference models in

heterogeneous refinement. 641,462 particles belonging to the well-resolved 2:2 CerS6-Nb

dimer complex class (48 % of input particles) were subjected to non-uniform refinement with

C2 symmetry imposed, yielding a 3.17 Å reconstruction. After CTF refinement, Bayesian

polishing and 3D classification without image alignment in RELION, the 154,239 particles

belonging to the three highest-resolution C2-symmetric classes were selected. These particles

were re-imported into cryoSPARC, and subjected to a final non-uniform refinement with C2

symmetry, yielding a 2.95 Å reconstruction (FSC = 0.143).

Atomic models were generated by fitting the CerS6 AlphaFold2[66] prediction and a Phyre2[67]

nanobody homology model into the cryo-EM maps, and subsequently manually adjusted in

Coot[68]. In both datasets, residues 72-119, corresponding to the Hox-like domains, were

poorly resolved in the sharpened maps. However, their position is evident in unsharpened and

blurred maps (Extended Data Fig. 2e; Extended Data Fig. 3a). In order to model this region,

we used tight restraints to the Alphafold2 prediction for this domain and docked it into the

envelope of the blurred maps ($B_{Blur} = 200 \text{ Å}^2$), and surface-exposed side-chains were

truncated at Cβ. The atomic models were refined using PHENIX real-space refinement[69] with

secondary structure and Ramachandran restraints, and NCS constraints. Restraint dictionaries

for FB_1 and the covalently attached C16:0 chain were generated using AceDRG[70]. The final

models comprise residues 2-330 (covalent intermediate state) or 2-334 (N-palmitoyl FB_1-

bound state) of CerS6, residues 1-124 of nanobody-22, the C16:0 chain covalently attached to

His211 of CerS6 or the N-palmitoyl FB_1 product, one phosphatidylcholine molecule and the

first N-acetylglucosamine residue of the N-linked glycan visible on Asn18. Weighted Fo −

Fc ligand difference maps were calculated using Servalcat [71], available in CCPEM[72], by

omitting the ligands from the atomic models.

Denaturing intact protein mass spectrometry

The intact masses of purified protein samples were analyzed by denaturing intact protein

mass spectrometry, conducted using an Agilent 1290 Infinity LC system in line with an

Agilent 6530 Accurate-Mass quadrupole time-of-flight mass spectrometer (Agilent

Technologies), as previously described[34]. Typically, 5-8 μg of purified protein (at 1.5-2.0 mg

mL^{-1}), diluted to 20 μL in 30 % methanol in 0.1 % formic acid, were used per injection. Data

were acquired between 100-3200 *m/z* and analyzed using MassHunter Qualitative Analysis

v.B.07.00 (Agilent Technologies) software. Peaks between 650-3200 *m/z* in the sum of the

mass spectra obtained during protein elution were deconvoluted using the maximum entropy

charge deconvolution algorithm.

The identity of the deconvoluted mass peaks was assigned initially based on the expected

mass of the purified proteins, and subsequently based on the observed mass shifts between

peaks in the deconvoluted mass spectra. The lower molecular weight CerS6 peak corresponds

to loss of the initiator methionine (theoretical mass shift of -131.20 Da), acetylation of the

new N-terminus (theoretical: +42.04 Da) and addition of an N-linked GlcNAc (theoretical:

+203.19 Da). However, the major glycosylation species observed contains the complete core

N-linked glycan (theoretical: +1217.05 Da), as expected for protein expressed in Expi293F

GnTI- cells. In addition, we observed +237.47 and +238.51 Da mass shifts relatively to these

two glycosylated species, respectively, which we interpreted as palmitoylation at a single site

(theoretical: +238.41 Da), corresponding to the covalent acyl-imidazole intermediate

observed in the cryo-EM structure. One additional modification was observed ($\sim$ +264 Da),

but its identity could not be assigned. This unknown modification occurs at a distinct site to

that of the palmitoylation, as both modifications can occur simultaneously in the same protein

molecule (Extended Data Fig. 1f,g), thus excluding the possibility of this unknown

modification occurring in the active site.

To monitor the reaction of the covalent intermediate species with the second substrates, prior

to denaturing intact protein mass spectrometry analysis the protein (2 mg mL^{-1}) was

incubated with 200 μM sphinganine (Avanti Polar Lipids), 200 μM fumonisin B$_1$ (Sigma-

Aldrich), or 600 μM FTY720 (Sigma-Aldrich) for 1h 30 min at 37 $^\circ$C. All intact mass

experiments were conducted at least twice using distinct biological samples. Replicate

deconvoluted mass spectra are shown in Extended Data Fig. 5.

Product detection by liquid chromatography high-resolution mass spectrometry (LC-

HRMS)

Product detection by LC-HRMS was conducted on a nanoElute LC system in line with a

TIMS TOF Pro 2 mass spectrometer (Bruker). The reactions were set up as described for the

denaturing intact protein MS experiments, diluted 1:40 (v/v) in 30 % methanol in 0.1 %

formic acid, and 1 μL of each sample was injected onto an IonOpticks C18 nano UHPLC

column (1.6 μm particle size; 0.075 x 250 mm).

The flow rate was set to 0.5 μL min^{-1}, and the solvent system consisted of 0.1 % Optima LC–

MS grade formic acid (Fisher Chemical) in high-performance liquid chromatography

electrochemical grade water (Fisher Chemical) (solvent A) and 0.1 % formic acid in Optima

LC–MS grade methanol (Fisher Chemical) (solvent B). Initial conditions were 60 % solvent

B, and a linear gradient from 60 % to 95 % solvent B was applied over 17.8 min to elute the

samples. This was then followed by a final 2.2 min isocratic elution with 95 % solvent B,

before the system was re-equilibrated between samples for 5 min with 60 % solvent B.

The mass spectrometer was operated in positive ion mode with a capillary voltage of 1,600

V, and the drying gas was supplied at 180 °C with a flow rate of 3 L min⁻1. Additional

parameters were: deflection 1 delta of 70 V, funnel 1 RF of 350 Vpp, funnel 2 RF of 600

Vpp, multipole RF of 500 Vpp. Data were acquired between 150-2200 *m/z*, and analyzed

using the Bruker Compass DataAnalysis 5.3.556 software. The extracted ion chromatograms

are presented in Fig. 3c and Fig. 4d, and correspond to the theoretical $[M+H]^+$ ions

(tolerance: $\pm$ 0.005 *m/z*) of the reaction products N-palmitoyl dihydrosphingosine (C16:0

ceramide) (theoretical *m/z*: 540.5350), N-palmitoyl FB_1 (theoretical *m/z*: 960.6254), and N-

palmitoyl FTY720 (theoretical *m/z*: 546.4881).

Liquid chromatography electrospray ionization tandem mass spectrometry (LC-ESI-

MS/MS) characterization of N-palmitoyl FTY720

To structurally characterize the proposed N-palmitoyl FTY720 reaction product, purified

protein was incubated with 600 μM FTY720 as before, and the reaction mixture was initially

analyzed by LC-ESI-MS on an Agilent 1290 Infinity LC System in line with an Agilent 6530

Accurate-Mass quadrupole time-of-flight mass spectrometer (Agilent Technologies) as

described above. This enabled the identification of the putative $[M+H]^+$ ion of the N-

palmitoyl FTY720 product (observed *m/z*: 546.4849; theoretical *m/z*: 546.4881). Its product

ion spectrum was then obtained by LC-ESI-MS/MS. For this purpose, the mass spectrometer

was operated in positive electrospray ionization (ESI+), 4 GHz mode. MS parameters were

capillary voltage of 4,000 V, fragmentor voltage of 175 V. Data were acquired between 100-

1700 *m/z*. The targeted parent ion (546.4881 *m/z*; retention time = 8.713 min) was fragmented

using a collision energy of 14 V.

Dihydroceramide synthase activity measurements

For activity assays, wild-type or mutant CerS6 proteins were over-expressed in Expi293F

cells, and membranes were prepared as follows. Cell pellet from 0.5 L of culture was thawed

in PBS, lysed using an Emulsiflex C5 homogeniser (Avestin), and cell debris was removed

by centrifugation. Membranes were subsequently isolated by ultracentrifugation at 160,000 x

g for 1h 30 min, resuspended in Assay Buffer (20 mM HEPES pH 7.5, 25 mM KCl, 1 mM

$MgSO_4$, 0.1 % (v/v) glycerol), flash-frozen in liquid N_2 and stored at -80 °C.

On the day of the assay, membranes were thawed, diluted to 0.25 mg mL^{-1}, and 20 µL of

diluted membranes were dispensed per well on flat bottom polystyrene 384-well Lumitrac

plates (Greiner Bio One). Then, 50 µM sphinganine (2.5 µL 500 µM sphinganine in Assay

Buffer containing 10% ethanol) and 50 µM palmitoyl-CoA (Sigma-Aldrich) (2.5 µL 500 µM

palmitoyl-CoA in Assay Buffer) were added to each well for a final reaction volume of 25

µL. For untreated controls, 5 µL of Assay Buffer were added instead of the substrates. The

plates were then incubated for 1h at room temperature, and the reaction was terminated by the

addition of 40 µL butanol spiked with N-palmitoyl(d9) dihydrosphingosine (Avanti Polar

Lipids) to yield a final concentration of 5 µM N-palmitoyl(d9) dihydrosphingosine as an

internal standard. The plates were then shaken at 1,800 rpm for 2 min, centrifuged at 1,000 x

g for 30 s, and 40 µL of the organic (upper) phase were transferred into 384-well

polypropylene deep well plates (Greiner Bio One) and diluted with 40 µL butanol.

The analytical sample handling was performed by a rapid-injecting RapidFire autosampler

system (Agilent, Waldbronn, Germany) coupled to a triple quadrupole mass spectrometer

(Triple Quad 6500, AB Sciex Germany GmbH, Darmstadt, Germany) as previously

described[73], with the following modifications. Liquid sample is aspirated by a vacuum pump

into a 10 μL sample loop for 6000 ms and subsequently flushed for 3000 ms onto a C4

cartridge (Agilent) with the aqueous mobile phase (99.5% water, 0.49% acetic Acid, 0.01%

Trifluoroacetic acid, flow rate 1.5 mL min^{-1}). The MRM transition for C16:0

dihydroceramide is 540.5 → 266.3 *m/z* (declustering potential 130V, collision energy 38 V)

and for the internal standard C16:0 (d9)dihydroceramide is 549.5 → 266.3 *m/z* (declustering

potential 130V, collision energy 38 V). The mass spectrometer was operated in positive

ionization mode (curtain gas 35 a.u., collision gas medium, ion spray voltage 4200 V,

temperature 550 °C, ion source gas 1 65 a.u., ion source gas 2 80 a.u.). MS data processing

was performed in GMSU (Alpharetta, GA, USA), and peak area ratios between C16:0

dihydroceramide and internal standard peaks were calculated.

Expression of each mutant in the membrane samples was evaluated through western blotting

using 10 ng mL^{-1} Strep-Tactin conjugated with horseradish peroxidase (IBA Lifesciences)

and using SYPRO Ruby (Thermo Fisher Scientific) staining as a loading control (Source

Data Fig. 1). Band intensity was quantified using the GeneTools software (Syngene) and

activity was normalized to protein expression.

Molecular dynamics simulations

Atomistic molecular dynamics simulations were performed using the Desmond software

package (D. E. Shaw Research, New York, NY) within the Maestro software suite

(Schrödinger). The simulation setup and analysis were carried out as follows: For each of the

two states, a CerS6 monomer was embedded in a lipid bilayer consisting of 99 POPC

molecules after undergoing the protein preparation step as implemented in Maestro. The

dimensions of the simulation cell were approximately 90x70x70 Å, with a minimum distance

of 10 Å between the protein and the cell boundaries. The systems contained 8,855 (covalent

intermediate bound state) or 9,368 (N-acyl FB1 bound state) water molecules and

correspondingly 8 or 6 chloride counterions to maintain overall charge neutrality. The

simulations were performed using the OPLS4 force field[74] for the protein and lipid

molecules, and the SPC water model for the water molecules. The system was set up using

the Maestro software suite (Schrödinger). The simulations were run in Desmond using an

NPT ensemble at a temperature of 300K. A timestep of 2 fs was employed for the integration

of equations of motion. Four independent simulations were performed, each with a duration

of 100 ns, resulting in a total simulation time of 400 ns. Snapshots of the system were

recorded every 100 ps for further analysis.

Figures

Figures depicting molecular models were generated using PyMOL[75] and ChimeraX[76].

Data availability

The cryo-EM maps have been deposited in the Electron Microscopy Data Bank (EMDB) under accession codes EMD-18770 (covalent acyl-enzyme intermediate state) and EMD-18771 (N-acyl FB_1-bound state). The atomic models have been deposited in the Protein Data Bank under accession codes 8QZ6 (covalent acyl-enzyme intermediate state) and 8QZ7 (N-acyl FB_1-bound state).

Acknowledgements

T.C.P. was supported by a Wellcome PhD studentship (102164/B/15/Z). Initial work on this project was funded by the Structural Genomics Consortium, a registered charity (number 1097737) that received funds from AbbVie, Bayer, Boehringer Ingelheim, the Canada Foundation for Innovation, Genome Canada, Janssen, Merck, Novartis, the Ontario Ministry of Economic Development and Innovation, Pfizer, and Takeda, as well as the Wellcome Trust (106169/Z/14/Z). D.B.S., A.C.W.P. and G.C. were supported by the Innovative Medicines Initiative 2 Joint Undertaking (JU) under grant agreement No 875510. The JU receives support from the European Union's Horizon 2020 research and innovation program and EFPIA and Ontario Institute for Cancer Research, Royal Institution for the Advancement of Learning McGill University, Kungliga Tekniska Hoegskolan, Diamond Light Source Limited. Electron microscopy was provided through the Oxford Particle Imaging Centre (OPIC), an Instruct-ERIC centre (funded by Wellcome Trust JIF award [060208/Z/00/Z] and equipment grant [093305/Z/10/Z]) and the Midlands Regional CryoEM Facility at the Leicester Institute of Structural and Chemical Biology (LISCB), major funding from MRC (MC_PC_17136). We thank Christos S. Savva (LISCB) for technical assistance with cryo-

EM data collection and William Greenland (Agilent) for technical advice on tandem mass

spectrometry. We thank Wyatt Yue and Christian Siebold for helpful discussions.

Author contributions

T.C.P. expressed and purified protein samples, prepared proteoliposome samples for alpaca

immunization and membranes for mutant activity testing, carried out biophysical

characterizations including bio-layer interferometry experiments, designed and generated

CerS6 mutants by site-directed mutagenesis, built and refined the atomic models, and

performed intact protein MS, product identification by high-resolution MS and small-

molecule structural characterization by LC-ESI-MS/MS. T.C.P. and G.C. prepared cryo-EM

grids. T.C.P. and A.C.W.P. collected and processed the cryo-EM data. C.S.T. carried out the

molecular dynamics simulations and analysis. A.Q. was involved in the early stages of the

project, including initial screening of expression and purification conditions. R.C. provided

access to mass spectrometry instruments for intact protein and small-molecule analysis, and

assisted with small-molecule MS experiments. S.Š. conducted alpaca immunizations,

nanobody library generation, panning and identification of unique nanobody sequences. M.T.

performed the CerS6 mutant activity assays, supervised by S.T.. T.C.P. and D.B.S. wrote the

original draft of the manuscript. All authors participated in the discussion and manuscript

editing. A.P., E.P.C., G.S. and D.B.S. supervised the research.

Competing interests

The authors declare no competing interests. C.S.T., M.T., S.T., A.P., and G.S. are employees

of Boehringer Ingelheim Pharma GmbH & Co. KG.

References

Summers, S. A., Chaurasia, B. & Holland, W. L. Metabolic Messengers: ceramides. *Nature Metabolism* **1**, 1051-1058, doi:10.1038/s42255-019-0134-8 (2019).

Turpin, S. M. *et al.* Obesity-induced CerS6-dependent C16:0 ceramide production promotes weight gain and glucose intolerance. *Cell Metab* **20**, 678-686, doi:10.1016/j.cmet.2014.08.002 (2014).

Apostolopoulou, M. *et al.* Specific Hepatic Sphingolipids Relate to Insulin Resistance, Oxidative Stress, and Inflammation in Nonalcoholic Steatohepatitis. *Diabetes Care* **41**, 1235-1243, doi:10.2337/dc17-1318 (2018).

Hajduch, E., Lachkar, F., Ferre, P. & Foufelle, F. Roles of Ceramides in Non-Alcoholic Fatty Liver Disease. *J Clin Med* **10**, 792, doi:10.3390/jcm10040792 (2021).

Hammerschmidt, P. & Brüning, J. C. Contribution of specific ceramides to obesity-associated metabolic diseases. *Cell Mol Life Sci* **79**, 395, doi:10.1007/s00018-022-04401-3 (2022).

Neeland, I. J. *et al.* Relation of plasma ceramides to visceral adiposity, insulin resistance and the development of type 2 diabetes mellitus: the Dallas Heart Study. *Diabetologia* **61**, 2570-2579, doi:10.1007/s00125-018-4720-1 (2018).

Błachnio-Zabielska, A. U. *et al.* Increased bioactive lipids content in human subcutaneous and epicardial fat tissue correlates with insulin resistance. *Lipids* **47**, 1131-1141, doi:10.1007/s11745-012-3722-x (2012).

Luukkonen, P. K. *et al.* Hepatic ceramides dissociate steatosis and insulin resistance in patients with non-alcoholic fatty liver disease. *J Hepatol* **64**, 1167-1175, doi:10.1016/j.jhep.2016.01.002 (2016).

Levy, M. & Futerman, A. H. Mammalian ceramide synthases. *IUBMB Life* **62**, 347-356, doi:10.1002/iub.319 (2010).

Ternes, P., Franke, S., Zähringer, U., Sperling, P. & Heinz, E. Identification and characterization of a sphingolipid delta 4-desaturase family. *J Biol Chem* **277**, 25512-25518, doi:10.1074/jbc.M202947200 (2002).

Kitatani, K., Idkowiak-Baldys, J. & Hannun, Y. A. The sphingolipid salvage pathway in ceramide metabolism and signaling. *Cell Signal* **20**, 1010-1018, doi:10.1016/j.cellsig.2007.12.006 (2008).

Raichur, S. *et al.* The role of C16:0 ceramide in the development of obesity and type 2 diabetes: CerS6 inhibition as a novel therapeutic approach. *Mol Metab* **21**, 36-50, doi:10.1016/j.molmet.2018.12.008 (2019).

Raichur, S. Ceramide Synthases Are Attractive Drug Targets for Treating Metabolic Diseases. *Front Endocrinol* **11**, 483, doi:10.3389/fendo.2020.00483 (2020).

Hammerschmidt, P. *et al.* CerS6-Derived Sphingolipids Interact with Mff and Promote Mitochondrial Fragmentation in Obesity. *Cell* **177**, 1536-1552.e1523, doi:10.1016/j.cell.2019.05.008 (2019).

Mizutani, Y., Kihara, A. & Igarashi, Y. Mammalian Lass6 and its related family members regulate synthesis of specific ceramides. *Biochemical Journal* **390**, 263-271, doi:10.1042/bj20050291 (2005).

Winter, E. & Ponting, C. P. TRAM, LAG1 and CLN8: members of a novel family of lipid-sensing domains? *Trends Biochem Sci* **27**, 381-383, doi:10.1016/s0968-0004(02)02154-0 (2002).

Spassieva, S. *et al.* Necessary role for the Lag1p motif in (dihydro)ceramide synthase activity. *J Biol Chem* **281**, 33931-33938, doi:10.1074/jbc.M608092200 (2006).

Kageyama-Yahara, N. & Riezman, H. Transmembrane topology of ceramide synthase in yeast. *Biochem J* **398**, 585-593, doi:10.1042/bj20060697 (2006).

Venkataraman, K. & Futerman, A. H. Do longevity assurance genes containing Hox domains regulate cell development via ceramide synthesis? *FEBS Lett* **528**, 3-4, doi:10.1016/s0014-5793(02)03248-9 (2002).

20 Mesika, A., Ben-Dor, S., Laviad, E. L. & Futerman, A. H. A new functional motif in Hox
domain-containing ceramide synthases: identification of a novel region flanking the Hox and
TLC domains essential for activity. *J Biol Chem* **282**, 27366-27373,
doi:10.1074/jbc.M703487200 (2007).
21 Sassa, T., Hirayama, T. & Kihara, A. Enzyme Activities of the Ceramide Synthases CERS2-6 Are
Regulated by Phosphorylation in the C-terminal Region. *J Biol Chem* **291**, 7477-7487,
doi:10.1074/jbc.M115.695858 (2016).
22 Tidhar, R. *et al.* Eleven residues determine the acyl chain specificity of ceramide synthases. *J*
Biol Chem **293**, 9912-9921, doi:10.1074/jbc.RA118.001936 (2018).
23 Laviad, E. L., Kelly, S., Merrill, A. H., Jr. & Futerman, A. H. Modulation of ceramide synthase
activity via dimerization. *J Biol Chem* **287**, 21025-21033, doi:10.1074/jbc.M112.363580
(2012).
24 Jensen, S. A. *et al.* Bcl2L13 is a ceramide synthase inhibitor in glioblastoma. *Proc Natl Acad*
Sci U S A **111**, 5682-5687, doi:10.1073/pnas.1316700111 (2014).
25 Kim, J. L. *et al.* Fatty acid transport protein 2 interacts with ceramide synthase 2 to promote
ceramide synthesis. *Journal of Biological Chemistry* **298**, 101735,
doi:10.1016/j.jbc.2022.101735 (2022).
26 Cingolani, F., Futerman, A. H. & Casas, J. Ceramide synthases in biomedical research. *Chem*
Phys Lipids **197**, 25-32, doi:10.1016/j.chemphyslip.2015.07.026 (2016).
27 Marasas, W. F. *et al.* Fumonisins disrupt sphingolipid metabolism, folate transport, and
neural tube development in embryo culture and in vivo: a potential risk factor for human
neural tube defects among populations consuming fumonisin-contaminated maize. *J Nutr*
134, 711-716, doi:10.1093/jn/134.4.711 (2004).
28 Riley, R. T. & Merrill, A. H., Jr. Ceramide synthase inhibition by fumonisins: a perfect storm of
perturbed sphingolipid metabolism, signaling, and disease. *J Lipid Res* **60**, 1183-1189,
doi:10.1194/jlr.S093815 (2019).
29 Riley, R. T. *et al.* Fumonisins (addendum). 415–573 (World Health Organization, Geneva,
Switzerland, 2018).
30 Merrill, A. H., Jr., van Echten, G., Wang, E. & Sandhoff, K. Fumonisin B1 inhibits sphingosine
(sphinganine) N-acyltransferase and de novo sphingolipid biosynthesis in cultured neurons in
situ. *J Biol Chem* **268**, 27299-27306 (1993).
31 Berdyshev, E. V. *et al.* FTY720 Inhibits Ceramide Synthases and Up-regulates
Dihydrosphingosine 1-Phosphate Formation in Human Lung Endothelial Cells. *Journal of*
Biological Chemistry **284**, 5467-5477, doi:10.1074/jbc.m805186200 (2009).
32 Lahiri, S. *et al.* Ceramide synthesis is modulated by the sphingosine analog FTY720 via a
mixture of uncompetitive and noncompetitive inhibition in an Acyl-CoA chain length-
dependent manner. *J Biol Chem* **284**, 16090-16098, doi:10.1074/jbc.M807438200 (2009).
33 Turner, N. *et al.* A selective inhibitor of ceramide synthase 1 reveals a novel role in fat
metabolism. *Nature Communications* **9**, 3165, doi:10.1038/s41467-018-05613-7 (2018).
34 Nie, L. *et al.* The structural basis of fatty acid elongation by the ELOVL elongases. *Nature*
Structural & Molecular Biology **28**, 512-520, doi:10.1038/s41594-021-00605-6 (2021).
35 Niu, Y. *et al.* Analysis of the mechanosensor channel functionality of TACAN. *eLife* **10**, e71188
, citation = eLife 72021;71110 e71188, doi:10.7554/eLife.71188 (2021).
36 Ke, M. *et al.* Cryo-EM structures of human TMEM120A and TMEM120B. *Cell Discov* **7**, 77,
doi:10.1038/s41421-021-00319-5 (2021).
37 Holm, L., Laiho, A., Törönen, P. & Salgado, M. DALI shines a light on remote homologs: One
hundred discoveries. *Protein Sci* **32**, e4519, doi:10.1002/pro.4519 (2023).
38 Ikeda, M. *et al.* Characterization of four mammalian 3-hydroxyacyl-CoA dehydratases
involved in very long-chain fatty acid synthesis. *FEBS Letters* **582**, 2435-2440,
doi:10.1016/j.febslet.2008.06.007 (2008).

39 Cleland, W. W. The kinetics of enzyme-catalyzed reactions with two or more substrates or
products. I. Nomenclature and rate equations. *Biochim Biophys Acta* **67**, 104-137,
doi:10.1016/0006-3002(63)91800-6 (1963).
40 Zollinger, M. *et al.* Absorption and disposition of the sphingosine 1-phosphate receptor
modulator fingolimod (FTY720) in healthy volunteers: a case of xenobiotic biotransformation
following endogenous metabolic pathways. *Drug Metab Dispos* **39**, 199-207,
doi:10.1124/dmd.110.035907 (2011).
41 Wedekind, J. E., Frey, P. A. & Rayment, I. The structure of nucleotidylated histidine-166 of
galactose-1-phosphate uridylyltransferase provides insight into phosphoryl group transfer.
Biochemistry **35**, 11560-11569, doi:10.1021/bi9612677 (1996).
42 Zhou, X. *et al.* Kinetic Mechanism of Human Histidine Triad Nucleotide Binding Protein 1.
Biochemistry **52**, 3588-3600, doi:10.1021/bi301616c (2013).
43 Trent, M. S., Worsham, L. M. S. & Ernst-Fonberg, M. L. HlyC, the Internal Protein
Acyltransferase That Activates Hemolysin Toxin: Roles of Various Conserved Residues in
Enzymatic Activity As Probed by Site-Directed Mutagenesis. *Biochemistry* **38**, 9541-9548,
doi:10.1021/bi9905617 (1999).
44 Dodds, A. W., Ren, X. D., Willis, A. C. & Law, S. K. The reaction mechanism of the internal
thioester in the human complement component C4. *Nature* **379**, 177-179,
doi:10.1038/379177a0 (1996).
45 Gadjeva, M. *et al.* The covalent binding reaction of complement component C3. *J Immunol*
161, 985-990 (1998).
46 Selvy, P. E., Lavieri, R. R., Lindsley, C. W. & Brown, H. A. Phospholipase D: enzymology,
functionality, and chemical modulation. *Chem Rev* **111**, 6064-6119, doi:10.1021/cr200296t
(2011).
47 Vanni, N. *et al.* Impairment of ceramide synthesis causes a novel progressive myoclonus
epilepsy. *Ann Neurol* **76**, 206-212, doi:10.1002/ana.24170 (2014).
48 Harrer, H., Laviad, E. L., Humpf, H. U. & Futerman, A. H. Identification of N-acyl-fumonisin B1
as new cytotoxic metabolites of fumonisin mycotoxins. *Mol Nutr Food Res* **57**, 516-522,
doi:10.1002/mnfr.201200465 (2013).
49 Harrer, H., Humpf, H. U. & Voss, K. A. In vivo formation of N-acyl-fumonisin B1. *Mycotoxin*
Res **31**, 33-40, doi:10.1007/s12550-014-0211-5 (2015).
50 Zelnik, I. D. *et al.* Computational design and molecular dynamics simulations suggest the
mode of substrate binding in ceramide synthases. *Nat Commun* **14**, 2330,
doi:10.1038/s41467-023-38047-x (2023).
51 Youssefian, L. *et al.* Autosomal recessive congenital ichthyosis: CERS3 mutations identified
by a next generation sequencing panel targeting ichthyosis genes. *Eur J Hum Genet* **25**, 1282-
1285, doi:10.1038/ejhg.2017.137 (2017).
52 Eckl, K. M. *et al.* Impaired epidermal ceramide synthesis causes autosomal recessive
congenital ichthyosis and reveals the importance of ceramide acyl chain length. *J Invest*
Dermatol **133**, 2202-2211, doi:10.1038/jid.2013.153 (2013).
53 Radner, F. P. *et al.* Mutations in CERS3 cause autosomal recessive congenital ichthyosis in
humans. *PLoS Genet* **9**, e1003536, doi:10.1371/journal.pgen.1003536 (2013).
54 Dougherty, D. A. The cation-π interaction. *Acc Chem Res* **46**, 885-893,
doi:10.1021/ar300265y (2013).
55 Merrill, A. H., Jr., Wang, E., Gilchrist, D. G. & Riley, R. T. Fumonisins and other inhibitors of de
novo sphingolipid biosynthesis. *Adv Lipid Res* **26**, 215-234 (1993).
56 Xie, T. *et al.* Structure and mechanism of a eukaryotic ceramide synthase complex. *Embo j*,
e114889, doi:10.15252/embj.2023114889 (2023).
57 Tidhar, R., Sims, K., Rosenfeld-Gur, E., Shaw, W. & Futerman, A. H. A rapid ceramide
synthase activity using NBD-sphinganine and solid phase extraction. *J Lipid Res* **56**, 193-199,
doi:10.1194/jlr.D052001 (2015).

58   Geertsma, E. R., Nik Mahmood, N. A., Schuurman-Wolters, G. K. & Poolman, B. Membrane
reconstitution of ABC transporters and assays of translocator function. *Nat Protoc* **3**, 256-
266, doi:10.1038/nprot.2007.519 (2008).
59   Hutter, C. A. J. *et al.* The extracellular gate shapes the energy profile of an ABC exporter.
Nature Communications **10**, 2260, doi:10.1038/s41467-019-09892-6 (2019).
60   Pardon, E. *et al.* A general protocol for the generation of Nanobodies for structural biology.
Nature Protocols **9**, 674-693, doi:10.1038/nprot.2014.039 (2014).
61   Zheng, S. Q. *et al.* MotionCor2: anisotropic correction of beam-induced motion for improved
cryo-electron microscopy. *Nat Methods* **14**, 331-332, doi:10.1038/nmeth.4193 (2017).
62   Punjani, A., Rubinstein, J. L., Fleet, D. J. & Brubaker, M. A. cryoSPARC: algorithms for rapid
unsupervised cryo-EM structure determination. *Nat Methods* **14**, 290-296,
doi:10.1038/nmeth.4169 (2017).
63   Punjani, A., Zhang, H. & Fleet, D. J. Non-uniform refinement: adaptive regularization
improves single-particle cryo-EM reconstruction. *Nat Methods* **17**, 1214-1221,
doi:10.1038/s41592-020-00990-8 (2020).
64   Asarnow, D., Palovcak, E. & Cheng, Y. *UCSF pyem v0.5*,
<https://doi.org/10.5281/zenodo.3576630}> (2019).
65   Zivanov, J. *et al.* New tools for automated high-resolution cryo-EM structure determination
in RELION-3. *Elife* **7**, doi:10.7554/eLife.42166 (2018).
66   Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature* **596**,
583-589, doi:10.1038/s41586-021-03819-2 (2021).
67   Kelley, L. A., Mezulis, S., Yates, C. M., Wass, M. N. & Sternberg, M. J. The Phyre2 web portal
for protein modeling, prediction and analysis. *Nat Protoc* **10**, 845-858,
doi:10.1038/nprot.2015.053 (2015).
68   Emsley, P., Lohkamp, B., Scott, W. G. & Cowtan, K. Features and development of Coot. *Acta
crystallographica. Section D, Biological crystallography* **66**, 486-501,
doi:10.1107/S0907444910007493 (2010).
69   Afonine, P. V. *et al.* Real-space refinement in PHENIX for cryo-EM and crystallography. *Acta
Crystallogr D Struct Biol* **74**, 531-544, doi:10.1107/s2059798318006551 (2018).
70   Long, F. *et al.* AceDRG: a stereochemical description generator for ligands. *Acta Crystallogr D
Struct Biol* **73**, 112-122, doi:10.1107/s2059798317000067 (2017).
71   Yamashita, K., Palmer, C. M., Burnley, T. & Murshudov, G. N. Cryo-EM single-particle
structure refinement and map calculation using Servalcat. *Acta Crystallogr D Struct Biol* **77**,
1282-1291, doi:10.1107/s2059798321009475 (2021).
72   Burnley, T., Palmer, C. M. & Winn, M. Recent developments in the CCP-EM software suite.
Acta Crystallogr D Struct Biol **73**, 469-477, doi:10.1107/s2059798317007859 (2017).
73   Tautermann, C. S. *et al.* Allosteric Activation of Striatal-Enriched Protein Tyrosine
Phosphatase (STEP, PTPN5) by a Fragment-like Molecule. *J Med Chem* **62**, 306-316,
doi:10.1021/acs.jmedchem.8b00857 (2019).
74   Lu, C. *et al.* OPLS4: Improving Force Field Accuracy on Challenging Regimes of Chemical
Space. *J Chem Theory Comput* **17**, 4291-4300, doi:10.1021/acs.jctc.1c00302 (2021).
75   Schrodinger, LLC. *The PyMOL Molecular Graphics System, Version 2.1* (2015).
76   Pettersen, E. F. *et al.* UCSF ChimeraX: Structure visualization for researchers, educators, and
developers. *Protein Sci* **30**, 70-82, doi:10.1002/pro.3943 (2021).

1006