Coevolution of longevity and female care of male gametes

Julian Baur[1], Mareike Koppik[1,2], Uroš Savković [3], Mirko Đorđević [3], Biljana Stojković [4] & David Berger[1],†

† Corresponding author: david.berger@ebc.uu.se.

1: Department of Ecology and Genetics, Uppsala University, Sweden.

2: Animal Ecology, Department of Zoology, Martin-Luther University Halle-Wittenberg, Halle (Saale), Germany; mareike.koppik@zoologie.uni-halle.de

3: Department of Evolutionary Biology, Institute for Biological Research "Siniša Stanković", National Institute of the Republic of Serbia, University of Belgrade, Bulevar despota Stefana 142, 11000 Belgrade, Serbia

    Uroš Savković – savkovic.uros@ibiss.bg.ac.rs +381112078376

    ORCID: https://orcid.org/0000-0001-9430-4619

    Mirko Đorđević – mirko.djordjevic@ibiss.bg.ac.rs +381112078376

    ORCID: https://orcid.org/0000-0002-5768-0081

4: Institute of Zoology, Chair of Genetics and Evolution, Faculty of Biology, Studentski trg 16, 11000 Belgrade, Serbia

    Biljana Stojković – bilja@bio.bg.ac.rs +381112635627

    ORCID: https://orcid.org/0000-0001-9978-2249

Keywords: Life-history, reproduction, female care, parental care, mutation rate, offspring quality, conflict, trade-off, germline

Abstract

An often-overlooked aspect of life-history optimization is the allocation of resources to protect the germline and secure safe transmission of genetic information. While failure to do so renders significant fitness consequences in future generations, germline maintenance comes with substantial costs. Thus, germline allocation should trade-off with other life history decisions and be optimized in accord with an organism's reproductive schedule. Here we tested this hypothesis by studying germline maintenance in lines of seed beetle, selected for early (E) or late (L) reproduction for 350 and 240 generations, respectively. Female animals provide maintenance of male gametes in their reproductive tract and oocytes. Here, we revealed the ability of young and old E and L-females to provide this form of germline maintenance by mating them to males with ejaculates with artificially elevated levels of protein and DNA damage. We find pronounced evolved differences in this form of germline maintenance between E and L females that accord with the age of reproduction in respective regime. These findings identify the central role of allocation to secure germline integrity in life history evolution and highlight how females can play a crucial role in mitigating effects of male germline decisions on mutation rate and offspring quality.

Introduction

The interplay between growth, reproduction, and survival is a central focus of life history theory (Roff, 1993; Stearns, 1992). Trade-offs among these three fundamental aspects of an organism's life play key roles in dictating the evolution of reproductive schedules, with cascading effects on organismal aging and demography (Flatt & Heyland, 2011; Monaghan et al., 2008; Promislow & Tatar, 1998). Indeed, some species invest heavily in early reproduction at the expense of longevity, while others prioritize growth and survival before reproducing, and these alternative strategies are expected outcomes of differences in mortality risk and age-related changes in the strength of natural selection (Roff, 1993; Stearns, 1992).

Another, often less explicitly defined, dimension of life history optimization involves the allocation of resources to protect the germline (Maklakov & Immler, 2016; Monaghan & Metcalfe, 2019). The integrity of the germline is essential for the transmission of genetic information to subsequent generations, and hence, the evolutionary consequences of germline maintenance strategies can be considerable both at the individual and population level (Agrawal & Whitlock, 2012; Lynch, 2010; Muller, 1950). The costs of germline maintenance may be substantial (Maklakov & Immler, 2016), in part due to the expensive repair machinery required to keep the rate of deleterious germline mutations at levels several magnitudes lower than the somatic mutation rate (Lynch, 2008). This sets the stage for trade-offs between the germline and other life history traits, where costly maintenance of the germline may secure the genetic quality of future generations at the cost of reduced reproduction and/or survival of the parent (Maklakov & Immler, 2016; Monaghan & Metcalfe, 2019).

Thus, theories of aging not only predict age-specific optimization of classic life history traits, but also concerted evolution of processes aimed at protecting the germline. For example, the mutation accumulation theory of aging (Medawar, 1952), from hereon: "MA", predicts a decline in germline maintenance past an organism's reproductive peak due to weakened selection and the accumulation of alleles with deleterious effects late in life. Indeed, ageing not only leads to somatic deterioration (Partridge & Barton, 1996), but also negatively

affects the maintenance of the germline (Monaghan & Metcalfe, 2019), shown, for example,

by the fact that *de novo* germline mutation rates in primates seem to be foremost

associated with advancing age (Kong et al., 2012; Rahbari et al., 2016a; Venn et al., 2014).

Ageing-trajectories can also be driven by antagonistic pleiotropy (Williams, 1957), from

hereon: "AP", whereby alleles with beneficial effects on germline maintenance at the peak

of reproduction (e.g. via increased allocation of energy to germline repair) are favoured by

selection despite potential deleterious side-effects on somatic maintenance (Chen et al.,

2020), reproduction (Baur & Berger, 2020; Berger et al., 2017), and/or germline

maintenance at other ages.

There is now growing evidence that germline decisions can be studied through a life-history

lens. For instance, germline maintenance strategies can differ between environments that

place different needs on maintenance versus reproduction (Berger et al., 2021; Chu et al.,

2018; Crean & Immler, 2021; Drake, 2009; Kurhanewicz et al., 2020; Puurtinen et al., 2016;

Svetec et al., 2016), or between individuals that differ in their overall energy budgets and

allocation decisions (Agrawal & Wang, 2008; Baur & Berger, 2020; Berger et al., 2017;

Koppik et al., 2023; Sharp & Agrawal, 2012). Interestingly, female animals possess the ability

to provide maintenance not only of their own germline cells, but also of male ejaculates and

haploid male DNA, by synthesizing anti-oxidants in the reproductive tract to neutralize

mutagenic reactive oxygen species, and by including mRNA and proteins in the oocyte to

detect and repair DNA damage in the early zygote (Horta et al., 2020; Jaroudi et al., 2009;

Stringer et al., 2018). This sets the stage for an intricate evolutionary dynamic, where care

over transferred ejaculates and newly formed zygotes is the ultimate decision of the female,

but the need for it might be dictated by the germline decisions of the male partner, or, like

in human populations, his age (Gao et al., 2019a; Rahbari et al., 2016b; Ségurel et al., 2014).

Even though there now is a good understanding of the mechanisms that cause age-

dependent variation in DNA-repair and maintenance of the germline, little is known about

the evolutionary potential of such mechanisms. In humans, for example, changes to

environmental (Klass, 1977), pharmaceutical (Johnson et al., 2013) and genetic (Guarente &

Kenyon, 2000) factors can increase longevity substantially within one or a few generations,

and the peak of reproduction has in many populations changed drastically over the last

century (Khandwala et al., 2017a; Mills et al., 2011a). However, the long-term genetic consequences of lifespan extension depend on whether selection on reproductive schedules are coupled with concomitant changes in germline repair and maintenance, which remain unexplored. Here we harnessed the power of long-term experimental evolution to investigate how the evolution of lifespan affects age-dependent germline maintenance in the seed beetle, *Acanthoscelides obtectus.* Using a set of lines that were established in 1989 and selected for early or late reproduction for more than 350 and 240 generations, respectively, we examined germline maintenance in young and old females receiving male ejaculates with artificially inflated levels of free radicals and DNA and protein damage (Fig. 1). This allowed us to explore i) whether female maintenance of male gametes evolves in concerted fashion with reproductive schedules and other life history traits, and ii) whether the AP (Williams, 1957) or the MA (Medawar, 1952) theory of ageing best explains the evolutionary patterns observed.

Methods

Study species

The seed beetle, *Acanthoscelides obtectus* (Coleoptera; Chrysomelidae; Bruchidae) is a pest of stored legumes, foremost, the common bean, *Phaseolus vulgaris* (Thakur, 2012). Females lay eggs either directly under or in close proximity of host seeds and the hatched larvae burrow into a seed where juvenile development takes place (Tucić et al., 1996). After pupation, adults emerge from the seed and typically become reproductively active within 24 hours. Adult beetles are aphageous, meaning they don´t need nutrition nor water to be able to reproduce and finish their lifecycle.

Experimental populations

The original base population, from which the experimental evolution lines were derived, was created by mass-mating wild-caught beetles from three locations in the area of Belgrade, Serbia, in 1983 (Tucić et al., 1996). Subsequently, the population was maintained on *P. vulgaris* in the laboratory for 3 years (27 generations) at large population size (ca. 5000 individuals), until the experimental evolution lines were created in 1986. Experimental

evolution was performed in dark incubators at 30°C. No food or water was offered to

experimental beetles.

During experimental evolution, one line was maintained under the same conditions as the

base population (from here on referred to as the "base" line); after 40 days of development,

400-500 adult beetles were chosen at random and provided with fresh host seeds and

allowed to reproduce for the next generation. Each of four replicate lines selected for early

reproduction (E-lines) was created by taking 400 adults at random on the peak of adult

emergence. To propagate the next generation, these adults were transferred to a new

bottle containing host seeds *ad libitum* and were there allowed to reproduce and lay eggs

for 48h, after which all adults were removed. Another four replicate lines were selected for

late reproduction (L-lines) by taking 100 newly emerged (0-24h) adults every day for up to

10 days. Each of these subgroups was kept in a vial without seeds for 10 days. On the 10th

day of each subgroup all the surviving beetles of a given replicate line were added to a

single bottle containing fresh host seeds to propagate the next generation. For more

information on the maintenance of the experimental evolution lines, see (Tucić et al., 1996,

1997). Development time of E-lines has evolved to be 29 days and the average virgin

lifespan around 13 days in aphagous conditions, while L-lines emerge after 32 days and

virgin beetles live on average 26 days for females and 22 days for males (Đorđević et al.,

2015). The regimes have also diverged in other traits tied to rate-dependent life-history,

such as the day of first oviposition (E: 1 day after eclosion, L: 4 days after eclosion) (Tucić et

al., 2004), metabolic rate, and body size (Arnqvist et al., 2017).

Experimental design

The experiment was started in 2020, after 240 (L-lines) and 350 (E-lines) generations of

experimental evolution. To estimate the evolution of age-dependent germline maintenance

we compared females of all lines mated to males when aged 1-3 days (young age class) or 6-

8 days (old age class). Note that 6-8 days is not truly an old age for L-lines and should be

interpreted in relative terms. However, since we wanted to compare E- and L females, and E

females age rapidly, 6-8 days were chosen to avoid any bias in form of selective

disappearance (van de Pol & Verhulst, 2006) of low quality individuals in the E-lines (Fig. 1).

A *Experimental Evolution of Reproductive Schedules*

B *Age-Dependent Germline Maintenance*

Figure 1: Experimental design.

A) The experiment included eight experimental evolution lines; four from the E- (early reproduction; salmon) and four from the L- (late reproduction; green) regime. The E-regime was propagated by harvesting eggs laid after only 2 days of adult reproduction, whereas the L-regime was propagated by providing individuals that had aged 10 days with host seeds and harvesting the eggs. The E and L regime had been kept for 350 and 240 generations, respectively, prior to the experiment. **B)** Females from all lines, that were either young (1-3 days) or old (6-8 days), were mated to standard males from the base population. These males were either serving as controls or they had been irradiated to generate a damaged ejaculate that would challenge maintenance in the female reproductive tract and oocytes. Offspring resulting from the matings to control males were counted to estimate age-dependent reproductive schedules of the F0 females in the experiment (Fig. 2). F1 offspring were then crossed within line and treatment group using a middle-class neighbourhood design to produce F2 offspring. From these F2 offspring counts, age-specific female care of male gametes could be deduced for each line by comparing counts from lineages deriving from young and old F0 females, mated with either irradiated or control males (Fig. 3). Photo credit: Mareike Koppik.

All lines were reared in staggered fashion by setting up several rearing containers of each line in the two generations prior to the experiment, so that emergence of females would happen continuously during the experiment and females assigned to the two age classes could be assayed simultaneously and mated to standard males from the same cohort (see below). We collected females on the day of their emergence, randomly assigned them to the young or old age class, and then isolated them in perforated 0.5ml Eppendorf tubes and left them to age. Females of both age classes were then mated to young (1-2 days) virgin males from the base population in 60 mm petri-dishes placed on heating plates kept at 30°C. Couples were observed to avoid multiple mating. Males were removed after a mating behaviour had been observed and females were allowed to lay eggs in a new 60 mm petri-dish containing host seeds *ad libitum*.

The base males were either untreated (controls) or had been exposed to a 10 Gy dose of γ-radiation, which causes substantial cellular damage to the male germline (Friedberg et al., 2005) and impairs male fertility in seed beetles (Baur & Berger, 2020; Berger et al., 2017; Koppik et al., 2023). Hence, the female repair and maintenance of damaged male gametes in each line could be assessed by comparing the offspring production in lineages stemming from females mated to control and irradiated males. γ-radiation produces free radicals with mutagenic properties and causes direct damage to DNA within sperm cells (Friedberg et al., 2005) and to proteins within the ejaculate (Kempner, 2001). Hence, our protocol captures a multitude of effects that occur naturally in male ejaculates (Dowling & Simmons, 2009; Maklakov & Immler, 2016), but that will have been greatly amplified by the irradiation treatment.

As our interest lies in capturing effects of variation in female care that transcend generations, we scored the fitness consequences of variation in female care by counting the F2 offspring produced by all descending lineages. To achieve this, F1 males and females stemming from the same treatment group and line were paired (while avoiding sib-mating) and allowed to mate and produce F2 offspring in petri-dishes with ad libitum host seeds. The emerging F2 offspring were counted to estimate each couple's fecundity, which was used to estimate germline maintenance of the their F0 grandmothers (see below: "Statistical analysis").

About half of the F0 couples produced no offspring. This was likely due to two reasons. First,

the mating latency of *A. obtectus* can be relatively long, and because the time in between

male irradiation and mating needed to be standardized, many couples did not mate in this

time. Secondly, even when a mating was thought to have been observed, several couples

did still not produce any offspring. Using a binomial model, we tested whether the incidence

of zero fertility was related to any of the factors of interest in our experiment (i.e.,

irradiation treatment, selection regime, and age). However, despite large sample size, no

factor had a significant effect on infertility. Indeed, all effects including male irradiation

treatment were very weak and non-significant (all $P > 0.3$), and the proportion of infertile

couples was relatively equal across treatments (E: young ctrl = 0.43; young irr = 0.48; old ctrl

= 0.59; old irr = 0.61. L: young ctrl = 0.50; young irr = 0.39; old ctrl = 0.48; old irr = 0.50).

Hence, we carried out our downstream analyses based on the remaining data from the 830

F0 couples that produced offspring. Each line and age-class was represented by 10-20

females mated to control males, and 20-30 females mated to irradiated males.

To reduce the effect of selection against epigenetic and DNA mutations induced by the

irradiation we applied a Middle-Class Neighbourhood (MCN) design (Shabalina et al., 1997).

The MCN design restricts each F0 couple to contribute the same number of F1 offspring for

future study, irrespective of their overall productivity. We note, however, that F0 couples

that produced less than four F1 offspring were not represented in the final data, and hence,

selection on very strongly deleterious effects of irradiation could not be excluded by the

MCN approach. For those F0 couples that did produce offspring, we aimed at propagating a

single male and female from each couple, so that each F0 couple would replace itself.

However, due to variability in the timing of eclosion of F1 individuals, additional F0 couples

fell out of the experiment as we did not want to age F1 beetles before setting them up in

assays. These strict criteria resulted in 490 F1 couples being analysed for offspring

production. Each line and age class was represented by ca 20 and 10 F1 couples that

originated from irradiated and control lineages, respectively. The experiment was

conducted in three temporal blocks, 3 days apart, corresponding to the three occasions on

which we irradiated F0 males from the base population. In the F1 generation, when

offspring productivity assays were set up, there was considerable overlap between blocks

due to variable timing of egg laying and juvenile development time.

Statistical analysis

We analyzed data using generalized linear mixed effect models implemented in the package MCMCglmm (Hadfield, 2010) for R (R Core Team, 2020). Offspring counts were analysed assuming Poisson distributed error. We used flat and uninformative priors for all models. We ran all models for 1.05 million iterations with an initial burn in of 50k iterations and a thinning factor of 500 to avoid autocorrelations, resulting in a posterior sample of 2000. P-values for all comparisons were calculated based on model posterior distributions, where significance of main effects was calculated by comparing posteriors of marginal means of two given groups (e.g. overall difference in offspring production in irradiated and control lineages). We used ggplot2 for graphical illustration (Wickham, 2016).

To provide an estimate of the extent of age-specific optimization of reproductive schedules at the current stage of experimental evolution, we first compared counts of offspring produced by young and old F0 females of all E- and L-lines, when mated to the standard base males belonging to the control (un-irradiated) treatment (N = 307). This model included female age (young vs. old), evolution regime (E vs. L), and their interaction, as fixed effects of interest. We also added the main effect of male age (1 or 2 days old) as a blocking factor. Replicate line, crossed with age, were added as random effects.

To analyse differences in female care of male gametes we compared the F2 offspring counts of control and irradiated lineages. This model included fully crossed fixed effects of evolution regime (E vs. L), F0 female age (young vs. old) and male radiation treatment (control vs. irradiated). Note that the three-way interaction between these effects tests our focal hypothesis that the E- and L-regime have diverged in age-specific female care of male gametes. Replicate line crossed with age and radiation treatment were included as random effects. To provide an intuitive measure of germline maintenance for reporting effect sizes, we also calculated the relative reduction in fertility due to unattended damage in the male ejaculate for each replicate line at young and old age, according to: $\Delta\zeta = 1 - \zeta_{IRR,\,i} / \zeta_{CTRL,\,i}$, where $\zeta_{IRR,\,i}$ and $\zeta_{CTRL,\,i}$ is the number of F2 offspring produced by lineages deriving from F0 females of age *i*, mated to irradiated and control males, respectively (see: Fig. 3B).

Results

Evolution of age-specific reproductive schedules

There was no difference in F1 offspring production between young E and L females mated to control males in the F0 generation (p_{MCMC} = 0.28). However, there were clear differences in aging trajectories. As expected, females originating from the E-regime showed a sharp decline in offspring production when mated at old age (p_{MCMC} < 0.001), while no such effect was seen in the old L-females. As follows, the interaction between regime and female age was highly significant (p_{MCMC} < 0.001) and accorded with predictions (Fig. 2).

Figure 2: **Evolution of female reproductive schedules.**

The number of F1 offspring produced by young and old F0 females of the E (salmon) and L (green) regime, when mated to standard control males from the base line. Shown are means and 95% confidence limits.

Coevolution of reproductive schedules and female care of male gametes

The irradiation treatment led to an overall decrease in F2 offspring numbers (effect of irradiation in L-regime: p_{MCMC} < 0.001; effect of irradiation in E-regime: p_{MCMC} = 0.002). Taken across both age-classes, there was no difference between evolution regimes in the effect of irradiation on offspring production (regime*treatment interaction: p_{MCMC} = 0.74), providing no evidence for strong divergence in female germline maintenance overall. However, in terms of the age-dependency of female care, the two regimes seemed to behave very differently (regime*age*treatment interaction; p_{MCMC} = 0.008, Fig. 3). This difference was due to that the effect of irradiation was stronger in the young age-class compared to the old age-class for the L-regime (p_{MCMC} = 0.004), while the aging trajectory showed the opposite pattern for E-lines (although the effect of ageing was not significant, p_{MCMC} = 0.35). These results confirm that female care of male gametes has evolved age-specific optimization.

Figure 3: **Evolution of age-dependent female germline maintenance of male ejaculates.**

A) F2 offspring production in lineages stemming from F0 grandmothers mated to either a control (light
bars) or irradiated (dark bars) male. Grandmothers were either young (1-3 days) or old (6-8 days) at
mating, and belonged to either the E (salmon, left panel) or L (green, right panel) regime. Shown are
means and standard errors. **B)** Line-specific estimates of the relative reduction in F2 offspring
production induced by the irradiation treatment in F0 males, calculated as : $\Delta\zeta = 1 - \zeta_{IRR, i} / \zeta_{CTRL, i}$, where
$\zeta_{IRR, i}$ and $\zeta_{CTRL, i}$ is the number of F2 offspring produced by lineages deriving from F0 females of age i,
mated to irradiated and control males, respectively. This estimate is thus inversely related to the
efficiency of germline DNA repair and maintenance in F0 mothers. Plain and striped bars represent
lineages originating from young and old F0 grandmothers, respectively.

Discussion

Here we have shown that selection on reproductive schedules can lead to the evolution of age-specific maintenance of male gametes in the female reproductive tract. Such germline maintenance is a somewhat underestimated form of parental care with genetic consequences that can transcend generations. Our findings echo previous studies that have provided evidence that germline maintenance is costly and can trade off with life-history traits (Agrawal & Wang, 2008; Baur & Berger, 2020; Berger et al., 2017; Chen et al., 2020; Maklakov et al., 2012), emphasizing that germline maintenance should be considered a life-history trait in its own right (Bartosch -Harlid et al., 2003; Berger et al., 2017; Bromham, 2011; Ellegren, 2007; Maklakov & Immler, 2016).

The observed evolution of aging trajectories follows the general expectation that repair and maintenance in the oocyte evolves to optimize function at peak reproduction. The mutation accumulation (MA) theory of ageing (Medawar, 1952) predicts that E-lines should show efficient maintenance early in life, but a sharp decline in maintenance at old age, as old E-beetles have not been exposed to selection for >350 generations. In L-lines, a constant and efficient maintenance at both age-classes is expected, given that mature gametes have already formed at adult eclosion (Biemont, 1973) and should need continuous maintenance. Indeed, the observed reproductive schedules of young and old E- and L-females mated to control (unirradiated) males were qualitatively consistent with the MA hypothesis (Fig. 2). However, for female care of male gametes, E-females showed only a modest drop at old age, and instead it was young L-females that showed the lowest care (Fig. 3). L- females have not been provided with egg-laying substrate at young age for over 240 generations and egg-laying at this age thus represents a novel condition, which means that accumulation of mutations with conditional deleterious effects only at young age cannot be excluded. However, while the dip in care was larger in young (relative to old) L-females compared to the dip in old (relative to young) E-females, the E-regime had undergone more generations of (potential) accumulation of mutations with age-specific effect. Thus, the observed aging trajectories of germline maintenance are not fully compatible with the MA hypothesis.

Another (not mutually exclusive) explanation for the evolved aging trajectories of germline

maintenance and female care is antagonistic pleiotropy (AP) (Williams, 1957). Indeed,

previous findings have shown that aging trajectories of fertility and longevity in the E- and L

lines of *A. obtectus* seem to be a combined result of both MA and AP (Stojković et al., 2015).

Pleiotropic effects of genes improving germline maintenance at young age could lead to

reproductive disadvantages at later ages (Maklakov & Immler, 2016; Nussey et al., 2006)

and/or reduced survival and longevity via trade-offs between somatic and germline DNA-

maintenance (Chen et al., 2020; Maklakov & Immler, 2016), and hence, would be

disfavoured by selection in the L-regime. One likely molecular candidate linking

development, longevity, and reproduction in insects is juvenile growth hormone and its

variants, which, in fruit flies is known to increase sexual maturation, vitellogenesis, and

reproduction, while decreasing lifespan (Flatt et al., 2005). The pronounced dip in germline

maintenance in young L-females is also mirrored by common observations of reduced

fertility due to reproduction too early in life across animal taxa (Jones et al., 2014).

The potential for females to care over male ejaculates and offspring already at the very

onset of fertilization opens up interesting ecological scenarios and the potential for sexual

conflict over parental care, even in species traditionally assigned as having little to no care

(Koppik et al., 2023; Maklakov et al., 2012; Schärer et al., 2012; Servedio et al., 2019). For

example, in most species with internal fertilization, healthy females may efficiently

compensate for poor ejaculates of older males, while the converse seems unlikely. This

asymmetry should have implications for assortative mating by age. For example, the

baseline germline mutation rate is several times greater in males compared to females in

most animal taxa (Ellegren, 2007; Hurst & Ellegren, 1998; Sayres & Makova, 2011) and, at

least in well-studied groups like primates, this bias increases with age (Gao et al., 2019b;

Wang et al., 2020). Thus, mating decisions based on partner age are likely to have

considerable consequences, and the form of maternal care studied here should be

intertwined in those decisions.

Germline maintenance decisions can also depend on the overall condition of the individual

and state-dependent allocation decisions between competing life-history demands (Berger

et al., 2017; Maklakov & Immler, 2016; Monaghan & Metcalfe, 2019; Sharp & Agrawal,

2012), with implications for sexual selection theory and mate choice processes (Baur & Berger, 2020; Beck & Promislow, 2007; Koppik et al., 2023; Segami et al., 2021). Viewing germline maintenance through a resource acquisition and allocation lens (Jong & Noordwijk, 1992) suggests that optimal mate choice is contingent on not only choosers' inferences of genetic quality in offspring based on the phenotype of their mating partner, but also of how this phenotype influences changes to the genetic quality of its gametes. For example, in many polyandrous species, strong postcopulatory sexual selection (sperm competition) can favour compromised germline maintenance in males (Bartosch -Harlid et al., 2003; Baur & Berger, 2020; Blumenstiel, 2007; Koppik et al., 2023). This suggests that males that are most successful in reproductive competition may sometimes pass on the greatest load of mutations to their offspring, questioning why females should prefer them in the first place (Baur & Berger, 2020; Beck & Promislow, 2007; Segami et al., 2021)? Interestingly, however, optimality theory predicts that females my tailor care depending on the assessed state of male gametes (Servedio et al., 2019), with the result that a male's mutation rate is not independent of his female mate and her potential compensation for his short-comings, which can complicate interpretations of both age- and sex-biases in mutation rate (Gao et al., 2019a).

Can mechanisms that protect the germline evolve fast enough to keep up with rapid environmental and demographic change? This question is of importance for understanding long-term evolutionary responses that depend on both demographic parameters and the germline mutation rate. It is also a question tied to ongoing debates in human reproductive biology against the background of the increasing age at first reproduction in western civilisations (Khandwala et al., 2017b; Mills et al., 2011b). While assisted reproduction technologies may help to mitigate immediate age-related fertility declines associated with such changes, less attention is allotted to the evolutionary implications of ever later reproduction in human populations. Our study provides compelling evidence that female care of male gametes can evolve in response to selection on reproductive schedules. The maintenance and repair capacity in oocytes peaked at the time of reproduction in E- and L-lines, and the observed evolution of aging trajectories is compatible with some role for antagonistic pleiotropy in the underlying genes. This suggest that balancing selection on concerted life-history syndromes might be responsible for the maintenance of genetic

variation in germline repair in the wild founding population, which likely contributed to the

observed evolutionary response. Our study highlights the evolutionary importance of this,

often underappreciated, type of maternal care and we hope to stimulate further research

aimed at elucidating the processes that govern the coevolution of germline maintenance,

mate choice, and other life-history traits.

Acknowledgements

We thank Johanna Liljestrand-Rönn for help in the lab, the kind people at BMC Uppsala for

providing access to the radiation source, and the members of the seed beetle lab group for

helpful discussions. This work was supported by grant no. 2019-05023 from the Swedish

Research Council (VR) to DB.

Author contributions

DB, JB and MK designed the study. JB and MK performed the study. JB and DB analyzed the

data, and DB and JB wrote the paper. US, MD and BS maintained, provided and prepared

the experimental evolution lines. All authors read and commented on drafts of the

manuscript.

References

Agrawal, A. F., & Wang, A. D. (2008). Increased transmission of mutations by low-condition females: Evidence for condition-dependent DNA repair. *PLoS Biology*, *6*(2), e30. https://doi.org/10.1371/journal.pbio.0060030

Agrawal, A. F., & Whitlock, M. C. (2012). Mutation Load: The Fitness of Individuals in Populations Where Deleterious Alleles Are Abundant. *Annual Review of Ecology, Evolution, and Systematics*, *43*(1), 115–135. https://doi.org/10.1146/annurev-ecolsys-110411-160257

Arnqvist, G., Stojković, B., Rönn, J. L., & Immonen, E. (2017). The pace-of-life: A sex-specific link between metabolic rate and life history in bean beetles. *Functional Ecology*, *31*(12), 2299–2309. https://doi.org/10.1111/1365-2435.12927

Bartosch -Harlid, A., Berlin, S., Smith, N. G. C., Mosller, A. P., & Ellegren, H. (2003). Life History and the Male Mutation Bias. *Evolution*, *57*(10), 2398–2406. https://doi.org/10.1111/j.0014-3820.2003.tb00251.x

Baur, J., & Berger, D. (2020). Experimental evidence for effects of sexual selection on condition-dependent mutation rates. *Nature Ecology & Evolution*, *4*(5), Article 5. https://doi.org/10.1038/s41559-020-1140-7

Beck, C. W., & Promislow, D. E. L. (2007). Evolution of Female Preference for Younger Males. *PLoS ONE*, *2*(9). https://doi.org/10.1371/journal.pone.0000939

Berger, D., Stångberg, J., Baur, J., & Walters, R. J. (2021). Elevated temperature increases genome-wide selection on de novo mutations. *Proceedings of the Royal Society B: Biological Sciences*, *288*(1944), 20203094. https://doi.org/10.1098/rspb.2020.3094

Berger, D., Stångberg, J., Grieshop, K., Martinossi-Allibert, I., & Arnqvist, G. (2017). Temperature effects on life-history trade-offs, germline maintenance and mutation rate under simulated climate warming. *Proc. R. Soc. B*, *284*(1866), 20171721. https://doi.org/10.1098/rspb.2017.1721

Biemont, J. C. (1973). Effects of the irradiations x on the reproductive power of Acanthoscelides obtectus Say (coleoptera: Bruchidae). *Ann. Zool. Ecol. Anim., v. 5, no. 4, pp. 581-591*. https://www.osti.gov/biblio/4119273-effects-irradiations-reproductive-power-acanthoscelides-obtectus-say-coleoptera-bruchidae

Blumenstiel, J. P. (2007). Sperm competition can drive a male-biased mutation rate. *Journal of Theoretical Biology*, *249*(3), 624–632. https://doi.org/10.1016/j.jtbi.2007.08.023

Bromham, L. (2011). The genome as a life-history character: Why rate of molecular evolution varies between mammal species. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *366*(1577), 2503–2513. https://doi.org/10.1098/rstb.2011.0014

Chen, H., Jolly, C., Bublys, K., Marcu, D., & Immler, S. (2020). Trade-off between somatic and germline repair in a vertebrate supports the expensive germ line hypothesis. *Proceedings of the National Academy of Sciences*, *117*(16), 8973–8979. https://doi.org/10.1073/pnas.1918205117

Chu, X.-L., Zhang, B.-W., Zhang, Q.-G., Zhu, B.-R., Lin, K., & Zhang, D.-Y. (2018). Temperature responses of mutation rate and mutational spectrum in an Escherichia coli strain and the correlation with metabolic rate. *BMC Evolutionary Biology*, *18*(1), 126. https://doi.org/10.1186/s12862-018-1252-8

Crean, A. J., & Immler, S. (2021). Evolutionary consequences of environmental effects on gamete performance. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *376*(1826), 20200122. https://doi.org/10.1098/rstb.2020.0122

de Vienne, C. M., Creveuil, C., & Dreyfus, M. (2009). Does young maternal age increase the risk of adverse obstetric, fetal and neonatal outcomes: A cohort study. *European Journal of Obstetrics & Gynecology and Reproductive Biology*, *147*(2), 151–156. https://doi.org/10.1016/j.ejogrb.2009.08.006

Đorđević, M., Savković, U., Lazarević, J., Tucić, N., & Stojković, B. (2015). Intergenomic Interactions in
Hybrids Between Short-Lived and Long-Lived Lines of a Seed Beetle: Analyses of Life History
Traits. *Evolutionary Biology*, *42*(4), 461–472. https://doi.org/10.1007/s11692-015-9340-9

Dowling, D. K., & Simmons, L. W. (2009). Reactive oxygen species as universal constraints in life-
history evolution. *Proceedings of the Royal Society B: Biological Sciences*, *276*(1663), 1737–
1745. https://doi.org/10.1098/rspb.2008.1791

Drake, J. W. (2009). Avoiding dangerous missense: Thermophiles display especially low mutation
rates. *PLoS Genetics*, *5*(6), e1000520.

Ellegren, H. (2007). Characteristics, causes and evolutionary consequences of male-biased mutation.
Proceedings of the Royal Society B: Biological Sciences, *274*(1606), 1–10.
https://doi.org/10.1098/rspb.2006.3720

Flatt, T., & Heyland, A. (Eds.). (2011). *Mechanisms of life history evolution: The genetics and*
physiology of life history traits and trade-offs. Oxford University Press.

Flatt, T., Tu, M.-P., & Tatar, M. (2005). Hormonal pleiotropy and the juvenile hormone regulation of
Drosophila development and life history. *BioEssays*, *27*(10), 999–1010.
https://doi.org/10.1002/bies.20290

Friedberg, E. C., Walker, G. C., Siede, W., & Wood, R. D. (2005). *DNA Repair and Mutagenesis*.
American Society for Microbiology Press.

Gao, Z., Moorjani, P., Sasani, T. A., Pedersen, B. S., Quinlan, A. R., Jorde, L. B., Amster, G., &
Przeworski, M. (2019a). Overlooked roles of DNA damage and maternal age in generating
human germline mutations. *Proceedings of the National Academy of Sciences*, *116*(19),
9491–9500. https://doi.org/10.1073/pnas.1901259116

Gao, Z., Moorjani, P., Sasani, T. A., Pedersen, B. S., Quinlan, A. R., Jorde, L. B., Amster, G., &
Przeworski, M. (2019b). Overlooked roles of DNA damage and maternal age in generating
human germline mutations. *Proceedings of the National Academy of Sciences*, *116*(19),
9491–9500. https://doi.org/10.1073/pnas.1901259116

Goldmann, J. M., Wong, W. S. W., Pinelli, M., Farrah, T., Bodian, D., Stittrich, A. B., Glusman, G.,
Vissers, L. E. L. M., Hoischen, A., Roach, J. C., Vockley, J. G., Veltman, J. A., Solomon, B. D.,
Gilissen, C., & Niederhuber, J. E. (2016). Parent-of-origin-specific signatures of de novo
mutations. *Nature Genetics*, *48*(8), 935–939. https://doi.org/10.1038/ng.3597

Guarente, L., & Kenyon, C. (2000). Genetic pathways that regulate ageing in model organisms.
Nature, *408*(6809), 255–262. https://doi.org/10.1038/35041700

Hadfield, J. D. (2010). MCMC Methods for Multi-Response Generalized Linear Mixed Models: The
MCMCglmm R Package. *Journal of Statistical Software*, *33*(2), 1–22.

Horta, F., Catt, S., Ramachandran, P., Vollenhoven, B., & Temple-Smith, P. (2020). Female ageing
affects the DNA repair capacity of oocytes in IVF using a controlled model of sperm DNA
damage in mice. *Human Reproduction*, *35*(3), 529–544.
https://doi.org/10.1093/humrep/dez308

Hurst, L. D., & Ellegren, H. (1998). Sex biases in the mutation rate. *Trends in Genetics*, *14*(11), 446–
452. https://doi.org/10.1016/S0168-9525(98)01577-7

Jaroudi, S., Kakourou, G., Cawood, S., Doshi, A., Ranieri, D. M., Serhal, P., Harper, J. C., & SenGupta,
S. B. (2009). Expression profiling of DNA repair genes in human oocytes and blastocysts using
microarrays. *Human Reproduction*, *24*(10), 2649–2655.
https://doi.org/10.1093/humrep/dep224

Johnson, S. C., Rabinovitch, P. S., & Kaeberlein, M. (2013). MTOR is a key modulator of ageing and
age-related disease. *Nature*, *493*(7432), 338–345. https://doi.org/10.1038/nature11861

Jones, O. R., Scheuerlein, A., Salguero-Gómez, R., Camarda, C. G., Schaible, R., Casper, B. B.,
Dahlgren, J. P., Ehrlén, J., García, M. B., Menges, E. S., Quintana-Ascencio, P. F., Caswell, H.,
Baudisch, A., & Vaupel, J. W. (2014). Diversity of ageing across the tree of life. *Nature*,
505(7482), Article 7482. https://doi.org/10.1038/nature12789

Jong, G. de, & Noordwijk, A. J. van. (1992). Acquisition and Allocation of Resources: Genetic (CO)
Variances, Selection, and Life Histories. *The American Naturalist*, *139*(4), 749–770.
http://www.jstor.org/stable/2462620

Katz, J., Khatry, S. K., LeClerq, S. C., Shrestha, S. R., West, K. P., & Christian, P. (2008). Miscarriage but
Not Stillbirth Rates Are Higher Among Younger Nulliparas in Rural Southern Nepal. *Journal of
Adolescent Health*, *42*(6), 587–595. https://doi.org/10.1016/j.jadohealth.2007.11.137

Kempner, E. S. (2001). Effects of high-energy electrons and gamma rays directly on protein
molecules. *Journal of Pharmaceutical Sciences*, *90*(10), 1637–1646.
https://doi.org/10.1002/jps.1114

Khandwala, Y. S., Zhang, C. A., Lu, Y., & Eisenberg, M. L. (2017a). The age of fathers in the USA is
rising: An analysis of 168 867 480 births from 1972 to 2015. *Human Reproduction*, *32*(10),
2110–2116. https://doi.org/10.1093/humrep/dex267

Khandwala, Y. S., Zhang, C. A., Lu, Y., & Eisenberg, M. L. (2017b). The age of fathers in the USA is
rising: An analysis of 168 867 480 births from 1972 to 2015. *Human Reproduction*, *32*(10),
2110–2116. https://doi.org/10.1093/humrep/dex267

Klass, M. R. (1977). Aging in the nematode Caenorhabditis elegans: Major biological and
environmental factors influencing life span. *Mechanisms of Ageing and Development*, *6*,
413–429. https://doi.org/10.1016/0047-6374(77)90043-4

Kong, A., Frigge, M. L., Masson, G., Besenbacher, S., Sulem, P., Magnusson, G., Gudjonsson, S. A.,
Sigurdsson, A., Jonasdottir, A., Jonasdottir, A., Wong, W. S. W., Sigurdsson, G., Walters, G. B.,
Steinberg, S., Helgason, H., Thorleifsson, G., Gudbjartsson, D. F., Helgason, A., Magnusson, O.
T., … Stefansson, K. (2012). Rate of de novo mutations and the importance of father's age to
disease risk. *Nature*, *488*(7412), 471–475. https://doi.org/10.1038/nature11396

Koppik, M., Baur, J., & Berger, D. (2023). Increased male investment in sperm competition results in
reduced maintenance of gametes. *PLOS Biology*, *21*(4), e3002049.
https://doi.org/10.1371/journal.pbio.3002049

Kurhanewicz, N. A., Dinwiddie, D., Bush, Z. D., & Libuda, D. E. (2020). Elevated Temperatures Cause
Transposon-Associated DNA Damage in C. elegans Spermatocytes. *Current Biology*, *30*(24),
5007-5017.e4. https://doi.org/10.1016/j.cub.2020.09.050

Lynch, M. (2008). The Cellular, Developmental and Population-Genetic Determinants of Mutation-
Rate Evolution. *Genetics*, *180*(2), 933–943. https://doi.org/10.1534/genetics.108.090456

Lynch, M. (2010). Evolution of the mutation rate. *Trends in Genetics*, *26*(8), 345–352.
https://doi.org/10.1016/j.tig.2010.05.003

Magnus, M. C., Wilcox, A. J., Morken, N.-H., Weinberg, C. R., & Håberg, S. E. (2019). Role of maternal
age and pregnancy history in risk of miscarriage: Prospective register based study. *BMJ*, *364*,
l869. https://doi.org/10.1136/bmj.l869

Maklakov, A. A., & Immler, S. (2016). The Expensive Germline and the Evolution of Ageing. *Current
Biology*, *26*(13), R577–R586. https://doi.org/10.1016/j.cub.2016.04.012

Maklakov, A. A., Immler, S., Lovlie, H., Flis, I., & Friberg, U. (2012). The effect of sexual harassment on
lethal mutation rate in female Drosophila melanogaster. *Proceedings of the Royal Society B:
Biological Sciences*, *280*(1750), 20121874–20121874.
https://doi.org/10.1098/rspb.2012.1874

Medawar, P. B. (1952). *An unsolved problem of biology*. Published for the College by H.K. Lewis.

Mills, M., Rindfuss, R. R., McDonald, P., te Velde, E., & on behalf of the ESHRE Reproduction and
Society Task Force. (2011a). Why do people postpone parenthood? Reasons and social policy
incentives. *Human Reproduction Update*, *17*(6), 848–860.
https://doi.org/10.1093/humupd/dmr026

Mills, M., Rindfuss, R. R., McDonald, P., te Velde, E., & on behalf of the ESHRE Reproduction and
Society Task Force. (2011b). Why do people postpone parenthood? Reasons and social
policy incentives. *Human Reproduction Update*, *17*(6), 848–860.
https://doi.org/10.1093/humupd/dmr026

Monaghan, P., Charmantier, A., Nussey, D. H., & Ricklefs, R. E. (2008). The evolutionary ecology of
senescence. *Functional Ecology*, *22*(3), 371–378. https://doi.org/10.1111/j.1365-
2435.2008.01418.x
Monaghan, P., & Metcalfe, N. B. (2019). The deteriorating soma and the indispensable germline:
Gamete senescence and offspring fitness. *Proceedings of the Royal Society B: Biological*
Sciences, *286*(1917), 20192187. https://doi.org/10.1098/rspb.2019.2187
Muller, H. J. (1950). Our load of mutations. *American Journal of Human Genetics*, *2*(2), 111–176.
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1716299/
Nussey, D. H., Kruuk, L. E. B., Donald, A., Fowlie, M., & Clutton-Brock, T. H. (2006). The rate of
senescence in maternal performance increases with early-life fecundity in red deer. *Ecology*
Letters, *9*(12), 1342–1350. https://doi.org/10.1111/j.1461-0248.2006.00989.x
Partridge, L., & Barton, N. H. (1996). On measuring the rate of ageing. *Proceedings of the Royal*
Society of London. Series B: Biological Sciences, *263*(1375), 1365–1371.
https://doi.org/10.1098/rspb.1996.0200
Promislow, D. E. L., & Tatar, M. (1998). Mutation and senescence: Where genetics and demography
meet. *Genetica*, *102*(0), 299–314. https://doi.org/10.1023/A:1017047212008
Puurtinen, M., Elo, M., Jalasvuori, M., Kahilainen, A., Ketola, T., Kotiaho, J. S., Mönkkönen, M., &
Pentikäinen, O. T. (2016). Temperature-dependent mutational robustness can explain faster
molecular evolution at warm temperatures, affecting speciation rate and global patterns of
species diversity. *Ecography*, *39*(11), 1025–1033. https://doi.org/10.1111/ecog.01948
R Core Team. (2020). *R: A Language and Environment for Statistical Computing*. R Foundation for
Statistical Computing. https://www.R-project.org/
Rahbari, R., Wuster, A., Lindsay, S. J., Hardwick, R. J., Alexandrov, L. B., Al Turki, S., Dominiczak, A.,
Morris, A., Porteous, D., Smith, B., Stratton, M. R., & Hurles, M. E. (2016a). Timing, rates and
spectra of human germline mutation. *Nature Genetics*, *48*(2), 126–133.
https://doi.org/10.1038/ng.3469
Rahbari, R., Wuster, A., Lindsay, S. J., Hardwick, R. J., Alexandrov, L. B., Al Turki, S., Dominiczak, A.,
Morris, A., Porteous, D., Smith, B., Stratton, M. R., & Hurles, M. E. (2016b). Timing, rates and
spectra of human germline mutation. *Nature Genetics*, *48*(2), 126–133.
https://doi.org/10.1038/ng.3469
Roff, D. (1993). *Evolution Of Life Histories: Theory and Analysis*. Springer Science & Business Media.
Sayres, M. A. W., & Makova, K. D. (2011). Genome analyses substantiate male mutation bias in many
species. *BioEssays*, *33*(12), 938–945. https://doi.org/10.1002/bies.201100091
Schärer, L., Rowe, L., & Arnqvist, G. (2012). Anisogamy, chance and the evolution of sex roles. *Trends*
in Ecology & Evolution, *27*(5), 260–264. https://doi.org/10.1016/j.tree.2011.12.006
Segami, J. C., Lind, M. I., & Qvarnström, A. (2021). Should females prefer old males? *Evolution*
Letters, *5*(5), 507–520. https://doi.org/10.1002/evl3.250
Ségurel, L., Wyman, M. J., & Przeworski, M. (2014). Determinants of Mutation Rate Variation in the
Human Germline. *Annual Review of Genomics and Human Genetics*, *15*(1), 47–70.
https://doi.org/10.1146/annurev-genom-031714-125740
Servedio, M. R., Powers, J. M., Lande, R., & Price, T. D. (2019). Evolution of sexual cooperation from
sexual conflict. *Proceedings of the National Academy of Sciences*, *116*(46), 23225–23231.
https://doi.org/10.1073/pnas.1904138116
Shabalina, S. A., Yampolsky, L. Y., & Kondrashov, A. S. (1997). Rapid decline of fitness in panmictic
populations of Drosophila melanogaster maintained under relaxed natural selection.
Proceedings of the National Academy of Sciences, *94*(24), 13034–13039.
http://www.pnas.org/content/94/24/13034
Sharp, N. P., & Agrawal, A. F. (2012). Evidence for elevated mutation rates in low-quality genotypes.
Proceedings of the National Academy of Sciences, *109*(16), 6142–6146.
https://doi.org/10.1073/pnas.1118918109

Shrim, A., Ates, S., Mallozzi, A., Brown, R., Ponette, V., Levin, I., Shehata, F., & Almog, B. (2011). Is Young Maternal Age Really a Risk Factor for Adverse Pregnancy Outcome in a Canadian Tertiary Referral Hospital? *Journal of Pediatric and Adolescent Gynecology*, *24*(4), 218–222. https://doi.org/10.1016/j.jpag.2011.02.008

Stearns, S. C. (n.d.). *THE EVOLUTION OF LIFE HISTORIES*.

Stojković, B., Đorđević, M., Janković, J., Savković, U., & Tucić, N. (2015). Heterosis in age-specific selected populations of a seed beetle: Sex differences in longevity and reproductive behavior. *Insect Science*, *22*(2), 295–309. https://doi.org/10.1111/1744-7917.12115

Stringer, J. M., Winship, A., Liew, S. H., & Hutt, K. (2018). The capacity of oocytes for DNA repair. *Cellular and Molecular Life Sciences*, *75*(15), 2777–2792. https://doi.org/10.1007/s00018-018-2833-9

Svetec, N., Cridland, J. M., Zhao, L., & Begun, D. J. (2016). The Adaptive Significance of Natural Genetic Variation in the DNA Damage Response of Drosophila melanogaster. *PLOS Genetics*, *12*(3), e1005869. https://doi.org/10.1371/journal.pgen.1005869

Thakur, D. R. (2012). *Taxonomy, Distribution and Pest Status of Indian Biotypes of Acanthoscelides obtectus (Coleoptera: Chrysomelidae: Bruchinae)—A New Record*. *44*(1), 189–195. https://doi.org/10.13140/RG.2.1.4209.3522

Tucić, N., Gliksman, I., Šešlija, D., Milanović, D., Mikuljanac, S., & Stojković, O. (1996). Laboratory evolution of longevity in the bean weevil (Acanthoscelides obtectus). *Journal of Evolutionary Biology*, *9*(4), 485–503. https://doi.org/10.1046/j.1420-9101.1996.9040485.x

Tucić, N., Šešlija, D., & Stanković, V. (2004). The short-term and long-term effects of parental age in the bean weevil (Acanthoscelides obtectus). *Evolutionary Ecology*, *18*(2), 187. https://doi.org/10.1023/B:EVEC.0000021090.51851.bd

Tucić, N., Stojković, O., Gliksman, I., Milanović, D., & Šešlija, D. (1997). Laboratory Evolution of Life-History Traits in the Bean Weevil (Acanthoscelides obtectus): The Effects of Density-Dependent and Age- Specific Selection. *Evolution*, *51*(6), 1896–1909. JSTOR. https://doi.org/10.2307/2411011

van de Pol, M., & Verhulst, S. (2006). Age-Dependent Traits: A New Statistical Model to Separate Within- and Between-Individual Effects. *The American Naturalist*, *167*(5), 766–773. https://doi.org/10.1086/503331

Venn, O., Turner, I., Mathieson, I., de Groot, N., Bontrop, R., & McVean, G. (2014). Strong male bias drives germline mutation in chimpanzees. *Science*, *344*(6189), 1272–1275. https://doi.org/10.1126/science.344.6189.1272

Wang, R. J., Thomas, G. W. C., Raveendran, M., Harris, R. A., Doddapaneni, H., Muzny, D. M., Capitanio, J. P., Radivojac, P., Rogers, J., & Hahn, M. W. (2020). Paternal age in rhesus macaques is positively associated with germline mutation accumulation but not with measures of offspring sociability. *Genome Research*, *30*(6), 826–834. https://doi.org/10.1101/gr.255174.119

Wickham, H. (2016). *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York. https://ggplot2.tidyverse.org

Williams, G. C. (1957). Pleiotropy, Natural Selection, and the Evolution of Senescence. *Evolution*, *11*(4), 398–411. JSTOR. https://doi.org/10.2307/2406060