1 Aberrant expression prediction across 2 human tissues

Florian R. Hölzlwimmer[1], Jonas Lindner[1], Georgios Tsitsiridis[1], Nils Wagner[1,2], Francesco Paolo
Casale[1,4,5], Vicente A. Yépez[1], Julien Gagneur[1,2,3,6]

[1] School of Computation, Information and Technology, Technical University of Munich, Garching,
Germany
[2] Helmholtz Association - Munich School for Data Science (MUDS), Munich, Germany
[3] Institute of Human Genetics, School of Medicine, Technical University of Munich, Munich, Germany
[4] Institute of AI for Health, Helmholtz Munich, Neuherberg, Germany
[5] Helmholtz Pioneer Campus, Helmholtz Munich, Neuherberg, Germany
[6] Institute of Computational Biology, Helmholtz Center Munich, Neuherberg, Germany

12 Abstract

Despite the frequent implication of aberrant gene expression in diseases, algorithms predicting
aberrantly expressed genes of an individual are lacking. To address this need, we compiled an
aberrant expression prediction benchmark covering 8.2 million rare variants from 633 individuals
across 49 tissues. While not geared toward aberrant expression, the deleteriousness score
CADD and the loss-of-function predictor LOFTEE showed mild predictive ability (1-1.6%
average precision). Leveraging these and further variant annotations, we next trained AbExp, a
model that yielded 12% average precision by combining in a tissue-specific fashion expression
variability with variant effects on isoforms and on aberrant splicing. Integrating expression
measurements from clinically accessible tissues led to another two-fold improvement.
Furthermore, we show on UK Biobank blood traits that performing rare variant association
testing using the continuous and tissue-specific AbExp variant scores instead of LOFTEE
variant burden increases gene discovery sensitivity and enables improved phenotype
predictions.

26 **Main**

Aberrant gene expression, gene expression levels outside the physiological range, is a frequent
cause of diseases. Aberrant underexpression of tumor suppressor genes and aberrant
overexpression of oncogenes are hallmarks of oncogenesis[1,2]. Moreover, aberrant gene
expression is a frequent cause of rare inheritable disorders[3–8] and contributes to risks for
common disease-associated traits[9].

Statistical methods to call expression outliers from RNA-seq data[10–13] applied to large cohorts
have enabled investigating the genetic basis of aberrant expression. Rare variants have been
found to be associated with expression outliers[14]. Specifically, rare variants including rare
structural variants and rare variants likely triggering nonsense-mediated decay such as
premature stop codons, frameshift, and splice-disrupting variants are enriched among
underexpression outliers[6,14,15]. Moreover, rare structural variants, notably gene duplication, were
found enriched among overexpression outliers[14,15].

Building on these findings, algorithms have been developed to prioritize which variants may be
the genetic cause of an expression outlier identified in an RNA-seq sample given the matched
genome[14,15]. However, there is no algorithm predicting aberrant expression caused by genetic
variants. A model predicting aberrant expression in multiple tissues and generalizing to unseen
variants could improve our ability to identify high-impact rare variants in large genomic cohorts
which in turn would aid in the identification of disease-associated genes and pinpointing
disease-causal variants.

To address this unmet need, we here establish a benchmark for rare-variant based prediction of
aberrant gene expression across human tissues (Fig. 1). While underexpression outliers can be
assumed to have strongly impaired or entirely lost function, the functional consequence of an
overexpression outlier is less clear, as it could as well result in a gain of function. We therefore
primarily focussed in this study on underexpression outliers and developed AbExp, a machine
learning model predicting aberrant underexpression in multiple tissues from an individual's rare
genetic variants. By integrating various variant annotations with tissue-specific isoform
proportions and expression variability, AbExp significantly outperforms other loss-of-function
annotators that were not explicitly developed for aberrant underexpression prediction. We
demonstrate how AbExp can be used in rare-variant association testing as well as phenotype
prediction on the UK Biobank dataset using 40 blood traits. Finally, we show that when gene
expression measurements from clinically accessible tissues are available, AbExp scores can be

integrated to improve the prediction of aberrant expression in non-accessible tissues.

Figure 1: **Study design. (a)** We aimed to predict whether protein-coding genes are aberrantly underexpressed
across 49 human tissues based on DNA and, optionally, RNA-seq data of clinically accessible tissues. Therefore, we
created a benchmark for aberrant underexpression prediction by processing 11,215 RNA-seq samples from 633
individuals across 49 tissues from GTEx. This yielded 11,200 underexpression outliers out of 99 million assessable
gene-sample pairs (0.01%). **(b)** Assessing various variant and tissue annotations, we found that predictions could be
significantly improved by weighting variant effects with tissue-specific isoform proportions and incorporating the
expression variability of a gene. Further integration of expression measurements from clinically accessible tissues led
to another two-fold improvement.

Results

68 A benchmark dataset of underexpression outliers across 49
69 human tissues

We first set out to predict which protein-coding genes are aberrantly underexpressed in which
tissues given an individual's genome. We focussed on protein-coding genes because they
represent the vast majority of transcripts implicated in human disease. Moreover, the

degradation mechanisms of messenger RNAs, which are strongly coupled with translation
notably via nonsense-mediated decay, substantially differ from the degradation mechanisms of
non-coding RNAs, a very heterogenous class of transcripts [16] whose degradation involves a
large variety of mechanisms[17–19]. To this end, we created a benchmark dataset using the
aberrant expression caller OUTRIDER[11] on 11,215 RNA-seq samples with paired whole-
genome sequencing data of the Genotype-Tissue Expression dataset[20] (GTEx v8), spanning 49
tissues and 633 individuals. For the whole-genome sequencing data, we selected GTEx version
7 since, unlike for version 8, structural variant calls were available. Although the choice against
GTEx version 8 reduced the number of underexpression outliers by 20%, it allowed us to
consider structural variants, which are important determinants of aberrant expression[15]. In each
tissue, we restricted the analysis to all protein-coding genes with an average read-pair count of
at least 450, an estimated minimal coverage required to detect 50%-reduction outliers[6]. Overall,
18,171 protein-coding genes out of 18,563 (97.9%) showed sufficient average coverage in at
least one tissue.

We defined a gene to be aberrantly underexpressed if OUTRIDER reported a false discovery
rate (FDR) lower than 0.05 and the expression was lower than expected (Methods). We further
removed samples with more than 20 outliers since such a high number of outliers could reflect
an unreliable fit of OUTRIDER due to technical reasons, for instance low sequencing depth[11], or
biological reasons, for instance an ancestry underrepresented in the cohort[6]. This exact cutoff
combination on FDR and on the number of outliers per sample was selected among 12 possible
combinations based on the generalisation performance of simple outlier predictors trained on
established outlier-associated features[15] (suppl. Fig. S1). Application of these cutoffs led to a
benchmark dataset of 11,200 underexpression outliers occurring in 3,240 genes and 10,999
samples (suppl. Fig. S2), amounting to nearly one underexpression outlier per sample on
average. With 99 million non-outliers in the dataset (gene-sample pairs assessed by
OUTRIDER but not identified as outliers), the proportion of underexpression outliers is
extremely low at just 0.01%. This creates a highly imbalanced prediction task.. A detailed
overview of dataset statistics including how many samples and genes remained after each
filtering step can be seen in suppl. table T1.

102 Integrating rare variant annotations to predict underexpression
103 outliers across tissues

We considered predicting aberrant expression for all 49 tissues for any protein-coding gene

from rare variants, here defined as variants occurring in at most two GTEx individuals and with a
gnomAD minor allele frequency of less than 0.1%. We reasoned that above 0.1% minor allele
frequency, variants are unlikely to cause an expression outlier since the average frequency
across samples of underexpression outliers among genes with sufficient RNA-seq coverage is
about 0.01%. Following observations from Ferraro and colleagues[15], we focussed on single-
nucleotide variants and short insertions and deletions located within the gene and up to 5,000
bp around the gene to fully cover promoter and transcription termination regions. Moreover, we
considered structural variants up to 1 megabase 5' of the gene[15]. We do not exclude that some
single nucleotide variants and short indels could also have strong effects when located further
than 5,000 bp of a gene. However, we did not consider this class of variants because sequence-
based models capturing distal enhancer effects are lacking[21].

We first investigated whether existing variant annotation tools, that were not developed for
predicting aberrant underexpression, showed informative signals. Among the variant
consequences annotated by Ensembl VEP, we found frameshifts, variants affecting start and
stop codons, and splicing variants to be strongly enriched among underexpression outliers,
consistent with previous reports[6,14,15] (Fig. 2a, suppl. Fig. S3). Variants including frameshifts,
splice-variants, and stop-gains, which introduce premature stop codons are known to be
strongly enriched within gene underexpression outliers as these often trigger nonsense-
mediated decay (NMD)[14]. LOFTEE is a tool that predicts a high-confidence subset of loss-of-
function associated variants, notably variants likely to trigger NMD, by implementing filters such
as removing stop-gained and frameshift variants that are within 50 bp of the end of the
transcript, or variants that affect splicing only in UTRs[22]. In GTEx more than 23% of the
aberrantly underexpressed genes had a LOFTEE variant, compared to non-outliers that had a
LOFTEE variant in less than 0.1% of the cases (Fig. 2a).

Moreover, we hypothesized that the deleteriousness score CADD[23] could also be predictive of
aberrant gene expression. The advantage of CADD, which was trained to distinguish between
simulated de novo variants and variants that have arisen and become fixed in human
populations, is that it provides a score for any variant. We found that the median CADD score of
rare GTEx variants was ~17 times higher among underexpressed genes than among non-
outliers (Fig. 2b).

Despite these enrichments, LOFTEE and CADD by themselves showed limited predictive value.

Using the sole LOFTEE-positive variants recalled 23.2% of the underexpression outliers at a
precision of 7.1% (Fig. 2c). Besides a small spike to 11.6% precision at 0.2% recall, CADD
never reached the same precision nor the same recall as LOFTEE (Fig. 2c). Next, we trained a
non-linear model that integrated all the above-mentioned features to quantitatively predict the
OUTRIDER z-score (Methods). Predicting the underlying quantitative z-scores turned out to
lead to better classifiers than directly predicting the binary classes of outliers and non-outliers.
Moreover, ranking based on the predicted z-scores uniformly outperformed ranking based on
CADD scores on held-out data (Fig. 2c). The integrative model reached the same precision at
the same recall as filtering for LOFTEE variants with the added value of providing a continuous
score, thereby allowing for applying more stringent cutoffs to yield a higher precision (up to
11%, Fig. 2c). Lastly, the advantage of the integrative model over CADD and LOFTEE was
observed in aggregate (Fig. 2c) as well as across individual tissues according to the average
precision (measured by the area under the precision-recall curve here and elsewhere, AUPRC,
Fig. 2d).
Additionally, we adapted Watershed, a method that was originally designed to identify the most
likely variant responsible for an observed expression outlier, to fit our task[15]. Watershed is a
Bayesian approach that relies on strong modeling assumptions. In particular, Watershed
represents the outlier status as a categorical variable and models variant features
independently. Using the Watershed features, we found that predicting tissue-specific z-scores
with a non-linear model largely outperforms the Watershed-derived outlier predictor (suppl. Fig.
S4). Altogether, these investigations demonstrate that outlier prediction benefits from a direct
and quantitative treatment and from flexible, non-linear machine learning models.

Figure 2: **Integrating rare variant annotations to predict underexpression outliers across tissues. (a)** Proportion of underexpressed outliers (red), and non-outliers (gray) with a rare variant of a given annotation (rows, Methods). Error bars mark 95% binomial confidence intervals. Notably, rare variants likely to trigger nonsense-mediated decay are strongly represented in underexpressed genes. **(b)** Distribution of CADD scores in different outlier classes. The numbers of data points are labeled for each box. **(c)** Precision-recall curve for all tissues combined. LOFTEE shows up as a single point because it is a binary filter. **(d)** Distribution of average precision (AUPRC) across 27 GTEx tissue types (Methods). *P*-values were obtained using a paired Wilcoxon test. A non-linear model based on LOFTEE, CADD and VEP annotations significantly outperforms existing methods. **For all boxplots:** Center line, median; box limits, first and third quartiles; whiskers span all data within 1.5 interquartile ranges of the lower and upper quartiles.

Accounting for tissue-specific isoform expression improves predictions

By construction, the predictions of this first model were independent of the tissue as neither the variant annotations nor the considered transcript isoforms were tissue-specific. However, since the transcript isoforms of a gene are often expressed at different proportions across tissues,

variants can have tissue-dependent effects[24]. For example, ENST00000358514, the canonical

transcript of *PSMB10*[25], was estimated to generate only about 4% of *PSMB10* total gene

expression in putamen (Methods). The vast majority (91%) of *PSMB10* gene expression in

putamen was attributed to another transcript, ENST00000570985. Conversely, in fibroblasts,

the canonical transcript contributed to nearly 48% of the total gene expression. Exon 4 is not

included in the transcript ENST00000570985 but is included in the canonical transcript

ENST00000358514, explaining why a frameshift variant in exon 4 was associated with a high

impact on gene expression in cultured fibroblasts but showed a limited effect in putamen (Fig.

3a,b).

Generally, we found that only 30% of the canonical transcripts contributed more than 90% of the

total expression for their respective genes and that as much as 18% of the canonical transcripts

contributed to less than 10% of the total expression for their respective genes (suppl. Fig. S5).

Therefore, relevant information is lost when considering the variant consequence assigned to a

single isoform, even if it is annotated as the canonical one. To address this issue, we calculated

the isoform composition in every tissue and weighted the VEP consequences and LOFTEE

classification of each variant by the proportion of affected transcripts per gene and tissue

(Methods). Training the model using these tissue-specific weighted annotations increased the

average precision by 58% to reach 3.2% in median across tissues (Fig. 3c).

Figure 3: **Tissue-specific features improve the model prediction. (a)** Sashimi plot of *PSMB10* for two individuals,
one carrying no rare variant in this region (control, upper tracks), and one carrying a heterozygous frameshift variant
(dashed line and lower tracks), in cultured fibroblasts (top) and putamen (bottom). The frameshift variant is located on
exon 4 which is included on the canonical transcript (ENST00000358514) but not on transcript ENST00000570985.
On the right, the bar plots show the transcript expression proportions on each tissue on average across GTEx. In
fibroblasts, the rare variant is associated with an approximately 25% reduction of RNA-seq coverage in this window
whereas in putamen no major RNA-seq coverage change is observed. **(b)** Fold change of gene expression against
normalized gene expression rank for *PSMB10* in fibroblasts and putamen (basal ganglia) brain tissues. *PSMB10* is
an expression outlier (red) in individual 2 in fibroblast but not in putamen, consistent with the rare variant triggering
nonsense-mediated decay and leading to a strong gene expression reduction in the tissue for which the exon 4-
containing transcript is the major isoform. **(c)** Left: Distribution of average precision (AUPRC) across 27 GTEx tissue
types. *P*-values were obtained using the paired Wilcoxon test. AbExp combines various variant and tissue
annotations to predict aberrant gene expression and outperforms LOFTEE by about sevenfold. Right: Precision-recall
curve for all tissues combined. **(d)** Fold change of gene expression against normalized gene expression rank for
LTBP3 in tibial artery, an autosomal recessive gene whose defect can lead to dental anomalies and short stature[26],
and for *OR2W3* in blood, an olfactory gene whose defect should not impair the viability of an individual[27]. Expression
outliers are highlighted in red. *LTBP3* is tightly regulated with a fold change range of ± 20% among non-outliers. The
individual marked in red carries a heterozygous frameshift variant that associates with 30% reduction and which is
detected as an outlier. In contrast, *OR2W3* shows very large variations where individuals with 30% reductions are not
outliers. **(e)** Distribution of gene expression fold changes for underexpression outliers (red) and non-outliers (grey)
harbouring a frameshift variant per decile of expression variability (biological coefficient of variation). **(f)** Left:
Proportion of underexpressed outliers (red; n=2,051), non-outliers (gray; n=2,749,843), and overexpressed outliers
(blue; n=1,177) with a rare variant within a certain distance of the transcription start site (TSS), normalized by the
length of each distance interval. Error bars mark 95% binomial confidence intervals. Right: Distribution of Enformer
scores in different outlier classes among rare variants within a certain distance of the TSS. **For all boxplots:** Center
line, median; box limits, first and third quartiles; whiskers span all data within 1.5 interquartile ranges of the lower and
upper quartiles.
218

219 Incorporating the tissue-specific gene expression variability
220 further improves predictions

Similar to other statistical models for RNA-seq data, OUTRIDER includes a measure of gene
expression variability called the biological coefficient of variation[11,28]. The biological coefficient
of variation is used instead of the plain coefficient of variation to account for sampling noise,
which particularly affects low RNA-seq read counts[28]. In our benchmark dataset, the biological
coefficient of variation captures the expression variability of genes per tissue across the GTEx
population. We reasoned that the same expression fold changes could cause an expression
outlier for a gene with low expression variability but not for a gene with high expression
variability. Indeed, we observed that the minimal fold-change among expression outliers

decreased with the biological coefficient variation (suppl. Fig. S6). Therefore, a given relative
reduction in gene expression can lead to aberrant expression in one gene or tissue but not
necessarily in another. For instance, a 30% reduction of the gene *LTBP3* in tibial artery
expression resulted in an outlier. In contrast, a 30% expression reduction in blood of *OR2W3*
would not lead to an outlier as *OR2W3* shows a very large gene expression variation in blood
ranging between 10% and 230% (Fig. 3d). It is not surprising that *OR2W3*, one of the over 800
human olfactory receptor genes[27] for which dysfunction is likely benign, exhibits more
expression variability than *LTBP3*, a gene whose dysfunction is associated with dental
anomalies and short stature[26].

Next, we aimed to improve our underexpression outlier predictor by adjusting for expression
variability. We first considered modeling expression fold-changes from variants and then
converting the predicted fold-changes into a *z*-score, under the assumption that variants affect
gene expression fold-changes independently of the expression variability. Using variants likely
triggering NMD to test this assumption, we noticed however that the same class of variants
associated with lower fold-changes among genes with lower expression variability (Fig. 3e,
suppl. Fig. S7), perhaps because genes with low expression variability are subject to regulatory
buffering mechanisms[29,30]. Therefore, we opted for a more general modeling approach in which
the biological coefficient of variation is provided as an input feature along with variant
annotations to a non-linear model predicting the *z*-score. This model increased the performance
by more than 63% to 5.0% average precision (median across tissues, Fig. 3c). Altogether, these
results show that gene expression variability must be taken into account when predicting
aberrantly expressed genes, and that predicting z-score rather than fold-change is more
relevant for variant interpretation.

252 Contribution of aberrant splicing variants, core promoter variants, 253 and structural variation

Aberrant splicing isoforms often contain a premature termination codon and as a consequence,
yet not always, are degraded by NMD[31]. Our model already included splice site information, as it
was based on LOFTEE, CADD, and tissue-specific isoform weighting of VEP annotations.
However, these features ignored splice sites that are not part of the genome annotations upon
which these tools are built. We have recently developed a model called AbSplice that predicts
aberrant splicing across tissues using a more comprehensive map of splice sites, including
unannotated weak splice sites, and their tissue-specific usage[32]. We found that AbSplice scores

were in median 10 times higher among underexpression outliers than among non-outliers
(suppl. Fig. S8a). Integrating AbSplice scores significantly increased the performance to 5.9%
average precision in median across tissues. In contrast, adding SpliceAI did not show any
significant improvement (suppl. Fig. S8b), in agreement with AbSplice improving over SpliceAI
for predicting tissue-specific aberrant splicing[32].

To better capture variants affecting transcription, we considered Enformer[33], a state-of-the-art
deep learning model that predicts thousands of transcription-related genome-wide assays
including 635 CAGE tracks of the ENCODE project, based on a 200-kb sequence context.
Building on previous work[21], we mapped Enformer-predicted CAGE tracks to gene expression
for the 49 GTEx tissues with regularized linear regression models (Methods). We found that
using elastic net models, rather than ridge regression, and modeling effects on canonical
isoforms, rather than on mixtures of tissue-specific isoforms, resulted in the best median
AUPRC across tissues (suppl. Fig. S9 a,b). This Enformer-based model was predictive for
expression outliers in the correct direction but only in the close vicinity of the transcription start
site (-50 bp to +200 bp, Fig. 3f and suppl. Fig. S9c,d). On the one hand, this observation may
reflect the biology of genetic determinants of aberrant expression, whereby variants with
extreme effects on gene expression may primarily lie in the core promoter, which extends to 50
bp on either side of the transcription start site in eukaryotes[34]. Consistently, we found a higher
proportion of outlier-associated variants in the core promoter region, independently of Enformer
predictions (Fig. 3f). On the other hand, the poor precision of Enformer beyond this narrow
window around the transcription start site may reflect the difficulty that Enformer has in
capturing long-range effects[21]. Despite the strong enrichment in underexpressed outliers, adding
Enformer to the AbExp feature set increased the median AUPRC only minimally to 6.0% across
tissues, albeit significantly.

We next considered the inclusion of structural variants overlapping transcripts (Methods).
Transcript ablations, which could be inferred in GTEx from the structural variant deletion calls,
were especially predictive of underexpression. Overall, 33 out of the 43 GTEx individuals
harboring transcript ablations also showed an expression outlier in at least one tissue. Including
structural variants into the model resulted in a large gain in precision among the top-ranked
predictions and increased the average precision to 10.2% in median across tissues (Fig. 3c).
We also considered structural variants up to 1 megabase 5' of the transcription start site, which
were enriched for expression outliers in agreement with Ferraro et al.[15] (suppl. Fig. S10a).

However, as the integration of these variants did not lead to a significant improvement (suppl.
Fig. S10b) we did not consider the distant structural variants further.

In the following, we refer to the model which integrates all significant features mentioned so far
as AbExp. AbExp takes as input a set of variants within 5,000 bp of any annotated transcript of
a protein-coding gene and returns a predicted z-score for each of the 49 tissues. For user
convenience, we furthermore suggest a high-confidence cutoff (AbExp < -3.84) corresponding
to 50% precision and 7.8% recall on our benchmark data, and a low-confidence cutoff (AbExp <
-1.64) corresponding to 20% precision and 21.5% recall. The high-confidence cutoff leads to
about 1 positive prediction every 6 GTEx samples, whereas the low-confidence cutoff leads to
about 1.2 positive predictions per GTEx sample, which is approximately the number of outliers
per sample (1.02). AbExp performs especially well in predicting the impact of NMD-associated
variants, outperforming other methods in these thanks to a more effective ranking (suppl. Fig.
S11). However, AbExp cannot reliably predict the impact of variants in categories rarely
associated with expression outliers in GTEx, such as missense and UTR variants. Larger
datasets and complementary models will be needed to improve on these variant categories.

Although our primary focus was on predicting underexpression outliers, we leveraged the fact
that AbExp quantitatively predicts z-scores to assess its performance for overexpression outliers
(suppl. Fig. S12). We found that the performance for overexpression outlier prediction was
substantially lower compared to underexpression outliers (median AUPRC 3.4% across
tissues). By far, the major predictive features were structural variants, yet Enformer and isoform
proportions also contributed significantly. Not surprisingly, LOFTEE, CADD, AbSplice, and other
loss-of-function associated variant effects were not predictive for overexpression. These results
align with the variant category enrichments reported by Ferraro et al., which showed that
duplications and variants near the TSS were the only categories specifically enriched among
overexpression outliers, but not among underexpression outliers[15]. Altogether, these results
show that AbExp can also be employed for over-expression outlier prediction, yet with a much
lower performance than for underexpression outliers.

320 AbExp replicates on independent datasets

We next assessed how AbExp performance replicated on two independent datasets. The first
dataset consisted of 295 individuals suspected to be affected by a mitochondrial disorder[6] with
whole-exome sequencing data paired with RNA-seq from fibroblasts. The second dataset

consisted of 233 whole-genome sequencing samples with RNA-seq from iPSC-derived motor
neurons from the AnswerALS research project[35]. Structural variant calls, and thus transcript
ablation calls, were not available on either dataset. Also, we did not compute Enformer scores
due to computational constraints and its limited added value for outlier prediction. Moreover, we
observed that the recall for all methods was twice as low on the ALS dataset than in GTEx and
in the mitochondrial disorder dataset (suppl. Fig. S13), perhaps because of poorer expression
outlier calls, a stronger role of epigenetic and trans-regulatory effects, or combinations thereof.
Taking these differences into account, our results on those two independent datasets were in
agreement with the evaluation on GTEx. We found that AbExp without transcript ablation
annotation significantly outperformed LOFTEE and CADD by two to three times larger average
precision (suppl. Fig. S13). Here too, AbExp without structural variants and Enformer allowed
for slightly better precision at the same recall than LOFTEE filtering, while offering a continuous
score allowing for reaching much higher precisions.
One concern of having used the biological coefficient of variations in the GTEx benchmark is
that they were computed on the very same RNA-seq data as those used to compute the ground
truth expression outliers. Nevertheless, including the biological coefficient of variation computed
on GTEx significantly improved the predictions on the mitochondrial disorder dataset and was
on par on the ALS dataset (suppl. Fig. S13), showing that the biological coefficient of variation
contributes with independent information to outlier prediction.

343 AbExp prioritizes deleterious variants

Having established and independently validated AbExp as an aberrant expression predictor, we
next assessed its potential for identifying deleterious variants. To this end, we first considered
the capacity of AbExp at distinguishing pathogenic from benign and likely benign variants as
reported in the ClinVar database (Methods). For each variant, we retained the minimum AbExp
score, i.e. the most impactful underexpression prediction across all tissues. We found that
AbExp improved over CADD, LOFTEE, and AbExp without biological coefficient of variation in
the high precision range with precision exceeding 99% up to 47% recall (Fig. 4a). However, at
lower precision and higher recall both CADD and LOFTEE outperformed AbExp.
One limitation of this benchmark is that ClinVar submitters are using established bioinformatics
tools including LOFTEE and CADD to identify pathogenic variants, introducing a bias in favor of
those tools. To circumvent this issue, we next examined a dataset devoid of human annotations
that could indicate that AbExp allows prioritizing deleterious variants. Reasoning that more
frequent variants in the human population are less likely to be deleterious, we stratified all

possible variants within 5 kb of protein-coding genes into four frequency categories: absent in

gnomAD, singletons (found only once), rare (more than once but less than 0.1%) or common

(more than 0.1%). AbExp was significantly and more strongly associated with these variant

categories than LOFTEE, CADD, and AbExp without biological coefficient of variation (Fig. 4b).

Moreover, focussing on variants absent in gnomAD, AbExp high-impact variants were more

significantly enriched than alternative methods among highly genetically constrained genes[22],

i.e. for those genes for which one can expect the strongest phenotypic impact (first LOEUF

deciles, Fig. 4c). Notably, when the biological coefficient of variation was not accounted for in

AbExp, the relative enrichment on constrained genes was reduced (Fig. 4c). This is consistent

with our observation that more genetically constrained genes showed lower expression

variability (Fig. 4d), aligning with previous findings in primates[36]. This interpretation was also

consistent when performing the enrichment analyses against LOEUF for further allele frequency

categories. At the extreme, common variants classified as high-impact by AbExp were depleted

in the first LOEUF decile and enriched in the last LOEUF decile (suppl. Fig. S14). Intermediate

behaviors were observed for singletons (variants observed only once in gnomAD) and rare

variants (non-singleton gnomAD variants with MAF <0.1%). Altogether, these results indicate

that AbExp can be used as an informative variant prioritization algorithm.

Fig. 4: AbExp prioritizes deleterious variants. (a) Precision-recall curve of AbExp (trained with and without using the biological coefficient of variation), CADD, and LOFTEE on distinguishing pathogenic from benign and likely begin variants in ClinVar. LOFTEE as a binary predictor is shown as a single point. The dashed vertical bars denote the high and low confidence cutoffs of AbExp. **(b)** Odds-ratio of high-impact variants among absent, singleton, rare (MAF < 0.1%) and common SNVs in gnomAD for the models shown in **a**. The analysis is restricted to variants within 5 kb of protein-coding genes. Error bars show Wald 95% confidence intervals from logistic regression fits. The high-impact cutoffs for CADD and AbExp without BCV were set to match the quantile of the high-impact cutoff of AbExp. **(c)** As in **b** among SNVs absent in gnomAD as a function of gene LOEUF decile[22]. Genes with a high LOEUF are more tolerant to loss of function. Error bars as in **b. (d)** BCV versus LOEUF across all genes and tissues. The black line shows a running median between LOEUF and BCV highlighting the two genes from Fig. 2d. The autosomal recessive gene *LTBP3* has a low LOEUF, denoting a low loss-of-function tolerance. In contrast, the olfactory gene OR2W3 has a high LOEUF, denoting a large loss-of-function tolerance.

AbExp improves rare variant association testing and phenotype prediction

The rise of large exome-sequencing and genome-sequencing cohorts empowers rare variant association testing (RVAT) which helps pinpoint causal genes for traits[37,38] and enables

improved phenotype predictions, particularly among individuals showing extreme phenotypes[39].
RVAT consists of identifying genes within which the occurrence of likely high-impact variants
are associated with a trait[40,41]. To this end, accurate prediction of high-impact variants can be
advantageous, suggesting that AbExp may have the potential to improve RVAT. To test this
hypothesis, we considered 40 continuous blood traits including high-density lipoprotein
cholesterol, glucose, and urate levels (suppl. table T2) from the UK Biobank 200k exome
release[42]. We chose blood traits for this proof-of-concept investigation since they are well-
studied and frequently measured to diagnose and monitor chronic disease conditions.
Moreover, RVAT methods are typically better calibrated on continuous traits as opposed to
binary traits[43,44].

To ease comparisons between variant annotations, we used linear regression as a common
framework for rare variant association testing (RVAT). As a realistic baseline, we considered
RVAT based on LOFTEE variants, similar to the Genebass study, a phenome-wide study
leveraging UK Biobank data[38]. For this baseline model, gene-trait association was tested by
regressing the trait against the number of LOFTEE variants. The second model leveraged that
AbExp is both quantitative and tissue-specific. To this end, we considered the lowest AbExp z-
score across all rare variants for each of the 49 tissues. The gene-trait association was tested
by regressing the trait against the resulting 49 values. Two further models were considered by
regressing against the minimum and against the median of the 49 values. To adjust for other
relevant factors and effects due to common genetic variation, all four models included as
covariates sex, age, the first 20 genetic principal components, a polygenic risk score predicting
the trait, and common variants reported to be associated with the trait and located 250,000 bp
around the gene (Methods). We fitted every model on two thirds of the dataset for gene-trait
association discovery. Phenotype permutation analysis indicated that all models were calibrated
(suppl. Fig. S15).

Using AbExp predictions for the 49 tissues, we identified in total 30% more gene-trait
associations compared to the LOFTEE-based model (Fig. 5a), showing that AbExp can
significantly (P = 5.2x10^{-4}) improve RVAT-based gene discovery. Notably, association testing
using the tissue-specific predictions outperformed aggregated forms of the AbExp score in most
cases by finding more gene-trait associations (suppl. Fig. S16). In some instances, we could
rationalize which tissues showed the most significant associations. This was the case for the

gene encoding Apolipoprotein B, a major constituent of triglyceride-rich lipoproteins synthesized
in the liver and whose predicted aberrant underexpression in the liver was found to be
negatively associated with blood triglyceride levels (suppl. Table T3). However, the high
correlation between the AbExp scores per gene across tissues can make the estimation of each
individual tissue-specific coefficient unstable, and, therefore, their interpretation should be done
with caution.

Having shown that AbExp can improve the gene-discovery sensitivity of RVAT, we next
assessed its utility in phenotype prediction. To this end, we used the remaining third of the
dataset which was not used for gene-trait association discovery. Specifically, we fitted gradient
boosted trees models predicting the traits given the AbExp scores on the one hand or the
number of LOFTEE variants on the other hand, of the genes discovered on the first two thirds of
the data. Compared to linear regression models, gradient boosted trees can capture non-linear
relationships through an ensemble of decision trees. These models were controlled for sex, age,
the first 20 genetic principal components, and a polygenic risk score predicting the trait
(Methods). The predictions rarely differed between the common variant-based model and the
model further including AbExp scores of rare variants, as exemplified for the Alanine
aminotransferase blood levels (Fig. 5b). For this trait, predictions for less than 0.5% of all
individuals differed between the two models by more than 1 standard deviation of the population
trait distribution (Fig. 5b). Remarkably, the trait values of those differing individuals tended to
deviate largely from the population average, suggesting that the model integrating AbExp
scores especially improves the predictions of individuals with extreme phenotypes that common
variants cannot explain (Fig. 5c).

On held-out data, the phenotype prediction model based on AbExp scores significantly
increased the amount of explained variation (R^2) over the model based on LOFTEE in 45% of
the traits and never significantly decreased R^2 (Fig. 5d). Considering the number of individuals
differing by more than 1 standard deviation to the common-variant based model, the AbExp
based model improved the prediction of 816 individuals across the 40 blood traits, while the
LOFTEE-based model only improved the prediction of 264 individuals (Fig. 5e). Moreover, the
advantage of using AbExp scores were similarly observed when predicting phenotypes with
regularized linear regression instead of gradient boosted trees (suppl. Fig. S17a,b), yet yielding
less accurate phenotype predictions (suppl. Fig. S17c). These results using regularized linear
regression, a method which like gradient boosted tree allows for robust fitting compared to

standard linear regression but that does not model non-linearities, further confirm the added
value of the tissue-specific predictions of AbExp for rare variant association testing.

Altogether, these results show that AbExp provides useful variant annotation for gene-trait
association discovery by rare variant association testing and for building improved genetic risk
scores.

Figure 5: AbExp improves rare variant association testing and phenotype prediction. (a) Number of genes
associating with different traits using a model based on LOFTEE or AbExp. **(b)** Alanine aminotransferase level
predicted using a model solely based on common variants (y-axis) against predictions using a model based on

common variants and AbExp scores (x-axis). Individuals whose predictions differed by more than 1 standard
deviation of the population trait distribution are marked in orange. **(c)** Alanine aminotransferase measurements
against predictions based on common variants and AbExp scores. Orange data points as in b). **(d)** Relative R^2
increase between AbExp-based and LOFTEE-based predictions across traits. Traits with a significant difference
between both models are marked red (two-sided paired t-test, nominal $P < 0.05$). Error bars show the standard
deviation among 5 cross-validation folds. **(e)** Positive bars show the number of individuals with an error reduced by at
least one standard deviation in the trait scale and therefore improved prediction, negative bars show the number of
individuals with an error increased by at least one standard deviation in the trait scale and therefore worse prediction
of the AbExp-based model (green) and the LOFTEE-based model (grey). All data presented in b-e) are computed on
held-out folds of a 5-fold cross-validation within a third of the UKBB data not used for the gene discovery shown in a).

Incorporating RNA-seq from clinically accessible tissues (CATs) boosts prediction performance

RNA sequencing is becoming increasingly popular for rare disease diagnostics as a
complementary assay to genome or exome sequencing as it allows the direct measurement of
aberrant gene regulation[4–6,45–47]. However, many rare disorders are suspected to originate from
tissues that can only be very invasively sampled such as the brain or the heart. We and others
[6,48] have shown that clinically accessible tissues (CATs), notably skin fibroblasts and to a lesser
extent whole blood, share a substantial fraction of expressed genes with non-CATs and,
therefore, are likely to capture aberrant expression occurring in non-CATs. The GTEx dataset –
a dataset of post-mortem samples – offers a unique opportunity to test the validity of this
assumption as it provides matched samples for CAT and non-CAT tissues. We found that the
mere ranking of genes according to their OUTRIDER z-score in fibroblast RNA-seq samples led
to an average precision in predicting underexpression outliers in non-CATs of 20.1% (median
across tissues), significantly larger than the genome-based predictor AbExp (10.1%, $P = 1.6$x10^-
[6], Fig. 6a). Next, we developed a model taking as input AbExp, whether the gene is expressed
in a CAT and, if so, its OUTRIDER z-score. Using RNA-seq from skin fibroblasts to predict
aberrant underexpression in all other tissues, this model reached an average precision of 23.8%
in median across tissues (Fig. 6a). Consistent with previous work based on shared expressed
genes[6,48] and our work on aberrant splicing prediction[32], fibroblasts turned out be more
informative than whole blood (median average precision 9.3% using RNA-seq only and 17.5%
when integrating AbExp, Fig. 6b), owing to blood expressing less genes than fibroblasts (See
e.g. ref. [6]). Altogether, we have established a method integrating direct measurements of
aberrant expression from RNA-seq data in a CAT along with genomic variant annotations to
predict aberrant underexpression in other tissues. Doing so, we showed that integrating RNA-
seq from fibroblasts yields a substantial improvement as it doubles the average precision over
using genomic variants alone.

Figure 6: Combining RNA-seq measurements from clinically accessible tissues with AbExp further improves the prediction performance. (a) Left: Distribution per predictor (rows) of average precision (AUPRC) across 26 tissue types excluding skin tissues (Methods). Center line, median; box limits, first and third quartiles; whiskers span all data within 1.5 interquartile ranges of the lower and upper quartiles. *P*-values were obtained using the paired Wilcoxon test. The "Gene expression (CAT)" predictor ranks genes according to their OUTRIDER *z*-score in fibroblasts RNA-seq data. Right: Precision-recall curve aggregated across the same GTEx tissues as in the left panel. LOFTEE as a binary predictor is shown as a single point. **(b)** as in (a) using Whole blood as CAT and all other tissues as non-CAT.

Discussion

Altogether, we established a benchmark dataset for aberrant gene underexpression prediction in 49 human tissues, addressing an unmet need in the area of high-impact variant effect prediction. We developed AbExp, a machine learning model predicting aberrant underexpression across tissues by integrating existing variant annotations with tissue-specific gene expression variability and transcript isoform composition. AbExp outperformed existing

variant annotation tools by up to 7-fold in average precision. Enhanced enrichments for
pathogenic variants and for variants not yet observed in gnomAD, particularly among
mutationally constrained genes, indicate that AbExp is a promising tool for variant prioritization
in clinical diagnostics. Using UK Biobank blood traits, we demonstrated that the continuous and
tissue-specific AbExp scores provide added information over the state-of-the-art putative loss of
function classifier LOFTEE for rare variant gene association testing as well as for phenotype
prediction. Finally, we showed that AbExp scores can be combined with gene expression
measurements from clinically accessible tissues to predict aberrant expression in other tissues
yielding an increased prediction performance by 2-fold over AbExp.

Refining the predictor, while our primary objective, also shed light into the biology of
underexpression outliers. We found that gene expression variability plays a dual role in this
context. On the one hand, altering expression of a gene is more likely to result in an outlier if the
expression of the gene varies little in the population than if it varies largely. On the other hand,
each variant category was associated with milder fold-changes for genes with lower expression
variability, indicating the involvement of regulatory mechanisms that confer robustness to
genetic perturbations. We modeled the outcome of these two counteracting phenomena with a
non-linear model trained from the data. Future biophysical investigations unraveling these
buffering mechanisms could help improving the predictions and more generally improve variant
interpretation. Our work also confirmed the importance of nonsense-mediated decay, which
underpins a substantial proportion of the outliers, and the need to take tissue-specific transcript
isoform into account when interpreting splice-affecting and nonsense variants as pioneered by
Cummings and colleagues[24].

This study has limitations. A basic assumption of AbExp is that an underexpression outlier is
caused by a rare variant. However, it is possible that a rare combination of frequent genetic
variants causes an expression outlier. Also, damage caused by one variant might be recovered
by another variant, e.g. a second frameshift variant recovering the frame after a first frameshift
variant. AbExp does not evaluate combinations of variants, which would require more complex
modeling in particular by taking phasing into account. Moreover, AbExp covers the variants up
to 5 kb away from transcript boundaries, missing middle-range and long-range enhancers.
Integrating structural variants located further 5' of the transcription start did not lead to
significant performance improvement. Future work could expand to distal variants, for instance if
sequence-based models of gene expression improved at this task[21]. Also, this work was

focused on cis-acting regulation by considering only variants located within or near the genes.
The effect of trans-acting gene regulation, which would need a very different modeling paradigm
than investigating here in order to capture regulatory networks, remains to be addressed. The
current recall remains modest. However, this can be an underestimation of the true recall since
it is unclear how many of the outliers that we aim to predict are artifacts, due to mosaicism, non-
genetic causes, or because they are measurement errors. Moreover, it is unclear whether a
gene expression outlier is caused by a change of transcript length or transcript abundance.
Indeed, our outlier classifications are based on read counts across annotated exons, which may
include shorter transcripts among down-regulated outliers[6]. Given that GTEx data relies on
short-read sequencing, accurately determining transcript length is challenging. Perhaps the
advance of long-read RNA-sequencing will allow to disambiguate these cases in the future.
Finally, we have used here an outlier calling method that was applied for each tissue separately
and that could not leverage statistical evidence across tissues. Future aberrant expression
predictors may be improved with better outlier calls.

Predicting gene expression from sequence is a long-standing goal of computational biology that
is still far from completion. While existing sequence-based models of cis-regulation are trained
across the whole range of expression levels[33,49,50], we have proposed here to focus on extreme
expression variations. Extremes may not well be captured by models trained to globally predict
gene expression, as evident from the limited contribution of Enformer[33] to our model. Also, the
biological mechanisms underpinning extreme expression variations may differ from those
governing moderate expression variations. However, the relevance of extreme expression for
clinical diagnostics and research is high. We hope that the benchmark and algorithms we have
developed will foster further research in this direction and aid in the development and validation
of methods predicting the impact of large-effect variants on the human transcriptome.

567 **Methods**

568 Underexpression outlier benchmark dataset

569 GTEx dataset

We downloaded the GTEx RNA-seq read alignment files in the BAM format from dbGaP
(phs000424.v8.p2). We excluded tissues with less than 60 RNA-seq samples due to insufficient
statistical power[11]. This filter discarded bladder, endocervix, ectocervix, fallopian tube, and

kidney medulla, leaving 49 tissues.

We obtained SNPs and small indels from the GTEx hg19 variant calls from the file

GTEx_Analysis_2016-01-

15_v7_WholeGenomeSeq_635Ind_PASS_AB02_GQ20_HETX_MISS15_PLINKQC.vcf.gz from

the dbGap entry phg000830.p1. Structural variants were obtained from Ferraro and

colleagues[15].

Expression outliers

Gene expression outlier analysis was performed following the aberrant expression module of

DROP v1.1.0[51] based on OUTRIDER[11]. To this end, we used as reference genome the GRCh38

primary assembly release 34 of the GENCODE project[52]. A fragment (reads pair) was assigned

to a gene if and only if both reads were entirely aligned within the gene, allowing for fragments

to be assigned to more than one gene. On each tissue separately, genes with an FPKM less

than 1 in 95% or more of the samples were considered to be not sufficiently expressed in the

tissue and filtered out, as previously described[11]. Since the variants were aligned to the hg19

reference genome, whereas the RNA-seq data was aligned to hg38, we lifted the gene

expression counts by their Ensembl gene IDs over to hg19, relying on Gencode[52] gene

mappings. Overall, the liftover had a minimal impact with only 76 out of 19,682 genes and 294

out of 87,748 transcripts that could not be mapped.

OUTRIDER is an expression outlier caller that uses an autoencoder to model RNA-seq

fragment count expectations with a negative binomial distribution. Specifically, OUTRIDER

models the probability of the observed fragment count $x_{s,g}$ for every gene g in a sample s as:

$$P(x_{s,g}|\mu_{s,g},\theta_{t(s),g}) = \text{NB}(x_{s,g}|\mu_{s,g},\theta_{t(s),g}) \ (1)$$

594 where:

595　　-　$\mu_{s,g}$ is the expected fragment count

596　　-　$\theta_{t(s),g}$ is the dispersion parameter for the gene g in the tissue of sample s $t(s)$

597 OUTRIDER further outputs:

598　　-　the biological coefficient of variation:

$$\text{BCV}_{t(s),g} = \frac{1}{\sqrt{\theta_{t(s),g}}}$$

599　　-　the log$_2$-transformed fold-change of the observed fragment count compared to the
600　　　expected fragment count:

$$\log_2 \mathrm{FC} = \log_2(x) - \log_2(\mu)$$

-     the nominal *p*-value

-     the False Discovery Rate using the Benjamini-Yekutieli method[53]

The resulting table was subsetted to individuals with an available whole genome sequencing

and to protein-coding genes. Some of the data points could not be detected as outliers due to

lack of statistical power. To reduce the proportion of these insufficiently powered data points in

our benchmark, we discarded observations with an expected fragment count $\mu_{s,g}$ less than 450,

a minimal value that was empirically estimated to allow recovering half of the two-fold reduction

outliers transcriptome-wide upon a FDR cutoff of 5%[6]. We labeled as gene expression outliers

all observations with an FDR less than 5%. Lastly, RNA-seq samples that contained more than

20 outliers were discarded because samples with numerous outliers may be samples for which

OUTRIDER could not adequately fit the data or for which gene expression is globally affected,

resulting in widespread expression aberrations throughout the genome that cannot be predicted

from local sequence variation.

614 OUTRIDER *z*-score computation

615 We quantile-mapped the OUTRIDER-fitted negative binomial distributions (Eq. 1) to the

616 standard normal distribution as follows:

617 $$z\text{-score} = \mathrm{CDF}^{-1}_{N(0,1)}(\mathrm{CDF}_{NB}(x|\mu,\theta))$$,

618 where $\mathrm{CDF}^{-1}_{N(0,1)}$ is the inverse cumulative distribution function of the standard normal

619 distribution and $\mathrm{CDF}_{NB}(x|\mu,\theta)$ the negative-binomial cumulative distribution function.

620 Precision-recall

621 We evaluated models using precision-recall curves due to the small proportion of outliers and

622 summarized them with the area under the precision-recall curve (AUPRC) as implemented in

623 'average_precision_score' function of the scikit-learn package v1.3.2:

$$P = \frac{T_P}{T_P + F_P}$$
$$R = \frac{T_P}{T_P + F_N}$$
$$AP = \sum_n (R_n - R_{n-1})P_n$$

, where P_n and R_n are the precision and recall at the n^{th} top prediction. The AUPRC is the

average precision for each cutoff weighted by the recall difference.

The average precision was computed on all held-out data together on the one hand, and on

held-out data for each of the 27 GTEx tissue types on the other hand. The GTEx tissue types

group together highly similar tissues, notably many regions of the brain. By calculating

performance within tissue types instead of reporting highly similar tissues separately, we avoid

biasing our evaluation for overrepresented tissue types.

631 Variant filtering, annotation, and gene-level aggregation

632 Filtering for rare high-quality variants

633 We considered a variant to be rare if it had a minor allele frequency in the general population

634 ≤0.001 based on the Genome Aggregation Database (gnomAD v2.1.1) and was found in at

635 most 2 individuals within GTEx. Variants had to be supported by at least 10 reads and had to

636 pass the conservative genotype-quality filter of GQ > 30. For structural variants, we only filtered

637 for the number of occurrences in the GTEx dataset (less than in 2 individuals).

638 Variant annotation with VEP and AbSplice

639 For all rare variants, we used Ensembl VEP[54] v108 to calculate consequences, LOFTEE loss-of-

640 function annotation, as well as CADD v1.6 scores. Tissue-specific aberrant splicing predictions

641 were generated with AbSplice[32].

642 Definition of gene-level features

For each combination of gene, individual, and tissue, we required a set of features to predict the

underexpression outlier label. Therefore, we constructed a set of features starting from the

annotations of the underlying rare variants. CADD predictions were max-aggregated per gene

across the rare variants. For AbSplice, we kept the maximum absolute score in the

corresponding gene and tissue.

Isoform-specific variant annotations, namely LOFTEE and VEP consequences, were

aggregated differently depending on whether isoform proportions should be taken into account.

When disregarding isoform proportions (i.e. the model "LOFTEE+CADD+consequences"), we

only incorporated variants affecting the canonical transcript (canonical according to VEP). The

gene was assumed to have an annotation (LOFTEE classification and each VEP consequence,

e.g. stop-gained) when any of the variants in the gene had this annotation. Since the VEP

canonical transcript does not differ between tissues, all tissues end up with the same set of
gene-level features for these annotations.

For all models incorporating isoform proportions, isoform-specific variant annotations were first
weighted by the total proportion of isoforms *i* in tissue *t* that they affect:

$$w_v = \sum_{i}^{\text{isoforms}(g)} \text{proportion}(i, t) \cdot \delta_{v \text{ affects } i}$$

658 , where $\delta_{v \text{ affects } i}$ is 1 if the variant affects the isoform, otherwise 0, the proportion of an isoform
659 *i* in a tissue *t* was estimated as the median TPM proportion across individuals among all isoform
660 of the same gene *g*:

$$\text{proportion}(i, t) = \text{median}_{s \in \text{individuals}} \left(\frac{tpm(i, t, s)}{\sum_{j \in \text{isoforms}(g)} tpm(j, t, s)} \right) \quad (2)$$

661 , and where tpm(*i, t, s*) is the transcript-level TPM obtained from GTEx v8 (dbGaP Accession
662 phs000424.v8.p2). All resulting variant annotations were then max-aggregated per gene and
663 tissue across variants.

664 Enformer variant effect prediction

We used Enformer[33] to evaluate the ability of deep-learning models to predict rare variant
effects in promoters. Given a transcript, we first extracted three 393,216 nt DNA sequences,
one centered at the transcription start site (TSS) and two others shifted by 43 nt upstream and
downstream of the TSS. Subsequently, Enformer was applied to each sequence, and we
extracted predictions for various subsets out of the 5,313 human assay tracks on the three
central bins from each sequence. We denote *p* as the number of selected tracks. To ensure that
regulatory elements falling on bin boundaries were accounted for, we averaged the predictions
over the sequence shifts *k* and central bins *l*:

$$\mathbf{u}(\text{seq}) = \text{mean}_{\substack{k \in \text{sequence-shifts,} \\ l \in \text{central-bins}}} (\mathcal{E}(\mathbf{X}(\text{seq}, k, l)))$$

Here, the *p*-dimensional vector $\mathbf{u}(\text{seq})$ contains the aggregated Enformer scores over the
central bins and shifts for all the *p* tracks, the matrix $\mathbf{X}(\text{seq}, k, l)$ is the one-hot encoding of the
shifted sequence *k* centered on bin *l*, and $\mathcal{E}$ is the Enformer model. Since Enformer does not
directly predict RNA-seq coverage for GTEx tissues, we trained different linear regression
models (e.g. ElasticNet, Ridge) taking the aggregated Enformer scores $\mathbf{u}(\text{seq})$ defined above
as input to predict the expression level for each GTEx tissue averaged across all GTEx

individuals, as previously[21].

Specifically, we introduced $y_{i,t} := \log_{10}(1 + \text{mean}_{s \in \text{individual}}(\text{tpm}(i,t,s)))$, where t is the

tissue, i is the transcript, $\text{tpm}(i,t,s)$ is the transcript-level TPM from GTEx v8 of individual s,

and aimed at predicting $y_{i,t}$ linearly from the aggregate scores. Hence, we modeled

$y_{i,t} = \boldsymbol{\beta}_t^\top \log_{10}(1 + \mathbf{u}(\text{seq}_i)) + \epsilon_{i,t}$, and estimated the coefficients $\boldsymbol{\beta}_t$ with different

regularized linear regression schemes (e.g. ElasticNet, Ridge).

Having fitted this model, we now have a function $\hat{y}(\text{seq},t) := \hat{\boldsymbol{\beta}}_t^T \log_{10}(1 + \mathbf{u}(\text{seq}))$ that

predicts gene expression from sequence for each tissue. We employed two different methods to

obtain a variant effect score for a given gene. First, we estimated the effect of the variant on the

canonical transcript of the gene, if the variant was in close proximity to its promoter, as follows:

$$\text{score\_canonical}(g,v,t) = \frac{\hat{y}(\text{alt},t) - \hat{y}(\text{ref},t)}{\log_{10}(2)}$$

689 Here, v is the variant of interest, t is the tissue, g is the gene of interest, and alt and ref are the

690 alternative and reference sequences of the canonical transcript respectively. Second, we

691 weighted all isoforms of the gene by their isoform proportions, using the following score:

$$\text{score\_weighted}(g,v,t) = \log_2 \frac{\sum_i^{\text{isoforms}(g)} \text{proportion}(i,t) \cdot 10^{\hat{y}(\text{alt}_i,t)}}{\sum_i^{\text{isoforms}(g)} \text{proportion}(i,t) \cdot 10^{\hat{y}(\text{ref}_i,t)}}$$

Here, v is the variant of interest, t is the tissue, g is the gene of interest, alt_i and ref_i are the

alternative and reference sequences of transcript i respectively, and $\text{isoforms}(g)$ returns the

isoforms of g that are located in close proximity to the variant.

For each GTEx individual, tissue, and gene combination, we took the signed maximum absolute

variant effect across all relevant variants as the variant effect score. Finally, we performed

cross-validation to optimize (1) the subset of Enformer tracks, (2) the linear model parameters

(e.g. L1 and L2 regularization), (3) the variant-effect function (i.e. weighted or canonical

isoforms), and (4) the shortest interval around the TSS, for which Enformer is predictive (suppl.

Fig. S9).

Variant type enrichment in underexpression outliers

To obtain the proportion of underexpression outliers explained by different classes of variants,

we first identified the variant consequences each gene-individual-tissue combination is affected

by. We then assigned each gene-individual-tissue combination to be caused by the most

impactful consequence among the following consequences: Downstream variant, Inframe CDS

(inframe deletion, inframe insertion, missense variant), Intron, NMD-like (NMD escaping,
Frameshift, Start lost, Stop gained, Stop lost), Promoter (Upstream gene variant), Splicing
disruption (Splice acceptor/donor/region variant), Structural ablation (Transcript ablation),
Synonymous, UTR (3' UTR, 5' UTR). The order of assignment was determined by performing a
Fisher's exact test for each variant consequence against the underexpression outlier class and
ordering the classes by decreasing significance. Afterward, the proportion of underexpression
outliers explainable by each variant type was calculated (suppl. Fig. S3).

713 Model training

All 633 individuals were split into six cross-validation groups with approximately equal numbers
of underexpression outliers and tissues.

The DNA-based models were trained to predict the OUTRIDER z-scores. To this end, we used
gradient-boosted trees[55] from the LightGBM[56] framework with default parameters:
boosting_type: gbdt, learning_rate: 0.1, max_depth: -1, min_child_samples: 20,
min_child_weight: 0.001, min_split_gain: 0, n_estimators: 100, num_leaves: 31, reg_alpha: 0,
reg_lambda: 0, subsample: 1, subsample_for_bin: 200000, subsample_freq: 0.

For training the model integrating CAT RNA-seq data, the same cross-validation scheme was
used as for the DNA-based models while excluding the CAT and highly related tissues thereof
from the predicted tissues. When using fibroblasts as CAT, we excluded the non sun-exposed
suprapubic skin, sun-exposed lower leg skin, and cultured fibroblasts. When using whole blood
as CAT, we excluded whole blood and EBV-transformed lymphocytes from the predicted
tissues. Unlike the DNA-based models, this model was trained to predict underexpressed
outliers using a logistic regression taking as input i) a binary variable indicating whether the
gene is expressed in the CAT, ii) the OUTRIDER z-score of the gene in the CAT, iii) the AbExp
prediction for the target tissue, and all 3 interaction terms between those 3 variables.

To compare Watershed with other machine learning methods, we trained Watershed models
using the following features: CADD, LOFTEE (high-confidence), LOFTEE (low-confidence), 5'
UTR variant, 3' UTR variant, downstream gene variant, intron variant, missense variant, non-
coding transcript exon variant, splice acceptor variant, splice donor variant, splice region variant,
stop gained, synonymous variant, upstream gene variant, sift score, polyphen score. As the
dataset was too large to be fitted in total with Watershed, we subsampled the non-outliers to ten
times the number of outliers. This change in class balance affects the posterior by constant but
not their ranking. Afterwards, we extracted the conditional probabilities P(Z|G) instead of the
standard posterior of P(Z|G,E) to predict the likelihood of an expression outlier on the validation

29   29

739 folds. Z is the latent variable, G is the vector of genomic annotations, and E is the outlier status.

Replication in independent datasets

In the two replication datasets, variants were filtered for genotype quality ≥ 30 and read depth ≥
10 reads. Moreover, rare variants were subsetted based on the gnomAD population with minor
allele frequency ≤ 0.001. Those high-quality and rare variants were used as candidates for
outlier prediction. Gene expression outliers were obtained with OUTRIDER and filtered for a
sufficiently large expected number of fragments ($\mu > 450$). Samples with more than 20 outliers
were removed.

The mitochondrial disease dataset[6] consisted of 311 whole-exome sequencing samples paired
with RNA-seq from fibroblasts. After filtering, this dataset contained 808 underexpression
outliers across 295 samples. A detailed overview of how many samples, genes, etc. remained
after each filtering step can be seen in suppl. table T4.

For the amyotrophic lateral sclerosis (ALS) dataset, we downloaded 253 transcriptomes with
matched whole-genome sequencing data from https://dataportal.answerals.org[35]. RNA-seq
measurements were obtained from iPSC-derived spinal motor neurons. After filtering, the
dataset contained 653 underexpression outliers across 233 samples (195 cases and 38
controls). The corresponding overview of how many samples, genes, etc. remained after each
filtering step can be seen in suppl. table T5.

Prediction of ClinVar pathogenic variants

To assess the efficacy of AbExp in distinguishing pathogenic from benign variants, we
annotated known pathogenic, likely benign, and benign variants from the ClinVar database with
both AbExp and CADD. Importantly, we used the official CADD pipeline to annotate all variants,
including those with missing pre-computed scores. Variants classified as both likely benign and
benign were retained solely as likely benign within the dataset. Furthermore, for each variant,
we retained only the minimum predicted AbExp score across all tissues.

GnomAD variant enrichments

For all possible SNVs within 5 kb of 19,018 protein-coding genes, we computed AbExp scores
with and without using the BCV, CADD scores, and LOFTEE high-impact loss of function
annotations. High-impact predictions of CADD and AbExp without BCV were identified by
matching score cutoffs to the quantile of the high-impact cutoff of AbExp. Using logistic

regression fits, we then computed the odds-ratio of high-impact variants among absent,
singleton, rare (MAF < 0.1%), and common SNVs in gnomAD for these models. Moreover, we
calculated odds-ratios for these variant types and methods as a function of gene LOEUF decile.
The loss-of-function observed/expected upper bound fraction (LOEUF) scores were
downloaded from https://gnomad.broadinstitute.org/downloads.

774 UK Biobank rare variant association testing and phenotype
775 prediction

776 We analyzed data from 200,593 European ancestry unrelated individuals in the UK Biobank
777 (using field 22011 and labeled as Caucasian in field 22006), all of whom had genotypes
778 available from exome-sequencing and microarrays as well as blood and urine measurements. A
779 detailed list of used phenotypes can be found in suppl. table T2. For every trait, trait values were
780 inverse rank normal transformed, as in the Genebass study[38].

781 Identification of lead trait-associated variants

782 To control for common variants in the vicinity of each gene, trait-associated variants were
783 obtained from the PanUKBB study[57]. Using plink v1.9[58,59], variants with a p-value ≤ 0.0001 were
784 clumped in a 250-kb window with an LD-cutoff of $r^2 < 0.5$ to identify independent lead variants
785 for every trait. The imputed genotypes were then subsetted for these lead variants in a 250-kb
786 window around each gene.

787 Application of polygenic risk scores

788 Polygenic risk scores were selected from the study by Privé et al.[60] if available and otherwise
789 from the study by Tanigawa et al.[61] (suppl. Table T2). Score files were obtained from the PGS
790 catalog database[62] and applied to the imputed genotypes using plink v2.0[58,63].

791 Calculation of AbExp and LOFTEE scores

792 The UKBB whole-exome sequencing data was subsetted for variants with a minor allele
793 frequency ≤0.001 based on gnomAD v3.1.1 and filtered for genotype quality ≥ 30 and read
794 depth ≥ 10 reads. The remaining variants were then annotated using Ensembl VEP[54] v108 with
795 the LOFTEE plugin[22] and AbExp.

796 Rare variant association testing

797 Gene association was tested using a likelihood ratio test between a restricted linear regression

798 model containing only covariates and four linear regression models with additional predictor
799 variables as described in Table 1.

Model	Predictor variables	Nr. of variables
LOFTEE	Number of LOFTEE variants in the gene of interest	1
AbExp all tissues	The set of minimal AbExp scores across all rare variants in the gene (0 in absence of any rare variant) for each tissue	49
Minimum AbExp	The minimum AbExp score across all rare variants in the gene and across all tissues, 0 in absence of any rare variant	1
Median AbExp	The median AbExp score across all rare variants in the gene and across all tissues, 0 in absence of any rare variant	1

Table 1: **Rare variant association test models.**

The following covariates were used for all models: sex, age, age², age times sex, age² times
sex, the 20 first genetic principal components (field 22009), lead associated variants in 250-kb
window around the gene of interest, and a polygenic risk score predicting the trait. The
polygenic risk scores and lead variants were based on the whole UK Biobank dataset. This mild
data leakage may have led to model overfitting. However, since these features were used as
covariates in both restricted and full models, the comparison between models remained valid. P-
value calibration of the models was assessed by permuting the phenotype once. Identification of
significantly trait-associated genes was performed on two thirds of the dataset.

Phenotype prediction

809
For the common-variant-based phenotype prediction model, the following features were used:
sex, age, age², age times sex, age² times sex, the 20 first genetic principal components (field
22009), and a polygenic risk score predicting the trait. In contrast to the rare variant association
testing, we did not include lead variants of the associating genes, as these would lead to a too
large number of predictor variables.
We compared two phenotype prediction models integrating rare and common variants. Both are

nonlinear models. To this end, we used gradient-boosted trees[55] from the LightGBM[56]

framework with default parameters: boosting_type: gbdt, learning_rate: 0.1, max_depth: -1,

min_child_samples: 20, min_child_weight: 0.001, min_split_gain: 0, n_estimators: 100,

num_leaves: 31, reg_alpha: 0, reg_lambda: 0, subsample: 1, subsample_for_bin: 200000,

subsample_freq: 0. Both models include the same features as the common-variant based

prediction models. The AbExp phenotype prediction model further included the AbExp scores of

significantly trait-associated genes in 49 tissues, while the LOFTEE phenotype prediction model

included the number of LOFTEE pLoF variants of significantly trait-associated genes. For each

trait, the two models can have a different set of significantly trait-associated genes as identified

by their corresponding RVATs. The phenotype prediction models were trained with 5-fold cross-

validation on the remaining third of the dataset. All evaluations were performed on the hold-out

folds.

Declarations

Ethics approval and consent to participate

No new data was generated for this study. The different ethics approvals can be found in the corresponding publications (GTEx[64], AnswerALS[35], mitochondrial disease[6]). The UK Biobank was approved by the North West Multi-centre Research Ethics Committee (21/NW/0157). Our reference number approved by the UK Biobank is 25214. All UK Biobank study participants gave written informed consent.

The research conformed to the principles of the Declaration of Helsinki.

Consent for publication

All individuals included or their legal guardians provided written consent to share pseudonymized patient data and analysis data, as described in the original publications.

Competing interests

The authors declare that they have no competing interests.

Acknowledgements

We thank Felix Brechtmann, Alexander Karollus, and Holger Prokisch for fruitful discussions and their valuable feedback on the manuscript. Figure 1 was created with BioRender.com. We thank Žiga Avsec and Alexander Karollus for advice on using the Enformer in this context.

Authors' contributions

J.G. conceptualized the project. F.R.H., F.P.C. and J.G. designed the methodology. F.R.H., N.W. and V.A.Y. curated the data. F.R.H., J.L., and G.T. performed investigation and developed the software. F.R.H., J.L., G.T., and V.A.Y. performed visualizations. F.R.H., V.A.Y. and J.G. wrote the original draft of the manuscript. All authors reviewed and edited the manuscript. J.G. supervised the project with the help of F.P.C. and V.A.Y.

Funding

The German Bundesministerium für Bildung und Forschung (BMBF) supported the study through the Model Exchange for Regulatory Genomics project (MERGE; grant no. 031L0174A

to F.R.H. and J.G.); the VALE (Entdeckung und Vorhersage der Wirkung von genetischen

Varianten durch Artifizielle Intelligenz für Leukämie Diagnose und Subtypidentifizierung) project

(031L0203B to F.R.H., J.G); and the ERA PerMed project PerMiM (01KU2016B to J.L., V.A.Y.

and J.G.). N.W. is supported by the Helmholtz Association under the joint research school

'Munich School for Data Science – MUDS'. F.P.C. was funded by the Free State of Bavaria's

Hightech Agenda through the Institute of AI for Health (AIH).

The Genotype-Tissue Expression (GTEx) project was supported by the Common Fund of the

Office of the Director of the National Institutes of Health, and by the NCI, NHGRI, NHLBI, NIDA,

NIMH and NINDS. This study was supported by data provided by the Answer ALS Consortium,

administered by the Robert Packard Center for ALS at Johns Hopkins. The funders had no role

in study design, data collection and analysis, decision to publish or preparation of the

manuscript. UK Biobank was established by the Wellcome Trust medical charity, Medical

Research Council, Department of Health, Scottish Government and the Northwest Regional

Development Agency. It has also had funding from the Welsh Government, British Heart

Foundation, Cancer Research UK and Diabetes UK. UK Biobank is supported by the National

Health Service (NHS).

870 Availability of data and materials

871 Data availability

No primary data were generated for this study. Rare variants from gnomAD v.2.1.1 are publicly

available at https://gnomad.broadinstitute.org. The GTEx v8 dataset is available at (under

dbGaP protection) https://gtexportal.org/home. The ALS dataset is available at

http://dataportal.answerals.org after a registration and approval process. The mitochondrial

dataset is described by Yépez et al.[6] The UK Biobank dataset is available at

https://biobank.ndph.ox.ac.uk/ukb/ after a registration and approval process. We provide the

aberrant expression benchmark dataset, isoform proportions, and the expected gene

expression in GTEx v8 as open-access in the Zenodo repository[65] (doi:

10.5281/zenodo.13895620).

881 Code availability

A Snakemake pipeline to calculate AbExp predictions can be found at

https://github.com/gagneurlab/abexp.

The source code of Watershed adapted for underexpression prediction can be found at

https://github.com/Hoeze/Watershed.

The source code for the UK Biobank rare-variant association study and phenotype prediction

can be found in the following repositories:

- Main analysis pipeline: https://github.com/gagneurlab/abexp-ukbb-trait-analysis

- Variant clumping: https://github.com/gagneurlab/abexp-ukbb-variant-clumping

- Polygenic risk score calculation: https://github.com/gagneurlab/abexp-ukbb-prs

References

1. Wang, L.-H., Wu, C.-F., Rajasekaran, N. & Shin, Y. K. Loss of Tumor Suppressor Gene Function in Human Cancer: An Overview. *Cell. Physiol. Biochem.* **51**, 2647–2693 (2018).

2. Kontomanolis, E. N. *et al.* Role of Oncogenes and Tumor-suppressor Genes in Carcinogenesis: A Review. *Anticancer Res.* **40**, 6009–6015 (2020).

3. Gonorazky, H. D. *et al.* Expanding the Boundaries of RNA Sequencing as a Diagnostic Tool for Rare Mendelian Disease. *Am. J. Hum. Genet.* **104**, 466–483 (2019).

4. Frésard, L. *et al.* Identification of rare-disease genes using blood transcriptome sequencing and large control cohorts. *Nat. Med.* **25**, 911–919 (2019).

5. Murdock, D. R. Enhancing Diagnosis Through RNA Sequencing. *Clin. Lab. Med.* **40**, 113–119 (2020).

6. Yépez, V. A. *et al.* Clinical implementation of RNA sequencing for Mendelian disease diagnostics. *Genome Med.* **14**, 38 (2022).

7. Dekker, J. *et al.* Web-accessible application for identifying pathogenic transcripts with RNA-seq: Increased sensitivity in diagnosis of neurodevelopmental disorders. *Am. J. Hum. Genet.* **110**, 251–272 (2023).

8. Deshwar, A. R. *et al.* Trio RNA sequencing in a cohort of medically complex children. *Am. J. Hum. Genet.* **110**, 895–900 (2023).

9. Smail, C. *et al.* Integration of rare expression outlier-associated variants improves polygenic risk prediction. *Am. J. Hum. Genet.* **109**, 1055–1064 (2022).

10. Li, B. *et al.* Evaluation of PrediXcan for prioritizing GWAS associations and predicting gene expression. in *Biocomputing 2018* 448–459 (WORLD SCIENTIFIC, 2017). doi:10.1142/9789813235533_0041.

11. Brechtmann, F. *et al.* OUTRIDER: A Statistical Method for Detecting Aberrantly Expressed Genes in RNA Sequencing Data. *Am. J. Hum. Genet.* (2018)

doi:10.1016/j.ajhg.2018.10.025.

12. Salkovic, E., Sadeghi, M. A., Baggag, A., Salem, A. G. R. & Bensmail, H. OutSingle: a novel

method of detecting and injecting outliers in RNA-Seq count data using the optimal hard

threshold for singular values. *Bioinformatics* **39**, btad142 (2023).

13. Segers, A., Gilis, J., Heetvelde, M. V., Baere, E. D. & Clement, L. Juggling offsets unlocks

RNA-seq tools for fast scalable differential usage, aberrant splicing and expression

analyses. 2023.06.29.547014 Preprint at https://doi.org/10.1101/2023.06.29.547014 (2023).

14. Li, X. *et al.* The impact of rare variation on gene expression across tissues. *Nature* **550**,

239–243 (2017).

15. Ferraro, N. M. *et al.* Transcriptomic signatures across human tissues identify functional rare

genetic variation. *Science* **369**, (2020).

16. Hombach, S. & Kretz, M. Non-coding RNAs: Classification, Biology and Functioning. *Adv.*

Exp. Med. Biol. **937**, 3–17 (2016).

17. Zhang, Z., Qin, Y.-W., Brewer, G. & Jing, Q. MicroRNA degradation and turnover: regulating

the regulators. *Wiley Interdiscip. Rev. RNA* **3**, 593–600 (2012).

18. Megel, C. *et al.* Surveillance and Cleavage of Eukaryotic tRNAs. *Int. J. Mol. Sci.* **16**, 1873–

1893 (2015).

19. Nair, L., Chung, H. & Basu, U. Regulation of long non-coding RNAs and genome dynamics

by the RNA surveillance machinery. *Nat. Rev. Mol. Cell Biol.* **21**, 123–136 (2020).

20. GTEx Consortium. The GTEx Consortium atlas of genetic regulatory effects across human

tissues. *Science* **369**, 1318–1330 (2020).

21. Karollus, A., Mauermeier, T. & Gagneur, J. Current sequence-based models capture gene

expression determinants in promoters but mostly ignore distal enhancers. *Genome Biol.* **24**,

56 (2023).

22. Karczewski, K. J. *et al.* The mutational constraint spectrum quantified from variation in

141,456 humans. *Nature* **581**, 434–443 (2020).

23. Rentzsch, P., Witten, D., Cooper, G. M., Shendure, J. & Kircher, M. CADD: predicting the

deleteriousness of variants throughout the human genome. *Nucleic Acids Res.* **47**, D886–

D894 (2019).

24. Cummings, B. B. *et al.* Transcript expression-aware annotation improves rare variant

interpretation. *Nature* **581**, 452–458 (2020).

25. Morales, J. *et al.* A joint NCBI and EMBL-EBI transcript set for clinical genomics and

research. *Nature* **604**, 310–315 (2022).

26. Dugan, S. L. *et al.* New recessive truncating mutation in LTBP3 in a family with oligodontia,

short stature, and mitral valve prolapse. *Am. J. Med. Genet. A.* **167**, 1396–1399 (2015).

27. Sharon, D., Kimchi, A. & Rivolta, C. OR2W3 sequence variants are unlikely to cause

inherited retinal diseases. *Ophthalmic Genet.* **37**, 366–368 (2016).

28. Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for

differential expression analysis of digital gene expression data. *Bioinformatics* **26**, 139–140

(2010).

29. Bader, D. M. *et al.* Negative feedback buffers effects of regulatory variants. *Mol. Syst. Biol.*

11, 785 (2015).

30. Einarsson, H. *et al.* Promoter sequence and architecture determine expression variability

and confer robustness to genetic variants. *eLife* **11**, e80943 (2022).

31. Teran, N. A. *et al.* Nonsense-mediated decay is highly stable across individuals and tissues.

bioRxiv 2021.02.03.429654 (2021) doi:10.1101/2021.02.03.429654.

32. Wagner, N. *et al.* Aberrant splicing prediction across human tissues. *Nat. Genet.* **55**, 861–

870 (2023).

33. Avsec, Ž. *et al.* Effective gene expression prediction from sequence by integrating long-

range interactions. *Nat. Methods* **18**, 1196–1203 (2021).

34. Haberle, V. & Stark, A. Eukaryotic core promoters and the functional basis of transcription

initiation. *Nat. Rev. Mol. Cell Biol.* **19**, 621–637 (2018).

35. Baxi, E. G. *et al.* Answer ALS, a large-scale resource for sporadic and familial ALS

combining clinical and multi-omics data from induced pluripotent cell lines. *Nat. Neurosci.*

25, 226–237 (2022).

36. Fair, B. J. *et al.* Gene expression variability in human and chimpanzee populations share

common determinants. *eLife* **9**, e59929 (2020).

37. Wang, Q. *et al.* Rare variant contribution to human disease in 281,104 UK Biobank exomes.

Nature **597**, 527–532 (2021).

38. Karczewski, K. J. *et al.* Systematic single-variant and gene-based association testing of

thousands of phenotypes in 394,841 UK Biobank exomes. *Cell Genomics* **2**, 100168 (2022).

39. Fiziev, P. P. *et al.* Rare penetrant mutations confer severe risk of common diseases.

Science **380**, eabo1131 (2023).

40. Wu, M. C. *et al.* Rare-Variant Association Testing for Sequencing Data with the Sequence

Kernel Association Test. *Am. J. Hum. Genet.* **89**, 82–93 (2011).

41. Lee, S., Abecasis, G. R., Boehnke, M. & Lin, X. Rare-Variant Association Analysis: Study

Designs and Statistical Tests. *Am. J. Hum. Genet.* **95**, 5–23 (2014).

42. Sudlow, C. *et al.* UK Biobank: An Open Access Resource for Identifying the Causes of a

Wide Range of Complex Diseases of Middle and Old Age. *PLOS Med.* **12**, e1001779

(2015).

43. Zhou, W. *et al.* SAIGE-GENE+ improves the efficiency and accuracy of set-based rare

variant association tests. *Nat. Genet.* **54**, 1466–1469 (2022).

44. Clarke, B. *et al.* Integration of variant annotations using deep set networks boosts rare

variant association genetics. 2023.07.12.548506 Preprint at

https://doi.org/10.1101/2023.07.12.548506 (2023).

45. Cummings, B. B. *et al.* Improving genetic diagnosis in Mendelian disease with transcriptome

sequencing. *Sci. Transl. Med.* (2017) doi:10.1126/scitranslmed.aal5209.

46. Kremer, L. S. *et al.* Genetic diagnosis of Mendelian disorders via RNA sequencing. *Nat.*

Commun. **8**, (2017).

47. Lunke, S. *et al.* Integrated multi-omics for rapid rare disease diagnosis on a national scale.

Nat. Med. **29**, 1681–1691 (2023).

48. Aicher, J. K., Jewell, P., Vaquero-Garcia, J., Barash, Y. & Bhoj, E. J. Mapping RNA splicing

variations in clinically-accessible and non-accessible tissues to facilitate Mendelian disease

diagnosis using RNA-seq. *Genet. Med. Off. J. Am. Coll. Med. Genet.* **22**, 1181–1190 (2020).

49. Zhou, J. *et al.* Deep learning sequence-based ab initio prediction of variant effects on

expression and disease risk. *Nat. Genet.* **50**, 1171–1179 (2018).

50. Linder, J., Srivastava, D., Yuan, H., Agarwal, V. & Kelley, D. R. Predicting RNA-seq

coverage from DNA sequence as a unifying model of gene regulation. 2023.08.30.555582

Preprint at https://doi.org/10.1101/2023.08.30.555582 (2023).

51. Yépez, V. A. *et al.* Detection of aberrant gene expression events in RNA sequencing data.

Nat. Protoc. **16**, 1276–1296 (2021).

52. Frankish, A. *et al.* GENCODE reference annotation for the human and mouse genomes.

Nucleic Acids Res. **47**, D766–D773 (2019).

53. Benjamini, Y. & Yekutieli, D. The control of the false discovery rate in multiple testing under

dependency. *Ann. Stat.* **29**, 1165–1188 (2001).

54. McLaren, W. *et al.* The Ensembl Variant Effect Predictor. *Genome Biol.* **17**, 122 (2016).

55. Hastie, T., Tibshirani, R. & Friedman, J. Boosting and Additive Trees. in *The Elements of*

Statistical Learning: Data Mining, Inference, and Prediction (eds. Hastie, T., Tibshirani, R. &

Friedman, J.) 337–387 (Springer, New York, NY, 2009). doi:10.1007/978-0-387-84858-

7_10.

56. Ke, G. *et al.* LightGBM: A Highly Efficient Gradient Boosting Decision Tree. in *Advances in*

Neural Information Processing Systems vol. 30 (Curran Associates, Inc., 2017).

57. Pan-UKB team. Pan-ancestry genetic analysis of the UK Biobank. (2020).

58. Chang, C. C. *et al.* Second-generation PLINK: rising to the challenge of larger and richer

datasets. *GigaScience* **4**, s13742-015-0047–8 (2015).

59. Shaun Purcell & Christopher Chang. PLINK 1.9. (2020).

60. Privé, F. *et al.* Portability of 245 polygenic scores when derived from the UK Biobank and

applied to 9 ancestry groups from the same cohort. *Am. J. Hum. Genet.* **109**, 12–23 (2022).

61. Tanigawa, Y. *et al.* Significant sparse polygenic risk scores across 813 traits in UK Biobank.

PLOS Genet. **18**, e1010105 (2022).

62. Lambert, S. A. *et al.* The Polygenic Score Catalog as an open database for reproducibility

and systematic evaluation. *Nat. Genet.* **53**, 420–425 (2021).

63. Shaun Purcell & Christopher Chang. PLINK 2.0. (2022).

64. GTEx Consortium. The Genotype-Tissue Expression (GTEx) project. *Nat. Genet.* **45**, 580–

585 (2013).

65. Hölzlwimmer, F. R. Aberrant gene expression prediction benchmark based on GTEx v8.

Zenodo https://doi.org/10.5281/zenodo.13895620 (2024).