Effect of composting and storage on the microbiome and resistome of cattle manure from a commercial dairy farm in Poland

Magdalena Zalewska[1*], Aleksandra Błażejewska[1], Mateusz Szadziul[1], Karol Ciuchciński[2], Magdalena Popowska[1*]

[1]Department of Bacterial Physiology, Institute of Microbiology, Faculty of Biology, University of Warsaw

[2]2Department of Environmental Microbiology and Biotechnology, Institute of Microbiology, Faculty of Biology, University of Warsaw

* Correspondence:
ma.popowska@uw.edu.pl, mm.zalewska10@uw.edu.pl

Keywords: antibiotic-resistant bacteria, antibiotic resistance genes, composting, dairy cattle manure, manure treatment strategies, storage

Abstract

Manure from food-producing animals, rich in antibiotic-resistant bacteria and antibiotic resistance genes (ARG), poses significant environmental and healthcare risks. Despite global efforts, most manure is not adequately processed before use on fields, escalating the spread of antimicrobial resistance. This study examined how different cattle manure treatments, including composting and storage, affect its microbiome and resistome. The changes occurring in the microbiome and resistome of the treated manure samples were compared with those of raw samples by high-throughput qPCR for ARGs tracking and metagenomic sequencing of the V3-V4 variable region of 16S rRNA gene to indicate microbial community composition. We identified 203 ARGs and mobile genetic elements (MGEs) in raw manure. Post-treatment reduced these to 76 in composted and 51 in stored samples. Notably, beta-lactam, MLSB, and vancomycin-resistance genes decreased, while genes linked to MGEs, integrons, and sulfonamide resistance increased after composting. Overall, total resistance gene abundance significantly dropped with both treatments. During composting, the relative abundance of genes was lower midway than at the end. Moreover, higher biodiversity was observed in samples after composting than storage. Our current research shows that both composting and storage effectively reduce ARGs in cattle manure. However, it's challenging to determine which method is superior, as different groups of resistance genes react differently to each treatment, even though a notable overall reduction in ARGs is observed.

1 Introduction

The increased demand for food of animal origin has resulted in growing numbers of food-producing animals producing large quantities of manure; indeed, a single dairy cow can produce 54 kg of wet manure per day (Girotto and Cossu, 2017). Such production has raised environmental concerns which are driving food animal producers to identify economically and environmentally-safe solutions for manure disposal.

Approximately 21% of global cattle milk production takes place in the European Union, with almost half of this EU production being concentrated in Germany, France, and Poland (FAO, 2020).

Many cases of antibiotic overuse or misuse have been reported in the animal production sector, and this has been associated with the emergence of antibiotic resistance in bacteria. After ingestion, only a small amount of the consumed antibiotics is absorbed or metabolized from the animal intestine, with about 75% of the applied antimicrobial being excreted into the feces or urine: the exact percentage depends on the antimicrobial class (Sarmah et al., 2006; Amarakoon et al., 2016; Spielmeyer, 2018; Filippitzi et al., 2019). In Europe, regulations limiting antimicrobial use in animal production are restrictive, and antibiotics are applied only to treat bacterial infections; the use of antibiotics as growth promoters has been limited. However, in other regions, antibiotics are still used as growth promoters (Dibner and Richards, 2005; Centner, 2016; Hu and Cowling, 2020).

Manure-borne antimicrobial resistance (AMR) determinants such as antibiotic-resistant bacteria (ARB), antibiotic-resistance genes (ARG), and mobile genetic elements (MGE) can contaminate surface water, groundwater, rivers, ponds, lakes, soil, or air through several pathways: (a) directly through runoff from animal production/breeding systems, (b) via the direct application of animal manure on arable fields for crop production, (c) through irrigation water from manure storage ponds and manure treatment facilities, and (d) via air particles (McEachran et al., 2015; Nguyen et al., 2020).

Due to the breeding programs applied in the dairy industry, modern dairy cows are now highly productive; however, they are also prone to a range of infections, particularly those related to the mammary gland (Zalewska et al., 2020). In such cases, bovine mastitis, dry cow therapy, and udder health disorders in dairy cattle are often treated using various antibiotics, including cephalosporins, penicillins, and macrolide-lincosamides (Barlow, 2011). In fact, these conditions are the most common cause of antibiotic use, followed by respiratory, reproductive, and gastrointestinal diseases (Ferroni et al., 2020). Antimicrobial prescription practices vary by country, and cattle are also often administered aminoglycosides, amphenicol, quinolones, and sulfonamides, often combined with trimethoprim (Merle et al., 2012; Diana et al., 2020; Ferroni et al., 2020; Olmos Antillón et al., 2020). In addition, dairy cattle demonstrate significantly higher mean annual antibiotic consumption than beef cattle (Merle et al., 2012; Ferroni et al., 2020).

The gastrointestinal tract of dairy cattle is inhabited by a diverse bacterial community, whose composition and function play key roles in animal wellness and production efficiency. The composition of the community varies along the gastrointestinal tract; however, the most common bacteria overall belong to the phyla *Firmicutes* (65%), *Bacteroidetes* (15%) and *Proteobacteria* (13%) (Mao et al., 2015). Unfortunately, this rich microbiota may also harbor ARG. Animal feces can also become a source of AMR spread, especially in more intensive livestock farming scenarios, characterized by both high amounts of bacteria and significant antimicrobial selective pressure, promoting the occurrence and enhancement of ARB and ARGs (Ferroni et al., 2020; Huygens et al., 2021). Antibiotic treatment of cattle induces shifts in the taxonomic structure and biodiversity of the gastrointestinal microbiota (Beyi et al., 2021); however, this effect varies between the compartments of the digestive tract, with the rumen being the most heavily impacted (Thomas et al., 2017).

Dairy cattle can produce as much as 7.3 kg of dry mass of manure per animal per day (compared to 5.5 kg in feedlot cattle), which is equivalent to 7-8% of the body weight of the cow (Font-Palma, 2019). In 2020, manure production in dairy cattle in Poland totaled approximately 149 million kg of nitrogen content, accounting for about 7.5% of production in the EU (FAO, 2020). The easiest disposal strategy for unprocessed manure is direct land application; however, such raw manure might contain antibiotic residues, ARB and ARG, thereby increasing the risk of AMR being spread into the environment. This poses a significant threat to both human and animal health. The persistence and leaching of antibiotics into the ground and surface waters depend on the physiochemical properties of

the antimicrobial compound and the fertilized soil. These compounds can persist for extended periods,
and eventually accumulate in crops (Zalewska et al., 2021). The continuous presence of low
concentrations of antibiotics creates selective pressure in the environment, promoting the growth of
resistant bacteria (Gullberg et al., 2011). The presence of ARG in the environment has even greater
implications compared to the presence of antibiotics alone (Ahmed et al., 2021). Consequently,
extensive research on the effects of manure management practices is necessary to understand their
effect on microbial communities and the spread of AMR.

The choice of manure management method can influence the microbial community of the manure
and reduce the survival of pathogens (Heinonen-Tanski et al., 2006). The most common strategy
involves storing the feces for a prolonged time, usually between 4 to 7.5 months; however, this requires
sufficient storage capacity (Loyon, 2018). An alternative strategy is the composting of manure, which
is both economically and environmentally beneficial, as it reduces solid organic waste volume while
eliminating pathogens, antibiotic residues and ARG (Dolliver et al., 2007; Tasho and Cho, 2016; Gou
et al., 2018). Composting transforms organic waste into a stable, humus-like product, that is easily
available for soil amendment. Although composting offers numerous advantages, estimates from 2011
show that only 0.8% (10.4 million tons) of total EU livestock manure was composted. Moreover, only
about 8% of the manure used in fields underwent any form of treatment (Foged et al., 2012). In the
Netherlands, 75% of manure from calves was applied to agricultural land without any treatment
(Berendsen et al., 2018). It is believed that the primary reason for omitting such processing is the lack
of adequate storage capacity (Köninger et al., 2021).

The structure of the microbial community in cattle manure and its resistome (pool of ARGs in
the microbial community) remain relatively unknown, with a lack of comprehensive data on the
subject. Until now, no studies have compared the effects of storage and composting of manure
originating from the same source; only a few have examined the effects of multiple manure treatment
strategies on the same raw manure batch. Therefore, the present study compares the effect of storage
and composting on a single manure batch on their ability to remove ARG. The analysis considers the
type of process as the independent variable, with ARG relative abundance and microbial phylum
relative abundance as the dependent variables. Hence, the aim of the study was to determine how
composting and storage of dairy cattle manure impact the ARG and microbiome of animal feces, with
the goal of limiting the potential risk of spreading ARG in the fields and compare these both manure
treatment techniques in term of better preparation manure for land application. The material was
collected simultaneously from the same animals, allocating one portion of the sample for storage and
another for composting. ARG distribution was identified by high-throughput qPCR and the microbial
community was characterized via metagenomic sequencing targeting of the V3-V4 variable regions of
the 16S rRNA gene.

2   Materials and methods

2.1   Cattle manure collection

The study was conducted on 50 Polish Holstein-Friesian dairy cows, Black and White variety.
The herd was located in the central part of Poland (Masovian voivodeship). The animals were kept in
a loose barn with free access to water, and fed the same total mixed ration (TMR) diet ad libitum,
consisting of corn silage (75%), concentrates (20%), and hay (5%), supplemented with a mineral and
vitamin mixture, according to the INRA system (Brzóska et al., 2014). The cattle manure samples were
collected in October 2019. The manure samples were deposited into two plastic drums measuring

approximately 20L each. One aliquot of manure was subjected to composting, and the other to storage
on a laboratory scale. The farm owner agreed to the manure being sampled and shared the history of
the use of antibiotics on the farm.

2.2   Manure management strategies

Defined amounts of cattle manure (approx. 5 kg) were mixed with Sitka spruce (4:1 ratio; w/w)
and placed in a bucket. The Sitka Spruce was sourced from a local hardware store. Containers with
manure were placed in the laboratory under a ventilated hood at room temperature. During composting,
the manure was mixed well each week for ten weeks, while the stored manure was left undisturbed for
four months. Although the duration of both processes differs, the composting and storage are both,
highly recommended for manure treatment before its land application and both were applied according
to commonly used practices - the composting process was completed within eight weeks (56 days), as
noted by Mc Carthy et al. (2011) and the four-month storage period was selected based on EU
legislation. Samples for the microbiological and molecular analysis were collected at the beginning
(raw/untreated manure), in the middle (a fifth week for composting - 5W; second month for storage -
2M), and at the end of the process (a tenth week for composting - 10W; fourth month for storage - 4M).
The experiment was conducted in triplicate. A total of 200 samples were collected during the entire
experiment. From each group, we gathered 40 samples, encompassing three technical repetitions. The
groups included control samples, composted samples at five and ten weeks, and stored samples at two
and four months. DNA isolated from these samples was grouped accordingly and then dispatched for
analysis.

The pH, moisture, and temperature were measured in both mixtures at approximately half of their
height (~ 15 cm) each week for ten weeks (enough time to observe the dynamics of changes during
composting and to stabilization of parameters during storage). The moisture content was adjusted to
50-60% for compost by adding sterile MilliQ water. The moisture, pH and temperature was measured
directly in the containers with treated samples along with the time. The moisture and pH was measured
by dual-purpose pH meter dedicated to environmental samples (pH-meter Stelzner 3000/Soiltester).
Temperature was measured by laboratory thermometer (Testo 905-T1).

2.3   DNA extraction

The total DNA was isolated using the commercially available FastDNA™ SPIN Kit for Feces
(MP Biomedicals) according to the manufacturer's recommendations. The quality and quantity of the
extracted DNA was analyzed with a Qubit 4.0 Fluorometer using the dsDNA high-sensitivity assay kit
(Invitrogen) and a Colibri spectrophotometer (Titertek Berthold). The DNA samples were isolated in
triplicate and then pooled to obtain a single DNA sample for each time point.

2.4   Metagenomic sequencing

The bacterial community structure was determined using Illumina technology on the Miseq
platform by sequencing the variable V3-V4 regions of bacterial 16S rRNA. The libraries were prepared
with Nextera® XT index Kit v2 Set A (Illumina). Once prepared, amplicons underwent sequencing,
2x300 bp paired-end sequences. PCR and sequencing was performed at the DNA Sequencing and
Oligonucleotide Synthesis Facility (Institute of Biochemistry and Biophysics Polish Academy of
Sciences). Raw sequences were processed and analyzed using Qiime2 software suite (Bolyen et al.,
2019) with the DADA2 option for sequence quality control and the newest release of the SILVA
(SILVA SSU database 138, accession date October 2021) ribosomal RNA sequence database for

taxonomy assignment (Quast et al., 2013; Yilmaz et al., 2014). The 99% cutoff of identified OUT were
applied.

2.5   High-throughput qPCR

High-throughput qPCR was performed by an outside firm (Resistomap, Finland) using the qPCR
SmartChip Real-Time PCR cycler (Takara). The qPCR cycling conditions and initial data processing
were carried out as previously described by Wang et al. (2014). The analysis used a 384-well template
of primers. All samples used for the analysis met the following criteria: 260/280 ratio of 1.8-2.0 (+/-
0.1) and a concentration of 10 ng/μl.

The threshold cycle (Ct) values were calculated using default parameters provided by the
SmartChip analysis software. All samples for each primer set underwent qPCR efficiency and melting
curve analysis. Amplicons with unspecific melting curves and multiple peak profiles were considered
false positives and excluded from further analysis. Samples meeting the following criteria were
selected for analysis: (1) Ct ≤ 27, (2) at least two replicates, (3) amplification efficiency 1.8 - 2.2. The
relative copy number was determined using an equation originally published by Chen et al. (2016) and
adapted to our study conditions. The gene copy numbers were calculated by normalizing the relative
copy numbers per 16S rRNA gene copy numbers.

$$\text{Relative gene copy number} = 10^{(27-Ct)/(10/3)}$$

The genes and their corresponding primer sequences can be found in Supplementary Materials
File 1. The gene names and groupings, as detailed and organized in the table from Supplementary
Materials File 1, are consistently referred to throughout the manuscript.

2.6   Statistical analysis

The minimal sample number was determined by G*Power software (version 3.1.9.7) (Faul et al.,
2007, 2009). The metagenomic data, including ARG distribution, was analyzed and visualized using
the MicrobiomeAnalyst web server (https://www.microbiomeanalyst.ca/)(Chong et al., 2020). Data
were not rarefied, not scaled and transformed. The differences in microbial community structure were
evaluated using the Qiime2 pipeline, based on the Kruskal-Wallis H-test (Shannon and Chao1 indexes).
The differences between ARG groups at each stage of the process were calculated using analysis of
variance (ANOVA) with post hoc Tukey's test (GraphPad Prism 9). The correlations between ARG
and MGE were calculated using Spearman's correlation coefficient with GraphPad Prism 9. The
networks representing correlations were built with Cytoscape (3.9.0 version) (Shannon et al., 2003).
The correlations were considered strong and significant when the absolute value of Spearman's rank
$|r|>0.7$ and $p < 0.05$ (Zhu et al., 2017).

3   Results

3.1   The changes in physicochemical properties of treated manure

206 The humidity, pH, and temperature were recorded weekly over a ten-week period, as detailed in
207 Table S1 of Supplementary File 2.

3.2   Microbiome composition

During the analysis we identified 28 OUT at phylum level (Table S2 Supplementary Material File 2 - mean abundance with standard deviation of each phylum across the treatment). In all samples, the microbial community composition was determined for the untreated (raw) cattle manure and the composted and stored manure; these determinations were performed at the beginning, middle and end of the process. The composition of bacterial communities at the phylum level is presented in Fig. 1. using the taxon bar plot, and the core microbiome is presented in Fig. S1 (Supplementary material file 2). The richness and diversity of the microbial communities were determined using the Shannon (comparable results, p-value 0.08) and Chao1 (higher value for composted samples, p-value 0.025) indexes (Fig. S2, S3 Supplementary Material File 2). The raw data obtained from sequencing was uploaded to the Sequence Read Archive (SRA) repository (BioProject ID: PRJNA1007935) (NCBI). The PCoA of taxon abundance in all sample groups, calculated based on the Bray–Curtis distance, revealed significant visual separation between the initial (untreated) and treated samples (p-value<0.01) (Fig. S4 Supplementary Material File 2; the results of pairwise PERMANOVA analysis are presented in Table S2 Supplementary Material File 2).

Figure 1. Relative abundance of phyla across samples; the samples were divided into five groups– control (cow manure raw; sample before treatment), cow manure composting 5W, cow manure composting 10W (composted samples after five weeks and ten weeks, respectively), cow manure storage 2M and cow manure storage 4M (stored samples after two months and four months, respectively); stacked barplots represents mean value for replicates.

3.3 Diversity and abundance of ARG

During the study, we found 202.67 analyzed genes (Aminoglycoside-, Beta-Lactam-, MLSB-, Phenicol-, Sulfonamide-, Tetracycline-, Trimethoprim-, Vancomycin-resistance genes together with Integrons, MDR, MGE, and 'Other' genes) in raw manure, 76 in composted samples after five weeks, 54 after ten weeks, and 51 in stored samples after two months and 39 after four months. The relative abundances of the analyzed ARG classes and MGE varied as storage and composting processed (Fig. S4 Supplementary Material File 2). Specifically, the relative prevalence of beta-lactam-, MLSB- and vancomycin-resistance genes fell in the treated manure samples compared to the untreated samples. In contrast, the relative abundance of genes encoding MGE, integrons and sulfonamide-resistance genes increases, but only after composting. However, the total gene count (i.e. total resistance gene abundance normalized per 16S rRNA gene) rapidly decreased following the application of both treatment strategies. Notably, for composting, the gene relative abundance was even lower in the middle of the process than at the end (Fig. 2).

Figure 2. The differences in the amounts of ARG and MGE detected during composting and storage compared with raw manure. The following cut-off points for significance within the same process were applied: the values differ significantly at $p < 0.01$ (indicated as *); the values differ significantly at $p < 0.05$ (indicated as **); the values do not differ significantly at $p > 0.5$. Relative abundances of genes were additionally compared at the end of both processes between them; the value with the same letters differs significantly: A, A – at $p<0.01$.; a, a at $p<0.05$ (indicated as A,B,C,D,E,F); the values differ significantly at $p < 0.05$ (indicated as d).

Composting efficiently reduced the quantity of aminoglycoside- and tetracycline-resistance genes, as well as MGE, within the initial five weeks (2-fold, 5-fold, 2.5-fold reduction respectively, $p<0.01$). However, a subsequent rise in their abundance was observed after ten weeks of treatment compared to their level after five weeks of treatment, compared to raw material it still remains lower (1,2-fold, 3-fold and 2-fold reduction respectively, $p<0.01$). Beta-lactam-, vancomycin-resistance genes, MLSB-coding genes, and genes responsible for MDR demonstrated a decline at both the five- (66-fold for beta-lactam-resistance genes, 74 for MLSB-coding genes, and 6-fold for MDR genes, with vancomycin-resistance genes being below detection limit; $p<0.01$) and ten-week (156-fold for beta-lactam-resistance genes, 80-fold for MLSB-coding genes, and 5-fold for MDR genes, with vancomycin-resistance genes being below detection limit; $p<0.01$) marks, with the rate of reduction being consistent between the two points. The abundance of Integron-coding genes decreased after five weeks of composting (1.5-fold, $p<0.01$), but remained unchanged after ten weeks compared to raw sample and sample after five weeks of composting. Neither phenicol- nor trimethoprim-resistance gene abundance varied between composting stages. However, the sulfonamide-resistance gene count increased after five weeks of composting (0.6 fold, $p<0.05$), and exhibited an even greater increase after ten weeks (0.33-fold, $p<0.01$).

Our results show that storage effectively diminishes the quantities of aminoglycoside-resistance genes (5-fold, $p<0.01$), MGE (9-fold, $p<0.01$), and genes responsible for MDR (12-fold, $p<0.01$) and MLSB-coding genes (8.5-fold, $p<0.01$) after two months. However, their numbers subsequently increased after four months of treatment, but still remains lower than in raw manure ($p<0.01$). The counts of Beta-lactam, vancomycin-resistance genes, and those labeled as 'Other' demonstrated a decline both at the two- (289-fold, 8.5-fold, respectively, with vancomycin-resistance genes being below detection limit; $p<0.01$) and four-month marks (579-fold, 7-fold, respectively, with vancomycin-resistance genes being below detection limit; $p<0.01$), yet the rate of reduction remained consistent between these points. The abundance of genes coding for integrons remained consistent after two months of storage but declined after four months (3-fold, $p<0.01$). No noticeable change in the amounts of phenicol-, sulfonamide-, and trimethoprim-resistance genes was observed across the storage periods. The tetracycline-resistance genes were lower after both the two (8-fold, $p<0.01$) and four month (4-fold, $p<0.01$) periods, but their concentration was notably higher at the end of the fourth month compared to the second.

In conclusion, both composting and storage processes reduce the abundance of aminoglycoside-resistance genes, with storage achieving a greater reduction. A similar trend can be observed for the abundance of MLSB-coding genes, MGE, and genes classified as 'Other'. For beta-lactam, tetracycline-, vancomycin-resistance genes, and genes responsible for MDR, a reduction in gene abundance was noted after both processes. However, composting and storage yielded comparable outcomes by the end of treatment. While storage resulted in a decreased abundance of integron-coding genes relative to untreated samples and ten-week composted manure, composting itself did not appear to affect these genes. For sulfonamide-resistance genes, composting increased their amount, while storage did not include any noticeable changes. Nevertheless, after two months of composting, the abundance of sulfonamide-resistance genes remained elevated compared to post-storage levels. The quantities of the phenicol- and trimethoprim-resistance genes did not demonstrate any significant variation after the two treatments compared to untreated manure (Fig. 2).

The relative abundances of ARG and MGE across all analyzed samples were subjected to principal coordinate analysis (PCA), using the Bray-Curtis distance method. The results indicate a significant visual separation between the untreated manure (control) and treated samples with regard to the relative abundances of resistance gene classes ($p<0.01$).

The core resistome across all analyzed samples consists of 29 genes. They belong to the
following groups: aminoglycoside-resistance genes (*aadA5_1, aadA_2, aac3-VI, aacC2, aadA1, strB,*
aadA2_3, aadA2_1), beta-lactam-resistance genes (*fox5*), phenicol-resistance genes (*floR_1*),
sulfonamide-resistance genes (*sul1_1, sul2_1*), tetracycline-resistance genes (*tetA_2, tetG_1, tetG_2*),
trimethoprim-resistance genes (*dfrA1_1, dfrA1_2*), integron-coding genes (*intI1_3, intI1_1, intI1_4*),
genes responsible for MDR (*mdtH_1, oprJ, acrR_3*), MGE (*tnpA_2, ISSm2, IncP_oriT, repA*), and
resistance genes classified as 'Other' (*qacEΔ1_2, qacEΔ1_3*). For both applied treatment strategies, a
decreased number of ARGs were observed post-treatment compared to raw manure (53 for compost
and 38 for storage) (Fig. S5 and Fig. S6 Supplementary material file 2).

In all composted and stored samples, almost all ARGs, excluding trimethoprim-resistance
genes, were found to be reduced at the end of both processes, which means lower diversity of genes
after treatment. In addition, the stored samples demonstrated lower numbers of aminoglycoside-, beta-
lactam-, phenicol-, sulfonamide-, tetracycline-resistance genes and MGE compared with composting.
However, the composted samples demonstrated lower numbers of genes conferring MLSB and MDR
resistance compared to the stored samples (Fig. 3).

Figure 3. Comparison between ARG (assigned to ARG classes) at the end of both treatment processes.

In the composted samples, the correlation network analysis revealed 124 nodes connected by
512 edges, representing strong and significant correlations (|r|>0,99; p<0.05) (Fig. 4). All correlations
were positive. The MGE (integrons included) exhibited more links than ARG, indicating their
significant role in network formulation. The MGEs (n=15) included in this network are *intI3_1,*
orf39_IS26, IncN_rep, Tn5, Tp614, tnpA_5, ISAba3, trfA, IS613, intI3_2, tnpA_6, intI1_3, ISSm2,
IS1111, tnpA_2.

In the stored samples, strong and significant correlations were found between MGE (integrons
included) and ARG levels, manifested as a network containing 144 nodes connected by 643 edges
(|r|>0.99; p<0.05) (Fig. 5). Every correlation in the network was positive. The MGE (integrons
included) had more connections than ARG, indicating they play a pivotal role in the formation of the
network. The MGE (n=16) presented in the network are *intI3_2, IS613, IS1111, ISAba3, orf37-IS26,*
tnpA_1, trfA, tnpA_5, orf39_IS26, Tn5, Tp614, tnpA_6, intI2_2, IncN_rep, intI1_1, intI3_1.

Figure 4. ARGs and MGEs interaction network in composted manure. The network is presented as an
'organic layout'. Based on Spearman's rank correlation, a strong and significant correlation is shown,
where |r|>0.99 and p < 0.05. The size of the nodes represents the degree of interaction. The gray edges
show positive correlations between ARGs and MGEs. The color of the nodes indicates ARG (white)
and MGE (gray).

Figure 5. ARGs and MGEs interaction network in stored manure. The network is presented as an 'organic layout'. Based on Spearman's rank correlation, a strong and significant correlation is shown, where |r|>0.99 and p < 0.05. The size of the nodes represents the degree of interaction. The gray edges show positive correlations between ARGs and MGEs. The color of the nodes indicates ARG (white) and MGE (gray).

4   Discussion

Farms often dispose of animal manure by direct application to the land. Many also employ a range of manure management technologies to reduce environmental pollution before application. These technologies not only reduce the total manure volume, but also can be used to produce organic fertilizer, or possibly facilitate biogas production. Two such technologies are storage and composting, after which the feces are often used as organic soil amendments with potential benefits for soil quality and crop yield (Epelde et al., 2018). Such amendments typically contain all nutrients essential for plant growth and can replace or complement inorganic fertilizers in agricultural production systems (Youngquist et al., 2016). However, cattle manure can also serve as a reservoir of environmental ARG pollution (Udikovic-Kolic et al., 2014; Zalewska et al., 2021), and the direct use of livestock manure without proper treatment could promote the dispersal of ARG in arable soil and cultivated plants (Ghosh and LaPara, 2007; Heuer et al., 2011; Udikovic-Kolic et al., 2014; Zalewska et al., 2023). Manure treatment strategies have not been specifically designed to mitigate AMR (Oliver et al., 2020).

Our present findings indicate that composting and storage are effective methods of reducing ARG from manure before use. It is hard to define which process accomplishes this task better, as despite a generally-observed significant reduction in the number of ARG, particular groups resistance genes responded differently to the applied strategies. For example, the levels of aminoglycoside-resistance genes, integrons and MGE, genes classified as 'other' and sulfonamide-resistance genes were lower after storage than composting. In contrast the MLSB levels in the stored samples were higher than those in the composted samples, but they were still lower than those found in the raw material. No differences were observed between the composted and stored samples for beta-lactam, MDR,

tetracycline, and vancomycin-resistance genes; those genes are often prevalent in animal manures. Similar results for composting were obtained by Gou et al. (2018), who found higher amounts and numbers of ARGs in unprocessed manure compared to composted samples.

As many of the antibiotic groups used in the animal production sector are considered critically important antimicrobials (CIA) or highly important antimicrobials (HIA) for human medicine (Zalewska and Popowska, 2020), i.e. the last-resort option for the treatment of certain bacterial infections, it is crucial to limit the spread of resistant bacteria or resistance genes encoding a particular resistance mechanism. Tetracyclines, penicillins, and macrolides, the most commonly used antibiotics in livestock, are categorized as HIA, CIA, and CIA respectively. Moreover, many of the antimicrobials used in dairy cattle also have significant treatment roles: first and second generation cephalosporins - HIA; third, fourth and fifth generation cephalosporins - CIA; penicillins - CIA; macrolides - CIA; lincosamides - HIA. In dairy cattle production sector, commonly used antimicrobial classes include aminoglycosides - CIA; amphenicol - HIA; quinolones - CIA; and sulfonamides (often combined with trimethoprim) – HIA. Therefore, the presence of resistance genes against those antimicrobial agents in the environment should be especially alarming.

Long-term storage and composting of dairy cattle manure results in a different prevalence of specific genes than observed in untreated manure. The observed reduction of ARG abundance in different composting processes may result from the elimination of ARG-bearing bacteria during the aerobic thermophilic phase (Wang et al., 2015; Xu et al., 2018). A well-executed aerobic compost treatment achieves a higher peak temperature during the thermophilic phase and is more effective in mitigating antibiotic residues, compared to static storage in the stockpiles. However, the efficiency strongly depends on the presence of antibiotic residues and their interactions (Youngquist et al., 2016). A study on ARG reduction in dairy manure storage in lagoons shows that higher incubation temperatures are required for significant reduction of ARG, with less efficient ARG mitigation observed at ambient temperature (Pei et al., 2007). Other studies on dairy manure storage reveal that depending on the local conditions, some ARG may decrease while others may increase (Ruuskanen et al., 2016). Typical concentrations of antibiotics in livestock manure are between 1 and 10 mg/kg, but could reach up to 200 mg/kg (Massé et al., 2014).

As tetracyclines and beta-lactam antibiotics are widely used in cattle production systems, their detection methods are well established. Tetracycline resistance genes are prevalent and persistent in cattle manure, unlike beta-lactam resistance genes, despite the common use of penicillins and cephalosporins. This suggests that beta-lactams are unstable in cattle feces: they have a very sensitive ring-shape structure, which is readily cleaved by ammonia, phosphate, and hydroxyl ions present in composted manure, resulting in their complete degradation (Kakimoto et al., 2007). Sulfonamide, macrolide, and lincosamide residues have also been found in animal manures, albeit in lower concentrations (usually below 1 ppb) (Oliver et al., 2020).

Zhang et al. (2019) found that approximately 64.7% of detectable veterinary antibiotics were efficiently removed from manure during composting; however, precise detection of antibiotics in complex matrices such as composting mixtures or feces remains challenging. Despite this, antibiotic residue degradation appears to be primarily influenced by abiotic factors, such as temperature, pH, hydrolysis, and sorption kinetics. Tetracyclines are also subject to biotic degradation processes (Youngquist et al., 2016). Although antibiotic levels universally decline during the manure treatment process, albeit to a varying degrees, its effect on ARB and ARG is less consistent. Compared to storage, composting appears to be more efficient at reducing ARG. However, ARG may persist after treatment and facilitate further proliferation in soil when manure is used as fertilizer. Moreover, it has been stated

409 that even though medically relevant ARB were eliminated during pretreatment, their ARGs may persist
410 and transfer to commensal soil bacteria, and continue to pose a risk after treatment (Edrington et al.,
411 2009).

4.1 Microbial community structure

Wang and Zeng (2018) reported that bacteria play a greater role in composting than fungi. Pitta et al.
(2016) identified several prevalent bacterial phyla in dairy cattle feces and raw manure: *Bacteroidetes,*
Firmicutes, Proteobacteria, Actinobacteria, Acidobacteria and *Spirochaetes*. This aligns with our
present findings, which also included *Spirochaetes* and *Acidobacteria*, though they were not the most
represented phyla. Moreover, Zhou and Yao (2020) report the presence of *Proteobacteria* and
Firmicutes as the main phyla in animal feces.

The decomposition of organic matter starts before composting commences, as microorganisms
are introduced unintentionally to the organic matter during grinding, mixing, cutting, and transporting.
The composting pile initially rests at an ambient temperature. It then gradually heats up during the
mesophilic phase to 25-45 °C, indicating increased microbial activity (Ezugworie et al., 2021). The
diversity of microbial community correlates with the metabolic functions of the microbes, e.g., lactic
acid-producing bacteria and acetate-oxidizing bacteria are usually detected at the mesophilic phase of
the composting process, where they utilize the easily degradable and soluble organic matter. Zhang et
al. (2022) have found that *Firmicutes, Proteobacteria, Bacteroidetes, Chloroflexi, Actinobacteria*, and
Planctomycetes were the dominant phyla during co-composting of cow manure and maize straw. In
addition, Zhang et al. (2020) report the presence of *Firmicutes, Bacterioidetes, Deinococcus-Thermus,*
Proteobacteria, and *Actinobacteria* in raw manure and composted samples; however, the abundance
of *Deinococcus-Thermus* and *Actinobacteria* were much higher in composted samples. Similarly, Zhou
and Yao (2020) noted that composting resulted in a shift in the bacterial community, with an increased
prevalence of *Actinobacteria* compared to raw manure. Our present findings also identified
Proteobacteria, Firmicutes, Planctomycetota, and *Actinobacteriota* as being the most prevalent during
composting; however, their ratio changed with time. Zhang et al. (2016) found the *Firmicutes,*
Proteobacteria, Bacteroidetes, and *Actinobacteria* to predominate in the thermophilic phase of
composting. Of these, *Firmicutes* were the dominant phylum due to their adaptability to aerobic
condition and heat resistance. Temperatures exceeding 70 °C terminates most microbial activities,
forcing certain bacteria to sporulate (Ezugworie et al., 2021).

While our sequencing results differed from those found in some previous studies, the identified
phyla remained consistent. Composting can be influenced by a range of physicochemical factors, such
as temperature, moisture, pH, aeration, and carbon-to-nitrogen ratio. As such, as there are no strict
guidelines on composting, composted samples can differ widely. A crucial role is played by
temperature: it governs the pace of biological processes conducted by microorganisms and determines
the presence of particular microorganisms; for example, fungi, which are not heat-resistant, are
eliminated at temperatures above 50 °C. In addition, a pH of 7-8 is needed for optimal composting,
which slows down at low pH. The C/N ratio should also be kept at optimum, as nitrogen is crucial for
microbial growth, and carbon is an energy source. Humidity should be kept at 40-60% to ensure the
transfer of dissolved nutrients necessary for microbial growth (Ezugworie et al., 2021). As such, the
differences in bacterial community structures obtained by different research teams may hence be due
to the differences in physicochemical parameters during the process. In our case, the temperature was
within the proper range, but the humidity and pH were slightly suboptimal (humidity above 80% and
pH below 7).

4.2 Changes in antibiotic resistance genes after treatment

4.2.1 Tetracycline-resistance genes

The tetracyclines are the most commonly used antibiotics in the livestock production sector; as such, tetracycline-resistance genes are the subject of intensive study. These genes appear to be unaffected by manure treatment strategies such as composting or storage, and the detection rate of t*etW* and *tetO* may account for up to 80% of fecal samples obtained from cattle raised in grassland-production systems, even without antibiotic usage (Santamaría et al., 2011). The following tetracycline-resistance genes were identified in the present study: efflux (*tetA, tetA/B, tetG, tetH, tetC, tetL, tetA(P)*), protection (*tet(36), tet(32), tetO, tetQ, tetW, tetM, tetPB*), deactivation (*tetX*) and regulation (*tetR*) in raw manure samples. Our findings are partially in line with those of other studies (Jauregi et al., 2021). Zhang et al. (Zhang et al., 2020) found *tetW, tet40, tetO, tetQ*, and *tetC* to be the most prevalent types across all analyzed raw dairy cattle manure samples, accounting for over 40% of detected ARGs. Marti et al. (2013) report the presence of *tetM, tetQ, tetS, tetT, tetA, tetB/P* and *tetW* in cattle manure, and Mu et al. (2015) found *tetM, tetO, and tetW* in cattle manure, and *tetM, tetO, tetQ*, and *tetW* in feces collected per rectum; the latter partially agrees with our present findings indicating these genes to be present in manure.

However, Zhou and Yao (Zhou and Yao, 2020) reported a general increase in the abundances of total tetracycline-resistance genes, which contradicts our results. Despite this, as in our present study, Zhang et al. (Zhang et al., 2020) failed to detect *tetA, tetW*, and *tetR*, and that total ARG abundance decreased after composting. Interestingly, Zhang et al. (2020) also indicate a reduction in *tetW, tetA*, and *tetR* abundance with composting time, together with a decrease in the concentration of the total amount of ARGs. Staley et al. (2021) did not identify any differences in the concentrations of *tetO* and *tetQ* genes between composted and stored samples; however, their level was found to change with sampling season and depth. In addition, Qian et al. (2016) report increased abundance of *tetC* and *tetX*, and decreased abundance of *tetQ, tetM* and *tetW* after composting.

Interestingly, while Wang et al. (2019) did not identify any decrease in *tetW, tetO* or *tetM*, Storteboom et al. (2007) indicate that the *tetO* and *tetW* genes appeared to gradually decrease during composting; this agrees with our present findings indicating that *tetO* was present in the middle of composting, but below the detection limit at the end of the process. However, our data indicate that *tetW*, together with *tetA, tetG, tetPB, tetQ, tetM, tetR, tetX,* and t*etA/B*, remained persistent during composting.

Another study examining the tetracycline-resistance genes *tetO* and *tetA* found no noticeable change in their abundance, even after one year of storage (Hurst et al., 2019). These findings are in line with ours, where *tetA* remained persistent throughout storage, contrary to *tetO*. Interestingly, Jauregi et al. (2021) found *tetW, tetO, tetM, tetB, tet32, tetN*, and *tetG* to be present in stored cattle manure even after six months; in contrast, *tet32, tetO, tetW,* and *tetM* were below the detection limit after two months of storage but reappeared after four months.

Tetracycline-resistance genes can be transmitted to gram-negative and gram-positive bacteria species, e.g., *tetM, tetW*, and *tetQ* originate from gram-positive bacteria and can be transmitted across Gram-positives and Gram-negatives (Chopra and Roberts, 2001). Moreover, tet genes are often placed on plasmids or inserted in transposons, facilitating their dissemination (Lima et al., 2020). Jauregi et al. (Jauregi et al., 2021) regard *tetM, tetO, tetW, tet32*, and *tetS* as being of particular concern due to their high dissemination potential associated with their co-occurrence with MGE; they also highlight *tet44* and *tetW/N/W*, which were not detected during our study. Our results partially agree, as our data

497 also indicates some tetracycline-resistance genes to be significantly associated with MGE: *tetM (tnpA,*
498 *IS613, trfA, orf37_IS26, ISAba3, IS1111, IntI3), tetO (tnpA, trfA, orf39_IS26, orf37_IS26, ISAba3,*
499 *IS1111, Tn5, IntI3, tp614, IS 613), tet32 (intI3)*; however, neither *tetW* or *tetS* were confirmed in
500 samples obtained from either process. We also found *tet36, tetPB, tetC, tetA(P), tetC, tetH,* and *tetL* to
501 have a high probability of being exchanged by HGT due to co-occurrence with MGE.

4.2.2 Beta-lactam-resistance genes

The following beta-lactam-resistance genes were identified in raw manure, classified as
deactivation (*ampC, blaMOX/blaCMY, blaOCH, blaSHV, blaROB, blaOXY, cphA, cfxA, blaCMY2,*
ampC/blaDHA, blaFOX, blaCTX-M, blaSFO, blaVIM, blaKPC, blaOXA, blaNDM, blaACC, bla1,
blaCMY, blaACT) and protection (*penA, pbp2b*). Our results are similar to those of Marti et al. (2013),
who identified *blaOXA, blaCTX-M, blaPSE* and *blaVIM,* as well as *blaTEM,* which was not found in
the present study. Pitta et al. (2016) report the presence of *pbp1A, pbp2*, and *pbp2B* in dairy cattle
manure; however, these were not present in our samples. While Zhou and Yao (2020) confirm our
observed general decrease in beta-lactam-resistance genes, Keenum et al. (2021) found the levels of
beta-lactam-resistance genes to be elevated after composting.

Hurst et al. (2019) did not identify *blaOXA, blaCTX-M,* and *blaVEB-1* after long-term storage of
dairy cattle manure. This is in line with our present findings, which did not identify any beta-lactam-
resistance genes present in raw manure in the samples subjected long-term storage and composting,
except for the *fox5* gene; however, *blaVEB* was not observed in our raw manure.

The *blaTEM* and *blaCTX-M* genes are often associated with *IncN* plasmids, which are believed
to play an essential role in beta-lactamase gene dissemination (Lima et al., 2020). Our study found *bla*
genes to be significantly associated with *IncN_rep* in composted manure samples but not in stored
ones. Moreover, beta-lactam-resistance genes are often located in insertion sequences such as *ISEcp1*
or *ISCR1* (Lima et al., 2020). Although our findings did not identify the insertion sequences mentioned
above, they did indicate the presence of other MGEs associated with *bla* genes viz. *IS613, trfA, ISAba3,*
IS1111, intI3, tnpA, Tp614, orf39-IS26, Tn5, IS613; this suggests they have high mobility in both
treatment strategies.

4.2.3 Aminoglycoside-resistance genes

The *aadA, aadA5, aadD, aadA9, aphA1/7, aac(6')I1, aac3-VI, aacC2, aadA1, aadE, strA, strB,*
acc, aacC4, aph6, aac(6')-Ib, aadA2, aph(2')-Ib, aphA3, aadA9, spcN, aac(6')-II, aphA3 genes were
observed in the present study, all of which are responsible for deactivation. Zhou and Yao (Zhou and
Yao, 2020) report a general decrease in aminoglycoside-resistance genes during treatment, which
agreed with our results.

Hurst et al. (2019) also found *aac-(6)-lb* and *aada1* to be present in manure after long-term
storage; however, our present findings indicate that *aadA1* was also persistent in manure after storage
as well as composting, but *aac(6)-lb* was not detected after either process. Although *aadA1* persisted
after composting, its concentration was reduced, as also noted by Zhang et al. (2020).

Moreover, Jauregi et al. (2021) found *aad6* and *ant6-lb* to be of particular concern due to their
co-occurrence with MGE and hence their high dissemination potential. However, although these genes
were not identified in raw manure in the present study, other genes of potential concern (i.e. co-
occurring with MGE) were noted: *aac(6")-Ib (intI3, IS613, tnpA, trfA, orf37-IS26, ISAba3, IS1111,*
intI3), aac(6")-Ib (intI3, IS613, tnpA, trfA, orf37-IS26, ISAba3, IS1111) and aphA3 (Tp614, IS613,
tnpA, trfA, orf37-IS26, orf39-IS26, ISAba3, IS1111, Tn5, intI3).

4.2.4 MLSB-resistance genes

541   Despite their wide application in livestock, often with lincosamides and streptogramin, the data
542 on resistance genes against macrolides is limited. Mu et al. (2015) identified *ermB* and *ermC* in cattle
543 feces (collected per rectum) and manure; however, the study only addressed macrolide-resistance
544 genes. Our present data did not confirm the presence of *ermB* and *ermC*, but did identify other genes
545 coding MLSB resistance phenotypes in cattle manure, such as *ermK, ermF, ereB, lnuB, vatE, erm36,*
546 *matA/mel, mphA, mphB, vgb, mefA, msrC, pncA, mphC, msrA, lnuC, oleC, carB*, and *pikR2*. In contrast,
547 Wang et al. (Wang et al., 2019) found *ermB* and *ermF* to be present in raw cattle manure, while Pitta
548 et al. (2016) found *carA, mefA, tlrC*, and *vgaA* genes to also be present in feces and manure.

549   Zhou and Yao (2020) report a general decrease in MLSB-resistance genes in total, which agreed
550 with our results. However, Hurst et al. (2019) found *ermB, ereC*, and *mefA* to remain persistent in
551 manure after long-term storage; our present findings did not indicate the presence of *ermB* in dairy
552 cattle manure, and *ereB* and *mefA* genes to be removed by long-term storage and composting. Wang et
553 al. (2019) did not identify any changes in *ermB* and *ermF* gene abundances during composting, since
554 their levels were too low at the beginning of the process.

555   Lima et al. (2020) report *ermB* to be frequently incorporated in the *Tn916-Tn1545* family of
556 conjugative transposons. Although *ermB* was not found in the present study, other MLSB phenotypes
557 were noted including genes co-occurring with MGE: *IS613, trfA, ISAba3, IS1111, intI3* and *tnpA* in
558 compost, and *IS613, tnpA, trfA, orf37-IS26, ISAba3, IS1111, intI3* and *IncN_rep* in storage. The
559 presence of *IncN_rep* in stored samples should be especially concerning because it suggests the
560 location of erm genes on a broad-range host plasmid from the IncN group.

4.2.5 Resistance genes against other antimicrobials

562   Although the tetracycline- and beta-lactam-resistance genes are considered to be most prevalent
563 in dairy and beef cattle feces, Marti et al. (2013) found chloramphenicol-resistance genes to be
564 predominant, with tetracycline-resistance genes in fourth place. The *cmlA* and, *floR* genes had high
565 dissemination potential, due to co-occurrence (Jauregi et al., 2021), but our data indicates that *cmlA*
566 was efficiently eliminated from animal manure after storage or composting. However, we agree that
567 the *floR* gene should be considered highly mobile as our findings indicate that it could be associated
568 with MGEs such as *IS613, tnpA_1, trfA, orf37-IS26, ISAba3, IS1111* and *intI3*, in both analyzed
569 treatment strategies.

570   The effect of composting on sulfonamide-resistance genes has also been analyzed. Zhou and Yao
571 (2020) report a general decrease in sulfonamide-resistance genes, which was in line with our present
572 results; in addition, Mu et al. (2015) also identified *sul1, sul2*, and *sul3* in cattle manure, as noted in
573 the present study. Additionally, our study also identified *sul4, folP*, and *folA*. The presence of *sul1* and
574 *sul2* was also confirmed (Wang et al., 2019); however, their study did not show a reduction in the
575 abundance of those genes after composting.

576   Furthermore, being often located on transposable elements of self-transposable or mobilizable
577 broad host-range plasmids, *sul* genes are highly prevalent in wide range of bacterial species (Lima et
578 al., 2020). The *sul2* gene should be especially concerning, because it is often embedded on small mobile
579 plasmid of the IncQ family with a broad host spectrum (Wang et al., 2019). The *sul3* gene has also
580 been found in many sources together with class 1 integrons (Lima et al., 2020). Our present findings
581 indicate the *sul* gene family to be associated with *intI3, tnpA, Tp614, orf39-IS26, Tn5, IncN_rep, IS613,*
582 *trfA, ISAba3, IS1111* in both processes; however, *intI1* was not noted during our study

In addition, *qnrA* and *qnrB* resistance genes were not identified in the present cattle manure
samples. Similarly, Mu et al. (2015) were also not able to detect the sought quinolone-resistance genes
(*qnrD* and *qnrS*). However, Wang et al. (2019) detected the presence of *qnrB* and *qnrS* genes in raw
cattle manure, but did not observe any reduction after composting. Pitta et al. (2016) confirm the
presence of *vanRA, vanRE*, and *vanRG* resistance genes in raw manure. While *vanRA* was found
together with *vanB, vanC, vanC2/vanC3, vanHB, vanRB, vanTC, vanTE, vanTG, vanYB* and *vanYD* in
the present study, *vanRE* and *vanRG* were absent. Finally, Zhou and Yao (2020) did not find any
change in total vancomycin-resistance gene levels after composting, which contradicts our present
results indicating a decrease in abundance.

Although significant reductions in ARG abundances grouped according to antibiotic type (Fig.
2) were noted after composting and storage, the effect of those treatment strategies on particular genes
is more complex, and many gaps still need to be addressed. The changes in ARG abundance during
storage or composting may vary depending on the location of the long-term storage pit or composting
pile, storage/composting time (age), manure removal techniques (scrap versus flush removal) or even
season of storage (Hurst et al., 2019). In addition, during composting, ARG and MGE removal can
also be influenced by temperature, pH, moisture, or type of waste and bulking material (Pezzolla et al.,
2021). In addition, ARG levels can also depend on the co-occurrence of ARG or MGE, and on various
factors affecting bacterial fitness, gene regulation and transfer, such as nutrient accessibility or heavy
metal concentration (Oliver et al., 2020). While ARG concentration also seems to be dependent on on-
site pressure, such as the presence of antibiotics or heavy metals, it also appears to be influenced by
the specific gene. Zhang et al. (2018) found lower levels of macrolide-resistance genes (*ermX, ermQ,*
ermF, ermT, and ermB) to be associated with lower tylosin concentrations, due to the effect of tylosin
degradation during composting. Guo et al. (2019) reported that adding cyromazine (a non-antibiotic
drug) to samples increases the prevalence of *blaCTX-M, blaVIM*, and *Tn916/1545* genes.

It should not be neglected that microorganisms themselves may generate selection pressure
regarding antibiotic resistance (Allen et al., 2010). Bacteria from the *Actinobacteria* produce diverse
secondary metabolites, including antibiotics, to promote their survival (Heul et al., 2018). Zhang et al.
(2016) observed a strong relationship between the availability of zinc and the presence of *tetQ, tetG,*
ermB, dfrA, and *intI1*. Similarly, the presence of bioavailable copper corresponded to the presence of
sul1, ermX ermB, dfrA7 and *aac(6')-ib-cr*; however, it should be noted that the study was performed
in a wastewater treatment plant.

In addition, the abundance of ARG appears to be contingent on microbial community structure
and its shift during processing. While composting is often considered more reliable than storage for
reducing pathogen levels in manure, it is also significantly more cost-, time- and labor-consuming
(Staley et al., 2021). Zhang et al. (2018) attribute the changes in ARG levels observed during treatment
to the dynamics of the potential host bacteria, with the mobility and type of gene influencing the host
range.

The increase in ARG noted during composting may occur due to the genes being harbored by
ARB during proliferation, or HGT between various ARB (Gou et al., 2018). Indeed, genes conferring
resistance to chloramphenicol and lincomycin have been found to undergo HGT from gram-positive
Actinobacteria to gram-negative *Proteobacteria*, despite the clear phylogenetic and ecological
boundaries between them (Jiang et al., 2017). HGT is promoted during composting due to it favoring
unfavorable conditions for bacterial growth which induce stress-response mechanisms (Lima et al.,
2020).

The differences between the ARG levels detected during composting or storage may be related
to the nature of bacterial succession and the survival of ARG-carrying bacteria during the thermophilic
phase. During storage, the bacterial strains harboring ARGs were initially suppressed, with their
number increasing later on; however, no differences in sample biodiversity were found between
treatment strategies. The scale of the study also matters. While lab-scale reactors are simple to prepare
and they have clear value in preliminary testing, they nevertheless cannot adequately simulate full-
scale processes, and their findings may not directly apply to field-scale studies (Staley et al., 2021).

The detected ARGs could confer resistance to all major antimicrobial classes, such as
aminoglycosides, beta-lactams, MLSB, tetracyclines, and vancomycin; all are critical antibiotic
groups, with some being even categorized as 'last-resort', life-saving antimicrobials. It is important to
remember that while the proposed manure management strategies reduce the risk of ARG spread, they
do not eliminate it entirely, and even if not apparent in treated samples, ARGs may persist in the
environment below the detection limit. Even trace amounts of genes may further spread with host
species succession or be disseminated via HGT.

641   5   Conclusions

Our findings prove that dairy cattle farms can serve as reservoirs of ARG and MGE. However,
both manure storage and composting, were able to reduce aminoglycoside-, beta-lactam-, tetracycline-
, MLSB- and vancomycin-resistance genes as well as MDR genes, MGE, and various "other" genes;
however they had less or no effect on phenicol-, sulfonamide or trimethoprim-resistance genes.

Composting resulted in greater richness and diversity of the bacterial community compared to
storage. In addition, while both proposed manure management strategies influence ARG and MGE
abundance, particular ARGs respond differently to treatment in highly gene-specific ways, and also
appear to depend on the dynamics of the bacterial community harboring ARG. It is difficult to compare
the effect of manure treatment strategies due to the range of possible influences. While both strategies
reduce ARG and MGE levels, composting provides fertilizer with higher biodiversity, and is devoid of
pathogens and parasite eggs, but is more labor- and time-consuming than storage.

There is a need to improve the strategies for manure management that offer more effective ARG
and MGE removal. This could be achieved by unifying the procedure (including process conditions
and additives), optimizing current methods, or by adopting new processes designed specifically to
remove or reduce ARG from animal feces, and not only to manage waste.

657   6   Conflict of Interest

658 *The authors declare that the research was conducted in the absence of any commercial or financial*
659 *relationships that could be construed as a potential conflict of interest.*

660   7   Author Contributions

661 M.Z. - Methodology, Formal analysis, Investigation, Data Curation, Writing - Original Draft,
662 Visualization; A.B. - Formal analysis, Investigation, Writing - Original Draft; M.S. – Investigation,
663 Writing - Original Draft; K.C. - Formal analysis, Data Curation; M.P. - Conceptualization, Writing -
664 Review & Editing, Project administration, Funding acquisition.

665   8   Funding

This work was supported by National Science Centre, Poland (2017/25/Z/NZ7/03026), grant under the European Horizon 2020, in the frame of the JPI-EC-AMR Joint Transnational Call (JPIAMR), JPI-EC-AMR JTC 2017, project INART–"Intervention of antibiotic resistance transfer into the food chain" to MP and partially in the frame of the "Excellence Initiative—Research University (2020–2026)" Program at the University of Warsaw.

9   Acknowledgments

This is a short text to acknowledge the contributions of specific colleagues, institutions, or agencies that aided the efforts of the authors.

10   Reference

Ahmed, W., Gyawali, P., Hamilton, K. A., Joshi, S., Aster, D., Donner, E., et al. (2021). Antibiotic Resistance and Sewage-Associated Marker Genes in Untreated Sewage and a River Characterized During Baseflow and Stormflow. *Frontiers in Microbiology* 12. Available at: https://www.frontiersin.org/articles/10.3389/fmicb.2021.632850 [Accessed November 14, 2023].

Allen, H. K., Donato, J., Wang, H. H., Cloud-Hansen, K. A., Davies, J., and Handelsman, J. (2010). Call of the wild: antibiotic resistance genes in natural environments. *Nat Rev Microbiol* 8, 251–259. doi: 10.1038/nrmicro2312.

Amarakoon, I. D., Zvomuya, F., Sura, S., Larney, F. J., Cessna, A. J., Xu, S., et al. (2016). Dissipation of Antimicrobials in Feedlot Manure Compost after Oral Administration versus Fortification after Excretion. *J Environ Qual* 45, 503–510. doi: 10.2134/jeq2015.07.0408.

Barlow, J. (2011). Mastitis therapy and antimicrobial susceptibility: a multispecies review with a focus on antibiotic treatment of mastitis in dairy cattle. *J Mammary Gland Biol Neoplasia* 16, 383–407. doi: 10.1007/s10911-011-9235-z.

Berendsen, B. J. A., Lahr, J., Nibbeling, C., Jansen, L. J. M., Bongers, I. E. A., Wipfler, E. L., et al. (2018). The persistence of a broad range of antibiotics during calve, pig and broiler manure storage. *Chemosphere* 204, 267–276. doi: 10.1016/j.chemosphere.2018.04.042.

Beyi, A. F., Brito-Goulart, D., Hawbecker, T., Slagel, C., Ruddell, B., Hassall, A., et al. (2021). Danofloxacin Treatment Alters the Diversity and Resistome Profile of Gut Microbiota in Calves. *Microorganisms* 9, 2023. doi: 10.3390/microorganisms9102023.

Bolyen, E., Rideout, J. R., Dillon, M. R., Bokulich, N. A., Abnet, C. C., Al-Ghalith, G. A., et al. (2019). Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. *Nat Biotechnol* 37, 852–857. doi: 10.1038/s41587-019-0209-9.

Brzóska, F., Kowalski, Z. M., Osięgłowski, S., and Strzetelski, J. (2014). *IZ PIB-INRA zalecenia żywieniowe dla przeżuwaczy i tabele wartości pokarmowej pasz*. Fundacja Instytutu Zootechniki Państwowego Instytutu Badawczego Patronus Animalium Available at: https://integro.bs.katowice.pl/32503383763/ksiazka/iz-pib-inra-zalecenia-zywieniowe-dla-przezuwaczy-i-tabele-wartosci-pokarmowej-pasz [Accessed October 21, 2022].

Centner, T. J. (2016). Recent government regulations in the United States seek to ensure the
effectiveness of antibiotics by limiting their agricultural use. *Environment International* 94,
1–7. doi: 10.1016/j.envint.2016.04.018.

Chen, Q., An, X., Li, H., Su, J., Ma, Y., and Zhu, Y.-G. (2016). Long-term field application of
sewage sludge increases the abundance of antibiotic resistance genes in soil. *Environment*
International 92–93, 1–10. doi: 10.1016/j.envint.2016.03.026.

Chong, J., Liu, P., Zhou, G., and Xia, J. (2020). Using MicrobiomeAnalyst for comprehensive
statistical, functional, and meta-analysis of microbiome data. *Nat Protoc* 15, 799–821. doi:
10.1038/s41596-019-0264-1.

Chopra, I., and Roberts, M. (2001). Tetracycline Antibiotics: Mode of Action, Applications,
Molecular Biology, and Epidemiology of Bacterial Resistance. *Microbiol. Mol. Biol. Rev.* 65,
232–260. doi: 10.1128/MMBR.65.2.232-260.2001.

Diana, A., Santinello, M., Penasa, M., Scali, F., Magni, E., Alborali, G. L., et al. (2020). Use of
antimicrobials in beef cattle: an observational study in the north of Italy. *Prev Vet Med* 181,
105032. doi: 10.1016/j.prevetmed.2020.105032.

Dibner, J. J., and Richards, J. D. (2005). Antibiotic growth promoters in agriculture: history and
mode of action. *Poult. Sci.* 84, 634–643. doi: 10.1093/ps/84.4.634.

Dolliver, H., Kumar, K., and Gupta, S. (2007). Sulfamethazine uptake by plants from manure-
amended soil. *J. Environ. Qual.* 36, 1224–1230. doi: 10.2134/jeq2006.0266.

Edrington, T. S., Fox, W. E., Callaway, T. R., Anderson, R. C., Hoffman, D. W., and Nisbet, D. J.
(2009). Pathogen Prevalence and Influence of Composted Dairy Manure Application on
Antimicrobial Resistance Profiles of Commensal Soil Bacteria. *Foodborne Pathogens and*
Disease 6, 217–224. doi: 10.1089/fpd.2008.0184.

Epelde, L., Jauregi, L., Urra, J., Ibarretxe, L., Romo, J., Goikoetxea, I., et al. (2018). Characterization
of Composted Organic Amendments for Agricultural Use. *Frontiers in Sustainable Food*
Systems 2. Available at: https://www.frontiersin.org/articles/10.3389/fsufs.2018.00044
[Accessed April 19, 2023].

Ezugworie, F. N., Igbokwe, V. C., and Onwosi, C. O. (2021). Proliferation of antibiotic-resistant
microorganisms and associated genes during composting: An overview of the potential
impacts on public health, management and future. *Science of The Total Environment* 784,
147191. doi: 10.1016/j.scitotenv.2021.147191.

FAO (2020). FAOSTAT. Available at: https://www.fao.org/faostat/en/#home [Accessed March 23,
2022].

Faul, F., Erdfelder, E., Buchner, A., and Lang, A.-G. (2009). Statistical power analyses using
G*Power 3.1: Tests for correlation and regression analyses. *Behavior Research Methods* 41,
1149–1160. doi: 10.3758/BRM.41.4.1149.

Faul, F., Erdfelder, E., Lang, A.-G., and Buchner, A. (2007). G*Power 3: A flexible statistical power
analysis program for the social, behavioral, and biomedical sciences. *Behavior Research*
Methods 39, 175–191. doi: 10.3758/BF03193146.

Ferroni, L., Lovito, C., Scoccia, E., Dalmonte, G., Sargenti, M., Pezzotti, G., et al. (2020). Antibiotic
Consumption on Dairy and Beef Cattle Farms of Central Italy Based on Paper Registers.
Antibiotics 9, 273. doi: 10.3390/antibiotics9050273.

Filippitzi, M. E., Devreese, M., Broekaert, K., Rasschaert, G., Daeseleire, E., Meirlaen, J., et al.
(2019). Quantitative risk model to estimate the level of antimicrobial residues that can be
transferred to soil via manure, due to oral treatments of pigs. *Preventive Veterinary Medicine*
167, 90–100. doi: 10.1016/j.prevetmed.2019.03.022.

Foged, H., Flotats Ripoll, X., Bonmatí Blasi, A., Palatsi Civit, J., Magrí Aloy, A., and Schelde, K. M.
(2012). Inventory of manure processing activities in Europe. Available at:
https://upcommons.upc.edu/handle/2117/18943 [Accessed May 9, 2023].

Font-Palma, C. (2019). Methods for the Treatment of Cattle Manure—A Review. *C* 5, 27. doi:
10.3390/c5020027.

Ghosh, S., and LaPara, T. M. (2007). The effects of subtherapeutic antibiotic use in farm animals on
the proliferation and persistence of antibiotic resistance among soil bacteria. *The ISME*
Journal 1, 191–203. doi: 10.1038/ismej.2007.31.

Girotto, F., and Cossu, R. (2017). "Animal Waste: Opportunities and Challenges," in *Sustainable*
Agriculture Reviews Sustainable Agriculture Reviews., ed. E. Lichtfouse (Cham: Springer
International Publishing), 1–13. doi: 10.1007/978-3-319-48006-0_1.

Gou, M., Hu, H.-W., Zhang, Y.-J., Wang, J.-T., Hayden, H., Tang, Y.-Q., et al. (2018). Aerobic
composting reduces antibiotic resistance genes in cattle manure and the resistome
dissemination in agricultural soils. *Science of The Total Environment* 612, 1300–1310. doi:
10.1016/j.scitotenv.2017.09.028.

Gullberg, E., Cao, S., Berg, O. G., Ilbäck, C., Sandegren, L., Hughes, D., et al. (2011). Selection of
Resistant Bacteria at Very Low Antibiotic Concentrations. *PLOS Pathogens* 7, e1002158.
doi: 10.1371/journal.ppat.1002158.

Guo, H., Gu, J., Wang, X., Yu, J., Nasir, M., Peng, H., et al. (2019). Responses of antibiotic and
heavy metal resistance genes to bamboo charcoal and bamboo vinegar during aerobic
composting. *Environmental Pollution* 252, 1097–1105. doi: 10.1016/j.envpol.2019.05.014.

Heinonen-Tanski, H., Mohaibes, M., Karinen, P., and Koivunen, J. (2006). Methods to reduce
pathogen microorganisms in manure. *Livestock Science* 102, 248–255. doi:
10.1016/j.livsci.2006.03.024.

Heuer, H., Schmitt, H., and Smalla, K. (2011). Antibiotic resistance gene spread due to manure
application on agricultural fields. *Curr. Opin. Microbiol.* 14, 236–243. doi:
10.1016/j.mib.2011.04.009.

Heul, H. U. van der, L. Bilyk, B., J. McDowall, K., F. Seipke, R., and Wezel, G. P. van (2018).
Regulation of antibiotic production in Actinobacteria: new perspectives from the post-
genomic era. *Natural Product Reports* 35, 575–604. doi: 10.1039/C8NP00012C.

Hu, Y. J., and Cowling, B. J. (2020). Reducing antibiotic use in livestock, China. *Bull. World Health*
Organ. 98, 360–361. doi: 10.2471/BLT.19.243501.

Hurst, J. J., Oliver, J. P., Schueler, J., Gooch, C., Lansing, S., Crossette, E., et al. (2019). Trends in
Antimicrobial Resistance Genes in Manure Blend Pits and Long-Term Storage Across Dairy
Farms with Comparisons to Antimicrobial Usage and Residual Concentrations. *Environ. Sci.*
Technol. 53, 2405–2415. doi: 10.1021/acs.est.8b05702.

Huygens, J., Daeseleire, E., Mahillon, J., Van Elst, D., Decrop, J., Meirlaen, J., et al. (2021).
Presence of Antibiotic Residues and Antibiotic Resistant Bacteria in Cattle Manure Intended
for Fertilization of Agricultural Fields: A One Health Perspective. *Antibiotics (Basel)* 10, 410.
doi: 10.3390/antibiotics10040410.

Jauregi, L., Epelde, L., González, A., Lavín, J. L., and Garbisu, C. (2021). Reduction of the resistome
risk from cow slurry and manure microbiomes to soil and vegetable microbiomes. *Environ*
Microbiol 23, 7643–7660. doi: 10.1111/1462-2920.15842.

Jiang, X., Ellabaan, M. M. H., Charusanti, P., Munck, C., Blin, K., Tong, Y., et al. (2017).
Dissemination of antibiotic resistance genes from antibiotic producers to pathogens. *Nat*
Commun 8, 15784. doi: 10.1038/ncomms15784.

Kakimoto, T., Osawa, T., and Funamizu, N. (2007). Antibiotic effect of amoxicillin on the feces
composting process and reactivation of bacteria by intermittent feeding of feces. *Bioresource*
Technology 98, 3555–3560. doi: 10.1016/j.biortech.2006.11.029.

Keenum, I., Williams, R. K., Ray, P., Garner, E. D., Knowlton, K. F., and Pruden, A. (2021).
Combined effects of composting and antibiotic administration on cattle manure–borne
antibiotic resistance genes. *Microbiome* 9, 81. doi: 10.1186/s40168-021-01006-z.

Köninger, J., Lugato, E., Panagos, P., Kochupillai, M., Orgiazzi, A., and Briones, M. J. I. (2021).
Manure management and soil biodiversity: Towards more sustainable food systems in the
EU. *Agricultural Systems* 194, 103251. doi: 10.1016/j.agsy.2021.103251.

Lima, T., Domingues, S., and Da Silva, G. J. (2020). Manure as a Potential Hotspot for Antibiotic
Resistance Dissemination by Horizontal Gene Transfer Events. *Veterinary Sciences* 7, 110.
doi: 10.3390/vetsci7030110.

Loyon, L. (2018). Overview of Animal Manure Management for Beef, Pig, and Poultry Farms in
France. *Frontiers in Sustainable Food Systems* 2, 36. doi: 10.3389/fsufs.2018.00036.

Mao, S., Zhang, M., Liu, J., and Zhu, W. (2015). Characterising the bacterial microbiota across the
gastrointestinal tracts of dairy cattle: membership and potential function. *Sci Rep* 5, 16116.
doi: 10.1038/srep16116.

Marti, R., Scott, A., Tien, Y.-C., Murray, R., Sabourin, L., Zhang, Y., et al. (2013). Impact of Manure
Fertilization on the Abundance of Antibiotic-Resistant Bacteria and Frequency of Detection

of Antibiotic Resistance Genes in Soil and on Vegetables at Harvest. *Appl Environ Microbiol*
79, 5701–5709. doi: 10.1128/AEM.01682-13.

Massé, D. I., Saady, N. M. C., and Gilbert, Y. (2014). Potential of Biological Processes to Eliminate
Antibiotics in Livestock Manure: An Overview. *Animals* 4, 146–163. doi:
10.3390/ani4020146.

Mc Carthy, G., Lawlor, P. G., Coffey, L., Nolan, T., Gutierrez, M., and Gardiner, G. E. (2011). An
assessment of pathogen removal during composting of the separated solid fraction of pig
manure. *Bioresour Technol* 102, 9059–9067. doi: 10.1016/j.biortech.2011.07.021.

McEachran, A. D., Blackwell, B. R., Hanson, J. D., Wooten, K. J., Mayer, G. D., Cox, S. B., et al.
(2015). Antibiotics, bacteria, and antibiotic resistance genes: aerial transport from cattle feed
yards via particulate matter. *Environ. Health Perspect.* 123, 337–343. doi:
10.1289/ehp.1408555.

Merle, R., Hajek, P., Käsbohrer, A., Hegger-Gravenhorst, C., Mollenhauer, Y., Robanus, M., et al.
(2012). Monitoring of antibiotic consumption in livestock: a German feasibility study. *Prev*
Vet Med 104, 34–43. doi: 10.1016/j.prevetmed.2011.10.013.

Mu, Q., Li, J., Sun, Y., Mao, D., Wang, Q., and Luo, Y. (2015). Occurrence of sulfonamide-,
tetracycline-, plasmid-mediated quinolone- and macrolide-resistance genes in livestock
feedlots in Northern China. *Environ Sci Pollut Res* 22, 6932–6940. doi: 10.1007/s11356-014-
3905-5.

Nguyen, B.-A. T., Chen, Q.-L., He, J.-Z., and Hu, H.-W. (2020). Microbial regulation of natural
antibiotic resistance: Understanding the protist-bacteria interactions for evolution of soil
resistome. *Science of The Total Environment* 705, 135882. doi:
10.1016/j.scitotenv.2019.135882.

Oliver, J. P., Gooch, C. A., Lansing, S., Schueler, J., Hurst, J. J., Sassoubre, L., et al. (2020). Invited
review: Fate of antibiotic residues, antibiotic-resistant bacteria, and antibiotic resistance genes
in US dairy manure management systems. *Journal of Dairy Science* 103, 1051–1071. doi:
10.3168/jds.2019-16778.

Olmos Antillón, G., Sjöström, K., Fall, N., Sternberg Lewerin, S., and Emanuelson, U. (2020).
Antibiotic Use in Organic and Non-organic Swedish Dairy Farms: A Comparison of Three
Recording Methods. *Frontiers in Veterinary Science* 7. Available at:
https://www.frontiersin.org/articles/10.3389/fvets.2020.568881 [Accessed April 26, 2023].

Pei, R., Cha, J., Carlson, K. H., and Pruden, A. (2007). Response of Antibiotic Resistance Genes
(ARG) to Biological Treatment in Dairy Lagoon Water. *Environ. Sci. Technol.* 41, 5108–
5113. doi: 10.1021/es070051x.

Pezzolla, D., Cucina, M., Proietti, P., Calisti, R., Regni, L., and Gigliotti, G. (2021). The Use of New
Parameters to Optimize the Composting Process of Different Organic Wastes. *Agronomy* 11,
2090. doi: 10.3390/agronomy11102090.

Pitta, D. W., Dou, Z., Kumar, S., Indugu, N., Toth, J. D., Vecchiarelli, B., et al. (2016). Metagenomic
Evidence of the Prevalence and Distribution Patterns of Antimicrobial Resistance Genes in

Dairy Agroecosystems. *Foodborne Pathogens and Disease* 13, 296–302. doi:
10.1089/fpd.2015.2092.

Qian, X., Sun, W., Gu, J., Wang, X.-J., Sun, J.-J., Yin, Y.-N., et al. (2016). Variable effects of
oxytetracycline on antibiotic resistance gene abundance and the bacterial community during
aerobic composting of cow manure. *Journal of Hazardous Materials* 315, 61–69. doi:
10.1016/j.jhazmat.2016.05.002.

Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., et al. (2013). The SILVA
ribosomal RNA gene database project: improved data processing and web-based tools.
Nucleic Acids Research 41, D590–D596. doi: 10.1093/nar/gks1219.

Ruuskanen, M., Muurinen, J., Meierjohan, A., Pärnänen, K., Tamminen, M., Lyra, C., et al. (2016).
Fertilizing with Animal Manure Disseminates Antibiotic Resistance Genes to the Farm
Environment. *Journal of Environment Quality* 45, 488. doi: 10.2134/jeq2015.05.0250.

Santamaría, J., López, L., and Soto, C. (2011). Detection and Diversity evaluation of tetracycline
resistance genes in grassland-based production systems in Colombia, South America.
Frontiers in Microbiology 2. Available at:
https://www.frontiersin.org/articles/10.3389/fmicb.2011.00252 [Accessed April 25, 2023].

Sarmah, A. K., Meyer, M. T., and Boxall, A. B. A. (2006). A global perspective on the use, sales,
exposure pathways, occurrence, fate and effects of veterinary antibiotics (VAs) in the
environment. *Chemosphere* 65, 725–759. doi: 10.1016/j.chemosphere.2006.03.026.

Shannon, P., Markiel, A., Ozier, O., Baliga, N. S., Wang, J. T., Ramage, D., et al. (2003). Cytoscape:
a software environment for integrated models of biomolecular interaction networks. *Genome*
Res 13, 2498–2504. doi: 10.1101/gr.1239303.

Spielmeyer, A. (2018). Occurrence and fate of antibiotics in manure during manure treatments: A
short review. *Sustainable Chemistry and Pharmacy* 9, 76–86. doi: 10.1016/j.scp.2018.06.004.

Staley, Z. R., Woodbury, B. L., Stromer, B. S., Schmidt, A. M., Snow, D. D., Bartelt-Hunt, S. L., et
al. (2021). Stockpiling versus Composting: Effectiveness in Reducing Antibiotic-Resistant
Bacteria and Resistance Genes in Beef Cattle Manure. *Applied and Environmental*
Microbiology 87, e00750-21. doi: 10.1128/AEM.00750-21.

Storteboom, H. N., Kim, S.-C., Doesken, K. C., Carlson, K. H., Davis, J. G., and Pruden, A. (2007).
Response of Antibiotics and Resistance Genes to High-Intensity and Low-Intensity Manure
Management. *Journal of Environmental Quality* 36, 1695–1703. doi: 10.2134/jeq2007.0006.

Tasho, R. P., and Cho, J. Y. (2016). Veterinary antibiotics in animal waste, its distribution in soil and
uptake by plants: A review. *Science of The Total Environment* 563–564, 366–376. doi:
10.1016/j.scitotenv.2016.04.140.

Thomas, M., Webb, M., Ghimire, S., Blair, A., Olson, K., Fenske, G. J., et al. (2017). Metagenomic
characterization of the effect of feed additives on the gut microbiome and antibiotic resistome
of feedlot cattle. *Sci Rep* 7, 12257. doi: 10.1038/s41598-017-12481-6.

Udikovic-Kolic, N., Wichmann, F., Broderick, N. A., and Handelsman, J. (2014). Bloom of resident antibiotic-resistant bacteria in soil following manure fertilization. *Proceedings of the National Academy of Sciences* 111, 15202–15207. doi: 10.1073/pnas.1409836111.

Wang, F.-H., Qiao, M., Chen, Z., Su, J.-Q., and Zhu, Y.-G. (2015). Antibiotic resistance genes in manure-amended soil and vegetables at harvest. *J. Hazard. Mater.* 299, 215–221. doi: 10.1016/j.jhazmat.2015.05.028.

Wang, F.-H., Qiao, M., Su, J.-Q., Chen, Z., Zhou, X., and Zhu, Y.-G. (2014). High throughput profiling of antibiotic resistance genes in urban park soils with reclaimed water irrigation. *Environ Sci Technol* 48, 9079–9085. doi: 10.1021/es502615e.

Wang, L., Zhao, X., Wang, J., Wang, J., Zhu, L., and Ge, W. (2019). Macrolide- and quinolone-resistant bacteria and resistance genes as indicators of antibiotic resistance gene contamination in farmland soil with manure application. *Ecological Indicators* 106, 105456. doi: 10.1016/j.ecolind.2019.105456.

Wang, S., and Zeng, Y. (2018). Ammonia emission mitigation in food waste composting: A review. *Bioresource Technology* 248, 13–19. doi: 10.1016/j.biortech.2017.07.050.

Xu, S., Amarakoon, I. D., Zaheer, R., Smith, A., Sura, S., Wang, G., et al. (2018). Dissipation of antimicrobial resistance genes in compost originating from cattle manure after direct oral administration or post-excretion fortification of antimicrobials. *Journal of Environmental Science and Health, Part A* 53, 373–384. doi: 10.1080/10934529.2017.1404337.

Yilmaz, P., Parfrey, L. W., Yarza, P., Gerken, J., Pruesse, E., Quast, C., et al. (2014). The SILVA and "All-species Living Tree Project (LTP)" taxonomic frameworks. *Nucleic Acids Research* 42, D643–D648. doi: 10.1093/nar/gkt1209.

Youngquist, C. P., Mitchell, S. M., and Cogger, C. G. (2016). Fate of Antibiotics and Antibiotic Resistance during Digestion and Composting: A Review. *Journal of Environmental Quality* 45, 537–545. doi: 10.2134/jeq2015.05.0256.

Zalewska, M., Błażejewska, A., Czapko, A., and Popowska, M. (2021). Antibiotics and Antibiotic Resistance Genes in Animal Manure – Consequences of Its Application in Agriculture. *Frontiers in Microbiology* 12. Available at: https://www.frontiersin.org/article/10.3389/fmicb.2021.610656 [Accessed July 1, 2022].

Zalewska, M., Błażejewska, A., Czapko, A., and Popowska, M. (2023). Pig manure treatment strategies for mitigating the spread of antibiotic resistance. *Sci Rep* 13, 11999. doi: 10.1038/s41598-023-39204-4.

Zalewska, M., Kawecka-Grochocka, E., Słoniewska, D., Kościuczuk, E., Marczak, S., Jarmuż, W., et al. (2020). Acute phase protein expressions in secretory and cistern lining epithelium tissues of the dairy cattle mammary gland during chronic mastitis caused by staphylococci. *BMC Veterinary Research* 16, 320. doi: 10.1186/s12917-020-02544-8.

Zalewska, M., and Popowska, M. (2020). "Antimicrobial/Antibiotic Resistance Genes Due to Manure and Agricultural Waste Applications," in *Antibiotics and Antimicrobial Resistance Genes: Environmental Occurrence and Treatment Technologies* Emerging Contaminants and

Associated Treatment Technologies., ed. M. Z. Hashmi (Cham: Springer International
Publishing), 139–161. doi: 10.1007/978-3-030-40422-2_6.

Zhang, L., Li, L., Sha, G., Liu, C., Wang, Z., and Wang, L. (2020). Aerobic composting as an
effective cow manure management strategy for reducing the dissemination of antibiotic
resistance genes: An integrated meta-omics study. *Journal of Hazardous Materials* 386,
121895. doi: 10.1016/j.jhazmat.2019.121895.

Zhang, L., Zhang, H., Wang, Z., Chen, G., and Wang, L. (2016). Dynamic changes of the dominant
functioning microbial community in the compost of a 90-m3 aerobic solid state fermentor
revealed by integrated meta-omics. *Bioresource Technology* 203, 1–10. doi:
10.1016/j.biortech.2015.12.040.

Zhang, Y., Chen, M., Guo, J., Liu, N., Yi, W., Yuan, Z., et al. (2022). Study on dynamic changes of
microbial community and lignocellulose transformation mechanism during green waste
composting. *Engineering in Life Sciences* 22, 376–390. doi: 10.1002/elsc.202100102.

Zhang, Y., Gu, A. Z., Cen, T., Li, X., He, M., Li, D., et al. (2018). Sub-inhibitory concentrations of
heavy metals facilitate the horizontal transfer of plasmid-mediated antibiotic resistance genes
in water environment. *Environ. Pollut.* 237, 74–82. doi: 10.1016/j.envpol.2018.01.032.

Zhang, Y.-J., Hu, H.-W., Chen, Q.-L., Singh, B. K., Yan, H., Chen, D., et al. (2019). Transfer of
antibiotic resistance from manure-amended soils to vegetable microbiomes. *Environ Int* 130,
104912. doi: 10.1016/j.envint.2019.104912.

Zhou, Z., and Yao, H. (2020). Effects of Composting Different Types of Organic Fertilizer on the
Microbial Community Structure and Antibiotic Resistance Genes. *Microorganisms* 8, 268.
doi: 10.3390/microorganisms8020268.

Zhu, B., Chen, Q., Chen, S., and Zhu, Y.-G. (2017). Does organically produced lettuce harbor higher
abundance of antibiotic resistance genes than conventionally produced? *Environment*
International 98, 152–159. doi: 10.1016/j.envint.2016.11.001.

11   Supplementary Material

Table S1: Humidity [%], pH, and temperature [°C] changes during manure treatments

Table S2. Mean abundance with standard deviation of each phylum across the treatment

Table S3. The results of pairwise PERMANOVA analysis. The multi-testing adjustment is based on
Benjamini-Hochberg procedure procedure (FDR)

Figure S1. Core microbiome across analyzed samples

Figure S2. Shannon indexes of microbial community diversity (p-value 0.08; Kruskal–Wallis
statistic)

Figure S3. Chao1 indexes of microbial community richness (p-value 0.025; Kruskal–Wallis statistic)

Figure S4. The PCoA of taxon abundance in all sample groups, calculated based on the Bray–Curtis
distance; [PERMANOVA] F-value: 7.1007; R-squared: 0.66983; p-value: 0.003

Figure S5. Comparison between resistomes of untreated cattle manure and resistomes at the end of
composting and storage

12   Data Availability Statement

969 The datasets generated for this study can be found in the Sequence Read Archive (SRA) repository
970 (NCBI) - BioProject ID: PRJNA1007935.