Comparison of the mechanism of antimicrobial action of the gold(I) compound

auranofin in Gram-positive and Gram-negative bacteria

Laísa Quadros Barsé[1*], Agnes Ulfig[1,5*], Marharyta Varatnitskaya[1], Melissa Vázquez-

Hernández[3], Jihyun Yoo[1], Astrid M. Imann[1,2], Natalie Lupilov[1], Marina Fischer[4], Katja

Becker[4], Julia E. Bandow[3], and Lars I. Leichert[1†]

[1] Ruhr University Bochum, Medical Faculty, Institute of Biochemistry and Pathobiochemistry
– Microbial Biochemistry, Universitätsstrasse 150, 44780 Bochum, Germany

[2] Westphalian University of Applied Sciences, Institute of Electrical Engineering and Applied
Sciences – Molecular Biology, August-Schmidt-Ring 10, 45665 Recklinghausen, Germany

[3] Ruhr University Bochum, Faculty of Biology and Biotechnology, Applied Microbiology,
Universitätsstrasse 150, 44780 Bochum, Germany

[4] Justus Liebig University Giessen, Chair of Biochemistry and Molecular Biology,
Interdisciplinary Research Center, Heinrich-Buff-Ring 26-32, 35392 Giessen, Germany

[5] Current address: University of Iceland, Faculty of Medicine, Louma G. Laboratory of
Epigenetic Research, Sturlugata 8, 101 Reykjavik, Iceland

[*] Shared first authorship.

[†]Correspondence: Lars I. Leichert
18   Ruhr University Bochum
19   Institute of Biochemistry and Pathobiochemistry
20   Microbial Biochemistry
21   Universitätsstrasse 150
22   44780 Bochum, Germany
23   Phone: +49 234 3224585
24   Fax: +49 234 3214332
25   E-mail: lars.leichert@ruhr-uni-bochum.de

27 **KEYWORDS:** auranofin, gold, drug repurposing, thioredoxin reductase, proteomics,
28 *Escherichia coli*, *Bacillus subtilis*

ABSTRACT

The antirheumatic gold(I) compound auranofin has been suggested to exhibit bactericidal activity by disrupting the thiol homeostasis through direct inhibition of bacterial thioredoxin reductase (TrxB). While highly effective at killing Gram-positive bacteria, it lacks significant activity against Gram-negative species for reasons that largely remain unclear. Here, we aimed to elucidate the molecular mechanisms underlying the low susceptibility of the Gram-negative model organism *Escherichia coli* to auranofin when compared to the Gram-positive model organism *Bacillus subtilis*. A change in the proteome of *E. coli* exposed to auranofin suggested that the effect of this gold compound is a combination of inactivation of thiol-containing enzymes, upregulation of proteins involved in basal metabolism, and the consequent induction of systemic oxidative stress. Susceptibility tests in *E. coli* mutants lacking the proteins upregulated upon auranofin treatment suggested that none of them are directly involved in *E. coli*'s high tolerance to auranofin. To elucidate factors that could make Gram-negative bacteria less susceptible to auranofin, we tested *E. coli* cells lacking the efflux pump component TolC. These cells were more sensitive to auranofin treatment than the wild type, but not to an extent that would fully explain the observed difference in susceptibility of Gram-positive and Gram-negative organisms. We thus tested if *E. coli*'s thioredoxin reductase (TrxB) is inherently less sensitive to auranofin than TrxB from *B. subtilis*, which was not the case. *E. coli* cells lacking components of the thioredoxin-system were also only marginally more susceptible to auranofin. However, *E. coli* strains lacking the low molecular weight thiol glutathione, but not glutathione reductase, showed a high susceptibility to auranofin. Bacterial cells expressing the genetically encoded redox probe roGFP2 allowed us to observe the oxidation of cellular protein thiols *in situ*. In line with their susceptibility, the kinetics of probe oxidation and the degree of oxidation promoted by auranofin is significantly higher in Gram-positive bacteria when compared to Gram-negative bacteria. Based on our findings, we hypothesize that auranofin leads to a global disturbance in the cellular thiol redox homeostasis in bacteria, but Gram-negative bacteria are inherently more resistant due to the presence of drug export systems and high cellular concentrations of glutathione.

INTRODUCTION

The evolutionary emergence and rapid spread of multidrug-resistant pathogenic bacteria is an ongoing and rising public health concern, as infections caused by these pathogens claim the lives of nearly 700,000 people across the globe each year (Humphreys and Fleck, 2016). This number is expected to increase dramatically if no urgent actions are taken. Unfortunately, the slow progress in the discovery and development of new antibiotic classes has led to a limited availability of novel antimicrobial compounds for clinical use and, thus, made the effective treatment of many bacterial infections more challenging (Hutchings et al., 2019). Accordingly, there is an urgent medical need to identify new and improved antibacterial agents with unprecedented modes of action to tackle the global antibiotic resistance crisis (Aslam et al., 2018). Since conventional *de novo* drug discovery and development are lengthy, expensive, and complex, repurposing approved drugs for antimicrobial therapy by re-evaluating their biological targets and mechanisms of action is considered a key strategy to overcome these issues (Farha and Brown, 2019). Although using known drugs for indications they were not originally developed for is far from new, it has gained considerable attention in the last decade, with about one-third of new US Food and Drug Administration (FDA) approvals being for repurposed drugs (Cha et al., 2018).

The use of metals, such as silver, gold, and platinum, and compounds based on these and other metals, in treating diseases dates to ancient times (Wang et al., 2012), but their full potential as therapeutic drugs has not been realized yet. With the discovery of the platinum-based compound cisplatin as an anticancer agent in the 20th century, metallodrugs have experienced an impressive renaissance of interest in modern medicine, leading to the development of several other metal complexes with antitumor and anti-inflammatory properties over the last decades (Alderden et al., 2006).

Auranofin, a gold(I)-containing triethyl phosphine, was initially approved by the FDA in 1985 as an anti-inflammatory drug for the treatment of rheumatoid arthritis and was the first orally active gold compound proven to slow the progression of the disease by inhibiting antibody-mediated cellular toxicity and stimulating cell-mediated immunity (Finkelstein et al., 1976; Messori and Marcon, 2004). Auranofin's primary mode of action in humans is thought to be the inhibition of thioredoxin reductase (TrxR) (Gromer et al. 1998). Auranofin's exceptionally high affinity toward TrxR is due to the reactivity of its thiol ligand with thiol groups and especially the selenol group of the selenocysteine present in eukaryotic TrxR (Angelucci et al., 2009; Jeon et al., 2003; Rackham et al., 2011; Saccoccia et al., 2012). TrxR was shown to be highly expressed in the inflamed joints of patients with rheumatoid arthritis, and it is assumed to be involved in disease activity and suppression of apoptosis in the affected

tissue (Kabuyama et al., 2008; Maurice et al., 1999). TrxR is a central component of the

antioxidant thioredoxin (Trx) system. It catalyzes the NADPH-dependent reduction of the

redox protein thioredoxin (Trx), which is crucial in maintaining the cellular environment in a

reduced state and protecting the cells against damage from oxidative stress (Holmgren and

Björnstedt, 1995; Lu and Holmgren, 2014).

More recent studies also indicate that TrxR is highly abundant in various types of human

cancer cells and, thus, represents a promising therapeutic target for anticancer drug

development. Gold-induced inactivation of mammalian TrxR by auranofin leads to severe

oxidative stress due to the overproduction of reactive oxygen species (ROS) and activation

of apoptotic signaling pathways, ultimately triggering cancer cell death (Bian et al., 2019;

Hwang-Bo et al., 2017).

Besides these effects in eukaryotes, auranofin, and related gold-based compounds have

also shown promising activity against a wide range of pathogenic microorganisms at a

clinically achievable nanomolar range. In a study by Debnath et al., auranofin was highly

efficacious against the intestinal parasite *Entamoeba histolytica,* providing the first evidence

that this gold drug could be repurposed as a novel antimicrobial agent (Debnath et al., 2012).

Indeed, more recent studies have reported remarkable bactericidal effects of auranofin

against various pathogenic Gram-positive bacteria *in vitro* and *in vivo*, including

Mycobacterium tuberculosis, certain *Bacillus* spp., *Enterococcus faecalis,* and methicillin-

resistant *Staphylococcus aureus* (MRSA) (Aguinagalde et al., 2015; Cassetta et al., 2014;

Harbut et al., 2015; Hokai et al., 2014). As a result, auranofin has become the reference

compound for developing new, structurally related organogold antimicrobial drugs with lower

toxicity, a broader spectrum of activity, and greater efficacy (Marzo et al., 2018).

Although progress has been made in understanding the underlying mechanistical aspects of

auranofin's antimicrobial activity, the precise mechanisms of its action and the respective

biological targets in bacteria are still largely unknown.

Recent studies point toward a complex multitarget mechanism of auranofin's action in

prokaryotic organisms that involves inhibition of several biosynthetic pathways, including

protein, DNA, and cell wall synthesis (Thangamani et al., 2016), and disruption of thiol-redox

homeostasis (Caroli et al., 2012; Harbut et al., 2015). Among these possible modes of action,

inhibition of thiol-containing redox enzymes essential for maintaining intracellular thiol-redox

balance is widely considered the primary mechanism through which auranofin exerts its

cytotoxic effects on bacteria (Harbut et al., 2015). A growing number of studies demonstrate

a strong inhibitory effect on bacterial thioredoxin reductase (TrxB). For instance, auranofin

has been shown to effectively inhibit TrxB of *S. aureus*, *M. tuberculosis,* and *H. pylori in vitro*,

with IC_{50} values in the low nanomolar range. Notably, the growth of these organisms was

also strongly impaired in the presence of auranofin (Harbut et al., 2015; Owings et al., 2016).

While auranofin is highly effective against Gram-positive bacteria, it is much less active

against Gram-negative bacteria, exhibiting a markedly higher minimal inhibitory

concentration (MIC) (Harbut et al., 2015). However, the reasons for the limited susceptibility

of Gram-negative bacteria to auranofin are still unknown.

In this work, we aimed to elucidate the molecular mechanisms underlying the increased

tolerance of Gram-negative bacteria to auranofin. We utilized *E. coli* and *B. subtilis* as

representative species of Gram-negative and Gram-positive bacteria, respectively, to

systematically uncover the mode of auranofin's action by testing its effect on TrxB activity

and growth of these organisms. We studied the proteomic response of *E. coli* exposed to

auranofin stress to uncover potential defense mechanisms that could mediate their increased

resistance to auranofin. We also focused on factors that differ between Gram-positive and

Gram-negative species, such as TolC, the outer membrane factor of an efflux pump in *E.*

coli. Cells lacking TolC were more susceptible to auranofin, however not to an extent that

would make TolC-dependent drug efflux the only factor explaining *E. coli*'s relative resistance

to auranofin. A systematic evaluation of mutants lacking components of the thioredoxin

system and the glutathione-dependent antioxidant system suggests that the presence of a

high concentration of the low molecular weight thiol glutathione is a major factor contributing

to *E. coli*'s tolerance. This is in line with cytoplasmic thiol oxidation caused by auranofin *in*

vivo, which we observed using *E. coli* and *B. subtilis* cells expressing the well-known redox

probe roGFP2. Overall, our data suggest that auranofin's mode of action against Gram-

negative species does not significantly differ from that against Gram-positive species.

Instead, it seems that the higher resistance of *E. coli* is caused in part by a combination of

drug efflux and, more importantly, the presence of high concentrations of the low-molecular

weight thiol glutathione, which attenuate the cytotoxic effects of auranofin.

MATERIAL AND METHODS

Determination of the Minimal Inhibitory Concentration (MIC) of Auranofin against Gram-negative and Gram-positive bacteria

Auranofin (Santa Cruz Biotechnology, Heidelberg, Germany) was tested for bactericidal activity against *Acinetobacter baumannii* DSM 30007, *Pseudomonas aeruginosa* DSM 50071, *Bacillus subtilis* 168, *Staphylococcus aureus* DSM 20231, *Staphylococcus aureus* ATCC 43300 (MRSA), *E. coli* DSM 30083, *E. coli* BW25113 and *E. coli* BW25113-derived mutant strains from the KEIO collection (Baba et al., 2006) listed in Table 1. *E. coli* DSM 30083, *A. baumannii*, *S. aureus*, and *B. subtilis* were grown in Mueller Hinton broth, and *P. aeruginosa* in cation-adjusted Mueller Hinton II. MIC assays with these strains were performed in a microdilution assay in 200 μL medium according to CLSI guidelines, as described by Albada et al. (Albada et al., 2012). *E. coli* BW25113 and the derived mutant strains from the KEIO collection were grown in reaction tubes containing 5 mL MOPS Minimal Medium (Teknova) (Neidhardt et al., 1974) supplemented with 10 μM thiamine.

Auranofin was dissolved in DMSO to a final concentration of 10 mg/mL. Serial dilutions of this stock solution were prepared in the various culture media with the Tecan Freedom Evo 75 liquid handling workstation (Tecan, Männedorf, Switzerland) for tests according to CLSI guidelines. They typically covered a range from 512 to 0.5 μg/mL. In *E. coli* BW25113 and the *E. coli* BW25113-derived mutants, a range of 256 μg/mL to 4 μg/mL was tested; these dilutions were prepared manually.

All bacterial strains were grown until the exponential growth phase was reached. At an OD_{600} of 0.4-0.5, 5×10^5 cells/mL of the exponentially growing cultures were used to inoculate the previously prepared auranofin dilutions in the same growth medium. After 16-18 hours of incubation at 37 °C, bacterial growth at various auranofin concentrations was assessed visually and photometrically by measuring the cultures' OD at 600 nm. The lowest auranofin concentration with no visually perceptible growth was recorded as MIC.

Construction of plasmids encoding bacterial TrxB

Plasmids and oligonucleotides used in this study are listed in Tables 1 and S1, respectively. Plasmid construction followed standard procedures. The correctness of all newly constructed plasmids was verified by DNA sequencing.

For the expression and subsequent purification of *E. coli* MG1655 and *B. subtilis* 168 TrxB, the respective *trxB* genes were fused to an N-terminal SUMO tag and cloned into pET22b(+)

vectors (Butt et al., 2005). The SUMO tag contains a SUMO (Small Ubiquitin-like Modifier)
protein sequence, which enhances expression and solubility of the fusion protein, a His$_6$ tag,
which allows for purification of the fusion protein using Ni–NTA chromatography, and an Ulp1
protease cleavage site (Butt et al., 2005).

The SUMO tag sequence was amplified from the pE-SUMO plasmid using the forward primer
AI003_SUMO_for carrying a NdeI restriction site and AI004_SUMO_rev or
MM004_SUMO_rev, containing an overhang complementary to the N-terminal portion of the
E. coli or *B. subtilis trxB* sequence, respectively (Table S1). In a second PCR, the genes
encoding TrxB were amplified from genomic DNA of *E. coli* MG1655 and *B. subtilis* 168
using the following primers: AI001_for and AI002_rev, and MM001_for and MM002_rev for
E. coli and *B. subtilis*, respectively. The resulting PCR fragments contained an overhang
sequence to the SUMO tag at the 5' end and an XhoI restriction site at the 3' end. Finally, an
overlap PCR was performed using the primers AI003_SUMO_for and AI002_rev (for *E. coli*
TrxB) or MM002_rev (for *B. subtilis* TrxB) to fuse the SUMO tag sequence to the N-terminus
of the respective *trxB* gene. The obtained PCR product (i.e., NdeI-SUMO-TrxB-XhoI) was
then restricted with NdeI and XhoI and cloned into a pET22b(+) vector, resulting in pAI (*E.
coli* MG1655 *trxB*) and pMM (*B. subtilis* 168 *trxB*). *E. coli* DH5α and *E. coli* BL21(DE3) cells
were then transformed with the plasmids for plasmid maintenance and heterologous protein
expression, respectively.

Heterologous protein expression

For heterologous protein expression, a single colony of *E. coli* BL21 (DE3) carrying pAI or
pMM was used to inoculate 50 mL LB broth containing 200 µg/L ampicillin, and the cells
were grown overnight at 37 °C and 125 rpm. The overnight culture was then used to
inoculate 5 L of LB medium, supplemented with ampicillin. Cells were incubated at 37 °C and
125 rpm until an OD$_{600}$ of 0.5-0.6 was reached. Subsequently, protein expression was
induced by adding 1 mM isopropyl 1-thio-ß-D-galactopyranoside (IPTG) to the cell culture.
After overnight incubation at 20 °C and 125 rpm, cells were harvested by centrifugation at
7800 x *g* and 4 °C for 45 min, and pellets were stored at -80 °C.

Purification of ecTrxB and bsTrxB

E. coli BL21 (DE3) cell pellets obtained after expression of *B. subtilis* 168 (*bs*TrxB) or
E. coli MG1655 TrxB (*ec*TrxB) and were resuspended in 30 mL lysis buffer 1 (50 mM sodium
phosphate, 300 mM NaCl, pH 8) containing 2 mL of EDTA-free protease inhibitor mixture
(Roche Applied Science, Penzberg, Germany). Cells were disrupted by passing the cell
suspension three times through a Constant Cell disruption system (TS benchtop; Constant

Systems, Daventry, UK) at 1.9 kbar and 4 °C, followed by the addition of the serine protease

inhibitor phenylmethylsulfonyl fluoride (PMSF) to a final concentration of 1 mM. The cell

lysate was centrifuged at 6700 x g and 4 °C for 1 h. The supernatant was vacuum filtered

through a 0.45 µm filter. Proteins were purified using Immobilized Metal Affinity

Chromatography (IMAC). The supernatant was loaded onto a Pierce Centrifuge Column pre-

packed with 2 mL HisPur™ Ni-NTA-Resin (Thermo Fischer Scientific) and pre-equilibrated

with lysis buffer 1. The column was then washed with lysis buffer 1, supplemented with 10

mM imidazole, followed by one washing step with lysis buffer 1 containing 20 mM imidazole.

Proteins were eluted by adding elution buffer 1 (50 mM sodium phosphate, 300 mM NaCl,

250 mM imidazole, pH 8.0). Elution fractions containing the respective protein were

combined, filled to 50 mL with lysis buffer 1, and concentrated to 1 mL using a Vivaspin 20

PES, MWCO 10000 concentrator system (Sartorius Stedim Biotech). The latter step was

repeated twice to reduce imidazole concentration in the buffer. To remove the SUMO tag,

ecTrxB and bsTrxB were digested with a His$_6$-tagged SUMO Ulp1 protease, using a 75-fold

excess (w/w) protein to protease. After overnight incubation at 4 °C, the reaction mixtures

were purified using IMAC as described above. Wash fractions containing the purified and

untagged TrxBs were concentrated using Vivaspin 20 PES, MWCO 10000 concentrator

system (Sartorius Stedim Biotech) to a final volume of 1 mL. Proteins were stored in lysis

buffer 1 supplemented with 10 % (v/v) glycerol at -80 °C.

Table 1: Bacterial strains and plasmids used in this study.

	Relevant properties	Source/reference
Bacterial strains		
A. baumannii DSM 30007	type strain	DSMZ (German collection of microorganisms and cell culture; Braunschweig, Germany).
B. subtilis 168	Wild type strain	DSMZ, strain DSM402
B. subtilis LQB001	*B. subtilis* 168 containing chromosomally inserted roGFP2 under the control of a xylose-inducible promotor in the *amyE* locus	This work.
E. coli DH5α	Cloning strain	New England Biolabs, Ipswich, MA, USA

E. coli BL21 (DE3)	Strain for heterologous protein expression from pET vectors	Stratagene, Santa Clara, CA.
E. coli DSM 30083	type strain	DSMZ
E. coli MG1655	Wild type strain	ATCC Manassas, VA, US, strain ATCC 700926
E. coli BW25113	Parental strain Keio collection (Baba et al., 2006)	The National BioResource Project (National Institute of Genomics, Japan)
E. coli JW5503	*E. coli* BW25113 Δ*tolC*	The National BioResource Project (National Institute of Genomics, Japan)
E. coli JW3467	*E. coli* BW25113 Δ*gor*	The National BioResource Project (National Institute of Genomics, Japan)
E. coli JW2663	*E. coli* BW25113 Δ*gshA*	The National BioResource Project (National Institute of Genomics, Japan)
E. coli JW5156	*E. coli* BW25113 Δ*trxA*	The National BioResource Project (National Institute of Genomics, Japan)
E. coli JW0871	*E. coli* BW25113 Δ*trxB*	The National BioResource Project (National Institute of Genomics, Japan)
E. coli AI001	DH5α pAI (*E. coli* trxB in pET22b(+))	This work
E. coli AI002	BL21 (DE3) pAI (E. coli trxB in pET22b(+))	This work
E. coli LQB001	BW25113 pCC_roGFP2	This work.
E. coli MM003	DH5α pMM (*B. subtilis* trxB in pET22b(+))	This work
E. coli MM004	BL21 (DE3) pMM (*B. subtilis* trxB in pET22b(+))	This work

P. aeruginosa DSM 50071	type strain	DSMZ (German collection of microorganisms and cell culture; Braunschweig, Germany).
S. aureus DSM 20231	type strain	DSM (German collection of microorganisms and cell culture; Braunschweig, Germany).
S. aureus ATCC 43300 (MRSA)	Methicillin-resistant	ATCC, Manassas, VA, US.
Plasmids		
pET22b(+)	IPTG-inducible plasmid (high copy number)	Novagen
pMM	pET22b(+) encoding *B. subtilis* 168 TrxR with an N-terminal SUMO tag	This study
pAI	pET22b(+) encoding *E. coli* MG1655 TrxR with an N-terminal SUMO tag	This study
pE-SUMO	SUMO tag	Lab collection
pCC_roGFP2	pCC encoding roGFP2. IPTG-inducible plasmid, amp^R	(Müller et al., 2017)
pSG1729_roGFP2	pSG1729 (Lewis and Marston, 1999) encodes roGFP2 under a xylose-inducible promotor for chromosomal insertion into the amyE locus of *B. subtilis*. amp^R, spec^R	This work
pMX_roGFP2	pMX encoding roGFP2 codon-optimized for *B. subtilis*.	Thermo Fischer Scientific

Analysis of TrxB inhibition by auranofin using a DTNB reduction assay

TrxB activity in the absence or presence of auranofin was determined by performing a 5,5'-dithiobis(2-nitrobenzoic) acid (DTNB) reduction assay. In this NADPH-dependent assay,

TrxB reduces DTNB directly to yield the spectrophotometrically detectable 5-thio-2-
nitrobenzoic acid (TNB). DTNB assays were performed in transparent, flat-bottom 96-well
plates (Sarstedt, Nümbrecht, Germany) as described by Lu et al. (Lu et al., 2013). Briefly,
1 µM *ec*TrxB or *bs*TrxB were added to TE buffer (50 mM Tris-HCl, 2 mM EDTA, pH 7.5)
containing 200 µM NADPH (Sigma-Aldrich, St. Louis, USA) for 5 min, in a total volume of
100 µL. The enzymatic reaction was then initiated by adding 100 µL TE buffer containing
2 mM DTNB (Sigma-Aldrich, St. Louis, USA). The amount of liberated TNB was then
quantified spectrophotometrically by measuring the absorbance at 412 nm for 15 min.
Experiments containing auranofin ranging from 0.1 to 2 µM were performed in triplicate. The
remaining TrxB activity after adding auranofin was normalized to activity in the absence of
auranofin, which was set to 100 %.

Proteomic profiling of auranofin-stressed *E. coli* cells using radioactive labeling and
2D-PAGE

2D PAGE-based proteomic analysis was performed as previously described (Wenzel *et al.*,
2011). For the analysis of the proteomic response of *E. coli* to auranofin exposure,
E. coli MG1655 was used. Briefly, a stock of auranofin was dissolved in DMSO to a final
concentration of 10 mg/mL. *E. coli* MG1655 cells were grown in Neidhardt MOPS minimal
medium (Neidhardt et al., 1974) at 37 °C and 200 rpm until an OD_{500} of 0.35 was reached.
Cultures were split into an untreated control and a culture treated with a sublethal
concentration of 120 µg/mL. After 10 minutes, newly synthesized proteins were labeled with
0.37 MBq/mL L-[^{35}S]-methionine (Hartmann Analytic, Braunschweig, Germany). 5 min. later,
protein synthesis was stopped by the addition of 100 µg/mL chloramphenicol and non-
radioactive L-methionine (to a final concentration of 1 mM L-methionine and 10 mM Tris
originally set to pH 7.5 in the stock solution). Cells were harvested by centrifugation and
washed twice with 100 mM Tris/1 mM EDTA, pH 7.5. The cells were disrupted by
ultrasonication with a VialTweeter (Hielscher, Teltow, Germany), and the supernatant was
recovered after centrifugation. The protein concentration was determined by a Bradford
assay (Roti NanoQuant, Roth, Karlsruhe, Germany). Protein biosynthesis rates were
determined by measuring the incorporation of radioactive methionine in auranofin-treated
and untreated samples after protein precipitation with 20 % (w/v) trichloro acetic acid using a
scintillation counter (Tri-Carb 2800TR, PerkinElmer). 55 µg of protein from pulse-labeled
samples for analytical gels and 300 µg of protein from unlabeled samples for preparative
gels were rehydrated into IPG strips (24 cm, pH 4-7; GE Healthcare, Chalfont St Giles, UK).
A Multiphor II (GE Healthcare) was used for isoelectric focusing, and for the second
dimension PAGE, the Ettan DALTwelve system (GE Healthcare) was used. The gels were
stained with ruthenium(II)tris(4,7-diphenyl-1, 10-phenantrolin disulfonate) and washed with

ethanol and acetic acid (Rabilloud et al., 2001). Fluorescent gel images were taken with the

Typhoon Trio[+] Variable Mode Imager (GE Healthcare) using an excitation wavelength of 532

nm and an emission filter of 600 nm. Radio-labeled analytical gels were dried onto Whatman

paper and exposed to Storage Phospho Screens (GE Healthcare). The phosphoscreens

were scanned with the Typhoon Trio[+] with an excitation wavelength of 633 nm and an

emission filter of 390 nm. For visualization and analysis of the 2D gels, the Decodon Delta

2D image analysis software was used (Bandow et al., 2008). Two biological replicates with

one technical replicate each were analyzed against their respective controls for quantitation.

Protein spots detected as two-fold upregulated compared to the untreated control based on

relative signal intensity were defined as marker proteins upon visual inspection (Wenzel et

al., 2011).

Protein identification from gel spots was performed as described previously (Wenzel et al.,

2013). Spots were excised from a non-radioactive 2D gel and destained with 20 mM

ammonium bicarbonate in 30 % acetonitrile. The protein spots were digested with either

12.5 ng/µL trypsin for low-intensity or 6.25 ng/µL trypsin for high-intensity spots (Promega,

Fitchburg, FI, USA) for 18 h at 37 °C. Peptides were eluted in 0.1 % trifluoroacetic acid in

MS-grade water and ultrasonicated for 15 min once (or twice if they were low-intensity

spots). For protein identification, the samples were analyzed using a nanoACQUITY-UPLC-

coupled SYNAPT G2-S high-definition mass spectrometer equipped with a NanoLockSpray

Source for electrospray ionization and a time-of-flight detector (Waters, Milford, USA). The

samples were loaded into a trap column (C_{18}, pore size 100 Å, particle diameter of 5 µm and

an inner diameter of 180 µm, length 20 mm) and eluted with 0.1 % formic acid/acetonitrile

gradient (350 µL/min, linear gradient 2-5 % acetonitrile in 2 min, 5-60 % acetonitrile in

20 min) from an analytical column (C_{18}, pore size 130 Å, particle diameter of 1.7 µm and an

inner diameter of 75 µm, length 150 mm) at 40 °C and subjected to mass spectrometry. The

spectra were recorded in a mass range of 50 to 1800 m/z with a 0.5 s/scan in positive mode.

The following parameters were used for the NanoLockSpray source: capillary voltage of

1.5 kV; sampling cone voltage of 30 V; source temperature of 100 °C; desolvation

temperature of 150 °C; cone glass flow 50 L/h; desolvation gas flow of 550 L/h. Leucine

enkephalin was used as a lock mass analyte using the lock spray channel (lock spray

capillary voltage, 2.5 kV). Analysis of spectra was performed by MassLynx V4.1 SCN813

(Waters).

For protein identification, a non-redundant version of *E. coli* MG1655 protein database (NCBI

Bio Project accession PRJNA647163) with 4740 protein entries (plus the sequences of

trypsin and human keratin) and the ProteinLynxGlobalServer (PLGS) software V3.5.3

(Waters) with the following parameters were used: chromatographic peak width, automatic;

MS ToF resolution, automatic; lock mass window, 0.3 Da; low energy threshold, 50 counts.

Other parameters were as follows: peptide tolerance, automatic; fragment tolerance,

automatic; minimal fragment ion matches per peptide, 2; minimal fragment ion matches per

protein, 5; minimal peptide matches per protein, 2; maximum protein mass, 250 kDa; primary

digest reagent, trypsin; secondary digest reagent, none; missed cleavages, 1; fixed

modifications, carbamidomethyl C; variable modifications, deamination N and Q, oxidation M;

false positive rate, 4 %. The following parameters were considered for protein assignment:

Coverage of identified protein was > 10 %, at least 3 peptide ions and 5 product ions were

assigned, and theoretical pI and MW were consistent with the spot's position on the 2 D gel.

RoGFP2 expression in bacteria.

To measure the redox state in the cytoplasm of *E. coli* MG1655 and *B. subtilis* 168, cells

were transformed with the pCC_roGFP2 plasmid as described (Müller et al., 2017) and

pSG1729_roGFP2, respectively. pSG1729_roGFP2 cannot replicate in *B. subtilis*, but part of

it integrates into the *amyE* locus on the chromosome through a double-crossover event. To

create pSG1729_roGFP2, the EGFP insert from pSG1729 (Lewis and Marston, 1999) was

removed using KpnI and HindIII restriction enzymes. The gene sequence of roGFP2, codon-

optimized for *B. subtilis*, was amplified from pMX_roGFP2 (Thermo Fischer Scientific) by

PCR. Restriction sites for KpnI and HindIII were introduced using the primers JY001_for and

JY001_rev. Plasmid construction followed standard procedures and was performed in *E. coli*

DH5α. After successful plasmid construction, confirmed by sequencing, pSG1729_roGFP2

was transformed into *B. subtilis* 168 using standard protocols (Anagnostopoulos and

Spizizen, 1961), and selection for chromosomal insertion of the plasmid was performed on

LB plates containing 100 µg/mL spectinomycin. Double crossover of positive clones was

confirmed by the inability to hydrolyze starch on LB plates containing 0.5 % starch and

subsequent PCR amplification of the chromosomal insert and sequencing of the PCR

product.

For roGFP2 expression in *E. coli* MG1655, cells were grown in MOPS medium at 37 °C and

125 rpm. 200 µM of IPTG were used to induce protein expression at an OD_{600} of 0.4-0.6, and

roGFP2 was subsequently expressed for 16 h at 20 °C and 125 rpm in MOPS medium. After

expression, cells were harvested, washed in HEPES buffer (40 mM, pH 7.4), and adjusted to

an OD_{600} of 1.2.

For roGFP2 expression in *B. subtilis* 168, a single colony was inoculated in DSM medium

(Bacto nutrient broth 0.8 % (w/v), KCl 13.4 mM, $MgSO_4$ 1 mM, $Ca(NO_3)_2$ 1 mM, $MnCl_2$ 10 µM,

$FeSO_4$ 1 µM) containing 0.1 % xylose (w/v), and cells were grown for 16 h at 30 °C and

200 rpm. Cells were harvested, washed in BMM medium (Stulke et al., 1993), and incubated

for 30 min at 25 °C and 800 rpm in a ThermoMixer (Eppendorf, Hamburg, Germany). The

final OD_{600} was adjusted to 5. Cells were treated with 200 µM aldrithriol-2 (Sigma-Aldrich)

(AT-2) as an oxidation control and 2 mM dithiothreitol (Sigma-Aldrich) as a reduction control.

0.5, 1, and 5 µM auranofin were used as treatment. As AT-2 and auranofin were dissolved in

DMSO, an equivalent volume of DSMO was added to all samples. 1 mL of cell suspension

was added to a 1500 µL Quartz Suprasil fluorescence cuvette (Hellma, Müllheim, Germany)

with a stirring bar. Measurements were done at 25 °C in a JASCO FP-8500 fluorescence

spectrometer equipped with the temperature-controlled sample holder 'EHC-813' at 25 °C for

60 min under continuous stirring. Measurement parameters in *E. coli* were as follows:

510 nm (Em), 350–500 nm (Ex), 5 nm slit width (Ex/Em), and medium sensitivity. After

stabilization, respective treatments were added at the indicated final concentrations, and a

time series of 61 additional spectra was recorded. Measurement parameters for *B. subtilis*

were as follows: 520 nm (Em), 350–500 nm (Ex), 10 nm slit width (Ex/Em), and high

sensitivity. After stabilization, respective treatments were added at the indicated final

concentrations, and a time series of 61 additional spectra was recorded. The ratios of the

fluorescence excitation intensities (405/488 nm) were used to calculate the probe's oxidation

state. All values were normalized to fully oxidized (AT-2-treated) and fully reduced (DTT-

treated) roGFP2 with the following equation [1]:

$$OxD = \frac{R - R_{red}}{\left(\frac{I_{488}ox}{I_{488}red}\right) * (R_{ox} - R) + (R - R_{red})}$$

R_{ox} is the 405/488 nm ratio of oxidized (AT-2 treated) and R_{red} of reduced (DTT-treated) cells

expressing roGFP2, respectively. $I_{488}ox$ and $I_{488}red$ are the fluorescence intensities at 488 nm

under oxidizing (i.e., AT-2) or reducing (i.e., DTT) conditions. R is the measured 405/488 nm

ratio of the sample.

For calculation and normalization of the probe's oxidation degree at each time point, the

means of all values recorded at all time points for R_{red}, R_{ox}, $I_{488}ox$, and $I_{488}red$ were used.

Data was processed using Microsoft Excel software and GraphPad Prism version 10.1.0.

RESULTS

Auranofin is antibacterial, but less effective against *E. coli* and other Gram-negative
species.

Previous studies have shown that auranofin possesses potent bactericidal activity against
many Gram-positive bacteria but seems to be markedly less effective against pathogenic
Gram-negative species (Cassetta et al., 2014; Harbut et al., 2015; Hokai et al., 2014).
However, the reasons for the poor susceptibility of Gram-negative bacteria to auranofin are
not entirely clear. We analyzed the antimicrobial activity of auranofin against Gram-negative
and Gram-positive bacterial strains by minimal inhibitory concentration (MIC) assays. For this
purpose, we added various concentrations of auranofin to a defined number of bacterial cells
of *B. subtilis* 168, *S. aureus* DSM 20231/ATCC 43300, *P. aeruginosa* DSM 50071,
A. baumanni DSM 30007 and *E. coli* DSM 30083 and examined their growth after overnight
incubation. The MIC was defined as the lowest auranofin concentration, at which no visibly
perceptible bacterial growth was observed. In line with expectations, auranofin showed
potent growth inhibition against Gram-positive bacteria, with an MIC of 0.5 µg/mL for each
Gram-positive strain tested (Table 2). The MIC values determined for auranofin against *S.*
aureus and *B. subtilis* correlate with those reported in previous studies (Cassetta et al., 2014;
Harbut et al., 2015; Hokai et al., 2014). In contrast, auranofin exhibited poor antimicrobial
activity against Gram-negative bacteria, with MICs of 128 µg/mL, 32 µg/mL, and 512 µg/mL
against *E. coli*, *A. baumannii,* and *P. aeruginosa*, respectively, which agrees with previous
reports (Harbut et al., 2015; Thangamani et al., 2016).

Table 2: MICs of Auranofin against various Gram-negative and Gram-positive bacterial species.
401

Strain	MIC, µg/mL
Gram-negative bacteria	
E. coli DSM 30083	128
A. baumannii DSM 30007	32
P. aeruginosa DSM 50071	512
Gram-positive bacteria	
B. subtilis 168	0.5
S. aureus DSM 20231	0.5
S. aureus ATCC 43300 (MRSA)	0.5

Auranofin causes upregulation of proteins involved in central metabolism and the defense against oxidative stress in *E. coli*.

To understand if auranofin exposure leads to a markedly different cellular response in Gram-negative bacteria, we performed proteomics experiments in *E. coli*. We compared our results to our results previously obtained for *B. subtilis* (Senges et al., 2020). In our experiments, we treated an exponentially growing culture of *E. coli* MG1655 with a sublethal concentration of 120 µg/mL auranofin. This concentration was found to partially inhibit bacterial growth by 26.27 ± 4.14 % (Fig. 1a). Upon exposure to auranofin for 10 min, the cells were pulse-labeled for 5 minutes with L-[^{35}S]-methionine, which was incorporated into newly synthesized proteins. Quantification of total L-[^{35}S]-methionine incorporation revealed no significant difference between the control and the auranofin-treated culture, suggesting that auranofin does not directly inhibit protein synthesis (Fig. 1b).

Newly synthesized proteins were quantified by densitometry of radioactive spots after two-dimensional PAGE. Auranofin-induced changes in protein synthesis patterns were visualized by overlaying the gel images of the L-[^{35}S]-methionine-labeled proteins under control conditions (green) and after treatment with auranofin (red). Protein spots synthesized only under control conditions appear in green; protein spots that were synthesized only upon auranofin treatment appear in red, whereas protein spots synthesized under both conditions appear yellow. Out of the three independent experiments, a representative dual-channel image produced from autoradiographs of 2D gels obtained for control and auranofin-treated cells is shown in Fig. 1c. Auranofin treatment caused a more than two-fold increase in relative synthesis rates of 30 protein spots (indicated by arrows). From non-radioactive gels, proteins could be identified by mass spectrometry in 20 of those spots (Fig. 1c, Tables 3 and S2).

Most identified marker proteins upregulated in response to auranofin treatment are associated with a specific response to oxidative stress and/or involved in cellular processes such as ROS detoxification, central metabolism, amino acid biosynthesis, and metal homeostasis.

Many of the upregulated proteins are enzymes of the central metabolic network, such as GpmM (phosphoglycerate mutase), PykF (pyruvate kinase I), Lpd (lipoamide dehydrogenase), and the alpha subunit of the ATP synthase F$_1$ complex of the electron transport chain AtpA. Auranofin treatment was also found to increase the synthesis of the enzyme dihydroorotase (PyrC), which functions in the pathway for the biosynthesis of pyrimidine nucleotides. Some other induced proteins, such as AhpF and AhpC, which are components of the alkyl hydroperoxide reductase, and Gor (glutathione oxidoreductase),

belong to the OxyR regulon and are involved in the protection against peroxide and in

maintaining redox balance (Storz et al., 1990; Storz and Tartaglia, 1992). Gor is responsible

for recycling the glutathione pool NADPH-dependently. NADPH is thought to be delivered by

the pentose pathway, of which RpiA was upregulated. A similar observation was made in our

laboratory when *E. coli* cells were subjected to diamide or the antimicrobial compound allicin,

both known to induce disulfide stress (Müller et al., 2013, 2016): here, central metabolic

enzymes and OxyR-dependent proteins were upregulated, as well. Additionally, auranofin

treatment led to the upregulation of a sulfur assimilation enzyme (CysN) for the biosynthesis

of sulfur-containing amino acids and the oxidative stress-protective multicopper oxidase

CueO.

To test if one of these upregulated proteins is directly involved in *E. coli*'s tolerance against

auranofin, we tested, if selected single mutants from the KEIO collection lacking these

proteins were more susceptible to auranofin in an MIC assay. The mutants lacking GpmM,

Pgk, Lpd, and PyrC did not grow in MOPS medium. None of the proteins missing in the

selected mutants that did grow, impacted the MIC when absent, except for NemA (Table 3).

The Δ*nemA* strain was found to be slightly more resistant to auranofin treatment and had an

MIC of 256 µg/mL (WT being 128 µg/mL).

Overall, the proteomic results indicate that auranofin-treated cells suffer systemic oxidative

stress. This could lead to the direct inactivation of essential thiol-containing enzymes, e.g., in

the central metabolism, and a depletion of free cysteine levels or, more generally, thiol levels.

We made a similar observation in *B. subtilis* (Senges et al., 2020), and based on our

proteome data, we could not differentiate between the mode of action of auranofin in Gram-

positive and Gram-negative bacteria.

461
462
Figure 1: Proteomic response of *E. coli* MG1655 stressed with 120 μg/mL Auranofin. (a) Growth
curves of *E. coli* MG1655 grown in MOPS media at 37 °C and 200 rpm without auranofin (■) and after
the addition of 120 μg/mL auranofin (□). An arrow indicates the timepoint of auranofin addition. **(b)**
Incorporation of L-[³⁵S]-methionine during a 5 min pulse starting 10 min after treatment with 120 μg/mL
auranofin. **(c)** 2D-PAGE analysis of the pulse-labeled proteomics response profile of *E. coli* MG1655
treated with 120 μg/mL auranofin. False-color overlay of treated and untreated radioactive gels shows
proteins synthesized after pulse labeling for 5 min after treatment with auranofin (red, Au) or in the
untreated control (green, Co). Upregulated proteins with a regulation factor ≥ 2 were considered marker
proteins and were labeled by arrows and the Spot ID Au_X. Marker proteins identified by mass
spectrometry are additionally labeled with the protein's name. For Spot ID, identified proteins, and
functions, see Table 3 and Supplemental Table S2. **(a, c)** Representative example from 3 **(a)** or 2 **(c)**
biological replicates. **(b)** Average and standard deviation of 3 biological replicates.

476 **Table 3: Identified marker proteins of auranofin-treated *E. coli* MG1655.** Protein functions and
477 accession number (EcoCyc ID) were taken from the EcoCyc database (Keseler et al., 2017). Regulation
478 factor (Reg. factor) is the average of 2 biological replicates. The MIC of the corresponding deletion
479 mutant is shown (n=3). Deletion mutants not viable in minimal medium are marked by n. v., MIC not
480 determined is marked by n. d.
481

Spot ID	Reg. factor	Protein (EcoCyc ID)	Function	MIC µg/mL
497	2.3	**AhpC** (EG11384)	alkyl hydroperoxide reductase, AhpC component	128
860	3.6	**AhpF** (EG11385)	alkyl hydroperoxide reductase, AhpF component	128
		LeuA (EG11226)	2-isopropylmalate synthase	n.d.
763	4.6	**Amn** (EG10039)	AMP nucleosidase	128
		AtpA (EG10098)	ATP synthase F1 complex subunit α	128
		CueO (EG12318)	multicopper oxidase CueO	128
756	4.0	**Amn** (EG10039)	AMP nucleosidase	128
		AtpA (EG10098)	ATP synthase F1 complex subunit α	128
		Lpd (EG10543)	lipoamide dehydrogenase	n.v.
758	15.0	**AtpA** (EG10098)	ATP synthase F1 complex subunit α	128
347	3.2	**Bfr** (EG10113)	bacterioferritin	128
690	2.1	**CysN** (EG10194)	sulfate adenylyltransferase subunit 1	128
		GpmM (EG12296)	2,3-bisphosphoglycerate-independent phosphoglycerate mutase	n.v.
782	4.0	**GlmM** (EG11553)	phosphoglucosamine mutase	n.d.
		Gor (EG10412)	glutathione reductase	128
		Mpl (EG12440)	UDP-N-acetylmuramate-L-alanyl-γ-D-glutamyl-meso-2,6-diaminoheptanedioate ligase	n.d.
		PhoA (EG10727)	alkaline phosphatase	n.d.
692	2.6	**GpmM** (EG12296)	2,3-bisphosphoglycerate-independent phosphoglycerate mutase	n.v
44	4.1	**HtpG** (EG10461)	chaperone protein HtpG	n.d.
754	7.2	**Lpd** (EG10543)	lipoamide dehydrogenase	n.v.
620	2.7	**MglB** (EG10593)	D-galactose/methyl-galactoside ABC transporter periplasmic binding protein	128
328	2.4	**MoaB** (EG11596)	protein MoaB	128
806	3.9	**NemA** (G6890)	N-ethylmaleimide reductase	256
718	6.1	**Pgk** (EG10703)	phosphoglycerate kinase	n.v.
859	3.6	**PykF** (EG10804)	pyruvate kinase 1	128
816	3.0	**PyrC** (EG10806)	dihydroorotase	n.v.
		YqhD (G7564)	NADPH-dependent aldehyde reductase YqhD	n.d.
239	2.0	**RpiA** (EG11443)	ribose-5-phosphate isomerase A	n.d.
25	3.2	**SucA** (EG10979)	2-oxoglutarate decarboxylase, thiamine-requiring	128
662	3.9	**YeaG** (G6969)	protein kinase YeaG	128

The higher resistance against auranofin can be partially explained by more effective efflux systems in Gram-negative bacteria.

Efflux pumps are critical players in antimicrobial resistance in Gram-negative bacteria. These pumps actively transport a variety of substrates, including antibiotics and heavy metals, out of the bacterial cell, thereby reducing the intracellular concentration of these compounds and conferring resistance to the cell. The outer membrane protein TolC, part of the small multidrug resistance (SMR) protein family, is a significant component involved in the export of various cytotoxic substances. Studies have shown that mutations in the TolC protein resulted in sensitivity to different antibiotics, such as vancomycin, erythromycin, and novobiocin (Augustus et al., 2004), and an increased susceptibility to metal ions (Munita and Arias, 2016).

We found that cellular susceptibility to auranofin was increased two-fold when we tested the MIC of an *E. coli* Δ*tolC* strain (Table 4). The MICs of *B. subtilis* and *S. aureus* were 0.5 μg/ml, i.e., still more than two orders of magnitude lower than that of the TolC-deficient *E. coli* strain. This suggests that TolC does play a role in the efflux of auranofin from the bacterial cell, albeit not to an extent that would fully explain the overall higher resistance of Gram-negative bacteria.

Table 4: MICs of Auranofin against *E. coli* BW25113 and *E. coli* BW25113 Δ*tolC*.

Strain	MIC, μg/mL
E. coli BW25113 (WT)	128
E. coli JW5503 (BW25113 Δ*tolC*)	64

Auranofin inhibits TrxB from both *E. coli* and *B. subtilis* to a similar extent.

Evidence suggests that auranofin primarily exerts its effect by targeting bacterial thiol-redox homeostasis by inhibiting thioredoxin reductase (TrxB) (Harbut et al., 2015; Hokai et al., 2014). Since TrxB is responsible for reducing oxidized thioredoxin (TrxA) and, thus, for maintaining a reducing intracellular environment, loss of TrxB activity can lead to severe oxidative stress in an organism with no alternative antioxidant system, such as the glutathione/glutaredoxin system.

While auranofin has been found to effectively inhibit TrxB of *S. aureus*, *M. tuberculosis,* and other Gram-positive bacterial species (Harbut et al., 2015; Owings et al., 2016), less is known about its inhibitory potency towards TrxB of Gram-negative bacteria. Hence, one possible explanation for auranofin's limited antibacterial activity against Gram-negative

bacteria might be that auranofin is a less effective inhibitor of TrxB in these organisms when

compared to Gram-positive bacteria. To explore this hypothesis, we tested auranofin's

effectiveness in a biochemical assay with TrxB from Gram-negative and Gram-positive

bacteria, with *E. coli* and *B. subtilis* as representative organisms, respectively.

The activity of purified TrxB from *E. coli* and *B. subtilis* (from now on referred to as "*ec*TrxB"

and "*bs*TrxB", respectively) was then determined in the presence of NADPH, and various

auranofin concentrations by performing a 5,5'-dithiobis(2-nitrobenzoic) acid (DTNB) reduction

assay (Arnér and Holmgren, 2005). The NADPH-dependent TrxB reduces DTNB in this

assay, yielding the detectable 5-thio-2-nitrobenzoic acid (TNB). The amount of liberated TNB

can be quantified spectrophotometrically by measuring its absorbance at 412 nm.

523
524
Figure 2: Auranofin inhibits *E. coli* (*ec*TrxB) and *B. subtilis* TrxR (*bs*TrxB) in a dose-dependent
manner. Purified *ec*TrxB and *bs*TrxB (1 µM) were preincubated with 200 µM NADPH in the absence (0
µM auranofin) or presence of various auranofin concentrations (0.1 to 2 µM) for 5 min. The addition of
2 mM DTNB initiated the reaction. Reaction progress was monitored by measuring the absorbance of
the liberated TNB chromophore at 412 nm over time. Experiments were performed in triplicates. Mean
(bars), individual values (dots) and SD are shown (error bars).
531

When auranofin was preincubated with 1 µM *ec*TrxB or *bs*TrxB, auranofin inhibited both

enzymes dose-dependently. The activity measurements showed potent and nearly complete

inhibition of *ec*TrxB and *bs*TrxB already at equimolarity (1 µM auranofin, corresponding to ~

0.68 µg/mL) (Fig. 2). While the auranofin concentration required to fully inhibit TrxB agreed

well with the previously determined MIC against *B. subtilis* (Table 1), it was far below the

level required to exhibit cytotoxicity towards *E. coli*. This observation strongly suggests that

the lower antimicrobial activity of auranofin against *E. coli* is not due to a lower affinity of

auranofin for *ec*TrxB.

Trx reductase is not the only target of Auranofin in Gram-negative bacteria.

Based on the results of our biochemical activity assays, we suspected that TrxB may not be

the only target of auranofin in *E. coli*. We speculated that other redox pathway components

might also be involved in the mechanism of action of auranofin. To test our hypothesis, we

analyzed auranofin's antimicrobial activity against additional mutants of the KEIO collection

(Baba et al., 2006). The mutants we chose lack components of the Trx antioxidant system,

i.e., the redox protein thioredoxin TrxA (Δ*trxA*; *E. coli* JW5156) or TrxB (Δ*trxB*; *E. coli*

JW0871) itself, respectively (Fig. 3). Growth of the selected *E. coli* strains in the absence of

auranofin was monitored until an OD_{600} of 0.5 was reached. *E. coli* mutant strains without a

functional Trx system showed a comparable or only slightly lower growth rate than the wild

type, excluding the possibility that potential differences in the susceptibility of the different

strains to auranofin are solely due to a generally impaired growth of the deletion mutants

(Fig. 3a). The cultures were then used to inoculate medium containing different amounts of

auranofin for an MIC assay. After overnight incubation, cultures were analyzed for growth. *E.*

coli deficient in TrxA or TrxB showed a four-fold increased sensitivity to auranofin compared

to wild-type *E. coli*, with MICs of 32 µg/mL (Δ*trxA* and Δ*trxB*) in contrast to 128 µg/mL (WT)

(Fig. 3b, c). The increased susceptibility of *E. coli* mutants lacking a functional Trx system,

especially the suspected target TrxB, indicates that auranofin leads to global oxidative stress

not only through TrxB inhibition but also in a direct manner or through inhibition of other thiol-

disulfide oxidoreductase systems.

The glutathione reductase confers no resistance to auranofin, while the glutathione

pool by itself has a substantial effect on the MIC.

Although auranofin showed a generally higher efficacy against *E. coli* strains deficient in the

functional Trx system, the MIC values were still markedly higher than those observed with

Gram-positive bacteria. In contrast to most Gram-negative bacteria, many Gram-positive

species lack glutathione. Glutathione by itself is considered an antioxidant, and with the small

redox protein glutaredoxin, it can compensate for the loss of a functional Trx system (Gon et

al., 2006). In Gram-positive organisms, on the other hand, the Trx system is essential (Lu

and Holmgren 2014; Uziel et al. 2004; Scharf et al. 1998). Hence, the additional

glutathione/glutaredoxin system in Gram-negative bacteria could be a reason for auranofin's

low potency against *E. coli* and other Gram-negative species. However, we already knew

from our MIC tests performed in mutants deficient in proteins upregulated in our proteomics

experiments that a Δ*gor* mutation did not affect the MIC. Thus, we argued that the absence

of a functional glutaredoxin-system does not play a role. We then tested if glutathione, the

tripeptide itself, could play an important role instead. Both, deletion of glutathione

oxidoreductase (Δ*gor*) and γ-glutamate-cysteine-ligase (Δ*gshA*), the enzyme catalyzing the

first step of glutathione biosynthesis, did not affect *E. coli*'s growth in the absence of

auranofin (Fig. 3a). But auranofin's activity against *E. coli* Δ*gshA* was markedly higher than

against the WT or Δ*gor*, which both had the same MIC. The MIC of Δ*gshA* decreased to 8

µg/mL (Fig. 3b, c), closer to the MIC observed in Gram-positive bacteria. The increase in *E.*

coli's sensitivity to auranofin upon depletion of the intracellular glutathione pool suggests that

this monothiol, typically present at concentrations in the millimolar range, confers substantial

protection against oxidative stress induced by auranofin.

Auranofin leads to thiol oxidation in the cytosol.

We then wanted to measure the intracellular thiol redox state in auronofin-treated bacteria *in situ*. roGFP2 is a genetically engineered variant of GFP containing two cysteines, which form a disulfide bridge upon oxidation, thereby changing the protein's fluorescent properties. This can be used to determine the oxidation degree of roGFP2 (Aller et al., 2013), independent of the probe's concentration (Dooley et al., 2004; Meyer and Dick, 2010; Müller et al., 2017). Because we assumed that auranofin interferes not only with TrxB but with the oxidation state

of thiols in general, we used roGFP2 as a tool to monitor the redox state of cysteines in

cellular proteins. We expressed roGFP2 in the cytoplasm of both *B. subtilis* and *E. coli*. We

then determined the oxidation degree of roGFP2 inside the cells. The oxidant AT-2 was used

to fully oxidize, and the reductant DTT to fully reduce roGFP2 within cells, setting the fully

reduced and fully oxidized baselines of our probe within the bacterial cell. In both strains, the

probe is more or less fully reduced in the cytosol of untreated cells. Using a concentration of

5 µM auranofin, the degree of oxidation of roGFP2 increases by 40 % after 60 min of

treatment in *E. coli* (Fig. 4b), whereas it increases by up to 90 % in less than 1 min in

B. subtilis (Fig. 4a). Lower concentrations of 1 µM or 500 nM auranofin still increased the

oxidation state of roGFP2 in *B. subtilis* to around 80 % within 5 min of treatment, while in

E. coli an oxidation state of only 10 to 20 % was reached after 60 min. The subsequenmt

addition of 2 mM DTT to the cells led to an immediate decrease in probe oxidation in all

cases, demonstrating that, in principle, the change in cellular oxidation imposed by auranofin

is reversible by thiol reductants in both organisms.

Figure 4: Auranofin exposure leads to roGFP2 oxidation *in vivo*. To evaluate the effects of auranofin
on protein thiols in bacteria *in situ*, *B. subtilis* **(a)** and *E. coli* **(b)** expressing roGFP2 were treated with
0.5, 1, or 5 µM auranofin. Aldrithiol-2 and dithiothreitol were used to fully oxidize and fully reduce the
probe within cells, respectively, and these baselines (not shown) were used to calculate the oxidation
degree (OxD) of roGFP2. OxD normalizes the ratiometrically determined redox state and accounts for
differences in probe synthesis and measurement settings. roGFP2 fluorescence was measured at an
emission wavelength of 510 nm, and the excitation wavelength was scanned from 350–500 nm. roGFP2
oxidation was monitored for 60 minutes after adding auranofin, and then DTT was added at a
concentration of 2 mM (indicated by an arrow).

DISCUSSION

The gold(I) compound auranofin, an FDA-approved antirheumatic drug, has recently been found to possess antimicrobial activity and, thus, is a promising candidate for drug repurposing for treating multidrug-resistant infections. Although auranofin shows high activity against a considerable range of Gram-positive bacteria, it seems less effective against Gram-negative species (Harbut et al., 2015; Owings et al., 2016). In the present study, we aimed to understand the underlying mechanism of the difference in susceptibility of Gram-negative and Gram-positive bacteria to auranofin.

We thus tested auranofin's ability to inhibit the growth of different Gram-positive and Gram-negative species. Growth of Gram-positive strains was prevented by an auranofin concentration as low as 0.5 µg/mL. These results agree with previously reported MICs against *B. subtilis* PY79*, M. tuberculosis,* and several methicillin-resistant *S. aureus* (MRSA) strains (Harbut et al., 2015; Hokai et al., 2014). As expected, a substantial difference in the effective concentration of auranofin was found for the Gram-negative species *A. baumannii* DSM 30007, *E. coli* DSM 30083, and *P. aeruginosa* DSM 50071 with MIC values of 32, 128, and 512 µg/mL, respectively.

Proteomic profiling of auranofin-stressed *E. coli* revealed that its cellular response is similar to the response of *B. subtilis*. *B. subtilis* was found to counteract auranofin stress by specifically upregulating enzymes that detoxify ROS, such as the catalase KatA, or replenish the cysteine pool, such as YrhB (Senges et al., 2020). Similarly, in *E. coli*, proteins belonging to the OxyR regulon, such as AhpC and AhpF, the two components of *E. coli*'s alkyl hydroperoxide reductase and glutathione oxidoreductase Gor, were upregulated significantly. The transcription factor OxyR regulates the expression of antioxidant genes in response to oxidative stress, particularly elevated levels of hydrogen peroxide (Storz et al., 1990; Storz and Tartaglia, 1992). We showed previously that OxyR-dependent genes are also upregulated in response to the oxidant diamide and the antimicrobial compound allicin, both of which were found to induce disulfide stress in *E. coli* (Müller et al., 2013, 2016). We then tested the susceptibility of mutants devoid of proteins upregulated in our proteomic experiments to auranofin. None of the mutants tested showed a lower MIC and, thus, higher susceptibility to auranofin, suggesting that these proteins are not the reason for *E. coli*'s high tolerance for auranofin.

Previous studies on auranofin as an antimicrobial drug suggested that, as in mammals, the flavoenzyme thioredoxin reductase (TrxB) is a primary target in bacteria (Harbut et al., 2015). Our studies showed that auranofin inhibits mammalian TrxR with a K_i of only 4 nM (Gromer et al. 1998). Along this line, auranofin has been shown to effectively inhibit bacterial TrxB

from numerous Gram-positive pathogens, including *M. tuberculosis* and *S. aureus* (Fuchs et
al., 2016; Harbut et al., 2015). However, it should be noted that bacterial TrxB is structurally
significantly different from mammalian TrxR and does not contain selenocysteine (Lu and
Holmgren, 2014). Selenocysteine is thought to play a major role in the inhibition of
mammalian TrxR by auranofin: Several lines of evidence suggest that selenocysteine is
either the direct binding site of auranofin or is essential for releasing the gold atom from its
ligands, given that the affinity of selenium for gold and its nucleophilic power are greater than
those of sulfur (Angelucci et al., 2009; Arscott et al., 1997; Dos Santos, 2014; Rigobello et
al., 2004, 2005). Indeed, there is direct evidence for auranofin binding to the selenocysteine
in human TrxR (Pickering et al., 2020). Additionally, TrxBs from Gram-positive and Gram-
negative bacteria are structurally much more similar to each other than to mammalian TrxR.
Bacterial TrxB has only a low degree of sequence homology (<30 %) to human TrxR (Arscott
et al., 1997).

Nevertheless, to uncover potential differences in TrxB inhibition in Gram-negative and Gram-
positive bacteria, which could explain the different susceptibilities of the species, we
biochemically tested the inhibition of TrxB from *E. coli* (*ec*TrxB) and *B. subtilis* (*bs*TrxB) by
auranofin. We found that inhibition of both *ec*TrxB and *bs*TrxB by auranofin is dose-
dependent, and in both cases, an equimolar amount is enough to inhibit their activity entirely.
These observations suggest that inhibition of bacterial TrxB by auranofin could be
irreversible. In any case, no significant difference in susceptibility could be observed between
TrxB from Gram-positive and Gram-negative species in our assay. This shows that, at a
target level, both Gram-positive and Gram-negative species are equally susceptible.

To test other properties that could make Gram-negative bacteria more resistant to auranofin,
we examined an *E. coli* strain lacking TolC. This efflux pump removes antibiotics and metals
from the cell (Koronakis, 2003). We found that the lack of TolC decreased auranofin's
resistance only by a factor of two, not fully explaining Gram-negative organisms' substantially
higher resistance than Gram-positive bacteria.

Unlike most Gram-negative bacteria, many Gram-positive bacteria do not possess the low
molecular weight thiol glutathione and the glutaredoxin system. Hence, TrxB and its
substrate thioredoxin (TrxA) are considered the key mediators of redox homeostasis in these
organisms (Lu and Holmgren, 2014), and the Trx system is essential for *B. subtilis* and *S.*
aureus' survival at normal growth conditions and under oxidative stress (Scharf et al., 1998;
Uziel et al., 2004). Conversely, *E. coli* strains deficient in TrxB or TrxA did show only
moderately higher susceptibility towards auranofin. These findings, too, suggest that TrxB
may not be the sole target of auranofin in *E. coli*. Following up on the idea that glutathione

and the associated glutaredoxins could compensate for the loss of TrxB in *E. coli*, as suggested by Harbut et al. (Harbut et al., 2015), we revisited auranofin's activity against an *E. coli* mutant devoid of the glutathione reductase (Gor). As there was no difference in the susceptibility to auranofin, in line with findings from Thangamani et al. (Thangamani et al., 2016), we concluded that the glutaredoxin system as a backup for thioredoxin is not the main reason for *E. coli*'s relative resistance to auranofin. However, the loss of the major intracellular low molecular weight thiol glutathione itself led to a substantial increase in sensitivity to auranofin. Therefore, several possible factors for the high tolerance of Gram-negative bacteria towards auranofin must exist. First, Gram-negative bacteria possess more effective mechanisms to avoid auranofin stress, i.e., the presence of efflux pumps, potentially in combination with the presence of an outer membrane. Second, the presence of glutathione was, in our experiments, the most effective contributor to auranofin resistance in *E. coli*. Glutathione's high concentration of up to 10 mM in the cell could make it a potent buffer against auranofin, depleting the gold compounds' ability to react with protein thiols. These findings corroborate that auranofin induces disulfide stress not only through TrxB inactivation but potentially through direct thiol modification in proteins and reaction with low molecular weight thiols in general.

The cysteine-containing redox probe roGFP2, synthesized in the cytosol of bacterial cells, revealed that auranofin indeed directly affected protein oxidation. A high oxidation degree was reached within minutes in Gram-positive bacteria. This quick oxidation is not observed in Gram-negative bacteria, presumably due to the existence of efflux pumps in combination with the presence of high concentrations of glutathione.

In clinical trials with auranofin, plasma concentrations of auranofin ranging between 0.8 and 1.5 µg/mL were reached (Capparelli *et al.*, 2017). While this therapeutically achievable auranofin concentration is sufficient to inhibit the growth of most Gram-positive pathogens, it is 20 to 340 times lower than the auranofin concentration necessary to prevent the growth of the tested Gram-negative strains, rendering auranofin virtually ineffective against these organisms in a clinical setting. Its limited activity against Gram-negative bacteria currently prevents its systemic clinical use as an antibiotic in treating infections with Gram-negative pathogens. The results presented here strongly suggest that, while auranofin's mechanism of action against *E. coli* does not substantially differ from that against *B. subtilis per se*, other factors unique to Gram-negative bacteria are attenuating auranofin's antimicrobial effectiveness. Thus, potential drugs based on auranofin's mechanism of action need to account for those factors, particularly the presence of high cellular concentrations of glutathione.

Our results indicate that the mode of action of auranofin in bacteria is a combination of the

inactivation of thiol-containing enzymes, a decrease of reduced Trx levels through TrxB

inactivation, and the induction of systemic oxidative stress. While most clinically used

antibiotics act on a single target, this broad-range effect of auranofin could prevent the

evolution of specific resistance mechanisms, and indeed several others reported that

spontaneous auranofin-resistant mutants cannot be generated (Harbut et al., 2015;

Thangamani et al., 2016; Elkashif and Seleem, 2020). This should make the development of

other gold-containing drugs, effective in Gram-negative bacteria, and based on auranofin's

mode of action a worthwhile endeavor.

AUTHOR CONTRIBUTIONS

J.E.B. and L.I.L. conceived and coordinated the study. A.U., A.M.I., M.V., and N.L. purified all proteins. M.F. and K.B. provided materials and developed protocols for the biochemical activity and inhibition assays. M.V-H. performed the experiments shown in Fig. 1. L.Q.B. performed the experiments shown in Fig. 2. A.M.I., M.V., and L.Q.B. performed the experiments shown in Fig. 3. L.Q.B and J. Y. performed the experiments shown in Fig. 4. A.U., L.Q.B., M.V., M.V-H., and L.I.L contributed to the analysis and interpretation of the results. A.U., L.Q.B. and L.I.L. wrote the manuscript. All authors approved the final version of the manuscript.

ACKNOWLEDGMENTS

L.I.L. and J.E.B. acknowledge funding from the German Research Foundation (DFG) through Research Training Group 2341 "Microbial Substrate Conversion (MiCon)".

L.I.L. received additional funding through the InnovationsFoRUM Host-Microbe-Interaction IF-018N-22-TP8, and J.E.B. gratefully acknowledges funding from the German federal state of North Rhine-Westphalia for the mass spectrometer (Forschungsgrossgeräte der Länder).

We gratefully acknowledge the excellent support of the technical staff at RUBION, Ruhr University Bochum.

REFERENCES

Aguinagalde, L., Díez-Martínez, R., Yuste, J., Royo, I., Gil, C., Lasa, Í., Martín-Fontecha, M., Marín-Ramos, N. I., Ardanuy, C., Liñares, J., García, P., García, E., and Sánchez-Puelles, J. M. (2015). Auranofin efficacy against MDR *Streptococcus pneumoniae* and *Staphylococcus aureus* infections. J Antimicrob Chemother, 70(9), 2608–2617. https://doi.org/10.1093/jac/dkv163

Albada, H. B., Chiriac, A.-I., Wenzel, M., Penkova, M., Bandow, J. E., Sahl, H.-G., and Metzler-Nolte, N. (2012). Modulating the activity of short arginine-tryptophan containing antibacterial peptides with N-terminal metallocenoyl groups. Beilstein J. Org. Chem, 8(1), 1753–1764. https://doi.org/10.3762/bjoc.8.200

Alderden, R. A., Hall, M. D., and Hambley, T. W. (2006). The Discovery and Development of Cisplatin. J Chem Educ, 83(5), 728. https://doi.org/10.1021/ed083p728

Aller, I., Rouhier, N., and Meyer, A. J. (2013). Development of roGFP2-derived redox probes for measurement of the glutathione redox potential in the cytosol of severely glutathione-deficient rml1 seedlings. Front Plant Sci, 4, 506. https://doi.org/10.3389/fpls.2013.00506

Anagnostopoulos, C. and Spizizen, J. (1961). Requirements for transformation in *Bacillus subtilis*. J. Bacteriol, 81(5), 741–746. https://doi.org/10.1128/jb.81.5.741-746.1961

Angelucci, F., Sayed, A. A., Williams, D. L., Boumis, G., Brunori, M., Dimastrogiovanni, D., Miele, A. E., Pauly, F., and Bellelli, A. (2009). Inhibition of *Schistosoma mansoni* thioredoxin-glutathione reductase by Auranofin. Structural and kinetic aspects. J Biol Chem, 284(42), 28977–28985. https://doi.org/10.1074/jbc.M109.020701

Arnér, E. S. J., and Holmgren, A. (2005). Measurement of Thioredoxin and Thioredoxin Reductase. Curr Protoc Toxicol, 24(1), 7.4.1-7.4.14. https://doi.org/10.1002/0471140856.tx0704s05

Arscott, L. D., Gromer, S., Schirmer, R. H., Becker, K., and Williams, C. H. (1997). The mechanism of thioredoxin reductase from human placenta is similar to the mechanisms of lipoamide dehydrogenase and glutathione reductase and is distinct from the mechanism of thioredoxin reductase from *Escherichia coli*. Proc Natl Acad Sci, 94(8), 3621–3626. https://doi.org/10.1073/pnas.94.8.3621

Aslam, B., Wang, W., Arshad, M. I., Khurshid, M., Muzammil, S., Rasool, M. H., Nisar, M. A., Alvi, R. F., Aslam, M. A., Qamar, M. U., Salamat, M. K. F., and Baloch, Z. (2018). Antibiotic resistance: a rundown of a global crisis. Infect Drug Resist, 11, 1645–1658. https://doi.org/10.2147/IDR.S173867

Augustus, A. M., Celaya, T., Husain, F., Humbard, M., and Misra, R. (2004). Antibiotic-sensitive TolC mutants and their suppressors. J Bacteriol, 186(6), 1851–1860. https://doi.org/10.1128/JB.186.6.1851-1860.2004

Baba, T., Ara, T., Hasegawa, M., Takai, Y., Okumura, Y., Baba, M., Datsenko, K. A., Tomita,
M., Wanner, B. L., and Mori, H. (2006). Construction of *Escherichia coli* K-12 in-frame,
single-gene knockout mutants: the Keio collection. Mol Syst Biol, 2, 2006.0008.
https://doi.org/10.1038/msb4100050

Bandow, J. E., Baker, J. D., Berth, M., Painter, C., Sepulveda, O. J., Clark, K. A., Kilty, I., and
VanBogelen, R. A. (2008). Improved image analysis workflow for 2-D gels enables
large-scale 2-D gel-based proteomics studies - COPD biomarker discovery study.
Proteomics, 8(15), 3030–3041. https://doi.org/10.1002/pmic.200701184

Bian, M., Fan, R., Zhao, S., and Liu, W. (2019). Targeting the Thioredoxin System as a
Strategy for Cancer Therapy. J Med Chem, 62(16), 7309–7321.
https://doi.org/10.1021/acs.jmedchem.8b01595

Butt, T. R., Edavettal, S. C., Hall, J. P., and Mattern, M. R. (2005). SUMO fusion technology
for difficult-to-express proteins. Protein Exp Purif, 43(1), 1–9.
https://doi.org/10.1016/j.pep.2005.03.016

Capparelli, E. V., Bricker-Ford, R., Rogers, M. J., McKerrow, J. H., and Reed, S. L. (2017).
Phase I Clinical Trial Results of Auranofin, a Novel Antiparasitic Agent. Antimicrob
Agents Chemother, 61(1). https://doi.org/10.1128/AAC.01947-16

Caroli, A., Simeoni, S., Lepore, R., Tramontano, A., and Via, A. (2012). Investigation of a
potential mechanism for the inhibition of SmTGR by Auranofin and its implications for
Plasmodium falciparum inhibition. Biochem Biophys Res Commun, 417(1), 576–581.
https://doi.org/10.1016/j.bbrc.2011.12.009

Cassetta, M. I., Marzo, T., Fallani, S., Novelli, A., and Messori, L. (2014). Drug repositioning:
auranofin as a prospective antimicrobial agent for the treatment of severe
staphylococcal infections. Biometals, 27(4), 787–791. https://doi.org/10.1007/s10534-
014-9743-6

Cha, Y., Erez, T., Reynolds, I. J., Kumar, D., Ross, J., Koytiger, G., Kusko, R., Zeskind, B.,
Risso, S., Kagan, E., Papapetropoulos, S., Grossman, I., and Laifenfeld, D. (2018).
Drug repurposing from the perspective of pharmaceutical companies. Br J Pharmacol,
175(2), 168–180. https://doi.org/10.1111/bph.13798

Debnath, A., Ndao, M., and Reed, S. L. (2012). Reprofiled drug targets ancient protozoans:
drug discovery for parasitic diarrheal diseases. Gut Microbes, 4(1), 66–71.
https://doi.org/10.4161/gmic.22596

Dooley, C. T., Dore, T. M., Hanson, G. T., Jackson, W. C., Remington, S. J., and Tsien, R. Y.
(2004). Imaging Dynamic Redox Changes in Mammalian Cells with Green Fluorescent
Protein Indicators. J Biol Chem, 279(21), 22284–22293.
https://doi.org/10.1074/jbc.M312847200

Dos Santos, H. F. (2014). Reactivity of auranofin with S-, Se- and N-containing amino acids.

Comput Theor Chem, 1048, 95–101. https://doi.org/10.1016/j.comptc.2014.09.005

Elkashif, A. and Seleem, M.N. (2020). Investigation of auranofin and gold-containing

analogues antibacterial activity against multidrug-resistant *Neisseria gonorrhoeae*. Sci

Rep, 10(1), 5602. https://doi.org/10.1038/s41598-020-62696-3.

Farha, M. A., and Brown, E. D. (2019). Drug repurposing for antimicrobial discovery. Nat

Microbiol, 4(4), 565–577. https://doi.org/10.1038/s41564-019-0357-1

Finkelstein, A. E., Walz, D. T., Batista, V., Mizraji, M., Roisman, F., and Misher, A. (1976).

Auranofin. New oral gold compound for treatment of rheumatoid arthritis. Ann Rheum

Dis, 35(3), 251–257. https://doi.org/10.1136/ard.35.3.251

Fuchs, B. B., RajaMuthiah, R., Souza, A. C. R., Eatemadpour, S., Rossoni, R. D., Santos, D.

A., Junqueira, J. C., Rice, L. B., and Mylonakis, E. (2016). Inhibition of bacterial and

fungal pathogens by the orphaned drug auranofin. Future Med Chem, 8(2), 117–132.

https://doi.org/10.4155/fmc.15.182

Gon, S., Faulkner, M. J., and Beckwith, J. (2006). *In Vivo* Requirement for Glutaredoxins and

Thioredoxins in the Reduction of the Ribonucleotide Reductases of *Escherichia coli*.

Antioxid Redox Signal, 8(5–6), 735–742. https://doi.org/10.1089/ars.2006.8.735

Gromer, S., Arscott, L. D., Williams, C. H., Schirmer, R. H., and Becker, K. (1998). Human

placenta thioredoxin reductase. Isolation of the selenoenzyme, steady state kinetics,

and inhibition by therapeutic gold compounds. J Biol Chem, 273(32), 20096–20101.

https://doi.org/10.1074/jbc.273.32.20096

Harbut, M. B., Vilchèze, C., Luo, X., Hensler, M. E., Guo, H., Yang, B., Chatterjee, A. K.,

Nizet, V., Jacobs, W. R., Schultz, P. G., and Wang, F. (2015). Auranofin exerts broad-

spectrum bactericidal activities by targeting thiol-redox homeostasis. Proc Natl Acad

Sci, 112(14), 4453–4458. https://doi.org/10.1073/pnas.1504022112

Hokai, Y., Jurkowicz, B., Fernández-Gallardo, J., Zakirkhodjaev, N., Sanaú, M., Muth, T. R.,

and Contel, M. (2014). Auranofin and related heterometallic gold(I)-thiolates as potent

inhibitors of methicillin-resistant *Staphylococcus aureus* bacterial strains. J Inorg

Biochem, 138, 81–88. https://doi.org/10.1016/j.jinorgbio.2014.05.008

Holmgren, A., and Björnstedt, M. (1995). Thioredoxin and thioredoxin reductase. Methods

Enzymol, 252, 199–208. https://doi.org/10.1016/0076-6879(95)52023-6

Humphreys, G., and Fleck, F. (2016). United Nations meeting on antimicrobial resistance.

Bull World Health Organ, 94(9), 638–639. https://doi.org/10.2471/BLT.16.020916

Hutchings, M. I., Truman, A. W., and Wilkinson, B. (2019). Antibiotics: past, present and

866   future. Curr Opin Microbiol, 51, 72–80. https://doi.org/10.1016/j.mib.2019.10.008

Hwang-Bo, H., Jeong, J.-W., Han, M. H., Park, C., Hong, S.-H., Kim, G.-Y., Moon, S.-K.,

Cheong, J., Kim, W.-J., Yoo, Y. H., and Choi, Y. H. (2017). Auranofin, an inhibitor of

thioredoxin reductase, induces apoptosis in hepatocellular carcinoma Hep3B cells by

generation of reactive oxygen species. Gen Physiol Biophys, 36(2), 117–128.

https://doi.org/10.4149/gpb_2016043

Jeon, K.-I., Byun, M.-S., and Jue, D.-M. (2003). Gold compound auranofin inhibits IκB kinase

(IKK) by modifying Cys-179 of IKKβ subunit. Exp Mol Med, 35(2), 61–66.

https://doi.org/10.1038/emm.2003.9

Kabuyama, Y., Kitamura, T., Yamaki, J., Homma, M. K., Kikuchi, S., and Homma, Y. (2008).

Involvement of thioredoxin reductase 1 in the regulation of redox balance and viability of

rheumatoid synovial cells. Biochem Biophys Res Commun, 367(2), 491–496.

https://doi.org/10.1016/j.bbrc.2007.12.178

Keseler, I. M., Mackie, A., Santos-Zavaleta, A., Billington, R., Bonavides-Martínez, C., Caspi,

R., Fulcher, C., Gama-Castro, S., Kothari, A., Krummenacker, M., Latendresse, M.,

Muñiz-Rascado, L., Ong, Q., Paley, S., Peralta-Gil, M., Subhraveti, P., Velázquez-

Ramírez, D. A., Weaver, D., Collado-Vides, J., … Karp, P. D. (2017a). The EcoCyc

database: reflecting new knowledge about *Escherichia coli* K-12. Nucleic Acids Res,

45(D1), D543–D550. https://doi.org/10.1093/nar/gkw1003

Koronakis, V. (2003). TolC - the bacterial exit duct for proteins and drugs. FEBS Letters,

555(1), 66–71. https://doi.org/10.1016/S0014-5793(03)01125-6

Lewis, P. J., and Marston, A. L. (1999). GFP vectors for controlled expression and dual

labelling of protein fusions in *Bacillus subtilis*. Gene, 227(1), 101–109.

https://doi.org/10.1016/S0378-1119(98)00580-0

Lu, J., and Holmgren, A. (2014). The thioredoxin antioxidant system. Free Radic Biol Med,

66, 75–87. https://doi.org/10.1016/j.freeradbiomed.2013.07.036

Lu, J., Vlamis-Gardikas, A., Kandasamy, K., Zhao, R., Gustafsson, T. N., Engstrand, L.,

Hoffner, S., Engman, L., and Holmgren, A. (2013). Inhibition of bacterial thioredoxin

reductase: An antibiotic mechanism targeting bacteria lacking glutathione. FASEB

Journal, 27(4), 1394–1403. https://doi.org/10.1096/fj.12-223305

Martinez, J. L., and Baquero, F. (2000). Mutation Frequencies and Antibiotic Resistance. J.

Antimicrob Agents, 44(7), 1771–1777. https://doi.org/10.1128/AAC.44.7.1771-

1777.2000

Marzo, T., Cirri, D., Pollini, S., Prato, M., Fallani, S., Cassetta, M. I., Novelli, A., Rossolini, G.

M., and Messori, L. (2018). Auranofin and its Analogues Show Potent Antimicrobial

Activity against Multidrug-Resistant Pathogens: Structure–Activity Relationships.

ChemMedChem, 13(22), 2448–2454. https://doi.org/10.1002/cmdc.201800498

Maurice, M. M., Nakamura, H., Gringhuis, S., Okamoto, T., Yoshida, S., Kullmann, F.,
Lechner, S., Van Der Voort, E. A. M., Leow, A., Versendaal, J., Muller-Ladner, U.,
Yodoi, J., Tak, P. P., Breedveld, F. C., and Verweij, C. L. (1999). Expression of the
thioredoxin-thioredoxin reductase system in the inflamed joints of patients with
rheumatoid arthritis. Arthritis Rheum, 42(11), 2430–2439. https://doi.org/10.1002/1529-
0131(199911)42:11<2430::AID-ANR22>3.0.CO;2-6

Messori, L., and Marcon, G. (2004). Gold complexes in the treatment of rheumatoid arthritis.
Met Ions Biol Syst, 41, 279–304. PMID: 15206120.

Meyer, A. J., and Dick, T. P. (2010). Fluorescent Protein-Based Redox Probes. Antioxid
Redox Signal, 13(5), 621–650. https://doi.org/10.1089/ars.2009.2948

Müller, A., Eller, J., Albrecht, F., Prochnow, P., Kuhlmann, K., Bandow, J. E., Slusarenko, A.
J., and Leichert, L. I. O. (2016). Allicin Induces Thiol Stress in Bacteria through S-
allylmercapto Modification of Protein Cysteines. J Biol Chem, 291(22), 11477-11490.
https://doi.org/10.1074/jbc.M115.702308

Müller, A., Hoffmann, J. H., Meyer, H. E., Narberhaus, F., Jakob, U., and Leichert, L. I.
(2013). Nonnative Disulfide Bond Formation Activates the 32-Dependent Heat Shock
Response in *Escherichia coli*. J Bacteriol, 195(12), 2807–2816.
https://doi.org/10.1128/JB.00127-13

Müller, A., Schneider, J. F., Degrossoli, A., Lupilova, N., Dick, T. P., and Leichert, L. I.
(2017). Systematic in vitro assessment of responses of roGFP2-based probes to
physiologically relevant oxidant species. Free Radic Biol Med, 106, 329–338.
https://doi.org/10.1016/j.freeradbiomed.2017.02.044

Munita, J. M., and Arias, C. A. (2016). Mechanisms of Antibiotic Resistance. Microbiol
Spectr., 4(2), 1–37. https://doi.org/10.1128/microbiolspec.VMBF-0016-
2015.Mechanisms

Neidhardt, F. C., Bloch, P. L., and Smith, D. F. (1974). Culture medium for enterobacteria. J
Bacteriol, 119(3), 736–747. https://doi.org/10.1128/jb.119.3.736-747.1974

Owings, J. P., McNair, N. N., Mui, Y. F., Gustafsson, T. N., Holmgren, A., Contel, M.,
Goldberg, J. B., and Mead, J. R. (2016). Auranofin and N- heterocyclic carbene gold-
analogs are potent inhibitors of the bacteria *Helicobacter pylori.* FEMS Microbiol Lett,
363(14), fnw148. https://doi.org/10.1093/femsle/fnw148

Pickering, I. J., Cheng, Q., Rengifo, E. M., Nehzati, S., Dolgova, N. V., Kroll, T., Sokaras, D.,
George, G. N., & Arnér, E. S. J. (2020). Direct Observation of Methylmercury and
Auranofin Binding to Selenocysteine in Thioredoxin Reductase. Inorg. Chem., 59(5),
2711–2718. https://doi.org/10.1021/acs.inorgchem.9b03072

Rabilloud, T., Strub, J.-M., Luche, S., Dorsselaer, A. van, and Lunardi, J. (2001). A

comparison between Sypro Ruby and ruthenium II tris (bathophenanthroline disulfonate)

as fluorescent stains for protein detection in gels. Proteomics, 1(5), 699–704.

https://doi.org/10.1002/1615-9861(200104)1:5<699::AID-PROT699>3.0.CO;2-C

Rackham, O., Shearwood, A.-M. J., Thyer, R., McNamara, E., Davies, S. M. K., Callus, B. A.,

Miranda-Vizuete, A., Berners-Price, S. J., Cheng, Q., Arnér, E. S. J., and Filipovska, A.

(2011). Substrate and inhibitor specificities differ between human cytosolic and

mitochondrial thioredoxin reductases: Implications for development of specific inhibitors.

Free Radic Biol Med, 50(6), 689–699.

https://doi.org/10.1016/j.freeradbiomed.2010.12.015

Rigobello, M. P., Folda, A., Baldoin, M. C., Scutari, G., and Bindoli, A. (2005). Effect of

Auranofin on the mitochondrial generation of hydrogen peroxide. Role of thioredoxin

reductase. Free Radic Res, 39(7), 687–695.

https://doi.org/10.1080/10715760500135391

Rigobello, M. P., Scutari, G., Folda, A., and Bindoli, A. (2004). Mitochondrial thioredoxin

reductase inhibition by gold(I) compounds and concurrent stimulation of permeability

transition and release of cytochrome c. Biochem Pharmacol, 67(4), 689–696.

https://doi.org/10.1016/j.bcp.2003.09.038

Saccoccia, F., Angelucci, F., Boumis, G., Brunori, M., Miele, A. E., Williams, D. L., and

Bellelli, A. (2012). On the mechanism and rate of gold incorporation into thiol-dependent

flavoreductases. J Inorg Biochem, 108, 105–111.

https://doi.org/10.1016/j.jinorgbio.2011.11.005

Scharf, C., Riethdorf, S., Ernst, H., Engelmann, S., Völker, U., and Hecker, M. (1998).

Thioredoxin is an essential protein induced by multiple stresses in Bacillus subtilis. J

Bacteriol, 180(7), 1869–1877. https://doi.org/10.1128/JB.180.7.1869-1877.1998

Senges, C. H. R., Stepanek, J. J., Wenzel, M., Raatschen, N., Ay, Ü., Märtens, Y.,

Prochnow, P., Hernández, M. V., Yayci, A., Schubert, B., Janzing, N. B. M., Warmuth,

H. L., Kozik, M., Bongard, J., Alumasa, J. N., Albada, B., Penkova, M., Lukežič, T.,

Sorto, N. A., … Bandow, J. E. (2020). Comparison of Proteomic Responses as Global

Approach to Antibiotic Mechanism of Action Elucidation. Antimicrob Agents Chemother,

65(1), e01373-20. https://doi.org/10.1128/AAC.01373-20

Storz, G., and Tartaglia, L. A. (1992). OxyR: a regulator of antioxidant genes. J Nutr, 122(3

Suppl), 627–630. https://doi.org/10.1093/jn/122.suppl_3.627

Storz, G., Tartaglia, L. A., and Ames, B. N. (1990). The OxyR regulon. Anton Leeuw Int J G,

58(3), 157–161. http://doi.org/10.1007/BF00548927

Stulke, J., Hanschke, R., and Hecker, M. (1993). Temporal activation of β-glucanase
synthesis in Bacillus subtilis is mediated by the GTP pool. J Gen Microbiol, 139(9),
2041–2045. https://doi.org/10.1099/00221287-139-9-2041

Thangamani, S., Mohammad, H., Abushahba, M. F. N., Sobreira, T. J. P., Hedrick, V. E.,
Paul, L. N., and Seleem, M. N. (2016). Antibacterial activity and mechanism of action of
auranofin against multi-drug resistant bacterial pathogens. Sci Rep, 6, 22571.
https://doi.org/10.1038/srep22571

Uziel, O., Borovok, I., Schreiber, R., Cohen, G., and Aharonowitz, Y. (2004). Transcriptional
regulation of the *Staphylococcus aureus* thioredoxin and thioredoxin reductase genes in
response to oxygen and disulfide stress. J Bacteriol, 186(2), 326–334.
https://doi.org/10.1128/jb.186.2.326-334.2004

Wang, H.-H., Su, C.-H., Wu, Y.-J., Lin, C.-A. J., Lee, C.-H., Shen, J.-L., Chan, W.-H., Chang,
W. H., and Yeh, H.-I. (2012). Application of Gold in Biomedicine: Past, Present and
Future. Int J Gerontol, 6(1), 1–4. https://doi.org/10.1016/j.ijge.2011.09.015

Wenzel, M., Patra, M., Albrecht, D., Chen, D. Y.-K., Nicolaou, K. C., Metzler-Nolte, N., and
Bandow, J. E. (2011). Proteomic signature of fatty acid biosynthesis inhibition available
for in vivo mechanism-of-action studies. Antimicrob Agents Chemother, 55(6), 2590–
2596. https://doi.org/10.1128/AAC.00078-11

Wenzel, M., Patra, M., Senges, C. H. R., Ott, I., Stepanek, J. J., Pinto, A., Prochnow, P.,
Vuong, C., Langklotz, S., Metzler-Nolte, N., and Bandow, J. E. (2013). Analysis of the
mechanism of action of potent antibacterial hetero-tri-organometallic compounds: a
structurally new class of antibiotics. ACS Chem Biol, 8(7), 1442–1450.
https://doi.org/10.1021/cb4000844

 SUPPLEMENTARY DATA

1001 **Supplemental Table S1: Oligonucleotides used in this study.**

Oligonucleotide	Sequence (5′ → 3′)
AI001_for	GAACAGATTGGAGGTGGCACGACCAAAC
AI002_rev	GCGACTCGAGTTATTTTGCGTCAGCTAAAC
AI003_SUMO_for	GCGTCATATGGGTCATCACCATCATC
AI004_SUMO_rev	GGTCGTGCCACCTCCAATCTGTTC
MM001_for	AACAGATTGGAGGTGTGTCAGAAGAAAAAATTTAT GACGTG
MM002_rev	GCGCCTCGAGTTATTTTAAGGTTTTCAGCGTTTCTTG GAGC
MM004_SUMO_rev	TTTTTCTTCTGACACACCTCCAATCTGTTCGCGGTGAG
MV001_for	CATCATATGAGCGATAAAATTATTCACC
MV002_rev	GAACTCGAGTTACGCCAGGTTAGCGTCGAGG
LQB001_for	AAACATATGATGGCTATCGTAAAAGCAA
LQB002_rev	AAAGGATCCTTAAAGATGTTTGTTTACAA
JY001_for	AAAGGTACCATGGTTAGCAAAGGC
JY002_rev	AAAAAGCTTTTACTTGTACAGTTCATCCATG

1002
1003
Supplemental Table S2: List of marker protein spots induced after treatment with Auranofin in
***E. coli* MG1655.** Bold proteins met criteria for MS identification, proteins denoted with (excl.) were
matched by the search engine, but were excluded, as they did not meet the criteria for Peptides (at least
3), Fragments (at least 5), Coverage % (at least 10), or gel position was inconsistent with theoretical
MW or pI. N/A: the search engine did not produce a match. EcoCyc Accession Number and function
obtained from the EcoCyc database (Keseler et al., 2017). Peptides: number of peptides annotated by
the search engine, Fragments: number of product ions annotated by the search engine. Coverage:
coverage of the protein sequence by the annotated peptides in %.
1012

Spot ID	Regulation factor	Protein	EcoCyc Accession Number	Function	theoretical MW [Da]	theoretical pI	Peptides	Fragments	Coverage %
25	3.2	**SucA**	EG10979	2-oxoglutarate decarboxylase, thiamine-requiring	104995	6.0	15	81	18.3
44	4.1	**HtpG**	EG10461	chaperone protein HtpG	71378	4.9	39	296	42.0
		CsiR (excl.)	EG12386	DNA-binding transcriptional repressor GlaR	24974	6.2	5	15	18.2
		YciH (excl.)	EG11128	putative translation factor	11389	9.8	4	13	26.9
		AroK (excl.)	EG10081	shikimate kinase 1	19526	5.1	2	5	17.3
239	2.0	**RpiA**	EG11443	ribose-5-phosphate isomerase A	22845	5.0	4	21	17.4
326	2.4	EptA (excl.)	EG11613	phosphoethanolamine transferase EptA	61627	6.4	2	9	9.5
328	2.4	**MoaB**	EG11596	protein MoaB	18653	5.7	6	83	25.3
347	3.2	**Bfr**	EG10113	bacterioferritin	18483	4.5	3	18	11.4

Spot ID	Regulation factor	Protein	EcoCyc Accession Number	Function	theoretical MW [Da]	theoretical pI	Peptides	Fragments	Coverage %
357	2.3	N/A							
374	2.3	FtsK (excl.)	G6464	cell division DNA translocase FtsK	146571	4.7	5	25	2.5
497	2.3	**AhpC**	EG11384	alkyl hydroperoxide reductase, AhpC component	20748	4.8	5	32	21.9
537	3.7	N/A							
606	4.4	YbcJ (excl.)	EG12879	putative RNA-binding protein YbcJ	7384	8.3	3	19	4.3
		XapR (excl.)	EG11146	DNA-binding transcriptional activator XapR	33605	9.1	2	7	12.2
620	2.7	**MglB**	EG10593	D-galactose/methyl-galactoside ABC transporter periplasmic binding protein	35690	5.6	23	156	50.3
638	2.7	N/A							
662	3.9	**YeaG**	G6969	protein kinase YeaG	74433	5.5	25	160	32.3
690	2.1	**CysN**	EG10194	sulfate adenylyltransferase subunit 1	52526	4.8	45	336	47.8
		GpmM	EG12296	2,3-bisphosphoglycerate-independent phosphoglycerate mutase	56158	5.0	8	35	13.6
		YedJ (excl.)	EG12710	putative HD superfamily phosphohydrolase YedJ	25888	5.9	4	14	17.3
692	2.6	**GpmM**	EG12296	2,3-bisphosphoglycerate-independent phosphoglycerate mutase	56158	5.0	25	168	21.6
		IhfA (excl.)	EG10440	integration host factor subunit α	11346	9.9	2	7	19.2
718	6.1	**Pgk**	EG10703	phosphoglycerate kinase	41092	4.9	7	41	17.6
752	7.0	CueO (excl.)	EG12318	multicopper oxidase CueO	56519	6.3	5	46	9.7
		Php (excl.)	G7731	putative hydrolase	32894	5.1	4	15	20.9
754	7.2	**Lpd**	EG10543	lipoamide dehydrogenase	50656	5.7	10	93	25.7
		MalI (excl.)	EG10557	DNA-binding transcriptional repressor MalI	36602	6.5	7	20	28.1
756	4.0	**Lpd**	EG10543	lipoamide dehydrogenase	50656	5.7	10	64	28.5
		Amn	EG10039	AMP nucleosidase	53961	5.9	11	49	19.8
		AtpA	EG10098	ATP synthase F1 complex subunit α	55187	5.7	6	38	10.7
757	4.1	Tcyj (excl.)	G7039	cystine ABC transporter periplasmic binding protein	29021	6.2	4	21	12.0
		MepM (excl.)	EG10013	peptidoglycan endopeptidase MepM	49027	9.8	6	20	13.9
758	15.0	**AtpA**	EG10098	ATP synthase F1 complex subunit α	55187	5.7	12	74	18.7
		Amn (excl.)	EG10039	AMP nucleosidase	53961	5.9	7	31	9.1
763	4.6	**Amn**	EG10039	AMP nucleosidase	53961	5.9	8	39	14.9
		AtpA	EG10098	ATP synthase F1 complex subunit α	55187	5.7	13	64	21.8
		CueO	EG12318	multicopper oxidase CueO	56519	6.3	3	25	11.0
		IbpB (excl.)	EG11535	small heat shock protein IbpB	16083	5.0	4	12	19.7

Spot ID	Regulation factor	Protein	EcoCyc Accession Number	Function	theoretical MW [Da]	theoretical pI	Peptides	Fragments	Coverage %
764	4.9	AtpA (excl.)	EG10098	ATP synthase F1 complex subunit α	55187	5.7	6	25	9.1
		ArgA (excl.)	EG10063	N-acetylglutamate synthase	49164	6.1	5	20	8.5
		PyrC (excl.)	EG10806	dihydroorotase	38802	5.8	3	14	8.6
782	4.0	**Gor**	EG10412	glutathione reductase	48741	5.6	16	91	35.1
		Mpl	EG12440	UDP-N-acetylmuramate—L-alanyl-γ-D-glutamyl-meso-2,6-diaminoheptanedioate ligase	49842	5.5	13	57	19.9
		GlmM	EG11553	phosphoglucosamine mutase	47513	5.6	9	50	20.0
		PhoA	EG10727	alkaline phosphatase	49408	5.7	11	45	31.2
806	3.9	**NemA**	G6890	N-ethylmaleimide reductase	39492	5.8	7	52	19.7
		YaiI (excl.)	G6230	DUF188 domain-containing protein YaiI	16958	5.4	5	18	22.4
816	3.0	**PyrC**	EG10806	dihydroorotase	38802	5.7	17	131	21.8
		YqhD	G7564	NADPH-dependent aldehyde reductase YqhD	42070	5.7	11	65	21.7
852	3.0	YoaA (excl.)	G6992	ATP-dependent DNA helicase YoaA	70333	6.7	5	21	6.4
		PstS (excl.)	EG10734	phosphate ABC transporter periplasmic binding protein	37001	9.0	3	12	9.0
		DnaE (excl.)	EG10238	DNA polymerase III subunit α	129822	5.0	5	22	3.2
		PaaE (excl.)	G6713	phenylacetyl-CoA 1,2-epoxidase, reductase subunit	39294	5.9	3	11	4.8
859	3.6	**PykF**	EG10804	pyruvate kinase 1	50697	5.7	10	61	17.2
		AhpF (excl.)	EG11385	alkyl hydroperoxide reductase, AhpF component	56142	5.3	3	17	5.8
860	3.6	**AhpF**	EG11385	alkyl hydroperoxide reductase, AhpF component	56142	5.3	22	168	36.5
		LeuA	EG11226	2-isopropylmalate synthase	57262	5.4	19	126	31.5
		PyrF (excl.)	EG10809	orotidine-5'-phosphate decarboxylase	26333	5.8	4	14	24.9

1013