Temperature dependence of mosquitoes: comparing mechanistic and machine learning approaches

Tejas S. Athni[1,2], Marissa L. Childs[3,4], Caroline K. Glidden[2,5], Erin A. Mordecai[2]

[1] Harvard Medical School, Boston, MA
[2] Department of Biology, Stanford University, Stanford, CA
[3] Emmett Interdisciplinary Program in Environment and Resources, Stanford University, Stanford, CA
[4] Center for the Environment, Harvard University, Cambridge, MA
[5] Stanford Institute for Human-centered Artificial Intelligence, Stanford University, Stanford, CA

Abstract

Mosquito vectors of pathogens (e.g., *Aedes*, *Anopheles*, and *Culex* spp. which transmit dengue, Zika, chikungunya, West Nile, malaria, and others) are of increasing concern for global public health. These vectors are geographically shifting under climate and other anthropogenic changes. As small-bodied ectotherms, mosquitoes are strongly affected by temperature, which causes unimodal responses in mosquito life history traits (e.g., biting rate, adult mortality rate, mosquito development rate, and probability of egg-to-adult survival) that exhibit upper and lower thermal limits and intermediate thermal optima in laboratory studies. However, it remains unknown how mosquito thermal responses measured in laboratory experiments relate to the realized thermal responses of mosquitoes in the field. To address this gap, we leverage thousands of global mosquito occurrences and geospatial satellite data at high spatial resolution to construct machine-learning based species distribution models, from which vector thermal responses are estimated. We apply methods to restrict models to the relevant mosquito activity season and to conduct ecologically-plausible spatial background sampling centered around ecoregions for comparison to mosquito occurrence records. We found that thermal minima estimated from laboratory studies were highly correlated with those from the species distributions ($r = 0.90$). The thermal optima were less strongly correlated ($r = 0.69$). For most species, we did not detect thermal maxima from their observed distributions so were unable to compare to laboratory-based estimates. The results suggest that laboratory studies have the potential to be highly transportable to predicting lower thermal limits and thermal optima of mosquitoes in the field. At the same time, lab-based models likely capture physiological limits on mosquito persistence at high temperatures that are not apparent from field-based observational studies but may critically determine mosquito responses to climate warming.

INTRODUCTION

Mosquito-borne diseases (e.g., malaria, dengue, Zika, chikungunya, West Nile, yellow fever) are responsible for a significant worldwide burden of infectious disease and represent a major threat to global public health (Bhatt et al., 2013; Gaythorpe et al., 2021; Murray et al., 2012; Puntasecca et al., 2021; Stanaway et al., 2016). As small-bodied ectotherms, mosquito vectors are sensitive to environmental conditions and, in particular, to abiotic factors such as temperature (Mordecai et al., 2019; Paaijmans et al., 2013). Climate change is likely to alter the climatic and habitat suitability for and the geographic distribution of mosquitoes, in turn affecting the distribution of pathogens they transmit (Kraemer et al., 2019; Messina et al., 2019; Parham & Michael, 2010; Rochlin et al., 2013). Understanding the limitations that temperature and other ecological conditions place on mosquito vectors is critically important for predicting how mosquitoes and vector-borne diseases will respond to climate change.

Previous laboratory experiments have measured the relationship between temperature and mosquito life history traits such as biting rate, adult mortality rate, mosquito development rate, and probability of egg-to-adult survival for many vector species (Focks et al., 1993; Rueda et al., 1990; Shapiro et al., 2017; Tun-Lin et al., 2000). Together, these trait thermal performance relationships can provide a mechanistic estimate of equilibrium mosquito abundance as a function of temperature (Mordecai et al., 2013, 2019; Parham & Michael, 2010). For mosquitoes for which these relationships have been estimated, the thermal response curves of individual traits follow predictable patterns: they decline to zero at lower thermal minima and upper thermal maxima and peak at intermediate thermal optima, as expected from first principles of physiology and enzyme kinetics (Amarasekare & Savage, 2012; Huey & Berrigan, 2001; Mordecai et al., 2019). In aggregate, modeled population-level mosquito abundance has a similar unimodal relationship with temperature (Mordecai et al., 2013), reflecting the underlying life history traits (Parham & Michael, 2010). These studies confirm a core tenet of the metabolic theory of ecology, which states that organismal physiology operates within and is restricted by thermal limits (Brown et al., 2004). Temperature-dependent effects of these life history traits in turn affect transmission of mosquito-borne disease (Craig et al., 1999; Johansson et al., 2014; Kushmaro et al., 2015; Lafferty, 2009; Liu-Helmersson et al., 2014; Mordecai et al., 2013, 2017; Morin et al., 2015; Paull et al., 2017; Shocket et al., 2018, 2020; Tesla et al., 2018; Villena et al., 2022; Vogels et al., 2017; Wesolowski et al., 2015). Despite these clear predictions from

ectotherm physiology and laboratory thermal performance experiments, it remains unknown to

what extent these experimental, laboratory-based mosquito thermal responses predict the realized

thermal responses of mosquito populations in the field. In particular: (1) is temperature an

important predictor of mosquito distributions? If so, (2) are effects of temperature on mosquito

occurrence nonlinear? and if so, (3) do the predicted thermal optima and limits from laboratory

experiments align with thermal optima and limits measured in the field?

The rapid rise of biodiversity data, machine learning, and open source satellite imagery in

the past decade has enabled new types of approaches necessary to address these questions.

Species distribution models offer a way to connect data on species occurrences with

environmental covariates to understand how environmental conditions influence the probability

of a species occurring in a given location (i.e., habitat suitability). Due to the challenges of

accurately detecting a species' absence, species distribution modeling typically compares

locations in which a species has previously been identified (occurrences) and an artificially-

generated set of background points (pseudo-absences) which approximate the potentially-

habitable and accessible area for a given species (Barve et al., 2011; Elith & Leathwick, 2009;

Peterson & Soberón, 2012; Soberon & Peterson, 2005). Species distribution models have been

used to characterize the ecological niches (i.e., habitat and climatic requirements) of organisms

across the tree of life, including elephants in South Asia (Kanagaraj et al., 2019), lynxes in

Canada (Peers et al., 2012), ants in Australia (Leahy et al., 2020), and rare plants in California

(Gogol-Prokurat, 2011). Beyond predicting geographic ranges, these models can indicate which

environmental covariates are most important for predicting species occurrences, and the

functional relationships between environmental conditions and species distributions.

Species distribution models often rely on remotely sensed satellite imagery to provide

information about biotic, abiotic, and anthropogenic variables. Recent advances in storage and

processing of remotely-sensed imagery, including publicly-available platforms like Google Earth

Engine (Gorelick et al., 2017), have increased the resolution and spatial extent over which these

models can feasibly be run. In parallel, the Global Biodiversity Information Facility (GBIF)

leverages crowdsourced data collection and digitized biodiversity data, where occurrence points

of many species are compiled within a central repository (Edwards, 2004). Together, these

innovations provide ripe new avenues and abundant data through which to tackle ecological

questions.

We focus on the vectors of the world's highest-burden mosquito-borne diseases (malaria,

dengue, chikungunya, Zika, West Nile, and other arboviruses)—*Aedes aegypti, Ae. albopictus,*

Anopheles gambiae, An. stephensi, Culex pipiens, Cx. quinquefasciatus, and *Cx. tarsalis*—each

of which has well-characterized thermal performances curves in laboratory settings and a wealth

of data on occurrences in the field (Kraemer et al., 2015; Messina et al., 2015, 2019; Mordecai et

al., 2013, 2017, 2019; Shocket et al., 2018). *Ae. aegypti* and *Ae. albopictus* are important vectors

of dengue, chikungunya, yellow fever, and Zika viruses. *An. gambiae* and *An. stephensi*

primarily vector the *Plasmodium* spp. protozoans that cause malaria. *Cx. pipiens, Cx.*

quinquefasciatus, and *Cx. tarsalis* transmit West Nile, St. Louis encephalitis, Western Equine

encephalitis, and other zoonotic viruses.

While previous mosquito species distribution models have focused on estimating habitat

suitability and predicting occurrence (Kraemer et al., 2019; Messina et al., 2015, 2019), our goal

is different: to specifically dissect the relationship between temperature and probability of

mosquito occurrence, while incorporating other drivers of occurrence, and to compare this

relationship among mosquito species and with laboratory-based model predictions. For this

reason we created new models, rather than using published ones, that use consistent methods

across multiple mosquito species. Further, we aimed to construct species distribution models for

each focal species using spatially- and temporally-similar data sources and consistent model

assumptions in order to compare thermal dependence across species.

Toward these objectives, this project leverages new computational and methodological

advances to address a long-standing question in disease ecology: does temperature predict

realized mosquito distributions in the field, and if so, what are the functional relationships? We

aim to compare the insights from a 'top-down' machine learning and statistical modeling

approach to results from 'bottom-up' models of mosquito population sizes from trait-based

experiments in the lab (Mordecai et al., 2019) to discern whether and how results from the lab

setting can translate into the real world. We make two important methodological advances to

ensure that our species distribution models capture true thermal limits on mosquito occurrence,

rather than sampling bias or restricted geographic ranges that covary with temperature. First, we

sample background pseudo-absence points only from the set of ecoregions in which the focal

mosquitoes occurred, as well as adjacent ecoregions. This ensures that comparator pseudo-

absence points are selected from regions where the focal mosquito species could have

realistically been found but were not, or in other words, zones that are ecologically plausible for

an occurrence. Second, we restrict temperature measurements to an 'activity season' during

which each mosquito species is blood-feeding and reproducing, and not in dormancy or torpor.

This captures temperature constraints within a physiologically realistic period, rather than

constraints due to overwintering or drought persistence, which are less likely to be comparable to

laboratory-based trait thermal response experiments. By using global mosquito occurrences from

2000-2019, geospatial satellite-derived covariates at high spatial and temporal resolution, bias-

reducing methods described above, and a gradient-boosted classification tree machine learning

algorithm well-suited for prediction tasks, we aim to provide evidence that spans across

continents and decades for the globally important vectors of human malaria and arboviral

disease. With a greater understanding of thermal constraints on vector species occurrence, we

can improve the ability to accurately project and mitigate the future impacts of climate change on

mosquito-borne disease distributions.

METHODS

Overview and study period

We focused on 2000-2019 to isolate recent land-use and climate patterns but provide a

long enough time period to capture mosquitoes' stable spatial distribution. The environmental

covariates and mosquito occurrences were both extracted for this time period, providing

temporal consistency in the analysis (Wisz et al., 2013). An overview of the methods can be seen

below (Fig. 1). Computation was performed in R v4.1.1, Google Earth Engine and the Sherlock

computing cluster at Stanford University.

a) species occurrence and background point selection

b) model covariates

c) critical thermal limits

154
Figure 1. Methods for comparing mosquito thermal performance derived from species distribution
models (a-c) and laboratory trait-based models (c). The analysis involved two major steps: statistical
modeling of mosquito occurrence using global species occurrences (a) and geospatial data (b) to identify
the relationship between temperature and species occurrence (solid line, c), and comparison with
mosquito abundance curves previously derived from laboratory life history traits (Mordecai et al., 2019)
(dashed line, c). Temperature-dependent mosquito abundance (M(T); dashed line in c) is modeled as a
function of temperature-dependent eggs laid per female mosquito per day (EFD(T)), probability of egg-
to-adult survival (pEA(T)), development rate from egg to adult (MDR(T) = 1/development time in days),
and daily adult mosquito mortality rate (μ(T)). Trait curves adapted from Mordecai et al., 2019 and
Villena et al., 2022; M(T) equation originally derived from Mordecai et al., 2013. Thermal minima and
optima of the probability of occurrence from the species distribution models were identified by
calculating the temperature at which the empirical derivative was first positive and stayed positive for the
next step (light blue point, c) and the temperature where the empirical derivative was zero and the
probability of occurrence was at its maximum (light orange, c). Sources of data for the stack of covariates
are described in Table S2.
170
171 **Environmental predictors**

172    Environmental covariates were defined *a priori* in order to avoid data-dredging (Rochlin

et al., 2013), and were based on a literature search of previous species distribution models.

Variables were informed by ecological and biological relevance, expectation to play a role in

mosquito vector ecology, and top contributor status in previous mosquito species distribution

models (Akpan et al., 2018; Conley et al., 2014; Fatima et al., 2016; Mweya et al., 2013).

Environmental covariates were computed using remotely sensed and reanalysis data, and

resampled using bilinear interpolation to consistent 1 km x 1 km resolution on Google Earth

Engine. This approach using global satellite data avoids the spatial interpolation limitations of

common climatic datasets such as WorldClim, in which data are sourced from ground

meteorological stations with uneven geographic distribution and nonrepresentative selection.

In order to verify minimal overlap between predictor variables, a pairwise correlation

analysis was performed with a Pearson's correlation coefficient $r < |0.8|$ threshold (Fig. S1).

Annual averages for covariates were computed across the study period, providing the

characteristic climatic conditions across the study's temporal range.

Temperature predictors

While many species are active year-round (Table 1), numerous species physiologically

undergo photoperiodic diapause in the winter months when ambient light is low in order to

conserve energy for life sustenance (Zhang et al., 2019) or aestivate during the dry season when

no breeding habitat is available (Lehmann et al., 2010). To capture the temperature that

mosquitoes experience while active, temperature covariates were calculated over the activity

season.

For the purposes of this study, the photoperiodic activity season was defined as the time

period encompassing days with at least 9 hours of sunlight (Fig. S2). This definition of the

activity season was chosen based on a variable separate from temperature and on laboratory

studies that found that either 8 or 9 hours of light can induce diapause in mosquitoes in the

absence of temperature change (Poelchau et al., 2013; Rinehart et al., 2006; Robich & Denlinger,

2005). Other tropical mosquito species (e.g., *Anopheles gambiae*) do not enter photoperiodic

dormancy due to their equatorial setting. Rather, these species' activity seasons are instead

constrained by lack of precipitation, when insufficient habitat is available for immature

development and survival. Studies have highlighted how Sahelian *Anopheles gambiae*

mosquitoes virtually disappear in the dry season then re-establish in the wet season through

aestivation behavior (Lehmann et al., 2010; Yaro et al., 2012), and climatological studies in the

Democratic Republic of the Congo characterize the dry season as having less than 50 mm of rain

in the past month (Kazadi & Kaoru, 1996). Precipitation activity season is therefore defined in

our study as the time period in which there is, on average, at least 50 mm of precipitation in the

past 30 days (Fig. S3).

Predictors capturing thermal central tendency include year round temperature mean,

photoperiod activity season temperature mean, and precipitation activity season temperature

mean. Measures of seasonal thermal variation include year round temperature standard deviation,

photoperiod activity season temperature standard deviation, and precipitation activity season

temperature standard deviation.

Table 1. Mosquito species by activity season.

Mosquito Species	Activity Season (Reference)	Raw Number of Occurrences	Unique Cell Centroids
Aedes aegypti	Year Round (Chang et al., 2007; Mitchell, 1995)	37,115	9,299
Aedes albopictus [#]	Photoperiod (Lacour et al., 2015; Poelchau et al., 2013)	39,488	8,688
Anopheles gambiae	Precipitation (Lehmann et al., 2010; Yaro et al., 2012)	13,650	903
Anopheles stephensi	Year Round (Sinka et al., 2011)	1,232	358
Culex pipiens	Photoperiod (Rinehart et al., 2006; Robich & Denlinger, 2005)	98,276	1,949
Culex quinquefasciatus	Year Round (Almirón & Brewer, 1996; Meuti et al., 2015)	30,978	2,670
Culex tarsalis	Photoperiod (Buth et al., 1990)	44,496	909

[#] This species enters winter diapause in some regions such as the Northeast US, but not other regions like the
Southeast US.

Other environmental predictors

Environmental variables other than temperature included measures of vegetation cover,

such as enhanced vegetation index mean, enhanced vegetation index standard deviation, and

forest cover percentage, to capture land cover that might affect mosquito habitat suitability

(Table S2, Fig. S4). Human population density and cattle density were included to capture

sources of blood meal, anthropophilic hotspots, and human-seeking behavior. Precipitation of the

driest quarter, precipitation of the wettest quarter, and surface water seasonality were included to

capture hydrology-driven characteristics influencing mosquito breeding site availability or

abundance, while wind speed was included to capture potential aerial dispersal limitation of

mosquitoes. Precipitation of the driest quarter and precipitation of the wettest quarter were

defined as the set of 3 consecutive months with the least and most precipitation, respectively,

across the 20-year study period.

Occurrence and background

Presence occurrences

Focal mosquito species were selected based on global disease burden, availability of life

history trait data from published laboratory studies, and abundance of publicly-accessible species

occurrences, resulting in seven species: *Ae. aegypti, Ae. albopictus, An. gambiae, An. stephensi,*

Cx. pipiens, Cx. quinquefasciatus, and *Cx. tarsalis*. Collectively, these seven species span nearly

all habitable continents (Fig. S5-S11). Species occurrence records for each species were obtained

from the Global Biodiversity Information Facility (GBIF) for the years 2000-2019. Supplemental

occurrence points were obtained from two papers for *An. stephensi* and *An. gambiae,* which had

low sample sizes from the GBIF database (Sinka et al., 2020; Wiebe et al., 2017). Occurrences

explicitly tagged in Africa were discarded from the supplemental *An. stephensi* data, as these are

part of an ongoing species invasion outside of the native range of the species in South Asia

(Sinka et al., 2020), and species distributions models have limitations in detecting environmental

suitability for invasive species as they do not yet occupy their environmental niche and are not

yet in equilibrium with their environment (Barbet-Massin et al., 2018).

Raw datasets were cleaned to remove occurrences with unknown basis of record and

those obtained through fossil records, due to the temporal and geographic uncertainty between

when and where the organism actually existed during its lifetime and its location of fossilization.

Uncertainty in the latitude and longitude coordinates of occurrence points were taken into

account through a two-step process. First, points with a coordinate uncertainty less than 1000

meters—the diameter of our environmental covariate grid cell size—and those points with

missing coordinate uncertainty were selected. Of the points with missing coordinate uncertainty,

we retain only those points with a latitude and a longitude with at least two significant decimal

points (i.e., each hundredth corresponds to approximately 1.11 km). The occurrence data were

then restricted to points that reside on landmasses and not in the ocean, those with non-missing

values for all covariates, and, when relevant, and those that occurred in areas where the species'

activity season length was greater than 0 (i.e., when the precipitation was above a certain value

in the grid cell, i.e., precipitation activity season, or when there was a certain amount of daylight

hours in the grid cell, i.e., photoperiod activity season) (Fig. S2-3). We filtered the occurrences

to the cell centroids of the unique 1 km x 1 km cells in which occurrence points were obtained,

which serves to spatially thin the occurrence points and reduce the effect of biased sampling

(Kramer-Schadt et al., 2013) (Table 1; Table S1).

Pseudo-absence background sampling

Species distribution models work by comparing the environmental covariates that best

predict locations with species occurrences to those in which the species does not occur, to

understand the environmental conditions that constrain a species' distribution. As the majority of

information is on species occurrences rather than species absences, methods for species

distribution modeling have developed to sample pseudo-absences: points that represent the area

in which a species could have been sampled yet was not reported. In order for these sampled

points to serve as plausible points where the species of interest was not found, they should be

within the species' accessible area, or the region that is reasonably reachable by the species

through dispersal or migration over the relevant time period (Barve et al., 2011; Peterson &

Soberón, 2012; Soberon & Peterson, 2005). Choosing pseudo-absences that occur within the

same ecoregions, i.e., geographic regions that possess similar species and community

assemblages, as occurrences provides a way to geographically restrict possible background

points. Moreover, these points should be chosen in a way that takes into account sampling bias

and collection effort.

To sample background points, we first created a bias mask to probability-weight the

sampling. 72 million occurrences from Class Insecta, filtered to the same criteria as mosquito

occurrence points (i.e., <1000m coordinate uncertainty or at least two significant decimal points

of latitude and longitude, 2000-2019 study period, no fossil record or literature points), were

extracted from GBIF. In order to capture the frequency of GBIF sampling conducted in a specific

geographic location, the number of insects were tallied for each 1 km x 1 km grid cell, and these

counts were converted into proportions of all insects. This proportion was regarded as the

probability weight for that given grid cell. To provide adequate sample size while avoiding a

skew of the overall sample towards disproportionately favoring pseudo-absences over

occurrences, we selected a number of unique background cell centroids equal to twice the

number of unique occurrence centroids (Liu et al., 2013). Background cells were sampled at

random without replacement and weighted by the bias mask from the geographical region that

was bounded by the focal species' ecoregions of occurrence. In addition, we include adjacent

ecoregions in the background sampling space to promote contiguity in the potential ecological

spaces from which pseudo-absences can be drawn and to better encapsulate the broad range of

environmental covariates that a given species may experience in nature. Ecoregions were defined

as RESOLVE Ecoregions that intersected with a given species' occurrence centroids, buffered

with a 200 km radius buffer that approximated an upper end of the wind-assisted geographic

mobility range of a mosquito (Chevillon et al., 1995; Verdonschot & Besse-Lototskaya, 2014).

By buffering occurrences and then choosing the set of ecoregions and adjacent ecoregions in

which these buffered points lie, we approximate what is known in ecological theory as the

'accessible area,' described above (Barve et al., 2011; Peterson & Soberón, 2012; Soberon &

Peterson, 2005). Because our focal questions centered on species' thermal breadth, we wanted to

ensure that as broad a temperature range as was reasonable was included in the background

sampling. Specifically, we checked whether our background sampling scheme provided a

temperature distribution at least as wide as that of the occurrences—a consideration important to

ensure that the statistical model can ascertain thermal minima or maxima.

Species distribution modeling

XGBoost model fitting

Raster values for each occurrence and background centroid were extracted from the stack

of covariates for a given species. The data was partitioned into a training and evaluation set,

where 80% of the data was randomly selected for the training set and 20% was randomly

selected for the validation set. We used stratified sampling so that there was an equal proportion

of presence points in each set. We used gradient boosted classification trees to model the training

data, as this class of algorithms fits flexible and nonlinear relationships (including thresholds)

between environmental conditions and probability of species occurrence, identifies interactions

between covariates due to the structure of the trees, and has been successfully used in other

species distribution models (Kraemer et al., 2015; Li et al., 2022; Messina et al., 2019). Gradient

boosted trees are a type of supervised learning algorithm that iteratively fits trees, each time

fitting to the residual errors from the previous predictions, effectively ensembling weaker

learners to accurately predict a target variable (Natekin & Knoll, 2013). We fit models using

eXtreme Gradient Boosting (XGBoost) (Chen et al., 2015) in R. Among the strengths of the

XGBoost algorithm are its rapid computational speed when dealing with large-scale data, ability

to handle class-imbalanced training sets (i.e., the number of observations are not equal across

each level of the outcome variable), and ability to tolerate highly-collinear predictor variables.

To quantify variation around model performance and output, we fit the model with 20 random

train and validation splits.

Bayesian hyperparameter optimization

Gradient boosted classification trees rely on hyperparameters that control how each tree

is fit and how individual trees' estimates are combined. Bayesian optimization was conducted to

tune key hyperparameters initialized with a range of user-inputted values, including learning rate

(range: 0.0001 to 0.3), maximum tree depth (range: 2 to 50), minimum child weight (range: 1 to

50), subsample of observations used in each tree (range: 0.25 to 1), subsample of columns to use

in each tree (range: 0.5 to 1), and minimum split loss (range: 0 to 100) (Chen et al., 2015;

XGBoost Parameters, n.d.). For each round of the Bayesian optimization (up to a maximum of

24 rounds), we use 5-fold cross validation where we split the training data into 5 disjoint sets,

and train the model on each combination of 4 folds while predicting out-of-sample for the fifth

fold. We use the average out-of-sample log loss averaged over all 5 folds to identify the optimal

number of boosting rounds (up to 10,000) for each iteration of the Bayesian optimization. After

all iterations of the Bayesian optimization, we select the set of hyperparameters with the lowest

out-of-sample log loss and train a final model using all 5 folds—representing all of the training

data, which is 80% of the full data set.

Interpreting the model

Receiver operating characteristic (ROC) curves and their area under the curve (AUC)

values, which graphically and numerically illustrate the discrimination ability of a binary

classifier, were computed for model predictions on both the training and evaluation sets. We

were particularly interested in two aspects of the resulting models: the inferred relationship

between the probability of occurrence and temperature, and the relative importance of

temperature in predicting a species distribution, compared to other environmental covariates. To

understand the relationship between temperature and occurrence, we plotted univariate partial

dependence plots for temperature mean and temperature standard deviation. Depending on the

species, we use annual temperature, temperature during the photoperiod-dictated activity season,

or precipitation-dictated activity season. Partial dependence plots depict the mean value of the

probability of occurrence given temperature across models fit to all combinations of other

variables at each value of temperature. To assess the importance of temperature in relation to

other environmental variables, measured in gain (i.e., the increase in accuracy that a given

variable provides when a regression tree branch is split upon that variable) (*XGBoost*

Parameters, n.d.), we computed variable importance scores for the full set of predictors and

compared them with temperature. We computed variable importance and PDPs for each

bootstrapping iterations described above.

365   Mosquito abundance curves from the lab

Physiological life history trait thermal performance curves were collected from published

literature (Mordecai et al., 2019; Shocket et al., 2020; Villena et al., 2022). Briefly, in previously

published work asymmetric responses like mosquito development rate (MDR) were modeled

using a Briere function, and both concave-down symmetric responses like eggs per female per

day (EFD) and probability of egg-to-adult survival (pEA) as well as concave-up symmetric

responses like mortality rate (μ) were modeled using quadratic functions. 5,000 posterior draws

were selected from each distribution, and mosquito abundance M(T) was calculated using

Equation (1), which was originally derived in Mordecai et al., 2013, adapted from Parham and

Michael, 2010. Median and 95% credible intervals were computed for each species' M(T) and

critical thermal limits.

378
$$M(T) \ = \ \frac{EFD(T)\ pEA\ (T)\ MDR(T)}{\mu(T)^2} \qquad\qquad (1)$$

379
380
Comparison of thermal relationships from XGBoost versus lab-based models of mosquito

abundance

We compared thermal minima and optima of the partial dependence plots from the

XGBoost model to the mosquito abundance M(T) curves from the lab trait trajectories. For

species distribution models, thermal minima were identified as the temperature at which the

partial dependence plot began increasing, which we operationalized as the first time the

empirical derivative was positive and stayed positive for the next step in temperature as well

(Fig. 1c). Thermal minima could also correspond to the threshold temperature at which species

probability of occurrence is increasing most rapidly, so we supplementarily identify thermal

minima as the temperature where the empirical derivative is largest (Fig. S14). Thermal optima

were identified as the point where the empirical derivative was zero and the partial dependence

plot was at its maximum (Fig. 1c). We did not identify thermal maxima, as few species had

partial dependence plots that clearly declined after the thermal optima and then reached a lower

plateau that was within the range of observed temperatures. For the mosquito abundance M(T)

curves derived from laboratory studies, to determine the thermal minima, we identified the

temperature at which M(T) exceeded zero for each M(T) curve calculated from posterior draws

and then calculated the median, 2.5th and 97.5th percentiles to produce a central estimate and

confidence interval for thermal minima. For thermal optima, we identified the temperature where

M(T) was highest for each curve, and similarly calculated median and 95% confidence intervals.

We calculate the Pearson correlation between the median species distribution model-based

thermal minima and the median lab-based thermal minima, and repeat for thermal optima to

quantify how well the two methods compare. In particular, because temperature varies

substantially in the field and M(T) curves are based on constant temperature measurements in the

lab, we expect nonlinear averaging to affect the exact thermal limits observed in the field.

RESULTS

The species distribution models predicted the occurrence of all focal species with high

accuracy: discrimination across both training (in-sample) and evaluation (out-of-sample) datasets

produced an area under the receiver operating curve (AUC) above 0.9, where 1 indicates perfect

discrimination of presence and absence and 0.5 is no better than a coin toss (Fig. 2, Fig. S12). Of

the focal species, *An. gambiae* recorded the lowest out-of-sample evaluation AUC of 0.938. The

fact that the AUCs in the evaluation set were comparable to the training set suggests that the

models are learning general patterns, differentiating between training and evaluation sets, and not

overfitting to the training data.

Measures of temperature, i.e., either annual or activity season-limited (as appropriate)

temperature mean and temperature standard deviation, were consistently important predictors of

mosquito occurrence, as measured by gain (Fig. 2). For all species, mean temperature was among

the top six predictors. For *Ae. aegypti* and *Ae. albopictus,* temperature mean was the top

predictor and temperature standard deviation was second (Fig. 2). *An. gambiae, Cx. pipiens,* and

Cx. tarsalis had temperature standard deviation as a more important predictor than temperature

mean (Fig. 2).

Beyond temperature, different predictors were important for different mosquito species.

Forest cover was the most important predictor for *An. stephensi, An. gambiae,* and *Cx. tarsalis*,

and precipitation of the driest quarter was the most important predictor for *Cx. quinquefasciatus*

(Fig. 2). Precipitation variables were among the top-5 predictor variables for all species.

Precipitation of the wettest quarter was among the top-5 predictors for *Ae.aegypti, Ae. albopictus*

and *Cx. pipiens* (Fig. 2). Precipitation of the driest quarter was among the top-5 predictors for

An. stephensi and *Cx. quinquefasciatus*, and both precipitation variables were among the top-5

predictors for *An. gambiae* and *Cx. tarsalis* (Fig. 2) Surface water seasonality, however, was the

least important predictor variable for all species distribution models (Fig. 2).

Figure 2. Temperature mean (red) and standard deviation (blue) were important predictors of mosquito occurrence for all focal species. Variable importance, measured in gain, is shown for each predictor variable by species. Temperature mean variables, including year-round temperature annual mean, photoseason activity season temperature mean, and precipitation activity season temperature mean, are colored in red; only the most biologically appropriate temperature mean variable was included for each species (Table 1). Likewise, temperature standard deviation variables (Table 1) are colored in blue. Other predictors (black bars) include cattle density, enhanced vegetation index mean, enhanced vegetation index standard deviation, forest cover percentage, human population density, precipitation of the driest quarter, precipitation of the wettest quarter, surface water seasonality, and wind speed. Points represent the median importance across the 20 bootstrapped model iterations and lines represent the range over all model iterations. Test and training AUC values are similarly medians and full ranges over the model iterations.

While mechanistic relationships between temperature and mosquito abundance, based on laboratory-derived M(T), were unimodal and hump-shaped in form, the temperature relationships inferred from the statistical models (univariate XGBoost partial dependence plots; PDPs) were typically nonlinear with steep thresholds and plateaus (Fig. 3). Both the mechanistic and statistical models showed steep increases as temperatures exceeded lower thermal limits (Figs. 3). By contrast, while M(T) consistently responded unimodally to temperature, relationships

from XGBoost PDPs mostly only showed lower thermal limits and thermal optima, but not upper

limits (Fig. 3). The difference in functional forms between M(T) and XGBoost PDPs is

consistent with the fact that they each describe distinct processes. The mechanistic model, M(T),

describes a relationship between temperature and mosquito *abundance* that continuously varies

with temperature and the underlying laboratory-measured traits. By contrast, the PDPs from

XGBoost capture the relationship between temperature and mosquito *occurrence probability*,

which we expect to rise and fall sharply and reach a plateau at intermediate, suitable

temperatures. The M(T) curves for all species showed clear thermal minima, optima, and

maxima because the underlying laboratory experiments captured the full range of temperatures at

which mosquito performance is optimized and at which it declines to zero.

467
Figure 3. Comparing mosquito temperature responses from mechanistic laboratory-based
mosquito abundance models and XGBoost statistical models. Mosquito abundance for each species is
estimated as the median M(T) curve with a shaded 95% credible interval band (blue lines and shaded
ribbons). XGBoost covariate responses are univariate partial dependence plots (PDPs) for annual mean
temperature, where mean temperature was bounded by photoperiod or precipitation for a subset of
species, that show the marginal effects of temperature on model prediction (black lines). Each black line
for the PDPs represents one of the 20 model iterations. Both M(T) and PDPSs are scaled to range from 0
to 1. Grey and red density plots in the bottom of each panel show the distribution of observed annual
temperatures for background and occurrence points, respectively.
477

Given that both the laboratory-based mechanistic approach (M(T)) and the statistical

approach (XGBoost PDPs) predicted thermal minima and optima for each species, we asked

whether the results of the two approaches were concordant. The estimates of thermal minima

between M(T) and PDP were highly correlated across species (r = 0.902; Fig. 4; Fig. S13, S14),

suggesting that M(T) captures key species-specific temperature constraints on occurrence in the

field. Mosquito species of the *Aedes* and *Culex* genera exhibited thermal minima that were 2-6°C

cooler in the field than predicted in the laboratory, while *Anopheles* mosquitoes exhibited

thermal minima 0.5°C higher in the field than in laboratory settings (Fig. 4; Fig. S13, S14). *Cx.*

tarsalis and *Cx. pipiens* consistently had the lowest and second-lowest thermal minima,

respectively, and *An. gambiae* consistently had the highest thermal minimum when viewed

across M(T) and PDP (Fig. 4; Fig. S13-S14). The alternative identification of thermal minima as

the threshold temperature with the greatest increase in species occurrence resulted in a similarly

high correlation between M(T) and PDP-based thermal minima (r = 0.897), but resulted in

thermal minima approximately 5°C warmer in the field than predicted by the laboratory for

Anopheles mosquitoes (Fig. S14). Thermal optima were similar, although more weakly

correlated, between M(T) and PDP estimates (r = 0.685), supporting, at least in part, the

conclusion that field based observations capture key components of lab-based non-linear

response to temperature. However, the relation appeared to deviate the most for temperature

species *Cx. pipiens* and *Cx. taralis,* for which thermal optima were 9.5°C and 10°C lower,

respectively, in the field versus laboratory (Fig. 4; Table S3; Fig. S13-S14). We could not

compare thermal maxima because our XGBoost models did not identify thermal maxima for any

focal species.

500
501

502
Figure 4. Mechanistic models (M(T); x-axis) captured species' thermal minima observed in the field
(XGBoost PDPs; y-axis). The dashed diagonal line is the 1:1 line, and mosquito species are colored by
genus. The Pearson correlation (r) was 0.902 for the PDP versus M(T) relationship for average T_{min} and
0.685 for the PDP versus M(T) relationship for average T_{opt}. For mechanistic estimates of T_{min} and T_{opt},
points indicate medians and horizontal line ranges are the 95% CI from the 5000 posterior samples used.
For statistical estimates of T_{min} and T_{opt}, points indicate medians and vertical lines are the full minimum to
maximum range over the 20 bootstrapped model iterations.
510

DISCUSSION

For all seven major vectors of human disease we investigated, species distribution models

captured occurrence probability with high discrimination and accuracy and temperature was an

important predictor (Fig. 2, Fig. S12). In particular, temperature mean and standard deviation

were highly important predictors across all focal species (Fig. 2). This finding echoes the

importance of temperature as a predictor in past models, although many such models define

temperature differently from our study (e.g., temperature suitability, temperature of the hottest

quarter, temperature of the coldest month) (Brady et al., 2013; Cianci et al., 2015; Fatima et al.,

2016; Kraemer et al., 2015). Four species (*Ae. aegypti*, *Ae. albopictus*, *Cx. pipiens,* and *Cx.*

quinquefasciatus) had either temperature mean or standard deviation as their most important

predictor (Fig. 2). Nonlinear relationships with temperature from 'bottom-up' models of

mosquito abundance from lab life-history traits were partially recapitulated in the 'top-down'

statistical models of global-scale occurrence data, particularly at the lower end of the temperature

range , consistent with previous work on other mosquito species (Fatima et al., 2016). In general,

thermal minima and optima were highly comparable across species between the trait-based and

statistical approaches (Fig. 4). However, thermal maxima were not comparable: in particular,

XGBoost did not consistently detect thermal maxima for occurrence in the field, despite

laboratory evidence that mosquito life history traits decline precipitously at high temperatures.

Thermal physiology theory and experiments have established that organismal

performance is constrained to an operative range of temperatures at which key functions can

occur and populations can stably persist (Angilletta & Angilletta, 2009; Bernhardt et al., 2018;

Dell et al., 2011; Deutsch et al., 2008; Somero, 2002). Yet, realized species distributions may not

clearly reflect these thermal constraints if other factors are also important for constraining

distributions, including rainfall, seasonality, biotic interactions, habitat, movement rates, and the

range of temperatures that occur within accessible regions (Buckley et al., 2010; Crozier &

Dwyer, 2006; Davis et al., 1998; Guisan & Thuiller, 2005; Heikkinen et al., 2007; Lounibos &

Juliano, 2018; Pulliam, 2000; Wisz et al., 2013). For example, aridity may constrain mosquito

distributions beyond direct high-temperature constraints. Likewise, average temperatures across

activity seasons may not reach levels high enough to exclude mosquito persistence (Bernhardt et

al., 2018; Lambrechts et al., 2011; Paaijmans et al., 2009, 2010, 2013). Alternatively, more

extreme temperature variation at low or high means can limit organismal performance, and even

moderate variation can reduce performance near optimal temperatures (i.e., where M(T) curves

are concave-down; Fig. 3). The impact of temperature variation could explain the observation

that for species with cooler thermal minima and optima, estimates at constant temperatures in the

lab were up to 10°C warmer than estimates in the field (Fig. 4).

This highlights the importance of climate change projections that consider both field-

based estimates of thermal and other constraints on species distributions and more mechanistic

estimates, particularly for thermal optima and maxima, which may be difficult to observe in the

field under current and historical conditions (Gamliel et al., 2020). Simply projecting species

distribution models like those derived here under future climate change scenarios is likely to

overlook the potential for warming temperatures to exceed thermal optima and limits for species

persistence. Moreover, it is critical for studies of vector-borne disease transmission under future

climate change to account for the gap between potential and realized future vector thermal

niches, as vectors may not immediately track geographic shifts in climate suitability. Approaches

that combine mechanistic thermal performance information (e.g., from laboratory studies and life

history models) with statistical inference based on current distributions are most likely to

accurately capture current and future environmental constraints on species persistence, including

for noxious species like disease vectors.

This study incorporates several methodological innovations from previously published

species distribution modeling methods for mosquitoes, particularly focused on estimating

temperature bounds, with applications for climate change models. First, we computed covariates

aligned to the time of collection of our mosquito occurrence data, in contrast to numerous studies

that use climatologies estimated from the years preceding the occurrence point sampling time

frame (Al Ahmed et al., 2015; Conley et al., 2014; Kraemer et al., 2015; Messina et al., 2019;

Mweya et al., 2013; Richman et al., 2018). Second, given our explicit goal of estimating thermal

limits, we restricted our measures of average temperature to the relevant activity season of each

focal species (Smeraldo et al., 2018), which ensures that we are using the temperature range of

the mosquito in the field when that species is active and non-dormant (and therefore comparable

to laboratory experimental measurements). Third, we selected background points from the

ecoregions in which a species occurs plus a 200 km buffer, as well as adjacent ecoregions, to

ensure that we are estimating temperature limitations based on regions that are plausible based

on habitat suitability. This method, grounded in ecological theory (Barve et al., 2011; Soberón,

2010; Soberon & Peterson, 2005) and similar to background limitation methods for non-

mosquito species in the literature (Gogol-Prokurat, 2011; Khoury et al., 2020; Lira-Noriega et

al., 2018; Mertens et al., 2021; Nuñez-Penichet et al., 2019), functionally decouples habitat and

temperature. This is largely a new advancement for mosquito species distribution modeling.

Finally, while most mosquito species distribution modeling studies focus on identifying

predictors of one or two mosquito species, here we explicitly aim to infer thermal limits and

optima and to compare these to mechanistic estimates (Brady et al., 2013) among multiple

medically-relevant mosquito vector species (Cianci et al., 2015; Cunze et al., 2016; Kraemer et

al., 2015; Kulkarni et al., 2010; Uusitalo et al., 2019).

However, our study also has several limitations. First, the lack of thermal breadth in the

background sampling may limit our ability to accurately infer thermal maxima.. Mosquito

species that occur in hotter climates, such as *Ae. aegypti, An. stephensi,* and *An. gambiae*, lack a

sufficient amount of ecoregion-matched background points that have comparably high, or higher,

temperatures than the occurrence points, making it difficult to estimate the temperature range

that would be hot enough to prevent occurrence (i.e., identify thermal maxima) (Merow et al.,

2013). Second, because our goal was to create accurate but comparable species distribution

models for seven mosquito species across their global extent, we primarily relied on publicly

available GBIF occurrence points (see Methods for data filtering criteria), which may not capture

the full extent of each species that other sources such as local vector surveillance data might

capture. Third, since we were focused on comparing laboratory-derived thermal performance

curves to field occurrence, we used temperatures measured during the activity season as

predictors in XGBoost models for four out of seven species. Despite the importance of activity

season temperature, other climatic variables such as winter low temperatures or dry-season

aridity may be equally or more important limitations on species distributions. Fourth, the species

distribution modeling framework assumes that species occupy their full climatic niche, but this

may not be the case for species that are actively expanding their ranges. Finally, our analysis is

based on a core assumption that using the 2000-2019 average of covariates can accurately

characterize the average ecological or habitat preferences of a mosquito reported in a specific

date and location.

CONCLUSIONS

As climate change shifts the geographic and seasonal distribution of environmental

conditions, it is critical to understand how temperature limits species ranges. Temperature in

particular affects vector-borne diseases because of its effects on vector biting rate, parasite

incubation rate, survival, and other life history traits. However, the thermal constraints on vector

occurrence are less well understood, and particularly how they vary among important vector

species. Here, we showed that temperature mean and variation during the activity season provide

important constraints on the ranges of seven important mosquito vectors, and that thermal

minima and, to a lesser degree, thermal optima observed in the field are closely correlated to

those measured in laboratory experimental studies. This finding suggests that species distribution

models can, to some extent, contribute to understanding the thermal biology of organisms that

cannot be studied in a laboratory setting. Importantly, statistical species distribution models

derived from field observations did not clearly identify upper thermal constraints even though

these have been directly demonstrated experimentally in the laboratory. Climate change is likely

to drive many regions past the currently observed range of temperatures. This highlights the

critical need for mechanistic, trait-based studies that capture temperature ranges at which

mosquito abundance and occurrence begin to decline (Mordecai et al., 2013; Ryan et al., 2015,

2019) and, at the very least, emphasizes that extrapolations from statistical models based on

current species distributions should be validated using physiological models (Elith & Leathwick,

2009; Hijmans & Graham, 2006; Khatchikian et al., 2011).

623
624

ACKNOWLEDGMENTS
TSA is supported by the National Institute of General Medical Sciences (grant no.
T32GM144273). MLC was supported by the Illich-Sadowsky Fellowship through the Stanford
Interdisciplinary Graduate Fellowship program at Stanford University and an Environmental
Fellowship at the Harvard University Center for the Environment. EAM and CKG were
supported by the National Science Foundation and the Fogarty International Center (grant no.
DEB-2011147). EAM was additionally supported by the National Institute of Allergy and
Infectious Diseases (grant nos R01AI168097 and R01AI102918), the National Institutes of
Health (grant no. R35GM133439), and by seed grants from the Stanford Woods Institute for the
Environment, King Center on Global Development, Center for Innovation in Global Health and
Terman Award. CKG was additionally supported by a Stanford Institute for Human-centered
Artificial Intelligence Postdoctoral Fellowship. The authors thank Marta Shocket and Oswaldo
Villena for providing mosquito life history trait data, and Marianne Sinka and Joshua
Longbottom for supplying additional species occurrence points for *Anopheles stephensi*.

REFERENCES

Abatzoglou, J. T., Dobrowski, S. Z., Parks, S. A., & Hegewisch, K. C. (2018). TerraClimate, a high-

resolution global dataset of monthly climate and climatic water balance from 1958–2015. *Scientific*

Data, *5*(1), 1–12.

Akpan, G. E., Adepoju, K. A., Oladosu, O. R., & Adelabu, S. A. (2018). Dominant malaria vector species

in Nigeria: Modelling potential distribution of Anopheles gambiae sensu lato and its siblings with

MaxEnt. *PloS One*, *13*(10), e0204233.

Al Ahmed, A. M., Naeem, M., Kheir, S. M., & Sallam, M. F. (2015). Ecological distribution modeling of

two malaria mosquito vectors using geographical information system in Al-Baha Province, Kingdom

of Saudi Arabia. *Pakistan Journal of Zoology*, *47*(6), 1797–1806.

Almirón, W. R., & Brewer, M. E. (1996). Winter biology of Culex pipiens quinquefasciatus say, (Diptera:

Culicidae) from Córdoba, Argentina. *Memorias Do Instituto Oswaldo Cruz*, *91*(5), 649–654.

Amarasekare, P., & Savage, V. (2012). A framework for elucidating the temperature dependence of

653   fitness. *The American Naturalist*, *179*(2), 178–191.

654 Angilletta, M. J., Jr, & Angilletta, M. J. (2009). *Thermal Adaptation: A Theoretical and Empirical*

655   *Synthesis*. OUP Oxford.

656 Barbet-Massin, M., Rome, Q., Villemant, C., & Courchamp, F. (2018). Can species distribution models

657   really predict the expansion of invasive species? *PloS One*, *13*(3), e0193085.

658 Barve, N., Barve, V., Jiménez-Valverde, A., Lira-Noriega, A., Maher, S. P., Peterson, A. T., Soberón, J.,

659   & Villalobos, F. (2011). The crucial role of the accessible area in ecological niche modeling and

660   species distribution modeling. *Ecological Modelling*, *222*(11), 1810–1819.

661 Bernhardt, J. R., Sunday, J. M., Thompson, P. L., & O'Connor, M. I. (2018). Nonlinear averaging of

662   thermal experience predicts population growth rates in a thermally variable environment.

663   *Proceedings. Biological Sciences / The Royal Society*, *285*(1886).

664   https://doi.org/10.1098/rspb.2018.1076

665 Bhatt, S., Gething, P. W., Brady, O. J., Messina, J. P., Farlow, A. W., Moyes, C. L., Drake, J. M.,

666   Brownstein, J. S., Hoen, A. G., Sankoh, O., Myers, M. F., George, D. B., Jaenisch, T., Wint, G. R.

667   W., Simmons, C. P., Scott, T. W., Farrar, J. J., & Hay, S. I. (2013). The global distribution and

668   burden of dengue. *Nature*, *496*(7446), 504–507.

669 Brady, O. J., Johansson, M. A., Guerra, C. A., Bhatt, S., Golding, N., Pigott, D. M., Delatte, H., Grech,

670   M. G., Leisnham, P. T., Maciel-de-Freitas, R., Styer, L. M., Smith, D. L., Scott, T. W., Gething, P.

671   W., & Hay, S. I. (2013). Modelling adult Aedes aegypti and Aedes albopictus survival at different

672   temperatures in laboratory and field settings. *Parasites & Vectors*, *6*, 351.

673 Brown, J. H., Gillooly, J. F., Allen, A. P., Savage, V. M., & West, G. B. (2004). TOWARD A

674   METABOLIC THEORY OF ECOLOGY. *Ecology*, *85*(7), 1771–1789.

675 Buckley, L. B., Urban, M. C., Angilletta, M. J., Crozier, L. G., Rissler, L. J., & Sears, M. W. (2010). Can

676   mechanism inform species' distribution models? *Ecology Letters*, *13*(8), 1041–1054.

677 Buth, J. L., Brust, R. A., & Ellis, R. A. (1990). Development time, oviposition activity and onset of

678   diapause in Culex tarsalis, Culex restuans and Culiseta inornata in southern Manitoba. *Journal of the*

American Mosquito Control Association, *6*(1), 55–63.

Chang, L.-H., Hsu, E.-L., Teng, H.-J., & Ho, C.-M. (2007). Differential Survival of Aedes aegypti and

Aedes albopictus (Diptera: Culicidae) Larvae Exposed to Low Temperatures in Taiwan. *Journal of*

Medical Entomology, *44*(2), 205–210.

Chen, T., He, T., Benesty, M., Khotilovich, V., Tang, Y., Cho, H., Chen, K., & Others. (2015). Xgboost:

extreme gradient boosting. *R Package Version 0. 4-2*, *1*(4), 1–4.

Chevillon, C., Pasteur, N., Marquine, M., Heyse, D., & Raymond, M. (1995). POPULATION

STRUCTURE AND DYNAMICS OF SELECTED GENES IN THE MOSQUITO CULEX

PIPIENS. *Evolution; International Journal of Organic Evolution*, *49*(5), 997–1007.

Cianci, D., Hartemink, N., & Ibáñez-Justicia, A. (2015). Modelling the potential spatial distribution of

mosquito species using three different techniques. *International Journal of Health Geographics*, *14*,

10.

Conley, A. K., Fuller, D. O., Haddad, N., Hassan, A. N., Gad, A. M., & Beier, J. C. (2014). Modeling the

distribution of the West Nile and Rift Valley Fever vector Culex pipiens in arid and semi-arid

regions of the Middle East and North Africa. *Parasites & Vectors*, *7*, 289.

Craig, M. H., Snow, R. W., & le Sueur, D. (1999). A climate-based distribution model of malaria

transmission in sub-Saharan Africa. *Parasitology Today* , *15*(3), 105–111.

Crozier, L., & Dwyer, G. (2006). Combining population-dynamic and ecophysiological models to predict

climate-induced insect range shifts. *The American Naturalist*, *167*(6), 853–866.

Cunze, S., Koch, L. K., Kochmann, J., & Klimpel, S. (2016). Aedes albopictus and Aedes japonicus - two

invasive mosquito species with different temperature niches in Europe. *Parasites & Vectors*, *9*(1),

573.

Davis, A. J., Jenkinson, L. S., Lawton, J. H., Shorrocks, B., & Wood, S. (1998). Making mistakes when

predicting shifts in species range in response to global warming. *Nature*, *391*(6669), 783–786.

Dell, A. I., Pawar, S., & Savage, V. M. (2011). Systematic variation in the temperature dependence of

physiological and ecological traits. *Proceedings of the National Academy of Sciences of the United*

States of America, *108*(26), 10591–10596.

Deutsch, C. A., Tewksbury, J. J., Huey, R. B., Sheldon, K. S., Ghalambor, C. K., Haak, D. C., & Martin,

P. R. (2008). Impacts of climate warming on terrestrial ectotherms across latitude. *Proceedings of*

the National Academy of Sciences of the United States of America, *105*(18), 6668–6672.

Didan, K. (2015). MOD13Q1 MODIS/Terra vegetation indices 16-day L3 global 250m SIN grid V006.

NASA EOSDIS Land Processes DAAC, *10*.

Dimiceli, C., Carroll, M., Sohlberg, R., Kim, D. H., Kelly, M., & Townshend, J. R. G. (2015). MOD44B

MODIS/terra vegetation continuous fields yearly L3 global 250m SIN grid V006. *NASA EOSDIS*

Land Processes Distributed Active Archive Center, *10*.

Edwards, J. L. (2004). Research and Societal Benefits of the Global Biodiversity Information Facility.

Bioscience, *54*(6), 485–486.

Elith, J., & Leathwick, J. R. (2009). Species Distribution Models: Ecological Explanation and Prediction

Across Space and Time. *Annual Review of Ecology, Evolution, and Systematics*, *40*(1), 677–697.

Fatima, S. H., Atif, S., Rasheed, S. B., Zaidi, F., & Hussain, E. (2016). Species Distribution Modelling of

Aedes aegypti in two dengue-endemic regions of Pakistan. *Tropical Medicine & International*

Health: TM & IH, *21*(3), 427–436.

Focks, D. A., Haile, D. G., Daniels, E., & Mount, G. A. (1993). Dynamic life table model for Aedes

aegypti (Diptera: Culicidae): analysis of the literature and model development. *Journal of Medical*

Entomology, *30*(6), 1003–1017.

Freire, S., MacManus, K., Pesaresi, M., Doxsey-Whitfield, E., & Mills, J. (2016). Development of new

open and free multi-temporal global population grids at 250 m resolution. *Proceedings of the 19th*

AGILE Conference on Geographic Information Science. https://agile-

online.org/conference_paper/cds/agile_2016/shortpapers/152_Paper_in_PDF.pdf

Gamliel, I., Buba, Y., Guy-Haim, T., Garval, T., Willette, D., Rilov, G., & Belmaker, J. (2020).

Incorporating physiology into species distribution models moderates the projected impact of

warming on selected Mediterranean marine species. *Ecography*, *43*(7), 1090–1106.

Gaythorpe, K. A., Hamlet, A., Jean, K., Garkauskas Ramos, D., Cibrelus, L., Garske, T., & Ferguson, N. (2021). The global burden of yellow fever. *eLife*, *10*. https://doi.org/10.7554/eLife.64670

Gilbert, M., Nicolas, G., Cinardi, G., Van Boeckel, T. P., Vanwambeke, S. O., Wint, G. R. W., & Robinson, T. P. (2018). Global distribution data for cattle, buffaloes, horses, sheep, goats, pigs, chickens and ducks in 2010. *Scientific Data*, *5*, 180227.

Gogol-Prokurat, M. (2011). Predicting habitat suitability for rare plants at local spatial scales using a species distribution model. *Ecological Applications: A Publication of the Ecological Society of America*, *21*(1), 33–47.

Gorelick, N., Hancher, M., Dixon, M., Ilyushchenko, S., Thau, D., & Moore, R. (2017). Google Earth Engine: Planetary-scale geospatial analysis for everyone. *Remote Sensing of Environment*, *202*, 18–27.

Guisan, A., & Thuiller, W. (2005). Predicting species distribution: offering more than simple habitat models. *Ecology Letters*, *8*(9), 993–1009.

Heikkinen, R. K., Luoto, M., Virkkala, R., Pearson, R. G., & Körber, J.-H. (2007). Biotic interactions improve prediction of boreal bird distributions at macro-scales. *Global Ecology and Biogeography: A Journal of Macroecology*, *16*(6), 754–763.

Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., … Jean-Noël Thépaut. (2020). The ERA5 global reanalysis. *Quarterly Journal of the Royal Meteorological Society*, *146*(730), 1999–2049.

Hijmans, R. J., & Graham, C. H. (2006). The ability of climate envelope models to predict the effect of climate change on species distributions. *Global Change Biology*, *12*(12), 2272–2281.

Huey, R. B., & Berrigan, D. (2001). Temperature, demography, and ectotherm fitness. *The American Naturalist*, *158*(2), 204–210.

Johansson, M. A., Powers, A. M., Pesik, N., Cohen, N. J., & Staples, J. E. (2014). Nowcasting the spread of chikungunya virus in the Americas. *PloS One*, *9*(8), e104915.

Kanagaraj, R., Araujo, M. B., Barman, R., Davidar, P., De, R., Digal, D. K., Gopi, G. V., Johnsingh, A. J.

T., Kakati, K., Kramer-Schadt, S., Lamichhane, B. R., Lyngdoh, S., Madhusudan, M. D., Ul Islam

Najar, M., Parida, J., Pradhan, N. M. B., Puyravaud, J.-P., Raghunath, R., Rahim, P. P. A., … Goyal,

S. P. (2019). Predicting range shifts of Asian elephants under global change. *Diversity &*

Distributions, *25*(5), 822–838.

Kazadi, S.-N., & Kaoru, F. (1996). Interannual and long-term climate variability over the Zaire River

Basin during the last 30 years. *Journal of Geophysical Research*, *101*(D16), 21351–21360.

Khatchikian, C., Sangermano, F., Kendell, D., & Livdahl, T. (2011). Evaluation of species distribution

model algorithms for fine-scale container-breeding mosquito risk prediction. *Medical and Veterinary*

Entomology, *25*(3), 268–275.

Khoury, C. K., Carver, D., Barchenger, D. W., Barboza, G. E., Zonneveld, M., Jarret, R., Bohs, L.,

Kantar, M., Uchanski, M., Mercer, K., Nabhan, G. P., Bosland, P. W., & Greene, S. L. (2020).

Modelled distributions and conservation status of the wild relatives of chile peppers (Capsicum L.).

Diversity & Distributions, *26*(2), 209–225.

Kraemer, M. U. G., Reiner, R. C., Jr, Brady, O. J., Messina, J. P., Gilbert, M., Pigott, D. M., Yi, D.,

Johnson, K., Earl, L., Marczak, L. B., Shirude, S., Davis Weaver, N., Bisanzio, D., Perkins, T. A.,

Lai, S., Lu, X., Jones, P., Coelho, G. E., Carvalho, R. G., … Golding, N. (2019). Past and future

spread of the arbovirus vectors Aedes aegypti and Aedes albopictus. *Nature Microbiology*, *4*(5),

854–863.

Kraemer, M. U. G., Sinka, M. E., Duda, K. A., Mylne, A. Q. N., Shearer, F. M., Barker, C. M., Moore, C.

G., Carvalho, R. G., Coelho, G. E., Van Bortel, W., Hendrickx, G., Schaffner, F., Elyazar, I. R. F.,

Teng, H.-J., Brady, O. J., Messina, J. P., Pigott, D. M., Scott, T. W., Smith, D. L., … Hay, S. I.

(2015). The global distribution of the arbovirus vectors Aedes aegypti and Ae. albopictus. *eLife*, *4*,

e08347.

Kramer-Schadt, S., Niedballa, J., Pilgrim, J. D., Schröder, B., Lindenborn, J., Reinfelder, V., Stillfried,

M., Heckmann, I., Scharf, A. K., Augeri, D. M., Cheyne, S. M., Hearn, A. J., Ross, J., Macdonald,

D. W., Mathai, J., Eaton, J., Marshall, A. J., Semiadi, G., Rustam, R., … Wilting, A. (2013). The

784   importance of correcting for sampling bias in MaxEnt species distribution models. *Diversity &*

785   *Distributions*, *19*(11), 1366–1379.

786   Kulkarni, M. A., Desrochers, R. E., & Kerr, J. T. (2010). High resolution niche models of malaria vectors

787   in northern Tanzania: a new capacity to predict malaria risk? *PloS One*, *5*(2), e9396.

788   Kushmaro, A., Awerbuch-Friedl, T., Levins, R., & Others. (2015). Temperature effects on the basic

789   reproductive number (R0) Of west nile virus, based on ecological parameters: endemic Vs. new

790   emergence regions. *Journal of Tropical Diseases & Public Health*.

791   https://www.researchgate.net/profile/Tamara-Awerbuch-Friedlander-

792   2/publication/291013223_Temperature_Effects_on_the_Basic_Reproductive_Number_R0_Of_West

793   _Nile_Virus_Based_On_Ecological_Parameters_Endemic_Vs_New_Emergence_Regions/links/577f

794   f38608ae69ab8824139e/Temperature-Effects-on-the-Basic-Reproductive-Number-R0-Of-West-

795   Nile-Virus-Based-On-Ecological-Parameters-Endemic-Vs-New-Emergence-

796   Regions.pdf?_sg%5B0%5D=started_experiment_milestone&origin=journalDetail

797   Lacour, G., Chanaud, L., L'Ambert, G., & Hance, T. (2015). Seasonal Synchronization of Diapause

798   Phases in Aedes albopictus (Diptera: Culicidae). *PloS One*, *10*(12), e0145311.

799   Lafferty, K. D. (2009). The ecology of climate change and infectious diseases. *Ecology*, *90*(4), 888–900.

800   Lambrechts, L., Paaijmans, K. P., Fansiri, T., Carrington, L. B., Kramer, L. D., Thomas, M. B., & Scott,

801   T. W. (2011). Impact of daily temperature fluctuations on dengue virus transmission by Aedes

802   aegypti. *Proceedings of the National Academy of Sciences of the United States of America*, *108*(18),

803   7460–7465.

804   Leahy, L., Scheffers, B. R., Williams, S. E., & Andersen, A. N. (2020). Diversity and Distribution of the

805   Dominant Ant Genus Anonychomyrma (Hymenoptera: Formicidae) in the Australian Wet Tropics.

806   *Diversity*, *12*(12), 474.

807   Lehmann, T., Dao, A., Yaro, A. S., Adamou, A., Kassogue, Y., Diallo, M., Sékou, T., & Coscaron-Arias,

808   C. (2010). Aestivation of the African malaria mosquito, Anopheles gambiae in the Sahel. *The*

American Journal of Tropical Medicine and Hygiene, *83*(3), 601–606.

Lira-Noriega, A., Soberón, J., & Equihua, J. (2018). Potential invasion of exotic ambrosia beetles

Xyleborus glabratus and Euwallacea sp. in Mexico: A major threat for native and cultivated forest

ecosystems. *Scientific Reports*, *8*(1), 10179.

Li, S. L., Acosta, A. L., Hill, S. C., Brady, O. J., de Almeida, M. A. B., Cardoso, J. da C., Hamlet, A.,

Mucci, L. F., Telles de Deus, J., Iani, F. C. M., Alexander, N. S., Wint, G. R. W., Pybus, O. G.,

Kraemer, M. U. G., Faria, N. R., & Messina, J. P. (2022). Mapping environmental suitability of

Haemagogus and Sabethes spp. mosquitoes to understand sylvatic transmission risk of yellow fever

virus in Brazil. *PLoS Neglected Tropical Diseases*, *16*(1), e0010019.

Liu, C., White, M., & Newell, G. (2013). Selecting thresholds for the prediction of species occurrence

with presence-only data. *Journal of Biogeography*, *40*(4), 778–789.

Liu-Helmersson, J., Stenlund, H., Wilder-Smith, A., & Rocklöv, J. (2014). Vectorial capacity of Aedes

aegypti: effects of temperature and implications for global dengue epidemic potential. *PloS One*,

9(3), e89783.

Lounibos, L. P., & Juliano, S. A. (2018). Where Vectors Collide: The Importance of Mechanisms

Shaping the Realized Niche for Modeling Ranges of Invasive Aedes Mosquitoes. *Biological*

Invasions, *20*(8), 1913–1929.

Merow, C., Smith, M. J., & Silander, J. A., Jr. (2013). A practical guide to MaxEnt for modeling species'

distributions: what it does, and why inputs and settings matter. *Ecography*, *36*(10), 1058–1069.

Mertens, A., Swennen, R., Rønsted, N., Vandelook, F., Panis, B., Sachter-Smith, G., Vu, D. T., &

Janssens, S. B. (2021). Conservation status assessment of banana crop wild relatives using species

distribution modelling. *Diversity & Distributions*, *27*(4), 729–746.

Messina, J. P., Brady, O. J., Golding, N., Kraemer, M. U. G., Wint, G. R. W., Ray, S. E., Pigott, D. M.,

Shearer, F. M., Johnson, K., Earl, L., Marczak, L. B., Shirude, S., Davis Weaver, N., Gilbert, M.,

Velayudhan, R., Jones, P., Jaenisch, T., Scott, T. W., Reiner, R. C., Jr, & Hay, S. I. (2019). The

current and future global distribution and population at risk of dengue. *Nature Microbiology*, *4*(9),

1508–1515.

Messina, J. P., Pigott, D. M., Golding, N., Duda, K. A., Brownstein, J. S., Weiss, D. J., Gibson, H.,

Robinson, T. P., Gilbert, M., William Wint, G. R., Nuttall, P. A., Gething, P. W., Myers, M. F.,

George, D. B., & Hay, S. I. (2015). The global distribution of Crimean-Congo hemorrhagic fever.

Transactions of the Royal Society of Tropical Medicine and Hygiene, *109*(8), 503–513.

Meuti, M. E., Short, C. A., & Denlinger, D. L. (2015). Mom Matters: Diapause Characteristics of Culex

pipiens-Culex quinquefasciatus (Diptera: Culicidae) Hybrid Mosquitoes. *Journal of Medical*

Entomology, *52*(2), 131–137.

Mitchell, C. J. (1995). Geographic spread of Aedes albopictus and potential for involvement in arbovirus

cycles in the Mediterranean basin. *Journal of Vector Ecology: Journal of the Society for Vector*

Ecology, *20*(1), 44–58.

Mordecai, E. A., Caldwell, J. M., Grossman, M. K., Lippi, C. A., Johnson, L. R., Neira, M., Rohr, J. R.,

Ryan, S. J., Savage, V., Shocket, M. S., Sippy, R., Stewart Ibarra, A. M., Thomas, M. B., & Villena,

O. (2019). Thermal biology of mosquito-borne disease. *Ecology Letters*, *22*(10), 1690–1708.

Mordecai, E. A., Cohen, J. M., Evans, M. V., Gudapati, P., Johnson, L. R., Lippi, C. A., Miazgowicz, K.,

Murdock, C. C., Rohr, J. R., Ryan, S. J., Savage, V., Shocket, M. S., Stewart Ibarra, A., Thomas, M.

B., & Weikel, D. P. (2017). Detecting the impact of temperature on transmission of Zika, dengue,

and chikungunya using mechanistic models. *PLoS Neglected Tropical Diseases*, *11*(4), e0005568.

Mordecai, E. A., Paaijmans, K. P., Johnson, L. R., Balzer, C., Ben-Horin, T., de Moor, E., McNally, A.,

Pawar, S., Ryan, S. J., Smith, T. C., & Lafferty, K. D. (2013). Optimal temperature for malaria

transmission is dramatically lower than previously predicted. *Ecology Letters*, *16*(1), 22–30.

Morin, C. W., Monaghan, A. J., Hayden, M. H., Barrera, R., & Ernst, K. (2015). Meteorologically Driven

Simulations of Dengue Epidemics in San Juan, PR. *PLoS Neglected Tropical Diseases*, *9*(8),

e0004002.

Murray, C. J. L., Rosenfeld, L. C., Lim, S. S., Andrews, K. G., Foreman, K. J., Haring, D., Fullman, N.,

Naghavi, M., Lozano, R., & Lopez, A. D. (2012). Global malaria mortality between 1980 and 2010:

a systematic analysis. *The Lancet*, *379*(9814), 413–431.

Mweya, C. N., Kimera, S. I., Kija, J. B., & Mboera, L. E. G. (2013). Predicting distribution of Aedes

aegypti and Culex pipiens complex, potential vectors of Rift Valley fever virus in relation to disease

epidemics in East Africa. *Infection Ecology & Epidemiology*, *3*.

https://doi.org/10.3402/iee.v3i0.21748

Natekin, A., & Knoll, A. (2013). Gradient boosting machines, a tutorial. *Frontiers in Neurorobotics*, *7*,

21.

Nuñez-Penichet, C., Cobos, M. E., Barro, A., & Soberón, J. (2019). Potential migratory routes of Urania

boisduvalii (Lepidoptera: Uraniidae) among host plant populations. *Diversity & Distributions*, *25*(3),

478–488.

Paaijmans, K. P., Blanford, S., Bell, A. S., Blanford, J. I., Read, A. F., & Thomas, M. B. (2010).

Influence of climate on malaria transmission depends on daily temperature variation. *Proceedings of*

the National Academy of Sciences of the United States of America, *107*(34), 15135–15139.

Paaijmans, K. P., Heinig, R. L., Seliga, R. A., Blanford, J. I., Blanford, S., Murdock, C. C., & Thomas,

M. B. (2013). Temperature variation makes ectotherms more sensitive to climate change. *Global*

Change Biology, *19*(8), 2373–2380.

Paaijmans, K. P., Read, A. F., & Thomas, M. B. (2009). Understanding the link between malaria risk and

climate. *Proceedings of the National Academy of Sciences of the United States of America*, *106*(33),

13844–13849.

Parham, P. E., & Michael, E. (2010). Modeling the effects of weather and climate change on malaria

transmission. *Environmental Health Perspectives*, *118*(5), 620–626.

Paull, S. H., Horton, D. E., Ashfaq, M., Rastogi, D., Kramer, L. D., Diffenbaugh, N. S., & Kilpatrick, A.

M. (2017). Drought and immunity determine the intensity of West Nile virus epidemics and climate

change impacts. *Proceedings. Biological Sciences / The Royal Society*, *284*(1848).

https://doi.org/10.1098/rspb.2016.2078

Peers, M. J. L., Thornton, D. H., & Murray, D. L. (2012). Reconsidering the specialist-generalist

887 paradigm in niche breadth dynamics: resource gradient selection by Canada lynx and bobcat. *PloS*

888 *One*, *7*(12), e51488.

889 Pekel, J.-F., Cottam, A., Gorelick, N., & Belward, A. S. (2016). High-resolution mapping of global

890 surface water and its long-term changes. *Nature*, *540*(7633), 418–422.

891 Peterson, A. T., & Soberón, J. (2012). Species distribution modeling and ecological niche modeling:

892 Getting the concepts right. *Natureza & Conservação*, *10*(2), 102–107.

893 Poelchau, M. F., Reynolds, J. A., Elsik, C. G., Denlinger, D. L., & Armbruster, P. A. (2013). RNA-Seq

894 reveals early distinctions and late convergence of gene expression between diapause and quiescence

895 in the Asian tiger mosquito, Aedes albopictus. *The Journal of Experimental Biology*, *216*(Pt 21),

896 4082–4090.

897 Pulliam, H. R. (2000). On the relationship between niche and distribution. *Ecology Letters*, *3*(4), 349–

898 361.

899 Puntasecca, C. J., King, C. H., & LaBeaud, A. D. (2021). Measuring the global burden of chikungunya

900 and Zika viruses: A systematic review. *PLoS Neglected Tropical Diseases*, *15*(3), e0009055.

901 Richman, R., Diallo, D., Diallo, M., Sall, A. A., Faye, O., Diagne, C. T., Dia, I., Weaver, S. C., Hanley,

902 K. A., & Buenemann, M. (2018). Ecological niche modeling of Aedes mosquito vectors of

903 chikungunya virus in southeastern Senegal. *Parasites & Vectors*, *11*(1), 255.

904 Rinehart, J. P., Robich, R. M., & Denlinger, D. L. (2006). Enhanced Cold and Desiccation Tolerance in

905 Diapausing Adults of Culex pipiens, and a Role for Hsp70 in Response to Cold Shock but Not as a

906 Component of the Diapause Program. *Journal of Medical Entomology*, *43*(4), 713–722.

907 Robich, R. M., & Denlinger, D. L. (2005). Diapause in the mosquito Culex pipiens evokes a metabolic

908 switch from blood feeding to sugar gluttony. *Proceedings of the National Academy of Sciences of the*

909 *United States of America*, *102*(44), 15912–15917.

910 Rochlin, I., Ninivaggi, D. V., Hutchinson, M. L., & Farajollahi, A. (2013). Climate change and range

911 expansion of the Asian tiger mosquito (Aedes albopictus) in Northeastern USA: implications for

912 public health practitioners. *PloS One*, *8*(4), e60874.

Rueda, L. M., Patel, K. J., Axtell, R. C., & Stinner, R. E. (1990). Temperature-dependent development

and survival rates of Culex quinquefasciatus and Aedes aegypti (Diptera: Culicidae). *Journal of*

Medical Entomology, *27*(5), 892–898.

Ryan, S. J., Carlson, C. J., Mordecai, E. A., & Johnson, L. R. (2019). Global expansion and redistribution

of Aedes-borne virus transmission risk with climate change. *PLoS Neglected Tropical Diseases*,

13(3), e0007213.

Ryan, S. J., McNally, A., Johnson, L. R., Mordecai, E. A., Ben-Horin, T., Paaijmans, K., & Lafferty, K.

D. (2015). Mapping Physiological Suitability Limits for Malaria in Africa Under Climate Change.

Vector Borne and Zoonotic Diseases , *15*(12), 718–725.

Shapiro, L. L. M., Whitehead, S. A., & Thomas, M. B. (2017). Quantifying the effects of temperature on

mosquito and parasite traits that determine the transmission potential of human malaria. *PLoS*

Biology, *15*(10), e2003489.

Shocket, M. S., Ryan, S. J., & Mordecai, E. A. (2018). Temperature explains broad patterns of Ross River

virus transmission. *eLife*, *7*. https://doi.org/10.7554/eLife.37762

Shocket, M. S., Verwillow, A. B., Numazu, M. G., Slamani, H., Cohen, J. M., El Moustaid, F., Rohr, J.,

Johnson, L. R., & Mordecai, E. A. (2020). Transmission of West Nile and five other temperate

mosquito-borne viruses peaks at temperatures between 23°C and 26°C. *eLife*, *9*.

https://doi.org/10.7554/eLife.58511

Sinka, M. E., Bangs, M. J., Manguin, S., Chareonviriyaphap, T., Patil, A. P., Temperley, W. H., Gething,

P. W., Elyazar, I. R. F., Kabaria, C. W., Harbach, R. E., & Hay, S. I. (2011). The dominant

Anopheles vectors of human malaria in the Asia-Pacific region: occurrence data, distribution maps

and bionomic précis. *Parasites & Vectors*, *4*, 89.

Sinka, M. E., Pironon, S., Massey, N. C., Longbottom, J., Hemingway, J., Moyes, C. L., & Willis, K. J.

(2020). A new malaria vector in Africa: Predicting the expansion range of Anopheles stephensi and

identifying the urban populations at risk. *Proceedings of the National Academy of Sciences of the*

United States of America, *117*(40), 24900–24908.

Smeraldo, S., Di Febbraro, M., Bosso, L., Flaquer, C., Guixé, D., Lisón, F., Meschede, A., Juste, J.,

Prüger, J., Puig-Montserrat, X., & Russo, D. (2018). Ignoring seasonal changes in the ecological

niche of non-migratory species may lead to biases in potential distribution models: lessons from bats.

Biodiversity and Conservation, *27*(9), 2425–2441.

Soberón, J. M. (2010). Niche and area of distribution modeling: a population ecology perspective.

Ecography, *33*(1), 159–167.

Soberon, J., & Peterson, A. T. (2005). Interpretation of models of fundamental ecological niches and

species' distributional areas. *Biodiversity Informatics*, *2*(0). https://doi.org/10.17161/bi.v2i0.4

Somero, G. N. (2002). Thermal physiology and vertical zonation of intertidal animals: optima, limits, and

costs of living. *Integrative and Comparative Biology*, *42*(4), 780–789.

Stanaway, J. D., Shepard, D. S., Undurraga, E. A., Halasa, Y. A., Coffeng, L. E., Brady, O. J., Hay, S. I.,

Bedi, N., Bensenor, I. M., Castañeda-Orjuela, C. A., Chuang, T.-W., Gibney, K. B., Memish, Z. A.,

Rafay, A., Ukwaja, K. N., Yonemoto, N., & Murray, C. J. L. (2016). The global burden of dengue:

an analysis from the Global Burden of Disease Study 2013. *The Lancet Infectious Diseases*, *16*(6),

712–723.

Tesla, B., Demakovsky, L. R., Mordecai, E. A., Ryan, S. J., Bonds, M. H., Ngonghala, C. N., Brindley,

M. A., & Murdock, C. C. (2018). Temperature drives Zika virus transmission: evidence from

empirical and mathematical models. *Proceedings. Biological Sciences / The Royal Society*,

285(1884). https://doi.org/10.1098/rspb.2018.0795

Tun-Lin, W., Burkot, T. R., & Kay, B. H. (2000). Effects of temperature and larval diet on development

rates and survival of the dengue vector Aedes aegypti in north Queensland, Australia. *Medical and*

Veterinary Entomology, *14*(1), 31–37.

Uusitalo, R., Siljander, M., Culverwell, C. L., Mutai, N. C., Forbes, K. M., Vapalahti, O., & Pellikka, P.

K. E. (2019). Predictive mapping of mosquito distribution based on environmental and

anthropogenic factors in Taita Hills, Kenya. *International Journal of Applied Earth Observation and*

Geoinformation, *76*, 84–92.

Verdonschot, P. F. M., & Besse-Lototskaya, A. A. (2014). Flight distance of mosquitoes (Culicidae): A

metadata analysis to support the management of barrier zones around rewetted and newly

constructed wetlands. *Limnologica* , *45*, 69–79.

Villena, O. C., Ryan, S. J., Murdock, C. C., & Johnson, L. R. (2022). Temperature impacts the

environmental suitability for malaria transmission by Anopheles gambiae and Anopheles stephensi.

Ecology, e3685.

Vogels, C. B. F., Hartemink, N., & Koenraadt, C. J. M. (2017). Modelling West Nile virus transmission

risk in Europe: effect of temperature and mosquito biotypes on the basic reproduction number.

Scientific Reports, *7*(1), 5022.

Wesolowski, A., Qureshi, T., Boni, M. F., Sundsøy, P. R., Johansson, M. A., Rasheed, S. B., Engø-

Monsen, K., & Buckee, C. O. (2015). Impact of human mobility on the emergence of dengue

epidemics in Pakistan. *Proceedings of the National Academy of Sciences of the United States of*

America, *112*(38), 11887–11892.

Wiebe, A., Longbottom, J., Gleave, K., Shearer, F. M., Sinka, M. E., Massey, N. C., Cameron, E., Bhatt,

S., Gething, P. W., Hemingway, J., Smith, D. L., Coleman, M., & Moyes, C. L. (2017).

Geographical distributions of African malaria vector sibling species and evidence for insecticide

resistance. *Malaria Journal*, *16*(1), 85.

Wisz, M. S., Pottier, J., Kissling, W. D., Pellissier, L., Lenoir, J., Damgaard, C. F., Dormann, C. F.,

Forchhammer, M. C., Grytnes, J.-A., Guisan, A., Heikkinen, R. K., Høye, T. T., Kühn, I., Luoto, M.,

Maiorano, L., Nilsson, M.-C., Normand, S., Öckinger, E., Schmidt, N. M., … Svenning, J.-C.

(2013). The role of biotic interactions in shaping distributions and realised assemblages of species:

implications for species distribution modelling. *Biological Reviews of the Cambridge Philosophical*

Society, *88*(1), 15–30.

XGBoost Parameters. (n.d.). Retrieved May 2, 2022, from

https://xgboost.readthedocs.io/en/stable/parameter.html

Yaro, A. S., Traoré, A. I., Huestis, D. L., Adamou, A., Timbiné, S., Kassogué, Y., Diallo, M., Dao, A.,

Traoré, S. F., & Lehmann, T. (2012). Dry season reproductive depression of Anopheles gambiae in

the Sahel. *Journal of Insect Physiology*, *58*(8), 1050–1059.

Zhang, C., Wei, D., Shi, G., Huang, X., Cheng, P., Liu, G., Guo, X., Liu, L., Wang, H., Miao, F., &

Gong, M. (2019). Understanding the regulation of overwintering diapause molecular mechanisms in

Culex pipiens pallens through comparative proteomics. *Scientific Reports*, *9*(1), 6485.

996
997