Spatially resolved single-cell analysis uncovers protein kinase Cδ-expressing microglia with anti-tumor activity in glioblastoma

Reza Mirzaei*[1,2,3], Reid McNeil[4, 5], Charlotte D'Mello[1,2], Britney Wong[1,2], Susobhan Sarkar[1,2], Frank Visser[2], Candice Poon[1,2,4,6], Pinaki Bose[4,5,6], V Wee Yong*[1,2,6]

[1]Department of Clinical Neurosciences, University of Calgary, Calgary, AB, Canada. [2]Hotchkiss Brain Institute, University of Calgary, Calgary, AB, Canada. [3]Tisch Cancer Institute, Icahn School of Medicine at Mount Sinai, New York, NY. [4]Arnie Charbonneau Cancer Institute, University of Calgary, Calgary, AB, Canada. [5]Department of Biochemistry and Molecular Biology, University of Calgary, Calgary, AB, Canada. [6]Department of Oncology, University of Calgary, Calgary, AB, Canada.

* Corresponding authors. Email: vyong@ucalgary.ca; reza.mirzaei@mssm.edu

Abstract

Glioblastoma (GBM) is a brain tumor that poses a formidable challenge to treatment options available. The tumor microenvironment (TME) in GBM is highly complex, marked by immunosuppression and cellular heterogeneity. Understanding the cellular interactions and their spatial organization within the TME is crucial for developing effective therapeutic strategies. In this study, we integrated single-cell RNA sequencing and spatial transcriptomics in a GBM mouse model to unravel the spatial landscape of the brain TME. We identified a previously unrecognized microglia subtype expressing protein kinase Cδ (PKCδ) associated with potent anti-tumor functions. The presence of PKCδ-expressing microglia was confirmed in resected human GBM specimens. Elevating tumoral PKCδ expression using niacin or adeno-associated virus in mice enhanced the phagocytosis of GBM cells by microglia in culture and increased the lifespan of mice with intracranial GBM. These findings were corroborated in analyses of the TCGA GBM datasets where low PKCδ samples showed negative pathway enrichment for apoptosis, phagocytosis, and immune signaling pathways, as well as lower levels of immune cell infiltration overall. Our study underscores the importance of integrating spatial context to unravel the TME, resulting in the identification of previously unrecognized subsets of microglia with anti-tumor functions. These findings provide valuable insights for advancing innovative immunotherapeutic strategies in GBM.

82
83 **Main**
84
Glioblastoma (GBM) remains the predominant primary malignant brain tumor among adults,
posing a significant challenge in the realm of oncology (1). The standard treatment approach
involves maximal safe resection using surgery, subsequent radiation, and chemotherapy utilizing
the oral alkylating agent temozolomide (1). However, despite these aggressive interventions and
other approaches such as tumor-treating fields, patients still face a median overall survival of less
than 21 months (2).
91
Improving prognosis is a formidable hurdle to overcome for GBM patients and the answer may
lie within the intricate landscape of the tumor microenvironment (TME) in the brain. The brain
TME exhibits two pivotal features: it is highly immunosuppressive and profoundly
heterogeneous, both of which significantly contribute to the dismal prognosis associated with
GBM (3-6). Unlike other tissues, the brain TME boasts a distinctive composition characterized
by a diverse array of functionally specialized organ-resident cells (7). Although single-cell
analysis has substantially deepened our understanding of the intricacies within the TME, it
grapples with inherent limitations, particularly concerning the spatial architecture of the TME (8-
10). This limitation restrains our ability to comprehensively explore the TME architecture and
identify subsets of immune cells based on their precise spatial coordinates. Single-cell analysis,
while invaluable, offers only indirect insights into cellular interactions due to the loss of spatial
organization information (11). Given that the positioning of immune cells within tumors dictates
their functions (12-14), unraveling the TME necessitates a focus on their anatomical locations.
105
Guided by this insight, our hypothesis posits that the GBM microenvironment manifests a dual
framework of both functional and spatial organization, with spatial architecture intricately
shaping the phenotypes of immune cells. Spatially resolved transcriptomics emerges as a new
technology that empowers us to unravel cellular interactions and organization in situ, thereby
deciphering the intricate ecosystem of GBM. To validate our hypothesis, we conducted
comprehensive analyses, employing both single-cell RNA sequencing (scRNA-seq) and spatial
transcriptomics techniques on a GBM mouse model. Additionally, we integrated published
scRNA-seq data of human GBM and performed immunofluorescent analyses on human
specimens, providing a comprehensive exploration of the brain TME. Our results identify a
previously unrecognized and spatially distributed microglia (MG) subset expressing protein
kinase Cδ (PKCδ). This cell type subset has potent anti-tumor functions including phagocytic
activity, and can be stimulated by a repurposed medication, niacin, to stimulate this MG subset
and curb GBM growth.

Results

Single-cell and spatially resolved transcriptomic profiling of the GBM tumor microenvironment

To develop a comprehensive understanding of the transcriptomic architecture in the brain TME, we employed the use of a rodent model of GBM (**Fig. 1A**). This model involved the intracranial implantation of mouse brain tumor-initiating cells (BTICs) isolated from NPcis mice, which carry mutations in Nf1 and Trp53, into syngeneic recipients (15). We previously confirmed that this model exhibited the prototypic phenotype of human GBM (16). In vivo bioluminescence imaging confirmed tumor growth in mice (**Supp Fig. 1A**). We obtained 11 brain samples from mice four weeks post-tumor implantation and generated single-cell suspensions from nine mice for scRNA-seq. After sorting of live cells using flow cytometry, we proceeded to process them for scRNA-seq utilizing a droplet-based platform (**Supp Fig. 1B**).

The transcriptomic profiles of 5440 cells were generated and grouped into 17 unique Seurat clusters based on gene expression profiles (GEPs) (**Fig. 1B**). Next, cell type deconvolution analysis was performed and identified 11 cell types based on differential expression of known cell type marker genes. This included acute neural stem cells (aNSCs), B cells, dendritic cells, erythrocytes, granulocytes, monocyte derived macrophages (MDMs), microglia (MG), natural killer cells, neurons, oligodendrocytes, and T cells (**Supp Fig. 1C**). Following the initial deconvolution, we noticed that cell type designation would encompass multiple default Seurat clusters, and this prompted us to perform a secondary refined deconvolution analysis. We separated the major cell types of monocytes and MG into subpopulations based on Seurat clustering and trajectory analysis using Monocle3 (**Sup Fig. 1D-H**). By employing this approach, we discerned variations in transcriptomic states among cell types along differentiation trajectories utilizing pseudotime values. This enabled the identification of five subsets of monocytes and four subsets of MG based on GEPs. This deconvolution of the immune contexture within the TME of GBM allowed us to pinpoint distinct subsets of immune cells and their impact on tumor biology. Consequently, we identified 21 cell type clusters ready for downstream analysis (**Sup Fig. 1C and H**).

The brain tissue from two additional tumor-bearing mice and one healthy brain without tumor implantation were subjected to Visium spatial transcriptomics as described in our previous study (5). After initial data processing, normalization and batch correction across samples, there was 18 default Seurat clusters identified with 12,885 spots ready for analysis (**Fig. 1C, D, Supp Fig. 2A and B**). To computationally identify spots containing tumor cells and their spatial locations we identified cells with copy-number variation (CNV) events using the InferCNV package. Notably, Seurat cluster 2 was enriched for CNV events compared to all other clusters, with common CNV events related to GBM with gains at chromosome 7 (17) (**Fig.1E, Supp Fig. 2C and D**). Cluster 2 was localized to known tumoral regions (**Fig. 2F**) aligning with the highly proliferative regions observed in H&E-stained slides reported in our previous study (5).

Differential expression analysis (DEA) identified differences in GEPs between tumoral spots identified by InferCNV compared to non-tumoral spots within spatial datasets. Volcano plots of differentially expressed genes (DEGs) highlighted statistically significant upregulation of cancer associated genes such as various extracellular matrix (ECM) molecules (*Col6a1*, *Col2a1, Bgn,*

Acan, Cspg4, *Vcan, Mgp, Mmp2*), cell cycle genes (*Ccnd3, Cdk4, Ccnd1*), immune-related genes
(*Ifitm2, Ifitm3, B2m, Lgals1, H2−D1, Tnfrsf1a, Cd9, Gpnmb*), and complement (*C4b*) in tumoral
spots (**Fig.1G** and **Supp Fig. 2E-H**). Ingenuity Pathway Analysis (IPA) of the DEGs from
cluster 2 in each mouse sample revealed enrichments in pathways related to metabolism, ECM
organization, and the neuronal system, among other significant pathways (**Fig. 1H** and **Supp
Fig. 3A**). Additionally, IPA identified notable enrichment of immune system and cell-cell
communication pathways in DEGs from cluster 2.
180
To assess the cellular distribution of the DEGs identified in cluster 2 of the mouse spatial data
within the human GBM microenvironment, we analyzed 44,712 cells from a published human
scRNA-seq dataset of seven GBM patients (4). After cellular deconvolution as described in our
previous study (5) (**Supp Fig. 3B**), we determined the expression profiles of genes from mouse
tumoral spots (cluster 2) from spatial data in human scRNA-seq data. We noticed that the ECM
genes such as *COL6A1*, *BGN*, *CSPG4*, *VCAN*, and *MGP* are mainly expressed by tumor cells
(**Supp. Fig. 3C**). On the contrary, immune related genes such as *IFITM2*, *IFITM3*, *LGALS1*,
TNFRSF1A, *CD9*, and *GPNMB* exhibited more specific expression by MDMs and MG compared
to other cell types in the human GBM microenvironment. This emphasizes the essential role of
MG and MDMs in orchestrating the immune response within both the human and mouse GBM
microenvironment.
192
Single-cell spatial heterogeneity in the GBM microenvironment
To identify the localization and distribution of different cell types with the GBM
microenvironment, we performed an integrative analysis using our single cell and spatial
transcriptomics datasets. Using the CARD package, we performed a spatially resolved cell type
deconvolution using our previously described mouse single cell data as reference. After spatial
deconvolution, we identified several patterns of cell type localization related to stromal and
immune cell types. High levels of GEP signatures related to neurons were distributed throughout
the spatial tissue slides, excluding known tumoral regions (**Fig. 2A** and **Supp Fig. 4A**). High
levels of GEP signatures related to oligodendrocytes were also identified, excluding the tumoral
regions. Furthermore, aNSC GEP signatures were identified within tumoral regions at the tumor
center; this may indicate their pivotal role in the tumor microenvironment related to tumor cell
origin.
205
Next, we unveiled the spatial distribution of immune cell types including subpopulations of
monocytes, MDMs, and MG from the scRNA-seq analysis. We also performed gene set
enrichment analysis (GSEA) using pathways from the Molecular Signatures DataBase
(MSigDB). The hallmark of cancer (HofC), KEGG and Reactome genesets were used to identify
enrichment of pathway activity using DEGs from cell types in the scRNA-seq data. Monocyte
subsets 1-5 were dispersed throughout the tumor and non-tumor regions, with tendency to
accumulate near tumoral margins (**Fig. 2A** and **Supp Fig. 4A**). Notably, monocyte subtype 5
exhibited a unique localization pattern, predominantly infiltrating the tumor core region, unlike
the other monocyte subtypes which localized to the tumor periphery. Monocyte subtype 5 was
marked by negative enrichment of several immune related pathways from HofC, KEGG and
REACTOME. These pathways included the negative enrichment of adaptive and innate immune
systems, interferon (IFN) gamma signaling, IFN alpha signaling, MHC class II antigen
presentation, interleukin (IL)10 and cytokine signaling within monocyte subset 5 suggesting an

immunologically inactive phenotype (**Supp Fig. 5** and **6**). Notably, monocyte subset 5 showed a
positive enrichment profile in the cell cycle pathway.
221
Additionally, MDMs cell populations were localized to the TME in spatial slides (**Fig. 2** and
Supp Fig. 4A). MDM1 and MDM2 subtypes displayed negative enrichment of HofC, KEGG,
and REACTOME pathways including IFN signaling, and cytokine signaling (**Supp Fig. 7**). The
pathway results imply that the GBM microenvironment was infiltrated by subtypes of monocyte
and MDM populations with immunologically dormant phenotypes, potentially contributing to
the unfavorable prognosis of GBMs.
228
Conversely, immune cell types that showed positive enrichment of immune related pathways
indicating immune activity and response were also localized to the TME region. These cell types
included monocyte subtypes 1, and 3 (**Fig. 2A** and **Supp. Fig. 4A**). These cell types were
positively enriched for HofC, KEGG, and REACTOME pathways including IFN alpha and
gamma signaling, cytokine signaling, antigen processing and presentation, IL10 signaling, and
JAK and STAT signaling (**Supp Fig. 5** and **6**). Enrichment of MG cell type signatures were
limited, but MG subset 2 signatures were present within the tumor area (**Fig. 2A** and **Fig. 4A**).
MG subset 2 showed positive enrichment of HofC, KEGG, and REACTOME pathways
including MHC class II antigen presentation, IFN alpha and gamma signaling, cytokine
signaling, and TCR signaling (**Supp Fig. 8** and **9**). These results may indicate a dynamic
relationship between immune cells that promote immune suppression or immune activity based
on the infiltration of specific subtypes of monocytes, MDMs, or MG and thus contribute to
cellular heterogeneity within the GBM TME.
242
In the realm of adaptive immune cells, including T and B cells as well as NK cells, we observed
significant accumulation in the tumor core compared to non-malignant areas (**Fig. 2A** and **Supp**
Fig. 4A). However, this pattern was not evident for DCs and granulocytes, which exhibited a
more uniform distribution across the tissue sections. Interestingly, the tumor areas were devoid
of dendritic cells. Altogether, these findings provide a comprehensive understanding of the
intricate spatial organization of various non-immune and immune cell populations within the
GBM tumor microenvironment.
250
Given the pivotal roles of MDMs, monocytes, and MG in GBM tumor development and their
higher accumulation in the TME, we decided to focus further on these immune cell populations
in this study. The tissues from a distinct set of mice implanted with the same tumor cells were
subjected to immunofluorescence (IF) staining, utilizing a combination of CD68 and TMEM119
markers to identify MDMs and MG. In alignment with our integrated transcriptomic analysis,
our IF data revealed that MDMs predominantly infiltrated near the tumor core (**Fig. 2B** and
Supp Fig. 4B). Conversely, MG were predominantly located at the outer edges of the tumor core
and exhibited dispersed accumulation throughout the entire brain. These findings corroborate
with the transcriptomic analyses, reinforcing the distinct spatial distribution of these immune cell
populations in the GBM microenvironment.
261
262 **Heterogeneous cellular interactions in the GBM microenvironment**
263
264 In our GBM mouse model, beyond just identifying patterns of cellular localization, we used the
265 CellChat package to decipher the global communication patterns among various cell types found

in the GBM TME. Within the deconvoluted cell types from the mouse scRNA-seq analysis, we
categorized cell-cell interactions into three distinct patterns. Specifically, cell types including
neurons 1, DCs, monocytes 1 and monocytes 2, as well as monocytes 3 and B cells, exhibited
pattern 1 of incoming communication (**Fig. 3A**). This pattern suggests that these cells received
similar signals from other cell types. On the other hand, MG3, MG4, MG2, MG1, aNSCs, and
oligodendrocytes displayed pattern 2 of incoming communication. Finally, pattern 3 was
observed in NK cells, granulocytes, T cells, monocytes 5, neurons 2, and MDM2.
272
We detected significant variations in incoming communication patterns, each intricately linked
with distinct signaling networks. Pattern 1 of incoming communications was found to be
associated with pathways such as macrophage migration inhibitory factor (MIF), chemokine
ligand (CCL), galectin, complement, secreted phosphoprotein 1 (SPP1), protein S (PROS),
visfatin, tumor necrosis factor (TNF), vascular endothelial growth factor (VEGF), CD40, IL4,
IL2, FASL among others (**Fig. 3B**). Pattern 2 exhibited a unique signature, involving different
pathways such as TGFβ, GAS, GRN, colony stimulating factor (CSF), IFN-II, oncostatin M
(OSM), B-cell activating factor (BAFF), IL6, leukemia inhibitory factor receptor (LIFR), and
insulin signaling networks. Lastly, incoming pattern 3 was associated with CXCL, protease-
activated receptors (PARs), IL1, KIT, IL12, epidermal growth factor (EGF), and CD137
networks. These findings provide a comprehensive view of the intricate molecular interactions
shaping the communication networks within the GBM microenvironment.
285
In the analysis of outgoing communication patterns, we observed a shared pattern 1 for exporting
signals within the GBM microenvironment among monocyte 4, granulocytes, MG2, DCs,
monocyte 3, MG1, monocyte 1, aNSCs, and oligodendrocytes (**Fig. 3C**). Neurons 1, neurons 2,
and B cells exhibited pattern 2, while NK cells, T cells, monocytes 2, monocytes 5, MDM, and
activated MDM displayed pattern 3 of outgoing communications. These distinct patterns
highlight the diverse strategies employed by different cell types in transmitting signals within the
complex GBM microenvironment. Pattern 1 of outgoing communications involved a diverse
array of signaling molecules, including MIF, galectin, complement, TGFβ, secreted
phosphoprotein 1 (SPP1), GRN, CXCL, visfatin, CSF, annexin, TNF among others (**Fig. 3D**).
Signaling in pattern 2 utilized different arrays of pathways such as GAS, midkine (MK),
angiopoietin-like protein (ANGPTL), Insulin-like growth factor (IGF), LIFR, and insulin. Lastly,
pattern 3 was associated with CCL, PARs, IFN-II, CD40, IL4, IL6, IL2, FASL, and FMS-like
tyrosine kinase 3 (FLT3).
299
Interestingly, immune cell subsets, particularly the monocytes and MDMs were highly
concentrated in tumor areas, including monocytes 3, 4, 5, MDM1, MDM2, and faint signatures
of MG1, MG2, which received a higher number of signaling networks associated with tumor
development, such as galectin, complement, TGFβ, SPP1, TNF, IFN-II, and IL4 (**Fig. 3E**).
Moreover, these infiltrating immune cell populations exported elevated levels of signaling
pathways, including CCL, GRN, CXCL, TNF, IL4, IL6, IL2, and IL12, compared to other
subsets (**Fig. 3F**). In contrast, MG3 and MG4, and monocytes 1, 2, with less concentrated
accumulation in tumor areas, expressed lower levels of signaling strength related to tumor
development or immune cell suppression, such as MIF, TNF, IL4, IL6, IFN-II, OSM, galectin,
and TGFβ, when compared to other subsets (**Fig. 3E** and **F**). These findings underscore the

significance of distinct immune cell populations with diverse spatial localizations in displaying
varying interaction profiles within the GBM microenvironment.
312
In our investigation of the primary contributors to both incoming and outgoing signaling
pathways among various cell types, we identified significant roles played by monocytes 3,
monocytes 4, MG1, B cells, and monocytes 1 in the incoming signaling pathways (**Fig. 3E**).
Regarding outgoing signaling patterns, key contributors included monocytes 3, MG1, neurons 1,
and monocytes 4, among others (**Fig. 3F**). In both the incoming and outgoing signaling
networks, pathways such as MIF, galectin, complement, TGFβ, and CCL emerged as the
strongest contributors for cell-to-cell interaction (**Supp Fig. 10**), highlighting their pivotal roles
in regulating cellular communications within the GBM microenvironment.
321
To validate these findings in human GBM and assess the significance of the top five identified
signaling networks from the mouse study, we examined the published scRNA-seq dataset of
human GBM. We explored all significant interactions (Ligand-Receptor pairs) associated with
MIF, galectin, complement, TGFβ, and CCL within the human GBM microenvironment. Except
for MIF, we noted the enrichment of all networks in the human GBM microenvironment.
Specifically, within the galectin network, *LGALS9* was implicated in interactions between
various subsets of tumor cells and immune cells (**Fig. 3G**). *LGALS9* expression was observed in
MG1, MG2, MDM3, and MDM5, and it played a role in their communication with T cells, MG3,
MDM2, and tumor cells 1 and 3. This communication was facilitated through different receptors,
including *CD44*, *PTPRC*, and *HAVCR2* on the target cells.
332
Another significant signaling network identified was the complement pathway. Several subsets
of human MG and MDMs, including MG1, MG2, MDM3, and MDM5, expressed C3
complement, facilitating their interaction with MG3 and MDM2 through *ITGAM_ITGB2*,
C3AR1, *ITGAX_ITGB2*, and *C3AR1*, respectively (**Fig. 3H**). Similarly, the TGFβ signaling
pathway played a significant role in mediating communication between MG1, MG2, MDM3,
and MDM5 with MG3, through engaging with *TGFbR1_R2* (**Fig. 3I**). Furthermore, in the CCL
network, *CCL3* and *CCL3L3* were primarily produced by MG1, MG2, tumor cells 4, tumor cells
6, oligodendrocytes, and MDM5, facilitating their interaction with MG3 through *CCR1* (**Fig.
3J**). Another chemokine, *CCL5*, was exclusively produced by MDM5 and participated in
crosstalk with MG3. In summary, the single-cell and spatial analysis of mouse GBM, in
conjunction with human GBM data, revealed critical signaling networks contributing to tumor
development within the tumor regions. We also discovered that different immune cells based on
their anatomical locations exhibited diverse interaction profiles with others cell types with the
possibility to modulate the GBM TME.
347
Identification of protein kinase Cδ (PKCδ)-positive myeloid cells in the GBM
microenvironment
After identifying significant immune related activity in the GBM microenvironment, we
examined individual DEGs in tumor spots more closely for important modulators of immunity.
Among the statistically significantly upregulated genes, increased expression of several genes
associated with immune regulation was evident in tumoral regions, including *H2-D1*, *C1qa*,
C1qc, *Ly86*, *Ifi27*, and *Prkcd* (which encodes PKCδ) in the spatial transcriptomics data (**Fig.**

4A). Given the limited information available about the immunological functions of PKCδ in the
TME, we conducted a subsequent focused investigation on this gene. Interestingly, spatial
transcriptomics data revealed the upregulation of *Prkcd* in both cluster 2 (tumor region) and
cluster 7 (tumor adjacent region) in tumor slide (**Fig. 4B** and **Supp Fig. 2H**). These two regions
corresponded to areas of the TME where infiltrating immune cells may play a significant role in
controlling tumor spread and ultimately influence disease prognosis.
361
Notably, the scRNA-seq analysis of the mouse GBM samples confirmed elevated *Prkcd*
expression in MG, monocytes, and MDMs, particularly in MG2 and monocytes 1 with the
highest levels of expression (**Fig. 4C**). Remarkably, based on spatial data, these immune cell
type clusters exhibited widespread infiltration across both tumor and non-tumoral areas, showing
a notable preference for infiltrating the tumor margin. To validate the cellular distribution of
PRKCD in human GBM, we analyzed the scRNA-seq data from human GBM samples, revealing
predominant expression of *PRKCD* in MG, with higher levels observed in MG1, MG2, and MG3
in that order (**Fig. 4D**).
370
Using TIMER2.0 (18,19) immune deconvolution analysis and the TCGA GBM RNA data the
correlation between *PRKCD* expression and TME purity revealed a significant negative
correlation; meanwhile a significant positive correlation was observed between *PRKCD*
expression and macrophages (M1-like) displaying an anti-tumor phenotype (**Fig. 4E**). The MG1
cluster of human scRNA-seq GBM also displayed less anti-inflammatory (M2-like) phenotypes
compared to other clusters based on lower levels of *IL10*, *CD163* and *ENTPD1* expression (**Fig.**
4F). Intriguingly, examination of outgoing and incoming signaling networks in the human
scRNA-seq GBM microenvironment using CellChat indicated that MG1 exhibited fewer
immunosuppressive signaling pathways, such as IL10, IL16, IL6, TNF, TGFβ, and galectin
(**Supp Fig. 11**). Additionally, as previously outlined, GSEA of MSigDB pathways from the
mouse scRNA-seq data showed that monocytes 1 and MG2 subtypes were significant positively
enriched for IFN gamma response, IFN alpha response, and inflammatory response (**Supp Fig.**
6-9).
384
To delve deeper into the phenotype of *Prkcd*-expressing cells, we performed differential gene
expression analysis between *Prkcd*[+] and *Prkcd*[-] MG, monocytes, and MDMs populations in our
mouse scRNA-seq data. Among the top-upregulated genes, we noted high expression of *Cybb* in
Prkcd[+] MDMs and MG (**Fig. 4G**). In human scRNA-seq GBM, we similarly observed high
expression of *CYBB* in MG1, MG2, and MG3, in that order (**Fig. 4H**). The *CYBB* gene encodes
the gp91-phox component of the phagocyte oxidase enzyme complex. This complex plays a
crucial role in producing superoxide and other reactive oxygen species, essential for microbial
killing (20). To corroborate these findings, we conducted IF staining for PKCδ, gp91-phox, and
IBA1 (MG and MDM marker) on GBM tissues from four human patients following surgical
resection. In all four patients, we observed co-expression of gp91-phox and PKCδ in MDMs and
MG within the tumor microenvironment (**Fig. 4I**). Altogether, these results suggest that MG
expressing high levels of *PRKCD* that infiltrate the tumor may possess the ability to phagocytose
tumor cells.
398
399
400

PKCδ in microglia contributes to phagocytosis of GBM cells: stimulation by niacin

Given the higher expression of *PRKCD* in MG as indicated by mouse and human scRNA-seq analysis, we conducted further validation experiments focusing on MG. We performed an in vitro phagocytosis experiment utilizing bioparticles labeled with the pHrodo dye which is sensitive to acidic pH environments in endosomes and lysosomes (**Fig. 5A**). pHrodo emits a distinct bright green fluorescence within phagosomes, allowing for the specific detection of intracellular phagocytosis.

In our previous research, we demonstrated that among various agents tested, vitamin B3 or niacin effectively stimulates phagocytosis in macrophages (21,22). We also observed significant elevation of phagocytosis in primary human MG treated with niacin compared with LPS (**Supp Fig. 12A-B**). Therefore, we utilized niacin as a stimulator of MG phagocytosis to investigate the role of PKCδ in the phagocytosis of tumor cells. MG were isolated from mouse brains and exposed to niacin to assess phagocytosis levels. Our results revealed a significant increase in bioparticle phagocytosis upon niacin treatment (**Fig. 5B, C** and **Supp Fig. 12C**). However, this elevation was abrogated when *Prkcd* was downregulated using two distinct shRNAs. To further validate the involvement of PKCδ in phagocytosis, we employed a pharmacological inhibitor of protein kinase C known as CRT 0066101 (23). It was evident that niacin, in the presence of CRT, failed to enhance phagocytosis (**Fig. 5D**), thus supporting the pivotal role of PKCδ in the phagocytosis process.

To better mimic conditions resembling the tumor microenvironment, we utilized patient derived BTICs labeled with pHrodo and co-cultured them with MG isolated from healthy individuals (**Fig. 5E**). We employed in vitro live imaging to monitor the phagocytosis of BTICs. In agreement with the findings from our in vitro mouse models, the downregulation of *PRKCD* in human MG substantially diminished the stimulatory effects of niacin on BTIC phagocytosis (**Fig. 5F** and **G**).

To obtain a comprehensive understanding of niacin's effects on the tumor microenvironment, we subjected mouse brain tissue isolated from niacin-treated mice to spatial transcriptomics and scRNA-seq. InferCNV analysis identified areas corresponding to tumor regions with chromosomal abnormalities as previously described (**Supp Fig. 13A-D**). After integrating with scRNA-seq data, we deconvoluted the cell types within the tumor microenvironment using our scRNA-seq mouse data and compared the results with those from earlier described tumor mice, collected at the same time point. Our analysis revealed that following niacin treatment, there was a higher level of infiltration of MG2 around the TME compared to untreated mice (**Fig. 5H, Fig. 2A** and **Supp Fig. 4A**). MG subtype 2 was characterized using the same GSEA and MSigDB pathway analysis. As described previously, this cell type was positively enriched for HofC, KEGG, and REACTOME pathways including MHC class II antigen presentation, IFN gamma and alpha signaling, cytokine signaling, and TCR signaling (**Supp Fig. 9**). In comparison to non-treated tumors, MG2 signatures were very limited in the TME. Beyond the cell infiltration, we also observed a significant upregulation of *Prkcd* in niacin-treated mice, especially in the tumor core margin, when compared to controls and normal brain tissue (**Fig. 5I** and **J**). Volcano plots of DEGs showed significant upregulation of *Prkcd* in cluster 2 (tumor region) and cluster 7 (tumor adjacent region) of spatial transcriptomics after niacin treatment (**Supp Fig. 13E** and **F**).

447 To validate these findings, we subjected additional tissues to IF staining for PKCδ and IBA1,
448 confirming higher PKCδ expression in IBA1-positive cells in niacin-treated mice compared to
449 controls (**Fig. 5K**).
450
451 **Overexpression of *Prkcd* in microglia and macrophages via adeno-associated virus**
452 **restrains intracranial tumor growth**
453
In a proof-of-concept experiment demonstrating the in vivo role of PKCδ in phagocytosis of
GBM cells and inhibition of tumor growth, we employed a targeted approach to upregulate
Prkcd specifically in MG and MDMs. Utilizing a newly developed adeno-associated virus
(AAV) named AAV-MG1.2 (24), we aimed to overexpress *Prkcd* in MG and MDMs in mice
(**Fig. 6A**). Following treatment of tumor-bearing mice with *Prkcd*-AAV, a substantial increase in
PKCδ protein expression was observed in IBA1-positive cells within the TME compared to
untreated control mice (**Fig. 6B** and **C**).
460
In a separate cohort, we administered treatment to tumor-bearing mice using either *Prkcd*-AAVs
or AAVs expressing the K376R point mutation (*Prkcd*-KR) in the ATP binding site of the *Prkcd*
gene, which exhibits deficient kinase activity (25). Then, brain tissues were subjected to IF
staining for arginase, a marker of immunosuppressive myeloid cells. AAV-*Prkcd* treatment led
to a significant reduction in double-positive IBA1 and arginase cells (**Fig. 6D** and **E**), indicating
a reprogramming of MG toward an anti-tumorigenic profile upon overexpression of wild-type
Prkcd. As phagocytosis is expected to result in increased apoptotic tumor cells, tissues from mice
treated with *Prkcd*-AAV were stained for cleaved caspase-3. Intriguingly, a significant
upregulation of cleaved caspase-3 was observed in cells proximal to MG (**Fig. F** and **G**).
Survival analysis further corroborated these findings, demonstrating that *Prkcd* overexpression
mediated by AAVs significantly prolonged the survival of mice compared to those treated with
AAVs expressing the mutant *Prkcd* (**Fig. 6H**).
473
474 **Analyses of *PRKCD* in the TCGA GBM database supports its anti-tumor function**
475
We stratified TCGA GBM bulk RNA-seq data based on *PRKCD* expression and conducted a
differential gene expression analysis between *PRKCD* low and high expression groups. Tumors
with *PRKCD* high expression exhibited upregulated levels of various ECM molecules, including
COL8A1, COL5A1, COL1A2, and *LAMA3* (**Supp Fig. 14A** and **B**). Moreover, genes associated
with inflammation and immune cell activation, such as *NLRP3*, *NFATC2*, and *PRAM1*, were
enriched in *PRKCD* high tumors. Notably, *PRKCD* high tumor samples exhibited significantly
higher immune scores using ESTIMATE when compared to low *PRKCD* samples (**Supp Fig.**
14C). Furthermore, using bulk deconvolution analysis of several immune cell types indicated
that high *PRKCD* samples had higher levels of immune cells including dendritic cells,
macrophages, and various subsets of T cells compared to *PRKCD* low samples (**Fig. 6I**). GSEA
and MSigDB analysis using KEGG and REACTOME pathways revealed enrichment of ECM
receptor interaction, cytokine receptor interaction, and chemokine signaling pathways in TCGA
GBM tumors with *PRKCD* high expression (**Fig. 6J** and **Supp Fig. 14D**). Notably, the apoptosis
pathway was also enriched in *PRKCD* high samples, indicating a possible connection to
increased apoptosis of tumor cells by PRKCD[+] MG. Conversely, low *PRKCD* samples showed
negative enrichment for apoptosis, phagocytosis, chemokine, and cytokine signaling pathways,

and ECM receptor interaction from IPA results of canonical pathway activity (**Supp Fig. 14E**
and **F**). Together the results from TCGA GBM analysis were concordant with scRNA-seq
analysis indicating that higher levels of *PRKCD* expression can increase anti-tumor response by
influencing immune cell infiltration in the TME.
496
497 **Discussion**
498
In recent years, breakthroughs in single-cell profiling technologies have transformed our
comprehension of tumor heterogeneity, plasticity, and potential therapeutic strategies. Despite
these advancements, the significance of the spatial organization within brain tumors has
remained largely unexplored. In this study, we pioneered integrative single-cell and spatial RNA
sequencing analyses in a GBM mouse model, unraveling the intricate spatial landscape of the
brain TME. By pinpointing crucial signaling pathways governing cell to cell interactions and
unveiling unique immune cell subsets, we extended our investigation to human GBM through
single-cell sequencing and bulk RNA TCGA data, corroborating to our mouse model findings.
This comprehensive approach not only validated our previous discoveries but also led to the
identification of a previously unknown MG subset with potent anti-tumor functions within the
brain microenvironment.
510
A significant finding in this study is the elucidation of the spatial landscape of the tumor immune
microenvironment in GBM. Distinct cell populations occupied specific regions within the brain,
and the spatial architecture of immune cells played a decisive role in shaping their phenotype and
functions. This was determined by their microenvironmental context and their interactions with
other cells. While a handful of studies have underscored the crucial role of cellular location
(17,26), the interaction profiles of immune cells with other cell types have been overlooked.
Notably, this aspect of the tumor immune microenvironment has been elucidated in only a
limited number of cancer types (14,27-30). Our study stands out as one of the few shedding light
on these nuances in GBM. Different immune cells, residing in diverse spatial locations,
displayed varied pathway activity and interaction profiles with neighboring cells in the brain
microenvironment. This underscores the significance of cellular localization indicating the
interacting partners of immune cells, a crucial aspect for cancer immunotherapy. Rejuvenating
immune cell activity requires a deep understanding of their interactome and their communication
with the surrounding environment, in addition to their cellular phenotype and localization.
Among various signaling networks, we observed that galectin, complement, TGFβ, and CCL are
among the pivotal pathways regulating cell-cell interactions in the GBM microenvironment.
527
Our integrative approach to studying the TME enabled us to unveil previously unidentified MG
exhibiting potent anti-tumor functions. These discoveries might have remained elusive if
traditional technologies requiring tissue disruption were employed. The identification of MG
with anti-tumor phenotypes and functions is crucial (31), as it opens avenues for the
development of tools to reprogram suppressed MG toward anti-tumor phenotypes. Through
spatial characterization of the GBM microenvironment, we pinpointed PKCδ$^+$ MG, present at a
very low frequency within the tumor microenvironment. Subsequently, we proposed two tools
including vitamin B3 (niacin) and AAV therapy with high potential for translation into clinical
applications, aiming to reprogram phenotypes of MG toward an anti-tumor phenotype with high
expression of PKCδ. PKCδ, functioning as a cytoplasmic kinase, plays diverse roles by

phosphorylating multiple target proteins involved in various cellular processes (32). Its
involvement in phagosome-mediated destruction of tumor cells, akin to its role in bacterial
infections (33,34), is plausible. The functions of PKCδ in tumor cells remain controversial (35-
37). However, in our study we found higher *PRKCD* expression in immune cells, specifically in
MDMs, monocytes, and MG, compared to tumor cells within the GBM microenvironment.
Intriguingly, niacin, a *PRKCD* inducer, did not directly impact GBM cell growth in our previous
study which showed that it affected myeloid cells (21). Moreover, we employed targeted
delivery of the *PRKCD* gene into MDMs and MG using an AAV strategy.
546
Notably, MG, being the earliest immune cells to combat tumors during the initial phases of
tumor growth in the central nervous system (38), offer a strategic window for intervention.
Moreover, MG and MDMs are the most prevalent immune cells in brain tumor
microenvironment (39). Thus, targeted therapies aimed at reprogramming these cells hold
substantial potential to hinder early-stage tumor growth in the CNS.
Reprogramming MG in the brain TME stands out as a promising strategy, especially when
contrasted with interventions in later stages characterized by expansive and uncontrollable tumor
growth. It is important to note that PKCδ is also expressed by other myeloid cells in GBM, such
as MDMs, monocytes and neutrophils. However, the focus on PKCδ in MG is particularly
intriguing due to their pivotal roles in brain tumor surveillance. Building upon our previous
findings (22) and the current results demonstrating the promotion of PKCδ expression by niacin,
along with the targeted in vivo elevation of PKCδ leading to MG reprogramming and tumor
growth inhibition, we have launched a phase 1/2a clinical trial to evaluate the safety and outcome
of niacin treatment in GBM patients (ClinicalTrials.gov Identifier: NCT04677049).
561
Collectively, our data indicate that the spatial localization of cells significantly influences the
phenotypes, interactome, and functions of immune cells within the GBM microenvironment. We
have elucidated the regional landscape of GBM and meticulously mapped its
microenvironmental terrain, encompassing the spatial localization of cells and their interactions.
This innovative approach to studying the tumor microenvironment has resulted in the
identification of a previously unrecognized subset of immune cells, along with the development
of two reprogramming tools poised for clinical translation. Altogether, these approaches have
proven effective in resolving tumor heterogeneity and generating a better understanding of GBM
and its immunosuppressive microenvironment.
571
Methods:

Animal experimental models
In this study, all mice utilized were of C57BL/6 background and were sourced from Charles
River as 6- to 8-week-old female wild-type mice. Ethical approval for all experiments involving
animals was obtained from the Animal Care Committee at the University of Calgary, following
the guidelines set forth by the Canadian Council of Animal Care.

578
Microglia isolation from human
MG were extracted from mixed preparations derived from brain tissue of human fetuses aged
between 18-22 weeks gestation, following established protocol (40). The purity of the MG
obtained exceeded 95%. The use of these specimens was approved by the University of Calgary

Research Ethics Board. The MG were diluted and fed using a medium composed of MEM, L-
glutamine, sodium pyruvate, 1% penicillin/streptomycin, 10% dextrose and 10% fetal bovine
serum (all culture constituents were from Invitrogen).
586

587 **Microglia isolation from mice**
588   Primary mouse MG, constituting over 95% pure non-adherent cells, were cultured from neonatal
589   brains using methods described in a previous publication (41).
590

591 **Microglia cell line**
592   CHME human MG cell line was used for in vitro validation experiments such as down regulation
593   of *PRKCD* in MG. This cell line was cultured under the same condition as primary human MG.
594

Human GBM tissues
For immunohistochemical analyses, human GBM tissues (n = 4) were obtained in accordance
with protocols approved by Indiana University's Institutional Review Board. Fresh tissues were
procured from patients through surgical procedures, immediately snap-frozen, and stored at -
80°C. Subsequently, the tissues were fixed in formalin and processed for histological
examination, leading to the creation of paraffin-embedded blocks. Hematoxylin and eosin
(H&E)–stained slides were meticulously examined by a neuropathologist. Written informed
consent was obtained from all patients prior to the collection of tumor biopsies. The information
of patients related to these specimens was described previously (5,42). We thank Dr. Chunhai
Hao, Indiana University, for providing the GBM samples.
605

Generation of GBM patient derived BTICs
Human BTIC lines were derived from resected tissues of GBM patients and cultured following
established protocols (21,43,44). These lines were sequentially cultured, preserved, and
authenticated within the University of Calgary BTIC Core. Previously reported identification of
two human BTIC lines (BT048 and BT073) was utilized in this study (5,42). Short tandem repeat
analysis was conducted for cell line authentication, and routine testing for Mycoplasma
contamination was performed using the Venor GeM Mycoplasma Detection Kit (Sigma,
MP0025).
614

Live cell imaging of in vitro phagocytosis
MG were seeded in a 96-well black-bottom plate at a density of 30,000 cells per well in complete
medium and incubated for 24 hours at 37°C with 5% CO_2. Following incubation, the medium
was replaced with DMEM containing 1% FBS. After 1 hour of incubation, niacin (100 μM) was
introduced to the cells, and they were further incubated for 12 hours at 37°C with 5% CO_2.
Subsequently, the medium was aspirated, and pHrodo S. aureus BioParticles (ThermoFisher,
P35361) or BTICs labeled with pHrodo Red, succinimidyl ester (ThermoFisher, P36600) were
added to the cells in live cell imaging solution. To stain cell nuclei, NucBlue Live ReadyProbes
Reagent (two drops per mL; Thermo Fisher, R37605) was also added. The cells were then
observed using an ImageXpress Micro Cellular Imaging and Analysis System (Molecular
Devices) for one hour. For continuous monitoring of phagocytosis, the Incucyte microscopy
system was employed. The percentage of NucBlue-positive cells that displayed pHrodo
positivity was quantified using MetaXpress or Incucyte software.
628

Intracranial tumor implantation

After dissociation of spheres, 25000-50000 viable cells from firefly luciferase–expressing or wild-type BTIC0309 were resuspended in 2 μL of PBS and stereotaxically implanted into the right striatum of mice, following previously described methods (45). In the niacin experiments, starting seven days after implantation, mice were treated with niacin at a dose of 100 mg/kg daily via intraperitoneal injection until the termination of the experiment. Control mice were injected with an identical vehicle (saline) solution.

Preparation of libraries for spatial transcriptomics

The spatial transcriptomics analysis was conducted using the Visium Spatial Gene Expression platform by 10x Genomics. Brain tissues, collected 40 days after tumor implantation, were frozen and sectioned into 10-μm slices, following the manufacturer's guidelines. Sections were placed on Visium slides and processed as per the user guide. Tissue optimization studies delineated an optimal permeabilization time of 12 mins. Brightfield images distinguishing tumor and non-tumor regions were captured using the EVOS FL Auto Imaging System. The subsequent library preparation steps were performed as per the user guide (Visium Spatial Gene Expression Reagent Kit; CG000239). The libraries were then sequenced on an Illumina NovaSeq 6000 system, employing a NovaSeq 200 Cycle S1 flow cell and specific read cycles for sequencing.

Data processing and batch correction of spatial transcriptomics

Spatial transcriptomics data was imported (images, counts, scale factors, and tissue position files) using Seurat and the CleanSpot R packages. Four tissue slides/samples were merged containing one normal tissue slide with no tumor, one tissue slide with tumor and niacin treatment, and two tissue slides with tumor and no niacin treatment. Initial data processing removed lowly expressed genes and spots with low feature counts; no spots were removed with mitochondrial contamination because removal of these spots compromised the overall number of spots available for analysis.

The merged spatial Seurat object was normalized (SCTransform, PCA components = 30), scaled, and then spots were clustered (res = 0.5) as per basic Seurat processing protocols. Next, spatial slides were checked for sample-based batch effects, and then these effects were removed using the Harmony R package. After batch correction, spots clustered based on biological gene expression and not by tissue slide/ samples. After initial data processing there were 12,588 spots ready for downstream analysis. After default Seurat clustering, differential expression analysis was performed between Seurat clusters using the FindAllMarkers function, thresholds for DEGs were set to log2fold change +/- 0.58 and p-adjusted value < 0.05.

InferCNV – Cancer/Tumor Spot Identification

Copy-Number Variation (CNV) events were used to identify tumor cells within spots of spatial slides. The InferCNV pipeline was run as per default, but importantly noting we used the Seurat clusters designated as "cancer spots – cluster 2" and "normal spots – all other clusters" to compare and identify CNV events in cancer vs normal spots. Tumor cells/spots were identified using the top duplication event identified, tumor spot identification overlapped nicely with visual images of tumor location on the tissue slide. Differential expression analysis was performed between identified cancer/ tumor spots and adjacent normal spots to validate InferCNV calls by identifying cancer-associated upregulated genes within cancer/ tumor spots.

Spot deconvolution in spatial transcriptomics
Spatial deconvolution was performed with the CARD R package to assign cell type identity to
spots by integrating our in-house mouse GBM single-cell data as a reference. CARD uses
autoregressive-based deconvolution to combine cell type specific expression information from
scRNA-seq data with correlation in cell type composition across tissue locations. Each spot on
the tissue slide was assigned a proportionality score of multiple cell types detected within each
spot. To validate and confirm CARD results we also performed deconvolution using the SingleR
to identify concordant calls between the two methods.
683

Analysis of public scRNA-seq datasets
The counts matrix of Richard et al (4) was downloaded and converted into a Seurat object using
the Seurat v3 package in R (46). Initial data filtering involved excluding cells with less than 50
genes and a percentage of mitochondrial genes exceeding 15%. Following this, data were
integrated and normalized using the SCTransform wrapper in Seurat. Principal component
analysis (PCA) reduction was employed, considering 15 significant PCA dimensions. Clusters
were identified using the FindNeighbours and FindClusters functions, with clustering executed at
a resolution of 0.5. Manual annotation of clusters was carried out based on the expression
patterns of lineage-specific signature genes. Cell clusters were visualized using the RunTSNE
function with PCA reduction. Differential expression genes (DEGs) for individual clusters
(versus all cells in other clusters) were identified using the FindMarkers function, including only
DEGs with a statistically significant P value of 0.05 for subsequent analysis. To visually
represent the results, the DoHeatmap functionality in Seurat was utilized to generate heat maps
of DEGs of interest for each cluster. Dot plots illustrating the average expression of hallmark
genes and genes of interest, expressed as a percentage in each cluster, were generated using the
DotPlot function. Additionally, the FeaturePlot and VlnPlot functions were employed to create
graphs depicting gene expression in specific cell clusters or sample groups.
701

Confocal immunofluorescence microscopy
We analyzed samples from GBM patients fixed in formalin and embedded in paraffin, as well as
frozen mouse sections, following established protocols (5,42). Following antibodies were used
for tissue staining: PKCδ (Abcam, ab182126, 1:200 dilution), IBA1 (ThermoFisher, PA5-18039,
1:100 dilution), Arginase (Cell Signaling, 93668, 1:50 dilution), Cleaved Caspase-3 (Cell
Signaling, 9661, 1:500 dilution), TMEM119 (Abcam, ab209064, 1:400 dilution), CD68
(Biolegend, 137001, 1:500 dilution) and gp91-phox (Novus Biologicals, NBP1-41012, 1:100
dilution). After primary antibody incubation, samples were treated with corresponding
fluorophore-conjugated secondary antibodies (1:500; Jackson ImmunoResearch Laboratories or
Thermo Fisher Scientific) and 4′,6-diamidino-2-phenylindole (DAPI; 1:1000). Laser confocal IF
imaging was performed at room temperature using the Leica TCS SP8 microscope with
consistent settings across samples, ensuring optimal contrast and minimal saturation. Each
sample generated three to four fields of view (FOV). Image acquisition was done using Leica
Application Suite X, and subsequent picture thresholding and three-dimensional (3D) rendering
were conducted with Imaris software (v9.9.1 Bitplane).
717

Confocal image analysis

Z-stacks were imported into Imaris in their original format. Cells expressing specific markers were identified in Imaris utilizing the surface tool. Background subtraction was applied to distinguish cells from the background environment. Accurate quantification was achieved through fluorescence thresholding and masking of undesired immuno-labeling. To identify positive signals, a consistent color brightness threshold was established using a value determined from tissues stained with negative secondary antibodies. The same threshold values for color brightness, along with size and circularity settings, were uniformly applied across all samples within each experimental set. Double positive cells were quantified using the parameter of the shortest distance between surfaces. Statistical data were exported to Excel for further analysis. Representative images were generated by merging and displaying the maximum-intensity projection of each channel/marker in a z-stack using pseudocolors in ImageJ. Adjustments to brightness and contrast were made consistently across samples to enhance image clarity.

Lentiviral vectors containing validated shRNA sequences targeting expression of human *PRKCD* (accession NM_006254) or a non-target control were obtained from Sigma (catalog numbers TRCN0000010193, TRCN0000195408, SHC002). Lentivirus production was performed using 293FT cells grown to 90% confluency on five, 15 cm^2 cell culture plates, by cotransfecting 112.5 µg lentiviral vector, 73 µg psPAX2 and 39.5 µg pMD2.G [gifts from Didier Trono (Addgene, plasmid # 12260) using PEIpro transfection reagent (Polyplus). The cell culture media was collected 48 hours post-transfection, centrifuged at 500xg for 5 min and 0.45 um filtered. The filtered media was underlaid with a 10% sucrose cushion and centrifuged at 12,000 xg for 4 hours at 4^{o}C (47). The resulting pellet was resuspended in sterile 1XDPBS and frozen at -80^{o}C as 20 µL aliquots. The lentivirus titer determined using the qPCR lentivirus titer kit was >10^9 IU/mL (Applied Biological Materials).

Plasmid construction

pAAV-CD68-hM4D(Gi)-mCherry, with the monocyte-specific promoter CD68 and was a gift from B. Roth (Addgene plasmid no. 75033(48)), was digested with SalI and HindIII to excise the hMD(Gi)-mCherry sequence. The cDNA sequence corresponding to mouse *Prkcd* (accession AF251036) was amplified from a mouse brain cDNA library and subcloned to generate pAAV-CD68-*Prkcd* or pAAV-CD68-*Prkcd*-KR (K376R mutant with ATP binding deficiency (25)) using the NEBuilder Hifi DNA assembly kit. All DNA constructs were verified by Sanger DNA sequencing.

AAV production for in vivo overexpression of *PRKCD* in microglia and macrophages

AAV viral vectors containing the MG1.2 capsid were generated using previous methods (49). Briefly, 293FT cells were grown to ~90% confluency in Corning Hyperflasks and co-transfected with 129 µg of pHELPER (Agilent), 238 µg of Rep-Cap plasmid encoding MG1.2, a gift from Minmin Luo (24) (Addgene plasmid # 184541) and 64.6 µg pAAV-*CD68-Prkcd* or pAAV-*CD68-Prkcd*-KR using the PEIpro transfection reagent (Polyplus). AAVs were precipitated from medium collected after 3 d and 5 d using a solution of 40% PEG/2.5 M sodium chloride and pooled with cells harvested after 5 d in buffer containing 500 mM sodium chloride, 40 mM Tris Base and 10 mM magnesium chloride. The lysate was incubated with 100 U ml−1 salt-active nuclease (Arcticzymes) at 37 °C for 1 h and then centrifuged at 2,000*g* for 15 min. AAVs were purified from the resulting lysate using an iodixanol step gradient containing 15, 25, 40 and 60%

iodixanol in Optiseal tubes (Beckman) followed by centrifugation at 350,000g using a Type 70
Ti ultracentrifuge rotor (Beckman). Following centrifugation, the AAVs were collected from the
40% layer using a 10-cc syringe and a 16-gauge needle, diluted in 1× PBS containing 0.001%
pluronic F68 (Gibco) and filtered using a 0.2-μm syringe filter. The AAVs were concentrated
and buffer exchanged by five rounds of centrifugation using Amicon Ultra-15 100-kDa-cutoff
centrifugal filter units (Millipore). The titer was determined using the qPCR Adeno-Associated
Virus Titration kit (Applied Biological Materials) and the purity was verified by SDS–PAGE and
total protein staining using InstantBlue reagent (Expedeon).
775
In vivo overexpression of *Prkcd* in myeloid cells with AAVs
To overexpress *Prkcd* in the brain of tumor-bearing mice, a solution of 5 μl saline containing 2.5
× 10^10 genome copies of AAV-MG1.2, designed with MG-targeting potential, was
intratumorally injected. The AAV-MG1.2 construct was engineered to overexpress *Prkcd* under
the control of the CD68 promoter. This procedure was performed 2 weeks after tumor injection.
Control mice were injected with 5 μl of saline containing 2.5 × 10^10 genome copies of AAV-
MG1.2 with a CD68 promoter designed to express a mutant version of *Prkcd* (*Prkcd*-KR).
783
Single cell isolation for scRNA-seq
Mice were euthanized four weeks post-tumor implantation, and brains were isolated. To enrich
for tumor tissue, brains were hemisected, and a 2 mm section around the tumor injection site was
collected into Hank's Balanced Salt Solution (HBSS, Gibco). Four brain tissues were pooled
together per sample to ensure an adequate number of cells for downstream scRNA-seq analysis
(Control: 5 mice, one replicates; niacin injection: 10 mice, two replicates). Subsequently, brain
tissues were homogenized using a Dounce homogenizer and filtered through a 40-μm cell
strainer. The cell suspension underwent Percol gradient separation to remove myelin debris.
After further filtration through a 35-μm cell strainer, cells were resuspended in PBS containing
1% heat-inactivated FBS (Sigma) and 0.1□mM EDTA.
794
scRNA-seq sample and library preparation
Once a single cell suspension was generated as described above, cells were labelled with
SYTOX™ Green Dead Cell Stain (ThermoFisher, S34860) to discriminate between live/dead
cells. Live cells were sorted on FACSAria III (BD) using a 100-μm nozzle. scRNA gene
expression libraries were prepared using the Chromium Next GEM single cell 3' reagent kits
v3.1 from 10x Genomics. Appropriate volume of cells, as determined from the user guide for
recovery of 3000 cells, was loaded on the Chromium single cell controller chip. Post cDNA
amplification QC and quantification as well as library construction QC was done using an
Agilent Bioanalyzer high sensitivity DNA chip for use with the Agilent 2100 Bioanalyzer. For
sequencing, all libraries were pooled and loaded at 300pM on the NovaSeq Ilumina system using
the NovaSeq S1 flowcell. A 28 base-pair Read 1 was used to sequence the cell barcode and UMI,
an 8 bp i7 index read was used to sequence the sample index and a 91bp Read 2 was used to
sequence the transcript using paired-end sequencing.
808
Single-cell RNAseq Data Processing
Sequencing depth obtained ranged from approximately 95,000-116,000 reads/cell. BCL files
obtained were processed using the Cell Ranger version 3.1 pipeline. Briefly, the fastq files were
processed with the cellranger count pipeline which uses STAR to align the reads to the mm10

mouse reference transcriptome. The cellranger aggregate pipeline was run to generate an
expression matrix with all combined libraries with normalization setting set to 'None'. The
Seurat R package was used to process single cell counts, cells with >200 nfeatures were
removed, as well as cells with <20% mitochondrial genes. In total, after filtering, there was 5440
cells ready for downstream analysis. Unfortunately, the tumor cells were lost during sample
processing and not available for analysis. The Seurat R package was used to perform data
normalization, scaling, and clustering (res = 0.5) of cells, resulting in 17 unique clusters.
Differential expression analysis was performed with the FindAllMarkers functions grouping by
Seurat clusters to identify top upregulated genes per cluster.
822
Cell type deconvolution of scRNA-seq data
scRNA-seq cell-type deconvolution analysis was performed using CellDex and SingleR R
packages with the mouse bulk RNA-seq cell type database used as a reference from CellDex. For
this analysis we were unable to identify a suitable "already labelled" single-cell mouse brain/
GBM dataset to deconvolute with, thus we settled for a bulk RNA-seq based approach. Cell type
labels determined by SingleR were validated using known marker genes for identified cell types.
After performing initial cell type deconvolution, we noticed that overall cell type assignments
encompassed multiple Seurat clusters. From this observation we performed a refined reclustering
analysis to identify subpopulations of cell types based on underlying Seurat clusters. We used the
Seurat clusters, DEGs between clusters, pseudotime/ trajectory analysis using the moncole3 R
package, and inferred biological pathway activity (GSEA and MSigDB – hallmark of cancer
gene sets, GO pathways, KEGG, and REACTOME using the ClusterProfiler R package). From
this analysis, we identified 5 subpopulations of monocytes (monocytes 1 and 3 were similar –
Group 1, monocytes 2 and 4 were similar – Group 2, and monocyte 5 – Group 3). From the
monocyte trajectory analysis, there were three distinct trajectories identified, along pseudo time,
Group 1 was timepoint 2, Group 2 were timepoint 3, and Group 3 were timepoint 1. There were
two populations of macrophages identified by SingleR, but these cell types did not contain any
subpopulations after re-clustering analysis and were labelled MDM1 and MDM2. Using the
same methods, we identified 4 subpopulations of MG. Based on DEGs and trajectory analysis,
each subpopulation was unique with some similarity between MG 1 and 2 subtypes. MG1 was
pseudotime point 2, MG2 was timepoint 3, MG3 was timepoint 1, and MG4 was timepoint 1.
844
CellChat – Cell to cell interaction analysis
The mouse GBM scRNA-seq dataset was used as input for a CellChat analysis, and cell type
identities were set to SingleR deconvolution/ reclustering analysis results. The mouse proportion
of the cell-to-cell interaction database was used, along with secreted ligand, ECM receptors, and
cell-cell contact databases. CellChat identified differentially expressed ligand-receptor genes
between cell types, and identified significant pathways related to cell-to-cell interactions. The
basic CellChat pipeline was used to identify interactions between cell types and incoming/
outgoing interactions was determined using non-negative matrix factorization (NMF).
853
Bulk RNA-seq TCGA GBM data download and processing
Glioblastoma (GBM) mRNA-seq expected counts and normalized transcripts per million (TPM)
counts were downloaded from the UCSCXena browser under the TCGA Hub. Expected counts
were used for DESeq2 differential expression analysis and TPM normalized counts were used
for all other downstream applications, including immune cell deconvolution analysis. For

DESeq2 differential expression analysis, GBM mRNA-seq counts were used to examine *PRKCD*
expression. GBM tumor samples were split into four quantiles of expression based on *PRKCD*,
as quantile 1 = low expression, and 4 = high expression. The GBM samples were filtered for
expression quantiles 1 and 4, and then DESeq2 differential expression analysis was performed
between these two expression groups with default settings. Differentially expressed gene
thresholds were set at a log2fold change +/- 0.58 and p-adjusted value < 0.05. DEGs were
visualized using volcano plots and heatmaps generated using ggplot2 and pheatmap functions.
866

Immune Contexture Analysis
The immune cell contexture of GBM high *PRKCD* verses *PRKCD* low expression of TCGA
samples were examined using two methods. Firstly, the ESTIMATE algorithm was used to
examine immune scores indicating overall levels of immune cell infiltration between the two
expression groups, which also included ESTIMATE stromal and purity scores. Secondly, a paper
published using immune cell markers validated by flow cytometry, immunohistochemistry, and
know cell type marker databases was used to deconvolute and assign tumor-infiltrating immune
(TIL) scores per TCGA sample including GBM samples. *PRKCD* expression groups (high = 4 vs
low = 1) were compared for differences in tumor-infiltrating immune scores using Wilcoxon T-
test (p-adj value < 0.05). The TIL score dataset came from Danaher et al., 2017 (50).
877

Qiagen Ingenuity Pathway Analysis of single cell and TCGA data
Differential expression results from scRNA-seq and TCGA analysis were used to run an IPA
core analysis and examine canonical pathway and upstream regulator activity between cell types
(scRNA-seq) and expression groups (*PRKCD* High vs Low – TCGA data). Differential
expression results from Seurat FindAllMarkers were used as input for IPA using previously
defined log2 foldchange and p-adjusted value for scRNA-seq data. Two differential expression
analysis were performed, one with DEGs in high *PRKCD* expression samples, and DEGs in low
PRKCD expression samples in TCGA data. Two separate IPA core analysis were run on the two
separate gene sets for TCGA. IPA results were downloaded and visualized using the ggplot2
package.
888

Pathway Analysis using ClusterProfiler R Package to perform GSEA of MSigDB Pathways
Secondary pathway analysis was performed using the Clusterprofiler R package, to perform
geneset enrichment analysis (GSEA) to identify enriched geneset/ pathways from the Mutational
Signatures Database (MSigDB). This database contains gene sets from various sources and for
this analysis included the hallmarks of cancer gene sets, KEGG, REACTOME, and Gene
Ontology (GO) pathways. The pathway analysis was used DEGs from DEAs to identify the
enrichment of pathway activity across scRNA-seq and bulk RNA-seq data. This method was
used to identify pathway activity in immune cell types in mouse and human scRNA-seq data
using DEGs of each cell type. This method was also used for TCGA GBM RNA data for DEGs
from *PRKCD* high vs low samples.
899

900　**Data availability**
901　Raw reads for the scRNA-seq, and spatial transcriptomics will be deposited in the Gene
902　Expression Omnibus (GEO) which will be available upon acceptance of the manuscript. All
903　other data are available upon request.
904

Code availability
Mouse scRNA-seq and spatial transcriptomics datasets discussed in this paper can be provide as per reasonable request from the authors.

Statistics and Reproducibility
Data were organized using Microsoft Excel, and graphs were created using GraphPad Prism 10.1.0. Each graph illustrates individual data points, with each point representing either an individual mouse (for in vivo experiments) or a distinct well in a 96-well plate (for in vitro experiments). The data points are accompanied by the mean ± SEM. The sample size was determined based on previous published studies (51,52), taking into account factors such as experimental cost, feasibility, and the availability of sex- and age-matched mice. Specific sample sizes are provided in the figure legends, and only one measurement was taken per sample. Littermate mice were randomly assigned to each experimental condition and treatment. Statistical analysis was performed using one-way ANOVA with Benjamini multiple-comparison test for comparisons involving two or more treatment groups against the control group. For data involving only two groups, significance was determined using a two-tailed, unpaired t-test. Blinding procedures were not implemented during the experiments.

Authors Contributions
R. Mirzaei: Conceptualization, data curation, software, analysis, validation, visualization, methodology, writing–original draft, writing–review and editing. R. McNeil: Data curation, analysis, visualization, methodology, writing–review and editing. B. Wong: Data curation, writing–review and editing. C. D'Mello: Software, analysis, Single-cell gene expression and spatial library preparation, writing–review and editing. S. Sarkar: Methodology, writing–review and editing. F. Visser: Methodology, writing–review and editing. C. Poon: Writing–review and editing. P. Bose: Supervision, software, writing–review and editing. V.W. Yong: Conceptualization, resources, supervision, funding acquisition, validation, project administration, writing–review and editing.

Acknowledgments
We thank members of the Yong laboratory for their thoughtful feedback on the study. The authors acknowledge the support and access to the Leica TCS SP8, ImageXpress Micro XLS High-Content Analysis System, and image analysis platforms provided by the Hotchkiss Brain Institute Advanced Microscopy Platform (HBI AMP) and the Cumming School of Medicine. The authors are thankful for the assistance and utilization of the IncuCyte Live-Cell Analysis System at the Snyder Institute's Live Cell Imaging Resource laboratory at the University of Calgary. The flow cytometry core and the UCDNA sequencing at the University of Calgary were also acknowledged for their support. R. Mirzaei received fellowship support from the University of Calgary's Eyes High program. Financial support for this study was provided by grants from the Canadian Institutes of Health Research and the Canadian Cancer Society.

Figure legends

Figure 1. Single-cell and spatially resolved transcriptomic profiling of the GBM tumor microenvironment. A. Schematic representation of the integrated scRNA-seq and spatial transcriptomics analysis conducted in the GBM mouse model. B and C. UMAP plots displaying Seurat clusters identified through scRNA-seq of mouse GBM (B) and spatial transcriptomics

analysis (C), respectively. D. Spatial Dimplot of Seurat clusters overlaid on tissue images. E. InferCNV analysis of spatial clusters highlighting malignant and non-malignant spots on tissue images. F. Spatial distribution depicting malignant and non-malignant areas. G. Volcano plots illustrating the differential expression of genes between malignant and non-malignant areas. H. Canonical pathways enriched in DEGs between malignant and non-malignant regions in mouse tumor 1.

Figure 2. Single-cell spatial heterogeneity in the GBM microenvironment. A. Deconvoluted cell type composition in spatial transcriptomics, determined using cell type-specific expression data obtained from the scRNA-seq dataset of mouse GBM. B. Representative immunofluorescence (IF) image of a mouse brain implanted with syngeneic tumor cells.

Figure 3. Heterogeneous cellular interactions in the brain tumor microenvironment. A. River plot illustrating patterns of incoming communication in target cells from scRNA-seq data of mouse GBM. B. Signaling networks enriched in three of the incoming communication patterns in mouse GBM. C. River plot displaying patterns of outgoing communication in secreting cells in mouse GBM. D. Signaling networks enriched in three of the outgoing communication patterns. E and F. Heatmaps indicating signals contributing significantly to outgoing or incoming signaling of specific cell groups, respectively in mouse GBM. G-J. Chord diagrams representing all significant signaling pathways from certain cell groups (source cells) to other cell groups (target cells) in human GBMs.

Figure 4. Identification of PKCδ-positive microglia in the GBM microenvironment. A. Volcano plot displaying DEGs between cluster 2 of mouse spatial transcriptomics compared to all other clusters. B. UMAP plot depicting the expression of *Prkcd* in spatial transcriptomics. C. Violin plot showing *Prkcd* expression in scRNA-seq clusters of mouse GBM. D. Violin plot displaying *PRKCD* expression in scRNA-seq clusters of human GBM. E. Correlation analysis between *PRKCD* expression, macrophage infiltration level, and phenotype based on Tumor Immune Estimation Resource (TIMER2.0). F. Dot plot depicting gene expression levels of M2-like macrophage markers in the human tumor microenvironment using human scRNA-seq data. G. Heatmap showing DEGs between *Prkcd*$^+$ and *Prkcd*$^-$ populations in scRNA-seq of mouse GBM. H. Violin plot illustrating the distribution of *CYBB* in different clusters of scRNA-seq of human GBM. L. Representative IF images of human GBM specimens stained for IBA1, PKCδ, and gp91-phox (CYBB) in four patients.

Figure 5. PKCδ in microglia contributes to phagocytosis of GBM cells. A. Schematic overview of the in vitro phagocytosis assay. B. Phagocytosis of pHrodo S. aureus BioParticles by CHME cells with downregulated *PRKCD* expression in the presence or absence of niacin. D. Phagocytosis assay depicting primary human MG stimulated with niacin in the presence or absence of the protein kinase inhibitor CRT 0066101. E. Schematic representation of in vitro phagocytosis of human BTICs conjugated with pHrodo by MG. F. pHrodo-conjugated human BTICs treated with CHME cells in which *PRKCD* expression was downregulated. G. Quantitative analysis of phagocytosis of human BTICs by CHME cells with downregulated *PRKCD* expression. H. Cellular deconvolution of Visium transcriptomics data for tumor mice receiving oral niacin. I. Comparison of *Prkcd* expression levels in niacin-treated mice versus control mice based on Visium transcriptomics. J. Quantification of *Prkcd* expression in Visium

997 transcriptomics. K. IF staining for PKCδ and IBA1 in mouse brains after niacin treatment
998 compared to control. One-way ANOVA with Benjamini multiple-comparison test for
999 comparisons involving two or more treatment groups against the control group. Significant
1000 differences in *Prkcd* expression between spatial slides were determined using Wilcoxon rank test
1001 ($p < 0.05$). All data presented as the mean ± SEM (error bars).
1002
Figure 6. Overexpression of *Prkcd* in microglia and macrophages via adeno-associated
virus restrains intracranial tumor growth. A. Schematic of intratumoral injection of AAVs
overexpressing *Prkcd* in MG and macrophages after tumor implantation. B. Representative IF
images showing expression of PKCδ in IBA1-positive cells after *Prkcd*-AAV injection. C.
Quantification of PKCδ-positive cells in the TME of *Prkcd*-AAV-treated mice versus control
mice injected with PBS. D. Representative IF images showing expression of arginase-1 in IBA1-
positive cells in *Prkcd*-AAV-treated mice compared to mice treated with AAVs expressing a
mutant version of *Prkcd* (K376R). E. Quantification of PKCδ-positive cells in the mouse TME.
F. Representative IF images of cleaved caspase-3 and IBA1. G. Quantification of cleaved
caspase-3 in the TME. H. Kaplan-Meier survival curve of mice treated with *Prkcd*-AAVs and
mutant *Prkcd*-AAVs encoding a deficient version of *PRKCD*. I. Tumor-infiltrating immune
score in GBM high *PRKCD* verses *PRKCD* low expression of TCGA samples. J. Enrichment
pathway analysis of GBM high *PRKCD* verses *PRKCD* low expression of TCGA samples.
Significance in panel C, E, and G was determined using a two-tailed, unpaired t-test. Kaplan-
Meier survival curves were analyzed for statistical differences between groups using the log-rank
Mantel-Cox test. Significant differences in immune scores were determined using Wilcoxon rank
test ($p < 0.05$).
1020
Supplementary Figure 1. Single-cell spatial analysis of GBM microenvironment. A.
Bioluminescence imaging of mice implanted with BTICs before harvesting tissues for scRNA-
seq. B. Schematic showing number of mice subjected to scRNA-seq and spatial transcriptomics.
C. UMAP plots displaying Seurat clusters identified through scRNA-seq. D. Pseudotime bar plot
of different monocyte and MDM sub-clusters showing differences in trajectories. E. UMAP of
Pseudotime analysis for monocytes and MDMs showing different trajectories. F. Pseudotime bar
plot of different MG sub clusters showing differences in trajectories. G. UMAP of Pseudotime
analysis for MG showing differences in trajectories. H. Heatmap of DEGs of scRNA-seq
clusters.
1030
1031 **Supplementary Figure 2. Spatial transcriptomics of mouse GBM.** A. Heatmap of DEGs in
1032 spatial clusters. B. Seurat spatial clusters overlaid on tissue image of normal brain. C and D.
1033 InferCNV plots of two tumor mice subjected to spatial transcriptomics. E-H. Volcano plots of
1034 DEGs in cluster 2 (tumor region) and cluster 7 (tumor adjacent region) for spatial transcriptomics
1035 of mouse GBM.
1036
1037 **Supplementary Figure 3. Transcriptomic profile of GBM microenvironment.** A. Canonical
1038 pathways enriched in DEGs between malignant and non-malignant regions in mouse tumor 2. B.
1039 UMAP plots displaying Seurat clusters identified through scRNA-seq of human GBM. C.
1040 Dotplot depicting gene expression levels in the human tumor microenvironment using human
1041 scRNA-seq data.
1042

Supplementary Figure 4. Spatial cellular composition in the brain of mouse GBM. A. Deconvoluted cell type composition in spatial transcriptomics. B. Representative IF image of mouse brain implanted with syngeneic tumor cells stained for TMEM119 and CD68.

Supplementary Figure 5. Gene set enrichment analysis of hallmark of cancer gene sets for monocyte subsets from scRNA-seq of mouse GBM.

Supplementary Figure 6. Gene set enrichment analysis of KEGG, and REACTOME for monocyte subsets from scRNA-seq of mouse GBM.

Supplementary Figure 7. Gene set enrichment analysis of KEGG, and REACTOME for MDM subsets from scRNA-seq of mouse GBM.

Supplementary Figure 8. Gene set enrichment analysis of hallmark of cancer gene sets for MG subsets from scRNA-seq of mouse GBM.

Supplementary Figure 9. Gene set enrichment analysis of KEGG, and REACTOME for MG subsets from scRNA-seq of mouse GBM.

Supplementary Figure 10. Signaling network in the GBM microenvironment. Contribution of each ligand-receptor pair to the overall signaling pathway calculated in scRNA-seq of mouse GBM.

Supplementary Figure 11. Network contributing to outgoing or incoming signaling in human GBM. Heatmaps showing signals contributing most to outgoing or incoming signaling of certain cell groups, respectively in the scRNA-seq of human GBM.

Supplementary Figure 12. Involvement of PKCδ in the phagocytosis of tumor cells. A. Representative flow cytometry plots of phagocytosis of pHrodo Bioparticles in human primary MG treated with LPS and niacin. B. Quantification of phagocytosis in human MG. C. Flow cytometry plots showing downregulation of *PRKCD* using two different shRNAs in CHME cells.

Supplementary Figure 13. Niacin treated spatial transcriptomic slide. A. Spatial Dimplot of Seurat clusters overlaid on tissue images of tumor mice treated with niacin. B. InferCNV analysis of spatial clusters highlighting malignant and non-malignant spots on tissue images. C. InferCNV heatmap showing CNV events across different chromosomes in cancerous spots (bottom) and normal spots (top). D. Spatial distribution depicting malignant and non-malignant areas. E-F. Volcano plots of DEGs between Seurat clusters for cluster 2 and cluster 7.

Supplementary Figure 14. Analysis of TCGA database for *PRKCD* expression. A. Volcano plot of DEGs between *PRKCD* high versus low samples using TCGA GBM RNA samples. B. Heatmap of top DEGs showing difference in gene counts between sample conditions. C. ESTIMATE analysis examining immune, stromal, and purity scores between *PRKCD* high and low samples, significant differences between conditions determined using Wilcoxon rank test (p < 0.05). D. GSEA and MSigDB analysis of KEGG and REACTOME pathways in *PRKCD* low

samples. E-F. Dot plots of IPA results on canonical pathways and upstream regulators enriched in either *PRKCD* high or low samples.

References

1. Stupp R, Mason WP, van den Bent MJ, Weller M, Fisher B, Taphoorn MJ, *et al.* Radiotherapy plus concomitant and adjuvant temozolomide for glioblastoma. N Engl J Med **2005**;352:987-96

2. Stupp R, Taillibert S, Kanner A, Read W, Steinberg D, Lhermitte B, *et al.* Effect of Tumor-Treating Fields Plus Maintenance Temozolomide vs Maintenance Temozolomide Alone on Survival in Patients With Glioblastoma: A Randomized Clinical Trial. JAMA **2017**;318:2306-16

3. Klemm F, Maas RR, Bowman RL, Kornete M, Soukup K, Nassiri S, *et al.* Interrogation of the Microenvironmental Landscape in Brain Tumors Reveals Disease-Specific Alterations of Immune Cells. Cell **2020**;181:1643-60.e17

4. Richards LM, Whitley OKN, MacLeod G, Cavalli FMG, Coutinho FJ, Jaramillo JE, *et al.* Gradient of Developmental and Injury Response transcriptional states defines functional vulnerabilities underpinning glioblastoma heterogeneity. Nat Cancer **2021**;2:157-73

5. Mirzaei R, D'Mello C, Liu M, Nikolic A, Kumar M, Visser F, *et al.* Single-Cell Spatial Analysis Identifies Regulators of Brain Tumor-Initiating Cells. Cancer Res **2023**;83:1725-41

6. LeBlanc VG, Trinh DL, Aslanpour S, Hughes M, Livingstone D, Jin D, *et al.* Single-cell landscapes of primary glioblastomas and matched explants and cell lines show variable retention of inter- and intratumor heterogeneity. Cancer Cell **2022**

7. Mirzaei R, Yong VW. Microglia-T cell conversations in brain cancer progression. Trends Mol Med **2022**;28:951-63

8. Yeo AT, Rawal S, Delcuze B, Christofides A, Atayde A, Strauss L, *et al.* Single-cell RNA sequencing reveals evolution of immune landscape during glioblastoma progression. Nat Immunol **2022**;23:971-84

9. González-Silva L, Quevedo L, Varela I. Tumor Functional Heterogeneity Unraveled by scRNA-seq Technologies. Trends Cancer **2020**;6:13-9

10. Wischnewski V, Maas RR, Aruffo PG, Soukup K, Galletti G, Kornete M, *et al.* Phenotypic diversity of T cells in human primary and metastatic brain tumors revealed by multiomic interrogation. Nat Cancer **2023**;4:908-24

11. Armingol E, Officer A, Harismendy O, Lewis NE. Deciphering cell-cell interactions and communication from gene expression. Nat Rev Genet **2021**;22:71-88

12. Laviron M, Petit M, Weber-Delacroix E, Combes AJ, Arkal AR, Barthélémy S, *et al.* Tumor-associated macrophage heterogeneity is driven by tissue territories in breast cancer. Cell Rep **2022**;39:110865

13. Ravi VM, Will P, Kueckelhaus J, Sun N, Joseph K, Salié H, *et al.* Spatially resolved multi-omics deciphers bidirectional tumor-host interdependence in glioblastoma. Cancer Cell **2022**;40:639-55.e13

14. Moldoveanu D, Ramsay L, Lajoie M, Anderson-Trocme L, Lingrand M, Berry D, *et al.* Spatially mapping the immune landscape of melanoma using imaging mass cytometry. Sci Immunol **2022**;7:eabi5072

15. Reilly KM, Loisel DA, Bronson RT, McLaughlin ME, Jacks T. Nf1;Trp53 mutant mice develop glioblastoma with evidence of strain-specific effects. Nat Genet **2000**;26:109-13

16. Pisklakova A, McKenzie B, Zemp F, Lun X, Kenchappa RS, Etame AB, *et al.* M011L-
deficient oncolytic myxoma virus induces apoptosis in brain tumor-initiating cells and
enhances survival in a novel immunocompetent mouse model of glioblastoma. Neuro
Oncol **2016**;18:1088-98
17. Ren Y, Huang Z, Zhou L, Xiao P, Song J, He P, *et al.* Spatial transcriptomics reveals
niche-specific enrichment and vulnerabilities of radial glial stem-like cells in malignant
gliomas. Nat Commun **2023**;14:1028
18. Li T, Fan J, Wang B, Traugh N, Chen Q, Liu JS, *et al.* TIMER: A Web Server for
Comprehensive Analysis of Tumor-Infiltrating Immune Cells. Cancer Res **2017**;77:e108-
e10
19. Li T, Fu J, Zeng Z, Cohen D, Li J, Chen Q, *et al.* TIMER2.0 for analysis of tumor-
infiltrating immune cells. Nucleic Acids Res **2020**;48:W509-W14
20. Rybicka JM, Balce DR, Khan MF, Krohn RM, Yates RM. NADPH oxidase activity
controls phagosomal proteolysis in macrophages through modulation of the lumenal
redox environment of phagosomes. Proc Natl Acad Sci U S A **2010**;107:10496-501
21. Sarkar S, Yang R, Mirzaei R, Rawji K, Poon C, Mishra MK, *et al.* Control of brain tumor
growth by reactivating myeloid cells with niacin. Sci Transl Med **2020**;12
22. Rawji KS, Young AMH, Ghosh T, Michaels NJ, Mirzaei R, Kappen J, *et al.* Niacin-
mediated rejuvenation of macrophage/microglia enhances remyelination of the aging
central nervous system. Acta Neuropathol **2020**;139:893-909
23. Harikumar KB, Kunnumakkara AB, Ochi N, Tong Z, Deorukhkar A, Sung B, *et al.* A
novel small-molecule inhibitor of protein kinase D blocks pancreatic cancer growth in
vitro and in vivo. Mol Cancer Ther **2010**;9:1136-46
24. Lin R, Zhou Y, Yan T, Wang R, Li H, Wu Z, *et al.* Directed evolution of adeno-
associated virus for efficient gene delivery to microglia. Nat Methods **2022**;19:976-85
25. Soh JW, Lee EH, Prywes R, Weinstein IB. Novel roles of specific isoforms of protein
kinase C in activation of the c-fos serum response element. Mol Cell Biol **1999**;19:1313-
24
26. Jain S, Rick JW, Joshi RS, Beniwal A, Spatz J, Gill S, *et al.* Single-cell RNA sequencing
and spatial transcriptomics reveal cancer-associated fibroblasts in glioblastoma with
protumoral effects. J Clin Invest **2023**;133
27. Guilliams M, Bonnardel J, Haest B, Vanderborght B, Wagner C, Remmerie A, *et al.*
Spatial proteogenomics reveals distinct and evolutionarily conserved hepatic macrophage
niches. Cell **2022**;185:379-96.e38
28. Sorin M, Rezanejad M, Karimi E, Fiset B, Desharnais L, Perus LJM, *et al.* Single-cell
spatial landscapes of the lung tumour immune microenvironment. Nature **2023**;614:548-
54
29. Karimi E, Yu MW, Maritan SM, Perus LJM, Rezanejad M, Sorin M, *et al.* Single-cell
spatial immune landscapes of primary and metastatic brain tumours. Nature
2023;614:555-63
30. Phillips D, Matusiak M, Gutierrez BR, Bhate SS, Barlow GL, Jiang S, *et al.* Immune cell
topography predicts response to PD-1 blockade in cutaneous T cell lymphoma. Nat
Commun **2021**;12:6726
31. Evans KT, Blake K, Longworth A, Coburn MA, Insua-Rodríguez J, McMullen TP, *et al.*
Microglia promote anti-tumour immunity and suppress breast cancer brain metastasis.
Nat Cell Biol **2023**

32.   Kikkawa U, Matsuzaki H, Yamamoto T. Protein kinase C delta (PKC delta): activation
1182   mechanisms and functions. J Biochem **2002**;132:831-9
1183   33.   Parihar SP, Ozturk M, Marakalala MJ, Loots DT, Hurdayal R, Maasdorp DB, *et al.*
1184   Protein kinase C-delta (PKCδ), a marker of inflammation and tuberculosis disease
1185   progression in humans, is important for optimal macrophage killing effector functions
1186   and survival in mice. Mucosal Immunol **2018**;11:496-511
1187   34.   Neehus AL, Moriya K, Nieto-Patlán A, Le Voyer T, Lévy R, Özen A, *et al.* Impaired
1188   respiratory burst contributes to infections in PKCδ-deficient patients. J Exp Med
1189   **2021**;218
1190   35.   Migliozzi S, Oh YT, Hasanain M, Garofano L, D'Angelo F, Najac RD, *et al.* Integrative
1191   multi-omics networks identify PKCδ and DNA-PK as master kinases of glioblastoma
1192   subtypes and guide targeted cancer therapy. Nat Cancer **2023**;4:181-202
1193   36.   Sarkar S, Yong VW. Reduction of protein kinase C delta attenuates tenascin-C stimulated
1194   glioma invasion in three-dimensional matrix. Carcinogenesis **2010**;31:311-7
1195   37.   Bessa C, Soares J, Raimundo L, Loureiro JB, Gomes C, Reis F, *et al.* Discovery of a
1196   small-molecule protein kinase Cδ-selective activator with promising application in colon
1197   cancer therapy. Cell Death Dis **2018**;9:23
1198   38.   Liu H, Sun Y, Zhang Q, Jin W, Gordon RE, Zhang Y, *et al.* Pro-inflammatory and
1199   proliferative microglia drive progression of glioblastoma. Cell Rep **2021**;36:109718
1200   39.   Khan F, Pang L, Dunterman M, Lesniak MS, Heimberger AB, Chen P. Macrophages and
1201   microglia in glioblastoma: heterogeneity, plasticity, and therapy. J Clin Invest **2023**;133
1202   40.   Giuliani F, Hader W, Yong VW. Minocycline attenuates T cell and microglia activity to
1203   impair cytokine production in T cell-microglia interaction. J Leukoc Biol **2005**;78:135-43
1204   41.   Mishra MK, Wang J, Keough MB, Fan Y, Silva C, Sloka S, *et al.* Laquinimod reduces
1205   neuroaxonal injury through inhibiting microglial activation. Ann Clin Transl Neurol
1206   **2014**;1:409-22
1207   42.   Mirzaei R, Gordon A, Zemp FJ, Kumar M, Sarkar S, Luchman HA, *et al.* PD-1
1208   independent of PD-L1 ligation promotes glioblastoma growth through the NFκB
1209   pathway. Sci Adv **2021**;7:eabh2148
1210   43.   Dzikowski L, Mirzaei R, Sarkar S, Kumar M, Bose P, Bellail A, *et al.* Fibrinogen in the
1211   glioblastoma microenvironment contributes to the invasiveness of brain tumor-initiating
1212   cells. Brain Pathol **2021**;31:e12947
1213   44.   Mirzaei R, Sarkar S, Dzikowski L, Rawji KS, Khan L, Faissner A, *et al.* Brain tumor-
1214   initiating cells export tenascin-C associated with exosomes to suppress T cell activity.
1215   Oncoimmunology **2018**;7:e1478647
1216   45.   Kelly JJ, Stechishin O, Chojnacki A, Lun X, Sun B, Senger DL, *et al.* Proliferation of
1217   human glioblastoma stem cells occurs independently of exogenous mitogens. Stem Cells
1218   **2009**;27:1722-33
1219   46.   Butler A, Hoffman P, Smibert P, Papalexi E, Satija R. Integrating single-cell
1220   transcriptomic data across different conditions, technologies, and species. Nat Biotechnol
1221   **2018**;36:411-20
1222   47.   Jiang W, Hua R, Wei M, Li C, Qiu Z, Yang X, *et al.* An optimized method for high-titer
1223   lentivirus preparations without ultracentrifugation. Sci Rep **2015**;5:13875
1224   48.   Grace PM, Strand KA, Galer EL, Urban DJ, Wang X, Baratta MV, *et al.* Morphine
1225   paradoxically prolongs neuropathic pain in rats by amplifying spinal NLRP3
1226   inflammasome activation. Proc Natl Acad Sci U S A **2016**;113:E3441-50

49. Challis RC, Ravindra Kumar S, Chan KY, Challis C, Beadle K, Jang MJ*, et al.* Systemic
AAV vectors for widespread and targeted gene delivery in rodents. Nat Protoc
2019;14:379-414
50. Danaher P, Warren S, Dennis L, D'Amico L, White A, Disis ML*, et al.* Gene expression
markers of Tumor Infiltrating Leukocytes. J Immunother Cancer **2017**;5:18
51. Ochocka N, Segit P, Walentynowicz KA, Wojnicki K, Cyranowski S, Swatler J*, et al.*
Single-cell RNA sequencing reveals functional heterogeneity of glioma-associated brain
macrophages. Nat Commun **2021**;12:1151
52. Rajendran S, Hu Y, Canella A, Peterson C, Gross A, Cam M*, et al.* Single-cell RNA
sequencing reveals immunosuppressive myeloid cell diversity during malignant
progression in a murine model of glioma. Cell Rep **2023**;42:112197
1238

Figure 1

Figure 2

A

Neurons 1 | Neurons 2 | Oligodendrocytes | aNSCs | MDM1 | MDM2 | Monocytes 1

Monocytes 2 | Monocytes 3 | Monocytes 4 | Monocytes 5 | Microglia 1 | Microglia 2 | Microglia 3

Signature strength
1
0.75
0.5
0.25
0

Microglia 4 | T cells | B cells | NK cells | Dendritic cells | Erythrocytes | Granulocytes

B

DAPI | Tumor cells (GFP)

TMEM119 | CD68

Scale bar=75

Figure 3

Figure 4

A

UMAP plot and volcano plot with labels DOWN (blue), NS (grey), UP (red); axes: -log10 (adjusted p-value) vs log2FC

B *Prkcd* — UMAP_1 vs UMAP_2

C *Prkcd* — Expression Level across: NSCs, B cells, Dendritic cells, Neurons 1, Neurons 2, Erythrocytes, Granulocytes, MDM1, MDM2, Microglia 1, Microglia 2, Microglia 3, Microglia 4, Monocytes 1, Monocytes 2, Monocytes 3, Monocytes 4, Monocytes 5, Oligodendrocytes, NK cells, T cells

D *PRKCD* — Expression Level across: MG1, MG2, Mac1, Oligodendrocytes1, Tumor cells1, Tumor cells2, Mac2, Tumor cells3, Tumor cells4, MG3, T cells, Tumor cells5, Tumor cells6, Mac3, Mac4, Oligodendrocytes2, Mac5, MG4

E Macrophage M1

p = 2.53e−08
Rho = −0.452 — Purity

Rho = 0.309
p = 2.37e−04 — XCELL

PRKCD Expression (log2 TPM), GBM; Purity vs Infiltration Level

F Dot plot — Average Expression; Percent Expressed (20, 40, 60); rows: MG4, MDM5, Oligodendrocytes2, MDM4, MDM3, Tumor cells6, T cells, Tumor cells5, MG3, Tumor cells4, Tumor cells2, MDM2, Tumor cells3, Tumor cells1, Oligodendrocytes1, MDM1, MG2, MG1; columns: IL10, CD163, ENTPD1

G *Prkcd* + cells | *Prkcd* − cells — heatmap; genes: Cebpb, Pim1, Polr2l, Itga4, Itgal, Tmsb10, Gpr141, Ciita, Cybb, Ighm, Hbb−bs, Ttr, Hba−a1, Igkc; Expression scale

H *CYBB* — Expression Level across: MG1, MG2, Mac1, Oligodendrocytes1, Tumor cells1, Tumor cells2, Mac2, Tumor cells3, Tumor cells4, MG3, T cells, Tumor cells6, Mac3, Mac4, Oligodendrocytes2, Mac5, MG4

I 100819, 101220, 110512, 110729

DAPI IBA1 PKCδ gp91-phox — Scale bar=50

Figure 5

Figure 6

A Injection of BTICs

4 weeks

Cd68   *Prkcd*

rAAV2_MG1.2

3-4 weeks

IF tissue staining

B DAPI IBA1 PKCδ

Scale bar=50

C
- ● Control
- ■ *Prkcd*-AAV

% PRKCD⁺ of IBA1⁺

0.0235

D Arginase-1 DAPI IBA1

Scale bar=50

E
- ● Mutant *Prkcd*-AAV
- ■ *Prkcd*-AAV

% Arginase-1 of IBA1⁺

0.0113

F Cleaved cas-3 DAPI IBA1

Scale bar=50

G
- ● Mutant *Prkcd*-AAV
- ■ *Prkcd*-AAV

% Cleaved caspase-3⁺ of DAPI⁺

0.0054

H
- Mutant *Prkcd*-AAV
- *Prkcd*-AAV

Percent survival

0.0265

Days

I

Tumor-infiltrating immune score

- Low *PRKCD*
- High *PRKCD*

B cells: p <= 0.01
DCs: p <= 0.01
Macrophages: p <= 0.0001
Exhausted T CD8: p <= 0.0001
T cells: p <= 0.0001
CD8T cells: p <= 0.05
Neutrophils: p <= 0.0001
Cytotoxic T cells: p <= 0.0001
Treg: p <= 0.0001
NK CD56dim cells: p <= 0.0001
Mast cells: ns
NK cells: p <= 0.0001
Th1 cells: p <= 0.05

J High *PRKCD* Samples

ECM_receptor_interaction
Cytokine_cytokine_receptor_interaction
neuroactive_ligand_receptor_interaction
Focal_adhesion
Calcium_signaling_pathway
Chemokine_signaling_pathway
Leukocyte_transendothelial_migration
Natural_killer_cell_mediated_cytotoxity
Wnt_signaling_pathway
FC_gamma_receptor_mediated_phagocytosis
ERBB_signaling_pathway
Apoptosis
Cell_adhesion_molecules_cams
Regulation_of_actin_cytoskeleton
VEGF_signaling_pathway
Pyrimidine_metabolism
Huntingtons_disease
Cell_cycle
Ribosome
Parkinsons_disease
Oxidative_phosphorylation

set Size
- 20
- 30
- 40
- 50
- 60
- 70

p.adjust
0.04
0.03
0.02
0.01

Normalized Enrichment Scores

Supplementary Figure 1

Supplementary Figure 2

Supplementary Figure 3

Supplementary Figure 4

Supplementary Figure 5

Supplementary Figure 6

Supplementary Figure 7

Supplementary Figure 8

Supplementary Figure 9

Supplementary Figure 11

Supplementary Figure 13