# Dystrophin deficiency impairs cell junction formation during embryonic myogenesis

**Elise Mozin[1], Emmanuelle Massouridès[2], Virginie Mournetas[3], Clémence Lièvre[1], Audrey Bourdon[1], Dana L Jackson[4], Jonathan S Packer[4], Cole Trapnell[4], Caroline Le Guiner[1], Oumeya Adjali[1], Christian Pinset[2], David L Mack[5] and Jean-Baptiste Dupont[1],***

Affiliations:

1: Nantes Université, CHU Nantes, INSERM, TARGET, F-44000 Nantes, France

2: Centre d'Etude des Cellules Souches, I-Stem, AFM, F-91100 Corbeil-Essonnes, France

3: ADLIN Science, F-91058 Evry, France

4: Department of Genome Sciences, University of Washington School of Medicine, Seattle, WA 98105, USA

5: Institute for Stem Cell and Regenerative Medicine, Department of Rehabilitation Medicine, University of Washington, Seattle, WA 98109, USA

**Contact info: *Correspondence: jean-baptiste.dupont@univ-nantes.fr; Twitter: @dpt_jb**

## Summary

Mutations in the *DMD* gene lead to Duchenne muscular dystrophy, a severe X-linked neuromuscular disorder which manifests itself as young boys acquire motor functions. DMD is diagnosed after 2 to 4 years, but the absence of dystrophin has an impact before symptoms appear in patients, which poses a serious challenge in the optimization of standards of care. In this report, we investigated the early consequences of dystrophin deficiency during skeletal muscle development. We used single-cell transcriptome profiling to characterize the myogenic trajectory of human pluripotent stem cells and showed that DMD cells bifurcate to an alternative branch when they reach the somite stage. Here, dystrophin deficiency was linked to marked dysregulations of cell junction families involved in the cell state transitions characteristic of somitogenesis. Altogether, this work demonstrates that *in vitro*, dystrophin deficiency has early consequences during myogenic development, which should be considered in future therapeutic strategies for DMD.

**Keywords: Duchenne muscular dystrophy, DMD, myogenesis, cell junctions, somite, hiPSCs**

## Introduction

Mutations in genes involved in skeletal muscle functions trigger a spectrum of diseases which can lead to

significant motor and respiratory impairments, sometimes shortly after birth[1–3]. The cellular and molecular

mechanisms of these diseases are complex and result in the breakdown of skeletal muscle homeostasis.

In Duchenne muscular dystrophy (DMD) patients, the lack of functional dystrophin causes systemic

pathological perturbations, which mostly affect striated muscles and the central nervous system. In muscle

cells, *DMD* mutations lead to plasma membrane fragility, calcium overload, oxidative stress, mitochondrial

impairments and the development of an inflammatory environment and of a fibro-adipogenic process[4]. As

muscles develop in the absence of dystrophin, capturing the pathological molecular cascade downstream

of the mutation and tracking progression over time are difficult using animal models. Early studies described

prenatal signs of muscle wasting in presymptomatic *mdx* mice and GRMD puppies[5,6]. Signs of muscle

damage such as variable myotube diameter, hyaline fibers and internal nuclei had also been observed in

fetuses at risk for DMD[7,8]. The most well-characterized functions of dystrophin have been described in fully

differentiated skeletal muscle cells, which express the longest isoform of the protein (Dp427m) to connect

the contractile machinery with the sarcolemma through the dystrophin-associated protein complex

(DAPC)[4]. However, the expression of the *DMD* locus gives rise to multiple other isoforms, and its regulation

during development is controlled by complex mechanisms. In the embryo, skeletal muscles emerge from

the paraxial mesoderm, which differentiates into transient metameric structures called somites, and then

into the dermomyotome[9]. This involves a succession of well-orchestrated cell division, migration and

transitions between epithelial and mesenchymal states, that later give rise to myogenic progenitors with

migratory properties[10–12]. In this context, the expression of dystrophin and the consequences of pathological

mutations during skeletal muscle development remain to be characterized in a reliable model.

Pluripotent stem cells represent a proxy for human embryonic development, and they allow the

characterization of early disease mechanisms in a dish[13,14]. Several groups have published protocols to

differentiate human induced pluripotent stem cells (hiPSCs) into paraxial mesoderm progenitors, somite

intermediates, dermomyotome and ultimately skeletal muscle progenitors[15–18]. Recently, our group has

used hiPSCs from DMD patients to demonstrate that marked transcriptome dysregulations occur prior to

skeletal muscle commitment[19]. Characterization of the *DMD* gene expression during early mesoderm

induction had led to the identification of an embryonic isoform of dystrophin (Dp412e) in mesoderm

progenitors and embryoid bodies[20]. This isoform possesses an alternative exon 1 and encodes an N-

truncated protein starting in exon 6 of the skeletal muscle dystrophin isoform (Dp427m). However, its

functions and molecular partners have remained unknown so far.

Transcriptomic analysis of hiPSCs and their derivatives at single-cell resolution offers the unprecedented

opportunity to shed light on complex biological processes such as embryonic development and to

investigate the consequences of disease-causing mutations in any given lineage intermediate. In particular,

single-cell RNA-sequencing (scRNA-Seq) have helped characterize the diversity of skeletal muscle cells

involved in muscle development[21,22], specification[23,24], ageing[25–27], and in various models of DMD[28–31].

Reconstruction of single-cell trajectories thanks to pseudotemporal ordering of cells have been used to

describe differentiation dynamics of myoblasts and the gene expression profiles associated with various

differentiation outcomes[32,33]. Studying the divergence of this trajectory when hiPSCs harbor a mutated *DMD*

will provide a deeper understanding of the molecular drivers of pathology and the cascade of compensatory

perturbations downstream of the mutation.

In this study, we established the single-cell trajectory of healthy and dystrophin-deficient hiPSCs subjected

to myogenic differentiation. We showed that hiPSCs derived from a DMD patient diverged from healthy

control cells as early as the somite stage (differentiation Day 10), and that the differences propagated as

differentiation progressed. By focusing on Day 10 and including an isogenic DMD line engineered from the

healthy control by CRISPR-Cas9, we demonstrated that several cell junction gene families were markedly

dysregulated as a consequence of dystrophin deficiency. Further characterization of the somite progenitors

derived from hiPSCs identified epithelial and mesenchymal populations coexisting *in vitro*, reflecting the

cell state transitions occurring in the embryo, and that the formation of epithelial islets is altered in the

absence of dystrophin. Finally, we confirmed our results in three additional DMD hiPSC lines by

immunostaining and analysis of previous bulk RNA-Seq data generated in our group, which strongly

suggests that *DMD* mutations have significant consequences during prenatal development, in relation to

the dynamics of cell junctions during somitogenesis.

## Results

**DMD hiPSCs bifurcate to an alternative branch of the myogenic trajectory**

Human iPSCs from a DMD patient harboring an exon 50 deletion in the *DMD* gene were subjected to directed differentiation into the skeletal muscle lineage in parallel with a healthy control[34], using a transgene-free and serum-free protocol[15]. Microscopic monitoring showed extensive cell proliferation and densification into multilayered cultures over time (Figure S1A). The differentiation was stopped at Day 28 when fields of thin and spindly cells resembling myotubes could be observed in the healthy control line. To investigate the dynamics of myogenic differentiation and the impact of dystrophin deficiency, cells were isolated at ten successive differentiation time points from Day 0 (D0) to Day 28 (D28), tagged with barcoded primers during *in situ* reverse-transcription and pooled for single-cell combinatorial indexing RNA-Seq (sci-RNA-Seq, Figure 1A)[35]. A total of 1917 individual cells could be retrieved from 20 distinct samples corresponding to the 2 hiPSC lines across 10 time points (Table S1), and used in the subsequent clustering analyses. After Uniform Manifold Approximation and Projection (UMAP) for dimension reduction, cells were distributed among 7 clusters expressing well-defined marker genes associated with pluripotency at Day 0 (*e.g. POU5F1*, clusters 4-5), primitive streak at Day 2 (*T,* cluster 5), paraxial mesoderm at Day 2-4 (*TBX6*, cluster 3), somite and dermomyotome at Day 7-10 (*PAX3*, clusters 2-7), and ultimately skeletal muscle progenitor cells from Day 22-28 (*PAX7*, *MYOD1*, cluster 1) (Figure S1B-C). Interestingly, DMD and healthy control cells were intermingled into superimposed clusters during the first week (Day 0 – 7), but from Day 10, they started to form distinct cell populations (Figure 1B-C). Single-cell trajectory reconstruction with Monocle[33] identified a branching point between Day 7 and Day 10, from which most of the cells segregated on two distinct branches in a genotype-dependent manner (Figure 1D-E). More precisely, 86 % of the healthy cells post branching point were on a single branch, while 86 % of the DMD cells were on the other branch (Figure S1D). Branch expression analysis modelling (BEAM) identified thousands of differentially-expressed genes (DEG) along the WT and the DMD-enriched branches. The most significant candidates (1987 genes with p-adj < 0.0001) were clustered based on their expression dynamics on the two branches for gene ontology analysis (Figure 1F, Table S2). Interestingly, modules of genes overexpressed along the Healthy branch were enriched in terms related to muscle development (*e.g.* Muscle cell differentiation, Muscle contraction, Actin cytoskeleton organization). In contrast, gene modules overexpressed along the

DMD branch matched with GO terms related to the development of alternative lineages, particularly

neurons (*e.g.* Neurogenesis, Synapse organization, Axon guidance). To gain insight into the regulation of

myogenesis in pseudotime, differential expression of paraxial mesoderm, somite and skeletal muscle

markers was assessed between the two branches. Significant differences in pseudotemporal dynamics and

expression were found for critical regulators of myogenesis, including *MYOD1*, *MYOG*, *MEF2C*. Notably,

the master regulators *MYOD1* and *MYOG* were almost unexpressed in DMD cells, in contrast with *PAX7*

and *MEF2C*, although for the latter, expression in DMD cells was significantly reduced as pseudotime

progressed (Figure 1G). In addition, genes coding for important structural proteins such as *MYH3*, *MYH8*,

*DES* or *TTN* were also found significantly dysregulated (Figure 1G). Thus, hiPSCs derived from a DMD

patient deviated from the myogenic trajectory followed by healthy control cells.

**DMD cells exhibit a marked dysregulation of cell junction genes at Day 10**

We focused our analysis at Day 10, the time point at which the deviation of DMD cells was first

evident on the single-cell trajectory. We observed that the sci-RNA-Seq data subset corresponding to the

"Day 10" time point formed a single UMAP cluster with a clear separation between DMD cells and healthy

controls (Figure 2A). Differential expression analysis using the regression model from Monocle 3 identified

94 genes significantly dysregulated in DMD cells (adjusted p-value < 0.01) (Table S3). Among these

differentially expressed genes (DEGs), we notably found multiple cell junction and extracellular matrix

genes, such as cadherins and proto-cadherins (*CDH11*, *PCDH9*), integrins (*ITGA1*, *ITGA4*, *ITGB1*, *ITGAV*),

fibronectin (*FN1*) and numerous collagens (*COL1A2*, *COL3A1*, *COL4A1*, *etc.*). Gene ontology (GO)

analysis confirmed significant enrichments in related biological processes, such as cell-matrix interaction

(p = 5.6E-4), cell-cell adhesion (p = 6.9E-7), extracellular matrix organization (p = 6.9E-4) and cell junction

organization (p = 4.0E-2) (Figure 2B). PANTHER Pathway enrichment analysis showed that the DEGs were

particularly enriched in genes involved in the integrin and cadherin signaling pathways (p = 2.4E-12 and p

= 4.2E-2, respectively) (Figure 2C). Of note, the GO analysis also indicated that DEGs included regulators

of key developmental processes in non-muscle lineages, such as neurogenesis and axon guidance

(*SEMA3A*, *EPHA3*, *NRP1*, *NEFL*, *UNC5C*), and angiogenesis (*ANGPT1*, *THSD7A*). Neuronal by-products

have already been observed when hiPSCs are differentiated with the protocol used in this first analysis[36].

In our study, the DMD hiPSC line might be more prone to such uncontrolled differentiation events, leading

to the formation of more alternative cell types.

Next, we assessed the robustness of our results using a distinct myogenic differentiation protocol.

We selected a commercial media formulation previously shown by our group and others to generate a

homogeneous muscle cell population[16,19]. hiPSCs were differentiated up to Day 10 with this new protocol

and analyzed by single-cell RNA-Seq. In addition to the DMD patient hiPSC line, we included an isogenic

DMD CRISPR line engineered from the healthy control line by deleting the entire exon 45 of the *DMD* gene

together with 17 base pairs in exon 54[37], leading to an absence of dystrophin protein at any differentiation

stage from hiPSCs to myotubes (Figure S2A). At Day 10, we confirmed that cells express somite marker

genes, and we showed that the DMD patient line expressed significantly higher levels of *PAX3*, *MEOX1*

and *NR2F1* than the healthy control, and lower levels of *MET*, *PTN* and *NR2F2* (Figure 2D). Interestingly,

the CRISPR line presented with a less pronounced dysregulation profile, yet statistically significant for all

the above-mentioned marker genes except *NR2F1*. After dimensional reduction, the UMAP plot revealed

that DMD cells and healthy controls were separated into distant clusters. The CRISPR clusters were found

in between, closer to the DMD cells with only a small fraction overlapping with healthy cells (Figure 2E).

We then conducted differential expression analysis using the Monocle 3 pipeline. DMD and CRISPR cells

showed 4,565 and 2,913 differentially expressed genes respectively when compared to the healthy control

line (p-adj < 0.01, Table S4). Among these, 1,885 genes were differentially expressed in both lines, which

represents the core gene set dysregulated at the somite stage as a direct consequence of the DMD

mutation (Figure 2F). Of note, this result also illustrated the influence of the genetic background on disease

manifestation at the transcriptome level, as 36 % of the CRISPR-induced dysregulations were not present

in the unrelated DMD patient line. Importantly however, the common DEGs also included 55 of the 94 genes

(59 %) found significantly dysregulated at Day 10 with the initial protocol, highlighting the overlap between

the two single-cell RNA-Seq data sets. To go further, we performed GO analysis on the 1,885 overlap

genes, and observed an enrichment in terms related to cell junctions, such as desmosome organization (p

= 7.2E-3), tight junction organization (p = 4.4E-4), and cell junction assembly (p = 2.2E-11) (Figure 2G).

Finally, we analyzed the detailed expression of key cell-cell junction gene families, and we observed that

tight junction genes such as claudins (*CLDN*), ocludin (*OCLN*) and tight junction proteins (*TJP*), but also

desmosome genes such as desmoplakin (*DSP*), desmogleins (*DSG*) and desmocollins (*DSC*) were

markedly down-regulated in the DMD line. In contrast, multiple genes from the protocadherin (*PCDH*) and

cadherin (*CDH*) families were up-regulated (Figure 2H). Similar to our previous observations, the CRISPR

line showed an intermediary dysregulation profile for all cell junction protein families (Figure 2I). In the

previous sci-RNA-Seq data analysis, adherens junctions were also found particularly affected in the BEAM

gene list, including members of the *CDH* and *PCDH* families, together with genes encoding catenins

(*CTNN*) (Figure S2B). Altogether, our results suggest a major role for cell junction gene families in the

initiation and progression of DMD during development.

**Dystrophin deficiency does not compromise myotube formation despite impaired upstream cell**

**state transitions**

In spite of earlier reports suggesting the existence of prenatal disease manifestations at the cellular

and molecular levels, DMD patients are born with functional muscles. We assessed the ability of DMD and

CRISPR hiPSCs to differentiate into multinucleated myotubes. Somite progenitors were amplified,

passaged in a Myoblast medium and further differentiated with a Myotube Fusion medium[16]. At Day 25,

cells were stained for two skeletal muscle proteins: myosin heavy chains and $\alpha$-actinin. Multinucleated

myotubes with striation patterns could be observed in the 3 hiPSC lines (Figure 3A). We quantified the area

positive for $\alpha$-actinin and obtained comparable fluorescent signals in the 3 hiPSC lines representing 34 %

of the total area on average (Figure 3B). Thus, the early dysregulations of the transcriptome observed at

the somite stage do not impact myotube formation *in vitro*. Prior to terminal differentiation, hiPSCs undergo

several rounds of amplification and passages as they transition through the successive developmental

intermediates. These culture conditions may introduce positive bias by retaining only the cells most capable

of differentiation, hereby masking prior developmental deficiency. Hence, we examined the appearance of

cell cultures at the somite stage, around Day 7 to Day 10. During paraxial mesoderm and somite

development in the embryo, progenitors undergo successive epithelial-mesenchymal and mesenchymal-

epithelial transitions, accompanied by a well-coordinated remodeling of the cell-cell and cell-matrix

junctions[10–12,38]. In our cell cultures at Day 10, we observed two cell populations with distinct features: (1)

spindly, refringent, tightly packed cells with a mesenchymal morphology and (2) larger, more flattened cells

in close contact to one another similar to squamous or cuboidal epithelia (Figure 3C). These larger cells

were first detected at Day 8 and progressively form "epithelial" islets surrounded by a "mesenchymal"

environment until Day 10 (Figure S3). We performed immunofluorescence targeting E-cadherin (E-cad), a

cell junction protein characteristic of epithelial cells and Vimentin (Vim), a cytoskeleton protein expressed

mostly in mesenchymal cells. As anticipated, the islets expressed high levels of E-Cad and were negative

for Vim, and the cells in between expressed Vim but no E-cad (Figure 3D). The fluorescent signal was

quantified in the 3 hiPSC lines at Day 10 and we found ~10% E-cad-positive cells in both the Healthy control

line and the CRISPR line, and less than 1% in the DMD line (Figure 3E). We next assessed the expression

of C-Met, a receptor tyrosine kinase expressed at the membrane of epithelial cells during dermomyotome

development from the somites and critical for subsequent myoblast migration[39]. Interestingly, we observed

a strong C-Met expression in epithelial islets, suggesting a dermomyotome identity, but no C-Met-positive

cell was found in the DMD line. Interestingly, differentiation of the CRISPR line resulted in C-Met expression

at Day 10, but at a level 5 times lower than in the healthy control (p-val = 0.029) (Figure 3F-G). This suggests

that somitogenesis and the associated cell state transitions are altered in the absence of dystrophin, but

that additional factors have an influence, depending on the genetic background.

**Somitogenesis and cell junction genes are dysregulated in three additional DMD hiPSC lines**

209            In order to increase the power of our study and extend it to additional DMD backgrounds, we

compared our findings with an RNA-Seq dataset previously generated in our group to describe the

myogenic differentiation of DMD hiPSCs at defined time points (https://muscle-dmd.omics.ovh/)[19]. More

precisely, hiPSC lines from three independent DMD patients and three healthy individuals were

differentiated in triplicates with a commercial media formulation[16] and cells were collected for bulk RNA-

Seq at several time points (Figure S4). Interestingly, the transcriptome of DMD hiPSCs was shown to

markedly diverge from healthy controls at Day 10, when somite marker genes are expressed[19].

Immunostaining at Day 10 showed the presence of epithelial islets expressing C-Met in the 6 hiPSC lines,

but at a level 4.7 lower in the 3 DMD patient lines (p = 0.0003), confirming our previous observation (Figure

4A-B). We analyzed the RNA-Seq data set and identified 1450 genes significantly dysregulated at Day 10

in the DMD cell lines (abs(log2 fold-change) ≥ 1 and adjusted p-value ≤ 0.01, Figure 4C, Table S5).

Importantly, GO analysis indicated a significant enrichment in genes involved in the formation of cell junctions (Figure 4D), which confirms their importance in the initial manifestation of DMD during somite differentiation. Particularly, desmosome organization (GO:0002934) was affected in DMD cells, with a significant downregulation of *GRHL1* (log FC = -2.5, p-val = 3.7E-06), *JUP*, (log FC = -1.0, p-val = 7.7E-04), *PERP* (log FC = -1.4, p-val = 1.2E-08), *PKP2* (log FC = -1.5, p-val = 2.6E-03) and *PKP3* (log FC = -2.9, p-val = 1.1E-03) (Figure 4E). Tight junction organization (GO:0120193) was also perturbed by the absence of dystrophin, with dysregulation of gene members of the claudin (CLDN), cadherin (CDH) and protocadherin (PCDH) families (Figure 4E). Similar to our previous findings, we noticed that most gene families were downregulated in DMD cells, with the exception of the protocadherin (*PCDH*) family, in which 8/9 genes were upregulated (*PCDH17*, *PCDHA2*, *PCDHA7*, *PCDHB18P*, *PCDHGA3*, *PCDHGA4*, *PCDHGA5*, *PCDHGB3*, mean log FC = 1.5) and only 1 was downregulated (*PCDHA13*, log FC = -1.7).

Altogether, our study combines sequencing and imaging data from four unrelated DMD patient hiPSC lines and four healthy controls plus one isogenic DMD mutant, and indicates that dystrophin deficiency leads to major dysregulations of cell junction gene families participating in cell state transitions during somite development.

## Discussion

Thanks to carefully defined differentiation protocols, hiPSCs and their derivatives can help better understand human embryonic development and investigate the early impact of mutations leading to specific genetic disorders without the use of embryos[13]. Using a published protocol recapitulating key developmental steps of human myogenesis, we provide here a temporally resolved dataset spanning ten time points along the differentiation of healthy and DMD hiPSCs from pluripotency to the skeletal muscle lineage. We observed that cells first go through intensive transcriptional changes as they leave pluripotency and adopt an early mesodermal fate. After Day 7, they stabilize as they become somite progenitors and their progenies (*i.e.* dermomyotome and skeletal muscle cells). The developmental trajectory computed from our pseudotime data showed that fractions of cells reach the end of the myogenic process as early as Day 10, reflecting the fact that cells differentiate in a heterogeneous and asynchronous manner[32,40,41].

During the second half of the differentiation process, cells that are "behind schedule" probably "catch up"

or remain stalled on an abortive developmental path. At the differentiation end stage, we observed that a

majority of cells did not express canonical myogenic markers such as *MYOD1*, *MYOG* or *PAX7*. However,

most of them were positive for the myocyte-specific enhancer factor 2C (*MEF2C*), known to be involved in

myogenesis but also in the development of a myriad of other tissues (*i.e.* heart, nervous system,

vasculature, bone and cartilage)[42]. Neural cells have already been identified as cellular by-products

generated alongside muscle cells with the myogenic differentiation protocol used in the first part of this

study[36]. For this reason, a commercial myogenic media known to produce more homogeneous cell

populations at the successive developmental steps was also used in this study[16,19,43]. This allowed us to

confirm that similar to what is seen in DMD patients, the absence of dystrophin does not compromise the

ability of hiPSCs to differentiate into myotubes. However, it may delay differentiation and / or deflect a

fraction of cells from the skeletal muscle into alternative or abortive lineages during embryogenesis, which

might go unnoticed *in vitro* due to the amplification and passage steps that are part of the differentiation

protocols.

Here, we discovered that dystrophin mutation leads to the dysregulation of cell junctions during the

successive cell state transitions which come with the *in vitro* differentiation of "somite-like" cell monolayers.

Somites are transient metameric structures emerging from the paraxial mesoderm in the embryo. They give

rise to multiple tissues such as the axial dermis, cartilages, bones and skeletal muscles of the axis and

limbs[44,45]. Individual somites are patterned along the dorsal-ventral axis into the sclerotome at the ventral

pole and the dermomyotome at the dorsal pole. Myogenic progenitors delaminate from the dermomyotome

to form primary multinucleated myofibers[12,46,47]. These developmental steps involve coordinated gene

expression and successive transitions between a mesenchymal and an epithelial state[10–12]. We showed

that the hiPSC model recapitulates key features of somitogenesis when subjected to myogenic

differentiation, both at the level of their morphology and gene expression program. As a monolayer, somite

progenitors derived from control hiPSCs spontaneously organize into discrete "epithelial islets", surrounded

by mesenchymal cells. In the embryo, somite epithelialization requires the activity of *Paraxis* and the

somites of *Paraxis-/-* murine embryos showed dysregulated somitogenesis and disorganized cell

junctions[11,48]. The activity of the Rho family small GTPases Cdc42 and Rac1 also plays an essential role in

the epithelialization during somitogenesis[49], possibly by controlling the architecture of the cytoskeleton and

the remodeling of cell junctions during the successive cell state transitions. One appealing hypothesis would

be that during somite development, the embryonic isoform of dystrophin serves as a stabilizing anchor in

newly formed epithelial cells. Testing this hypothesis will require further investigations in appropriate model

systems, as Dp412e is specific to a subgroup of anthropoids, including humans.

Multiple protein isoforms have been described from the *DMD* locus, each with a defined expression

pattern[50–52], but their function remains to be precisely characterized. In skeletal and cardiac muscles, the

Dp427m dystrophin isoform is known to interact with the dystrophin-associated protein complex (DAPC),

but in other tissues, the protein partners of other isoforms are less characterized. Recent studies have

described association of dystrophin with alternative partners in specific cells and tissues, such as aquaporin

4, calcium and potassium channels[53–57]. However, the exact molecular mechanisms driven by these

interactions reman to be investigated. Dp412e was identified as an "embryonic" isoform induced during the

formation of early mesoderm progenitors and embryoid bodies from hiPSCs[20]. Here, our results strongly

suggest that Dp412e has specific roles during embryonic development and the cell transitions occurring

when somites develop, which remain to be precisely determined.

Altogether, our study leads us to consider DMD even more as a developmental disease, even though

patients are born without apparent symptoms. A better understanding of the "invisible" DMD initiation events

during development and early postnatal life will help identify biomarkers early in the course of disease

progression, which in turn could accelerate diagnosis and pinpoint new therapeutic targets.

## Acknowledgments

This work was funded by the INSERM ATIP Avenir program, Nantes Université, the University Hospital of Nantes, the Association Française contre les Myopathies (AFM) Téléthon, Genopole, and by the Wellstone Muscular Dystrophy Cooperative Research Center supported by the National Institute of Health (NIH, grant number U54 AR065139). We thank the GenoA Genomics Core Facility and the BiRD Bioinformatics Core Facility (Nantes Université, SFR Bonamy, UMS Biocore, Biogenouest), which is part of the Institut Français de Bioinformatique (ANR-11-INBS-0013) for the access to the computing and storage infrastructure and their help with sequencing and processing the single-cell libraries. We acknowledge the IBISA MicroPICell facility (Nantes Université, SFR Bonamy, UMS Biocore, Biogenouest), member of the national infrastructure France-Bioimaging supported by the French national research agency (ANR-10-INBS-04).

## Author contributions

Conceptualization: CP, DLM, JBD; Methodology: EMo, CL, JBD; Software: VM, JSP, CT, JBD; Validation: EMo, CL, JBD; Formal analysis: EMo, CL, JBD; Investigation: EMo, EMa, VM, CL, AB, DLJ, JBD; Resources: AB, CT, OA, CP, DLM, JBD; Data Curation: EMo, VM, JSP, JBD; Writing – Original Draft: EMo, CL, JBD; Writing – Review & Editing: EMo, EMa, VM, CL, CT, CLG, OA, CP, DLM, JBD; Visualization: EMo, CL, JBD; Supervision: CT, CLG, OA, CP, DLM, JBD; Project administration: JBD; Funding acquisition: EMa, VM, OA, CP, DLM, JBD.

## Declaration of interests

The authors declare no competing interests

## Inclusion and diversity

Our research group supports inclusive, diverse and equitable conduct of research.

**Figures & Figure Legends**

**Figure 1:** Myogenic differentiation of DMD and healthy control hiPSCs at the single-cell resolution. (A) Cell

collection timeline along hiPSC myogenic differentiation using the combination of 5 defined media (M#1 to

M#5) described previously[15]. D0 to D28: Day 0 to Day 28. (B) UMAP plot showing the 1917 individual cells

colored by hiPSC line of origin. (C) Deconvolution of the UMAP plot by collection time point (D0 to D28).

(D) Single-cell trajectory capturing the dynamics of myogenic differentiation as DMD and Healthy hiPSCs

progress along the initial common branch (hiPSCs) to the bifurcation point (1) and one of the two alternative

branches. (E) Deconvolution of the single-cell trajectory by collection time point (D0 to D28). (F) Branched

expression analysis modeling identifying modules of genes with branch-dependent expression. Key Gene

Ontology terms associated with each individual module are indicated on the right-hand side. (G) Gene

expression kinetic in pseudotime for myogenic markers along the Healthy (dotted line) and the DMD (full

line) branch. Each dot shows an individual cell with its computed pseudotime value. Significant differences

in branch-dependent expression are indicated with *** (p-adj < 0.001).

**Figure 2:** DMD cells exhibit a marked dysregulation of cell junction genes at Day 10. (A) UMAP plot of the

individual cells collected at Day 10 from the sci-RNA-Seq data set (Figure 1) colored by their hiPSC line of

origin. (B-C) Gene Ontology (B) and Pathway enrichment (C) analyses performed with the 94 genes

differentially expressed in DMD cells at Day 10 in the sci-RNA-Seq data set. (D) Violin plots showing the

expression of somite marker genes in single cells from Healthy, CRISPR and DMD hiPSCs at Day 10. (E)

UMAP plot showing the 3566 individual cells colored by their hiPSC line of origin. (F) Venn diagram of the

differentially expressed genes in DMD and CRISPR hiPSC lines at Day 10. The absolute numbers of genes

are indicated in the appropriate sections. (G) Gene ontology analysis with the 1885 genes differentially

expressed in both the DMD and the CRISPR hiPSC lines at Day 10, showing a selection of significantly

enriched biological processes. Legend: * false discovery rate (FDR) < 0.05; ** FDR < 0.01; *** FDR < 0.001;

(X/Y): number of genes from the GO category found differentially expressed in the dataset / total number

of genes in the GO category. (H-I) Differential expression of cell junction genes and potential implications

at the protein level in the DMD (H) and CRISPR (I) hiPSC lines. Fold change estimates from single-cell

data are indicated between brackets and color-coded for each protein. CLDN: claudin; OCLN: occludin;

TJP: tight junction protein; CDH: cadherin; CTNN: contactin; PCDH: protocadherin; DSG: desmoglein; DSP:

desmoplakin; DSC: desmocollin; PKP: plakophilin; JUP: plakoglobin; PERP: p53 apoptosis effector related

to PMP22.

**Figure 3:** DMD does not compromise myotube formation despite altered upstream cell state transitions.

(A) Fluorescent staining of myosin heavy chains and a-actinin (red) in myotubes derived from Healthy

hiPSCs. Arrowheads indicate multinucleation and the right panel focuses on a striation pattern. (B)

Quantification of the a-actinin fluorescent area in the three cell lines, expressed as a percentage of total

area (n = 3 to 5 panels by cell line). (C) Optical microscopy with phase contrast of somite progenitors

derived from hiPSCs. Insets 1 and 2 highlight the "epithelial-like" and "mesenchymal-like" cell populations.

Scale bar = 200μm. (D) Detection of E-Cad and Vim in somite progenitors derived from hiPSCs by

immunofluorescence and confocal microscopy. (E) Quantification of the percentage of E-Cad-positive cells

in the Healthy, DMD and CRISPR hiPSC lines at Day 10 in three replicate experiments (N = 10 panels by

experiment). n.s. not significant; *** p-value < 0.001. (F) Detection of C-Met in the three hiPSC lines by

immunostaining and confocal microscopy at differentiation Day 10. (G) Quantification of the C-Met

fluorescent area normalized by the number of nuclei. Imaging was performed on four large mosaic panels

per cell line.

**Figure 4:** Somitogenesis and cell junction genes are dysregulated in three additional DMD hiPSC lines. (A)

Fluorescent staining of C-Met in DMD hiPSCs and healthy control lines at Day 10 of the myogenic

differentiation. One representative picture is shown per individual line (Healthy_1 to 3 and DMD_1 to 3). (B)

Quantification of the C-Met fluorescent area normalized by the number of nuclei. Imaging was performed

on three independent pictures per line. (C) Volcano plot of the differentially expressed genes in DMD somite

progenitor cells at Day 10 (thresholds: $|log2\ FC| > 1$ & p-adj < 0.01). (D) Gene Ontology analysis using the

differentially expressed genes as input and showing key biological processes. Legend: * false discovery

rate (FDR) < 0.05; ** FDR < 0.01; *** FDR < 0.001; (X/Y): number of genes from the GO category found

differentially expressed in the dataset / total number of genes in the GO category. (E) Differential expression

of cell junction genes and potential implications at the protein level. Fold change values (DMD vs. Healthy)

are indicated between brackets and color-coded for each protein. CLDN: claudin; OCLN: occludin; TJP:

tight junction protein; CDH: cadherin; CTNN: contactin; PCDH: protocadherin; DSG: desmoglein; DSP:

desmoplakin; DSC: desmocollin; PKP: plakophilin; JUP: plakoglobin; PERP: p53 apoptosis effector related

to PMP22.

## Material & Methods

**RESOURCE AVAILABILITY**

*Lead contact*

Further information and requests for resources and reagents should be directly addressed to and will be fulfilled by the lead contact, Jean-Baptiste Dupont (jean-baptiste.dupont@univ-nantes.fr).

*Materials availability*

This study did not generate new unique reagents

*Data and code availability*

- The two single-cell RNA-seq data sets have been deposited at GEO and are publicly available as of the date of publication. Accession numbers are listed in the key resources table. This paper also analyzes publicly available data. These accession numbers are listed in the key resources table. Original western blot images and microscopy data reported in this paper have been deposited at Zenodo repository. DOIs are listed in the key resources table.

- All original code has been deposited as R Notebook files at Zenodo Repository. DOIs are listed in the key resources table.

- Any additional information required to reanalyze the data reported in this paper is available from the lead contact upon request.

**EXPERIMENTAL MODEL AND STUDY PARTICIPANT DETAILS**

Ethical statement: The Healthy, DMD and CRISPR lines have been described, characterized and published previously[34,37]. Participants gave informed consent for the generation of urine-derived hiPSC lines, as required by the Institutional Review Board (IRB). The experiments performed in this manuscript fall into the

Category 1A described in the Guidelines for Stem Cell Research and Clinical Translation of the International

Society for Stem Cell Research (ISSCR, https://www.isscr.org/guidelines).

<u>hiPSC lines:</u> The Healthy UC3-4 line was generated from an adult male participant with no known skeletal

muscle disease. The DMD 72039 line was generated from an adult male with a declared DMD pathology

and carrying an exon50 deletion in the *DMD* gene. These individuals provided urine samples, from which

urine cells were expanded and reprogrammed to generate hiPSCs by Guan X. *et al.* at the University of

Washington, Seattle (USA)[34]. The DMD CRISPR line was generated from the UC3-4 line by CRISPR Cas

9 gene editing by Smith A. *et al.* in collaboration with the Institute for Stem Cell and Regenerative Medicine

Ellison Stem Cell Core[37]. It carries an exon 45 deletion and a 17 bp deletion in exon 54 of the *DMD* gene.

The absence of dystrophin was confirmed by Western blot after myogenic differentiation, in comparison

with the Healthy UC3-4 control line (Figure S2A). Cell lines were not otherwise authenticated. The three

hiPSC lines were extensively characterized, expanded and shared with the TaRGeT INSERM laboratory at

Nantes Université through a dedicated Material Transfer Agreement.

<u>hiPSC maintenance:</u> All cell culture experiments were performed at 37 °C and 5 % $CO_2$ in a standard tissue

culture incubator. The three hiPSC lines were expanded in the TaRGeT laboratory as Master Cell Banks

(MCB) at various passages (Healthy: passage 34; DMD: passage 25; CRISPR: passage 46). Cells were

thawed at 37°C and seeded as clusters on Matrigel-coated plates (1:60 dilution) after spinning 3 min at 300

g and resuspension in mTeSR Plus culture medium supplemented in 10 µM ROCK inhibitor. Fresh media

without ROCK inhibitor was renewed the day after seeding and then every other day. Cultures were

manually cleaned from abnormally looking clusters and were passaged with Versene when the overall

confluence reached 70 – 80 %. Working Cell Banks (WCB) were stored after two passages from the MCBs.

Cells were detached with Versene, spun down 3 min at 300 g, and clusters were gently resuspended in

Cryostor freezing media for long term cryopreservation in liquid nitrogen.

<u>Rat Muscle biopsies:</u> The protein samples used as positive controls in the dystrophin western-blot were

obtained from rat pectoral muscle biopsies from a previous study[58]. Dmd[mdx] rats and healthy controls were

handled and housed in the UTE IRS2 from Nantes Université, according to a protocol approved by the

Institutional Animal Care and Use Committee of the Région des Pays de la Loire (University of Angers,

France) as well as the French Ministry for National Education, Higher Education and Research

(authorization #2018102616384887).

**METHOD DETAILS**

hiPSC myogenic differentiation: This study combines the use of two myogenic differentiation protocols:

• Establishment of the myogenic trajectory: hiPSCs were differentiated with a succession of five

defined media, as published previously[15]. After seeding on Matrigel (1:60 dilution) as single cells at

a density of 30,000 cells / cm², hiPSCs were grown for 2 – 3 days in mTeSR Plus media. ROCK

inhibitor was added upon seeding but removed by a fresh media change after one day. The cells

were then incubated in five successive differentiation media based on Dulbecco's Modified Eagle's

Medium (DMEM) / F12 supplemented with Non-Essential Amino Acids (NEAA) and additional

molecules detailed below:

○ Media 1 (3 days): Insulin transferrin selenium (ITS) 1X + 3 µM CHIR99021 + 0.5 µM LDN

○ Media 2 (3 days): Insulin transferrin selenium (ITS) 1X + 3 µM CHIR99021 + 0.5 µM LDN

+ basic Fibroblast Growth Factor (bFGF) at 20 µg / ml

○ Media 3 (2 days): 15 % Knockout serum replacement (KSR) + 0.5 µM LDN + bFGF at 20

µg / ml + Hepatocyte Growth Factor (HGF) at 10 µg / ml + Insulin-like Growth Factor (IGF)

1 at 2 ng / ml.

○ Media 4 (4 days): 15 % KSR + IGF-1 at 2 ng / ml

○ Media 5 (16 days): 15 % KSR + IGF-1 at 2 ng / ml + HGF at 10 µg / ml

Cells were collected at Day 0, 1, 2, 4, 7, 10, 14, 18, 22 and 28 after incubation with Trypsin-EDTA

alone, or with a combination of Trypsin-EDTA + collagenase IV at 50 U / µl for 10 min at 37 °C,

mechanical dissociation and passage through a 70 µm cell strainer to remove debris and

extracellular matrix.

• Post hoc analysis of the somite differentiation stage at Day 10: in parallel, hiPSCs were

differentiated with a commercially available media previously shown as capable of reproducing the

somite stage with high accuracy[16,19]. This protocol uses lower initial seeding densities and allows

for an easier visualization and imaging of the cultures as differentiation progresses. Briefly, hiPSCs

were first amplified and dissociated as single cells as described above, and then seeded on

collagen I-coated plates at 3,500 cells / cm² in Skeletal Muscle Induction Medium (amsbio SKM01).

The media was changed every 2 to 3 days. At Day 7, cells were dissociated with Trypsin-EDTA

and cryopreserved in CryoStor CS10 medium. They were seeded on new collagen-coated plates

at 20,000 cells / cm² in SKM01 until Day 10.

single-cell RNA-Seq:

• sci-RNA-Seq: the myogenic trajectory followed by hiPSCs was determined by single-cell

combinatorial indexing RNA-Seq (sci-RNA-Seq) as previously published[35]. Cells were collected as

differentiation progressed (5 to 10 million per sample, from Day 0 to Day 28) and spiked with 10 %

murine cells of the NIH/3t3 cell line for estimation of doublet proportions. after centrifugation for 5

min at 300 g and 4 °C, pellets were washed in DPBS and resuspended in pre-chilled methanol for

fixation and permeabilization. They were then stored at – 20°C until all samples were processed at

the end of the differentiation. Fixed cells were pelleted by centrifugation (same settings), washed

twice in 1 ml DPBS + 1 % Diethylpyrocarbonate (DEPC), and another three times in 1 ml of cell

wash buffer containing 1 % SUPERase In RNase Inhibitor and 1 % BSA in ice-cold DPBS. The

final resuspension was made in 100 μl of cell wash buffer before counting with a hemocytometer.

The Reverse Transcription step was performed 10 min at 55 °C *in situ* on 2,000 cells per sample,

with the Superscript IV RT kit and barcoded oligo dT primers dispatched in a Lo-bind 96-well plate.

To increase the diversity of barcode combinations obtained after indexing, six distinct barcodes

were used for the Day 0, Day 1 and Day 2 samples and four barcodes for the remaining samples

(Day 4, Day 7, Day 10, Day 14, Day 18, Day 22, Day 28). The RT reaction was stopped with 40

mM EDTA and 1 mM spermidine (5 μl per well). Cells with barcoded cDNA were then pooled in a

flow cytometry tube, stained with 300 μM DAPI and sorted in a new 96-well Lo-bind plate (25 cells

per well) containing Elution buffer (5 μl per well). Sorting plates can be stored at – 80 °C after brief

centrifugation. Second strand synthesis was performed in each well with the NEBNext® Ultra™ II

Non-Directional RNA Second Strand Synthesis Module (0.5 μl of Buffer + 0.25 μl of enzyme per

well). After incubation at 16 °C for 150 min, the reaction was terminated at 75 °C for 20 min.

| 478 | Tagmentation was performed in each well with the illumina Tagment DNA TDE1 Enzyme (0.5 µl |
| 479 | per well) and buffer (5 µl per well) kit, after addition of human genomic DNA (0.25 µl per well). The |
| 480 | plate was incubated at 55 °C for 5 min and the reaction was stopped with 12 µl DNA binding buffer |
| 481 | per well and incubation at room temperature (RT) for 5 min. AMPure XP beads (36 µl per well) |
| 482 | were added for purification with standard protocol. The elution step was performed in a final volume |
| 483 | of 17 µl. Libraries were amplified by PCR using the NEBNext High-Fidelity 2X PCR Master Mix and |
| 484 | barcoded P5 primers (2 µl of 10 µM primer per well), in addition to a 10 µM P7 primer (2 µl per |
| 485 | well). Amplification was carried out in a standard thermal cycler with the following program: 5 min |
| 486 | at 72 °C + 30 sec at 98 °C + 18 cycles of (10 sec at 98 °C + 30 sec at 66 °C + 30 sec at 72 °C) + |
| 487 | 5 min at 72 °C. Samples were collected from each well and pooled in a single tube. The library was |
| 488 | then purified using 0.8 volume of AMPure XP beads according to manufacturer's instructions. |
| 489 | Libraries were quantified by Qubit and visualized by electrophoresis on a 6 % TBE-PAGE gel. |
| 490 | Sequencing was performed on the NextSeq 500 platform using a V2 75 cycles kit and the following |
| 491 | settings: Read 1: 18 cycles, Read 2: 52 cycles, Index 1: 10 cycles, Index 2: 10 cycles. Here, two |
| 492 | sorting plates were generated and used to generated independent libraries sequenced |
| 493 | successively with distinct P7 primers. The first sorting plate was full and thus 96 barcoded P5 |
| 494 | primers were used in the PCR amplification step; the second sorting plate only used 86 wells (and |
| 495 | thus 86 barcoded P5 primers). The sequences of the barcoded primers used in this protocol can |
| 496 | be found in Table S6. Raw data analysis involved base calling (bcl2fastq), demultiplexing based |
| 497 | on P5 barcodes (1 mismatched base allowed), adaptor trimming (trim_galore), alignment to the |
| 498 | human (hg19) or the mouse (mm10) genome (STAR), removing of UMI duplicates and |
| 499 | demultiplexing based on RT barcodes. Percentages of reads mapping uniquely to the human and |
| 500 | the mouse genome were quantified and cells with over 90 % of reads assigned to the human |
| 501 | genome were kept for subsequent analysis. Secondary analysis of the cell data set object was |
| 502 | performed with the Monocle 3 analysis pipeline (Trapnell 2014, Qiu 2017), whose code is freely |
| 503 | available on the Cole Trapnell lab Github and in the Zenodo Repository with the DOI listed in the |
| 504 | key resources table. |

• split-pool barcoding: the *post hoc* analysis of the somite differentiation step at Day 10 was
performed with the split-pool barcoding kit commercialized by parse Biosciences, according to
manufacturer's instructions. Cells were collected at Day 10 and counted with a hemocytometer to
isolate 500,000 cells for subsequent fixation and permeabilization with the Cell Fixation Kit.
Samples were stored at − 80°C and thawed immediately prior to library preparation with the
Evercode Whole Transcriptome Mini Kit. Primary analysis of the raw data including quality control,
alignment to the human genome (hg19), demultiplexing and generation of the matrix, feature
annotation and cell annotation files were carried out by the proprietary Parse Biosciences analysis
suite. Secondary analysis used the Monocle 3 pipeline as previously indicated. The code has been
deposited at Zenodo Repository with the DOI listed in the key resources table.

Immunofluorescence:

• Somite progenitors: hiPSCs were first differentiated into somite progenitor cells with the amsbio
commercial protocol to constitute a working cell bank at Day 7. Subsequently, cryotubes containing
one million progenitors were thawed and seeded at 23,500 cells/cm² on Lab-Tek™ 4-well chamber
slides (Thermofisher Cat# 154526PK) coated with collagen I (1:60 dilution). Cells were maintained
in Skeletal Muscle Induction medium supplemented with 2% Pen/Strep and incubated at 37°C, 5%
$CO_2$ until confluency.

• Myotubes: myogenic progenitors were differentiated from hiSPCs with the amsbio commercial
protocol and banks were made at Day 17. Subsequently, cryotubes containing one million
progenitors were thawed and seeded at 23,500 cells/cm² on Lab-Tek™ 4-well chamber slides
(Thermofisher Cat# 154526PK) coated with collagen I (1:60 dilution). They were maintained in
Skeletal Muscle Myoblast medium supplemented with 2% Pen/Strep and incubated at 37°C, 5%
$CO_2$ until confluency. Differentiation was induced with the Skeletal Muscle Myotube medium
supplemented with 2% Pen/Strep for 8 days, with a fresh media change every 2 to 3 days.

Cells were fixed in 4 % paraformaldehyde (PFA) for 1 hr and then permeabilized in PBS + Triton 1X + 2.5%
bovine serum albumin (BSA). Primay antibodies were diluted in permeabilization buffer and incubated on

the cells overnight at 4 °C (E-Cadherin Alexa Fluor™ 488, Vimentin eFluor™ 570 and C-Met: 1:50 dilution;

Myosin Heavy Chain: 1:300 dilution; $\alpha$-actinin: 1:500 dilution). The next day, nuclei were stained with

1:10.000 DAPI and for myotubes, the anti-mouse or anti-goat (C-Met) secondary antibodies were diluted

at 1:1000 in PBS and added for 1 hr at RT. Coverslips were mounted on the slides using ProLong Gold

Antifade Reagent after removing the Lab-Tek™ walls. Images were taken with a 20 X oil immersion

objective on a NIKON® A1 RSi confocal microscope.

<u>Western-blot:</u> Proteins were extracted from frozen cell pellets in RIPA buffer during 1 hour (10 mM Tris +

150 Mm NaCl + 1mM protease inhibitor cocktails + 1% Igepal + 0.1% SDS). Protein extracts diluted at 1:10

were quantified with the DC Protein Assay kit at 750 nm on the Thermo Scientific™ Multiskan™ GO. Protein

samples (50 µg per sample) were denatured by addition of NuPAGE LDS sample buffer 4X and

Dithiothreitol (DTT), and loaded onto a Nupage 3-8% TA Gel before migrating for 2 hrs at 100 V in 1X

NuPAGE tris acetate SDS buffer. Protein extracts from Healthy and DMD$^{mdx}$ rat pectoral muscles were

loaded as controls. The Bio-Rad Trans-Blot Turbo Transfer System was used for protein transfer. After

overnight saturation at 4 °C in saturation buffer (5 % milk + 0.1 % Tween 20 + 1 % NP40), membranes

were incubated with primary Mouse anti-dystrophin NCL-DYS2 antibody diluted at 1:250 for 1 hr. The $\alpha$-

tubulin protein was labeled with a Mouse Monoclonal Anti-α-Tubulin antibody diluted at 1:10000. All

membranes were then washed three times for 5 minutes in PBS + 0.1 % Tween 20 before incubation with

the corresponding secondary antibodies (Goat Anti-Mouse antibody/HRP diluted at 1 :5000 during 1 hr).

The ECL kit was used for detection of the HRP enzyme activity, with exposure times of 2 minutes ($\alpha$-tubulin)

and 2 hours (Dystrophin).

**QUANTIFICATION AND STATISTICAL ANALYSIS**

<u>Immunofluorescence:</u> Each hiPSC line was differentiated and imaged as 4 by 4 panels or individual

pictures, and analyzed with the QuPath software. C-Met and Ecad-positive areas were determined by

thresholding on the Alexa488 fluorescent channel (ECad: Threshold = 40 or regions manually drawn if low

signal, C-Met: Threshold = 600, minimal object size= 2,500 µm²), and the number of nuclei was determined

in the positive and negative areas using the Cell Detection tool with the following parameters: C-Met: DAPI

threshold = 100, object size between 10 and 400 µm², background radius = 8 µm, sigma = 1.5 µm; E-Cad:

DAPI threshold = 100, object size between 75 and 2000 µm², background radius = 8 µm, sigma = 2 µm. E-

Cad pictures were acquired at higher magnification, explaining why different parameters had to be used.

MF20- and a-actinin-positive areas were determined by thresholding on the TRITC channel (MF20:

threshold = 100, $\alpha$-actinin: threshold = 1000). Data were represented and analyzed with GraphPad PRISM

8.0.1. Non-parametric statistical tests were used to compare groups as sample size were low and we could

not assume gaussian distributions nor homoscedasticity. The Mann-Whitney statistics was used when only

two groups were compared, and the Kruskal-Wallis statistics for comparison of the 3 groups.

**KEY RESOURCES TABLE**

| REAGENT or RESOURCE | SOURCE | IDENTIFIER |
|---|---|---|
| Antibodies | | |
| Myosin heavy chain | DSHB | Cat# MF20 RRID:AB_2147781 |
| alpha-Actinin (Sarcomeric) | Sigma-Aldrich | Cat# A7811; RRID:AB_476766 |
| CD324 (E-cadherin) Alexa Fluor™ 488 | Thermo Fisher | Cat# 53-3249-80 RRID:AB_10671270 |
| Vimentin eFluor™ 570 | Thermo Fisher | Cat# 41-9897-80 RRID:AB_11220476 |
| Human HGFR/c-MET | R&D Systems | Cat# AF276 RRID:AB_355289 |
| Goat anti-Mouse IgG2b Alexa Fluor™ 555 | Thermo Fisher | Cat# A-21147 RRID:AB_2535783 |
| Donkey anti-Goat IgG Alexa Fluor™ 488 | Thermo Fisher | Cat# A-11055 RRID:AB_2534102 |
| Dystrophin | Leica | Cat# NCL-DYS2 RRID:AB_442081 |
| GAPDH | Novus | Cat# NB300-320 RRID:AB_10001796 |
| Alpha tubulin | Sigma-Aldrich | Cat# T5168 RRID:AB_477579 |
| Goat Anti-Mouse Immunoglobulins/HRP | Agilent | Cat# P0447 RRID:AB_2617137 |
| Polyclonal Rabbit Anti-Goat Immunoglobulins/HRP | Agilent | Cat# P0449 RRID:AB_2617143 |
| Chemicals, peptides, and recombinant proteins | | |
| mTeSR™ Plus | STEMCELL Technologies | Cat# 100-0276 |
| Matrigel® Matrix | Corning | Cat# 354234 |
| Versene solution | Thermo Fisher | Cat# 15040066 |
| CryoStor CS10 cryopreservation medium | STEMCELL Technologies | Cat# 07959 |
| DPBS | Thermo Fisher | Cat# 14190144 |
| Y-27632 ROCK inhibitor | Selleckchem | Cat# S1049 |

| DMEM/F12 | Thermo Fisher | Cat# 11320033 |
|---|---|---|
| CHIR99021 | STEMCELL Technologies | Cat# 72054 |
| StemMACS™ LDN-193189 | Miltenyi Biotec | Cat# 130-103-925 |
| Recombinant Human FGF2 | R&D Systems | Cat# 233-FB-010 |
| Recombinant Human IGF-1 | R&D Systems | Cat# 291-G1-200 |
| Recombinant Human HGF | R&D Systems | Cat# 294-HG-005 |
| KnockOut™ serum replacement | Thermo Fisher | Cat# 10828010 |
| Collagenase, type IV | Thermo Fisher | Cat# 1710419 |
| Trypsin-EDTA | Sigma-Aldrich | Cat# T4174 |
| Skeletal Muscle. Induction Medium | amsbio | Cat# SKM01 |
| Skeletal Muscle. Myoblast Medium | amsbio | Cat# SKM02 |
| Skeletal Muscle. Myotube Medium | Amsbio | Cat# SKM03 |
| Cultrex Rat Collagen I | R&D Systems | Cat# 3440-100-01 |
| Paraformaldehyde | Thermo Fisher | Cat# #28906 |
| Triton 100X | Eurobio | Cat# GAUTTR001 |
| BSA (immunofluorescence) | Sigma-Aldrich | Cat# A3059 |
| ProLong™ Gold Antifade reagent | Thermo Fisher | Cat# P36934 |
| Tris base | Sigma-Aldrich | Cat# 10708976001 |
| cOmplete™ Protease Inihibitor Cocktail | Merck | Cat# 11873580001 |
| Protease inhibitor cocktail | Sigma-Aldrich | Cat# P8340 |
| Igepal CA-630 | Sigma-Aldrich | Cat# I3021 |
| SDS solution | Sigma-Aldrich | Cat# 05030 |
| NuPAGE™ LDS sample buffer | Thermo Fisher | Cat# NP0007 |
| Dithiothreitol | Thermo Fisher | Cat# D1532 |
| NuPAGE™ Tris-acetate Gel | Thermo Fisher | Cat# EA03785BOX |
| NuPAGE™ Tris acetate SDS buffer | Thermo Fisher | Cat# LA0041 |
| Tween 20 | Merck | Cat# 8170721000 |
| Pierce™ ECL Western Blotting Substrate | Thermo Fisher | Cat# 32106 |
| Nuclease free water | Ambion | Cat# AM9932 |
| Methanol | Fisher Scientific | Cat# 10285131 |
| SUPERase In RNase inhibitor | Invitrogen | Cat# AM2694 |
| BSA (single-cell RNA-Seq) | New England Biolabs | Cat# B9000 |
| 1M Tris-HCl pH 7.5 | Thermo Fisher | Cat# 15567027 |
| 5M NaCl (single-cell RNA-Seq) | Thermo Fisher | Cat# AM9759 |
| 1M MgCl2 | Thermo Fisher | Cat# AM9530G |
| IGEPAL CA-360 | Sigma-Aldrich | Cat# 18896 |
| Diethyl pyrocarbonate | Sigma-Aldrich | Cat# D5758 |
| 10mM dNTP | Thermo Fisher | Cat# 18427013 |
| Superscript IV reverse transcriptase | Thermo Fisher | Cat# 18090200 |
| RNaseOUT Recombinant Ribonuclease Inhibitor | Thermo Fisher | Cat#10777019 |
| 0.5M EDTA pH 8.0 | Thermo Fisher | Cat# 15575020 |
| Spermidine | MP Biomedicals | Cat# 0219485201 |
| Elution Buffer | Qiagen | Cat# 19086 |
| DAPI | Thermo Fisher | Cat# D1306 |
| DNA binding buffer | Zymo Research | Cat# D4003-1 |
| AMPure XP beads | Beckman Coulter | Cat# A63880 |
| Ethanol | Decon Labs | Cat# 2716 |
| NEBNext® High Fidelity 2X PCR Master Mix | New England Biolabs | Cat# M0541 |

| Novex® TBE-PAGE gels, 6% | Thermo Fisher | Cat# EC6265BOX |
|---|---|---|
| Quick-Load 2-log DNA Ladder | New England Biolabs | Cat# N0550 |
| | | |
| **Critical commercial assays** | | |
| Nextseq 500/550 V2 75 cycle kit | illumina | Cat# 20024906 |
| DC Protein Assay | Bio-Rad | Cat# 5000111 |
| Tagment DNA TDE1 Enzyme and Buffer Kit | illumina | Cat# 20034197 |
| Evercode™ Whole Transcriptome Mini Kit | Parse Biosciences | Cat# EC-W01010 |
| Cell Fixation Kit | Parse Biosciences | Cat# SB1001 |
| Qubit dsDNA HS Kit | Thermo Fisher | Cat# Q32851 |
| NEBNext® Ultra™ II Non-Directional RNA Second Strand Synthesis Module | New England Biolabs | Cat# E6111 |
| | | |
| **Deposited data** | | |
| sciRNA-Seq myogenesis time course data | This paper | GEO accession number: GSE233605 |
| Bulk RNA Seq Day 10 data | Mournetas et al.[1] | ArrayExpress: E-MTAB-8321 |
| Split-pool barcoding scRNASeq Day 10 data | This paper | GEO accession number: GSE233606 |
| Supplementary Tables S1 to S6 | This paper | Zenodo DOI: 10.5281/zenodo.8414351 |
| Full Western blots films related to Figure S2A | This paper | Zenodo DOI: 10.5281/zenodo.7937310 |
| Immunofluorescence pictures related to Figure 3A-B | This paper | Zenodo DOI: 10.5281/zenodo.8325281 |
| Immunofluorescence pictures related to Figure 3D-E | This paper | Zenodo DOI: 10.5281/zenodo.7950222 |
| Immunofluorescence pictures related to Figure 3F-G | This paper | Zenodo DOI: 10.5281/zenodo.8325290 |
| Immunofluorescence pictures related to Figure 4A-B | This paper | Zenodo DOI: 10.5281/zenodo.8325317 |
| **Experimental models: Cell lines** | | |
| UC3-4 Healthy hiPSC line, passage 32 | Laboratory of David L Mack | N/A |
| DMD 72039 D3 hiPSC line, passage 22 | Laboratory of David L Mack | N/A |
| UC3-4 DMD CRISPR hiPSC line, passage 41 | Laboratory of David L Mack | N/A |

| NIH/3T3 cell spike (single-cell RNA-Seq) | ATCC | N/A |
|---|---|---|
| **Oligonucleotides** | | |
| sciRNA-Seq barcoded oligonucleotides (see Table S1) | IDT | N/A |
| Custom P5 and P7 primers (see Table S1) | IDT | N/A |
| | | |
| **Software and algorithms** | | |
| Fiji (imageJ) | Schindelin *et al.*[2] | https://fiji.sc/ |
| QuPath version 0.4.3 | Bankhead *et al.*[3] | https://qupath.github.io/ |
| PRISM version 8 | Graphpad | https://www.graphpad.com/features |
| bcl2fastq version 2.16.0.10 | illumina | https://emea.support.illumina.com/sequencing/sequencing_software/bcl2fastq-conversion-software.html |
| trim_galore version 0.4.5 | Felix Krueger | https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/ |
| STAR version 2.5.2b | Dobin *et al.*[4] | https://github.com/alexdobin/STAR |
| R version 4.2.2 | https://cran.r-project.org/ | |
| dplyr version 1.1.0 | Wickham *et al.* | https://dplyr.tidyverse.org/ |
| monocle version 1.3.1 | Cao *et al.*[5,6] | https://cole-trapnell-lab.github.io/monocle3/ |
| ggplot2 version 3.1.4 | Wickham *et al.* | https://ggplot2.tidyverse.org/ |
| sci-RNA-Seq and split-pool barcoding analysis pipelines | This paper | Zenodo DOI : 10.5281/zenodo.8348000 |

## Supplemental references:

1.      Mournetas, V., Massouridès, E., Dupont, J.-B., Kornobis, E., Polvèche, H., Jarrige, M., Dorval, A.R.L., Gosselin, M.R.F., Manousopoulou, A., Garbis, S.D., et al. (2021). Myogenesis modelled by human pluripotent stem cells: a multi-omic study of Duchenne myopathy early onset. J Cachexia Sarcopenia Muscle *12*, 209–232. 10.1002/jcsm.12665.

2.      Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source platform for biological-image analysis. Nat Methods *9*, 676–682. 10.1038/nmeth.2019.

3.      Bankhead, P., Loughrey, M.B., Fernández, J.A., Dombrowski, Y., McArt, D.G., Dunne, P.D., McQuaid, S., Gray, R.T., Murray, L.J., Coleman, H.G., et al. (2017). QuPath: Open source software for digital pathology image analysis. Sci Rep *7*, 16878. 10.1038/s41598-017-17204-5.

4.      Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner. Bioinformatics *29*, 15–21. 10.1093/bioinformatics/bts635.

5.      Cao, J., Packer, J.S., Ramani, V., Cusanovich, D.A., Huynh, C., Daza, R., Qiu, X., Lee, C., Furlan, S.N., Steemers, F.J., et al. (2017). Comprehensive single-cell transcriptional profiling of a multicellular organism. Science *357*, 661–667. 10.1126/science.aam8940.

6.      J, C., M, S., X, Q., X, H., Dm, I., Aj, H., F, Z., S, M., L, C., Fj, S., et al. (2019). The single-cell transcriptional landscape of mammalian organogenesis. Nature *566*. 10.1038/s41586-019-0969-x.

# Supplemental information

589    **Supplemental Figure S1:** Myogenic differentiation of hiPSCs at the single-cell resolution. (A) Observation

590    of hiPSC cultures along the differentiation at process by phase contrast microscopy. (B) UMAP plot showing

591    the 1917 individual cells colored by cluster (left panel) and by collection time point from Day 0 to D28 (right

592    panel). (C) Expression of successive developmental markers in hiPSCs along the myogenic differentiation.

593    (D) Number of DMD and Healthy cells on the three branches of the developmental trajectory.

**Supplemental Figure S2:** (A) Expression of dystrophin along the myogenic differentiation of CRISPR hiPSCs (right panel) and healthy control cells (left panel). Proteins were extracted from cell samples collected from Day 0 to Day 25 or from the pectoral muscles of a healthy rat as positive controls ("Muscle") and a Dmd$^{mdx}$ rat as a negative control ("-"), and subjected to anti-dystrophin western-blot analysis. Alpha tubulin (a-tubulin) was used as a control. (B) Pseudotime expression of *CDH* and *PCDH* along the Healthy and the DMD branches in the sci-RNA-Seq data set. Individual cells are colored by their hiPSC line of origin and ordered along the pseudotime axis. BEAM statistics: \*\*\*: adjusted p-value < 0.001.

**Supplemental Figure S3:** Evolution of hiPSC cultures between Day 7 and Day 10 of the myogenic

differentiation protocol. Arrowheads indicate "round" cells appearing from Day 8 and at the core of the

epithelial islets visible from Day 9 – Day 10 (white rectangle).

**Supplemental Figure S4:** Differentiation of DMD and Healthy hiPSCs and cell collection protocol in the

analyzed study[19]. Each individual line is indicated by a blue (Healthy) or a red (DMD) dot, and the analysis

at Day 10 is highlighted by a dotted square.

**Supplemental Table S1:** cell_data file obtained after the initial sciRNA-Seq experiment. The genotype,

time point, number of expressed genes and cluster of origin are indicated for each individual cell.

**Supplemental Table S2:** Differentially expressed genes (p-adj < 0.0001) obtained after branch expression

analysis modeling (BEAM) and the cluster to which they belong on the heatmap shown in Figure 1F.

**Supplemental Table S3:** Differentially expressed genes (p-adj < 0.01) detected at Day 10 after sciRNA-

Seq.

**Supplemental Table S4:** Differentially expressed genes (p-adj < 0.01) detected at Day 10 after split-pool

barcoding RNA-Seq.

**Supplemental Table S5:** Differentially expressed genes (abs(log2(fold-change)) > 1 & p-adj < 0.01)

detected at Day 10 after bulk RNA-Seq in our previous study[19].

**Supplemental Table S6:** List of barcoded primers used in the RT reaction and in the illumine library

preparation of the sciRNA-Seq experiment.

The Supplementary Tables S1 to S6 have been deposited on the Zenodo database, and are available using

the following DOI: 10.5281/zenodo.8414351

## References:

1.      Bushby, K., Finkel, R., Birnkrant, D.J., Case, L.E., Clemens, P.R., Cripe, L., Kaul, A., Kinnett, K., McDonald, C., Pandya, S., et al. (2010). Diagnosis and management of Duchenne muscular dystrophy, part 1: diagnosis, and pharmacological and psychosocial management. Lancet Neurol *9*, 77–93. 10.1016/S1474-4422(09)70271-6.

2.      Wang, C.H., Bonnemann, C.G., Rutkowski, A., Sejersen, T., Bellini, J., Battista, V., Florence, J.M., Schara, U., Schuler, P.M., Wahbi, K., et al. (2010). Consensus statement on standard of care for congenital muscular dystrophies. J. Child Neurol. *25*, 1559–1581. 10.1177/0883073810381924.

3.      Wang, C.H., Dowling, J.J., North, K., Schroth, M.K., Sejersen, T., Shapiro, F., Bellini, J., Weiss, H., Guillet, M., Amburgey, K., et al. (2012). Consensus statement on standard of care for congenital myopathies. J. Child Neurol. *27*, 363–382. 10.1177/0883073812436605.

4.      Duan, D., Goemans, N., Takeda, S., Mercuri, E., and Aartsma-Rus, A. (2021). Duchenne muscular dystrophy. Nat Rev Dis Primers *7*, 1–19. 10.1038/s41572-021-00248-3.

5.      Nguyen, F., Cherel, Y., Guigand, L., Goubault-Leroux, I., and Wyers, M. (2002). Muscle lesions associated with dystrophin deficiency in neonatal golden retriever puppies. J. Comp. Pathol. *126*, 100–108. 10.1053/jcpa.2001.0526.

6.      Merrick, D., Stadler, L.K.J., Larner, D., and Smith, J. (2009). Muscular dystrophy begins early in embryonic development deriving from stem cell loss and disrupted skeletal muscle formation. Dis Model Mech *2*, 374–388. 10.1242/dmm.001008.

7.      Toop, J., and Emery, A.E. (1974). Muscle histology in fetuses at risk for Duchenne muscular dystrophy. Clin. Genet. *5*, 230–233.

8.      Emery, A.E. (1977). Muscle histology and creatine kinase levels in the foetus in Duchenne muscular dystrophy. Nature *266*, 472–473. 10.1038/266472a0.

9.      Pourquié, O., Al Tanoury, Z., and Chal, J. (2018). The Long Road to Making Muscle In Vitro. Curr Top Dev Biol *129*, 123–142. 10.1016/bs.ctdb.2018.03.003.

10.     Linker, C., Lesbros, C., Gros, J., Burrus, L.W., Rawls, A., and Marcelle, C. (2005). beta-Catenin-dependent Wnt signalling controls the epithelial organisation of somites through the activation of paraxis. Development *132*, 3895–3905. 10.1242/dev.01961.

11.     Rowton, M., Ramos, P., Anderson, D.M., Rhee, J.M., Cunliffe, H.E., and Rawls, A. (2013). Regulation of mesenchymal-to-epithelial transition by PARAXIS during somitogenesis. Developmental Dynamics *242*, 1332–1344. 10.1002/dvdy.24033.

12.     Zhou, Y., Zhang, Y., and Zhu, D. (2018). Myostatin promotes the epithelial-to-mesenchymal transition of the dermomyotome during somitogenesis. Developmental Dynamics *247*, 1241–1252. 10.1002/dvdy.24681.

13.     Avior, Y., Sagi, I., and Benvenisty, N. (2016). Pluripotent stem cells in disease modelling and drug discovery. Nat Rev Mol Cell Biol *17*, 170–182. 10.1038/nrm.2015.27.

14.     Karagiannis, P., Takahashi, K., Saito, M., Yoshida, Y., Okita, K., Watanabe, A., Inoue, H., Yamashita, J.K., Todani, M., Nakagawa, M., et al. (2019). Induced Pluripotent Stem Cells

and Their Use in Human Models of Disease and Development. Physiological Reviews *99*, 79–114. 10.1152/physrev.00039.2017.

15.     Chal, J., Al Tanoury, Z., Hestin, M., Gobert, B., Aivio, S., Hick, A., Cherrier, T., Nesmith, A.P., Parker, K.K., and Pourquié, O. (2016). Generation of human muscle fibers and satellite-like cells from human pluripotent stem cells in vitro. Nat Protoc *11*, 1833–1850. 10.1038/nprot.2016.110.

16.     Caron, L., Kher, D., Lee, K.L., McKernan, R., Dumevska, B., Hidalgo, A., Li, J., Yang, H., Main, H., Ferri, G., et al. (2016). A Human Pluripotent Stem Cell Model of Facioscapulohumeral Muscular Dystrophy-Affected Skeletal Muscles. Stem Cells Transl Med *5*, 1145–1161. 10.5966/sctm.2015-0224.

17.     Choi, I.Y., Lim, H., Estrellas, K., Mula, J., Cohen, T.V., Zhang, Y., Donnelly, C.J., Richard, J.-P., Kim, Y.J., Kim, H., et al. (2016). Concordant but Varied Phenotypes among Duchenne Muscular Dystrophy Patient-Specific Myoblasts Derived using a Human iPSC-Based Model. Cell Rep *15*, 2301–2312. 10.1016/j.celrep.2016.05.016.

18.     Xi, H., Fujiwara, W., Gonzalez, K., Jan, M., Liebscher, S., Van Handel, B., Schenke-Layland, K., and Pyle, A.D. (2017). In Vivo Human Somitogenesis Guides Somite Development from hPSCs. Cell Rep *18*, 1573–1585. 10.1016/j.celrep.2017.01.040.

19.     Mournetas, V., Massouridès, E., Dupont, J.-B., Kornobis, E., Polvèche, H., Jarrige, M., Dorval, A.R.L., Gosselin, M.R.F., Manousopoulou, A., Garbis, S.D., et al. (2021). Myogenesis modelled by human pluripotent stem cells: a multi-omic study of Duchenne myopathy early onset. J Cachexia Sarcopenia Muscle *12*, 209–232. 10.1002/jcsm.12665.

20.     Massouridès, E., Polentes, J., Mangeot, P.-E., Mournetas, V., Nectoux, J., Deburgrave, N., Nusbaum, P., Leturcq, F., Popplewell, L., Dickson, G., et al. (2015). Dp412e: a novel human embryonic dystrophin isoform induced by BMP4 in early differentiated cells. Skelet Muscle *5*, 40. 10.1186/s13395-015-0062-6.

21.     J, C., M, S., X, Q., X, H., Dm, I., Aj, H., F, Z., S, M., L, C., Fj, S., et al. (2019). The single-cell transcriptional landscape of mammalian organogenesis. Nature *566*. 10.1038/s41586-019-0969-x.

22.     Xi, H., Langerman, J., Sabri, S., Chien, P., Young, C.S., Younesi, S., Hicks, M., Gonzalez, K., Fujiwara, W., Marzi, J., et al. (2020). A Human Skeletal Muscle Atlas Identifies the Trajectories of Stem and Progenitor Cells across Development and from Human Pluripotent Stem Cells. Cell Stem Cell *27*, 158-176.e10. 10.1016/j.stem.2020.04.017.

23.     Giordani, L., He, G.J., Negroni, E., Sakai, H., Law, J.Y.C., Siu, M.M., Wan, R., Corneau, A., Tajbakhsh, S., Cheung, T.H., et al. (2019). High-Dimensional Single-Cell Cartography Reveals Novel Skeletal Muscle-Resident Cell Populations. Mol Cell *74*, 609-621.e6. 10.1016/j.molcel.2019.02.026.

24.     Dos Santos, M., Backer, S., Saintpierre, B., Izac, B., Andrieu, M., Letourneur, F., Relaix, F., Sotiropoulos, A., and Maire, P. (2020). Single-nucleus RNA-seq and FISH identify coordinated transcriptional activity in mammalian myofibers. Nat Commun *11*, 5102. 10.1038/s41467-020-18789-8.

25.    De Micheli, A.J., Spector, J.A., Elemento, O., and Cosgrove, B.D. (2020). A reference
single-cell transcriptomic atlas of human skeletal muscle tissue reveals bifurcated muscle stem
cell populations. Skelet Muscle *10*, 19. 10.1186/s13395-020-00236-3.

26.    Kimmel, J.C., Yi, N., Roy, M., Hendrickson, D.G., and Kelley, D.R. (2021). Differentiation
reveals latent features of aging and an energy barrier in murine myogenesis. Cell Rep *35*,
109046. 10.1016/j.celrep.2021.109046.

27.    Mj, P., Co, S., C, S., K, C., X, C., Mt, W., N, S., and Dp, M. (2020). Single-nucleus RNA-
seq identifies transcriptional heterogeneity in multinucleated skeletal myofibers. Nature
communications *11*. 10.1038/s41467-020-20063-w.

28.    Chemello, F., Wang, Z., Li, H., McAnally, J.R., Liu, N., Bassel-Duby, R., and Olson, E.N.
(2020). Degenerative and regenerative pathways underlying Duchenne muscular dystrophy
revealed by single-nucleus RNA sequencing. Proceedings of the National Academy of Sciences
*117*, 29691–29701. 10.1073/pnas.2018391117.

29.    Saleh, K.K., Xi, H., Switzler, C., Skuratovsky, E., Romero, M.A., Chien, P., Gibbs, D.,
Gane, L., Hicks, M.R., Spencer, M.J., et al. (2022). Single cell sequencing maps skeletal muscle
cellular diversity as disease severity increases in dystrophic mouse models. iScience *25*,
105415. 10.1016/j.isci.2022.105415.

30.    Scripture-Adams, D.D., Chesmore, K.N., Barthélémy, F., Wang, R.T., Nieves-Rodriguez,
S., Wang, D.W., Mokhonova, E.I., Douine, E.D., Wan, J., Little, I., et al. (2022). Single nuclei
transcriptomics of muscle reveals intra-muscular cell dynamics linked to dystrophin loss and
rescue. Commun Biol *5*, 989. 10.1038/s42003-022-03938-0.

31.    Taglietti, V., Kefi, K., Bronisz-Budzyńska, I., Mirciloglu, B., Rodrigues, M., Cardone, N.,
Coulpier, F., Periou, B., Gentil, C., Goddard, M., et al. (2022). Duchenne muscular dystrophy
trajectory in R-DMDdel52 preclinical rat model identifies COMP as biomarker of fibrosis. Acta
Neuropathol Commun *10*, 60. 10.1186/s40478-022-01355-2.

32.    Trapnell, C., Cacchiarelli, D., Grimsby, J., Pokharel, P., Li, S., Morse, M., Lennon, N.J.,
Livak, K.J., Mikkelsen, T.S., and Rinn, J.L. (2014). The dynamics and regulators of cell fate
decisions are revealed by pseudotemporal ordering of single cells. Nat. Biotechnol. *32*, 381–
386. 10.1038/nbt.2859.

33.    Qiu, X., Mao, Q., Tang, Y., Wang, L., Chawla, R., Pliner, H.A., and Trapnell, C. (2017).
Reversed graph embedding resolves complex single-cell trajectories. Nat. Methods *14*, 979–
982. 10.1038/nmeth.4402.

34.    Guan, X., Mack, D.L., Moreno, C.M., Strande, J.L., Mathieu, J., Shi, Y., Markert, C.D.,
Wang, Z., Liu, G., Lawlor, M.W., et al. (2014). Dystrophin-deficient cardiomyocytes derived from
human urine: new biologic reagents for drug discovery. Stem Cell Res *12*, 467–480.
10.1016/j.scr.2013.12.004.

35.    Cao, J., Packer, J.S., Ramani, V., Cusanovich, D.A., Huynh, C., Daza, R., Qiu, X., Lee,
C., Furlan, S.N., Steemers, F.J., et al. (2017). Comprehensive single-cell transcriptional profiling
of a multicellular organism. Science *357*, 661–667. 10.1126/science.aam8940.

36.    Kim, J., Magli, A., Chan, S.S.K., Oliveira, V.K.P., Wu, J., Darabi, R., Kyba, M., and
Perlingeiro, R.C.R. (2017). Expansion and Purification Are Critical for the Therapeutic

Application of Pluripotent Stem Cell-Derived Myogenic Progenitors. Stem Cell Reports *9*, 12–22.
10.1016/j.stemcr.2017.04.022.

37.     Smith, A.S., Luttrell, S.M., Dupont, J.-B., Gray, K., Lih, D., Fleming, J.W., Cunningham,
N.J., Jepson, S., Hesson, J., Mathieu, J., et al. (2022). High-throughput, real-time monitoring of
engineered skeletal muscle function using magnetic sensing. J Tissue Eng *13*,
20417314221122127. 10.1177/20417314221122127.

38.     Thiery, J.P., Acloque, H., Huang, R.Y.J., and Nieto, M.A. (2009). Epithelial-
Mesenchymal Transitions in Development and Disease. Cell *139*, 871–890.
10.1016/j.cell.2009.11.007.

39.     Dietrich, S., Abou-Rebyeh, F., Brohmann, H., Bladt, F., Sonnenberg-Riethmacher, E.,
Yamaai, T., Lumsden, A., Brand-Saberi, B., and Birchmeier, C. (1999). The role of SF/HGF and
c-Met in the development of skeletal muscle. Development *126*, 1621–1629.

40.     Zhang, H., and Miller, R.H. (1995). Asynchronous differentiation of clonally related spinal
cord oligodendrocytes. Mol Cell Neurosci *6*, 16–31. 10.1006/mcne.1995.1003.

41.     Pauklin, S., and Vallier, L. (2013). The Cell-Cycle State of Stem Cells Determines Cell
Fate Propensity. Cell *155*, 135–147. 10.1016/j.cell.2013.08.031.

42.     Dong, C., Yang, X.-Z., Zhang, C.-Y., Liu, Y.-Y., Zhou, R.-B., Cheng, Q.-D., Yan, E.-K.,
and Yin, D.-C. (2017). Myocyte enhancer factor 2C and its directly-interacting proteins: A
review. Progress in Biophysics and Molecular Biology *126*, 22–30.
10.1016/j.pbiomolbio.2017.02.002.

43.     Mérien, A., Tahraoui-Bories, J., Cailleret, M., Dupont, J.-B., Leteur, C., Polentes, J.,
Carteron, A., Polvèche, H., Concordet, J.-P., Pinset, C., et al. (2021). CRISPR gene editing in
pluripotent stem cells reveals the function of MBNL proteins during human in vitro myogenesis.
Hum Mol Genet, ddab218. 10.1093/hmg/ddab218.

44.     Brent, A.E., and Tabin, C.J. (2002). Developmental regulation of somite derivatives:
muscle, cartilage and tendon. Current Opinion in Genetics & Development *12*, 548–557.
10.1016/S0959-437X(02)00339-8.

45.     Weldon, S.A., and Münsterberg, A.E. (2022). Somite development and regionalisation of
the vertebral axial skeleton. Seminars in Cell & Developmental Biology *127*, 10–16.
10.1016/j.semcdb.2021.10.003.

46.     Ordahl, C.P., Berdougo, E., Venters, S.J., and Denetclaw, W.F. (2001). The
dermomyotome dorsomedial lip drives growth and morphogenesis of both the primary myotome
and dermomyotome epithelium. Development *128*, 1731–1744. 10.1242/dev.128.10.1731.

47.     Gros, J., Manceau, M., Thomé, V., and Marcelle, C. (2005). A common somitic origin for
embryonic muscle progenitors and satellite cells. Nature *435*, 954–958. 10.1038/nature03572.

48.     Burgess, R., Rawls, A., Brown, D., Bradley, A., and Olson, E.N. (1996). Requirement of
the paraxis gene for somite formation and musculoskeletal patterning. Nature *384*, 570–573.
10.1038/384570a0.

49.    Nakaya, Y., Kuroda, S., Katagiri, Y.T., Kaibuchi, K., and Takahashi, Y. (2004).
Mesenchymal-epithelial transition during somitic segmentation is regulated by differential roles
of Cdc42 and Rac1. Dev Cell *7*, 425–438. 10.1016/j.devcel.2004.08.003.

50.    Feener, C.A., Koenig, M., and Kunkel, L.M. (1989). Alternative splicing of human
dystrophin mRNA generates isoforms at the carboxy terminus. Nature *338*, 509–511.
10.1038/338509a0.

51.    Bies, R.D., Phelps, S.F., Cortez, M.D., Roberts, R., Caskey, C.T., and Chamberlain, J.S.
(1992). Human and murine dystrophin mRNA transcripts are differentially expressed during
skeletal muscle, heart, and brain development. Nucleic Acids Research *20*, 1725–1731.
10.1093/nar/20.7.1725.

52.    Muntoni, F., Torelli, S., and Ferlini, A. (2003). Dystrophin and mutations: one gene,
several proteins, multiple phenotypes. The Lancet Neurology *2*, 731–740. 10.1016/S1474-
4422(03)00585-4.

53.    T, F., T, Y., H, T., and K, I. (2020). Dystroglycan regulates proper expression,
submembranous localization and subsequent phosphorylation of Dp71 through physical
interaction. Human molecular genetics *29*. 10.1093/hmg/ddaa217.

54.    Romo-Yáñez, J., Rodríguez-Martínez, G., Aragón, J., Siqueiros-Márquez, L., Herrera-
Salazar, A., Velasco, I., and Montanez, C. (2020). Characterization of the expression of
dystrophins and dystrophin-associated proteins during embryonic neural stem/progenitor cell
differentiation. Neurosci Lett *736*, 135247. 10.1016/j.neulet.2020.135247.

55.    Belmaati Cherkaoui, M., Vacca, O., Izabelle, C., Boulay, A.-C., Boulogne, C., Gillet, C.,
Barnier, J.-V., Rendon, A., Cohen-Salmon, M., and Vaillend, C. (2021). Dp71 contribution to the
molecular scaffold anchoring aquaporine-4 channels in brain macroglial cells. Glia *69*, 954–970.
10.1002/glia.23941.

56.    Fujimoto, T., Stam, K., Yaoi, T., Nakano, K., Arai, T., Okamura, T., and Itoh, K. (2023).
Dystrophin Short Product, Dp71, Interacts with AQP4 and Kir4.1 Channels in the Mouse
Cerebellar Glial Cells in Contrast to Dp427 at Inhibitory Postsynapses in the Purkinje Neurons.
Mol Neurobiol. 10.1007/s12035-023-03296-w.

57.    Leyva-Leyva, M., Sandoval, A., Morales-Lázaro, S.L., Corzo-López, A., Felix, R., and
González-Ramírez, R. (2023). Identification of Dp140 and α1-syntrophin as novel molecular
interactors of the neuronal CaV2.1 channel. Pflugers Arch *475*, 595–606. 10.1007/s00424-023-
02803-1.

58.    Bourdon, A., François, V., Zhang, L., Lafoux, A., Fraysse, B., Toumaniantz, G., Larcher,
T., Girard, T., Ledevin, M., Lebreton, C., et al. (2022). Evaluation of the dystrophin carboxy-
terminal domain for micro-dystrophin gene therapy in cardiac and skeletal muscles in the
DMDmdx rat model. Gene Ther *29*, 520–535. 10.1038/s41434-022-00317-6.