Recovery from social isolation requires dopamine in males, but not the autism-related gene *nlg3* in either sex

Ryley T. Yost[1], Andrew M. Scott[3], Judy M. Kurbaj[1#], Brendan Walshe-Roussel[1], Reuven
Dukas[2] and Anne F. Simon[1]

[1]Department of Biology, Western University, London, ON, Canada
[2]Animal Behaviour Group, Department of Psychology, Neuroscience and Behaviour, McMaster
University, Hamilton, ON, Canada
[3]St. Joseph's Healthcare, Hamilton, ON, Canada
[#]Author was undergraduate at the time of data collection

* Correspondence:
Anne F. Simon
asimon28@uwo.ca

Keywords: Social Behaviour, Loneliness, Social Isolation, Social Space, Neuroligin,
Dopamine, Recovery.

Abstract

Social isolation causes profound changes in social behaviour in a variety of species including
humans, monkeys, mice, bees, and vinegar flies. However, the genetic and molecular
mechanisms modulating behavioural responses to both social isolation and social recovery
remain to be elucidated. In this study, we quantified the behavioural response of vinegar flies to
social isolation through the use of two distinct protocols, one involving flies' social space
preference and the other assessing flies' sociability, defined as their spontaneous tendencies to
form groups. We found that social isolation increased social space and reduced sociability. These
effects of social isolation, however, were reversible and could be reduced after 3 days of group
housing. Flies with a loss of function of *neuroligin3* (ortholog of autism-related *neuroligin*
genes) with known increased social space in a socially enriched environment, were still able to
recover from social isolation. Using a *UAS-TH*-RNAi driven in all neurons, we show that
dopamine is important for a response to social isolation and recovery in males but not in females.
Furthermore, only in males, dopamine levels are reduced after isolation and are not recovered
after group housing. Finally, in socially enriched flies with a loss of function of *neuroligin3*,
dopamine levels are reduced in males, but not in females. We propose a model to explain how
dopamine and *neuroligin3* are involved in the behavioural response to social isolation and its
recovery in a dynamic and sex-specific manner.

33
34
35
36

1   Introduction

The environment an individual has been exposed to can influence social behaviour by affecting
the interactions between individuals (Rubenstein and Hofmann, 2015). Social isolation, or the
absence of interactions with others, has profound effects on mental and physical health in
humans (Holt-Lunstad et al., 2010; Cacioppo et al., 2011; Meyer-Lindenberg and Tost, 2012).
During the COVID-19 times, forced social isolation had negative consequences on families and
individuals (Pellicano and Stears, 2020). Higher risk individuals such as children and people
with neuropsychiatric disorders, like Autism Spectrum Disorders, face more challenges to their
physical and mental health because of changes to their behavioural and environment support
(Bellomo et al., 2020; Summers et al., 2021). It is important to understand the interactions
between genetic predispositions and altered social environment, as well as the possibility to
recover from social isolation.
49
Social isolation causes changes to the social behaviour in several organisms, including a variety
of behavioural deficits in monkeys (Harlow et al., 1965;Mitchell and Clark, 1968;Harlow and
Suomi, 1971;Suomi et al., 2008), deficits in social interactions, aggression, fear and anxiety in
mice (Wongwitdecha and Marsden, 1996; Heidbreder et al., 2000; Fone and Porkess, 2008;
Zhang et al., 2012; An et al., 2017; Endo et al., 2018; Medendorp et al., 2018; Zhang et al., 2018)
and decreased social affiliation in bees (Hewlett *et al.*, 2018). In *Drosophila melanogaster*, social
isolation leads to physiological and behavioural changes, including decreased fiber number in the
mushroom bodies (Technau, 1984), changes to neural excitability (Ueda and Wu, 2009),
decreased lifespan (Leech *et al.*, 2017), chemical communication (Kent *et al.*, 2008), olfactory
memory (Chabaud *et al.*, 2009), courtship and courtship memory (Krupp *et al.*, 2008), sleep
patterns (Ganguly-Fitzgerald *et al.*, 2006), circadian rhythm (Krupp *et al.*, 2008), aggression
(Ueda and Kidokoro, 2002; Ueda and Wu, 2009;Ramin et al., 2014), sociability (Scott et al.,
2018) and social network structure (Bentzur *et al.*, 2020). Isolated male flies are much more
territorial and defend food patches and access to mates more than socially enriched flies
(Hoffman, 1990). Our work and others' have identified that isolated flies have increased social
space (Simon et al., 2012; Xie et al., 2018; Yost et al., 2020), a measure of inter-individual
distances (Simon et al., 2012; Burg et al., 2013).
67
Plastic changes in behaviour are important for an organism to adapt to a changing environment.
Changes in social behaviour because of isolation can be partially or fully recovered in monkeys
(Harlow et al., 1965; Harlow and Suomi, 1971), mice (Chen et al., 2016; An et al., 2017; Endo et
al., 2018) and bees (Hewlett *et al.*, 2018). Flies are no exception. In *Drosophila*, sleep and
aggression levels were recovered after group housing (Ganguly-Fitzgerald et al., 2006; Wang et
al., 2008). However, more research needs to be conducted on the recovery from isolation of other
Drosophila social behaviours.
75
Previous studies have identified sensory modalities, neurocircuitry and synaptic proteins to be
important for social space (Brenman-Suttner *et al.*, 2018). One identified synaptic protein is
neuroligin3, a post-synaptic cell adhesion molecule that regulates synaptic development and
function and is an ortholog to a human autism candidate gene (Knight et al., 2011; Sindi et al.,
2014; Yost et al., 2020). We have previously shown *nlg3* to be important for social space (Yost
et al., 2020) and sociability (the tendency to engage in non-aggressive interactions with
conspecifics; Scott *et al.*, 2018). We also showed *nlg3* is required for a typical response to social

isolation, but that the NLG3 protein levels are unchanged after social isolation (Yost *et al.*,
2020). However, the role of *nlg3* in social recovery has not been studied.
85
Another molecule identified as being important for social behaviour in the fly is dopamine (DA),
a monoamine neurotransmitter (Monastirioti, 1999; Waddell, 2010; Van Swinderen and
Andretic, 2011). We and others have previously shown that DA modulates social space
(Fernandez *et al.*, 2017) and is important for a response to social isolation. Specific
dopaminergic neural circuity is in part responsible for the behavioural modification after
isolation (Xie *et al.*, 2018) and DA levels in males are reduced after isolation (Ganguly-
Fitzgerald *et al.*, 2006), however the effect of DA on social recovery has not been identified. In
addition, evidence suggests that *nlg3* and DA are part of a similar neurocircuitry or involved in a
similar pathway responsible for behaviour regulation (Izquierdo et al., 2013; Uchigashima et al.,
2016; Bariselli et al., 2018), however their dual involvement in the regulation of social space
needs more attention.
97
In this study we examined the effect of social isolation on two measures of social behaviour,
social space (Simon *et al.*, 2012) and sociability (Scott *et al.*, 2018). We further investigated the
role of *nlg3* and DA in the recovery from social isolation. Finally, we propose a model for the
integrated regulation of *nlg3* and DA in the behavioural response to the social environment.

2   Methods

2.1   Fly stocks and husbandry

All fly lines were maintained in mixed sex groups in bottles on JazzMix media or our own food
made following the same recipe (brown sugar, corn meal, yeast, agar, benzoic acid, methyl
paraben and propionic acid; Fisher Scientific, Whitby, ON, Canada) at 25°C, 50% relative
humidity on a 12:12h light:dark cycle. All flies reared in bottles for experimental use were a
maximum of 14 days old to avoid variation in behaviour resulting from older parents (Brenman-
Suttner *et al.*, 2018). Fly lines were obtained from the following places: CS (Canton-S) was
obtained from the laboratory of Seymour Benzer in 1998, *nlg3*^{Def1} are from Dr. Brian Mozer
(Yost *et al.*, 2020), *w;; TH-GAL4* was provided by Dr. Serge Birman, RNAi against *tyrosine*
hydroxylase (*w;; UAS-THmiR-G*) was a gift from Dr. Mark Wu (Xie *et al.*, 2018). All lines
except w;; *UAS-ThmiR-G* were outcrossed five times to our control line, CS, to reduce variation
in behaviour caused by genetic background. Crosses used to generate *TH>ThmiR-G* and their
appropriate genetic controls can be found in **Supplemental Figure 1**.

2.2   Generation of isolation and recovery treatments

Mated flies used in all experiments were collected from bottles at 1 day old and remained
mixed sex in new bottles for one more day (2 days total) to allow mating (see confirmation of
mating status below) to avoid the effects of mating status on social space (Simon *et al.*, 2012).
Following, socially isolated flies were transferred to individual vials using cold anesthesia and
remained singly housed for 2, 4, or 7 days. Group housed age-matched control flies were kept in
mixed sex bottles for the same duration as the isolated flies, such that the group-housed control
flies for isolation were 4, 6 and 9 days old respectively. To test for a recovery after social
isolation, flies isolated for 7 days were transferred to mixed sex bottles for either 2 or 3 days.

Group housed flies used as a control for the recovery treatment were maintained in mixed sex
bottles until tested with the recovery flies, such that the group-housed control flies for isolation
then recovery were 11 and 13 days old respectively. To test the effect of group housing density
on social space, we sorted the group housed flies into vials mixed-sex containing 2, 6 or 16 flies
and a separate uncontrolled amount (random) in a bottle.

To test for the effects of virginity on social isolation, pupae were collected from bottles and
placed individually into vials and allowed to eclose over 24 hours. Isolated virgin flies remained
in individual vials while group-housed virgin flies were transferred into vials containing 15
same-sex flies to maintain virginity.

Females Mating Status at 2 and 4 days old: one-day-old female flies were collected and
maintained with males in bottles either 1 day (24 hours), or 3 days. Females were then sexed
under cold anesthesia and each female was placed alone in a fresh vial. Females were scored as
having been mated when 3^{rd} instar larvae were observed.
Males Mating Status at 2 and 4 days old: Single 1-day old male flies, sexed under cold
anesthesia, were placed with one 4-day old virgin female in vials (based on the females results,
more than 90% of females that age can mate). After either 1 day (24 hours), or 3 days, the males
were removed, and females remained in isolation until third instar larvae were counted. For both
sexes, vials with dead flies were not counted. We cannot ensure all flies have mated, even in the
group housing treatment, so all flies exposed to initial group housing are used in the behaviour
experiments, including the small percentage of non-mated flies.
145

146   **2.3   Social behaviour assays**

147   **Sociability tests** were performed at McMaster university. All other experiments took place at
148   Western University under similar settings.
149
Fly handling prior to behaviour: Twenty-four hours prior to all behaviour assays, 15-17 flies
that were group-housed (either lifelong or after a few days of recovery) were collected using cold
anesthesia and placed in vials. Single housed flies were kept isolated until right before testing.
The morning of the experiment, all flies were transferred to new vials and allowed to habituate to
the testing conditions of 25°C and 50% relative humidity for at least two hours. **The social space**
assay was conducted under uniform light in the same room between 1:00 PM and 5:00 PM (ZT
4-8) to decrease behavioural variation linked to diel periodicity.
157
The social space assay: Performed as previously described (Simon et al., 2012;McNeil et al.,
2015;Yost et al., 2020). In short: once flies have settled at their preferred inter-individual
distance, a photo was taken (after 20-50 minutes, depending on their genotype). Using the open
access software ImageJ (RRID:SCR_003070 - Schneider *et al.*, 2012), the number of flies within
4 body lengths (~1cm) was determined for each fly in the chamber. The number of flies within 4
body lengths was averaged using all flies in the chamber, representing one individual replicate.
This metric has been used in previous studies (Xie et al., 2018; Yost et al., 2020). The routines
for image analysis are publicly available (Yost *et al.*, 2020). All data sets are the combination of
1-3 replicates per day and 3-5 independent days of testing. Each independent day was separated
by at least a week to control for environmental factors beyond our control.

Sociability assay: The sociability chamber consisted of a circular arena (90 mm wide by 20 mm
high) divided into 8 compartments with a hole in the center to allow flies to enter any
compartment. We added to each compartment a patch of fresh food coated with a layer of
grapefruit-yeast suspension (3 g yeast in 100 mL grapefruit juice) to enhance attractiveness. We
modified the chamber and performance of the assay from Scott, Dworkin, and Dukas (Scott *et*
al., 2018). We transferred by mouth aspiration sixteen same-sex flies to the chamber through a
hole in the lid and allowed them to acclimate for one hour. Experimenters blind to treatment then
counted the number of flies in each chamber and calculated an aggregation index (sample
variance divided by the mean number of flies in each chamber). The variance could take values
between 0 and 32. For example, the least sociable option would have 8 chambers of 2 flies each,
with a variance of 0 and the corresponding aggregation index of 0. The most sociable situation
would have 7 chambers of 0, and 1 chamber with all 16 flies. This would have a variance of 32,
and therefore an aggregation index of 16 (32/2). We tested flies in sessions of four replicate
arenas per treatment over three consecutive weeks, for twelve arenas total per treatment (total N
= 96).

2.4   Dopamine quantification

Adults were separated by sex, and dopamine extracted from their heads using the following
procedure. Extraction occurred by flash freezing flies in liquid nitrogen followed by manual
decapitation and homogenization of heads in 5 mM of ammonium acetate in 90% acetonitrile
using microtissue grinders (Kimble Chase, USA). The supernatant was transferred and filtered
through a 0.65 µm filter (Millipore) at 4°C and samples were stored at -80 °C before LC/MS
detection.
190
LC/MS analysis was performed using an Agilent 1260 Infinity LC system coupled to an Agilent
6230 TOF system. A XBridge C-18 column Rapid Resolution HT was used (4.6 Å~150 mm, 3.5
µm, 600 Bar, Waters) at 25 °C and samples were eluted with a gradient of CH_3CN (Solvent B:
90% CH_3CN in H_2O, containing 0.1% formic acid) in H_2O (Solvent A: containing 0.1% formic
acid). The UV lamp was set at 282 nm, the injection volume was 10 µL. The flow rate was set to
0.4 mL/min and infused into an Agilent 6230 TOF-MS through a Dual Spray ESI source with a
gas temperature of 325 °C flowing at 8 L/min, and a nebulizer pressure of 35 psi. The fragmentor
voltage was set to 175 V with a capillary voltage of 3500 V and a skimmer voltage of 65 V. The
instrument was set in positive ESI mode, and quantification occurred using a standard curve of
known dopamine concentrations (**Supplemental Figure 2**). Total ion count was extracted using
Agilent MassHunter Qualitative Analysis software (version B.05.00).
202

2.5   Statistics

Data were stored in Excel files, and statistically analyzed using GraphPad Prism
(RRID:SCR_002798, version 7.0a for Mac, GraphPad Software, La Jolla California USA,
www.graphpad.com). We first assessed the normality and homoscedasticity of the data
distribution prior to applying the appropriate parametric tests (Simple *t*-test or ANOVA or
Welch- *t*-test/ANOVA when the standard deviations were different among groups). One-way and
Two-way ANOVAs were used, followed by Tukey's or Sidak's post hoc test (as appropriate) to

correct for multiple comparisons (see Table 1 for all the Two-Way ANOVA results). We used
the traditional alpha level of 0.05 for all statistical tests (Wasserstein and Lazar, 2016).
212

213 **3   Results**

214 **3.1   Isolation for seven days leads to increased social space and decreased sociability**

215 Seven days of social isolation leads to an increased social space (Simon *et al.* 2012). We wanted
216 to determine whether we could reduce the duration of social isolation and still influence social
217 space.
218

Because mating status influences social space, and virgin flies are further apart than mated flies
(Simon *et al.*, 2012), we needed to first assess how long flies should be group housed post-
emergence to avoid the confounding effect of virginity on social space. We found that 101 out of
112 (90.2%) of females were mated and 55 out of 72 (76.4%) of males were able to mate after
spending 4-5 days, mixed with females (**Supplemental Table 1**). In comparison, 94/103 (91.2%)
of females were mated 2 days after adult emergence spent with males, and 69/81 (85.2%) of
males of the same age were able to mate. The ages at which the flies were tested had no
significant effect on their mating capabilities ($\chi^2(1) = 0.022$, $p=0.88$ for females, and ($\chi^2(1) =$
1.92, $p=0.166$ for males). We thus decided to expose our flies to two days of group housing post-
emergence, before socially isolating them for 2, 4 or 7 days, as we now know that the same
percentage of flies have mated by that age, as when they are 4-5 days. We then assessed their
social space by recording the number of flies present within 4-body lengths.
231

Canton-S (CS, our control line) males had fewer flies within 4 body lengths after social isolation
compared to group housed flies at all three time points (2, 4 or 7 days) (**Figure 1A**; Two-way
ANOVA results in **Table 1**; Effect of isolation: $F_{1,46} = 22.01$, $p < 0.0001$). There was no effect of
2 days of social isolation in CS females. Only after 4 and 7 days of isolation did CS females have
fewer flies within 4 body lengths and the increase in social space was larger with increasing
number of days flies were isolated (**Figure 1B**; Two-way ANOVA results in **Table 1**; Effect of
isolation: $F_{1,47} = 6.653$, $p = 0.0131$; Effect of days of isolation: $F_{2,47} = 3.683$, $p = 0.0327$). The
effect of isolation after 7 days was similar to the effects found in earlier studies (Simon *et al.*,
2012). Moving forward we used seven days of isolation for the rest the experiments as seven
days had the largest increase in social space in males and females.
242

We next used another measure of social behaviour, sociability (Scott *et al.*, 2018), to examine the
effect of isolation. Both males and females had a lower aggregation index in isolated flies
compared to those group housed (**Figure 1C, Table 2**; Males – Welch's T-test: $t_{12.08} = 4.039$, $p =$
0.0016; Females – Welch's T-test: $t_{13.37} = 2.578$, $p = 0.0225$).

3.2   Males and females recover from isolation after three days of group-housing

As recovery from isolation has been noted in various organisms including *Drosophila*
melanogaster (see introduction), we tested whether social space and sociability could also be
recovered in flies that were group housed following isolation. We started with 2 days of group
housing following isolation and found that males were not different in flies within 4 body lengths

indicating they had recovered in social space, however females still had a lower number of flies
within 4 body lengths in the recovery compared to group housed flies (**Figure 1D, Table 2**;
Welch's T-test: $t_{15.13} = 2.626$, $p = 0.0189$). Because females had not fully recovered after two
days of group housing post isolation, we tested social space in flies that had three days of group
housing after seven days of isolation. Again, males were not different in flies within 4 body
lengths, however females were now also not different in flies within 4 body lengths in recovery
and group housed flies indicating that females also can recover from isolation but take longer
than males (one more day in this case – **Figure 1E, Table 3**). A period of three days of group
housing following seven days of isolation was then used for all other experiments investigating
recovery. Finally, we tested sociability in males and females after 3 days of recovery. Males and
females did not differ in aggregation index in recovery and group housed flies, indicating that
both sexes recovered (**Figure 1F, Table 3**).

3.3   Mating status does not alter social space in response to isolation

We have previously reported that virgin flies have an increased social space (Simon *et al*., 2012)
as do isolated flies (Yost *et al*., 2020 and Figure 1), however we have not tested virgin flies in
response to isolation, so we next tested the effect of isolation on social space in mated and virgin
flies. There were less flies within 4 body lengths after isolation in both mated and virgin males
(**Figure 2A**; Two-way ANOVA results in **Table 1**; Effect of isolation: $F_{1,52} = 34.69$, $p <$
0.0001). In addition, virgin males had less flies within 4 body lengths when group and single
housed compared to mated group and single housed (**Figure 2A**; Two-way ANOVA results in
Table 1; Effect of mating status: $F_{1,52} = 4.223$, $p = 0.0449$). In females we found that in mated
flies, single housed flies had less flies within 4 body lengths than group housed individuals
(**Figure 2B**; Two-way ANOVA results in **Table 1**; Effect of isolation: $F_{1,52} = 9.924$, $p =$
0.0027), however, group and single housed virgins had similar flies within 4 body lengths ($p =$
0.4308). Similar to males, virgin females had less flies within 4 body lengths when group and
single housed compared to mated group and single housed **Figure 2B**; Two-way ANOVA results
in **Table 1**; Effect of mating status: $F_{1,52} = 5.269$, $p = 0.0258$).
279
280

3.4   Group housed density does not alter social space

We next tested what happens when group housed flies are reared at different densities ranging
from 2 flies up to an uncontrolled or random number of flies. We found that single housed flies
have less flies within 4 body lengths compared to group housed flies at all densities in males
(**Figure 2C**, **Table 3**; One-way ANOVA with Holm-Sidak *post hoc*: $F_{4,40} = 30.87$, $p < 0.0001$)
and females (**Figure 2D**, **Table 3**; One-way ANOVA with Holm-Sidak *post hoc*: $F_{4,40} = 31.35$, p
< 0.0001).

3.5   The lack of *nlg3* does not prevent from recovering from social isolation

We have previously shown that *nlg3* is important for a typical social space response to isolation
(Yost *et al.*, 2020). We wondered whether *nlg3* is also important for the recovery from social
isolation. In males, both CS and *nlg3^{Def1}* had less flies within 4 body lengths in single housed
compared to group housed flies and *nlg3^{Def1}* had less flies within 4 body lengths than CS (**Figure
3A**, Two-way ANOVA results in **Table 1**; Effect of social experience: $F_{1,32} = 46.48$, $p < 0.0001$;

Effect of genotype: $F_{1,32} = 10.10, p = 0.0033$). When testing the recovery, CS males did not
differ in the number of flies within 4 body lengths, however $nlg3^{Def1}$ males had increased flies
within 4 body lengths in the recovery treatment compared to group housed $nlg3^{Def1}$ flies (**Figure**
3B, Two-way ANOVA results in **Table 1**; Effect of social experience: $F_{1,37} = 17.72, p = 0.0002$;
Effect of genotype: $F_{1,37} = 34.94, p > 0.0001$; Interaction of social experience and genotype:
$F_{1,37} = 13.62, p = 0.0007$).

In females, CS had less flies within 4 body lengths when single housed, however $nlg3^{Def1}$
females had no difference in the number of flies within 4 body lengths compared to group
housed flies (**Figure 3C**, Two-way ANOVA results in **Table 1**; Effect of social experience: $F_{1,32}$
$= 27.97, p > 0.0001$; Effect of genotype: $F_{1,32} = 13.71, p = 0.0008$; Interaction of social
experience and genotype: $F_{1,32} = 5.370, p = 0.0270$). In the recovery from isolation, CS flies
were not different in the number of flies within 4 body lengths compared to flies always group
housed however $nlg3^{Def1}$ flies had increased number of flies within 4 body lengths in the
recovery treatment compared to group housed flies (**Figure 3D**, Two-way ANOVA and Tukey
posthoc results in **Table 1**; Effect of social experience: $F_{1,34} = 0.3197, p = 0.5755$; Effect of
genotype: $F_{1,34} = 6.178, p = 0.0180$; Interaction of social experience and genotype: $F_{1,34} =$
$10.60, p = 0.0026$).

This recovery experiment is repeated in **Supplemental Figure 3** (those data were in fact
collected at the same time as the isolation treatments published in Yost *et al.*, 2020).

Combined, the data indicate that $nlg3$ is not required for recovery to occur.

**3.6   Dopamine is required for a response to isolation and decreases after isolation in a sex-
        specific manner**

Dopamine is important not only for social space in both sex (Fernandez *et al.*, 2017) but also for
the response to social isolation in males (Xie *et al.*, 2018). We tested to see if we could
recapitulate these results after isolation and determine if dopamine was important for the
recovery from isolation. We drove an RNAi against the gene encoding for Tyrosine Hydroxylase
(*TH*), the rate limiting enzyme for dopamine biosynthesis (*TH*-RNAi, specifically the *UAS-
THmiR-G*) using a *TH-gal4* driver.

In males *TH>THmiR-G*, there was no effect of social experience. We did observe a significant
reduction in the number of flies within 4 body lengths with isolation in flies with *TH*-Gal4/+ and
UAS-THmiR-G/+ (**Figure 4A**; Two-way ANOVA results in **Table 1**; Effect of isolation: $F_{1,42} =$
$19.13, p < 0.0001$; Genotype and isolation interaction: $F_{2,35} = 8.635, p = 0.0009$). When
investigating the recovery, we observed no difference in any genotype between group housed
and recovery treatments (**Figure 4B**; Two-way ANOVA results in **Table 1**).

In females, we observed a decrease in the number of flies within 4 body lengths for all three
genotypes after isolation and a decrease in *TH>THmiR-G* flies compared to their genetic controls
(**Figure 4C**; Two-way ANOVA results in **Table 1**; Effect of genotype: $F_{2,37} = 4.009, p = 0.0265$;
Effect of isolation: $F_{1,37} = 21.91, p < 0.0001$). Female *TH>THmiR-G* also displayed a decreased
number of flies within 4 body length; however, no genotype was different comparing group

housed to recovery treatments (**Figure 4D**; Two-way ANOVA results in **Table 1**; Effect of
genotype: $F_{2,35}$ = 5.124, p = 0.0112).
340
To establish the importance of dopamine in a response to isolation and recovery, we performed
LC/MS on the heads of males and females to quantify changes in dopamine levels after isolation
and recovery. After isolation, males had decreased dopamine levels while females' dopamine
levels remained similar to those of group housed flies. Further, male dopamine levels returned to
a similar level as group housed after a recovery period. Females' dopamine levels remained
unchanged (**Figure 4E**; Two-way ANOVA results in **Table 1**; Effect of social experience: $F_{3,27}$
= 3.043, p = 0.0459).
348
Taken together, these results indicate that dopamine is important for social space in response to
isolation and recovery in males but not females, and that dopamine levels are influenced by
previous social experience in a sex-specific manner.

3.7   *nlg3^{Def1}* flies have less dopamine, but its levels are unchanged after isolation.

To determine if an interaction was occurring between DA and *nlg3* we tested DA levels in group
housed CS and *nlg3^{Def1}*. In both males and females, DA was decreased in *nlg3^{Def1}* compared to
CS and females had lower DA levels compared to males (**Figure 5A**; Two-way ANOVA results
in **Table 1**; Effect of sex: $F_{1,7}$ = 6.107, p = 0.0428; Effect of genotype: $F_{1,7}$ = 17.21, p = 0.0043).
However, when we tested the effect of social experience, no difference in DA levels was
observed between *nlg3^{Def1}* group housed, isolated and recovery males and females (**Figure 5B**;
Two-way ANOVA results in **Table 1**).
360

4   Discussion

Here we report the effects of social isolation on social space and sociability. In addition, for the
first time, we report the effect of social recovery on the fly social behaviour. Social space was
increased, and sociability decreased after isolation but was recovered after three days of group
housing following isolation. We also show that mating status affected the response to isolation in
a sex-specific and that as soon as 2 flies or more are present, the size of the group prior to
assaying social space as no effect on social spacing. In addition, we show that an autism
candidate gene, *nlg3,* important for the response to social experience, is not required for recovery
from social isolation to occur. We also further confirm the role of dopamine in social space. But
we show for the first time the sex-specific role of DA in social spacing: DA is not only required
for a response to isolation but also for recovery from isolation in males, but not females.
Furthermore, DA levels also respond to the social environment by decreasing in males after
isolation and returning to normal levels when recovery occurs, but not in females. Finally, we
show that a *nlg3* loss of function mutant, *nlg3^{Def1}*, inhibits a change in DA in males and females,
with no further changes added with social isolation.
376
We first looked at how long flies needed to be isolated to effect social space. We saw a sex-
specific response to isolation where males only needed two days of isolation while females
needed closer to seven days. The females were less effected in the early days of isolation and
seemed to be more resilient to the effects of isolation than males (**Figure 1A & B**). Our results

for both males and females isolated for seven days phenocopy the results of Simon *et al.*, (2012).
With regard to recovery, males also responded faster, and recovered in two days versus three
days in females (**Figure 1D & E**). In summary, although males were affected sooner by
isolation, they were able to recover faster than females. Finally, we showed that sociability is
reduced in males and females but recovers after group housing (**Figure 1 C & F**), indicating that
isolation is affecting multiple social behaviours in the fly.

As we have previously shown that mating status and isolation are important for social space
(Simon et al., 2012; Yost et al., 2020), we investigated the role of mating status in social space in
response to isolation. In both males and females, social space was increased in virgin group
housed flies compared to mated group housed flies as previously reported (Simon et al., 2012;
Figure 2A). Both virgin and mated male flies had increased social space after isolation. We then
wanted to know if the number of flies present while group housed affected social space. We
tested single housed flies and flies reared group housed with 2, 6, 16, and an uncontrolled
random number of flies. The isolated flies had increased social space as expected (**Figure 2C &**
D). Group housed flies had lower social space than isolated flies, however, the social space was
similar amongst all the densities indicating the presence of even one other fly is enough to avoid
the negative consequences of isolation on social space.

Next, we looked at social space in the Drosophila homolog of an autism candidate gene, *nlg3*,
after isolation and recovery. We showed that males CS and *nlg3^{Def1}* flies had increased social
space after isolation. We have previously reported a diminished response to isolation in *nlg3^{Def1}*
flies (Yost *et al.*, 2020), however this time, we observed that male responded to isolation
similarly to their genetic controls (**Figure 3A**). However, isolated *nlg3^{Def1}* females did not,
similarly to what we had reported previously (**Figure 3C**; Yost *et al.*, 2020). When testing the
recovery after isolation, male and female CS flies had similar social space to flies always group
housed, indicating recovery had occurred (**Figure 3B, D**). Interestingly, in *nlg3^{Def1}* flies, both
male and females had decreased social space in the recovery treatment, which appears as an over
compensatory response. In an independent repeat of those data, both CS and *nlg3^{Def1}* flies
recovered similarly from isolation (**Supplemental Figure 3**). In both cases, lack of *nlg3* did not
prevent the recovery from social isolation.

Using an RNAi against tyrosine hydroxylase, we were able to show that DA is important for a
response to isolation in males since the isolated flies did not respond to isolation and even had a
slightly closer social space in isolated compared to group housed flies (**Figure 4A**). A similar
result was reported by Xie *et al.* (2018), where group housed flies acted as if isolated and
isolated flies acted as if group housed. We do not see as strong of effect as those authors
reported, but we also used a pan-TH driver, while Xie *et al.* (2018) focused on a few TH cells
only. Our work and that of Xie *et al.* (2018) demonstrate the importance of DA in males in the
response to the social environment. However, we show for the first time that females do not
require DA for a response to isolation and recovery to occur (**Figure 4C & D**). Another
neurotransmitter or neuromodulator could be at play in females, in response to social experience.
It has been previously shown that both males and females respond similarly to reduced DA
levels, when group housed, with an increase in social space (Fernandez *et al.*, 2017). But DA
might not be the neurotransmitter involved in responding to social experience in females.

When we isolated our control line, CS, we confirmed that there is a decrease in DA in males but
not females and that the DA levels in males returned to the level of group housed flies after
recovery (**Figure 4E**). Ganguly-Fitzgerald *et al.* (2006) also saw a decrease in DA in males after
isolation. Taken together these results indicate DA in males is extremely important for social
space in response to isolation, but not in females. Finally, we report that DA levels in *nlg3^{Defl}*
remain unchanged after isolation, providing further evidence that *nlg3* and DA are both
contributing to the modulation of social space after isolation (**Figure 5A & B**). In both CS and
nlg3^{Defl}, decreases in DA were observed between 9- and 12-day old flies regardless of social
experience, consistent with age related decreases in DA previously reported (Neckameyer et al.,
2000).
437
As for recovering from social isolation, what does this mean for dopamine and an autism related
gene? Dopamine in males is required for a response to the environment, however the flies need
nlg3 for DA levels to change in response to the social environment and subsequently modulate
behaviour. Furthermore, DA levels decrease in response to social isolation in males, but the
NLG3 protein itself is not responding to social experience (Yost *et al.*, 2020). We predict that
nlg3 and DA are part of a pathway responsible for the modulation of social behaviour after
isolation in males. *Nlg3* would be required downstream of DA, for proper DA signaling in
response to the social environment, with some feedback regulation, since DA did not change
after isolation when *nlg3* was absent. Females do require *nlg3* for a typical response to isolation
(Yost *et al.*, 2020), but other neurotransmitters or neuromodulators, responsive to social
experience in females, may be interacting with *nlg3* (**Figure 6**). A role of Nlgs in DA signaling,
and their common influence on behaviour, including social behaviour, has been observed in mice
(Rothwell *et al.*, 2014, Bariselli *et al.*, 2018), the northern swordtail fish (Wong and Cummings,
2014) and the worm *C. elegans* (Izquierdo *et al.*, 2013). What remains unknown is whether Nlg3
is directly involved in the post-synaptic recruitment of one of the dopamine receptors in flies, as
has been suggested, based on work in mice, by Uchigashima *et al*. (2016). Finally, whether
Cyp20 is also involved in the response to social experience, in both males and females, to
modulate changes in social space and sociability still needs to be investigated (Wang *et al.,*
2008).
457
In conclusion, for the first time, we have demonstrated that social space and sociability are
recoverable after isolation in a sex-specific manner. We have also shown that DA is required for
the recovery from social isolation in a sex-specific manner and that *nlg3* is required for DA
levels to respond to the social environment. If those gene by environment by sex interactions are
evolutionary conserved, it might help better understanding the molecular basis of the social
difficulties encountered by humans with neurodevelopmental disorders in response to changing
social environments.
465

466   **5   Conflict of Interest**

467   The authors declare that the research was conducted in the absence of any commercial or
468   financial relationships that could be construed as a potential conflict of interest.
469
470

6   Author Contributions

Data acquisition and analysis were carried out by RTY, AMS, JMK, and BW-R; conception:
RTY and AFS. The first draft was written by RTY and the final draft edited by RTY, AMS,
JMK, BW-R, RD and AFS.

7   Funding

This project was funded by an internal and provincial grant to RTY, NSERC Fellowship to RTY
and AMS, NSERC Discovery grants 05420-2019 to RD, and 05054-2022 to AFS.

8   Acknowledgments

We would like to thank Dr. Mark Bernards, Dr. Repon Saha, and Karina Kaberi for their
knowledge and expertise obtaining the dopamine level quantification.

9   References

An, D., Chen, W., Yu, D.Q., Wang, S.W., Yu, W.Z., Xu, H., Wang, D.M., Zhao, D., Sun, Y.P.,
Wu, J.C., Tang, Y.Y., and Yin, S.M. (2017). Effects of social isolation, re-socialization
and age on cognitive and aggressive behaviors of Kunming mice and BALB/c mice.
Anim Sci J 88**,** 798-806.

Bariselli, S., Hornberg, H., Prevost-Solie, C., Musardo, S., Hatstatt-Burkle, L., Scheiffele, P., and
Bellone, C. (2018). Role of VTA dopamine neurons and neuroligin 3 in sociability traits
related to nonfamiliar conspecific interaction. *Nat Commun* 9**,** 3173.

Bellomo, T.R., Prasad, S., Munzer, T., and Laventhal, N. (2020). The impact of the COVID-19
pandemic on children with autism spectrum disorders. *J Pediatr Rehabil Med* 13**,** 349-
354.

Bentzur, A., Ben-Shaanan, S., Benichou, J.I.C., Costi, E., Levi, M., Ilany, A., and Shohat-Ophir,
G. (2020). Early life experience shapes male behavior and social networks in Drosophila.
Curr Biol.

Brenman-Suttner, D.B., Long, S.Q., Kamesan, V., De Belle, J.N., Yost, R.T., Kanippayoor, R.L.,
and Simon, A.F. (2018). Progeny of old parents have increased social space in
Drosophila melanogaster. *Sci Rep* 8**,** 3673.

Burg, E.D., Langan, S.T., and Nash, H.A. (2013). Drosophila social clustering is disrupted by
anesthetics and in *narrow abdomen* ion channel mutants. *Genes Brain Behav* 12**,** 338-
347.

Cacioppo, J.T., Hawkley, L.C., Norman, G.J., and Berntson, G.G. (2011). Social isolation. *Ann N*
Y Acad Sci 1231**,** 17-22.

Chabaud, M.A., Isabel, G., Kaiser, L., and Preat, T. (2009). Social facilitation of long-lasting
memory retrieval in Drosophila. *Curr Biol* 19**,** 1654-1659.

Chen, W., An, D., Xu, H., Cheng, X., Wang, S., Yu, W., Yu, D., Zhao, D., Sun, Y., Deng, W.,
Tang, Y., and Yin, S. (2016). Effects of social isolation and re-socialization on cognition
and ADAR1 (p110) expression in mice. *PeerJ* 4**,** e2306.

Endo, N., Ujita, W., Fujiwara, M., Miyauchi, H., Mishima, H., Makino, Y., Hashimoto, L.,
Oyama, H., Makinodan, M., Nishi, M., Tohyama, C., and Kakeyama, M. (2018).
Multiple animal positioning system shows that socially-reared mice influence the social
proximity of isolation-reared cagemates. *Commun Biol* 1**,** 225.

Fernandez, R.W., Akinleye, A.A., Nurilov, M., Feliciano, O., Lollar, M., Aijuri, R.R., O'donnell,
J.M., and Simon, A.F. (2017). Modulation of social space by dopamine in *Drosophila*
melanogaster, but no effect on the avoidance of the *Drosophila* stress odorant. *Biol Lett*
13**,** 20170369.

Fone, K.C., and Porkess, M.V. (2008). Behavioural and neurochemical effects of post-weaning
social isolation in rodents-relevance to developmental neuropsychiatric disorders.
Neurosci Biobehav Rev 32**,** 1087-1102.

Ganguly-Fitzgerald, I., Donlea, J., and Shaw, P.J. (2006). Waking experience affects sleep need
in *Drosophila*. *Science* 313**,** 1775-1781.

Harlow, H.F., Dodsworth, R.O., and Harlow, M.K. (1965). Total social isolation in monkeys.
PNAS 54**,** 90-97.

Harlow, H.F., and Suomi, S.J. (1971). Social recovery by isolation-reared monkeys. *PNAS* 68**,**
1534-1538.

Heidbreder, C.A., Weiss, I.C., Domeney, A.M., Pryce, C., Homberg, J., Hedou, G., Feldon, J.,
Moran, M.C., and Nelson, P. (2000). Behavioral, neurochemical and endocriminological
characterization of the early social isolation syndrome. *Neurosci* 100**,** 749-768.

Hewlett, S.E., Wareham, D.M., and Barron, A.B. (2018). Honey bee (*Apis mellifera*) sociability
and nestmate affiliation are dependent on the social environment experienced post-
eclosion. *J Exp Biol* 221.

Hoffman, A.A. (1990). The influence of age and experience with conspecifics on territorial
behaviour in *Drosophila melanogaster*. *Journal of Insect Behavior* 3**,** 1-12.

Holt-Lunstad, J., Smith, T.B., and Layton, J.B. (2010). Social relationships and mortality risk: a
meta-analytic review. *PLoS Med* 7**,** e1000316.

Izquierdo, P.G., Calahorro, F., and Ruiz-Rubio, M. (2013). Neuroligin modulates the locomotory
dopaminergic and serotonergic neuronal pathways of *C. elegans*. *Neurogenetics* 14**,** 233-
242.

Kent, C., Azanchi, R., Smith, B., Formosa, A., and Levine, J.D. (2008). Social context influences
chemical communication in *D. melanogaster* males. *Curr Biol* 18**,** 1384-1389.

Knight, D., Xie, W., and Boulianne, G.L. (2011). Neurexins and neuroligins: recent insights
from invertebrates. *Mol Neurobiol* 44**,** 426-440.

Krupp, J.J., Kent, C., Billeter, J.C., Azanchi, R., So, A.K., Schonfeld, J.A., Smith, B.P., Lucas,
C., and Levine, J.D. (2008). Social experience modifies pheromone expression and
mating behavior in male *Drosophila melanogaster*. *Curr Biol* 18**,** 1373-1383.

Leech, T., Sait, S.M., and Bretman, A. (2017). Sex-specific effects of social isolation on ageing
in *Drosophila melanogaster*. *J Insect Physiol* 102**,** 12-17.

Mcneil, A.R., Jolley, S.N., Akinleye, A.A., Nurilov, M., Rouzyi, Z., Milunovich, A.J.,
Chambers, M.C., and Simon, A.F. (2015). Conditions affecting social space in
Drosophila melanogaster. *J Vis Exp*, e53242.

Medendorp, W.E., Petersen, E.D., Pal, A., Wagner, L.M., Myers, A.R., Hochgeschwender, U.,
and Jenrow, K.A. (2018). Altered behavior in mice socially isolated during adolescence
corresponds with immature dendritic spine morphology and impaired plasticity in the
prefrontal cortex. *Front Behav Neurosci* 12**,** 87.

Meyer-Lindenberg, A., and Tost, H. (2012). Neural mechanisms of social risk for psychiatric
disorders. *Nat Neurosci* 15**,** 663-668.

Mitchell, G.D., and Clark, D.L. (1968). Long-term effect of social isolation in nonsocially
adapted rhesus monkeys. *Journal of Genetic Psychology* 113**,** 117-128.

Monastirioti, M. (1999). Biogenic amine systems in the fruit fly *Drosophila melanogaster*.
Microscopy Research and Technique 45**,** 106-121.

Neckameyer, W.S., Woodrome, S., Holt, B., and Mayer, A. (2000). Dopamine and senescence in
Drosophila melanogaster. *Neurobiology of Aging* 21**,** 145-152.

Pellicano, E., and Stears, M. (2020). The hidden inequalities of COVID-19. *Autism* 24**,** 1309-
1310.

Ramin, M., Domocos, C., Slawaska-Eng, D., and Rao, Y. (2014). Aggression and social
experience: genetic analysis of visual circuit activity in the control of aggressiveness in
Drosophila. *Mol Brain* 7**,** 55.

Rubenstein, D.R., and Hofmann, H.A. (2015). Proximate pathways underlying social behavior.
Current Opinion in Behavioral Sciences 6**,** 154-159.

Schneider, C.A., Rasband, W.S., and Eliceiri, K.W. (2012). NIH Image to ImageJ: 25 years of
image analysis. *Nat Methods* 9**,** 671-675.

Scott, A.M., Dworkin, I., and Dukas, R. (2018). Sociability in fruit flies: genetic variation,
heritability and plasticity. *Behavior Genetics* 48**,** 247-258.

Simon, A.F., Chou, M.T., Salazar, E.D., Nicholson, T., Saini, N., Metchev, S., and Krantz, D.E.
(2012). A simple assay to study social behavior in Drosophila: measurement of social
space within a group. *Genes Brain Behav* 11**,** 243-252.

Sindi, I.A., Tannenberg, R.K., and Dodd, P.R. (2014). Role for the neurexin-neuroligin complex
in Alzheimer's disease. *Neurobiol Aging* 35**,** 746-756.

Summers, J., Baribeau, D., Mockford, M., Goldhopf, L., Ambrozewicz, P., Szatmari, P., and
Vorstman, J. (2021). Supporting children with neurodevelopmental disorders during the
COVID-19 pandemic. *J Am Acad Child Adolesc Psychiatry* 60**,** 2-6.

Suomi, S.J., Van Der Horst, F.C., and Van Der Veer, R. (2008). Rigorous experiments on
monkey love: an account of Harry F. Harlow's role in the history of attachment theory.
Integr Psychol Behav Sci 42**,** 354-369.

Technau, G.M. (1984). Fibre number in the mushroom bodies of adult *Drosophila melanogaster*
depends on age, sex and experience. *Journal of Neurogenetics* 21**,** 183-196.

Uchigashima, M., Ohtsuka, T., Kobayashi, K., and Watanabe, M. (2016). Dopamine synapse is a
neuroligin-2-mediated contact between dopaminergic presynaptic and GABAergic
postsynaptic structures. *Proc Natl Acad Sci U S A* 113**,** 4206-4211.

Ueda, A., and Kidokoro, Y. (2002). Aggressive behaviours of female *Drosophila melanogaster*
are influenced by their social experience and food resources. *Physiol Entomol* 27**,** 21-28.

Ueda, A., and Wu, C.F. (2009). Effects of social isolation on neuromuscular excitability and
aggressive behaviors in *Drosophila*: altered responses by *Hk* and *gsts1*, two mutations
implicated in redox regulation. *J Neurogenet* 23**,** 378-394.

Van Swinderen, B., and Andretic, R. (2011). Dopamine in Drosophila: setting arousal thresholds
in a miniature brain. *Proc Biol Sci* 278**,** 906-913.

Waddell, S. (2010). Dopamine reveals neural circuit mechanisms of fly memory. *Trends*
Neurosci 33**,** 457-464.

Wang, L., Dankert, H., Perona, P., and Anderson, D.J. (2008). A common genetic target for
environmental and heritable influences on aggressiveness in *Drosophila*. *Proc Natl Acad*
Sci U S A 105**,** 5657-5663.

Wongwitdecha, N., and Marsden, C.A. (1996). Social isolation increases aggressive behaviour
and alters the effects of diazepam in the rat social interaction test. *Behav Brain Res* 75**,**
27-32.

Xie, T., Ho, M.C.W., Liu, Q., Horiuchi, W., Lin, C.C., Task, D., Luan, H., White, B.H., Potter,
C.J., and Wu, M.N. (2018). A genetic toolkit for dissecting dopamine circuit function in
Drosophila. *Cell Rep* 23**,** 652-665.

Yost, R.T., Robinson, J.W., Baxter, C.M., Scott, A.M., Brown, L.P., Aletta, M.S., Hakimjavadi,
R., Lone, A., Cumming, R.C., Dukas, R., Mozer, B., and Simon, A.F. (2020). Abnormal
social interactions in a Drosophila mutant of an autism candidate gene: *neuroligin 3*. *Int J*
Mol Sci 21.

Zhang, B., Sato, K., and Yamamoto, D. (2018). Ecdysone signaling regulates specification of
neurons with a male-specific neurite in Drosophila. *Biol Open* 7**,** bio029744.

Zhang, Y., Zu, X., Luo, W., Yang, H., Luo, G., Zhang, M., and Tang, S. (2012). Social isolation
produces anxiety-like behaviors and changes PSD-95 levels in the forebrain. *Neurosci*
Lett 514**,** 27-30.

10   Data Availability Statement

618 The datasets analyzed for this study can be found in the Dryad repository
619 (https://doi.org/10.5061/dryad.59zw3r2fs)
620
621
622
623

Experiment	Figure Number	Effect	*Df*	*F*	*p*	*Sidak post hoc*		
Social space						**Comparison Group**	**Compared with**	*p*
Days of Isolation-Males	1A	Days	(2,46)	2.412	0.1009	Group housed vs. single housed	2 days	0.0529
		Isolation	**(1,46)**	**22.01**	**<0.0001**		**4 days**	**0.0424**
		Days * Isolation	(2,46)	0.1534	0.8582		**7 days**	**0.0093**
Days of Isolation-Females	1B	**Days**	**(2,47)**	**3.683**	**0.0327**	Group housed vs. single housed	2 days	0.9992
		Isolation	**(1,47)**	**6.653**	**0.0131**		4 days	0.1404
		Days * Isolation	(2,47)	1.453	0.2441		7 days	0.0746
Mating Status-Males	2A	**Isolation**	**(1,52)**	**35.69**	**0.0327**	Group housed vs. single housed	**Mated**	**<0.0001**
		Mating Status	**(1,52)**	**4.223**	**0.0131**		**Virgin**	**0.0007**
		Isolation * Mating Status	(1,52)	0.2036	0.2441			
Mating Status-Females	2B	**Isolation**	**(1,52)**	**9.924**	**0.0027**	Group housed vs. single housed	**Mated**	**0.0033**
		Mating Status	**(1,52)**	**5.269**	**0.0258**		Virgin	0.4308
		Isolation * Mating Status	(1,52)	2.131	0.1504			
Cs & *nlg3^{Def1}* Isolated - Males	3A	**Genotype**	**(1,32)**	**10.10**	**0.0033**	Group housed vs. single housed	**Cs**	**=0.0001**
		Isolation	**(1,32)**	**46.48**	**<0.0001**		*nlg3^{Def1}*	**<0.0001**
		Genotype * Isolation	(1,32)	0.0086	0.9267			
Cs & *nlg3^{Def1}* Recovery - Males	3B	**Genotype**	**(1,37)**	**39.94**	**0.0002**	Group housed vs. recovery	Cs	0.9213
		Recovery	**(1,37)**	**17.72**	**<0.0001**		*nlg3^{Def1}*	**<0.0001**
		Genotype * Recovery	**(1,37)**	**13.62**	**0.0007**			
Cs & *nlg3^{Def1}* Isolated - Females	3C	**Genotype**	**(1,32)**	**13.71**	**0.0008**	Group housed vs. single housed	**Cs**	**<0.0001**
		Isolation	**(1,32)**	**27.97**	**<0.0001**		*nlg3^{Def1}*	0.0769
		Genotype * Isolation	**(1,32)**	**5.370**	**0.0270**			
Cs & *nlg3^{Def1}* Recovery - Females	3D	**Genotype**	**(1,34)**	**6.178**	**0.0180**	Group housed vs. recovery	Cs	0.1326
		Recovery	(1,34)	0.3197	0.5755		*nlg3^{Def1}*	**0.0197**
		Genotype * Recovery	**(1,34)**	**10.60**	**0.0026**			
TH miR Isolation-Males	4A	Genotype	(2,35)	1.633	0.2099	Group housed vs. single housed	*th-gal4/+*	0.9851
		Isolation	**(1,35)**	**4.631**	**0.0384**		***UAS-th-miR-G/+***	**0.0003**
		Genotype * Isolation	**(2,35)**	**8.635**	**0.0009**		*th > th-miR-G*	0.5283
TH miR Recovery-Males	4B	**Genotype**	**(2,37)**	**9.018**	**0.0006**	Group housed vs. recovery	*th-gal4/+*	>0.9999
		Recovery	(1,37)	4.631	0.7555		*UAS-th-miR-G/+*	0.9990
		Genotype * Recovery	(2,37)	8.635	0.8263		*th > th-miR-G*	0.8707
TH miR Isolation-Females	4C	**Genotype**	**(2,37)**	**4.009**	**0.0265**	Group housed vs. single housed	***th-gal4/+***	**0.0188**
		Isolation	**(1,37)**	**21.91**	**<0.0001**		***UAS-th-miR-G/+***	**0.0025**
		Genotype * Isolation	(2,37)	0.8892	0.4196		*th > th-miR-G*	0.2928
TH miR Recovery-Females	4D	**Genotype**	**(2,35)**	**5.124**	**0.0112**	Group housed vs. recovery	*th-gal4/+*	0.8403
		Recovery	(1,35)	2.562	0.1184		*UAS-th-miR-G/+*	0.0667
		Genotype * Recovery	(2,35)	2.036	0.1457		*th > th-miR-G*	0.9661
Dopamine levels						**Comparison Group**	**Compared with**	*p*
Cs	4E	Sex	(1,27)	1.665	0.2078	Males Grouped housed (GH) – 9 days old (9do)	***Males SH 9do***	**0.0262**
		Social Experience	**(3,27)**	**3.043**	**0.0459**		*Males GH 12do*	0.2372
		Sex * Social experience	(3,27)	1.130	0.3545		*Males Rec 12 do*	0.3747
							Females SH 9do	0.9613
							Females GH 12do	0.9913
							Females Rec 12 do	0.4623
Cs and *nlg3^{Def1}*	5A	**Sex**	**(1,7)**	**6.107**	**0.0428**	Cs males	*Cs females*	0.1531
		Genotype	**(1,7)**	**17.21**	**0.0043**		***Males nlg3^{Def}***	**0.0347**
		Sex * Genotype	(1,7)	2.797	0.1384		***Females nlg3^{Def}***	**0.0179**
						Cs females	*Males nlg3^{Def}*	0.8203
							Females nlg3^{Def}	0.4884
						nlg3^{Def} males	*Females nlg3^{Def}*	0.9935
nlg3^{Def1}	5B	Sex	(1,16)	0.007	0.999	Males Grouped housed (GH) – 9 days old (9do)	*Males SH 9do*	0.9922
		Social Experience	(3,16)	1.354	0.2922		*Males GH 12do*	0.7168
		Sex * Social experience	(3,16)	0.921	0.3513		*Males Rec 12 do*	0.9969
							Females SH 9do	0.9798
							Females GH 12do	0.6796
							Females Rec 12 do	0.9989

625   **Table 1. ANOVA table for all Two-way ANOVAs performed.** Statistically significant results are bolded.

Experiment	Figure Number	Comparison	Df	t	p
Sociability Isolation – Males	1C	Group housed vs. single housed males	**12.08**	**4.039**	**0.0016**
Sociability Isolation – Females	1C	Group housed vs. single housed females	**13.37**	**2.578**	**0.025**
Social Space Recovery 2 day – Males	1D	Group housed vs recovery males	13.11	0.3097	0.7617
Social Space Recovery 2 day – Females	1D	Group housed vs. recovery females	**15.13**	**2.2626**	**0.0189**

Table 2. All t-tests performed. Statistically significant results are bolded

Experiment	Figure Number	Df	F	p	Sidak post hoc		
					Comparison Group	Compared with	p
Social Space Recovery – Males & Females	1E	(3,28)	0.9434	0.4329	Group housed Females	Recovery Females	0.3881
Sociability Recovery – Males & Females	1F	(3,44)	2.102	0.1136	Group housed Males	Recovery Males	0.8841
					Group housed Females	Recovery Females	0.2599
Group Housed Density – Males	2C	**(4,40)**	**30.87**	**<0.0001**	Single Housed	2	**<0.0001**
						6	**<0.0001**
						16	**<0.0001**
						Random	**<0.0001**
Group Housed Density – Females	2C	**(4,40)**	**31.35**	**<0.0001**	Single Housed	2	**<0.0001**
						6	**<0.0001**
						16	**<0.0001**
						Random	**<0.0001**

Table 3. ANOVA table for all One-way ANOVAs performed. Statistically significant results are bolded.

<u>**Figure captions**</u>

Figure 1. Isolation for seven days leads to increased social space and decreased sociability,

but flies recover after 3 days of group housing. A-B: Social space presented in terms of mean

number of flies within 4 body lengths +/- s.e.m. in males (A) and females (B) after 2, 4 or 7 days

of isolation. **A.** Male flies had fewer flies within 4 body lengths after 2, 4 and 7 days of isolation

(Two-way ANOVA and Sidak post hoc test - $F_{1,46} = 22.01$, $p < 0.0001$). **B.** Females had fewer

flies within 4 body lengths after isolation and that number decreased further with increasing days

of isolation (Two-way ANOVA - Effect of Isolation: $F_{1,47} = 6.653$, $p = 0.0131$; Effect of Days of

Isolation: $F_{2,47} = 3.687$, $p = 0.0367$). **C.** Aggregation index in males and females after 7 days of

isolation. Both males (Welch's T-test: $t_{12.08} = 4.039$, $p = 0.0016$) and females (Welch's T-test:

$t_{13.37} = 2.578$, $p = 0.0225$) had a lower aggregation index after 7 days of isolation. **D-E:** Males

and females after 2 (D) and 3 (E) days of group housing (recovery) following 7 days of isolation.

D. Males showed no difference in the number of flies within 4 body lengths after 2 days of group

housing compared to males continuously group housed. However, females had fewer flies within

4 body lengths compared to females group housed continuously (Welch's T-test: $t_{15.13} = 2.626$, p

$= 0.0189$). **E.** Both males and females were not different in the number of flies within 4 body

lengths comparing group housed and recovery flies (One-way ANOVA: $F_{3,28} = 0.9434$, $p =$

0.4329). **F.** Sociability presented as the mean aggregation index +/- s.e.m. of males and females

group housed and recovery. Males and females did not differ in aggregation index between

group housed and recovery flies (One-way ANOVA: $F_{3,44} = 2.102$, $p = 0.1136$). N = 7-12 for all

treatments. *$p < 0.05$. ****$p < 0.0001$. Grey shade: Group Housed (GH); blue shades: single

housed (SH) and recovery from social isolation (REC) males; red shades: SH and REC females.

Figure 2. Mating status does not change social space in response to isolation and group

housed with a single fly is enough to avoid a larger social space. A-B: Number of flies within

4 body lengths in males (A) and females (B) group and single housed while mated or virgin. **A.**

Male flies had less flies within 4 body lengths after isolation (Two-way ANOVA – Effect of

isolation: $F_{1,52} = 34.69$, $p < 0.0001$) regardless of mating status. Males had less flies within 4

body lengths when virgin compared to mated flies (Two-way ANOVA – Effect of mating status:

$F_{1,52} = 4.223$, $p = 0.0449$). **B.** Female flies had less flies within 4 body lengths after isolation

(Two-way ANOVA – Effect of isolation: $F_{1,52} = 9.924$, $p = 0.0027$) regardless of mating status.

Females had less flies within 4 body lengths when virgin compared to mated flies (Two-way

ANOVA – Effect of mating status: $F_{1,52} = 5.269$, $p = 0.0258$). **C-D:** Number of flies within 4

body lengths in males (C) and females (D) reared single housed or group housed with 2, 6, 16 or

a random number of flies. Single housed flies had less flies within 4 body lengths than group

housed flies that were reared with any number of individuals in males (C; One-way ANOVA

with Holm-Sidak post hoc: $F_{4,40} = 30.87$, $p < 0.0001$) and females (D; One-way ANOVA with

Holm-Sidak post hoc: $F_{4,40} = 31.35$, $p < 0.0001$). N = 9-13 for all treatments. GH: Group housed,

SH: Single housed. Bars: mean +/- s.e.m.

Figure 3: $nlg3^{Def1}$ flies have decreased social space after a recovery from isolation. **A-B:**

Number of flies within 4 body lengths in males group and single housed (A) and group housed

with a recovery (B). **A.** Single housed male flies had decreased flies within 4 body lengths

compared to group housed flies in both CS and $nlg3^{Def1}$ (Two-way ANOVA - Effect of social

experience: $F_{1,32} = 46.48$, $p < 0.0001$) and $nlg3^{Def1}$ flies had less flies within 4 body lengths in

both group housed and single housed (Two-way ANOVA - Effect of genotype: $F_{1,32}$ = 10.10, p =
0.0033). **B.** The number of flies within 4 body lengths for CS males was not different in group
housed vs recovery flies, however $nlg3^{Def1}$ flies had an increased number of flies within 4 body
lengths in recovery compared to group housed flies (Two-way ANOVA - Effect of social
experience: $F_{1,37}$ = 17.72, p = 0.0002; Effect of genotype: $F_{1,37}$ = 34.94, p < 0.0001; Interaction
of social experience and genotype: $F_{1,37}$ = 13.62, p = 0.0007). **C-D:** Number of flies within 4
body lengths in group and single housed females (C) and group housed with a recovery females
(D). **C.** CS females had less flies within 4 body lengths in single housed compared to group
housed flies, however $nlg3^{Def1}$ flies were not different between group and single housed flies
(Two-way ANOVA - Effect of social experience: $F_{1,32}$ = 27.97, p < 0.0001; Effect of genotype:
$F_{1,32}$ = 13.71, p = 0.0008; Interaction of social experience and genotype: $F_{1,32}$ = 5.370, p =
0.0270). **D.** The number of flies within 4 body lengths for CS females was not different in group
housed vs recovery flies, however $nlg3^{Def1}$ females had an increased number of flies within 4
body lengths in recovery compared to group housed flies (Two-way ANOVA - Effect of social
experience: $F_{1,34}$ = 0.3197, p = 0.5755; Effect of genotype: $F_{1,34}$ = 6.178, p = 0.0180; Interaction
of social experience and genotype: $F_{1,34}$ = 10.60, p = 0.0026). N = 7-12 for all treatments. GH:
Group housed. GH: Group housed, SH: Single housed, REC: Recovery.
695
696
Figure 4. Dopamine is important for a response to social isolation and decreases after
isolation in males but not females. Social space measured in terms of number of flies within 4
body lengths**.** Bars: mean +/- s.e.m. **A-B:** Social space in isolated (A) and recovered (B) male
flies. **A.** The social space in male *TH > THmiR-G* was not different in isolated vs group housed
flies, however *TH-GAL4/+* and *UAS-THmiR-G/+* males had lower flies within 4 body lengths
when isolation (Two-way ANOVA - Effect of Isolation: $F_{1,43}$ = 19.13, p < 0.0001; Interaction
between isolation and genotype: $F_{2,42}$ = 8.254, p = 0.0009). **B.** Male recovery flies in all
genotypes were not different compared to group housed males. **C-D:** Social space in isolated (C)
and recovered (D) female flies. **C.** All three genotypes had a decrease in the number of flies
within 4 body lengths after isolation and *TH > THmiR-G* flies had the lowest number of flies
within 4 body lengths compared to their genetic controls (Two-way ANOVA – Effect of
Isolation: $F_{1,37}$ = 21.91, p < 0.0001; Effect of Genotype: $F_{2,37}$ = 4.009, p = 0.0265). **D.** All three
genotypes were similar when comparing group housed vs. recovery treatments, however *TH >*
THmiR-G females had a less flies within 4 body lengths compared to the controls (Two-way
ANOVA – Effect of Genotype: $F_{2,35}$ = 5.124, p = 0.0112). **E.** Dopamine levels (pg/head) in
males decreased in isolated compared to group housed flies but do not differ in recovery flies
compared to group housed in males or females (Two-Way ANOVA – Effect of Social
Experience: $F_{3,27}$ = 3.043, p = 0.0459, but the only significant difference after correcting for
multiple comparison through a Dunnett test is indicated on graph, p=0.0132). Appropriate age
comparisons were made (see methods for details). *p < 0.05. ****p < 0.0001. Bars: mean +/-
s.e.m. N=5-9 for all treatments in social space. N=4 for dopamine quantification. GH: Group
Housed, light grey shade; SH: Single Housed, dark grey shade; REC: Recovery from social
isolation, medium grey shade.
720
Figure 5. $nlg3^{Def1}$ flies have less dopamine, and those levels are not altered by social
experience A. DA levels (pg/head) in CS (grey shade) and $nlg3^{Def1}$ males (blue shade) and
females (red shade). $nlg3^{Def1}$ flies had less DA when group housed and isolated compared to

group housed CS, with females showing a lesser amount overall (Two-way ANOVA, Effect of
Genotype: $F_{1,7}$= 17.21, p = 0.0043 and Effect of Sex: $F_{1,7}$ = 6.107, p = 0.0428). **B.** DA levels
(pg/head) in male (blue shade) and female (red shade) group housed and recovery *nlg3^{Def1}* flies
are unchanged with social experience. Appropriate age comparisons were made (see methods for
details). Bars: mean +/- s.e.m. N=3-6 for all treatments. GH: Group Housed. SH: Single Housed.
REC: Recovery from social isolation. ***P < 0.001. **P < 0.01. *P < 0.05.
730
Figure 6: A. Manipulations performed in this study. B. Working model: Nlg3 and DA are
part of a pathway responsible for the modulation of social behaviour after isolation in males.
Nlg3 would be required downstream of DA, for proper DA signaling in response to the social
environment, with some feedback regulation. Females do require Nlg3 for a typical response to
isolation, but other neurotransmitters or neuromodulators, responsive to social experience in
females, may be interacting with Nlg3. Blue outline: Drosophila brain outline. Created with
BioRender.com.
738
739
740

741   **Figure 1**

Isolation

Recovery

742
743

744   <u>**Figure 2:**</u>

747 <u>**Figure 3**</u>

748
749

750   **Figure 4**

752   **Figure 5**

753
754

755   **Figure 6**

756
757
758