1

A non-conducting role of the Ca$_v$1.4 Ca^{2+} channel drives homeostatic plasticity at the cone

photoreceptor synapse

J. Wesley Maddox[1,6], Gregory J. Ordemann[1,6], Juan de la Rosa Vázquez [1], Angie Huang,[1] Christof

Gault,[1] Serena R. Wisner[2,3], Kate Randall[1], Daiki Futagi[4], Steven H. DeVries[4], Mrinalini Hoon[2,5]

and Amy Lee[1*]

[1]Dept of Neuroscience, University of Texas-Austin, Austin, TX 78712, USA

[2]Dept. of Ophthalmology and Visual Sciences, University of Wisconsin- Madison, Madison, WI,

53706, USA

[3]Neuroscience Training Program, University of Wisconsin-Madison, Madison WI 53706 USA

[4]Dept. of Ophthalmology, Northwestern University Feinberg School of Medicine, Chicago, IL,

60611, USA

[5] McPherson Eye Research Institute, Madison WI 53706 USA

[6]These authors contributed equally

[*]Corresponding author, email: amy.lee1@austin.utexas.edu

ABSTRACT

In congenital stationary night blindness type 2 (CSNB2)—a disorder involving dysfunction of the $Ca_v1.4$ Ca^{2+} channel—visual impairment is relatively mild considering that $Ca_v1.4$ mediates synaptic transmission by rod and cone photoreceptors. Here, we addressed this conundrum using a $Ca_v1.4$ knockout (KO) mouse and a knock-in (KI) mouse expressing a non-conducting $Ca_v1.4$ mutant. Surprisingly, aberrant Ca_v3 currents were detected in cones of the KI and KO but not wild-type mice. Cone synapses, which fail to develop in KO mice, are present but enlarged in KI mice. Moreover, light responses in cone pathways and photopic visual behavior are preserved in KI but not in KO mice. In CSNB2, we propose that Ca_v3 channels maintain cone synaptic output provided that the Ca^{2+}-independent role of $Ca_v1.4$ in cone synaptogenesis remains intact. Our findings reveal an unexpected form of homeostatic plasticity that relies on a non-canonical role of an ion channel.

KEYWORDS: photoreceptor, Ca^{2+} channel, ribbon synapse, retina, synaptogenesis

INTRODUCTION

At the first synapse in the visual pathway, the light-dependent graded electrical signals produced in rod and cone photoreceptors gates the release of glutamate onto postsynaptic neurons. To accomplish this task, photoreceptor synapses are specialized with a ribbon organelle, which helps prime synaptic vesicles [1, 2] and postsynaptic dendrites from horizontal and bipolar cells that invaginate deep within the terminal [3]. A variety of proteins interact with the ribbon and synaptic vesicles near release sites (i.e., active zones) [4]. The importance of these proteins for vision is illustrated by the numerous inherited retinal diseases linked to mutations in their encoding genes [5].

One such gene is *CACNA1F*, which encodes the voltage-gated Ca^{2+} (Ca_v) channel expressed in retinal photoreceptors, $Ca_v1.4$ [6-8]. Among the sub-family of $Ca_v1.x$ L-type channels,

$Ca_v1.4$ exhibits unusually slow inactivation that is well-matched for supporting the tonic, Ca^{2+}-

dependent release of glutamate from photoreceptor synaptic terminals in darkness [9, 10]. More than

200 mutations in *CACNA1F* cause vision disorders including congenital stationary night blindness

type 2 (CSNB2) [11, 12]. These mutations are broadly categorized as producing a gain of function or

loss of function in $Ca_v1.4$ [13]. How these mutations in *CACNA1F* lead to the variable clinical

phenotypes of CSNB2 is largely unknown. Symptoms may include strabismus, low visual acuity,

and in many cases, night blindness [14, 15]. The latter suggests a primary defect in rod pathways,

which is surprising given that knockout (KO) mice are completely blind and lack any evidence of

either rod or cone synaptic responses [6, 7, 16]. A major caveat is that rod and cone synapses fail to

form in $Ca_v1.4$ KO mice [7, 16-18]. Thus, $Ca_v1.4$ KO mice are not suitable for studies of how *CACNA1F*

mutations differentially affect rod and cone pathways or for efforts to uncover how the biophysical

properties of $Ca_v1.4$ shape photoreceptor synaptic release properties.

Here, we overcome this hurdle with a knock-in mouse strain (G369i KI) expressing a non-

conducting mutant form of $Ca_v1.4$ [19]. We show that cone ribbon synapses in G369i KI mice are

largely preserved and that downstream signaling through cone pathways, although greatly

impaired, can support visual function. This novel mechanism requires the ability of the $Ca_v1.4$

protein, independent of its Ca^{2+} conductance, to nucleate the assembly of cone ribbon synapses

and involves an aberrant Ca_v3 (T-type) conductance that appears when $Ca_v1.4$ Ca^{2+} signals are

compromised.

RESULTS

Ca^{2+} currents in cones are mediated by Ca_v3 channels upon $Ca_v1.4$ loss-of-function

A prevailing yet unsupported hypothesis regarding the relatively mild visual phenotypes in

CSNB2 is that additional Ca_v subtypes may compensate for $Ca_v1.4$ loss of function in cones. If

so, then Ca^{2+} currents (I_{Ca}) mediated by these subtypes should be evident in cones of $Ca_v1.4$ KO

and G369i KI mice. The G369i mutation is an insertion of a glycine residue in a transmembrane

domain, which prevents Ca^{2+} permeation through the channel [19]. Rods of G369i KI mice lack any

evidence of I_{Ca}, despite the normal presynaptic clustering of the mutant channel [19]. To test if this

might differ in G369i KI cones, we first analyzed the localization of the mutant G369i $Ca_v1.4$

channels in cones by immunofluorescence with antibodies against $Ca_v1.4$, as well as cone

arrestin (CAR) and CtBP2 to label cone terminals (i.e., pedicles) and ribbons, respectively (Fig.

1). In $Ca_v1.4$ KO mouse retinas, cone pedicles were shrunken and retracted into the outer nuclear

layer (ONL, Fig. 1a) and had malformed ribbons (Fig. 1b). In contrast, cone pedicles in G369i KI

mice were normally localized in the outer plexiform layer (OPL, Fig. 1a) and were populated by

multiple ribbons (Fig. 1b). As in WT mice, labeling for $Ca_v1.4$ was clustered near elongated

ribbons in cones of G369i KI mice (Fig. 1b). Thus, unlike in $Ca_v1.4$ KO mice, the mutant $Ca_v1.4$

protein is normally localized and supports the integrity of cone pedicles and ribbons in G369i KI

mice.

We next compared patch clamp recordings of cones in retinal slices of adult WT, G369i

KI, and $Ca_v1.4$ KO mice under conditions designed to isolate I_{Ca} (Fig. 2). In WT cones, there was

a large, sustained I_{Ca} that activated around -50 mV and peaked near -20 mV, consistent with the

properties of $Ca_v1.4$ (Fig. 2a-d). A small-amplitude I_{Ca} was detected in cones of G369i KI and

$Ca_v1.4$ KO mice but activated at significantly more negative voltages than in WT cones (Fig. 2a-

d; Table 1). Moreover, I_{Ca} in the G369i KI and $Ca_v1.4$ KO cones activated around -60 mV and

peaked near -35 mV. This aberrant I_{Ca} was not sustained as in WT cones but inactivated rapidly

during 50-ms (Fig. 2b) and 500-ms (Fig. 2e) step depolarizations. Overlay of the conductance-

voltage (Fig. 2d) and inactivation curves (Fig. 2f) revealed a sizeable window current. These

features of I_{Ca} in G369i KI and $Ca_v1.4$ KO cones resembled those of Ca_v3 T-type channels rather

than $Ca_v1.4$ [20].

Although Ca_v3 channels were reported in patch clamp recordings of cone pedicles in WT

mouse cones [21], we did not observe a low voltage-activated component of I_{Ca} in the I-V curve from

our recordings of WT mouse cone somas, which would be indicative of a Ca_v3 subtype (Fig. 2a,c).

Moreover, I_{Ca} in WT cones was blunted by the Ca_v1 antagonist isradipine (Fig. 3a) but not the

Ca_v3 antagonist, ML 218 (Fig. 3b; ML 218 caused a minor suppression of I_{Ca} in some WT cones,

which could be attributed to weak activity on $Ca_v1.4$, Supp.Fig. S1). In contrast, I_{Ca} in G369i KI

cones was significantly suppressed by ML 218 but not by isradipine (Fig. 3a,b). To further test for

a Ca_v3 contribution to I_{Ca}, we analyzed cones of ground squirrel retina, where the large amplitude

of I_{Ca} facilitates pharmacological analyses. Consistent with its actions on $Ca_v1.4$ in WT mouse

cones (Fig. 3c) and transfected HEK293T cells (Supp.Fig. S1), ML 218 caused an insignificant

inhibition of peak I_{Ca} (-7.0 ± 20.8%, n = 6 cones) as well as a negative shift in the voltage-

dependence of activation in ground squirrel cones (Supp.Fig. S2). As a positive control, we

confirmed that ML 218 blocked a prominent Ca_v3-type current in ground squirrel type 3a OFF

cone bipolar cells (Supp.Fig. S2). Application of isradipine followed by ML 218 resulted in a time-

and voltage-dependent suppression of the residual I_{Ca}, which is a hallmark of Ca_v1 inhibition by

dihyropyridine antagonists such as isradipine (Fig. 3d, Supp.Fig. S2)[22]. Finally, we also performed

patch clamp recordings of cone terminals in macaque retina where there was no evidence of a

Ca_v3 current (Supp.Fig. S3). We conclude that Ca_v3 channels contribute significantly to I_{Ca} in

cones only when $Ca_v1.4$ Ca^{2+} signals are absent.

Cone synaptogenesis relies on the $Ca_v1.4$ protein but not its Ca^{2+} conductance

As shown in previous studies [7, 16-18], CtBP2 labeled stubby, sphere-like structures

resembling immature ribbon material in $Ca_v1.4$ KO mice (Fig. 1). The presence of elongated

ribbons in G369i KI mice suggested that, as in rods [19], the non-conducting mutant $Ca_v1.4$ protein

may support the molecular assembly of cone synapses. To this end, we analyzed cone synapses

by immunofluorescence and confocal microscopy. Along with the major constituents of the ribbon,

CtBP2 and RIBEYE [23], presynaptic proteins such as bassoon and members of the postsynaptic

signaling complex in depolarizing (ON) cone bipolar cells (GPR179, mGluR6, and TRPM1 [24])

were enriched near cone ribbons in G369i KI mice as in WT mice (Fig. 4a). Compared to WT

mice, the labeled structures occupied a larger volume (Fig. 4b,c), albeit at a much lower density

(Fig. 4d), which increased linearly with the volume of the pedicle in G369i KI mice (Fig. 4e-i),

perhaps as a homeostatic response to $Ca_v1.4$ loss-of-function.

The cone synapse is structurally complex, with the dendritic tips of two horizontal cells

and an intervening ON cone bipolar cell invaginating deeply into the pedicle near the ribbon [3]. To

test how the switch in Ca_v subtypes might affect this arrangement of postsynaptic partners, we

generated 3D reconstructions of cone synapses by serial block-face scanning electron

microscopy (SBFSEM; Fig. 5). As with the enlarged synaptic contacts (Fig. 4), there was some

evidence of structural modifications in G369i KI pedicles. Compared to WT pedicles, ribbons

appeared disorganized in G369i KI pedicles which extended telodendria laterally rather than

apically (Fig. 5a). In addition, a slightly larger fraction of synaptic sites in the G369i KI pedicle

(35% vs 14% in WT) formed incorrect postsynaptic partnerships (Fig.5b-d'; Table 2).

Nevertheless, the number of ribbons is normal in G369i KI cones and about half of the ribbons

making invaginating contacts with the appropriate cell types (*i.e.,* both horizontal cells and cone

bipolar cells; Table 2). Therefore, while necessary for the structural refinement of the cone

synapses, $Ca_v1.4$ Ca^{2+} signals are largely dispensable for cone synapse assembly.

Cone signaling to a postsynaptic partner is intact in G369i KI mice

The preservation of cone synapses in G369i KI mice allowed the unique opportunity to

test whether a Ca_v subtype other than $Ca_v1.4$ could support ribbon-mediated synaptic release. To

this end, we compared synaptic transmission between cones and horizontal cells (HCs, Fig. 6) in

retinal slices of WT and G369i KI mice. In darkness, glutamate released from cones depolarizes

horizontal cells via activation of α-amino-3-hydroxy-5-methyl-4-isoxazolepropionic

(AMPA)/kainate receptors [25, 26]. The resulting excitatory postsynaptic current (EPSC) undergoes

a decline in response to light stimuli that hyperpolarize cones [27]. In WT HCs, a 1 s light pulse

(λ=410 nm) inhibited the standing EPSC, which is reflected as an outward (hyperpolarizing "ON")

current (Fig. 6a). Upon termination of the light pulse, an inward (depolarizing "OFF") current

signaled the resumption of the EPSC (Fig. 6a). Both the ON and OFF components of the EPSCs

in WT HCs increased with light intensity, reflecting the impact of luminance on the presynaptic

membrane potential of cones and the subsequent change in glutamate release from their

terminals (Fig. 6b). The ON and OFF components of EPSCs in G369i KI HCs were measurable,

although lower in amplitude than in WT (Fig. 6a,b). The EPSC in darkness and the light response

were abolished by the AMPA/kainate receptor antagonist, DNQX, in both WT and G369i KI

horizontal cells (Fig. 6c,d). Thus, while greatly impaired, cone synapses can transmit light

information to postsynaptic partners in the retina of G369i KI mice.

Light responses of bipolar cells and visual behavior is spared in G369i KI but not Ca$_v$1.4 KO mice

While horizontal cells act as inhibitory interneurons that modulate photoreceptor output,

the vertical dissemination of visual information from cones to the inner retina is relayed by

glutamate to ON and OFF cone bipolar cells (CBCs) which express mGluR6 and AMPA/kainate

receptors, respectively. To test whether cone synaptic transmission to CBCs is intact in G369i KI

mice, we recorded electroretinograms (ERGs) under light-adapted conditions using Ca$_v$1.4 KO

mice as a negative control (Fig. 7a-d). In these recordings, the light-induced response of

photoreceptors and the postsynaptic response of ON CBCs corresponds to the a- and b-waves,

respectively. Unlike in Ca$_v$1.4 KO mice, the a-waves of G369i KI mice were like those in WT mice,

which indicates that cones do not degenerate in this mouse strain. While reduced in amplitude,

the b-wave was measurable in G369i KI mice and significantly larger than in Ca$_v$1.4 KO mice at

the highest light intensities (Fig. 7a,b). We also recorded flicker ERGs using 10 Hz light stimuli

that can isolate cone pathways involving both ON and OFF CBCs. In WT mice, flicker responses

exhibited two peaks, one at a lower irradiance (-2 log cd·s/m^2) and one at higher irradiance (0.5

log cd·s/m^2). The peak at the lower irradiance is attributed to responses in both rod and cone

pathways, and the peak at higher irradiance is attributed to responses exclusively in cone

pathways (Fig. 7c,d)[28, 29]. While significantly lower in amplitude than in WT mice, flicker responses

in G369i KI mice showed a similar non-monotonic relation. Compared to $Ca_v1.4$ KO mice, G369i

KI mice showed peak flicker responses that did not differ at -2 log $cd \cdot s/m^2$, but were significantly

higher at 0.5 log $cd \cdot s/m^2$ (Fig.7d). The inverted flicker responses at higher illuminations in G369i

KI mice (Fig. 7c) were absent in $Ca_v1.4$ KO mice and may result from the hyperpolarizing

contribution of cone-to-OFF CBC transmission. These results suggest that cone-to-CBC signaling

is intact in G369i KI mice.

To validate these results with respect to vision-guided behavior, we used a swim test that

assesses the ability of mice to identify a visible platform [30]. In darkness, G369i KI and $Ca_v1.4$ KO

mice took significantly longer to find the platform compared to WT mice (Fig. 7e,f). These results

are consistent with our flicker response assays (Fig.7c,d) as well as the absence of I_{Ca} in rods

and rod-to-rod bipolar cell synaptic transmission in G369i KI mice [19]. In daylight conditions, G369i

KI mice but not $Ca_v1.4$ KO mice performed as well as WT mice (Fig. 7e,f). Thus, G369i KI mice

are unique in retaining visual function under photopic but not scotopic conditions. Collectively, our

results suggest that Ca_v3 channels can support cone synaptic responses and visual behavior in

G369i KI but not $Ca_v1.4$ KO mice.

DISCUSSION

The nervous system has remarkable abilities to adapt to pathological perturbations in

neuronal activity. In the retina, ablation or degeneration of photoreceptors triggers various

postsynaptic mechanisms that maintain some level of visual function in rod or cone pathways.

These include remodeling of bipolar cell dendrites and their synapses[31-33] as well as changes in

the sensitivity of bipolar cells to photoreceptor input and inhibitory modulation[33, 34]. To our

knowledge, this study provides the first evidence for a presynaptic form of homeostatic plasticity

that originates within photoreceptors. Using $Ca_v1.4$ KO and G369i KI mice, we identify the

upregulation of a Ca_v3 conductance as a common response to $Ca_v1.4$ loss-of-function in cones.

However, Ca_v3 channels can only compensate for $Ca_v1.4$ loss-of-function when cone synapse

structure is maintained. Thus, our results also highlight a crucial, non-conducting role for the

$Ca_v1.4$ protein that allows cone synapses to function in the absence of $Ca_v1.4$ Ca^{2+} signals.

A non-canonical role for $Ca_v1.4$ in regulating cone synapse assembly

A major finding of our study is that cone synapse formation requires the $Ca_v1.4$ protein but

not Ca_v-mediated Ca^{2+} influx. As shown for rod synapses [19], ribbons and other components of the

pre- and post-synaptic complex assemble normally at cone synapses in G369i mice (Figs. 1,4,5).

Ca_v3 Ca^{2+} signals are dispensable for this process since their presence in $Ca_v1.4$ KO cones is not

accompanied by any semblance of ribbon synapses (Fig.1B). An intimate relationship between

$Ca_v1.4$ and ribbons is supported by the colocalization of $Ca_v1.4$ and RIBEYE puncta resembling

ribbon precursor spheres in the developing OPL[7]. Moreover, light adaptation, which decreases

the size of the ribbon, leads to a reduction in $Ca_v1.4$ labeling in mouse retina[2]. While evidence for

the binding of RIBEYE to $Ca_v1.4$ is lacking, such an interaction could support the oligomerization

of RIBEYE A and B domain within a macromolecular complex [35]. Alternatively, $Ca_v1.4$ could

pioneer sites of ribbon assembly perhaps by serving as a docking or nucleation site for the active

zone. Regardless of the mechanism, our findings show that the formation of photoreceptor

synaptic complexes does not require Ca^{2+} influx through $Ca_v1.4$ channels.

Despite a normal molecular organization, G369i KI cone synapses were enlarged and

made errors in postsynaptic partner selection. While I-V curves predict similar peak I_{Ca} amplitudes

in cones of WT and G369i KI mice near the membrane potential of cones in darkness (-45 to -50

mV [36], Fig.1c), the strong inactivation of Ca_v3 channels predicts greatly diminished Ca^{2+} signals

in G369i KI pedicles. Paradoxically, reductions in presynaptic Ca^{2+} in rod photoreceptors are

thought to cause illumination-dependent shrinkage of ribbons *in vivo* and *in vitro* [2, 37, 38]. However,

an inverse correlation between presynaptic Ca^{2+} influx and ribbon synapse size is seen in sensory

hair cells in zebrafish [39]. The mechanism involves decreased mitochondrial Ca^{2+} uptake and

increase in the redox state of nicotinamide adenine dinucleotide (NAD^+/NADH ratio) [40]. If a similar

mechanism applied in cones, Ca_v3 Ca^{2+} signals in G369i KI cones may decay too quickly to

enable mitochondrial Ca^{2+} uptake mechanisms that trim ribbons. The size of ribbons and the

postsynaptic specialization increased linearly with pedicle volume in G369i KI but not in WT mice

(Fig. 4e-i), which could represent a form of homeostatic synaptic scaling as has been

demonstrated at the neuromuscular junction [41, 42]. Similarly, the presence of some non-

invaginating contacts with incorrect partner pairings in G369i KI mice (Fig.5, Table 2) could

represent a compensatory response to weakened synaptic output. Considering that only a subset

of CBC subtypes re-wire correctly following partial ablation of cones in immature mice [32], the

structurally normal cone synapses of G369i KI mice could involve contacts with CBCs that are

unusually resilient to loss of presynaptic input.

While it is well-established that $Ca_v1.4$ is the main Ca_v subtype in mouse cones, Ca_v3

channels, in particular $Ca_v3.2$, have been reported to be expressed in mouse cones by

electrophysiology [21] and single cell RNA-seq studies [21, 43, 44]. By pharmacological and other

criteria, we found no evidence for a functional contribution of Ca_v3 in our recordings of cones in

WT mice (Figs. 2,3), ground squirrels, or macaque (Supp.Figs. S2,S3). A caveat of using

dihydropyridine antagonists such as isradipine to isolate the contribution of Ca_v1 from other Ca_v

subtypes is the relatively low sensitivity of Ca_v1 channels to these drugs and their strong voltage-

dependence. At a holding voltage of -90 mV, isradipine at 1 μM causes only ~80% inhibition of

$Ca_v1.4$ with greater block at depolarized voltages [22], which agrees with our recordings of cones

in WT mice (Fig.3a) and ground squirrels (Fig.3d, Supp.Fig.S2). Moreover, Ca_v3 blockers such

as Z944 [45] and ML218 (Supp.Fig. S2) have additional activity on Ca_v1 channels at micromolar

concentrations including effects on current amplitude and activation voltage. Thus, currents

mediated by $Ca_v1.4$ that are sensitive to Ca_v3 blockers (e.g, >5 µM ML218) and spared by

dihydropyridines at negative voltages may be mistaken as being mediated by Ca_v3. In our

experiments, the biophysical properties of the isradipine-sensitive I_{Ca} in cones of WT mice and

ground squirrels resembled only those of Ca_v1, whereas those of the ML218-sensitive I_{Ca} in cones

of G369i KI mice resembled only those of Ca_v3. Therefore, we favor the interpretation that Ca_v3

contributes to I_{Ca} in mouse cones only upon silencing of the $Ca_v1.4$ Ca^{2+} conductance.

Consistent with the diminutive Ca_v3-mediated I_{Ca}, light responses of HCs were evident but

smaller in G369i KI than in WT mice. Due to their slow activation and strong inactivation [46], Ca_v3

channels may have a reduced ability to fuel the Ca^{2+} nanodomains that support fast and sustained

components of release, both of which occur only at ribbon sites in cones[1, 47]. Ca_v3 channels may

also be located further from the ribbon than $Ca_v1.4$, thus lowering the efficiency of coupling to

exocytosis. Unfortunately, it was not possible to test this with commercially available antibodies

against $Ca_v3.2$, which yielded identical patterns of immunofluorescence in the OPL of WT and

$Ca_v3.2$ KO mice (data not shown). A detailed analysis of the Ca_v3 subtype(s) and their subcellular

localization in G369i KI cones is required to unravel the shortcomings of Ca_v3 channels with

respect to cone synaptic release.

Based on extremely heterogeneous clinical presentations, CSNB2 manifests as a

spectrum of visual disorders that originate from various mutations in $CACNA1F$[14, 48]. Even though

some CSNB2 mutant $Ca_v1.4$ channels may traffic normally to the plasma membrane, many of

these mutations are expected to produce non-functional, non-conducting $Ca_v1.4$ channels.[49, 50]

Yet, the visual phenotypes of CSNB2 patients are not as severe as the complete blindness in

$Ca_v1.4$ KO mice, which lack any $Ca_v1.4$ protein expression and exhibit no signs of visual behavior

(Fig.7e,f)[51]. Collectively, our results suggest that G369i KI mice accurately model $Ca_v1.4$

channelopathies in CSNB2 patients that are characterized by a greater impairment in rod than in

cone pathways. This interpretation is supported by our findings that G369i KI mice exhibit

horizontal cell responses to bright but not dim illumination (Fig. 6), ERG responses under

conditions of light adaptation (Fig.7a-d) but not dark-adaptation [19], and visual behavior under

photopic but not scotopic conditions (Fig. 7d). Together with the enlargement of synaptic sites,

modest levels of synaptic release from cones of G369i KI mice may be sufficient to support nominal transmission of visual information through cone pathways. Furthermore, we acknowledge that G369i KI mice could also exhibit homeostatic alterations in the inner retina, which are known to support visual function when photoreceptor input is severely compromised [32-34]. Future studies of the retinal circuitry and visual behavior of G369i KI mice could identify compensatory pathways that are recruited upon Ca$_v$1.4 loss-of-function and how they might be targeted in novel therapies for CSNB2 and related disorders.

ACKNOWLEDGEMENTS

J.W.M., G.J.O., J.D.V., A.H., K.R., and A.L. were supported by NIH grants EY026817 (to A.L.), EY029953 (to J.W.M.), and startup funds from the University of Texas-Austin (to A.L.); S.R.W. and M.H. were supported by NIH grant EY031677 (to M.H.), an unrestricted grant from RPB to UW Madison Dept. of Ophthalmology and a McPherson ERI professorship to M.H.; D.F. and S.H.D. were supported by NIH grants EY012141 and EY032506, an unrestricted grant from RPB to the Northwestern University Dept. of Ophthalmology, and an RPB International Travel Award. The authors thank S. Knecht and R. Wong for assistance with serial EM image collection, E. Seidemann and B. Shukla for the generous donation and collection of macaque retinal tissue, respectively, and M. McCall and F. Vinberg for advice on ERGs.

AUTHOR CONTRIBUTIONS

Conceptualization, J.W.M., G.O., A.L.; Methodology, J.W.M., G.O., J.D.V., A.L., M.H., S.H.D., D.F.; Investigation, J.W.M., G.O., J.D.V., A.H., C.G., K.R., S.R.W., S.H.D., D.F.; Writing—original draft, A.L.; Writing—review and editing, J.W.M., G.J.O., M.H., S.H.D., A.L.; Funding acquisition, J.W.M., M.H., S.H.D., A.L.

313 **DECLARATION OF INTERESTS**

314 The authors declare no competing interests.

316 **Table 1. Comparison of parameters from electrophysiological recordings of cones.**

	C_M (pF)	P-value	n	R_M (MΩ)	P-value	n
WT	4.88 (3.86, 5.01)	--	11	3.57 (2.81, 4.65)	--	11
G369i KI	4.08 (3.88, 4.34)	0.73[a]	9	5.06 (4.04, 8.24)	0.04[a]	9
Ca_v1.4 KO	3.31 (3.00, 3.76)	0.001[a], 0.04[b]	6	8.21 (7.26, 9.06)	0.001[a], 0.47[b]	6
G-V	V_h (mV)	P-value	n	k	P-value	n
WT	-37.54 (-38.91, -35.13	--	13	2.75 (2.47, 4.32)		13
G369i KI	-47.59 (-50.07, -46.72)	0.0001[a]	9	2.99 (2.68, 4.16)	0.99[a]	9
Ca_v1.4 KO	-44.98 (-50.15, -43.27)	0.085[a], 0.99[b]	3	4.33 (3.73, 5.22)	0.34[a], 0.49[b]	3
Steady state inactivation	V_h (mV)	P-value	n	k	P-value	n
WT	-49.06 (-57.95, -41.82)	--	8	-7.25 (-12.30, -4.97)	--	8
G369i KI	-59.40 (-62.73, -58.01)	0.01[a]	8	-3.81 (-5.23, -2.46)	0.12[a]	8
Ca_v1.4 KO	-56.62 (-61.65, -56.56)	0.88[a], 0.88[b]	3	-3.67 (-5.17, -1.19)	0.14[a], 0.99[b]	3
	Tau activation	P-value	n	Tau deactivation	P-value	n
WT	2.04 (1.26, 2.71)	--	11	0.88 (0.62. 2.09)	--	6
G369i KI	3.32 (2.99, 4.06)	0.001[a]	9	3.40 (2.62, 4.34)	0.004[a]	6

Values represent median (25[th], 75[th] quartiles). V_h and k were determined from Boltzmann fits of
the G-V and steady-state inactivation curves. Time constant (tau) for activation was obtained from
exponential fit of the rising phase of I_{Ca} evoked by a 50-ms test pulse to a voltage near the peak
of the I-V. Tau deactivation was determined from exponential fit of the decay of the tail current
evoked by repolarization to -90 mV from +20 mV. C_M, membrane capacitance; R_M, input
resistance. P-values were determined by Kruskal Wallis test. [a], relative to WT. [b], relative to G369i
KI.
324

325 **Table 2. Comparison of parameters for cone synapse organization**

	WT		G369i KI	
	Cone 1	Cone 2	Cone 1	Cone 2
# of ribbons	12	8	8	6
Type of contact: % total (fraction of total)				
Invaginating	58.3 (7/12)	87.5 (7/8)	50 (4/8)	50 (3/6)
Non-invaginating	41.7	12.5	50	50

	(5/12)	(1/8)	(4/8)	(3/6)
Partner composition: % total (fraction of total)				
HC/BC	83.3	87.5	50.0	66.7
HC only	8.3	12.5	25.0	33.3
BC only	8.3	0	12.5	0
Glia only	0	0	12.5	0

326 Results represent analysis of n = 2 pedicles reconstructed by serial block face scanning electron
327 microscopy in images obtained from WT or G369i KI mice (N= 1 animal each). Sites were counted
328 as 'invaginating' if postsynaptic partners were enveloped on all sides by the cone terminal for at
329 least a couple of consecutive image planes.
330

331
332
333
FIGURE LEGENDS

Figure 1. Cone pedicles and the mutant Ca$_V$1.4 channel are normally localized in the OPL

of G369i KI mice but not Ca$_V$1.4 KO mice. Confocal images of the ONL and OPL of WT, G369i

KI, and Ca$_V$1.4 KO mice labeled with antibodies against cone arrestin (CAR), CtBP2, and Ca$_V$1.4.

a, Inverted images of CAR labeling. Lower panels correspond to boxed region of the upper panels

and depict pedicles labeled by CAR antibodies (dotted outlines). Cone pedicles remain within the

OPL of WT and G369i KI retina (arrows) but are misshapen and retracted in the ONL of the Ca$_V$1.4

KO retina (arrowheads). *b*, High magnification, deconvolved images show Ca$_V$1.4 labeling near

cone ribbons in WT and G369i KI pedicles (arrows) and ribbon spheres without Ca$_V$1.4 labeling

in the Ca$_V$1.4 KO pedicle (arrowheads).

Figure 2. A low-voltage activated I_{Ca} is present in cones of G369i KI and Ca$_V$1.4 KO but

absent in WT mice. *a,b*, Representative traces of I_{Ca} evoked by voltage ramps (a) and voltage

steps (b). *c,d*, I-V (c) and G-V (d) were plotted against test voltage for I_{Ca} evoked by 50-ms voltage

steps from a holding voltage of -90 mV. Numbers of cells: WT, n = 13; G369i KI, n = 9; CaV1.4

KO, n = 3. *e*, Representative I_{Ca} traces (top) and voltage-protocol (bottom) for steady-state

inactivation. I_{Ca} was evoked by a conditioning pre-pulse from -90 mV to various voltages for 500

ms followed by a test pulse to -30 mV for 50 ms. *f*, I/I_{max} represents the current amplitude of the

test pulse normalized to current amplitude of the pre-pulse and was plotted against pre-pulse

voltage. Numbers of cells: WT, n = 8; G369i KI, n = 8; Ca$_v$1.4 KO, n = 3. In graphs in c,d, and f,

smooth lines represent Boltzmann fits, symbols and bars represent mean ± SEM, respectively. In

graph in f, line without symbols represents G-V curve for G369i KI cones replotted from d. Shaded

region indicates window current for G369i KI cones.

Figure 3. Pharmacological characterization of I_{Ca} in cones of WT and G369i KI mice and

ground squirrel.

a,b, Analysis of mouse cones. *Left*, representative traces for I_{Ca} evoked by voltage ramps in cones

of WT or G369i KI mice before (baseline) and after exposure to 1 μM of isradipine (ISR, *a*, WT, n

= 4; G369i KI, n = 5) or 5 μM ML 218 (b, WT, n = 5; G369i KI, n = 5). *Right*, I_{Ca} amplitudes before

(-) and during (+) perfusion of the blocker on the same cells. Each point represents a different

cell. *, p < 0.05 by paired t-test. *c*,*d*, Analysis of ground squirrel cones. Representative traces

corresponding to baseline-corrected I_{Ca} evoked by voltage ramps (*c,d*, left) and corresponding G-

V plot (*c*, right) before and during application of ML 218 (c) or ISR alone or ISR+ML218 (d). In *d,*

I_{Ca} amplitudes are plotted before (-) and after (+) block for cones in the superior (sup.; n = 7

cones), middle (mid.; n = 5 cones), and inferior (inf.; n = 5 cones) thirds of the retina. The I_{Ca}

blocked by ISR alone was measured at the peak of the control I-V curve between -40 and -20

mV. The I_{Ca} amplitude blocked by adding ML218 was measured as the average current between

-50 and -45 mV before ML218 addition relative to zero current following the addition of ML218.

The changes produced by ML218 were small but nonetheless significant (sup., n = 7, ISR: p <

0.0001, ML218: p = 0.0047; mid., n = 5, ISR: p < 0.0001, ML218: p = 0.0001; inf., n =5, ISR: p =

0.0019, ML218: p = 0.0460; two-tailed t test).

Figure 4. Immunofluorescence characterization of cone synapses in WT and G369i KI mice.

a, Confocal images of the OPL of WT and G369i KI mice labeled with antibodies against cone arrestin (CAR) and proteins that are presynaptic (CtBP2, bassoon) or postsynaptic (GPR179, TRPM1, mGluR6). Every other panel shows high-magnification, deconvolved images of single pedicles labeled with cone arrestin (rod spherule-associated signals were removed for clarity). Arrows indicate ribbon synapses, which appear enlarged in the G369i KI pedicles. ***b-d***, Violin plots represent volume occupancy of labeling for each synaptic protein normalized to their respective CAR-labeled pedicles. ***e-i***, Dependence of synapse size on pedicle size. Volumes corresponding to labeling of CtBP2 (*e*: $p = 0.051$, $r = 0.4$ for WT; $p < 0.0001$, $r = 0.88$ for G369i KI), $Ca_V1.4$ (*f*: $p = 0.8$, $r = 0.06$ for WT; $p = 0.002$, $r = 0.88$ for G369i KI), bassoon (*g*: $p = 0.1$, $r = 0.34$ for WT; $p < 0.0001$, $r = 0.75$ for G369i KI), mGluR6 (*h*: $p = 0.32$, $r = 0.35$ for WT; $p = 0.002$, $r = 0.73$ for G369i KI) and TRPM1 (*i*: $p = 0.32$, $r = 0.35$ for WT; $p = 0.007$, $r = 0.66$ for G369i KI) are plotted against pedicle volume. Dashed and solid lines represent fits by linear regression for WT and G369i KI, respectively.

Figure 5. Cone synapses form incorrect pairings with postsynaptic partners in G369i KI mice. 3D reconstructions of WT and G369i KI pedicles (n = 2 each) were obtained by SBFSEM. ***a***, 3D renderings showing ribbons (magenta) within cone pedicles (gray) from WT and G369i KI mice. ***b,b'***, 3D renderings (b) show ribbon sites in a WT cone pedicle contacting one horizontal (HC1; yellow) and three bipolar cells (BC1-3; purple). The raw image (*b'*) shows a single plane example of BC1-2 and HC1 contacting the ribbon site. **c-d'**, Single plane raw images (left panels, *c; d*) and 3D reconstructions (right panels, *c; d'*) show ribbon sites within the G369i KI cone pedicle contacting in *c*: horizontal cells (HC1-2) only, CBCs only (BC1-2); and in *d,d'*: glia (the G369i KI cone pedicle contacts a glial cell (orange) and an unknown partner); the glial cell completely envelops the pedicle. Inset in *d'* shows glial-contacting ribbon site (arrow). In other panels, arrows indicate points of contact between ribbons and other postsynaptic elements.

Figure 6. Light responses of horizontal cells are impaired in G369i KI mice. *a-b,* Representative traces from whole-cell patch clamp recordings of horizontal cells held at -70 mV (*a*) and quantified data (*b*) for currents evoked by 1 s pulses of light (λ = 410 nm) plotted against light intensities. In *b*, peak current amplitudes during (ON current) and after (OFF current) the light stimuli were plotted against photon flux per μm^2 ($\Phi_q/\mu m^2$). Data represent mean ± SEM. WT, n = 8; G369i KI, n =9. *c-d,* Representative traces from horizontal cells held at -70 mV (*c*) and quantified data (*d*) for currents evoked by 1-s pulses of light (λ = 410 nm, 1.2 x 10^5 $\Phi_q/\mu m^2$) before, during, and after washout of DNQX (20 μM). In *d,* symbols represent responses from individual cells, n = 5 cells for WT and 3 cells for G369i KI, bars represent mean ± SEM. **, p < 0.01 by paired t-tests.

Figure 7. Photopic vision is reduced but not lost in G369i KI mice. *a*, Representative traces of photopic ERGs recorded in the presence of background green light (20 cd · s/m^2) in WT, G369i KI and Ca$_V$1.4 KO mice. Flash intensities are shown at left. Arrows and arrowheads depict the a- and b-waves, respectively. *b*, a-wave (left) and b-wave (right) amplitudes are plotted against light intensity. Symbols and bars represent mean ± SEM. WT, n = 7; G369i KI, n = 5; Ca$_V$1.4 KO, n = 6. For a-waves, there was a significant effect of light intensity (p < 0.001) and genotype (p = 0.0086) by a mixed-effects model; by post-hoc Tukey test, there was no significant difference between WT and G369i KI at any light intensity. For b-waves at the highest light intensity, WT vs G369i KI, p = 0.0001; WT vs. Ca$_V$1.4 KO, p < 0.0001; G369i KI vs. Ca$_V$1.4 KO, p = 0.0038, Two-way ANOVA with Tukey's post hoc analysis. *c,d*, Representative traces (*c*) and quantified data (*d*) for 10 Hz flicker responses evoked by white light flashes of increasing luminance (from -4 to 2 log cd· s/m^2). Arrows in *c* depict inverted waveform responses in G369i KI mice that are absent in CaV1.4 KO mice. Symbols and bars represent mean ± SEM. WT, n = 4; G369i KI, n = 5; Ca$_V$1.4

KO, n = 5. At each of the 3 highest light intensities there was a significant difference ($p < 0.05$) in

b-waves of WT vs G369i KI and G369i KI vs $Ca_V1.4$ KO by Two-way ANOVA with Tukey's post

hoc analysis. *e,* Representative swim path traces of WT, G369i KI and $Ca_V1.4$ KO mice from the

visible platform swim tests performed in the dark (upper traces) and light (lower traces). *f,*

Quantified latency to platform. Symbols represent the average of the last 3 swim trials for each

mouse of each genotype for both dark and light conditions. Dotted lines represent the mean. WT,

n = 11; G369i KI, n = 10; $Ca_V1.4$ KO, n = 9. *, $p < 0.05$; ***, $p < 0.001$; ****, $p < 0.0001$; Kruskal-

Wallis one-way ANOVA with Dunn's post hoc analysis.

Methods

Table 3. Key resources used in this study.

Primary Antibodies				
Antibody Name (clone/designation)	**Working Concentration**	**Source**	**Identifier**	**RRID**
Bassoon (SAP7F407/Ms monoclonal)	2 µg/mL	ThermoFisher Scientific	MA1-20689	AB_2066981
$Ca_V1.4$ (Rb Polyclonal)	5 µg/mL	Dr. Amy Lee (Lui et al., 2013)	Ab167	AB_2650487
Cone Arrestin (Rb Polyclonal)	1 µg/mL	Millipore	AB15282	AB_1163387
Cone Arrestin-CF647 (Rb Polyclonal)	5 µg/mL	Millipore	AB15282	AB_1163387
CtBP2 (Clone 16/Ms Monoclonal)	1 µg/mL	BD Biosciences	612044	AB_399431
GPR179 (mAB 1a/Ms Monoclonal)	2 µg/mL	Millipore	MAB427	AB_2069582
mGluR6 (366/Ms Monoclonal)	1 µg/mL	Dr. Theodore Wensel (Agosto et al., 2014, 2018)	N/A	N/A
TRPM1 (545H5/Ms Monoclonal)	1 µg/mL	Dr. Theodore Wensel (Agosto et al., 2018)	N/A	N/A
Secondary Antibodies				
Antibody Name and Conjugate	**Working Concentration**	**Source**	**Identifier**	**RRID**
Goat Anti-Mouse IgG1 Alexa Fluor 488	4 µg/mL	ThermoFisher	A-21121	AB_2535764
Goat Anti-Mouse IgG2a Alexa Fluor 555	4 µg/mL	ThermoFisher	A-21137	AB_2535776
Goat Anti-Rabbit Alexa Fluor 555	4 µg/mL	ThermoFisher	A-21429	AB_2535850
Goat Anti-Rabbit Alexa Fluor 647	4 µg/mL	ThermoFisher	A-21245	AB_2535813

Chemicals		
Reagent or Resource	**Source**	**Identifier**
Adenosine 5'-triphosphate magnesium salt	SIGMA	A9187
Agarose, low gelling temperature	Sigma-Aldrich	A0701
AMES	Sigma, US Biological	Sigma: A1420; US Biological: A1372
Antisedan	Zoetis US LLC	10000449
1,2-Bis(2-aminophenoxy)ethane-N,N,N′,N′-tetraacetic acid (BAPTA)	Sigma-Aldrich	A4926
Cadmium Chloride	Sigma	202908
Calcium chloride dihydrate	Sigma-Aldrich	223506
Cesium Chloride	Fisher Bioreagents	BP210
Cesium methanesulfonate	Sigma-Aldrich	C1426
Creatine phosphate, dipotassium salt	EMD Millipore	237911
D-Glucose	Fisher	D16
DL-*threo*-β-Benzyloxyaspartic acid (DL-TBOA)	Tocris	1223
6,7-Dinitroquinoxaline-2,3-dione disodium salt (DNQX)	Abcam	ab120169
Dimethyl sulfoxide	SIGMA	D2650
Dulbecco's Modified Eagle Medium, high glucose	Gibco	11965-092
Dulbecco's Phosphate Buffered Saline, no calcium, no magnesium	Gibco	14190-144
Ethylene glycol-bis(2-aminoethylether)-N,N,N',N'-tetraacetic acid (EGTA)	Sigma-Aldrich	03777
Fetal Bovine Serum (FBS)	Gibco	16140-071
FuGENE HD Transfection reagent	Promega	E2311
Guanosine 5'-triphosphate sodium salt hydrate	Sigma-Aldrich	G8877
4-(2-Hydroxyethyl)piperazine-1-ethanesulfonic acid (HEPES)	SIGMA	H3375
Systane Hypromellose lubricant eye gel	Alcone	300650474016
Isradipine	Sigma-Aldrich	I6658
Ketamine Injection C IIIN 100 mg/mL	Dechra Veterinary Products, LLC	17033-100-10
Magnesium Chloride hexahydrate	SIGMA	M2393
Methanesulfonic acid	Sigma-Aldrich	471356
Mix-n-Stain Antibody Labeling Kit	Biotium	92238
ML218 Hydrochloride	Tocris	4507
N-Methyl-D-glucamine (NMDG)	Sigma-Aldrich	66930
Nickel Chloride	Sigma	339350
Optimal Cutting Temperature compound	Sakura	4583
Penicillin-Streptomycin	SIGMA	P0781
Picrotoxin	Tocris	1128
Sodium Bicarbonate	MACRON	7412-12
Sodium phosphate monobasic anhydrous	Sigma	S3139
Sodium phosphate dibasic anhydrous	Sigma	RDD022
Sodium Chloride, Crystal	MACRON	7581-12
Potassium Chloride	Fisher Chemical	P217
Sodium hydroxide	Fisher chemical	S318
Sodium lactate	Sigma-Aldrich	1614308
Sodium malate	Sigma-Aldrich	1613881
Sodium pyruvate	Sigma-Aldrich	P2256
Sodium succinate	Sigma-Aldrich	S9637
Strychnine HCl	Sigma	S8753

(3S)-3-[[3-[[4-(Trifluoromethyl)benzoyl]amino]phenyl]methoxy]-L-aspartic acid (TBF-TBOA)	Tocris	2532	
Tetraethylammonium chloride (TEA-Cl)	Sigma-Aldrich	86614	
TRIS	VWR	0497	
Tropicamide ophthalmic solution 1%	Akorn, Inc.	17478-0102-12	
Trypsin-EDTA (0.05%), phenol red	Gibco	25300054	
Anased (Xylazine) Injection 100 mg/mL	Pivetal	46066-0750-02	
4-Ethylphenylamino-1,2-dimethyl-6-methylaminopyrimidinium chloride (ZD7288)	Tocris	1000	
Experimental Models: Organisms/Strains			
Reagent or Resource	**Source**	**Identifier**	**RRID**
HEK293T	ATCC	CRL-3216	CVCL_0063
Mouse: C57BL/6J	Jackson Labs	000664	IMSR_JAX:000664
Mouse: G369i KI	A. Lee [19]	N/A	N/A
Mouse: $Ca_V1.4$ KO	Jackson Labs	017761	IMSR_JAX:017761
Ground Squirrel	TLC	--	--
Software and Algorithms			
Reagent or Resource	**Source**	**Identifier**	
Amira	ThermoFisher Scientific	https://www.thermofisher.com/amira	
cellSens	Olympus	https://www.olympus-lifescience.com/en/software/cellsens	
Espion software	Diagnosys, Inc.	https://info.diagnosysllc.com/software	
FLUOVIEW	Olympus	https://www.olympus-lifescience.com/en/laser-scanning/fv3000	
GraphPad Prism	GraphPad	https://www.graphpad.com	
IGOR Pro	WaveMetrics	https://www.wavemetrics.com	
ImageJ	NIH	https://imagej.nih.gov/ij	
pClamp	Molecular Devices	https://www.moleculardevices.com	
Patchmaster	Harvard Bioscience, Inc.	https://www.heka.com	

Animals

All mouse and macaque experiments were performed in accordance with guidelines

approved by the National Institutes of Health and the Institutional Animal Care and Use

Committees at the University of Texas at Austin. All ground squirrel (*Ictidomys tridecemlineatus*)

procedures performed at Northwestern University were approved by the Institutional Animal Care

and Use Committee. The G369i KI [19] and $Ca_v1.4$ KO mouse strains were bred on the C57BL6/J

background strain for at least 10 generations. Adult male and female mice were used (6-12 weeks

old), and aged-matched C57BL6/J mice were used as the control (WT) animals.

Immunofluorescence

Mice between 6-8 weeks of age were anesthetized using isoflurane and euthanized by

cervical dislocation. Eyes were enucleated and hemisected. The eye cups with retina were fixed

on ice in 4% paraformaldehyde in 0.1 M phosphate buffer (PB) for 30 min. Fixed eye cups were

then washed three times with 0.1 M PB containing 1% glycine followed by infusion of 30% sucrose

at 4°C overnight. The eye cups were orientated along their dorsal-ventral axis and frozen in a 1:1

(wt/vol) mixture of Optimal Cutting Temperature compound and 30% sucrose in a dry

ice/isopentane bath. Eye cups were cryosectioned at 20 µm on a Leica CM1850 cryostat (Leica

Microsystems), mounted on Superfrost plus Micro Slides (VWR), dried for 5 to 10 min at 42°C,

and stored at -20°C until used. Slides with mounted cryosections were warmed to room

temperature, washed with 0.1 M PB for 30 min to remove the OCT/sucrose mixture and blocked

with dilution solution (DS, 0.1 M PB/10% goat serum/0.5% Triton-X100) for 15 min or overnight

at room temperature. All remaining steps were carried out at room temperature. All primary

antibodies and appropriate secondary antibodies were diluted in DS at concentrations specified

in the Key Resources Table (Table 3). Sections were incubated with primary antibodies for 1 h or

overnight and then washed five times with 0.1 M PB. Sections were then incubated with secondary

antibodies for 30 min and then washed five times with 0.1 M PB. Trace 0.1 M PB was removed,

and sections were then mounted with #1.5H coverslips (ThorLabs) using ProLong Glass Antifade

Mountant with or without NucBlue (Thermo Fisher Scientific).

For double labeling with other rabbit polyclonal antibodies, CAR antibodies were

conjugated with the CF647 fluorophore (CAR-647) using the Mix-n-Stain antibody labeling kit

according to the manufacturer's protocol (Biotium). Sections were processed first with rabbit

polyclonal $Ca_V1.4$ or EAAT2 antibodies and corresponding secondary antibodies as described

above. To prevent CAR-647 from binding to any available sites on the anti-rabbit secondary

antibodies previously added to the sections, $Ca_V1.4$ or EAAT2 antibodies were readded to the

sections and incubated for 30 min. After washing, the sections were incubated with CAR-647 for

1 h, followed by wash steps and cover glass mounting.

Immunofluorescence in labeled retinal sections was visualized using an Olympus FV3000

confocal microscope (Tokyo, Japan) equipped with an UPlanApo 60x oil HR objective (1.5 NA).

Images were captured using the Olympus FLUOVIEW software package. Acquisition settings

were optimized using a saturation mask to prevent signal saturation prior to collecting 16-bit. All

confocal images presented are maximum z-projections. Images (256 x 256 pixels) used in

analyses of synaptic proteins were collected using a 30X optical zoom, 0.6 Airy disk aperture, and

voxel size of $0.028 \, \mu m \times 0.028 \, \mu m \times 0.2 \, \mu m \; (X \times Y \times Z)$. Amira segmentation was used for

generating 3D binary masks, and Amira 3D label analysis was used for quantification of

immunofluorescent and masked images. Non-deconvolved images were used for all analyses.

3D binary masks of individual CAR-labeled pedicles were made by setting the threshold 1

standard deviation above mean fluorescence intensity of each 3D image. 3D binary masks of

presynaptic labels (CtBP2, Ca$_V$1.4, and Bassoon) or postsynaptic labels (GPR179, mGluR6, and

TRPM1) within or associated with the pedicle, respectively, were made by setting the threshold 3

standard deviations above the mean fluorescence intensity. To aid in presentation, high-mag

images displayed in Figs.3f, 4a, and 6f were deconvolved using cellSens software (Olympus),

and any immunofluorescence corresponding to rod synaptic proteins was subtracted from the

image.

Molecular biology and transfection

The cDNAs for Ca$_V$1.4 (GenBank: NM_019582), β_{2X13} (GenBank: KJ789960), and $\alpha_2\delta$-

4 (GenBank: NM_172364) were previously cloned into pcDNA3.1 [52]. The cDNA for Ca$_V$3.2

(GenBank: AF051946) was a gift from Dr. Edward Perez-Reyes, University of Virginia. All

constructs were verified by DNA sequencing before use. Human embryonic kidney 293T

(HEK293T) cells were cultured in Dulbecco's Modified Eagle's Medium with 10% FBS at 37°C in

5% CO_2. At 70–80% confluence, the cells were co-transfected with cDNAs encoding human

Ca$_V$1.4 (1.8 μg) β$_{2X13}$ (0.6 μg), α$_2$δ-4 (0.6 μg), and enhanced GFP in pEGFP-C1 (Clonetech, 0.1

μg) or Ca$_V$3.2 (2 μg) and pEGFP-C1 (0.1 μg) using FuGENE 6 transfection reagent according to

the manufacturer's protocol. Cells treated with the transfection mixture were incubated at 37°C

for 24 hr, dissociated using Trypsin-EDTA, and replated at a low density to isolate single cells.

Replated cells were then incubated at 30°C or 37°C for an additional 24 hr before beginning

experiments.

Solutions for patch clamp recordings

HEK293T extracellular recording solution contained the following (in mM): 140 Tris, 20

CaCl$_2$, 1 MgCl$_2$, pH 7.3 with methansulfonic acid, osmolarity 309 mOsm/kg. HEK293T internal

recording solution contained the following: 140 NMDG, 10 HEPES, 2 MgCl$_2$, 2 Mg-ATP, 5 EGTA,

pH 7.3 with methansulfonic acid, osmolarity 358 mOsm/kg.

For recordings of I_{Ca} in mouse retina, extracellular recording solution contained the

following (in mM): 115 NaCl, 2.5 KCl, 22.5 NaHCO$_3$, 1.25 NaH$_2$PO$_4$, 2 CaCl$_2$, 1 MgCl$_2$, 5 HEPES,

5 CsCl, 5.5 Glucose, osmolarity 290 mOsm/kg. Mouse cone intracellular solution contained the

following (in mM): 105 CsMeSO$_4$, 20 TEA-Cl, 1 MgCl$_2$, 11 HEPES, 10 EGTA, 4 Mg-ATP, 10

phosphocreatine, 0.3 Na-GTP, pH 7.4 with CsOH, osmolarity 300 mOsm/kg.

For recordings of I_{Ca} in ground squirrel retina, extracellular recording solution contained

the following (in mM): 10 HEPES, 85 NaCl, 3.1 KCl, 2.48 MgSO$_4$, 6 Glucose, 1 Na-succinate, 1

Na-malate, 1 Na-lactate, 1 Na-pyruvate, 2 CaCl$_2$, 25 NaHCO$_3$, and 20 TEA-Cl, osmolarity 285 ±

5 mOsm/kg. Intracellular solution contained the following (in mM): 80 CsCl, 10 BAPTA, 2 MgSO$_4$,

10 HEPES, 20 TEA-Cl, 5 Mg-ATP, and 0.5 Na-GTP, pH 7.35 with CsOH, osmolarity 285 ± 5

mOsm/kg. For recordings of I_{Aglu}**,** extracellular recording solution contained (in mM): 125 NaCl,

3.0 KCl, 1.25 NaH$_2$PO$_4$, 25 NaHCO$_3$, 2 CaCl$_2$, 1 MgCl$_2$, 3 dextrose, 3 sodium pyruvate, 0.1

picrotoxin and 0.02 DNQX, and in indicated experiments 0.0013 TFB-TBOA. Intracellular I$_{Aglu}$

contained (in mM): 125 KSCN, 10 TEA-Cl, 10 HEPES, 1 $CaCl_2$, 2 $MgCl_2$, 0.3 Na-GTP, 4 Mg-

ATP, 10 K_2 phosphocreatine, 0.02 ZD7288. Mouse and ground squirrel extracellular slice

recording solutions were equilibrated with 5% CO_2 / 95% O_2 to a pH of ~7.5.

For recordings of light responses in horizontal cells of mouse retina, extracellular

recording solution consisted of Ames' media supplemented with 100 U/mL penicillin, 0.1 mg/mL

streptomycin and 22.6 mM $NaHCO_3$, osmolarity 280 ± 5 mOsm/kg. The intracellular recording

solution contained the following (in mM): 135 K-Aspartate, 10 KCl, 10 HEPES, 5 EDTA, 0.5

$CaCl_2$, 1 Mg-ATP, 0.2 Na-GTP, pH 7.35 with KOH, osmolarity 305 ± 5 mOsm/kg.

Table 4. Pharmacological drugs and their concentrations used in electrophysiological
recordings of cones.
535

Compound	Concentration (µM) for mouse recordings	Concentration (µM) for ground squirrel recordings
Cadmium	200	N/A
DL-TBOA	N/A	375
DNQX	20	N/A
Isradipine	1	2
ML218	5	5
Nickel	100	N/A
Picrotoxin	N/A	50
Strychnine	N/A	10
ZD7288	N/A	50
Glutamate	1000	N/A

Patch clamp electrophysiology

Whole-cell voltage clamp recordings of transfected HEK293T cells were performed 48 to

72 hrs after transfection using an EPC-10 amplifier and Patchmaster software (HEKA Elektronik,

Lambrecht, Germany). Patch pipette electrodes with a tip resistance between 4 and 6 MΩ were

pulled from thin-walled borosilicate glass capillaries (World Precision Instruments, Sarasota, FL)

using a P-97 Flaming/Brown Puller (Sutter Instruments, Novato, CA). A reference Ag/AgCl wire

was placed into the culture dish mounted on an inverted Olympus IX70 microscope. Recordings

were performed at room temperature. A pressurized perfusion pencil multi-barrel manifold

controlled with Valve Bank II (AutoMate Scientific, Inc., Berkeley, CA) was used to deliver

extracellular solutions. ML218 of different concentrations (0.5, 1, 5, 25 and 100 mM) was added

to the extracellular solution the day of experiments. Series resistance was compensated up to

70%, and passive membrane leak subtraction was conducted using a P/−4 protocol. Whole-cell

Ca^{2+} currents (I_{Ca}) of transfected HEK293T cells were evoked for 50 ms with incremental +5 mV

steps from -80 mV to +65 mV. Current-Voltage (IV) data were fit with a single Boltzmann equation:

$I_{Ca} = G_{max}(V_m\text{-}V_r)/(1\text{+}exp[\text{-}(V_m\text{-}V_h))/k])$, where G_{max} is the maximal conductance, V_m is the test

voltage, V_r is the Ca^{2+} reversal potential, V_h is the membrane potential required to activate 50%

of G_{max}, and k is the slope factor. Data were sampled at 100 kHz, filtered at 3 kHz, and analyzed

using custom programs written in IgroPro (WaveMetrics).

Whole-cell voltage clamp recordings of mouse cones, macaque cone terminals, and

mouse horizontal cells were performed using an EPC-10 amplifier and Patchmaster software

(HEKA). Patch pipette electrodes with a tip resistance between 10 and 14 MΩ for cones and 6-8

for horizontal cells were pulled from thick-walled borosilicate glass (1.5 mm outer diameter; 0.84

mm inner diameter; World Precision Instruments).

To prepare retinal slices, adult mice (6 – 8 weeks old) were anesthetized using isoflurane

and euthanized by cervical dislocation. Eyes were enucleated, placed into cold Ames' media

slicing solution, and hemisected. Following removal of the vitreous, the eye cup was separated

into dorsal and ventral halves using a scalpel. Ventral retina was isolated, molded into low-melt

agarose and mounted in a Leica VT1200s vibratome (Leica Biosystems). Mouse retina slicing

solution was continuously bubbled with 100% O_2 and contained the following: Ames' Medium with

L-glutamine supplemented with (in mM) 15 NaCl, 10 HEPES, 10 U/mL penicillin, 0.1 mg/mL

streptomycin, pH 7.4, osmolarity 300 mOsm/kg. Vertical (~200 μm) or horizontal (~160 μm) retinal

slices were anchored in a recording chamber, placed onto a fixed stage, and positioned under an

upright Olympus BX51WI microscope equipped with a 60X water-immersion objective (1.0 NA),

and superfused with extracellular solution (flow rate of ~1-2 ml/min) at room temperature. Slices

were visualized using IR-DIC optics and an IR-2000 (Dage MTI, Michigan City, IN) or SciCam Pro

(Scientifica, Uckfield, United Kingdom) CCD camera controlled by the IR-capture software

package or µManager, respectively [53, 54]. Drugs used in these experiments were added to the

mouse extracellular solution the day of experiments at the concentration described in Table 4. A

reference Ag/AgCl pellet electrode was placed directly into the recording chamber solution. Data

from whole-cell recordings with a series resistance >20 MΩ were discarded.

Cone somas were identified based on their morphology and location (outer ONL cell

layer). Cone identity was confirmed by the whole-cell capacitance (~3-4 pF), which is larger than

rod whole-cell capacitance (~0.7-1 pF). For cone voltage ramp recordings, cones were held at -

90 mV for 200 ms followed by a ramp of +0.5 mV/ms to +40 mV. To determine voltage activation

of I_{Ca}, whole-cell Ca^{2+} currents in cones were evoked for 50 ms with incremental +5 mV steps

from -80 mV to +40 mV. The activation voltage of I_{Ca} is reported as G/G_{max}, where G is the

conductance at each test voltage and G_{max} is the maximum peak conductance for each cone.

Conductance was calculated using the equation $I_{Ca} = G(V_m - V_r)$, where V_r is +60 mV. To determine

steady state inactivation of I_{Ca}, currents in cones were evoked for 500 ms with incremental +5 mV

steps from -90 mV and -30 mV followed by a final step to -30 mV for 50 ms after each test voltage.

The steady state inactivation of I_{Ca} is reported as I/I_{max}, where I is the peak current in the final

voltage step to -30 mV and I_{max} is the maximum peak current for each cone. Data were sampled

between 20 and 60 kHz and filtered at 3 kHz.

For horizontal cell light responses, horizontal slices were prepared from central mouse

retina. The identity of horizontal cells was determined based on their larger soma diameter (~15

µm) compared to bipolar cell somas (~6 µm). During whole cell patch clamp recordings, horizontal

cells were held at -70 mV. Light stimuli (1 s) at 410 nm (630 x 830 µm) were presented onto the

retina (at a minimum of 5 s intervals) through the microscope's condenser using a Polygon1000

DMD pattern illuminator (Mightex, Pleasanton, CA, USA) and a custom-built light path. Light

intensity (in watts) was measured at the point on the microscope stage where the retina is placed

using a power meter (Thorlabs, Newton, New Jersey, USA). Photon flux Φ_q (photons/s) within the

light stimulus area was calculated using the measured light intensities in the formula:

$$\Phi_q \text{ per } \mu m^2 = \frac{measured\ light\ intensity\ (W)}{\frac{hc}{\lambda}}/light\ stimulus\ area\ (\mu m^2)$$

where h is Planck's constant (J*s), c is the speed of light (m/s), and λ is the wavelength (m). Light

stimulus intensity was increased in Log2 steps from 4.9×10^2 to 2.1×10^5 Φ_q/µm². For each light

intensity step, both ON and OFF current amplitudes were measured from baseline (averaged 5

ms of current prior to light onset) to the maximum positive (after light onset) or maximum negative

(after light offset) current, respectively. To confirm the identity of these horizontal cell light

responses as AMPA-mediated currents, 1 s light stimuli (410 nm, 1.2×10^5 Φ_q/µm²) were

continuously delivered every 10 s before, during and after bath application of 20 µM DNQX.

For voltage clamp recordings of 13-lined ground squirrel cones, retinal slices were

prepared as previously described [55]. The eyecup was divided along the dorsal to ventral axis into

superior, middle, and inferior parts. The dorsal area above the line of the optic nerve head was

defined as superior, the central region with a width of about 5 mm just ventral to the optic nerve

head was middle, and the remaining ventral area was inferior. Isradipine and ML218, alone or in

combination, were applied from separate puffer pipettes whose orifices were aimed at the cone

synaptic region. Recordings were made with an Axopatch 200B amplifier (Molecular Devices).

Signals were electronically filtered at 5 kHz and digitized at a rate of 10 kHz. Additional Gaussian

filtering was added (cutoff frequency of 500 Hz). Tissue was viewed through a 63x water

immersion objective on a Zeiss Axioskop FS2 microscope and superfused with extracellular

solution at room temperature. Drugs and their concentrations used during these experiments are

described in Table 4. Membrane potential was continuously maintained at -85 mV. During ramp

stimulation, the membrane potential was depolarized to -85 to +35 mV at a rate of 1 mV/ms.

For rhesus macaque cone terminal recordings, a 12-year-old male was sedated with

ketamine (5 mg/kg I.M.) and dexmedetomidine (0.015 mg/kg I.M.). Post-sedation, the animal

received buprenorphine (0.02 mg/kg I.M.), atropine (0.02 mg/kg I.M.), and maropitant citrate (1

mg/kg S.Q.). The animal was intubated and maintained with inhaled isoflurane (0.75-2.0%) and

propofol (7-8 mg/kg/hr I.V.). Crystalloid fluids (5 mL/kg/hr) were administered I.V., and

phenylephrine (5-10 mcg/kg/hr I.V.) was used for blood pressure support. The animal was under

anesthesia for approximately 3 hours prior to perfusion. Transcardial perfusion was approached

through a midline thoracotomy. Immediately prior to perfusion, 5 mL of Euthasol® (sodium

pentobarbital 390 mg/mL/phenytoin sodium 50 mg/mL) was administered I.V. The descending

thoracic aorta was clamped, the pericardium was opened, and the right atrium was cut. The apex

of the left ventricle was sharply incised, and a large bore cannula (Yankauer suction handle, 5

mm internal diameter) was inserted through the left ventricle until it could be palpated in the

ascending aorta. The cannula was clamped in place at the apex of the left ventricle. The animal

was perfused with 4000 mL of cold phosphate-buffered saline at ~500 mL/min. Eyes were

removed approximately 1 hour following perfusion. Eyes were dissected, and eye cups were

allowed to dark adapt for ~30 min and stored in bicarbonate buffered Ames' media at 32°C

equilibrated with 95% O_2/5% CO_2 prior to slice preparations. Vertical sections of central retina

were prepared from 5 mm retina punches as described for mice.

639   Serial block-face scanning electron microscopy and 3D reconstructions

Eye cups were prepared from P42 WT and G369i KI littermates and fixed using 4%

glutaraldehyde in 0.1M cacodylate buffer, pH 7.4, for 4 hours at room temperature followed by

additional fixation overnight at 4°C. Glutaraldehyde-fixed eye cups were then washed 3 times in

0.1M cacodylate buffer. Retinas were thereafter isolated and embedded in Durcupan resin after

staining, dehydration, and embedding as described previously (Della Santina et al., 2016). A

Thermo Scientific VolumeScope serial block face scanning electron micrscope was used to image

embedded retinas. Retinal regions comprising a 2x2 montage of 40.96 μm tiles were imaged at a

resolution of 5nm/pixel and section thickness of 50 nm. Image stacks were aligned, and cone

photoreceptor terminals reconstructed using TrakEM2 (NIH). Postsynaptic partners at cone

ribbons were followed to the inner nuclear layer to determine their identity. Amira software was

used for 3D visualization of reconstructed profiles.

Electroretinography

Retinal function was assessed using the Celeris system (Diagnosys, Inc.) paired with the

Espion software (Diagnosys, Inc.). Mice were anaesthetized under red light (660 nm) via

intraperitoneal (I.P.) injection with ketamine/xylazine mixture (100 mg/kg ketamine, 10 mg/kg

xylazine). Tropicamide ophthalmic solution (1 %) and hypromellose lubricant eye gel (0.3 %) were

administered topically to both eyes before the mouse was secured to a heated (37 °C) platform

to maintain body temperature. Ag/AgCl corneal stimulators were placed on each eye. After

collecting data from individual mice, atipamezole (Antisedan, 1-2 mg/kg) was I.P. injection

administered to reverse the effects of the ketamine and xylazine. For photopic ERGs, eyes were

light adapted using a background green light at 20 cd·s/m^2 for 10 min. Following light adaptation,

eight different light intensity pulses (-0.5, 0, 0.5, 1, 1.5, 2, 2.3 and 2.6 log cd·s/m^2) were delivered

on top of the background green light. ERG a-waves were measured from baseline to the peak of

the negative potential. ERG b-waves were measured from the peak of the a-wave to the peak of

the positive potential. For flickering ERGs, mice were dark adapted (>12 hrs). White light pulses

from -4 to 2.5 log cd·s/m^2 were delivered in 0.5 log unit steps at 10 Hz. Response amplitudes

were measured from the trough to the peak of each response at all light intensities. Each intensity

stimulus was delivered 10 times, with a 3 s interval between each stimulus, and averaged. The

mean response amplitudes recorded in the right and left eye of each mouse are reported for all

quantified ERG data.

Visible platform swim test

The visible platform swim test was performed as has been previously described [30]. WT,

G369i KI, and $Ca_v1.4$ KO male and female mice (6- 9 weeks old) were adapted to the procedure

room for at least 1 h prior to beginning the experiments. A water-filled, 4-foot diameter galvanized

steel tank and a visible white platform with a diameter of 10 cm was used for the swim test. Mice

were subjected to 6 swim trials per day. Assays conducted under photopic (55 lux) and scotopic

conditions (0 lux) were performed on days 1 and 2, respectively. Light intensity for photopic and

scotopic conditions was measured at the platform using an Extech HD450 light meter (FLIR

Systems, Nashua, New Hampshire). Mice were given 90 s to find the platform before being

removed. After one trial was performed on all mice, the platform was moved to one of three

different locations, top left, top center, and top right in relation to the initial site of mouse placement

in tank. Each trial was recorded using an infrared camera (Basler AG, Ahrensburg, Germany) and

EthoVision XT16 software (Noldus Information Technology). Locating the platform was

considered successful when mice contacted the platform with a head-on approach, even if mice

failed to escape onto the platform. Mice were allowed to rest on the platform for 15 s at the end

of each trial before being returned to a pre-warmed cage. Latency to find the platform was

manually recorded and confirmed using recorded videos. Male and female mice were tested

separately, and no sex differences in performance were identified post-hoc using two-way RM-

ANOVA (Supplementary Table 1). Male and female data were combined into a single group for

each genotype. Latency to platform was calculated by averaging the final 3 trials under scotopic

and photopic conditions for each mouse and compared using Kruskal-Wallis one-way ANOVA

with Dunn's posthoc multiple comparison test.

Data analysis

Electrophysiological data were analyzed by custom routines written in IgorPro software (Wavemetrics) and statistical analysis was performed using Prism software (GraphPad). Data were analyzed for normality by Shapiro Wilk test followed by parametric (t-test) or non-parametric methods (Kruskal Wallis or Mann-Whitney).

References

1.   Snellman, J*., et al.* Acute destruction of the synaptic ribbon reveals a role for the ribbon in vesicle priming. *Nat Neurosci* **14**, 1135-1141 (2011).

2.   Dembla, E., Dembla, M., Maxeiner, S. & Schmitz, F. Synaptic ribbons foster active zone stability and illumination-dependent active zone enrichment of RIM2 and Cav1.4 in photoreceptor synapses. *Sci Rep* **10**, 5957 (2020).

3.   Haverkamp, S., Grunert, U. & Wassle, H. The cone pedicle, a complex synapse in the retina. *Neuron* **27**, 85-95 (2000).

4.   Mercer, A.J. & Thoreson, W.B. The dynamic architecture of photoreceptor ribbon synapses: cytoskeletal, extracellular matrix, and intramembrane proteins. *Vis Neurosci* **28**, 453-471 (2011).

5.   Frederick, C.E. & Zenisek, D. Ribbon Synapses and Retinal Disease: Review. *Int J Mol Sci* **24** (2023).

6.   Mansergh, F*., et al.* Mutation of the calcium channel gene Cacna1f disrupts calcium

signaling, synaptic transmission and cellular organization in mouse retina. *Hum Mol Genet* **14**,

3035-3046 (2005).

7.   Liu, X*., et al.* Dysregulation of $Ca_v1.4$ channels disrupts the maturation of photoreceptor

synaptic ribbons in congenital stationary night blindness type 2. *Channels* **7**, 514-523 (2013).

8.   Chang, B*., et al.* The nob2 mouse, a null mutation in Cacna1f: anatomical and functional

abnormalities in the outer retina and their consequences on ganglion cell visual responses. *Vis*

Neurosci **23**, 11-24 (2006).

9.   Singh, A*., et al.* C-terminal modulator controls Ca^{2+}-dependent gating of $Ca_v1.4$ L-type

Ca^{2+} channels. *Nat Neurosci* **9**, 1108-1116 (2006).

10.   Wahl-Schott, C*., et al.* Switching off calcium-dependent inactivation in L-type calcium

channels by an autoinhibitory domain. *Proc. Natl. Acad. Sci. U. S. A.* **103**, 15657-15662 (2006).

11.   Bech-Hansen, N.T.*, et al.* Loss-of-function mutations in a calcium-channel a_1-subunit gene

in Xp11.23 cause incomplete X-linked congenital stationary night blindness. *Nature Genet.* **19**,

264-267 (1998).

12.   Strom, T.M*., et al.* An L-type calcium-channel gene mutated in incomplete X-linked

congenital stationary night blindness. *Nature Genet.* **19**, 260-263 (1998).

13.   Waldner, D.M., Bech-Hansen, N.T. & Stell, W.K. Channeling Vision: CaV1.4-A Critical Link

in Retinal Signal Transmission. *Biomed Res Int* **2018**, 7272630 (2018).

14.   Boycott, K.M., Pearce, W.G. & Bech-Hansen, N.T. Clinical variability among patients with

incomplete X-linked congenital stationary night blindness and a founder mutation in CACNA1F.

Can J Ophthalmol **35**, 204-213 (2000).

15.   Zeitz, C., Robson, A.G. & Audo, I. Congenital stationary night blindness: an analysis and

update of genotype-phenotype correlations and pathogenic mechanisms. *Progress in retinal and*

eye research **45**, 58-110 (2015).

16.   Regus-Leidig, H*., et al.* Photoreceptor degeneration in two mouse models for congenital

stationary night blindness type 2. *PLoS One* **9**, e86769 (2014).

17.   Zabouri, N. & Haverkamp, S. Calcium channel-dependent molecular maturation of

photoreceptor synapses. *PLoS One* **8**, e63853 (2013).

18.   Raven, M.A*., et al.* Early afferent signaling in the outer plexiform layer regulates

development of horizontal cell morphology. *J Comp Neurol* **506**, 745-758 (2008).

19.   Maddox, J.W*., et al.* A dual role for Cav1.4 Ca(2+) channels in the molecular and structural

organization of the rod photoreceptor synapse. *Elife* **9**, e62184 (2020).

20.   Perez-Reyes, E. Molecular physiology of low-voltage-activated t-type calcium channels.

Physiol Rev **83**, 117-161 (2003).

21.   Davison, A., Lux, U.T., Brandstatter, J.H. & Babai, N. T-Type Ca(2+) Channels Boost

Neurotransmission in Mammalian Cone Photoreceptors. *J. Neurosci.* **42**, 6325-6343 (2022).

22. Koschak, A*., et al.* Cav1.4alpha1 subunits can form slowly inactivating dihydropyridine-

sensitive L-type Ca2+ channels lacking Ca2+-dependent inactivation. *J. Neurosci.* **23**, 6041-6049

(2003).

23. Schmitz, F., Konigstorfer, A. & Sudhof, T.C. RIBEYE, a component of synaptic ribbons: a

protein's journey through evolution provides insight into synaptic ribbon function. *Neuron* **28**, 857-

872 (2000).

24. Martemyanov, K.A. & Sampath, A.P. The Transduction Cascade in Retinal ON-Bipolar

Cells: Signal Processing and Disease. *Annu Rev Vis Sci* **3**, 25-51 (2017).

25. Hack, I., Frech, M., Dick, O., Peichl, L. & Brandstatter, J.H. Heterogeneous distribution of

AMPA glutamate receptor subunits at the photoreceptor synapses of rodent retina. *Eur J Neurosci*

13, 15-24 (2001).

26. Brandstatter, J.H., Koulen, P. & Wassle, H. Selective synaptic distribution of kainate

receptor subunits in the two plexiform layers of the rat retina. *J. Neurosci.* **17**, 9298-9307 (1997).

27. Feigenspan, A. & Babai, N. Functional properties of spontaneous excitatory currents and

encoding of light/dark transitions in horizontal cells of the mouse retina. *Eur J Neurosci* **42**, 2615-

2632 (2015).

28. Nusinowitz, S., Ridder, W.H., 3rd & Ramirez, J. Temporal response properties of the

primary and secondary rod-signaling pathways in normal and Gnat2 mutant mice. *Exp Eye Res*

84, 1104-1114 (2007).

29.   Vinberg, F., Wang, T., Molday, R.S., Chen, J. & Kefalov, V.J. A new mouse model for

stationary night blindness with mutant Slc24a1 explains the pathophysiology of the associated

human disease. *Hum Mol Genet* **24**, 5915-5929 (2015).

30.   Bimonte-Nelson, H., Daniel, J. & Koebele, S. *The maze book: theories, practice, and*

protocols for testing rodent cognition (Springer, 2015).

31.   Beier, C*., et al.* Deafferented Adult Rod Bipolar Cells Create New Synapses with

Photoreceptors to Restore Vision. *J. Neurosci.* **37**, 4635-4644 (2017).

32.   Shen, N., Wang, B., Soto, F. & Kerschensteiner, D. Homeostatic Plasticity Shapes the

Retinal Response to Photoreceptor Degeneration. *Curr Biol* **30**, 1916-1926 e1913 (2020).

33.   Leinonen, H*., et al.* Homeostatic plasticity in the retina is associated with maintenance of

night vision during retinal degenerative disease. *Elife* **9** (2020).

34.   Care, R.A*., et al.* Mature Retina Compensates Functionally for Partial Loss of Rod

Photoreceptors. *Cell rep* **31**, 107730 (2020).

35.   Magupalli, V.G*., et al.* Multiple RIBEYE-RIBEYE interactions create a dynamic scaffold for

the formation of synaptic ribbons. *J. Neurosci.* **28**, 7954-7967 (2008).

36.   Ingram, N.T., Sampath, A.P. & Fain, G.L. Voltage-clamp recordings of light responses

from wild-type and mutant mouse cone photoreceptors. *J. Gen. Physiol.* **151**, 1287-1299 (2019).

37.   Spiwoks-Becker, I., Glas, M., Lasarzik, I. & Vollrath, L. Mouse photoreceptor synaptic

ribbons lose and regain material in response to illumination changes. *Eur J Neurosci* **19**, 1559-

1571 (2004).

38.   Regus-Leidig, H., Specht, D., Tom Dieck, S. & Brandstatter, J.H. Stability of active zone

components at the photoreceptor ribbon complex. *Mol Vis* **16**, 2690-2700 (2010).

39.   Sheets, L., Kindt, K.S. & Nicolson, T. Presynaptic CaV1.3 channels regulate synaptic

ribbon size and are required for synaptic maintenance in sensory hair cells. *J. Neurosci.* **32**,

17273-17286 (2012).

40.   Wong, H.C.*, et al.* Synaptic mitochondria regulate hair-cell synapse size and function. *Elife*

8 (2019).

41.   Goel, P.*, et al.* Homeostatic scaling of active zone scaffolds maintains global synaptic

strength. *J Cell Biol* **218**, 1706-1724 (2019).

42.   Hong, H.*, et al.* Structural Remodeling of Active Zones Is Associated with Synaptic

Homeostasis. *J. Neurosci.* **40**, 2817-2827 (2020).

43.   Williams, B., Maddox, J.W. & Lee, A. Calcium Channels in Retinal Function and Disease.

Annu Rev Vis Sci (2022).

44.   Macosko, E.Z.*, et al.* Highly Parallel Genome-wide Expression Profiling of Individual Cells

Using Nanoliter Droplets. *Cell* **161**, 1202-1214 (2015).

45.   Tringham, E*., et al.* T-type calcium channel blockers that attenuate thalamic burst firing

and suppress absence seizures. *Sci Transl Med* **4**, 121ra119 (2012).

46.   Perez-Reyes, E., Van Deusen, A.L. & Vitko, I. Molecular pharmacology of human Cav3.2

T-type Ca2+ channels: block by antihypertensives, antiarrhythmics, and their analogs. *The*

Journal of pharmacology and experimental therapeutics **328**, 621-627 (2009).

47.   Choi, S.Y., Jackman, S., Thoreson, W.B. & Kramer, R.H. Light regulation of Ca2+ in the

cone photoreceptor synaptic terminal. *Vis Neurosci* **25**, 693-700 (2008).

48.   Bijveld, M.M*., et al.* Assessment of night vision problems in patients with congenital

stationary night blindness. *PLoS One* **8**, e62927 (2013).

49.   Peloquin, J.B., Rehak, R., Doering, C.J. & McRory, J.E. Functional analysis of congenital

stationary night blindness type-2 CACNA1F mutations F742C, G1007R, and R1049W.

Neuroscience **150**, 335-345 (2007).

50.   Hoda, J.C., Zaghetto, F., Koschak, A. & Striessnig, J. Congenital stationary night

blindness type 2 mutations S229P, G369D, L1068P, and W1440X alter channel gating or

functional expression of Cav1.4 L-type Ca2+ channels. *J. Neurosci.* **25**, 252-259 (2005).

51.   Kerov, V*., et al.* alpha2delta-4 Is Required for the Molecular and Structural Organization

of Rod and Cone Photoreceptor Synapses. *J. Neurosci.* **38**, 6145-6160 (2018).

52.   Lee, A*., et al.* Characterization of Cav1.4 complexes ($\alpha_1$1.4, β_2, and $\alpha_2\delta-4$) in HEK293T

cells and in the retina. *J. Biol. Chem.* **290**, 1505-1521 (2015).

53.   Edelstein, A., Amodaj, N., Hoover, K., Vale, R. & Stuurman, N. Computer control of

microscopes using microManager. *Curr Protoc Mol Biol* **Chapter 14**, Unit14 20 (2010).

54.   Edelstein, A.D.*, et al.* Advanced methods of microscope control using muManager

software. *J Biol Methods* **1** (2014).

55.   DeVries, S.H. & Schwartz, E.A. Kainate receptors mediate synaptic transmission between

cones and 'Off' bipolar cells in a mammalian retina. *Nature* **397**, 157-160 (1999).

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Supplementary Figure S1. Effect of ML218 on HEK293T cells transfected with Ca$_v$1.4, β$_{2x13}$, and α$_2$δ−4 or Ca$_v$3.2. a,b, Traces (*a*) and I-V plot (*b*) for I_{Ca} evoked by 50 ms depolarizations from -90 mv to various voltages without (0 µM) and in the presence of the indicated concentrations of ML218. **c,d**, Traces (*c*) and graph (*d*) depicting inhibitory effects of ML218 (5 µM) on Ca$_v$3.2-mediated ICa evoked by voltage ramp in individual transfected cells. **, p < 0.01 by paired t-test.

Supplementary Figure S2. Adult ground squirrel cones lack a Ca$_v$3-like conductance. a, Representative traces for I_{Ca} evoked by voltage steps from a continuous holding voltage of -85 mV to various voltages. Following recording of I_{Ca} bathed in control solution (baseline), cone terminals were perfused sequentially with ISR (2 µM) and then ISR (2 µM) + ML218 (5 µM). Voltage protocol is shown below current traces. Dashed line indicates zero current. **b**, Traces show the I_{Ca} sensitive to ISR (top) and ISR+ML218 (middle). I_{Ca} was evoked by steps from -85 mV to voltages between -65 and -5 mV in increments of +10 mV. The I_{Ca} recorded in ISR was subtracted from the baseline I_{Ca} (top). I_{Ca} recorded in ISR+ML218 was subtracted from I_{Ca} recorded in ISR (middle). The small amplitude and noninactivating properties of the current blocked by adding ML218 to ISR suggest that it is mediated by Cav1.4 channels that continue to undergo a time-dependent block at the concentration of ISR used here. **c**, Top, Current-voltage (I-V) plots of peak and steady-state I_{Ca} from *b*. Bottom, normalized conductance (G/G$_{max}$) vs. voltage relationship (G-V) of the I_{Ca} blocked by ISR and ISR + ML218. Smooth lines represent Boltzmann fits; symbols and bars represent mean ± SEM, respectively; n = 4 cones. Due a negative shift in the activation properties caused by ML218 on a residual Ca$_v$1 current that is assumed to remain at the end of the experiment, the G-V curve of the I_{Ca} isolated through subtraction by applying ISR+ML218 underwent a statistically insignificant shift to the right. Data presented in *a-c* are from the same cone. **d**, Voltage ramp current response in an OFF cb3a bipolar cell before and after the application of ML218 (5 µM). V$_{1/2}$ was shifted to the right by 13.1 ± 4.3 mV (n = 3 cb3a cells, mean ± SD) consistent with the block of a Ca$_v$3-like conductance. In this experiment, the amplitude of an I_{Ca} component that had a more depolarized activation range was slightly increased. A similar voltage ramp applied to OFF cb3b bipolar cells produced only a slight leftward shift in an I_{Ca} that had a more depolarized activation range, consistent with the exclusive expression a Ca$_v$1-type current (V$_{1/2}$ was shifted by -1.9 ± 0.4 mV, n = 3 cb3b cells; data not shown). **e**, Neurobiotin filled OFF cb3a. Following recordings of cells in *d*, retina slices were fixed and co-immunolabeled for bassoon and choline acetyltransferase (CHAT) to depict synapses and the OFF and ON sublamina of the innerplexiform layer (IPL), respectively. Streptavidin was used to label the neurobiotin to depict the OFF cb3a axon stratification and morphology.

Supplementary Figure S3. Characterization of I_Ca in cones of macaque retina. (a-c) Representative current traces (a) and I-V (b), and G-V (b) for I_Ca evoked by 200-ms steps from a holding voltage (V_h) of -90 mV or -50 mV (which should isolate the Ca_v1-mediated I_Ca from any contribution of Ca_v3 channels). n=3 cells. (d) $V_{1/2}$ and slope factor (k) obtained from Boltzmann fit of data in *c*. There was no significant effect of holding voltage on these parameters, arguing against a contribution of Ca_v3 channels to the whole-cell I_Ca. (e) Lack of effect of inactivating voltage on *I_Ca*. Test currents were evoked from a holding voltage of -100 V to -30 mV before (P1) and after (P2) a 500 ms step to -60 mV. Voltage protocol shown above overlay of P1 and P2 current traces. Boxed region is shown with expanded timescale; the similar activation kinetics of P1 and P2 currents supports the contribution of a single Ca_v subtype. Graph at right shows lack of significant difference in amplitudes of P1 and P2 currents, arguing against the contribution of Ca_v3 channels to the P1 current.

Supplementary Table 1

Male vs. Female WT (Photopic)	SS	DF	MS	F (DFn, DFd)	P value
Trial x Sex	1125	5	224.9	F (5, 45) = 0.6060	P=0.6956
Trial	3564	5	712.9	F (5, 45) = 1.921	P=0.1096
Sex	273.6	1	273.6	F (1, 9) = 0.6616	P=0.4370
Subject	3722	9	413.5	F (9, 45) = 1.114	P=0.3727
Residual	16703	45	371.2		
Male vs. Female G369i KI (Photopic)	SS	DF	MS	F (DFn, DFd)	P value
Trial x Sex	1457	5	291.4	F (5, 45) = 0.5362	P=0.7477
Trial	3516	5	703.2	F (5, 45) = 1.294	P=0.2833
Sex	1150	1	1150	F (1, 9) = 0.8168	P=0.3897
Subject	12675	9	1408	F (9, 45) = 2.592	P=0.0168
Residual	24451	45	543.4		
Male vs. Female Cav1.4 KO (Photopic)	SS	DF	MS	F (DFn, DFd)	P value
Trial x Sex	1736	5	347.1	F (5, 35) = 0.6597	P=0.6563
Trial	993	5	198.6	F (3.056, 21.39) = 0.3775	P=0.7737
Sex	78.37	1	78.37	F (1, 7) = 0.03788	P=0.8512
Subject	14481	7	2069	F (7, 35) = 3.932	P=0.0029
Residual	18416	35	526.2		
Male vs. Female WT (Scotopic)	SS	DF	MS	F (DFn, DFd)	P value
Trial x Sex	862	5	172.4	F (5, 45) = 0.6907	P=0.6331
Trial	914.5	5	182.9	F (5, 45) = 0.7328	P=0.6027
Sex	634.7	1	634.7	F (1, 9) = 2.121	P=0.1793
Subject	2693	9	299.3	F (9, 45) = 1.199	P=0.3193
Residual	11232	45	249.6		
Male vs. Female G369i KI (Scotopic)	SS	DF	MS	F (DFn, DFd)	P value
Trial x Sex	7648	5	1530	F (5, 45) = 2.581	P=0.0390
Trial	1644	5	328.8	F (5, 45) = 0.5547	P=0.7339
Sex	4712	1	4712	F (1, 9) = 4.716	P=0.0580
Subject	8993	9	999.2	F (9, 45) = 1.686	P=0.1206
Residual	26671	45	592.7		
Male vs. Female Cav1.4 KO (Scotopic)	SS	DF	MS	F (DFn, DFd)	P value
Trial x Sex	1727	5	345.4	F (5, 35) = 1.470	P=0.2246
Trial	12215	5	2443	F (2.802, 19.62) = 10.40	P=0.0003
Sex	96.33	1	96.33	F (1, 7) = 0.1346	P=0.7246
Subject	5011	7	715.9	F (7, 35) = 3.046	P=0.0131
Residual	8225	35	235		