Early life stress induces sex-specific changes in behavior and parallel locus coeruleus

neuron excitability

Savannah Brannan[1], Ben D Richardson[1]

[1]Department of Pharmacology, Southern Illinois University – School of Medicine, Springfield, IL

62702

Corresponding Author: Ben D Richardson, PhD: brichardson29@siumed.edu

Abstract

Many psychiatric disorders are associated with specific risk factors, including biological

sex, chronic stress, and adversity in childhood, but mechanisms underlying these relationships

are unknown. The locus coeruleus (LC) is a brain area that contains adrenergic norepinephrine

(NE)-releasing neurons with established sex differences in excitability and stress sensitivity. To

understand how adversity in early life affects cognitive/affective behavior and LC physiology, we

exposed C57BL/6J mice to a dual-phase (early development and adolescence) early life variable

stress (ELVS) paradigm and assessed behavior and LC physiology in early adulthood. ELVS

caused females, but not males, to display increased novel environment exploration and reduced

preference for sucrose. In addition, ELVS caused elevated activity in a familiar environment,

modest deficits in Y-maze performance, and altered attention in an operant task, regardless of

sex. A reduction in LC neuron excitability, partly due to an increase in action potential delay time,

was also found only in female mice exposed to ELVS, paralleling robust behavioral changes.

Pharmacological compensation of these changes in LC activity with reboxetine corrected some

ELVS-induced behavioral changes. CRF-induced changes in LC neuron activity were also

mediated by different preferential signaling pathways in male and female mice, a potential

mechanism for lasting sex-specific changes in LC physiology in response to ELVS. Through this

animal model of early life stress, we identified sex differences in behavior and parallel changes in

LC neuron excitability and CRF sensitivity, identifying mechanisms involved in determining how

stress and sex interact to cause LC activity dysregulation and related behavioral changes.

Introduction

Adverse childhood experiences (ACEs) have been shown to increase risk for many neurodevelopmental and affective disorders such as attention deficit hyperactivity disorder (ADHD), autism, anxiety and depression. Although mature neurological systems are often resistant to acute stress, prolonged periods of stress in more malleable and immature systems can alter neurodevelopment [1–5]. The more ACEs a child has, the greater the risk for adverse outcomes. Sex is a biological variable involved in ACEs, leading to adverse psychiatric outcomes, with females showing an increased risk for stress-related disorders in comparison to males [6]. Epidemiological studies indicate women may have increased stress-related disorder development risk/symptom exacerbation during life-phases characterized by hormonal changes that include puberty, menses, pregnancy, postpartum, and menopause [2, 3].

Increased ACE exposures leading to attention, memory, affective, and/or mood disorders represent deficits in behavioral domains is shaped by multiple circuits and modulatory neurotransmitters distributed throughout the central nervous system. For several reasons, we hypothesize that the noradrenergic locus coeruleus system (LC) may be a key neuromodulatory brain area responsible for ACE-related changes in behavior. Optimal activity and processing by multiple circuits to shape multiple behaviors depend upon optimal norepinephrine (NE) release from widely-projecting LC_{NE} neurons (e.g. prefrontal cortex, hippocampus, amygdala, etc. [7–15] to modulate attention [16, 17], cognitive flexibility [18, 19], learned behavior [9, 20], anxiety [10, 21] and reward [10, 14], among others [7]. Specifically, work in animal models indicates that LC-NE neuronal projections to the prefrontal cortex (PFC) and hippocampus are involved in maintaining attention, arousal, and aspects of memory formation [11–13, 22, 23] while those to the amygdala (basolateral and central) shape anxiety and emotional learning behaviors [10, 24, 25], and others modulate sensorimotor integration, associative fear learning, response inhibition, and working memory via projections to the cerebellum [26]. Norepinephrine (NE) release is highly dynamic, as appropriate tonic and phasic [16, 17, 27, 28] activity of LC-NE

neurons follow an inverted U-shaped function [7, 29, 30]. As such, low and high NE levels both

lead to negative outcomes and behavioral changes in cognitive and affective processes.

It is also established that corticotropin-releasing factor (CRF) directly alters the activity of

adrenergic neurons in the LC [31–34] and optimal LC neuron activity and related NE release in

the prefrontal cortex, amygdala, and hippocampus shape attention, anxiety, and memory [35–39].

Animal model data indicate that, compared to males, female LC neurons have different basal

physiological properties [40, 41] and elevated CRF sensitivity that can acutely increase basal NE

release from LC neurons [42–44]. These types of basal sex differences in CRF-activated signaling

may underpin females' increased stress-related disorder prevalence [45, 46]. Despite the

established role of NE in cognitive and affective behaviors and sex differences in LC and NE

dynamics, how early stress interacts with the sexually-dimorphic LC to shape behaviors

associated with LC function (i.e. affect, cognition, and locomotion) is poorly understood. To

identify and understand the mechanistic relationship between sex and affective, cognitive, and

locomotive behavioral changes associated with stress, we developed a two-phase model of stress

exposure early in life to determine how these behaviors are affected and if they are related to sex-

specific LC physiology changes. Our results suggest an interaction between sex and stress in

affective and cognitive behaviors that involve the LC in C57bl6/J (B6J) mice.

Materials and Methods

Animals

Male and female C57bl6/J (Jackson Labs, #000664) or Dbh-tdTomato mice (described

below) were purchased from Jackson Laboratory and/or bred in-house. To generate mice

expressing tdTomato in locus coeruleus noradrenergic neurons under control of dopamine β-

hydroxylase (Dbh-tdTomato mice), Dbh-Cre knock-in mice (B6.Cg-*Dbh*$^{tm3.2(cre)Pjen}$*/J*; stock

#033951) provided by Patrician Jensen, NIEHS [47] were crossed with homozygous Ai14

tdTomato reporter mice (B6.Cg-*Gt(ROSA)26Sor*$^{tm14(CAG-tdTomato)Hze}$/J, Stock #007914) provided by

Hongkui Zeng, Allen Institute for Brain Science [48] . All mice were group housed under a reversed

12/12-hour light-dark cycle (lights on from 21:00 to 9:00) with *ad libitum* access to food and water.

All manipulations and tests were performed during the animals' active dark phase (9:00 to 21:00),

unless otherwise stated. All procedures involving animals were conducted in accordance with

protocols approved by the Institutional Animal Care and Use Committee (#2022-100) of Southern

Illinois University - School of Medicine.

Early Life Variable Stress (ELVS) Animal Model

Male and female C57bl6/J mice were subject to either a two-phase (postnatal and

adolescent) early life variable stress (ELVS) paradigm or treated as controls with mice in a litter

either exposed to ELVS or treated as controls. The ELVS paradigm consisted of two phases of

manipulations, first from postnatal day 6-17 and the second from postnatal day 28-42. First, mice

in the ELVS group were isolated together from dams for 4hrs/day during postnatal day 6-10 (P6-

10) with access to a warming pad, and then weaned early at postnatal day 17 (P17). Pups were

supplemented with nutrient gel (Nutra-gel, Bio-Serv, Flemington, NJ, USA) from P17 to P21.

ELVS mice were exposed to sequential variable stressors from P28-42, with each stressor

conducted twice on their designated day starting with tape restraint (20 min), forced ice-cold swim

in a 40 cm x 10 cm filled container (20 min), 2,3,5-Trimethyl-3-thiazoline (TMT) synthetic fox odor

exposure (20 min), light cycle disruption by omission of one night of dark phase, and 48 hours of

social isolation. Control male and female C57bl6/J mice were not separated from dams and

weaned at P21 and were briefly handled 5 times per week for 10-15 minutes beginning at the

same time as the stressors (P28) for the ELVS group.

Behavioral Tests

Behavioral tests were done during the animals' active dark phase in a dimly lit room (~15-

20 lux) for testing. Mice were habituated to the testing room for 30 minutes prior to completion of

behavioral testing. Mouse performance on all behavioral assays, except for the rotarod, were

digitally recorded for analysis. Each apparatus was cleaned thoroughly with scent-free

disinfectant between animals.

Open Field Test

Mice were placed in a white acrylic chamber (40 cm x 40 cm wide x 30 cm high square

open top chamber) for 30 min with total distance, cumulative time in and number of entrances into

center (20 cm x 20 cm) area relative to edges (10 cm border) were analyzed using the EthoVision

XT 17 software (Noldus, Wageningen, the Netherlands).

Y-Maze

Mice were placed in a 3-armed "Y" shaped apparatus with symmetrical arms 35 cm in

length, 30 cm wall height and 10 cm lane widths at 120-degree angles from one another. Total

alternations, or the number of times the mouse enters 3 arms, were recorded and compared to

correct (sequential) alternations, where every arm entered within the 3 entrances had not been

entered in the immediate previous entrance.

Elevated Zero Maze

Mice were placed in the open arm of an elevated circular platform (50 cm diameter with 5

cm wide track) suspended 61 cm from the floor, with two opposing walled (20 cm high) segments

and two open segments in between. The amount of time spent in and entries into the open versus

closed segments were calculated using Ethovision XT 17.

Novel Object Recognition and Novel Object Placement Testing

The novel object recognition (NOR) assay was used to assess spatial memory. Following

exposure to the open field for 30 min as the chamber acclimation period, the NOR task was

performed in the same open field chamber with two phases, each with a 10-minute duration. The

first phase was the object habituation period, wherein two identical objects were placed

equidistant apart on opposite sides of the chamber. In the second phase, one familiar object from

the previous phase was preplaced with a novel object at the same location. Each phase was done

on separate days, with the novel object phase occurring 24 h after the same object phase. More

time spent with the novel object, was interpreted as increased recall of the familiar object and

related to memory and novelty seeking. The amount of time spent with the novel or moved object

versus the familiar object is calculated based on the time each mouse spent within 2 cm of the

object using Ethovision XT 17. Because this assay was performed in the same arena used as the

open field assay, the total distance covered during the second phase (novel object phase) of the

assay was also used to determine exploratory activity in a familiar environment to which the

mouse had been exposed two previous times.

Sucrose Preference Test

Two weeks before sucrose preference testing (SPT), mice were progressively water-

restricted, first to 4 hours of *ad libitum* water access per day for 2 days and then for 2 hours of

access per day for 2 days, followed by two days of 24 hr access. This was repeated twice before

limiting water access for 22 hrs prior to sucrose preference testing. For SPT, mice were placed in

a clean empty cage for one hour with 2 bottles, one bottle containing 1% sucrose solution and a

bottle containing water. The sucrose preference index was determined by comparing the amount

of sucrose solution versus water consumed by each individual mouse.

Rotarod

Motor and vestibular function was evaluated using the accelerating rotarod (3.17 cm

diameter, IITC life Sciences, Inc., Woodland Hills, CA, USA). This rod accelerated in rotations per

minute (RPM) from 4 10 40 RPM during the span of 5 minutes. Each mouse received 3 trial

sessions per day for two consecutive days for a total of 6 trials. A 10 min rest period was given to

the mouse between each trial. The time the mouse fell from the rotating rod onto the platform

below was recorded. If the mouse spent a reduced fall time latency, it indicated disrupted motor

or vestibular function.

Operant Five-Choice Serial Reaction Time Task (5CSRTT)

For 5SCRTT of attention and learning behaviors, mice were trained using standard

5CSRTT protocols (Asinof and Paine, 2014; Humby et al., 2005) and the apparatus was controlled

using ABET software (Lafayette Instruments). Two weeks before training, mice will be water

restricted starting at 4 hours of ad libitum access per day, reducing to 2 hours per day. During

training sessions, the mice were given sweetened condensed milk as a reward for correctly

performing the task and weighed to ensure they were at the proper weight after water restriction.

The operant conditioning chamber (30.5 x 24.1 x 29.2 cm) consisted of 2 Plexiglas sidewalls, an

aluminum front wall and back wall, and a stainless-steel grid floor will be used to carry out this

experiment (Lafayette Instruments). The front wall contained five nose poke apertures (2.5 x 2.2

x 2.2 cm each) with each aperture containing a light-emitting diode (LED) and an infrared sensor

cable to detect mouse nose insertions. The mice were trained by progressively shortening the

intertrial interval and stimulus duration for 30 minutes daily for 5-7 days per week to reach baseline

performance with intertrial interval (ITI) of 5sec and the stimulus duration of 0.8sec. Once mice

reached the criterion threshold of 80% correct trials with <30% of trials omitted, they were

assessed again on ten different variations with either shorter or longer ITI, shorter or longer

stimulus duration, aperture light brightness, sound and light distractor manipulations, and

combinations of these changes relative to the standard variables used during training. Response

accuracy (% correct) and response omissions (# of premature responses during ITI) for each

variable condition, along with the number of premature responses during specific training stages,

were evaluated for differences based on mouse sex and treatment as control or exposure to

ELVS.

Reboxetine Administration

In a separate cohort of 7-9-week-old control and ELVS mice, 2 mg/kg reboxetine diluted

into apple juice or apple juice alone (100 µl) was given to control and ELVS C57bl6/J mice one

hour prior to behavioral testing on SPT, open field, and y-maze. To determine reboxetine effects

on these behaviors, one behavior was evaluated each day on consecutive days with pre-test

apple juice/reboxetine administered each day 1 hr prior to each assay. The order of the behavioral

assays was kept the same for all mice.

Ex vivo Acute Brian Slice Preparation

Animals were anesthetized with 4% isoflurane, followed by intracardial perfusion with ice-

cold oxygenated N-Methyl-D-glucamine (NMDG) artificial cerebrospinal fluid (NMDG-ACSF),

which contained (in mM): 92 NMDG, 2.5 KCl, 0.5 $CaCl_2$, 10 $MgCl_2$, 1.2 NaH_2PO_4, 30 $NaHCO_3$, 20

HEPES, 25 D-glucose, 2 ascorbic acid, 2 thiourea, and 3 sodium pyruvate and had an osmolarity

of 300–310 mOsm with pH adjusted to 7.3–7.4 with HCl. Coronal slices (250 μm thick) that

included the LC were sectioned in ice-cold NMDG-ACSF using a Compresstome vibrating

microtome (Processionary Instruments, LLC) and were transferred to a holding chamber

containing NMDG-ACSF at 35 °C where the NaCl concentration increased steadily over 25 min.

After 30 min, slices were transferred to a modified HEPES-based ACSF (HEPES-ACSF, 35 °C),

which contained (in mM): 92 NaCl, 2.5 KCl, 2 $CaCl_2$, 2 $MgCl_2$, 1.2 NaH_2PO_4, 30 $NaHCO_3$, 20

HEPES, 25 D-glucose, 2 ascorbic acid, 2 thiourea, 3 sodium pyruvate, and 3 myo-inositol and

had an osmolarity of 300–310 mOsm with pH 7.3–7.4. After incubating slices in HEPES-ACSF at

35 °C for 1 hr, slices were maintained in HEPES-ACSF at room temperature until being

transferred to the recording chamber where they were continuously perfused at a 3-5 ml/min with

oxygenated ACSF, which contained (in mM): 125 NaCl, 2.5 KCl, 2 $CaCl_2$, 2 $MgCl_2$, 1NaH_2PO_4,

26 $NaHCO_3$, 20 D-glucose, 2 ascorbic acid, and 3 myo-inositol and had an osmolarity of 310–320

mOsm with pH 7.3–7.4.

Ex vivo Acute Brain Slice Whole-Cell Electrophysiology

Neurons in the LC were visualized using an upright microscope (Olympus BX51WI) with

a 40x water-immersion objective. The microscope was able to detect interference contrast and

imaged via an infrared-sensitive camera connected to a video monitor. Whole-cell patch-clamp

recordings were performed at 32 °C maintained with an in-line solution heater from visually

identified LC neurons using the 4[th] ventricle as a landmark. Noradrenergic neurons in the LC were

identified by the presence of tdTomato in Dbh-tdTomato mice and action potential properties of

putative noradrenergic LC neurons in C57bl6/J mice were verified to be similar to Dbh-tdTomato+

neurons.

Using a horizontal puller (P1000, Sutter Instruments, Novato, CA, USA), patch pipettes

were pulled from filamented borosilicate glass capillaries (outer diameter 1.5 mm, inner diameter

0.86, Sutter Instruments), having a tip resistance of 5–7 MΩ when filled with potassium gluconate-

based internal solution that contained (in mM): 139.6 mM K-gluconate, 0.4 mM KCl, 4 mM NaCl,

0.5 mM $CaCl_2$, 10 mM Hepes, 5 mM EGTA free acid, 4 mM Mg-ATP, 0.5 mM Na-GTP with an

osmolality 285–290 mOsm and the pH adjusted to 7.2-7.3 with KOH (E_{Cl} = -85 mV). The

calculated liquid junction potential of 15.5 mV was left uncorrected. For experiments assessing

the role of ATP-sensitive potassium channels, all nucleotide salts were removed from the internal

solution. For voltage-clamp recordings of synaptic activity, 5 mM QX-314 was included in the

internal solution (iso-osmotically replaced NaCl) to block voltage-gated Na^+ channels. All signals

were acquired at 20 kHz and low-pass filtered at 10 kHz via a MultiClamp 700B amplifier. Data

were collected from the neurons with an input resistance >100 MΩ. If the series resistance for a

given recording was >30 MΩ or changed by more than 20%, the data were rejected for analysis.

The current and voltage signals were recorded and digitized with a MultiClamp 700B amplifier

and Digitdata 1440 controlled by Clampex 10 data acquisition software (Molecular Devices, San

Jose, CA, USA). For 5-6 weeks old LC excitability and CRF response recordings, baseline

excitability was collected 5 minutes after whole-cell and after evoked current injections (from -30

pA, increasing by 10 pA up to 100 pA with 10 seconds between each sweep) to ensure stable

recordings and complete dialysis of the cytosolic space with intracellular solution. CRF responses

were collected within 10 minutes following 500 nM CRF (human, rat, AnaSpec, Fremont, CA,

USA) applied in the ACSF bath application. For recordings from 7-8-week old mice, spontaneous

firing rates were collected first during a 3-5-minute free run, with sweeps excluded if they were

20% different from the average firing rate. Evoked firing responses to current injection steps were

next performed using the same protocol as previously described. IN current-clamp mode, the

neuronal membrane potential was kept at -60 mV by injecting an additional holding current, then

the same current injection protocol was performed. The action potential threshold was determined

by taking the average potential at which the dV/dt reached 20mV/ms when neurons fired

spontaneously. To determine the resting membrane, an all-points histogram was generated for

the spontaneous firing recording and the peak of the histogram (0.5 mV bins) was identified as

the resting potential in highly active neurons. Neuronal membrane resistance was determined

based on the holding current required for 5 mV step in voltage-clamp mode.

Quantification and statistical analysis

All data were evaluated with two- or three-way ANOVAs or repeated measures ANOVAs

(RMANOVAs), results of which are reported in **Supplementary Table 1**. If Spearman's test for

heteroscedasticity and the Shapiro-Wilk test for normality were not significant ($p > 0.05$), paired

t-tests (drug effects), unpaired t-tests (stress effects), or Fisher's Exact tests were used for

pairwise comparisons as indicated. If either Spearman's or the Shapiro-Wilk test were significant

($p < 0.05$), a Wilcoxon Signed Rank Test for paired data (dug effects) or Mann-Whitney U test for

unpaired data were used for pairwise comparisons. Alpha levels of $p < 0.05$ were considered

significant for all main effects and pairwise comparisons, and $p < 0.10$ was considered significant

only for interactions when the main effects of stress had $p < 0.05$ in ANOVAs (**Figure 1c** and **2f**).

All results are expressed as the mean ± SEM and individual data points represent individual

animals in behavioral assays or neurons in cellular assays. Analyses and graphing were

conducted in Clampfit 10.0 (Molecular Devices, San Jose, CA, USA), Easy Electrophysiology

(Easy Electrophysiology Ltd, London, UK), Prism (10.0.2, Graphpad, Boston, MA, USA), and Igor

Pro 8 (WaveMetrics, Lake Oswego, OR, USA)

Results

ELVS induces hyperactivity in a non-novel environment and impaired short-term memory with

female-specific increased exploratory and anhedonia-like behavior.

To determine how early life stress impacts affective, cognitive, and locomotive behaviors

of male and female mice, we exposed C57Bl/6J (B6J) mice to a two-phase stress paradigm during

early life (P6-16) and adolescence (P28-P42) as described. At 6-7 weeks of age, male and female control (handled daily) and ELVS mice were evaluated on a behavioral battery, including open field (OF), Y-maze, elevated zero maze (EZM), novel object recognition (NOR), sucrose preference (SP), and rotarod. Female ELVS mice displayed an increase in OF distance moved (**Figure 1a-c**) when compared to female controls that persisted across the 30 min trial and was absent in male mice. In contrast, total locomotion in the familiar open field after the third exposure (habituation phase of NOR) was significantly increased in ELVS-exposed mice, regardless of sex, indicating the development of hyperactivity in a familiar environment due to ELVS exposure. Female ELVS mice also entered the center of the open field arena more frequently than control females (**Figure 1e**), consistent with total OF locomotion. Mice of both sexes spent similar amounts of time in the center of the OF, regardless of ELVS exposure (**Figure 1f**), suggesting a lack of increased anxiety-like behavior despite ELVS exposure. Consistent with OF center time and absence of anxiety-like behavior, there were no significant differences between groups in the cumulative duration of time spent in the open arms of the elevated zero maze (**Figure 1g, h**). In the Y-maze (**Figure 1i-k**), ELVS-exposed mice of both sexes demonstrated increased arm entries (**Figure 1j**) and decreased correct arm alternation sequences (**Figure 1k**), indicating modestly impaired short-term memory relative to control mice. There were no significant sex- or stress-related differences observed in 24 hr novel object recognition (NOR, **Supp. Figure 1a-d**) or rotarod latency to fall times (**Supp. Figure 1e-g**), which tested long-term (24 hr) memory and motor function/coordination, respectively.

Following innate behavioral testing at 6 weeks of age, 7-week-old mice were water restricted for two weeks, then completed sucrose preference testing. Only female ELVS mice displayed a significant reduction in sucrose preference (**Figure 1m**) relative to control mice, which suggests ELVS induces an anhedonia-like state in female mice. Overall, in mice of both sexes at 1.5 – 2 months of age, ELVS exposure led to increased exploratory activity and impaired short-term memory, both ADHD-relevant behavioral domains. In parallel, hyperactivity and anhedonia-

like behavior were observed specifically in female mice, indicating changes in affective behavior

due to a previous exposure to dual-phase ELVS.

ELVS exposure results in male-specific increases in impulsivity and female-specific deficits in

attention.

Mice continued to be water restricted for the five-choice serial reaction time task (5CSRTT)

training and testing. There was no effect of stress on the time necessary for mice to reach the

criterion threshold (>80% accuracy and <30% omitted trials; **Figure 1a**). Although there were

group effects of stress and sex on the total time to complete all training and testing sessions

(**Figure 1b**), there were no significant ELVS effects on these metrics for either sex. As a measure

of impulsivity, we evaluated the number of premature nosepoke responses for the first training

trial when the stimulus duration and intertrial interval were shortened (**Figure 1c**), where we found

that premature responses increased in males exposed to ELVS in contrast to being decreased in

ELVS-exposed females. Once the mice had been trained to criterion on standard parameters,

mice were evaluated in 5CSRTT sessions with varied parameters across ten different test

sessions. Accuracy and trial omissions were similar for all groups on most test sessions (**Supp.**

Figure 2), including those with short stimulus duration (**Figure 1d**, **g**). However, when the intertrial

interval was reduced, ELVS-exposed mice of both sexes performed with similar accuracy (**Figure**

2e), but omitted significantly more trials (**Figure 2h**). 5CSRTT response accuracy was also

reduced in ELVS female mice when session parameters were challenging with longer intertrial

intervals in the presence of an audio distractor and the target brightness was varied (**Figure 2f**),

with no changes in omissions (**Figure 2i**). These behavioral data demonstrate how ELVS

exposure leads to male-specific increases in impulsivity while impairing specific attention

parameters in ELVS females, suggesting sexually dimorphic effects of ELVS on specific neural

circuits responsible for these behavioral changes.

Corticotropin-releasing factor (CRF) directly excites noradrenergic neurons based on ELVS
exposure and via different signaling pathways in male and female mice.

Based on previous studies showing the LC's involvement in affective and attentive behaviors [35, 49, 50] and sex differences in second messenger pathways mediating CRF effects in LC neuron [32], we further probed this sex difference during adolescence to determine the degree to which CRF-evoked changes in LC excitability is dependent on sex and ELVS exposure. To determine the accuracy of targeting putative noradrenergic LC neurons recorded in wildtype B6J mice, we assess CRF (500 nM) effects on LC neuron activity using transgenic mice expressing tdTomato driven by noradrenergic marker, dopamine β-hydroxylase (Dbh-tdTomato mice) (**Figure 3a** & **b**). Baseline and CRF-induced changes in firing behavior of tdTomato+ noradrenergic LC neurons were similar to baseline and CRF-induced changes in B6J mice of both sexes (**Figure 3c-g**). Because the generation of Dbh-tdTomato mice relies on a knock-in of Cre in place of one copy of the *Dbh* gene, mechanistic studies were not performed in these mice due to potential compensatory changes that may occur in the absence of one copy of *Dbh*. In B6J mice, we controlled for whole cell configuration alterations in the internal cell conditions by performing cell-attached recordings of responses to CRF bath applications, with CRF responses similar to whole cell results (**Supp. Figure 3a**). There were also no significant changes in spontaneous firing rate when cells were held in the whole cell condition (**Supp. Figure 3b**), suggesting that this did not confound the observed firing changes upon CRF application.

To determine how ELVS exposure impacts CRF-induced LC neuron excitability changes, we performed the same protocol of bath-applying CRF (500 nM) onto LC neurons of ELVS mice at 6 weeks of age after completion of the ELVS protocol. While ELVS male LC neurons were still responsive to CRF, there was a significant interaction between sex and stress on the effects of CRF on spontaneous firing, with ELVS female LC neurons no longer responding by a change in firing rate (**Figure 3e-g**). Sex differences in LC neuron CRF sensitivity may depend on sexually

dimorphic variations in CRF actions on LC neurons, which may shift after exposure to stress of

different types and timing in early life.

Although it has previously been reported that female LC CRFR1 receptor is predominantly

coupled to the G_s pathway [51–53], since CRFR1 can couple to G_s, G_q, and G_i pathways, we went

on to determine the role of G_q and G_s signaling in CRF-induced increases in LC neuron excitability

in both sexes. To determine G_q pathway dependence of CRF effects on excitability, a protein

lipase C (PLC) inhibitor 10 µM U73122 was added to the internal solution. In the presence of the

PLC inhibitor, bath application of 500 nM CRF no longer increased spontaneous firing of male

noradrenergic neurons (**Figure 3h** & **i**). In contrast, the inclusion of the PLC inhibitor had no effect

on CRF-induced changes in the spontaneous firing behavior of female noradrenergic LC neurons

(**Figure 3h & i**). 10 µM NF 449, a $G\alpha_s$ inhibitor [54], was next added to the internal pipette solution

to determine dependence of G_s-mediated signaling in CRF-induced excitability changes. CRF no

longer significantly increased female LC neurons' spontaneous firing in the presence of NF 449,

but CRF still significantly increased spontaneous firing in LC neurons from males (**Figure 3j** & **k**).

CRF did not have a robust effect on evoked firing in any groups or pharmacological treatments,

except in naïve males with NF449 added to the internal (**Supp. Figure 4**), which is consistent with

changes in spontaneous firing in response to CRF. Together, these data suggest that CRF

preferentially signals through G_q signaling pathways in males and G_s signaling pathways in

females, which may explain how stress signaling may lead to sexually dimorphic long-term

changes.

ELVS Female LC Neurons are less excitable after stress and show increased action potential

delay attributed to A-type potassium channels.

Given the sexual dimorphism of the behavioral phenotype produced by ELVS, sex

differences in LC neurons' CRF sensitivity, female-specific ELVS effects on LC neuronal CRF

sensitivity, and the lack of excitatory or inhibitory synaptic activity differences after stress (**Supp.**

Figure 5), we next assessed current clamp recordings from putative LC noradrenergic neurons

in *ex vivo* acute brain slices to determine whether ELVS produced persistent sex-specific changes

in excitability. At 6 weeks of age, immediately following stress, there were no clear changes

observed in LC neuron excitability (**Supp. Figure 6**), although at 7-8 weeks of age we found

robust ELVS-induced female-specific changes in LC neuron activity (**Figure 4**). Noradrenergic LC

neurons from female ELVS mice, but not males, displayed reduced spontaneous firing rates when

compared to controls (**Figure 4a** & **b)** with no changes in resting membrane potential (V_m; **Figure**

4c). Surprisingly, the membrane resistance was higher and the action potential (AP) threshold

more depolarized in ELVS female LC neurons (**Figure 4d** & **e**). Both male and female ELVS LC

neurons displayed modest, but significantly increased AP half-width (**Figure 4g**). LC neurons from

ELVS females also elicited fewer action potentials in response to current injection compared to

controls, a change which was absent in males (**Figure 4h-l**).

Since the resting potential of LC neurons was quite variable, in the same population of

neurons we also recorded evoked action potential firing when constant current was injected to

maintain the membrane potential at -60 mV to identify specifically how excitability may change

when controlling for V_m. From an initial resting potential of -60 mV in all neurons, ELVS females

continued to show reduced evoked excitability (**Figure 4o-q**), while absent in males (**Figure 4o**)

this excitability change appeared to be due to an increase in the delay time to the first action

potential following current injection in females (**Figure 4s & t**), at least in part. Although the delay

time to the first action potential was variable in all groups, almost all LC neurons in female ELVS

mice displayed a notable delay (>40-50 ms) in response to 70 pA current injection, while LC

neurons from animals in all other groups displayed both no delay and long delays (**Figure 4t**).

This highlights a female-specific LC physiology change that could contribute to the observed

sexually dimorphic LC-involved behavioral differences, potentially due to changes in distal release

of NE throughout the brain.

To determine the mechanism involved in reducing excitability and increased action

potential delay time in ELVS female LC neurons, we first investigated the role of ATP-sensitive

K+ channels, which could account for changes in excitability, by removing both cyclic nucleotides

(ATP and GTP) from the internal solution to promote maximal K_{ATP} activity and reduce variations

in activity that could alter membrane resistance and V_m (**Supp. Figure 7**). Although removing ATP

altered some stress effects on certain action potential properties, such as the action potential

threshold, the medium after hyperpolarization (mAHP) amplitude, and the action potential

halfwidth, female ELVS LC neurons remained less excitable both at their resting potential (**Supp.**

Figure 4i, **k**, **l**) and at -60 mV (**Supp. Figure 4n**, **p**, **q**). Importantly, the sex-specific ELVS-induced

increase in action potential delay time also persisted (**Supp. Figure 7r-t**). It is worth noting the

ELVS males showed a significant reduction in spontaneous firing rate when compared to controls

when ATP and GTP were excluded, but did not show any significant differences in evoked firing

at rest or at -60 mV (**Supp. Figure 7**).

Since action potential delays and suppression of firing rate in response to depolarizing

current are due to the activation of transient A-type voltage-gated potassium (K_v) channel-

mediated currents (I_A) known to be present in LC neurons and modifiable [55–57], we

pharmacologically assessed the I_A-dependence of this delay with 4-aminopyridine (4-AP, 1 mM).

To determine if this difference in A-type potassium channels contributes to the increase in action

potential delay time in ELVS females, we recorded current-injection evoked action potentials from

a V_m of -60 mV before and after bath application of 4-AP (1 mM) to block these transiently

activated channels (**Figure 5a**). In nearly all neurons from each group, 4-AP significantly reduced

the action potential delay time (**Figure 5b-f**), except for male ELVS mice where the delay time

was already short. In the presence of 4-AP, the action potential delay times were comparable in

all groups (**Figure 5f**). These data indicate that the 4-AP-sensitive increased action potential delay

time in ELVS female LC neurons is due to some change in expression or modulation of the K_v

channel or accessory protein mediating I_A that occurs due to ELVS exposure.

Restoration of extracellular NE levels by reboxetine ameliorated ELVS-induced changes in

behavior.

Electrophysiological recordings of LC neurons indicate a reduced excitability and

increased delay to respond to current injection, both of which would lead to reduced distal NE

release. Therefore, to determine the impact of these physiological changes on behavior, we

assessed whether increasing norepinephrine levels at noradrenergic synapses using reboxetine,

a selective norepinephrine reuptake inhibitor, would restore behaviors found to be altered in

separate cohorts of control and ELVS mice (**Figure 6**). We found that 2mg/kg oral reboxetine

administration restored female-specific effects of ELVS on sucrose preference behavior (**Figure**

6a & **b**). Oral reboxetine also significantly reduced exploratory behavior in ELVS mice (**Figure 6c**

& **d**), but did not alter y-maze performance (**Figure 6e** & **f**). Overall, this indicates that

manipulating the LC-NE system in a manner that is targeted based on the sex- and stress-

dependent changes in physiology may be beneficial in restoring specific affective behavioral

changes.

Discussion

Summary of Findings

In our two-phase animal model developed to mimic ACE-relevant timing of and systematic

variation in exposure to threat and deprivation in early life in mice, we identified altered exploratory

locomotion in a familiar environment, short-term memory in early adulthood, and evidence of

reduced ability to maintain attention, regardless of sex (**Figures 1 & 2**). We also found male-

specific increases in impulsivity and evidence of female-specific reduced impulsivity, increased

exploratory behavior, and an anhedonia-like state (**Figures 1 & 2**). Notably, changes in basal

anxiety-like behaviors were absent due to ELVS exposure in the B6J mice used here, in contrast

to Balb/C mice used in our previous work [58], where a prominent increase in anxiety-like behavior

was observed in female mice. Motivated by the role of norepinephrine release from the LC in

shaping these behaviors, our evaluation of LC neurons in these mice using electrophysiological

recordings identified a reduction in LC neuron excitability that was due in part to a prolongation of

the onset action potential delay (**Figure 4**), which was mediated by a 4-AP-sensitive A-type Kv

channel (**Figure 5**). Unlike male Balb/C mice exposed to a similar ELVS protocol [58], basal

properties of LC neurons from B6J male mice exposed to ELVS were remarkably similar to those

from control mice. In addition to blunted CRF sensitivity specifically in female mice exposed to

ELVS, in LC neurons from naïve mice, CRF-induced excitability changes were due to involvement

of sex-specific signaling pathways (**Figure 3**), which may partly explain why exposure to stressful

stimuli during adolescence leads to sex-specific changes in basal LC neuron excitability. Finally,

pharmacologically compensating for this loss of LC neuron excitability/NE release by blocking NE

reuptake with reboxetine [59] restored a subset of ELVS-induced behavioral changes (**Figure 6**),

a drug that has been used clinically to treat depression, panic disorder, and ADHD [60–62] with

variable success. Together, these data highlight the dynamics and importance of understanding

the complex interplay between sex, age (critical periods), genotypic variation, stress exposure,

and neuromodulator circuit changes.

The Impact of Stress and Sex on LC Function Related to Affect and Cognition

The LC-NE system is also well-documented to be stress-sensitive, as it is activated by an

array of environmental stressors, including social and predator stress [19, 21, 63–69]. Largely

limited to stress exposure in males, acute stress-induced changes in LC_{NE} activity is proposed to

occur primarily through the activation of $CRFR_1$ by CRF released from several brain areas,

including the central amygdala and paraventricular nucleus of the hypothalamus [19, 64, 70–73].

At least in male rats, stress and stress-related CRF release shift LC activity towards a high tonic

firing mode (3-8 Hz) while simultaneously decreasing phasic firing events [72, 74, 75]. These

stress-induced increases in CRF and modulation of LC-NE neuron activity are paralleled by CRF-

induced changes in multiple processes and behaviors known to be heavily shaped by LC-NE

neuron activity, including attention [76], cognitive flexibility [19], anxiety [10, 15], and memory [77].

The data presented here shed light on how the LC-NE system changes with multiple early life

stress exposures, and likely related increase in CRF release, depending on sex.

The nature of the LC physiology changes of female mice due to ELVS exposure has the

potential to affect both tonic and phasic firing, each of which is proposed to be related to specific

behavioral events [16, 17, 28, 78]. First, the reduced LC spontaneous and persistent evoked firing

rate observed only in ELVS-exposed females will affect tonic firing. Second, by reducing the

likelihood of generating an action potential in response to short depolarizing events (i.e. the

beginning of current injection steps), the prolongation of the action potential delay time will act as

a filter of fast synaptic inputs to reduce the likelihood of phasic firing and subsequent distal NE

release. By disrupting both types of LC neuron firing modes, these changes observed in females

could result in disrupted NE signaling in multiple brain regions involved in attention, arousal,

memory, and/or locomotion (i.e. prefrontal cortex, hippocampus, ventral tegmental area, and

lateral cerebellar nucleus).

Although LC-NE neurons were previously thought to be a homogenous population, recent

data suggest that they are much more heterogeneous in their molecular [79] or functional

properties that relate to specific LC-NE projection targets [80, 81]. Given the range of delay times,

resting potentials, and spontaneous firing rates observed in this population of LC neurons, it

appears that ELVS exposure leads to a shift in the distribution of these key properties across the

population, likely due to sex-specific variation in CRF-activated signaling. However, it is not clear

how a change in these functional properties may relate to changes in specific molecularly-,

anatomically- (target brain area), or functionally-defined LC neuron sub-populations. Whether

acute or persistent changes in the physiological properties of LC-NE neurons and/or their synaptic

properties, like those observed here, may impact NE release throughout the central nervous

system differentially based on specific circuits or related behavior is unclear. Since there is a

smaller distribution of firing rates yet lager distribution of action potential delay times across LC

neurons in ELVS females when compared to controls, this property-dependent shift in the

functional diversity of LC neurons could impact how firing modes required for optimal projecting

brain region states. For instance, distinct LC ensembles projecting to the cortex may modulate

particular cortical states with members of each ensemble having definitive firing rates [82]. Further

effort to determine exactly how these changes in LC-NE system excitability alter widespread

noradrenergic signaling *in vivo* during specific behaviors would provide support for the importance

of this physiological change and suggest the LC-NE system to be a female-specific target of

interest for stress-related disorders, particularly with onset after adolescence.

The specific sexually dimorphic molecular mechanisms underpinning LC physiology

changes after exposure to the ELVS are still unclear. However, our results in naïve mice (**Figure**

3) establish that there are basal sex differences in how CRF exposure affects the firing of LC

neurons, at least in the signaling pathways involved in CRF-induced increases in firing rate,

indicating preferential G_q signaling in males and G_s in females. These data are in line with other

reports of female CRFR1 in the LC being predominantly G_s-coupled and males having increased

β-arrestin-related internalization of CRFR1 after acute stress [32]. Bridging the gap in future work

to understand how two-phase early life stress and acute stress affect CRF-dependent changes in

LC physiology and related signaling based on sex will be important to understand fundamental

sex-specific processes that shape responses to stress on both timescales. Based on these data,

we predict that the predominant G_s-dependent CRF actions in females, leading to PKA activation,

will cause unique subsequent and persistent changes in LC physiology (excitability, delay) in

females that will be absent in males. In parallel, the G_q coupling predominance could be protective

in males due to its transient Ca^{2+}-dependent action versus a prolonged upregulation of a signaling

cascade. Given recent reports of persistent LC physiology changes occurring only when stress

exposure happens at adolescence and not adulthood [83], determining the degree to which this

applies to our dual-phase ELVS paradigm and the duration of any lasting behavioral and

physiological changes will also be important to understand dynamics that shape the broader

impacts of stress. Finally, at a molecular level, it is unclear how exactly I_A is modulated or affected

by stress, but clarifying the channels and auxiliary subunit changes over longer time scales may

reveal novel mechanisms of broader and LC-specific responses to critically-timed stress to shape

novel prevention and treatment strategies to improve specific behavioral outcomes.

Clinical Impact of Findings

The treatment of stress- or specifically ACE-related disorders should consider sex and the

timing of stress exposure as biological variables influencing patient outcomes. These behavioral

results suggest females may be more sensitive to depression-like behaviors and ADHD-relevant

deficits as a result of early life stress during comparable time periods used in this study. Based

on these data, individuals exposed to ACEs as adolescents may be particularly sensitive to

developing negative neuropsychiatric outcomes later in life, based on sex differences in LC

neuron physiology and central stress (CRF) sensitivity at this time. Regardless, the specific role

of sex hormones in development and neurophysiological changes in response to stress is unclear.

Estrogen receptor α (ERα) and β (ERβ) are differentially expressed in the amygdala, prefrontal

cortex, and hippocampus responsible for stress, anxiety, and cognitive behaviors, respectively

[84]. ERβ has been associated with non-reproductive behaviors, with its activation improving

memory and reducing anxious behaviors in females [85, 86]. In contrast, testosterone signaling

has been suggested to be a factor supporting resilience to stress-induced changes, which is in

line with the behavioral data presented here [87, 88].

This study may implicate the adrenergic system, particularly the LC in the brain, as a

region to explore biomarkers of dysfunction in female patients with a history of stress in early life.

According to the DSM-V, cognitive issues are a core symptom of depression that include aspects

of memory and attention being disrupted [89, 90]. These symptoms are in line with the female-

specific behavioral impacts of our dual-phase stress model, pointing to increasing noradrenergic

tone as a therapeutic avenue for treatment. There are many drugs that target this brain area,

particularly second-line therapies for attention deficit hyperactivity disorder (ADHD), such as

reboxetine [61] used in this study, which may prove to be effective in managing cognitive

impairments in this patient group with depression and/or PTSD. These data provide a foundation

for further research into neuro-stress regulation differences in males and females, relevant to

many stress-related disorders, from PTSD to substance abuse disorder, that have sex differences

in their prevalence ratios [46]. Perhaps a more robust individualized understanding of how

patients' genetic and environmental risk factors interact with the type of timing of major stressful

life events may increase the likelihood of therapeutic success when these variables are used to

optimize drug choice.

Acknowledgements

This work was supported by institutional start-up funds and a Research Seed Grant from SIU-

School of Medicine provided to BDR.

Conflict of Interest

Neither author has any conflicts of interest to declare.

References

1.   Pitts BL, Eisenberg ML, Bailey HR, Zacks JM. PTSD is associated with impaired event

processing and memory for everyday events. Cognitive Research: Principles and

Implications. 2022;7:35.

2.   Vesga-López O, Blanco C, Keyes K, Olfson M, Grant BF, Hasin DS. Psychiatric Disorders in

Pregnant and Postpartum Women in the United States. Archives of General Psychiatry.

2008;65:805.

3.   Hickey M, Bryant C, Judd F. Evaluation and management of depressive and anxiety

symptoms in midlife. Climacteric. 2012;15:3–9.

4.   Aston-Jones G, Rajkowski J, Kubiak P, Alexinsky T. Locus coeruleus neurons in monkey are

selectively activated by attended cues in a vigilance task. The Journal of Neuroscience :

The Official Journal of the Society for Neuroscience. 1994;14:4467–4480.

5.   Kuo C-C, Hsieh J-C, Tsai H-C, Kuo Y-S, Yau H-J, Chen C-C, et al. Inhibitory interneurons

regulate phasic activity of noradrenergic neurons in the mouse locus coeruleus and

functional implications. The Journal of Physiology. 2020;598:4003–4029.

6.   Borodovitsyna O, Duffy BC, Pickering AE, Chandler DJ. Anatomically and functionally

distinct locus coeruleus efferents mediate opposing effects on anxiety-like behavior.

Neurobiology of Stress. 2020;13.

7.   Poe GR, Foote S, Eschenko O, Johansen JP, Bouret S, Aston-Jones G, et al. Locus coeruleus:

a new look at the blue spot. Nat Rev Neurosci. 2020;21:644–659.

8.   Chandler DJ, Jensen P, McCall JG, Pickering AE, Schwarz LA, Totah NK. Redefining

Noradrenergic Neuromodulation of Behavior: Impacts of a Modular Locus Coeruleus

Architecture. J Neurosci. 2019;39:8239–8249.

9.   Breton-Provencher V, Drummond GT, Sur M. Locus Coeruleus Norepinephrine in Learned

Behavior: Anatomical Modularity and Spatiotemporal Integration in Targets. Frontiers in

Neural Circuits. 2021;15.

10. McCall JG, Siuda ER, Bhatti DL, Lawson LA, McElligott ZA, Stuber GD, et al. Locus coeruleus

to basolateral amygdala noradrenergic projections promote anxiety-like behavior. eLife.

2017;6.

11. Ishida Y, Shirokawa T, Miyaishi O, Komatsu Y, Isobe K. Age-dependent changes in

projections from locus coeruleus to hippocampus dentate gyrus and frontal cortex. The

European Journal of Neuroscience. 2000;12:1263–1270.

12. Lim EP, Tan CH, Jay TM, Dawe GS. Locus coeruleus stimulation and noradrenergic

modulation of hippocampo-prefrontal cortex long-term potentiation. The International

Journal of Neuropsychopharmacology. 2010;13:1219–1231.

13. Mair RD, Zhang Y, Bailey KR, Toupin MM, Mair RG. Effects of clonidine in the locus

coeruleus on prefrontal- and hippocampal-dependent measures of attention and memory

in the rat. Psychopharmacology. 2005;181:280–288.

14. Dornellas APS, Burnham NW, Luhn KL, Petruzzi MV, Thiele TE, Navarro M. Activation of

locus coeruleus to rostromedial tegmental nucleus (RMTg) noradrenergic pathway blunts

binge-like ethanol drinking and induces aversive responses in mice. Neuropharmacology.

2021;199.

15.   Borodovitsyna O, Duffy BC, Pickering AE, Chandler DJ. Anatomically and functionally

distinct locus coeruleus efferents mediate opposing effects on anxiety-like behavior.

Neurobiology of Stress. 2020;13.

16.   Rajkowski J, Kubiak P, Aston-Jones G. Locus coeruleus activity in monkey: phasic and tonic

changes are associated with altered vigilance. Brain Research Bulletin. 1994;35:607–616.

17.   Howells FM, Stein DJ, Russell VA. Synergistic tonic and phasic activity of the locus

coeruleus norepinephrine (LC-NE) arousal system is required for optimal attentional

performance. Metabolic Brain Disease. 2012;27:267–274.

18.   McBurney-Lin J, Vargova G, Garad M, Zagha E, Yang H. The locus coeruleus mediates

behavioral flexibility. Cell Rep. 2022;41:111534.

19.   Snyder K, Wang W-W, Han R, McFadden K, Valentino RJ. Corticotropin-releasing factor in

the norepinephrine nucleus, locus coeruleus, facilitates behavioral flexibility.

Neuropsychopharmacology. 2012;37:520–530.

20.   Su Z, Cohen JY. Two types of locus coeruleus norepinephrine neurons drive reinforcement

learning. 2022.

21.   McCall JG, Al-Hasani R, Siuda ER, Hong DY, Norris AJ, Ford CP, et al. CRH Engagement of

the Locus Coeruleus Noradrenergic System Mediates Stress-Induced Anxiety. Neuron.

2015;87:605–620.

22.   Pasquier DA, Reinoso-Suarez F. The topographic organization of hypothalamic and brain

stem projections to the hippocampus. Brain Research Bulletin. 1978;3:373–389.

23.   Fitzpatrick CM, Runegaard AH, Christiansen SH, Hansen NW, Jørgensen SH, McGirr JC, et

al. Differential effects of chemogenetic inhibition of dopamine and norepinephrine

625  neurons in the mouse 5-choice serial reaction time task. Progress in Neuro-

626  Psychopharmacology & Biological Psychiatry. 2019;90:264–276.

627 24. Uematsu A, Tan BZ, Ycu EA, Cuevas JS, Koivumaa J, Junyent F, et al. Modular organization

628  of the brainstem noradrenaline system coordinates opposing learning states. Nature

629  Neuroscience. 2017;20:1602–1611.

630 25. Campese VD, Soroeta JM, Vazey EM, Aston-Jones G, LeDoux JE, Sears RM. Noradrenergic

631  Regulation of Central Amygdala in Aversive Pavlovian-to-Instrumental Transfer. eNeuro.

632  2017;4.

633 26. Carlson ES, Hunker AC, Sandberg SG, Locke TM, Geller JM, Schindler AG, et al.

634  Catecholaminergic Innervation of the Lateral Nucleus of the Cerebellum Modulates

635  Cognitive Behaviors. The Journal of Neuroscience : The Official Journal of the Society for

636  Neuroscience. 2021;41:3512–3530.

637 27. Feng J, Dong H, Lischinsky JE, Zhou J, Deng F, Zhuang C, et al. Monitoring norepinephrine

638  release in vivo using next-generation GRABNE sensors. Neuron. 2024;112:1930-1942.e6.

639 28. Vazey EM, Moorman DE, Aston-Jones G. Phasic locus coeruleus activity regulates cortical

640  encoding of salience information. Proc Natl Acad Sci U S A. 2018;115:E9439–E9448.

641 29. Devilbiss DM, Page ME, Waterhouse BD. Locus Ceruleus Regulates Sensory Encoding by

642  Neurons and Networks in Waking Animals. J Neurosci. 2006;26:9860–9872.

643 30. Devilbiss DM, Waterhouse BD. The Effects of Tonic Locus Ceruleus Output on Sensory-

644  Evoked Responses of Ventral Posterior Medial Thalamic and Barrel Field Cortical Neurons

645  in the Awake Rat. J Neurosci. 2004;24:10773–10785.

646   31.   Butler PD, Weiss JM, Stout JC, Nemeroff CB. Corticotropin-releasing factor produces fear-

647   enhancing and behavioral activating effects following infusion into the locus coeruleus. J

648   Neurosci. 1990;10:176–183.

649   32.   Bangasser DA, Curtis A, Reyes BAS, Bethea TT, Parastatidis I, Ischiropoulos H, et al. Sex

650   differences in corticotropin-releasing factor receptor signaling and trafficking: potential

651   role in female vulnerability to stress-related psychopathology. Molecular Psychiatry.

652   2010;15:877, 896–904.

653   33.   Valentino RJ, Foote SL, Aston-Jones G. Corticotropin-releasing factor activates

654   noradrenergic neurons of the locus coeruleus. Brain Research. 1983;270:363–367.

655   34.   Wang Q, Liu Y, Zhang J, Wang W. Corticotropin-Releasing Factor Receptors in the Locus

656   Coeruleus Modulate the Enhancement of Active Coping Behaviors Induced by Chronic

657   Predator Odor Inoculation in Mice. Frontiers in Psychology. 2020;10.

658   35.   Seo D, Zhang ET, Piantadosi SC, Marcus DJ, Motard LE, Kan BK, et al. A locus coeruleus to

659   dentate gyrus noradrenergic circuit modulates aversive contextual processing. Neuron.

660   2021;109:2116-2130.e6.

661   36.   Breton-Provencher V, Drummond GT, Sur M. Locus Coeruleus Norepinephrine in Learned

662   Behavior: Anatomical Modularity and Spatiotemporal Integration in Targets. Frontiers in

663   Neural Circuits. 2021;15.

664   37.   Lim EP, Tan CH, Jay TM, Dawe GS. Locus coeruleus stimulation and noradrenergic

665   modulation of hippocampo-prefrontal cortex long-term potentiation. The International

666   Journal of Neuropsychopharmacology. 2010;13:1219–1231.

38.   McCall JG, Siuda ER, Bhatti DL, Lawson LA, McElligott ZA, Stuber GD, et al. Locus coeruleus

to basolateral amygdala noradrenergic projections promote anxiety-like behavior. eLife.

2017;6.

39.   Fortress AM, Hamlett ED, Vazey EM, Aston-Jones G, Cass WA, Boger HA, et al. Designer

receptors enhance memory in a mouse model of Down syndrome. The Journal of

Neuroscience : The Official Journal of the Society for Neuroscience. 2015;35:1343–1353.

40.   Mariscal P, Bravo L, Llorca-Torralba M, Razquin J, Miguelez C, Suárez-Pereira I, et al. Sexual

differences in locus coeruleus neurons and related behavior in C57BL/6J mice. Biol Sex

Differ. 2023;14.

41.   Lee J, Wang Z, Messi ML, Milligan C, Furdui CM, Delbono O. Sex differences in single

neuron function and proteomics profiles examined by patch-clamp and mass spectrometry

in the locus coeruleus of the adult mouse. Acta Physiologica. 2024. 8 March 2024.

https://doi.org/10.1111/apha.14123.

42.   Bangasser DA, Zhang X, Garachh V, Hanhauser E, Valentino RJ. Sexual dimorphism in locus

coeruleus dendritic morphology: a structural basis for sex differences in emotional

arousal. Physiology & Behavior. 2011;103:342–351.

43.   Bangasser DA, Wiersielis KR, Khantsis S. Sex differences in the locus coeruleus-

norepinephrine system and its regulation by stress. Brain Research. 2016;1641:177–188.

44.   Mulvey B, Bhatti DL, Gyawali S, Lake AM, Kriaucionis S, Ford CP, et al. Molecular and

Functional Sex Differences of Noradrenergic Neurons in the Mouse Locus Coeruleus. Cell

Reports. 2018;23:2225–2235.

45.   Bangasser DA, Valentino RJ. Sex differences in stress-related psychiatric disorders:

neurobiological perspectives. Frontiers in Neuroendocrinology. 2014;35:303–319.

46.   Swaab DF, Bao A-M. Sex differences in stress-related disorders: Major depressive disorder,

bipolar disorder, and posttraumatic stress disorder. Handbook of Clinical Neurology.

2020;175:335–358.

47.   Tillage RP, Sciolino NR, Plummer NW, Lustberg D, Liles LC, Hsiang M, et al. Elimination of

galanin synthesis in noradrenergic neurons reduces galanin in select brain areas and

promotes active coping behaviors. Brain Struct Funct. 2020;225:785–803.

48.   Madisen L, Mao T, Koch H, Zhuo J, Berenyi A, Fujisawa S, et al. A toolbox of Cre-dependent

optogenetic transgenic mice for light-induced activation and silencing. Nature

Neuroscience. 2012;15:793–802.

49.   Aston-Jones G, Rajkowski J, Cohen J. Role of locus coeruleus in attention and behavioral

flexibility. Biological Psychiatry. 1999;46:1309–1320.

50.   Hallock HL, Adiraju SS, Miranda-Barrientos J, McInerney JM, Oh S, DeBrosse AC, et al.

Electrophysiological correlates of attention in the locus coeruleus-prelimbic cortex circuit

during the rodent continuous performance test. Neuropsychopharmacology : Official

Publication of the American College of Neuropsychopharmacology. 2024;49:521–531.

51.   Bangasser DA, Curtis A, Reyes BAS, Bethea TT, Parastatidis I, Ischiropoulos H, et al. Sex

differences in corticotropin-releasing factor receptor signaling and trafficking: potential

role in female vulnerability to stress-related psychopathology. Molecular Psychiatry.

2010;15:877, 896–904.

52.   Bangasser DA, Wiersielis KR. Sex differences in stress responses: a critical role for

corticotropin-releasing factor. Hormones. 2018;17:5–13.

53.   Bangasser DA, Valentino RJ. Sex differences in stress-related psychiatric disorders:

neurobiological perspectives. Frontiers in Neuroendocrinology. 2014;35:303–319.

54.   Hohenegger M, Waldhoer M, Beindl W, Böing B, Kreimeyer A, Nickel P, et al. Gsalpha-

selective G protein antagonists. Proc Natl Acad Sci U S A. 1998;95:346–351.

55.   Itri JN, Vosko AM, Schroeder A, Dragich JM, Michel S, Colwell CS. Circadian Regulation of

A-Type Potassium Currents in the Suprachiasmatic Nucleus. Journal of Neurophysiology.

2010;103:632–640.

56.   Burdakov D, Ashcroft FM. Cholecystokinin Tunes Firing of an Electrically Distinct Subset of

Arcuate Nucleus Neurons by Activating A-Type Potassium Channels. J Neurosci.

2002;22:6380–6387.

57.   Wu R-N, Hung W-C, Chen C-T, Tsai L-P, Lai W-S, Min M-Y, et al. Firing activity of locus

coeruleus noradrenergic neurons decreases in necdin-deficient mice, an animal model of

Prader–Willi syndrome. J Neurodevelop Disord. 2020;12.

58.   Brannan S, Garbe L, Richardson BD. Early life stress induced sex-specific changes in

behavior is paralleled by altered locus coeruleus physiology in BALB/cJ mice. Neurobiology

of Stress. 2024;33.

59.   Page ME. The promises and pitfalls of reboxetine. CNS Drug Rev. 2003;9:327–342.

60.   Riahi F, Tashakori A, Izadi-Mazidi S, Salehi-Veysi M. Effectiveness of reboxetine in

treatment of outpatient children and adolescents with attention deficit-hyperactivity

disorder with comorbid anxiety disorders. Iran J Psychiatry. 2013;8:195–200.

61. Cohen-Yavin I, Yoran-Hegesh R, Strous RD, Kotler M, Weizman A, Spivak B. Efficacy of

Reboxetine in the Treatment of Attention-Deficit/Hyperactivity Disorder in Boys With

Intolerance to Methylphenidate. Clinical Neuropharmacology. 2009;32:179–182.

62. Block SG, Nemeroff CB. Emerging antidepressants to treat major depressive disorder.

Asian J Psychiatr. 2014;12:7–16.

63. Curtis a L, Bello NT, Valentino RJ. Evidence for functional release of endogenous opioids in

the locus ceruleus during stress termination. The Journal of Neuroscience : The Official

Journal of the Society for Neuroscience. 2001;21:RC152.

64. Curtis AL, Leiser SC, Snyder K, Valentino RJ. Predator stress engages corticotropin-releasing

factor and opioid systems to alter the operating mode of locus coeruleus norepinephrine

neurons. Neuropharmacology. 2012;62:1737–1745.

65. Bingham B, McFadden K, Zhang X, Bhatnagar S, Beck S, Valentino R. Early adolescence as a

critical window during which social stress distinctly alters behavior and brain

norepinephrine activity. Neuropsychopharmacology. 2011;36:896–909.

66. Borodovitsyna O, Flamini MD, Chandler DJ. Acute Stress Persistently Alters Locus

Coeruleus Function and Anxiety-like Behavior in Adolescent Rats. Neuroscience.

2018;373:7–19.

67. Cassens G, Roffman M, Kuruc A, Orsulak PJ, Schildkraut JJ. Alterations in brain

norepinephrine metabolism induced by environmental stimuli previously paired with

inescapable shock. Science. 1980;209:1138–1140.

68. Francis DD, Caldji C, Champagne F, Plotsky PM, Meaney MJ. The role of corticotropin-

releasing factor–norepinephrine systems in mediating the effects of early experience on

the development of behavioral and endocrine responses to stress. Biological Psychiatry.

1999;46:1153–1166.

69. George SA, Knox D, Curtis AL, Aldridge JW, Valentino RJ, Liberzon I. Altered locus

coeruleus–norepinephrine function following single prolonged stress. Eur J of

Neuroscience. 2013;37:901–909.

70. Valentino RJ, Van Bockstaele E. Convergent regulation of locus coeruleus activity as an

adaptive response to stress. Eur J Pharmacol. 2008;583:194–203.

71. Borodovitsyna O, Flamini MD, Chandler DJ. Acute Stress Persistently Alters Locus

Coeruleus Function and Anxiety-like Behavior in Adolescent Rats. Neuroscience.

2018;373:7–19.

72. Reyes BAS, Glaser JD, Bockstaele EJV. Ultrastructural evidence for co-localization of

corticotropin-releasing factor receptor and mu-opioid receptor in the rat nucleus locus

coeruleus. Neuroscience Letters. 2007;413:216–221.

73. Reyes B a. S, Carvalho AF, Vakharia K, Van Bockstaele EJ. Amygdalar peptidergic circuits

regulating noradrenergic locus coeruleus neurons: linking limbic and arousal centers. Exp

Neurol. 2011;230:96–105.

74. Reyes B a. S, Carvalho AF, Vakharia K, Van Bockstaele EJ. Amygdalar peptidergic circuits

regulating noradrenergic locus coeruleus neurons: linking limbic and arousal centers. Exp

Neurol. 2011;230:96–105.

75. Reyes BAS, Zitnik G, Foster C, Van Bockstaele EJ, Valentino RJ. Social Stress Engages

Neurochemically-Distinct Afferents to the Rat Locus Coeruleus Depending on Coping

Strategy,,. eNeuro. 2015;2.

76. Cole RD, Kawasumi Y, Parikh V, Bangasser DA. Corticotropin releasing factor impairs

sustained attention in male and female rats. Behavioural Brain Research. 2016;296:30–34.

77. Paretkar T, Dimitrov E. The Central Amygdala Corticotropin-releasing hormone (CRH)

Neurons Modulation of Anxiety-like Behavior and Hippocampus-dependent Memory in

Mice. Neuroscience. 2018;390:187–197.

78. Grimm C, Duss SN, Privitera M, Munn BR, Karalis N, Frässle S, et al. Tonic and burst-like

locus coeruleus stimulation distinctly shift network activity across the cortical hierarchy.

Nat Neurosci. 2024;27:2167–2177.

79. Hughes AC, Pittman BG, Xu B, Gammons JW, Webb CM, Nolen HG, et al. A single-vector

intersectional AAV strategy for interrogating cellular diversity and brain function. Nature

Neuroscience. 2024;27:1400–1410.

80. Waterhouse BD, Predale HK, Plummer NW, Jensen P, Chandler DJ. Probing the structure

and function of locus coeruleus projections to CNS motor centers. Front Neural Circuits.

2022;16:895481.

81. Chandler DJ, Gao W-J, Waterhouse BD. Heterogeneous organization of the locus coeruleus

projections to prefrontal and motor cortices. Proc Natl Acad Sci U S A. 2014;111:6816–

6821.

82. Noei S, Zouridis IS, Logothetis NK, Panzeri S, Totah NK. Distinct ensembles in the

noradrenergic locus coeruleus are associated with diverse cortical states. Proceedings of

the National Academy of Sciences. 2022;119.

83.   Borodovitsyna O, Tkaczynski JA, Corbett CM, Loweth JA, Chandler DJ. Age- and Sex-

796   Dependent Changes in Locus Coeruleus Physiology and Anxiety-Like Behavior Following

797   Acute Stressor Exposure. Frontiers in Behavioral Neuroscience. 2022;16:808590.

798   84.   Cover KK, Maeng LY, Lebrón-Milad K, Milad MR. Mechanisms of estradiol in fear circuitry:

799   implications for sex differences in psychopathology. Translational Psychiatry. 2014;4:e422.

800   85.   Oyola MG, Portillo W, Reyna A, Foradori CD, Kudwa A, Hinds L, et al. Anxiolytic effects and

801   neuroanatomical targets of estrogen receptor-β (ERβ) activation by a selective ERβ agonist

802   in female mice. Endocrinology. 2012;153:837–846.

803   86.   Jacome LF, Gautreaux C, Inagaki T, Mohan G, Alves S, Lubbers LS, et al. Estradiol and ERβ

804   agonists enhance recognition memory, and DPN, an ERβ agonist, alters brain monoamines.

805   Neurobiology of Learning and Memory. 2010;94:488–498.

806   87.   Pfau ML, Russo SJ. Peripheral and Central Mechanisms of Stress Resilience. Neurobiology

807   of Stress. 2015;1:66–79.

808   88.   Russo SJ, Murrough JW, Han M-H, Charney DS, Nestler EJ. Neurobiology of resilience.

809   Nature Neuroscience. 2012;15:1475–1484.

810   89.   Perini G, Cotta Ramusino M, Sinforiani E, Bernini S, Petrachi R, Costa A. Cognitive

811   impairment in depression: recent advances and novel treatments. Neuropsychiatr Dis

812   Treat. 2019;15:1249–1258.

813   90.   Mattingly G, Anderson RH, Mattingly SG, Anderson EQ. The impact of cognitive challenges

814   in major depression: the role of the primary care physician. Postgrad Med. 2016;128:665–

815   671.

819 *Figure 1*

Figure 1. ELVS causes sex-independent and female-specific changes in cognitive and affective behavior. (**a-c**) Total distance (meters, m) moved in the open field (OF) assay (**a**) across the entire 30 min session (**b**; females: U = 48.0 – 92.0, *p* = 0.00030 – 0.045) and in total (**c**; male: *t*(30) = 0.70, *p* = 0.49; female: *t*(33) = 4.08, *p* = 0.0003). (**d**) Total distance (m) moved in OF apparatus during the third exposure (3x OF) for 10 minutes (combined: *t*(47) = 3.13, *p* = 0.0030). (**e**, **f**) The number of entries into (**e**; male: *t*(30) = 0.63, *p* = 0.53; female: *t*(33) = 4.63, *p* < 0.0001) and total time spent (**f**) in the center of the OF arena. (**g**, **h**) Time (seconds) spent (**h**) in the elevated zero maze (EZM, **g**) open arm. (**i-k**) Y-Maze (**i**) total number of arm entries (**j**; combined: *t*(63) = 2.83, *p* = 0.0063) and correct arm alternation percentages (**k**; combined: *t*(63) = 2.94, *p* = 0.0046). (**l**, **m**) Sucrose preference (**l**) index values as the ratio of 1% sucrose/H_2O only (**m**: male: *t*(14) = 1.89, *p* = 0.08; female: *t*(14) = 5.99, *p* < 0.0001). All results are presented as mean ± SEM for each group with circles representing data from individual mice. **a-k**: N = 10-18 mice (6 weeks) per group. **m**: N = 8 mice (9 weeks) per group. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$ with two-tailed unpaired t-test or Mann-Whitney U test for pairwise comparisons.

Figure 2

Figure 2. ELVS leads to sexually dimorphic changes in impulsivity and female-specific distractibility in 5-choice serial reaction time task (5CSRTT). (**a**) Days needed to meet criteria during 5CSRTT training. (**b**) Days required to complete training and all session variation tests (male: $t(16) = 0.98$, $p = 0.34$; female: $t(18) = 2.02$, p 0.059). (**c**) Number of premature responses during the first training transition session in 5CSRTT training with reduced stimulus duration and intertrial interval (males: U = 15.0, $p = 0.024$; females: U = 22.0, p=0.033). Trial accuracy % (**d-f**) and omission % (**g-i**) in test sessions with shortened stimulus duration in **d** and **g**, shortened intertrial interval in **e** and **h** (combined: $t(36) = 2.97$, $p = 0.0053$), and lengthened intertrial interval with a simultaneous sound distractor in **f** (male: U = 30.5, $p = 0.42$; female: U = 9.0, p=0.0010) and **i**.

Figure 3

Figure 3. CRF excites LC neurons via different pathways in male and female mice and is lost in female mice after stress exposure. (**a**) Schematic of animals used and brain slice approach for recording form LC neurons in mice expressing tdTomato in dopamine β-hydroxylase (DβH)-expressing neurons, control, ELVS-exposed, or naïve mice. (**b**) DIC (top) and fluorescent tdTomato (bottom) image of the LC below the fourth ventricle (left) and a high magnification image of LC neurons in DβH-tdTomato mice. (**c**, **e**, **f**, **h**, **j**) Representative paired membrane voltage traces of spontaneous action potentials generated by LC neurons in ACSF and after bath application of 500nM CRF from DβH-tdTomato (**c**), control C57 (**e**), and ELVS C57 (**f**) mice, or naïve mice with 10µM U73122 (**h**) or 10µM NF 449 (**j**) in the intracellular solution. (**d**, **g**, **i**, **k**)

Group spontaneous action potential frequency of LC neurons in ACSF and after bath application

of 500nM CRF from DβH-tdTomato (**d**; male: $t(10) = 2.55$, $p = 0.029$; female: $t(5) = 2.61$; $p =$

0.048), control and ELVS C57 mice (**g**; Control male – CRF: $p = 0.044$; ELVS male – CRF: $p =$

0.0015; Control female – CRF: $p = 0.020$; female control CRF vs. ELVS CRF: $p = 0.045$; ELVS

female- CRF: p=0.17), or naïve mice with 10µM U73122 (**i**; male: $W = 3$, $p = 0.93$; female: $W = -$

91, $p = 0.0002$) or 10µM NF 449 (**k**; male: $t(10) = 3.44$, $p = 0.0063$; female: $t(12) = 1.53$; $p = 0.15$)

in the intracellular solution. In **d**, **g**, **i**, and **k**, the left y-axis corresponds to mean ± SEM group

bars and right y-axis corresponds to individual paired values for each individual neuron. All group

data are presented as mean ± SEM (bars) for each group, with connected circles representing

paired pre- and post-CRF values from individual neurons. Two- (**d**, **i**, **k**) or three-way (**g**)

RMANOVAs were used for group comparisons, followed by a paired t-test, Wilcoxon Signed-Rank

test for single pairwise comparisons in two-way RMANOVAs or Bonferroni posthoc in three-way

RMANOVAs to assess CRF effects. n = 6-11 neurons from N = 2-4 mice (6-8 weeks old) for **c**

and **d,** or n = 11-16 neurons from N = 3 mice (5-7 weeks old) for **e**-k. *$p < 0.05$, **$p < 0.01$, ***$p <$

0.001 for pairwise comparisons.

Figure 4

Figure 4. ELVS causes reduced LC neuron excitability and increased AP delay time in

female mice. (**a, h, i, m, n**) Representative traces of male and female control and ELVS putative

LC neurons when firing spontaneously (**a**), in response to +80/-30 pA current injection (1 s) when

cells are at their natural resting membrane potential (**h**, **i**) or at -60 mV (**m**, **n**). Note that the same

example recordings for both Vm states are provided from the same neuron, some of which are

the same in both resting potential states because the natural Vm was at -60 mV. (**b-g**) Action

potential spontaneous rate (**b**; males: U = 479.5, *p* = 0.82; females: U = 271.0, p=0.014), resting

membrane potential (**c**), membrane resistance (**d**; males: U = 443.0, *p* = 0.14; females: U = 253.5,

$p < 0.0001$), action potential (AP) threshold (**e**; males: U = 410.0, p = 0.61; females: U = 256.0, p

= 0.0069), medium after hyperpolarization (mAHP) amplitude (**f**), and AP half-width (**g**; males: U

= 314.0, p = 0.020; females: U = 295.0, p = 0.023) for neurons from each group. (**j**, **k**, **o**, **p**) Number

of action potentials elicited by putative LC neurons in response to 1 sec current injection steps

from -30 to +100 pA for male (**j**, **o**) and female (**k**, **p**) mice with inter-step membrane potential

varying naturally (**j**, **k**; females: U = 293.0 – 370.0, p = 0.00030 – 0.031) or maintained near -60

mV (**o**, **p**; females: U = 253.0 – 322.0, p = 0.0019 – 0.038). (**l**, **q**) Number of action potentials

elicited by putative LC neurons in response to maximum 100 pA for male and female mice with

inter-step membrane potential varying naturally (**l**; males: U = 403.5, p = 0.11; females: U = 310.0,

p = 0.025) or maintained near -60 mV (**q**; males: U = 284.0, p = 0.12; females: U = 253.0, p =

0.0019). (**r**, **s**) Delay time to first action potential after current injection onset present in putative

LC neurons in response to 1 sec current injection steps from +70 to +100 pA for male (**r**) and

female (**s**; U = 118.0 – 148.5, $p < 0.0001$) mice with inter-step membrane potential maintained

near -60 mV. (**t**) Delay time to first action potential after current injection onset present in putative

LC neurons in response to 70 pA current injection (1 sec) for male (U = 258.0, p = 0.31) and

female (U = 124.0, $p < 0.0001$) mice. All group data are presented as mean ± SEM (bars or

shading) for each group, with circles representing data from individual neurons. Two- or three-

way ANOVAs were used for group comparisons, followed by a t-test or Mann-Whitney U test for

single pairwise comparisons to assess stress effects. n = 26-35 neurons from N=4-5 mice (7-8

weeks old) for each group. For **b-g**, **l**, **q**, **r-t**, *p < 0.05, **p < 0.01, ***p < 0.001 with two-tailed

unpaired t-test or Mann-Whitney U test. For clarity in **j**, **k**, **o**, **p**, only *p < 0.05 with Mann-Whitney

U test for each current injection.

Figure 5. Blockade of K_A (A-type) potassium channels reduces action potential delay time and normalizes ELVS-induced action potential delay time prolongation. (**a**) Representative recordings from LC neurons in male (top) and female (bottom) control and ELVS mice before (left) and after (right) 5 min bath application of 20 mM 4-aminopyridine (4-AP). **b-e**) Delay time to first action potential after current injection onset present in putative LC neurons in response to 1 sec current injection steps from +70 to +100 pA before (**b**, **d**) and after 4-AP application (**c**, **e**) for male (**b**, **c**) and female (**d**,**e**; control: U = 40.0 − 50.0, *p* = 0.023 − 0.047; ELVS: U = 64.0 - 80.0, *p* = 0.47 − 0.84) control (blue) and ELVS (gold) mice with inter-step membrane potential maintained near -60 mV. (**f**) Delay time to first action potential after current injection onset present in putative LC neurons in response to 70 pA current injection (1 sec) for male (control: *W* = 80, *p* = 0.0024; ELVS: *W* = 24, *p* = 0.38) and female (control: *W* = 79, *p* = 0.010; ELVS: *W* = 74, *p* = 0.0015) mice. All group data are presented as mean ± SEM (bars or shading) for each group, with connected circles representing paired data from individual neurons. Three-way ANOVAs and repeated measures ANOVAs were used for group comparisons, followed by a Mann-Whitney U test for single pairwise comparisons to assess stress effects or Wilcoxon Rank-Sum for paired 4-AP

928   effects. n = 11-14 neurons from N=3 mice (7-8 weeks old) for each group. *$p < 0.05$, **$p < 0.01$

929   for each indicated pairwise comparison.

931   *Figure 6*

933
Figure 6. Norepinephrine transporter blockade ameliorates ELVS-induced reduced

sucrose preference and increased exploratory activity in female mice. Performance of

control and ELVS mice after oral consumption of vehicle (apple juice) or the norepinephrine

transporter inhibitor reboxetine (2 mg/kg) in the sucrose preference (**a**, **b**; ELVS female vehicle

vs. reboxetine: $p = 0.0001$; Control vs. ELVS female: $p < 0.0001$; ELVS male vs. female: $p =$

0.010), open field (**c**, **d**; control vehicle vs. reboxetine: U = 79.0, $p = 0.18$; ELVS vehicle vs.

reboxetine: U = 52.0, $p = 0.011$), or y-maze (**e**, **f**) assays. All group data are presented as mean

± SEM (bars) for each group, with circles representing data from individual mice. Three- or two-

way ANOVAs were used for group comparisons, followed by a Bonferroni posthoc or Mann-

Whitney U test for pairwise comparisons to assess effects of oral reboxetine treatment on group

performance. N = 6-10 mice (7-9 weeks old) for each group and sex. *$p < 0.05$, ***$p < 0.001$ for

pairwise comparisons.