Local tree cover predicts mosquito species richness and disease vector presence in a

tropical countryside landscape

Johannah E. Farner[1], Meghan Howard[1], Jeffrey R. Smith[2], Christopher B. Anderson[3], Erin A.

Mordecai[1]

[1]Department of Biology, Stanford University, Stanford, CA, USA; [2]Department of Ecology and

Evolution, Princeton University, Princeton, NJ, USA; [3]Planet Labs, San Francisco, CA, USA.

December 2023

Abstract

Land use change is an important driver of both biodiversity loss and zoonotic disease transmission in tropical countryside landscapes. Developing solutions for protecting biodiversity, public health, and livelihoods in working landscapes requires understanding the spatial scales at which habitat characteristics such as land cover shape biodiversity, especially for arthropods that transmit pathogens. A growing body of evidence shows that species richness for many taxa correlates with tree cover at small spatial scales of <100 m, indicating that local tree cover management is a promising conservation tool. To investigate whether mosquito species richness, community composition, and presence of specific disease vector species respond to tree cover— and if so, whether at spatial scales similar to other taxa—we surveyed mosquito communities along a tree cover gradient and across agricultural, residential, and forested land uses in rural southern Costa Rica. We found that tree cover was both positively correlated with mosquito species richness and negatively correlated with the presence of the common invasive dengue vector *Aedes albopictus*, particularly at small spatial scales of 80 – 200m. Beyond tree cover, land use type predicted community composition and *Ae. albopictus* presence, but not species richness. The results suggest that preservation and expansion of tree cover at local scales can protect biodiversity for a wide range of taxa and also confer protection against disease vector occurrence.

Introduction

Although land use change is a leading cause of loss and endangerment of both biodiversity

and ecosystem services (e.g., Sala, 2000; Pereira *et al.*, 2010; Maxwell *et al.*, 2016; Giam, 2017),

human-dominated landscapes also have high potential and a pressing need to contribute to

conservation (Norris, 2008; Chazdon *et al.*, 2009). Countryside landscapes characterized by

natural habitat remnants patchworked with human residential and agricultural infrastructure are

globally dominant, and thus represent both an exceptional threat to and exceptional opportunity

for conservation (Norris, 2008; Maxwell *et al.*, 2016). A critical requirement for leveraging the

conservation potential of countrysides is understanding which landscape features help to retain

biodiversity and biodiversity-associated ecosystem services, such as crop pollination and pest

control (Frishkoff *et al.*, 2019). Percent tree cover at small spatial scales of <100m has emerged

as a reliable predictor of biodiversity for taxa in Latin American tropical countryside landscapes

including birds, non-flying mammals, and bats (Mendenhall *et al.*, 2016). These findings suggest

that local tree cover management holds promise as a practicable conservation tool, but major

knowledge gaps remain surrounding how local tree cover relates to biodiversity and ecosystem

services. A key open question relevant to biodiversity conservation is the extent to which local

tree cover correlates with species richness across taxa, particularly for invertebrates.

Additionally, because tree cover management may influence human – wildlife contact patterns,

developing best practices for this conservation tool requires understanding how tree cover in

working landscapes relates to vector-borne and zoonotic disease risk.

The spatial scales at which mosquito communities (family Culicidae) respond to tree cover

present a knowledge gap relevant to both biodiversity and human health. In addition to playing

roles as prey, predators, and detritivores in aquatic and terrestrial food webs (Addicott, 1974;

Heard, 1994; Daugherty, Alto and Juliano, 2000; Poulin, Lefebvre and Paz, 2010), several

mosquito species are important vectors of wildlife and human diseases including malaria,

dengue, chikungunya, Zika, yellow fever, West Nile fever, and arboviral encephalitis

(Garmendia, Van Kruiningen and French, 2001; Lemon, 2008; LaPointe, Atkinson and Samuel,

2012; World Health Organization, 2020). Mosquito community composition and abundance are

fundamental to when, where, and how widely such diseases are transmitted (reviewed in

Franklinos *et al.*, 2019). Landscape context is likely to have multifaceted effects on mosquito

community composition through its influence on biotic and abiotic conditions that affect

mosquito life cycles. Differences in temperature, habitat structural complexity, and biotic context

associated with tree cover are all likely to affect the presence and abundance of mosquito species

that vary in their thermal niches, aquatic breeding habitat requirements, and preferred groups of

vertebrates for blood meals (Laird, 1988; Gutman *et al.*, 2004; Mordecai *et al.*, 2019; Prevedello

et al., 2019). Indeed, previously documented shifts in mosquito communities across habitats

show associations between forest conversion and low mosquito biodiversity, and suggest that

high rates of land use change and active invasions by major vector species are together reshaping

mosquito communities in ways that increase disease risk (Ferraguti *et al.*, 2016; Meyer Steiger,

Ritchie and Laurance, 2016; Burkett-Cadena and Vittor, 2018; Chaves *et al.*, 2021). Determining

the spatial scales at which tree cover shapes mosquito communities is a critical next step towards

understanding how tree cover in countryside landscapes can be managed to balance biodiversity

conservation, public health, and economic needs.

The global range expansions of the dengue vectors *Aedes aegypti* and *Aedes albopictus*

exemplify ongoing rapid change to mosquito communities that may be linked to concurrent

changes in both land cover and disease transmission (Rezza, 2012; Kraemer *et al.*, 2015). In

Central America, the first suspected dengue outbreaks occurred in the 1600s; both the disease

and the historically common introduced vector *Aedes aegypti* and dengue were eradicated in the

1940s; and both vector and virus were reintroduced in the 1970s (Brathwaite Dick *et al.*, 2012).

Dengue in this region has increased dramatically in recent decades: from 378,469 cases reported

between 1990 and 2000 to 1,338,330 cases reported between 2007 and 2017 (WHO, 2021).

Concurrent with this increase, *Ae. albopictus* was introduced to and spread throughout Central

America (Benedict *et al.*, 2007). However, how *Ae. albopictus* colonization relates to dengue in

this region is unknown, because its distribution in its new range is poorly documented, and its

role in disease transmission must be disentangled from that of *Ae. aegypti* (Rezza, 2012;

Kraemer *et al.*, 2015). Globally, *Ae. albopictus* and *Ae. aegypti* are predominantly associated

with rural and urban human settlements, respectively, suggesting that *Ae. albopictus* and its

responses to tree cover may play particularly important roles in shaping disease risk in

countryside landscapes where dengue can be prevalent (Braks *et al.*, 2003; Tsuda *et al.*, 2006).

The countryside of Costa Rica is an ideal system in which to study relationships between

land cover, mosquito community characteristics, and disease vector occurrence because this

system is the subject of intensive long-term research into links between landscape context,

biodiversity, and ecosystem services (Burkett-Cadena and Vittor, 2018). This presents a unique

opportunity to improve scientific understanding of mosquito community assembly, and resulting

disease risk, in comparison with a rich understanding of other taxonomic diversity and ecosystem

services. This is particularly relevant to this region because mosquito-borne diseases, including

dengue virus, impose a substantial public health burden. *Aedes aegypti* is considered the primary

dengue vector in Costa Rica, but the ongoing, patchily described invasion by the globally

important vector species *Aedes albopictus* is potentially reshaping disease risk (Troyo *et al.*,

2006; Calderon-Arguedas *et al.*, 2010; Calderón-Arguedas *et al.*, 2015; Rojas-Araya *et al.*,

2017). Finally, a pioneering national Payment for Ecosystem Services program offers a well-

established venue for translating research findings into land management policy, and evidence

for additional ecological and human health benefits of local tree cover may help motivate

enforcement of an existing law that protects riparian buffer zones up to 50m around rivers, which

is expected to increase access to clean water for vulnerable populations, and pertains to a spatial

scale at which tree cover has previously been shown to protect biodiversity for other taxa

(Sánchez-Azofeifa *et al.*, 2007; Langhans *et al.*, 2022).

Here, we combine field observations of mosquito community composition in forested,

agricultural, and residential settings in a rural area of southern Costa Rica with remotely-sensed

land cover data in order to ask: (Q1) Does mosquito species richness decline with declining tree

cover, consistent with patterns previously observed for a wide range of taxa in a tropical

countryside landscape? (Q2) Is the presence of *Aedes albopictus*, a key mosquito vector of

human disease, negatively correlated with tree cover? (Q3) How does the spatial scale at which

tree cover is observed affect the answers to (Q1) and (Q2)? We additionally ask (Q4) How do

mosquito communities respond to environmental gradients versus land use types? Based on

evidence that species richness for birds, non-flying mammals, bats, reptiles, and amphibians, and

plants increases with tree cover (Mendenhall *et al.*, 2014, 2016), we hypothesize that as local tree

cover increases, (H1) mosquito species richness increases, (H2) the presence of human-

associated *Aedes albopictus* decreases and (H3) these relationships are strongest at spatial scales

of <100 m. We additionally hypothesize that (H4) shifts in mosquito community composition

along tree cover gradients are reflected in distinct forest and residential mosquito species

assemblages that are bridged by intermediate agricultural assemblages. By improving scientific

understanding of how landscape context influences mosquito community assembly in Costa

Rica, the results of this study contribute to a more general understanding of links between land

cover, biodiversity, and ecosystem services through the lens of entomological factors that

influence disease risk.

Methods

Study area

This study was conducted in the cantons of Coto Brus, Corredores, and Golfito (8°43′14″N,

82°57′20″W), located in the southern Puntarenas region of Costa Rica along the border with

Panama (Figure 1). The region ranges from coastal lowland tropical rainforest (0 m above sea

level) to high elevation cloud forest (1500 m above sea level) and has distinct wet and dry

seasons. The study area is predominately composed of rural communities surrounded by

agriculture interspersed with forest patches, and also includes the protected Las Cruces forest

reserve. Dengue reintroduction in Costa Rica was first reported from the Puntarenas region in

1993, where the disease has since been endemic (World Health Organization, 1994; Troyo *et al.*,

2009). *Ae. aegypti* is common and *Ae. albopictus* has a growing presence in this area (Troyo *et*

al., 2008; Rojas-Araya *et al.*, 2017).

Figure 1. Tree cover and land use varied across study sites in Costa Rica. *Darker green indicates higher percent tree cover, remotely observed at 30 m resolution. Inset shows study sites, categorized by land use type. Blue, green, and yellow points denote agricultural, forest, and residential land uses, respectively.*

Study sites

With landowner permission, we accessed 37 sites located in San Vito, Sabalito, Copa Buena,

Ciudad Neilly, and Pavones (Figure 1). The sites represented three broad land use classes that

were determined on-site by the survey team: residential (N = 17), agricultural (N = 12), and

forest (N = 8) (Figure 1). Residential sites included urban and peri-urban areas; agricultural sites

included coffee plantations, an oil palm plantation, a pine plantation, one mixed agricultural

field, and one pasture; forested sites included primary and secondary forest edges, interiors, and

fragments.

Environmental variables

To quantify percent tree cover, we used a 30 m resolution map of tree cover in Costa Rica

created by Echeverri et. al (2022) from multi-sensor satellite observations and fine-scale tree

cover maps (Figure 1). From this map, we calculated percent tree cover at different spatial scales

surrounding each site using the R package "raster" (Hijmans *et al.*, 2015). Specifically, we began

by calculating percent tree cover within a radius of 30 m; we then increased this radius by

increments of 10 m up to 200 m, and by increments of 50 m for radii between 200 m and 1000 m

(following Mendenhall *et. al*, 2014). To account for mosquito interspecific variation in

sensitivity to thermal conditions (Mordecai *et al.,* 2019), which could affect observed

relationships between land cover and mosquito community characteristics along the 1500 m

elevational gradient surveyed, we additionally extracted mean annual temperature data from

1970 – 2000 for each study site from the WorldClim 1 km^2 resolution mean annual temperature

dataset (Fick and Hijmans, 2017).

Sample collection

We trapped mosquitoes at all sites twice between June 19 and August 9, 2017, excepting

the sites in Pavones, which were trapped once. At each site, we placed a total of four traps

overnight for a 12-16 hour period within an area of 30 m in radius: one unlighted CDC trap

baited with carbon dioxide produced by a mixture of Fleischmann's Dry Active Yeast, household

refined sugar, and water; one BG Sentinel baited with a BG-Lure and octanol; and two BG-GAT

traps furnished with yellow sticky cards, corn oil, and a mixture of water and local leaf litter.

Trap locations within sites were chosen per BioGents recommendations, and square metal frames

covered in large black plastic bags were placed over BG Sentinel and BG-GAT traps for

protection from rainfall. To supplement the overnight trapping, consistent members of the field

team carried out 20 minutes of direct aspiration at each site during each trapping session.

Mosquito identification

Trained personnel identified, sexed, and counted any *Ae. albopictus* and *Ae. aegypti*

mosquitoes collected at each site. All other mosquitoes were counted, stored, and transported to

Stanford University for molecular identification. We extracted and amplified DNA from the

mitochondrial CO1 gene from pooled samples of mosquitoes from each site using MyTaq

RedMix (Meridian Bioscience, Cincinnati, OH), following the protocol provided by the

manufacturer. Amplified DNA was sequenced via Illumina next-generation sequencing, with

samples containing *Aedes albopictus* and *Culex tarsalis* DNA as positive controls. We used the

R package "dada2" to filter and trim the DNA sequences to 473 bp, with a minimum overlap of

20 bases and a maximum of 5 expected errors (Callahan *et al.*, 2016). We estimated taxonomic

placement for the sequenced mosquitoes by using the R packages "Biostrings" and

"DECIPHER" to group DNA sequences into operational taxonomic units (OTUs, henceforth

referred to as species) of 97% sequence similarity, and comparing representative sequences for

each species to the BOLD and GenBank database records (Altschul *et al.*, 1990; Ratnasingham

and Hebert, 2007; Wright, 2016; Pagès *et al.*, 2022). Species were identified based on top

matches with sequence similarity ≥ 97% (Hardulak *et al.*, 2020). When sequence similarity to the

top match was < 97%, a higher level of taxonomic identification (e.g., genus) was assigned based

on placement within a phylogenetic tree of the BOLD database sequences.

Statistical analyses

We described mosquito communities in terms of species richness and species

composition by combining presence data from the morphologically identified *Aedes* data and the

sequencing data. To quantify relationships between species richness and percent tree cover, we

used generalized linear models (GLMs) with negative binomial error corrections for

overdispersion and mean-centered independent variables. To assess the spatial scales at which

tree cover best predicted species richness, we compared AIC values for GLMs that included

percent tree cover surrounding each site calculated at radii ranging from 30 m to 1000 m. We

used binomial logistic regression in order to analyze relationships between *Ae. albopictus* disease

vector presence/absence and percent tree cover across spatial scales. For the 1000 m spatial scale

where climate data were available, we additionally assessed the relative influence of mean

annual temperature on species richness and *Aedes* vector presence with GLMs including mean

annual temperature and its interaction with tree cover.

To compare species richness and *Aedes* vector presence between forest, agricultural, and

residential land uses, we used Kruskal-Wallis tests with Bonferroni p-value adjustments to

account for multiple comparisons.

To compare species composition among land use types and along environmental

gradients, we first calculated the Jaccard coefficient of community similarity for each pair of

sites for use in statistical tests and ordination. We then tested for differences in community

similarity among land uses with permutational analysis of variation (PERMANOVA), first for all

land use types, and then with pairwise adonis functions. Because PERMANOVA is sensitive to

heterogeneity in dispersion among groups (Anderson and Walsh, 2013), we additionally tested

whether dispersion differed among land use types using Tukey's Honest Significant Differences

method with betadisper() calculations of group average distances to the median. To visualize

community similarity across land uses, we used non-metric multidimensional scaling (NMDS).

All the above analyses of compositional similarity among land uses were run using the R

package "vegan" (Oksanen *et al.*, 2013); for the pairwise PERMANOVA analyses, we used the

R package "ecole" which provides wrapper functions for "vegan" (Smith, 2022). Finally, to

quantify compositional turnover along environmental gradients of tree cover, mean temperature,

elevation, and geographic distance, we used the R package "gdm" for generalized dissimilarity

modeling (GDM), a form of nonlinear matrix regression that is robust to collinearity (Fitzpatrick

et al., 2022). As above, we compared GDM models that incorporated tree cover at radii ranging

from 30 – 1000 m to identify the spatial scale at which tree cover best explained compositional

turnover.

All analyses were performed in R version 4.2.1. In addition to the R packages cited

above, we used the packages "tidyverse" (Wickham *et al.*, 2019), "cowplot" (Wilke *et al.*, 2019),

"MASS" (Ripley *et al.*, 2013), "interactions" (Long, 2019), "gridExtra" (Auguie *et al.*, 2017),

and "reshape2" (Wickham, 2007) for data analysis and figure generation.

Results

Across sites, tree cover ranged from 0% to 100% (mean = 27.9%, SD = 34.8%) at the

smallest spatial scale we considered, a 30 m radius. Surrounding tree cover within a 1000 m

radius, the largest spatial scale considered, ranged from 8.2% to 74.8% (mean = 33.4%, SD =

20.3%). The lowest and highest site elevations were 12 m and 1451 m above sea level,

respectively (mean = 776 m, SD = 473.2 m). Mean annual temperature ranged from 18.8 to 26.4

℃ (mean = 22.6 ℃, SD = 2.3 ℃).

A total of 1,283 mosquitoes representing 48 species in 13 genera were collected from 34

sites (Figure 2). The number of mosquitoes collected at a site ranged from 0 to 244 (mean = 35,

SD = 64). Of these, 108 individuals from 12 residential and four agricultural sites were

morphologically identified as *Ae. albopictus*, and 5 individuals from single sites within each land

use category were identified as *Ae. aegypti*. *Ae. albopictus* DNA was detected in the pooled

samples of molecularly identified mosquitoes from 7 sites where *Ae. albopictus* individuals were

also morphologically identified. The five most common species were *Culex quinquefasciatus*,

Ae. albopictus, *Cx. nigripalpus*, *Wyeomyia adelpha/Guatemala,* and *Limatus durhamii* (Figure 2,

Table S1).

Figure 2. The commonness of 48 observed mosquito species varied both overall and

among land use types. *Blue bars show the number of presences observed for each species in*

agricultural sites, green bars show forest sites, and yellow bars show residential sites. Asterisks

indicate species known to vector human diseases.

Site-level species richness ranged from one to 19 (mean = 5.2, SD = 4.1). Overall species

counts for forest, agricultural, and residential land uses were 33, 29, and 21, respectively. Ten

species (21%) were observed in all three land uses. Nineteen species (40%) were shared among

forest and agricultural land uses, 13 species (27%) were shared among agricultural and

residential land uses, and twelve species (25%) were shared among forest and residential land

uses (Figure 2, Table S1). Eleven species (23%) were found only in forested settings, six species

(13%) were found only in agricultural settings, and six species (13%) were found only in

residential settings (Figure 2, Table S1). Two species, *Ae. albopictus* and *Culex*

quinquefasciatus, were common (observed at >50% of sites) in residential settings, no species

were common in agricultural settings, and three species—*Culex nigripalpus*, *Wyeomyia*

complosa, and *Wyeomyia adelpha/guatemala*—were common in forested settings (Table S1).

At least five of the mosquito species observed are known vectors of human diseases.

Three of these— the dengue and chikungunya virus vector *Ae. albopictus* (present at 15 sites)

and the St. Louis Encephalitis virus vectors *Cx. quinquefasciatus* (present at 17 sites) and *Cx.*

nigripalpus (present at 13 sites)—were the three most frequently observed species (Figure 2,

Table S1) (Reisen, 2003; Simmons *et al.*, 2012). Rarely observed vector species included the

dengue, chikungunya, yellow fever, and Zika virus vector *Ae. aegypti* (present at three sites

spanning all three land use types) and the malaria vector *Anopheles albimanus* (present at one

agricultural site) (Figure 2, Table S1) (Zimmerman, 1992; Simmons *et al.*, 2012). In contrast to

Ae. aegypti, Cx. nigripalpus, and *Cx. quinquefasciatus*, which were observed in all land use

types, *Ae. albopictus* was observed only in residential and agricultural settings associated with

intensive human modification (Figure 2, Table S1).

Mosquito species richness was explained by tree cover, but not by land use type.

Comparisons of GLMs using tree cover calculated for radii ranging between 30 m and 1000 m

surrounding each site indicated that species richness was positively correlated with tree cover at

radii between 80 m and 600 m, and tree cover at a 250 m radius had the largest effect size

(estimated effect = 1.29 x 10^{-2}, SE = 4.75 x 10^{-3}, z-value = 2.71, p-value = 6.65 x 10^{-3}) (Figure

3a, Table S2). At the 1000 m spatial scale where both tree cover and climate data were available,

the interaction between tree cover and mean annual temperature had a significant effect on

species richness (estimated effect = 7.87 x 10^{-3}, SE = 3.65 x 10^{-3}, z-value = -2.16, p-value = 3.11

x 10^{-2} (Table S3). Specifically, at high temperatures, species richness was low even when tree

cover was high (Figure S1). By contrast, Kruskal-Wallis test results indicated that species

richness did not differ significantly among forested, agricultural, and residential sites (chi-

squared = 2.60, df = 2, p-value = 0.28) (Figure 3b). Notably, the highest species richness was

observed at a site in the Coto Brus forest reserve (Figure 3b).

Figure 3. Species richness is correlated with tree cover surrounding survey sites for radii
between 80 m and 600 m. *(A) The estimated effect of surrounding tree cover calculated across*
spatial scales on species richness. Radii where the relationship between tree cover and species
richness is significant (p < 0.05) are shown in black. Values above and below the dashed line
are positive and negative, respectively. (B) Species richness increases with percent tree cover at
the 250 m radius: the scale identified as having the strongest effect. Land use (colored points)
and species richness are not significantly correlated. The site with the highest species richness
and the site with the highest surrounding tree cover were both located in the Las Cruces forest
reserve (arrows). In both panels, gray shading shows +/- 1 SE.
322

From the *Aedes* disease vector survey, we present results only for *Ae. albopictus* because

observations of *Ae. aegypti* were insufficient for statistical analysis. Both tree cover and land use

type predicted *Ae. albopictus* presence. Comparisons of GLMs using tree cover calculated for

radii ranging between 30 m and 1000 m surrounding each site indicated that *Ae. albopictus*

presence was negatively correlated with tree cover at radii between 50 m and 200 m, and was

best explained by tree cover at a 110 m radius (estimated effect = -4.23 x 10^{-2}, SE = 1.96 x 10^{-2},

z-value = -2.16, p-value = 3.08 x 10^{-2}) (Figure 4a, Table S3). At the 1000 m spatial scale where

we additionally assessed the influence of climate, *Ae. albopictus* presence was negatively

correlated with tree cover and positively correlated with temperature (tree cover estimated effect

= -7.76 x 10^{-2}, SE = 3.74 x 10^{-2}, z-value = -2.07, p-value = 3.81 x 10^{-2}; mean annual temperature

estimated effect = 6.62 x 10^{-1}, SE = 3.35 x 10^{-1}, z-value = 1.97, p-value = 4.814 x 10^{-2}) (Table

S5). Land use type also predicted *Ae. albopictus* presence (Kruskal-Wallis chi-squared = 9.58, p-

value = 8.311 x 10^{-3}). Specifically, *Ae. albopictus* was significantly more likely to be observed in

residential settings (present at 13/17 sites) than in forested settings (present at 0/8 sites) (Kruskal-

Wallace chi-squared = 9.03, Bonferroni-adjusted p-value = 7.98 x 10^{-3}), and its presence in

agricultural settings (present at 4/12 sites) did not differ significantly compared to either

residential (Kruskal-Wallace chi-squared = 2.41, adjusted p-value = 3.6 x 10^{-1}) or forested

(Kruskal-Wallace chi-squared = 3.04, adjusted p-value = 2.43 x 10^{-1}) settings. Fourteen of the

15 sites where *Ae. albopictus* was present were surrounded by less than 25% tree cover within a

50 m radius. The only site within a pine plantation was a clear outlier, where *Ae. albopictus* was

present under 85% tree cover (Figure 4B).

Figure 4. Aedes albopictus is most likely to be found at low tree cover levels and in
residential settings. *(A) The estimated effects of tree cover across spatial scales on Ae.*
albopictus presence, where black lines indicate statistically significant (p < 0.05) effects and
pink lines indicate non-significant (p > 0.05) relationships. Values above and below the dashed
line are positive and negative, respectively. (B) Land use type (colored points), percent tree
cover, and Ae. albopictus presence/absence at the highest-significance 110 m radius. In both
panels, gray shading shows +/- 1 SE.

In contrast to species richness, community composition and dispersion were predicted by

land use type. PERMANOVA results comparing all three land uses showed that land use

significantly affected community composition (Sum of squares = 2.113, R^2 = 0.16, F-value =

3.03, p-value = 0.001), and pairwise PERMANOVAs showed that agricultural and forest

mosquito communities differed significantly from residential communities (Table 1). Wider

dispersion among agricultural compared to residential mosquito communities (average distance

to the median: agriculture = 0.612, forest = 0.572, residential = 0.497; Tukey test adjusted p-

values: residential – agricultural = 0.0422, residential – forested = 0.303, forested – agricultural

= 0.736) likely contributed to the community dissimilarity detected between these land uses

(Anderson and Walsh, 2013) (Figure 5). NMDS visualization of communities grouped by land

use type suggests that more variable agricultural communities bridge relatively distinct forest and

residential communities (NMDS stress = 0.12), in agreement with the statistical test results

(Figure 5, Table 1). The wider variation among agricultural sites is also evident from the species

observation table: no single species was observed at more than 1/3 of all agricultural sites,

whereas *Ae. albopictus* and Unidentified Culicidae 1 were both observed at >70% of residential

sites, and *Culex nigripalpus*, *Wyeomyia adelpha/guatemala*, and *Wyeomyia complosa* were each

observed at >60% of forested sites (Table S1).

Figure 5. Distinct mosquito communities were observed in forest and residential land uses,
while communities in agricultural settings were more variable. *NMDS ordination*
visualization groups sites by community similarity, with colors indicating land use types. Each
point represents the community at one study site, and the distance between points is smaller for
more similar communities. Ellipses show 95% confidence intervals for ordination of agriculture,
forest, and residential mosquito communities.

Finally, generalized dissimilarity modeling (GDM) indicated that environmental

gradients explained little of the species turnover among sites. Among the spatial scales for which

tree cover was calculated, the model using the 50m radius explained the highest amount of

species turnover among sites (Table S6). The model that included mean annual temperature,

geographic distance, and tree cover at the 50m radius explained 7% of species turnover among

sites. Elevation showed no relationship with species turnover. Whereas increasing tree cover was

associated with a consistent increase in community turnover, increasing temperature was

associated with a steep increase in community turnover up to a plateau around 22 ℃ (Figure 6).

Figure 6. Generalized dissimilarity modelling (GDM) of community dissimilarity indicated that (A) percent tree cover and (B) mean annual temperature explained 7% of deviation from the null. (C) Geographic distance and (D) elevation did not significantly contribute to community turnover. In each panel, the x-axis shows the environmental gradient and the y-axis shows the amount of compositional turnover, measured as partial ecological distance. The maximum height the spline reaches on the y-axis indicates the total amount of compositional turnover the gradient is associated with, and the slope shows how the rate of compositional turnover varies along the environmental gradient. The difference in height between any two points along the I-spline corresponds to the modeled contribution of that predictor variable to the difference between those points. Grey shading shows +/- 1 standard deviation when 70% of sites are sampled 10 times.

Discussion

We found that local tree cover, but not land use (residential, agricultural, or forest),

predicted mosquito species richness: more diverse communities occurred at higher tree cover

(supporting H1) (Figure 3, Table S2). By contrast, community composition was more predictable

for forested and residential land uses, and more variable among agricultural sites (supporting H4)

(Figure 5, Table 1). Environmental gradients of tree cover, climate, and geographic distance

explained 7.2% of species turnover among sites (Figure 6, Table S6). *Ae. albopictus* presence

varied significantly with both tree cover and land use, but in the opposite direction from

mosquito diversity: *Ae. albopictus* occurrence probability increased with lower tree cover and in

residential compared to forested sites (supporting H2) (with intermediate probability in

agricultural sites; Figure 4, Table S4). Overall, our results add to support from other taxa for the

value of both natural and semi-natural habitat in sustaining biodiversity and ecosystem

services—here, in the form of protection against an invasive mosquito that is a major vector of

human disease.

Mosquito diversity

The spatial scales at which tree cover predicted mosquito species richness were small,

and comparable to previous findings for other taxa in the same study area, highlighting the

disproportionately positive impact of small patches of trees on biodiversity. The radius at which

tree cover best predicted species richness was 250 m; by comparison, biodiversity was correlated

with tree cover at small spatial scales for non-flying mammals (70 m), bats (50 - 60 m), birds (30

m), reptiles (50 m), and amphibians (80 m) in the same region of Costa Rica (Mendenhall *et al.,*

2014, 2016; Frank *et al.,* 2017). Although the radius at which mosquito species richness

responded most strongly to tree cover was larger compared to previously studied taxa, the spatial

scale remained local, and significant effects of tree cover were found at radii as small as 80 m

(Figure 3). Our observation of a positive relationship between species richness and tree cover

aligns with those of many other studies of mosquito diversity along land cover gradients in

locations including Latin America, Asia, and Europe (e.g., Johnson, Gómez and Pinedo-

Vasquez, 2008; Thongsripong *et al.*, 2013; Ferraguti *et al.*, 2016; Chaverri *et al.*, 2018), and

clarifies the spatial scales at which tree cover shapes mosquito communities.

Our observation that mosquito community composition was distinct among different land

uses is consistent with patterns observed both for other taxa in this system, and for mosquitoes in

other regions (Mendenhall *et al.*, 2014, 2016; Meyer Steiger, Ritchie and Laurance, 2016; da

Silva Pessoa Vieira *et al.*, 2022). In agricultural settings, relatively high species richness and

community similarity with forested settings support the argument that farmlands can contribute

substantially to biodiversity conservation (Figure 5, Table 1) (e.g., Norris, 2008). However, the

high proportion of species unique to forest habitats and the high species richness observed inside

the large Las Cruces forest reserve also reaffirm the singular importance of forests and protected

areas as refugia for biodiversity (Coetzee, Gaston and Chown, 2014; Mendenhall *et al.*, 2016)

(Figure 3, Table S1). Additionally, the compositional variability among agricultural settings and

the close community resemblance between some agricultural and residential sites indicate a need

for additional research on how mosquito communities respond to specific land uses, crop

assemblages, or management practices that can result in similar levels of tree cover (Figure 5,

Table 1). For example, organic farming methods are associated with higher arthropod diversity

globally compared to conventional methods, and Kenyan ricelands that rely on natural rather

than artificial irrigation have higher mosquito species richness Kenya (Muturi *et al.*, 2006;

Lichtenberg *et al.*, 2017). Costa Rican croplands that are less intensively farmed may similarly

support greater mosquito biodiversity, a pattern already observed for bird species richness

(Hendershot *et al.*, 2020).

457   Environmental gradients of tree cover and temperature shaped species turnover, but

explained only 7% of variance in community composition, suggesting that additional habitat

characteristics may play important roles in determining species composition. Such factors might

include local microclimates, differences among types of tree cover (e.g., agricultural types,

primary versus secondary forest), and/or the presence of vertebrate hosts preferred by different

mosquito species. Differences in species abundances and community evenness, which were not

quantified here, might also respond more strongly to gradual environmental change than the

identities of the species present. However, our result that land use predicts species composition,

while land cover predicts diversity, aligns with patterns of abundance-based Dipteran and

Culicid diversity observed in the tropical Australian countryside (Smith and Mayfield, 2015;

Meyer Steiger, Ritchie and Laurance, 2016).

Disease vectors

470   The most frequently observed disease vector, *Ae. albopictus*, was more likely to be

observed in sites with lower surrounding tree cover and agricultural or residential land uses,

suggesting that rural landscapes with more forest and tree cover may be more resistant to

invasion by this species. These observations align with this species' well-established preferences

for taking blood meals from humans and livestock (Niebylski *et al.*, 1994; Richards *et al.*, 2006),

and its association with rural, agricultural, suburban, and/or deforested settings in the Americas,

Asia, and Africa (Gilotra, Rozeboom and Bhattacharya, 1967; Braks *et al.*, 2003; Young *et al.*,

2017; Câmara *et al.*, 2020; Canelas *et al.*, 2023). The 40 – 200 m distances where tree cover

negatively affected *Ae. albopictus* presence fell within the 80 – 600 m range where tree cover

positively affected species richness, suggesting promise for local tree cover management as a

means of supporting both public health and biodiversity conservation. Protection against

mosquito disease vectors conferred by tree cover may extend beyond *Ae. albopictus* to include at

least 16 other significant vectors of human diseases that are favored by deforestation, including

Ae. aegypti, multiple Anopheline malaria vectors, and the Amazonian malaria vector

Nyssorhynchus darlingi (Burkett-Cadena and Vittor, 2018; Chaves *et al.*, 2021).

Our finding that *Ae. albopictus* was associated with, but inconsistently observed in,

agricultural settings (present in 33% of agriculture sites) (Figure 4, Table S1) reinforces that

agricultural lands have the potential to either harbor or resist invasive species, and suggests

vector associations with agricultural subtypes as a key future research direction. In our survey,

factors that differentiated the high tree-cover pine plantation and two of the six surveyed coffee

plantations as suitable habitat for *Ae. albopictus* are of particular interest (Figure 4).

Understanding vector responses to agriculture is particularly important because this land use is

the most likely candidate for local tree cover management in Costa Rica due to its spatial extent,

the established Payment for Ecosystem Services program infrastructure for incentivizing

landowner forest retention and tree planting, and a previous finding that urban tree cover is

correlated with dengue incidence in this region (Sánchez-Azofeifa *et al.*, 2007; Troyo *et al.*,

2009).

However, further research is needed to understand how tree cover management could be

applied for such public health protections. For example, future studies should aim to capture the

seasonality of vector presence and abundance along tree cover gradients (Calderón-Arguedas *et*

al., 2008; Troyo *et al.*, 2008; Romero-Vega *et al.*, 2023); assess relationships between vector

presence and other factors in habitats that have similar tree cover but differ in aspects such as

crop type or tree cover geometry; and use reforestation efforts or experimental tree cover

additions at relevant spatial scales to test mosquito community responses. In the study region,

tree cover responses of *Ae. aegypti* and *An. albimanus* require clarification, as we observed these

regionally important vectors too rarely for statistical analysis, and saw unexpected rarity and

apparent habitat generalism in the canonically widespread and urban vector *Ae. aegypti* (Figure

2, Table S1) (Troyo *et al.*, 2006; Cáceres *et al.*, 2012). Additionally, the potential disease control

benefits of local tree cover should be tested with human pathogen surveillance in field-captured

mosquitoes from different environments, and by comparison of disease case data to mosquito

community data. In addition to dengue, Zika, and malaria, St. Louis Encephalitis Virus should be

considered for surveillance in rural areas where humans and animals live in close proximity,

because two potential vectors—*Cx. quinquefasciatus* and *Cx. nigripalpus*—were common, and

this unmonitored disease is already widespread among both domesticated and wild animal hosts

in Costa Rica (Medlin *et al.*, 2016; A. Chaves *et al.*, 2021; Piche-Ovares *et al.*, 2023).

Conclusions

Overall, our results join a large body of evidence for the importance of local tree cover to

maintaining both biodiversity and ecosystem services, adding prevention of disease vector

proliferation to a list of benefits including water quality, crop pollination, carbon sequestration,

and soil health, among others (Millennium Ecosystem Assessment, 2005; Jose, 2009; Barrios *et*

al., 2018; Frishkoff *et al.*, 2019). Our findings follow patterns observed repeatedly across the

globe associating tree cover with higher mosquito species richness and lower disease vector

presence, suggesting broad applicability of local tree cover management as a combined vector

control and biodiversity conservation method. Further, we shed light on the contexts in which

tree cover both increases mosquito species richness and decreases *Ae. albopictus* presence,

namely at spatial scales of 80 – 200m (Figures 2,3; Tables S2, S4). We also find that at larger

spatial scales of 1000m, warm mean annual temperatures increase habitat suitability for *Ae.*

albopictus and limit tree cover contributions to species richness, but note that other factors that

covary with climate across the study region may contribute to this result (Figure S1, Tables S3,

S5). Although the specific mechanisms and characteristics by which tree cover inhibits disease

vectors remain unclear, the regularity with which simple measures of local tree cover correlate

with both biodiversity and ecosystem services represents a practicable solution for fostering both

human and ecosystem health in countryside landscapes.

Acknowledgements

We thank the many landowners who generously provided access to research sites; Henry Sandi

Amador, Esteban Calderon, and Erick Barrantes of the Organization for Tropical Studies for

field assistance; Jamieson O'Marr for field and laboratory assistance; Gretchen Daily for advice

on the project design; and Mordecai lab members for laboratory assistance and helpful feedback

throughout all stages of the project. This work was funded by National Institutes of Health 331

(R35GM133439, R01AI168097, R01AI102918), the National Science Foundation

(DEB3322011147, with Fogarty International Center), the Stanford Center for Innovation in

Global Health, King Center on Global Development, Woods Institute for the Environment, the

Ward Wilson Woods Jr Environmental Study Fund, the Lewis and Clark Fund from the

References

Addicott, J.F. (1974) 'Predation and prey community structure: An experimental study of the

effect of mosquito larvae on the protozoan communities of pitcher plants', *Ecology*, 55(3), pp.

475–492. Available at: https://doi.org/10.2307/1935141.

Altschul, S.F. *et al.* (1990) 'Basic local alignment search tool', *Journal of Molecular Biology*,

215(3), pp. 403–410. Available at: https://doi.org/10.1016/S0022-2836(05)80360-2.

Anderson, M.J. and Walsh, D.C.I. (2013) 'PERMANOVA, ANOSIM, and the Mantel test in the

face of heterogeneous dispersions: What null hypothesis are you testing?', *Ecological*

Monographs, 83(4), pp. 557–574. Available at: https://doi.org/10.1890/12-2010.1.

Auguie, B., Antonov, A. and Auguie, M.B., 2017. 'Package 'gridExtra'', *Miscellaneous*

Functions for "Grid" Graphics.

Barrios, E. *et al.* (2018) 'Contribution of trees to the conservation of biodiversity and ecosystem

services in agricultural landscapes', *International Journal of Biodiversity Science, Ecosystem*

Services & Management, 14(1), pp. 1–16. Available at:

https://doi.org/10.1080/21513732.2017.1399167.

Benedict, M.Q. *et al.* (2007) 'Spread of the tiger: Global risk of invasion by the mosquito *Aedes*

albopictus', *Vector-Borne and Zoonotic Diseases*, 7(1), pp. 76–85. Available at:

https://doi.org/10.1089/vbz.2006.0562.

Braks, M.A.H. *et al.* (2003) 'Convergent habitat segregation of *Aedes aegypti* and *Aedes*

albopictus (Diptera: Culicidae) in Southeastern Brazil and Florida', *Journal of Medical*

Entomology, 40(6), pp. 785–794. Available at: https://doi.org/10.1603/0022-2585-40.6.785.

Brathwaite Dick, O. *et al.* (2012) 'The history of dengue outbreaks in the Americas', *The*

American Journal of Tropical Medicine and Hygiene, 87(4), pp. 584–593. Available at:

https://doi.org/10.4269/ajtmh.2012.11-0770.

Brosi, B.J., Daily, G.C. and Ehrlich, P.R. (2007) 'Bee community shifts with landscape context

in a tropical countryside', *Ecological Applications*, 17(2), pp. 418–430. Available at:

https://doi.org/10.1890/06-0029.

Burkett-Cadena, N.D. and Vittor, A.Y. (2018) 'Deforestation and vector-borne disease: Forest

conversion favors important mosquito vectors of human pathogens', *Basic and Applied Ecology*,

26, pp. 101–110. Available at: https://doi.org/10.1016/j.baae.2017.09.012.

Cáceres, L. *et al.* (2012) 'Characterization of *Plasmodium vivax* malaria transmission at the

border of Panamá and Costa Rica', *Biomedica*, 32(4), pp. 557–569.

Calderón-Arguedas, O. *et al.* (2008) 'Urban mosquito species (Diptera: Culicidae) of dengue

endemic communities in the Greater Puntarenas area, Costa Rica', *Revista de Biología Tropical*,

57(4). Available at: https://doi.org/10.15517/rbt.v57i4.5459.

Calderon-Arguedas, O. *et al.* (2010) 'Expansion of *Aedes albopictus* skull in Costa Rica', *Rev.*

Ibero-Latinoam. Parasitol., 69(2), pp. 220–222.

Calderón-Arguedas, O. *et al.* (2015) 'Dengue viruses in *Aedes albopictus* Skuse from a pineapple

plantation in Costa Rica', *Journal of Vector Ecology*, 40(1), pp. 184–186. Available at:

https://doi.org/10.1111/jvec.12149.

Callahan, B.J. *et al.* (2016) 'DADA2: High-resolution sample inference from Illumina amplicon

data', *Nature Methods*, 13(7), pp. 581–583. Available at: https://doi.org/10.1038/nmeth.3869.

Câmara, D.C.P. *et al.* (2020) 'Diversity of mosquito (Diptera: Culicidae) vectors in a

heterogeneous landscape endemic for arboviruses', *Acta Tropica*, 212, p. 105715. Available at:

https://doi.org/10.1016/j.actatropica.2020.105715.

Canelas, T. *et al.* (2023) 'Demographic and environmental factors associated with the

distribution of *Aedes albopictus* in Cameroon', *Medical and Veterinary Entomology*, 37(1), pp.

143–151. Available at: https://doi.org/10.1111/mve.12619.

Chaverri, L.G. *et al.* (2018) 'Mosquito species (Diptera: Culicidae) diversity from ovitraps in a

Mesoamerican tropical rainforest', *Journal of Medical Entomology*, 55(3), pp. 646–653.

Available at: https://doi.org/10.1093/jme/tjx254.

Chaves, A. *et al.* (2021) 'Serosurvey of nonhuman primates in Costa Rica at the human–wildlife

interface reveals high exposure to Flaviviruses', *Insects*, 12(6), p. 554. Available at:

https://doi.org/10.3390/insects12060554.

Chaves, L.S.M. *et al.* (2021) 'Anthropogenic landscape decreases mosquito biodiversity and

drives malaria vector proliferation in the Amazon rainforest', *PLOS ONE*, 16(1), p. e0245087.

Available at: https://doi.org/10.1371/journal.pone.0245087.

Chazdon, R.L. *et al.* (2009) 'Beyond reserves: A research agenda for conserving biodiversity in

human-modified tropical landscapes', *Biotropica*, 41(2), pp. 142–153. Available at:

https://doi.org/10.1111/j.1744-7429.2008.00471.x.

Coetzee, B.W.T., Gaston, K.J. and Chown, S.L. (2014) 'Local scale comparisons of biodiversity

as a test for global protected area ecological performance: A meta-analysis', *PLOS ONE*, 9(8), p.

e105824. Available at: https://doi.org/10.1371/journal.pone.0105824.

Daugherty, M.P., Alto, B.W. and Juliano, S.A. (2000) 'Invertebrate carcasses as a resource for

competing *Aedes albopictus* and *Aedes aegypti* (Diptera: Culicidae)', *Journal of Medical*

Entomology, 37(3), pp. 364–372. Available at: https://doi.org/10.1093/jmedent/37.3.364.

Echeverri, A. *et al.* (2022) 'Biodiversity and infrastructure interact to drive tourism to and within

Costa Rica', *Proceedings of the National Academy of Sciences*, 119(11), p. e2107662119.

Available at: https://doi.org/10.1073/pnas.2107662119.

Ferraguti, M. *et al.* (2016) 'Effects of landscape anthropization on mosquito community

composition and abundance', *Scientific Reports*, 6(1), p. 29002. Available at:

https://doi.org/10.1038/srep29002.

Fick, S.E. and Hijmans, R.J. (2017) 'WorldClim 2: New 1 km spatial resolution climate

surfaces for global land areas', *International journal of climatology*, 37(12), pp. 4302–4315.

Fitzpatrick, M. *et al.* (2022) 'gdm: Generalized Dissimilarity Modeling', *R package*. Available

at: https://CRAN.R-project.org/package=gdm.

Frank, H.K. *et al.* (2017) 'Phylogeny, traits, and biodiversity of a Neotropical bat assemblage:

Close relatives show similar responses to local deforestation', *The American Naturalist*, 190(2),

pp. 200–212. Available at: https://doi.org/10.1086/692534.

Franklinos, L.H.V. *et al.* (2019) 'The effect of global change on mosquito-borne disease', *The*

Lancet Infectious Diseases, 19(9), pp. e302–e312. Available at: https://doi.org/10.1016/S1473-

3099(19)30161-6.

Frishkoff, L.O. *et al.* (2019) 'Countryside biogeography: The controls of species distributions in

human-dominated landscapes', *Current Landscape Ecology Reports*, 4(2), pp. 15–30. Available

at: https://doi.org/10.1007/s40823-019-00037-5.

Garmendia, A.E., Van Kruiningen, H.J. and French, R.A. (2001) 'The West Nile virus: Its recent

emergence in North America', *Microbes and Infection*, 3(3), pp. 223–229. Available at:

https://doi.org/10.1016/S1286-4579(01)01374-0.

Giam, X. (2017) 'Global biodiversity loss from tropical deforestation', *Proceedings of the*

National Academy of Sciences, 114(23), pp. 5775–5777. Available at:

https://doi.org/10.1073/pnas.1706264114.

Gilotra, S.K., Rozeboom, L.E. and Bhattacharya, N.C. (1967) 'Observations on possible

competitive displacement between populations of *Aedes aegypti* Linnaeus and *Aedes albopictus*

Skuse in Calcutta', *Bulletin of the World Health Organization*, 37(3), pp. 437–446.

Gutman, G. *et al.* (eds) (2004) *Land Change Science: Observing, Monitoring and Understanding*

Trajectories of Change on the Earth's Surface. Dordrecht: Springer Netherlands (Remote

Sensing and Digital Image Processing). Available at: https://doi.org/10.1007/978-1-4020-2562-4.

Hardulak, L.A. *et al.* (2020) 'DNA metabarcoding for biodiversity monitoring in a national park:

Screening for invasive and pest species', *Molecular Ecology Resources*, 20(6), pp. 1542–1557.

Available at: https://doi.org/10.1111/1755-0998.13212.

Heard, S.B. (1994) 'Pitcher-plant midges and mosquitoes: A processing chain commensalism',

Ecology, 75(6), pp. 1647–1660. Available at: https://doi.org/10.2307/1939625.

Hendershot, J.N. *et al.* (2020) 'Intensive farming drives long-term shifts in avian community

composition', *Nature*, 579(7799), pp. 393–396. Available at: https://doi.org/10.1038/s41586-

020-2090-6.

Hijmans, R.J., Van Etten, J., Cheng, J., Mattiuzzi, M., Sumner, M., Greenberg, J.A., Lamigueiro,

O.P., Bevan, A., Racine, E.B., Shortridge, A. and Hijmans, M.R.J., 2015. Package 'raster'. *R*

package, *734*, p.473.

Hughes, J.B., Daily, G.C. and Ehrlich, P.R. (2002) 'Conservation of tropical forest birds in

countryside habitats', *Ecology Letters*, 5(1), pp. 121–129. Available at:

https://doi.org/10.1046/j.1461-0248.2002.00294.x.

Johnson, M.F., Gómez, A. and Pinedo-Vasquez, M. (2008) 'Land use and mosquito diversity in

the Peruvian Amazon', *Journal of Medical Entomology*, 45(6), pp. 1023–1030. Available at:

https://doi.org/10.1603/0022-2585(2008)45[1023:LUAMDI]2.0.CO;2.

Jose, S. (2009) 'Agroforestry for ecosystem services and environmental benefits: An overview',

Agroforestry Systems, 76(1), pp. 1–10. Available at: https://doi.org/10.1007/s10457-009-9229-7.

Karp, D.S. *et al.* (2013) 'Forest bolsters bird abundance, pest control and coffee yield', *Ecology*

Letters, 16(11), pp. 1339–1347. Available at: https://doi.org/10.1111/ele.12173.

Kraemer, M.U. *et al.* (2015) 'The global distribution of the arbovirus vectors *Aedes aegypti* and

Ae. albopictus', *eLife*, 4, p. e08347. Available at: https://doi.org/10.7554/eLife.08347.

Laird, M. (1988) *The natural history of larval mosquito habitats*. Academic Press Ltd, London,

UK. ISBN: 9780124340053. Available at:

http://www.cabdirect.org/cabdirect/abstract/19892056420 (Accessed: 5 July 2021).

Langhans, K.E. *et al.* (2022) 'Modeling multiple ecosystem services and beneficiaries of riparian

reforestation in Costa Rica', *Ecosystem Services*, 57, p. 101470. Available at:

https://doi.org/10.1016/j.ecoser.2022.101470.

LaPointe, D.A., Atkinson, C.T. and Samuel, M.D. (2012) 'Ecology and conservation biology of

avian malaria: Ecology of avian malaria', *Annals of the New York Academy of Sciences*, 1249(1),

pp. 211–226. Available at: https://doi.org/10.1111/j.1749-6632.2011.06431.x.

Lemon, S.M., Institute of Medicine (U.S.), and Forum on Microbial Threats (eds) (2008) *Vector-*

borne diseases - understanding the environmental, human health, and ecological connections:

workshop summary. Washington, D.C.: National Academies Press.

Lichtenberg, E.M. *et al.* (2017) 'A global synthesis of the effects of diversified farming systems

on arthropod diversity within fields and across agricultural landscapes', *Global Change Biology*,

23(11), pp. 4946–4957. Available at: https://doi.org/10.1111/gcb.13714.

Long, J.A. and Long, M.J.A., 2019. Package 'interactions'. *See https://interactions.jacob-*

long.com.

Maxwell, S.L. *et al.* (2016) 'Biodiversity: The ravages of guns, nets and bulldozers', *Nature*,

536(7615), pp. 143–145. Available at: https://doi.org/10.1038/536143a.

Medlin, S. *et al.* (2016) 'Serosurvey of selected arboviral pathogens in free-ranging, two-toed

sloths (*Choloepus hoffmanni*) and three-toed sloths (*Bradypus variegatus*) in Costa Rica, 2005–

07', *Journal of wildlife diseases*, 52(4), pp. 883–892.

Mendenhall, C.D. *et al.* (2014) 'Countryside biogeography of Neotropical reptiles and

amphibians', *Ecology*, 95(4), pp. 856–870. Available at: https://doi.org/10.1890/12-2017.1.

Mendenhall, C.D. *et al.* (2016) 'Quantifying and sustaining biodiversity in tropical agricultural

landscapes', *Proceedings of the National Academy of Sciences*, 113(51), pp. 14544–14551.

Available at: https://doi.org/10.1073/pnas.1604981113.

Meyer Steiger, D.B., Ritchie, S.A. and Laurance, S.G.W. (2016) 'Mosquito communities and

disease risk influenced by land use change and seasonality in the Australian tropics', *Parasites &*

Vectors, 9(1), p. 387. Available at: https://doi.org/10.1186/s13071-016-1675-2.

Millennium ecosystem assessment, M. (2005) *Ecosystems and human well-being*. Island press

Washington, DC.

Mordecai, E.A. *et al.* (2019) 'Thermal biology of mosquito☐borne disease', *Ecology Letters*.

Edited by J. (Jeb) Byers, p. ele.13335. Available at: https://doi.org/10.1111/ele.13335.

Muturi, E.J. *et al.* (2006) 'Mosquito species diversity and abundance in relation to land use in a

riceland agroecosystem in Mwea, Kenya', *Journal of Vector Ecology*, 31(1), pp. 129–137.

Available at: https://doi.org/10.3376/1081-1710(2006)31[129:MSDAAI]2.0.CO;2.

Niebylski, M.L. *et al.* (1994) 'Blood hosts of *Aedes albopictus* in the United States', *Journal of*

the American Mosquito Control Association, 10(3), pp. 447–450.

Norris, K. (2008) 'Agriculture and biodiversity conservation: opportunity knocks', *Conservation*

Letters, 1(1), pp. 2–11. Available at: https://doi.org/10.1111/j.1755-263X.2008.00007.x.

Oksanen, J. *et al.* (2013) 'Package "vegan"', *Community ecology package, version*, 2(9), pp. 1–

295.

Pagès, H. *et al.* (2022) 'Biostrings: Efficient manipulation of biological strings'. (R package).

Available at: https://bioconductor.org/packages/Biostrings.

Pereira, H.M. *et al.* (2010) 'Scenarios for global biodiversity in the 21st Century', *Science*,

330(6010), pp. 1496–1501. Available at: https://doi.org/10.1126/science.1196624.

Piche-Ovares, M. *et al.* (2023) 'Serosurvey in two dengue hyperendemic areas of Costa Rica

evidence active circulation of WNV and SLEV in peri-domestic and domestic animals and in

humans', *Pathogens*, 12(1), p. 7. Available at: https://doi.org/10.3390/pathogens12010007.

Poulin, B., Lefebvre, G. and Paz, L. (2010) 'Red flag for green spray: adverse trophic effects of

Bti on breeding birds', *Journal of Applied Ecology*, 47(4), pp. 884–889. Available at:

https://doi.org/10.1111/j.1365-2664.2010.01821.x.

Prevedello, J.A. *et al.* (2019) 'Impacts of forestation and deforestation on local temperature

across the globe', *PLOS ONE*. Edited by Y. Xu, 14(3), p. e0213368. Available at:

https://doi.org/10.1371/journal.pone.0213368.

Ratnasingham, S. and Hebert, P.D.N. (2007) 'bold: The Barcode of Life data system

(http://www.barcodinglife.org)', *Molecular Ecology Notes*, 7(3), pp. 355–364. Available at:

https://doi.org/10.1111/j.1471-8286.2007.01678.x.

Reisen, W.K. (2003) 'Epidemiology of St. Louis encephalitis virus', *Advances in virus research*,

61, pp. 139–184.

Rezza, G. (2012) '*Aedes albopictus* and the reemergence of dengue', *BMC Public Health*, 12(1),

p. 72. Available at: https://doi.org/10.1186/1471-2458-12-72.

Richards, S.L. *et al.* (2006) 'Host-feeding patterns of *Aedes albopictus* (Diptera: Culicidae) in

relation to availability of human and domestic animals in suburban landscapes of central North

Carolina', *Journal of Medical Entomology*, 43(3), pp. 543–551. Available at:

https://doi.org/10.1093/jmedent/43.3.543.

Ricketts, T.H. *et al.* (2004) 'Economic value of tropical forest to coffee production', *Proceedings*

of the National Academy of Sciences, 101(34), pp. 12579–12582. Available at:

https://doi.org/10.1073/pnas.0405147101.

Ripley, B., Venables, B., Bates, D.M., Hornik, K., Gebhardt, A., Firth, D. and Ripley, M.B.,

2013. Package 'mass'. *Cran r*, *538*, pp.113-120.

Rojas-Araya, D. *et al.* (2017) 'New records of *Aedes albopictus* (Skuse) in four locations of

Costa Rica', *Revista Biomédica*, 28(2), pp. 79–88. Available at:

https://doi.org/10.32776/revbiomed.v28i2.575.

Romero-Vega, L.M. *et al.* (2023) 'Seasonal changes in the diversity, host preferences and

infectivity of mosquitoes in two arbovirus-endemic regions of Costa Rica', *Parasites & Vectors*,

16(1), p. 34. Available at: https://doi.org/10.1186/s13071-022-05579-y.

Sala, O.E. (2000) 'Global biodiversity scenarios for the year 2100', *Science*, 287(5459), pp.

1770–1774. Available at: https://doi.org/10.1126/science.287.5459.1770.

Sánchez-Azofeifa, G.A. *et al.* (2007) 'Costa Rica's Payment for Environmental Services

program: intention, implementation, and impact', *Conservation Biology*, 21(5), pp. 1165–1173.

Available at: https://doi.org/10.1111/j.1523-1739.2007.00751.x.

da Silva Pessoa Vieira, C.J. *et al.* (2022) 'Land-use effects on mosquito biodiversity and

potential arbovirus emergence in the Southern Amazon, Brazil', *Transboundary and Emerging

Diseases*, 69(4), pp. 1770–1781. Available at: https://doi.org/10.1111/tbed.14154.

Simmons, C.P. *et al.* (2012) 'Dengue', *New England Journal of Medicine*, 366(15), pp. 1423–

1432. Available at: https://doi.org/10.1056/NEJMra1110265.

Smith, R.J. (2022) *R package 'ecole', phytomosaic/ecole: ecole: School of Ecology Package*.

Available at: https://rdrr.io/github/phytomosaic/ecole/.

Smith, T.J. and Mayfield, M.M. (2015) 'Diptera species and functional diversity across tropical

Australian countryside landscapes', *Biological Conservation*, 191, pp. 436–443. Available at:

https://doi.org/10.1016/j.biocon.2015.07.035.

Thongsripong, P. *et al.* (2013) 'Mosquito vector diversity across habitats in central Thailand

endemic for dengue and other arthropod-borne diseases', *PLOS Neglected Tropical Diseases*,

7(10), p. e2507. Available at: https://doi.org/10.1371/journal.pntd.0002507.

Troyo, A. *et al.* (2006) 'Dengue in Costa Rica: the gap in local scientific research', *Revista*

Panamericana de Salud Pública, 20, pp. 350–360. Available at: https://doi.org/10.1590/S1020-

49892006001000012.

Troyo, A. *et al.* (2008) 'Seasonal profiles of <u>Aedes aegypti</u> (Diptera: Culicidae) larval habitats in

an urban area of Costa Rica with a history of mosquito control', *Journal of Vector Ecology*,

33(1), pp. 76–88. Available at: https://doi.org/10.3376/1081-

1710(2008)33[76:SPOAAD]2.0.CO;2.

Troyo, A. *et al.* (2009) 'Urban structure and dengue incidence in Puntarenas, Costa Rica',

Singapore Journal of Tropical Geography, 30(2), pp. 265–282. Available at:

https://doi.org/10.1111/j.1467-9493.2009.00367.x.

Tsuda, Y. *et al.* (2006) 'Different spatial distribution of *Aedes aegypti* and *Aedes albopictus*

along an urban-rural gradient and the relating environmental factors examined in three villages in

northern Thailand', *Journal of the American Mosquito Control Association*, 22(2), pp. 222–228.

Available at: https://doi.org/10.2987/8756-971X(2006)22[222:DSDOAA]2.0.CO;2.

Wickham, H (2007). "Reshaping data with the reshape package." *Journal of Statistical*

Software, **21**(12), 1–20. http://www.jstatsoft.org/v21/i12/.

Wickham, H. *et al.* (2019) 'Welcome to the Tidyverse', *Journal of Open Source Software*, 4(43),

p. 1686. Available at: https://doi.org/10.21105/joss.01686.

Wilke, C.O., Wickham, H. and Wilke, M.C.O., 2019. Package 'cowplot'. *Streamlined plot theme*

and plot annotations for 'ggplot2, *1*.

World Health Organization (1994) 'Outbreak of classic dengue, Costa Rica', *Weekly*

Epidemiological Record, 69, pp. 85–86.

World Health Organization (2021) *Dengue data application*. Available at:

https://ntdhq.shinyapps.io/dengue5.

Wright, E. (2016) 'Using DECIPHER v2.0 to analyze big biological sequence data in R', *The R*

Journal, 8(1), pp. 352–359.

Young, K.I. *et al.* (2017) 'Abundance and distribution of sylvatic dengue virus vectors in three

different land cover types in Sarawak, Malaysian Borneo', *Parasites & Vectors*, 10(1), p. 406.

Available at: https://doi.org/10.1186/s13071-017-2341-z.

Zimmerman, R.H. (1992) 'Ecology of malaria vectors in the Americas and future direction',

Memórias do Instituto Oswaldo Cruz, 87, pp. 371–383. Available at:

https://doi.org/10.1590/S0074-02761992000700064

Tables

827
828   **Table 1. PERMANOVA results for community composition compared among land use**

829   **types. Asterisks indicate p < 0.05.**

Land use pair	Sum of squares	F-Value	R^2	Bonferroni-adjusted p-value
agriculture vs forest	0.591	1.45	0.0785	0.291
agriculture vs residential	0.788	2.22	0.0846	0.021 *
forest vs residential	1.56	4.63	0.181	0.003 *

830
831