Queuosine Salvage in *Bartonella henselae* Houston 1: A Unique Evolutionary Path

Samia Quaiyum[a%], Yifeng Yuan[a%], Guangxin Sun[b], R. M. Madhushi N. Ratnayake[c#] Geoffrey Hutinet[a@],

Peter C. Dedon[b], Michael F. Minnick[d] and Valérie de Crécy-Lagard[a, e*]

[a] Department of Microbiology and Cell Science, University of Florida, Gainesville, FL 32611; [@]Current

address: Department of Biology, Haverford College, 370 Lancaster Ave, Haverford, PA

[b] Department of Biological Engineering, Massachusetts Institute of Technology, Cambridge, MA 02139

[c] Department of Chemistry, University of Florida, Gainesville, FL 32611; [#]Current address: Department of

Biochemistry, School of Medicine, Vanderbilt University, Nashville, TN 37232

[d] Division of Biological Sciences, University of Montana, Missoula, Montana, MT 59812

[e] Genetic Institute, University of Florida, FL 32611

[%]Contributed equally

*Corresponding author: vcrecy@ufl.edu

Short title: Queuosine salvage in Bartonella

Abstract

Queuosine (Q) stands out as the sole tRNA modification that can be synthesized via salvage pathways. Comparative genomic analyses identified specific bacteria that showed a discrepancy between the projected Q salvage route and the predicted substrate specificities of the two identified salvage proteins: 1) the distinctive enzyme tRNA guanine-34 transglycosylase (bacterial TGT, or bTGT), responsible for inserting precursor bases into target tRNAs; and 2) Queuosine Precursor Transporter (QPTR), a transporter protein that imports Q precursors. Organisms like the facultative intracellular pathogen *Bartonella henselae*, which possess only bTGT and QPTR but lack predicted enzymes for converting preQ$_1$ to Q, would be expected to salvage the queuine (q) base, mirroring the scenario for the obligate intracellular pathogen *Chlamydia trachomatis*. However, sequence analyses indicate that the substrate-specificity residues of their bTGTs resemble those of enzymes inserting preQ$_1$ rather than q. Intriguingly, mass spectrometry analyses of tRNA modification profiles in *B. henselae* reveal trace amounts of preQ$_1$, previously not observed in a natural context. Complementation analysis demonstrates that *B. henselae* bTGT and QPTR not only utilize preQ$_1$, akin to their *Escherichia coli* counterparts, but can also process q when provided at elevated concentrations. The experimental and phylogenomic analyses suggest that the Q pathway in *B. henselae* could represent an evolutionary transition among intracellular pathogens—from ancestors that synthesized Q *de novo* to a state prioritizing the salvage of q. Another possibility that will require further investigations is that the insertion of preQ$_1$ has fitness advantages when *B. henselae* is growing outside a mammalian host.

Keywords: Queuosine, *Bartonella henselae*, tRNA modification, intracellular parasite, TGT enzyme, YhhQ

Author summary

Transfer RNAs (tRNAs) are adaptors that deliver amino acids to ribosomes during translation of messenger RNAs (mRNAs) into proteins. tRNA molecules contain specially-modified nucleotides that affect many aspects of translation, including regulation of translational efficiency, as modified nucleotides primarily occur near the portion of tRNA (anticodon) that directly interacts with the coding sequence (codon) of the

mRNA while it is associated with a ribosome. Queuosine (Q) is a modified tRNA nucleotide located in the

anticodon that can be synthesized or uniquely imported from the environment as Q or a precursor using a

salvage mechanism. Free-living bacteria, e.g., *E. coli*, can synthesize Q or salvage precursors from the

environment, but many obligate intracellular pathogens, e.g., *Chlamydia trachomatis*, cannot synthesize Q

and must import a precursor from eukaryotic hosts. In this study, we determined that *Bartonella henselae*,

a facultative intracellular bacterial pathogen of vascular cells, falls somewhere in the middle, as it is unable

to synthesize Q but can salvage Q or certain precursors. The unusual nature of *Bartonella*'s system suggests

different evolutionary scenarios. It could be a snapshot of the transition from Q synthesis to strict Q salvage

or represent a unique adaptation to a complex multi-host lifestyle.

Introduction

tRNA modifications fine-tune translation by various mechanisms such as modulating the efficiency or

accuracy of translation or affecting tRNA stability (1). Recent studies have revealed that modifications can

play key roles in bacterial pathogenesis (2,3). Queuosine (Q) is a modification of the wobble base in tRNAs

with GUN anticodons of many bacteria and eukaryotes that can affect both translational efficiency and

accuracy depending on the organism (4,5). The *in vivo* significance of this modification has remained

enigmatic for decades as it has been lost repeatedly during evolution (6), but recent studies have suggested

that it may act as a regulatory component in the translation of proteins derived from genes enriched in TAT

codons compared with TAC codons (7,8). In bacteria, Q modification was shown to have roles in oxidative

stress, metal homeostasis, and virulence (9–12).

Q is synthesized from guanosine triphosphate (GTP) by bacteria in a complex eight-step pathway

fully elucidated in *E. coli* (Fig. 1A). Four enzymes (GCHI, QueD, QueE, and QueC) convert GTP into 7-

cyano-7-deazaguanine (preQ$_0$). QueF then reduces preQ$_0$ to 7-aminomethyl-7-deazaguanine (preQ$_1$) that is

inserted into tRNAs by tRNA guanine-34 transglycosylase (bacterial TGT, or bTGT) (4). The inserted base

preQ$_1$ is converted to Q by two additional steps involving QueA and QueG or QueH, depending on the

organism (4,13). It should be noted that Q is the only tRNA modification that can be salvaged or recycled

(5). Eukaryotes only use the salvage route, and their TGT enzyme (eTGT), a heterodimeric QTRT1/QTRT2

complex, incorporates the queuine (q) base directly into the target tRNAs (5). The salvage routes in bacteria

vary greatly. Some organisms lack the $preQ_0$ or $preQ_1$ pathway genes but encode all the downstream genes

and import these precursor bases to finalize Q synthesis *in vivo* (14). This salvage route is also observed in

organisms like *E. coli* that can synthesize Q *de novo*. Other bacteria lack all canonical synthesis genes

except *tgt* and can salvage q like eukaryotes. Two q salvage routes have been identified in bacteria, to date

(15), including the direct salvage route found in the intracellular pathogen *Chlamydia trachomatis*, or the

indirect salvage route found in the gut microbe *Clostridioides difficile* (Fig. 1B and C, respectively). In the

direct route, the substrate specificity of the *C. trachomatis* bTGT enzyme has shifted to insert q instead of

$preQ_1$ like most bacterial homologs (15). In the indirect route, a recently discovered radical enzyme (QueL)

can regenerate the $preQ_1$ intermediate from a q precursor that is imported directly or derived from the

hydrolysis of the Q nucleoside by QueK (15). Only a few transporters involved in Q salvage pathways have

been identified and experimentally characterized. The first was the YhhQ /COG1738 family, now renamed

QPTR (Queuosine Precursor Transporter), involved in the transport of $preQ_0$ and $preQ_1$ in *E. coli* and q in

C. trachomatis (14,15). Members of the Energy-Coupling Factors (ECFs) family had been predicted to be

involved in $preQ_0$ and $preQ_1$ transports (15)) and two of the *C. difficile* specificity components (or ECF-

QueT) were shown to transport a variety of Q precursors in a reconstituted system (15).

The proportion of Q-modified tRNAs can change with the developmental stage in several

eukaryotic parasites that undergo complex cycles that switch between hosts such as *Trypanosoma cruzei*

and *Entamoeba histolytica* (16,17). Very little is known about the role of Q in bacterial pathogens that also

switch between mammalian and insect hosts such as the facultative intracellular pathogen *Bartonella*

henselae. This bacterium uses fleas and possibly ticks as vectors during blood feeding. Feces of these insects

can also infect cats when they are scratched into a break in the skin. Once inside the cat, the bacteria enter

the bloodstream, primarily residing within endothelial cells, where they multiply (18). *B. henselae* exhibits

a high level of heterogeneity (19) and 16S rRNA sequence analyses led to the identification of two

serotypes, Houston-1, and Marseille (20,21). Metabolic reconstruction of the Q synthesis pathway of *B.*

henselae Houston 1 suggested it utilized a direct q salvage pathway like *C. trachomatis*. However, analysis

of *B. henselae* bTGT and QTPR sequences did not match this prediction as these were more similar to the

bTGT and QPTR enzymes that recognize $preQ_1$. To resolve this discrepancy, we set out to characterize

experimentally the Q salvage enzymes in this facultative intracellular pathogen.

Results

Metabolic reconstruction and sequence analyses of queuosine salvage genes give contradictory

results.

We previously showed that QPTR proteins have different substrate specificities, shifting from $preQ_0$ and

$preQ_1$ in *E. coli* to q in *C. trachomatis* (14,15). To better understand the molecular determinants that drive

this change in specificity, we constructed a Sequence Similarity Network (SSN) of the QPTR family

(PF02592). We then colored the SSN based on the presence/absence of the Q synthesis genes in the

corresponding genome as an indirect way to predict the substrate specificity of a given QPTR protein. As

shown in Fig. 2, we were able to generate an SSN that separated the QPTR's predicted to salvage $preQ_0$

and $preQ_1$ (in yellow or red) from those predicted to salvage q (in blue). However, a few exceptions stood

out even at a stringent alignment score of 70. The QPTRs found in organisms that encode only *tgt*, and

hence predicted to salvage q, cluster with QPTR proteins predicted to salvage $preQ_0$/ $preQ_1$ because their

genomes encode QueA and QueG or QueH (circled in Fig. 2). These include QPTR proteins from

Bartonella species such as *B. henselae* Houston-1 (UniProt id A0A0H3M726_BARHE).

115   Following up on this discrepancy, we retrieved all Q synthesis proteins from the InterPro database

with their IPR protein family ID (see Materials and Methods) and counted the numbers of each protein

encoded in each genome at the level of different taxonomic ranks, including order, class, and family (Table

S1). Then, we extracted the TGT sequences from all orders, classes, families, and genera that encode only

tgt as described in the methods section, aligned them, and compared the predicted substrate-binding

residues (Table S2). As we previously reported (15), TGTs that salvage q typically contains GG[LS][AS]G

in the substrate-binding pocket (Fig. 3). Interestingly, bTGTs in *Bartonella* and *Pelagibacter* contain a

GGLAVG site like the *E. coli* enzyme (Fig. 3) and are phylogenetically distant from other bTGTs found in

genomes that only encode *tgt* (Fig. 3). We then examined the substrate-binding pocket in a modeled

structure of *B. henselae* bTGT (Bh bTGT, UniProt ID A0A0R4J8M4_BARHE) aligned with the structure

of the human TGT catalytic subunit QTRT1 in complex with q (PDB ID: 6H45) (22). The aspartate residues

and G216GLAVGE222 that are conserved in bTGT proteins of the *Hyphomicrobiales* order are in

proximity to queuine (Fig. S1B and C). The predicted distance between V220 and the cis-diol groups of q

is less than 1 Angstrom (Fig. S1C), suggesting that it may prevent the binding of q to *B. henselae* bTGT as

in the *Zymomonas mobilis* TGT (23). We previously showed that *C. trachomatis* TGT salvages q and that

its substrate-binding pocket can accommodate the larger substrate (15). Here, we propose that the bTGT

protein and the QPTR transporter in *Bartonella* and *Pelagibacter* species salvage $preQ_1$, even though they

lack, like *C. trachomatis*, the enzymes that make Q-tRNAs from $preQ_1$-tRNAs.

B. henselae bTGT and QPTR proteins preferentially salvage $preQ_1$ but also q and Q with low affinity.

The gene encoding the *B. henselae* Bh bTGT protein complemented the Q- phenotype of a *queDF tgt*

deletion mutant of *E. coli* when expressed *in trans* and in the presence of exogenous $preQ_1$ even in low

concentrations (down to 10 nM) (Fig. 4A). Similarly, Bh QPTR transported both $preQ_1$ and $preQ_0$ in an *E.*

coli strain auxotrophic for $preQ_1$ and $preQ_0$ (Fig. 4B). These results confirmed our predictions based on the

SSN that the *Bartonella* bTGT and QPTR proteins use $preQ_1$ as a substrate but do not match the metabolic

reconstruction that predicted q salvage in *B. henselae*. Hence, we tested if this organism's bTGT and QPTR

could have evolved a broader substrate specificity and use q as a substrate.

To test whether the QPTR and bTGT proteins of *B. henselae* can use q as a substrate, we used an

E. coli strain that expresses the QPTR and/or the bTGT proteins of *C. trachomatis* that can only use q as a

substrate (Fig. 1) (15). Salvage of q was observed when expressing both Bh *tgt* and *yhhQ* genes at

concentrations of q over 500 nM (Fig. 4C). When overexpressing the *E. coli yhhQ* gene, no such salvage

of q was observed even at concentrations of 5 μM (Fig. 4C). However, if the *C. trachomatis* bTGT and

QPTR proteins can salvage q when present at 100 nM, the Bh QPTR cannot (Fig. S2). These results showed

that the Bh QPTR protein has acquired the capacity to transport q while retaining the $preQ_1/preQ_0$

specificity, but it is still not as efficient as the *C. trachomatis* QPTR transporter that is specific for q.

150   We then tested whether the Bh QPTR could transport the Q ribonucleoside by modeling with an *E.*

coli strain expressing the *C. difficile* Q hydrolase (Cd *queK*) and q lyase (Cd *queL*) genes that allow Q to

be salvaged by *E. coli* (Fig. 1) (15). Expressing only Bh QPTR allowed the salvage of Q only at extremely

high concentrations (5 μM) (Fig. 4D). When the *C. difficile* Q transporter (Cd ECF_TAA', QueT) is

expressed in this strain, Q can be salvaged when present at concentrations of 100 nM [(15) and (Fig. 4D].

Traces of $preQ_1$ can be detected in endogenous *B. henselae* tRNAs

The natural habitat of this intracellular pathogen (mammalian vasculature) should be richer in q than in

$preQ_1$. We, therefore, analyzed by LC-MS/MS the tRNA modification profile of bulk tRNA extracted from

B. henselae cells grown in sheep blood agar (HIBB) medium. The experiments were performed three times

independently with conflicting results. The first two experiments were done to test the tRNA extraction

protocols with an intracellular bacterium, using only one sample each time. Small amounts of $preQ_1$ and

minute amounts of Q were detected the first time (but neither was present the second time (data not shown).

Because cells were grown in the presence of sheep blood and serum, we had little control over the sources

of Q or $preQ_1$; we, therefore, repeated the experiment a third time with 5 independent samples, adding 100

nM $preQ_1$ in three of the samples. As shown in Fig. 5, Fig. S3 and Table S3, small amounts of $preQ_1$ and

Q were detected in all samples (around 1000 times less than in a typical *E. coli* sample). The exogenous

addition of 100 nM $preQ_1$ did not make any difference. It is not possible to determine if the observed Q was

derived from *B. henselae* tRNAs or from contaminating mammalian tRNAs, as eukaryotic-specific tRNA

modifications such as m^2_2G, are detected in similar quantities (Table S3) and great variations in Q levels

were observed between samples. Nevertheless, the presence of preQ$_1$ cannot be explained by any

contamination as mammalian host tRNAs never harbor this modification and other bacteria would not

accumulate it, as preQ$_1$ had only been detected previously in *queA* mutants of *E. coli* (24). The low amounts

detected suggest that the tRNAs are not fully modified, as the preQ$_1$ levels are 11%-40% of cmnm^{5}s^{2}U

levels and 1.1%-2.8% of k^{2}C levels; two well-conserved bacterial modifications (25)(Fig. S4 and Table

S3).

Early loss of Q pathway genes in the *Bartonellaceae* family within the *Hyphomicrobiales* order

B. henselae belongs to the Alphaproteobacteria class (25). This is a diverse Gram-negative taxon comprised

of several phototrophic genera, several genera metabolizing C1-compounds (e.g., *Methylobacterium* spp.),

symbionts of plants (e.g., *Rhizobium* spp.), endosymbionts of arthropods (*Wolbachia*) and intracellular

pathogens (e.g., *Rickettsia*) (26). To better understand the evolution of the Q synthesis and salvage pathway

in Alphaproteobacteria, we performed a phylogenetic distribution analysis of the corresponding genes in

2,127 different species with complete genome sequences from the class in the BV-BRC database as

described in the methods section (Fig. S5). The tree suggests that there were three events involving the loss

of preQ$_1$ synthesis genes: one occurred after the split between *Brucella* and *Bartonella* species (Fig. 6 and

Fig. S5 node 1), one occurred after the split of *Pelagibacteraceae* (Fig. S5 and Fig S6 node 2), and one

occurred after the split of *Anaplasmataceae* (Fig. S5 and Fig S6 node 3). This analysis revealed that most

Alphaproteobacteria are prototrophic for Q, as 1,414 (64%) of the species analyzed encode the complete

pathway. In addition, if the loss of the preQ$_1$ synthesis genes occurred sporadically in different branches,

all the organisms analyzed, with the exception of *Bartonella*, harbored *tgt*, *queA*, and *queG* genes and hence

were not predicted to salvage q. Focusing more specifically on the *Bartonella* genus using a similar analysis

revealed a very different pattern (Fig. 6).

Among the 65 organisms in the *Bartonella* genus with available complete genomes, as of October

2023, only *Bartonella* sp. HY038, branching at the root of the genus, encoded the canonical *de novo* Q

synthesis pathway. Most, like *B. henselae*, have lost all the genes but *tgt*. In addition, a more in-depth

analysis revealed fragmented *tgt* genes of several *B. quintana* strains (boxed in Fig. 6), suggesting these

organisms have totally lost the capacity to make Q-modified tRNAs. This evolutionary scenario seems to

be a recurring theme in intracellular bacteria like the *Rickettsiales*. While nearly all rickettsiae branching

closer to the root retain the full pathway except for *queD* (collapsed in Fig. S6), other rickettsiae such as

Anaplasma, Ehrlichia, and *Wolbachia*, have lost nearly the full pathway. However, cases of fragmented *tgt*

genes are rare and only observed in a *Wolbachia* endosymbiont of *Cimex lectularius* (box in Fig. S5).

In summary, the phylogenic distribution analysis suggests that the direct ancestor of *Bartonella*

species must have harbored the full Q pathway but that it was lost very early in the evolution of this clade.

Most bacteria in this clade are predicted to transport and insert preQ$_1$ but without further conversion to Q.

In addition, some species like *B. quintana* have lost the pathway completely (Fig. 6).

Discussion

Q is an ancient modification predicted to be present in the ancestors of bacteria (27). It is still present in

most extant bacteria even if minimalist genetic codes can exist without this complex modification (28).

Independent analyses of the genomes of bacteria in the human microbiome have shown that 90 to 95% of

these organisms maintain the capacity to synthesize or salvage Q with around half encoding the full

synthesis pathway (15,29). Many of the bacteria that have lost Q are organisms that have undergone a

genome reduction process, where their genetic material has been streamlined over evolutionary time such

as the parasitic *Mycoplasma* spp. or insect endosymbionts such as *Riesia pediculicola* (30,31).

Obligate intracellular human pathogens tend to have reduced genomes compared to their free-living

ancestors as their metabolisms have adapted to a nutritionally rich niche (32). Regarding Q, the scenarios

that one can envision in the transition to a strict intracellular lifestyle with access to the queuine precursor

from the mammalian host are: 1) keeping the ancestral pathway; 2) losing the modification; or 3) switching

to a queuine salvage route. We performed the metabolic reconstruction of Q metabolism in genera of strict

intracellular human pathogens such as *Chlamydia* spp., members of the order Rickettsiales (*Anaplasma*

spp., and *Rickettsia* spp.) and *Coxiella burnetii* (33,34). Indeed, examples of these three possible paths were

observed (Fig. 7): *Rickettsia* (Fig. 7A) and *Coxiella* (Table S5 line 267) have kept the full Q synthesis

pathway; *Anaplasma* spp. (Fig. 7A), *Borrelia* (Fig. 7B), *Ehrlichia* spp. (Table S5 line 52), and *Chlamydia*

(Table S5 line 212), kept only *tgt*. Most species in Mycobacteriales have totally lost the pathway genes

except *tgt* (Fig. 7C).

The situation seen in *B. henselae* is not commonly observed in other intracellular bacteria and no

other organisms in the Hyphomicrobiales order seem to follow the same pattern (Fig. 7D). Indeed, the

presence of bTGT and QPTR homologs and the absence of QueA and QueG or QueH (Fig. 6) would suggest

that these organisms salvage q like *C. trachomatis*, but the corresponding enzymes have retained their

substrate specificity towards preQ$_1$ (Figs. 3 and 4). PreQ$_1$ is not a molecule found in mammalian cells (Brian

Green, personal communication), and the fact that we were able to detect a small proportion of tRNAs

carrying that modification when extracted from *B. henselae* cells grown in HIBB suggests this pathway is

functional even if the source of preQ$_1$ remains a mystery and could be due to contamination with a preQ$_1$

synthesizing organism, an unknown source of preQ$_1$ in the culture medium, or from the activity of a yet-to-

be-discovered q hydrolase (intracellular or possibly extracellular as q transport is not efficient as discussed

below) (Fig. 1D). The low amount of preQ$_1$ modification (Table S3) suggests it does not play an important

role in decoding accuracy under these conditions. The fact that *tgt* gene decay is observed in several

organisms in this clade such as *B. quintana* reinforces this idea. A primary difference between *B. quintana*

and *B. henselae* is their reservoir ecology. *B. quintana* uses only humans as a reservoir, whereas *B. henselae*

is more promiscuous and frequently isolated from both cats and humans (35).

Another intriguing finding of this study is that the QPTR and bTGT enzymes of *B. henselae* can

use q as a substrate when present at high concentrations (>500 nM), whereas the *E. coli* orthologs cannot.

It is difficult to establish if such concentrations could be physiological, and we could not show with certainty

that *B. henselae* tRNA extracted from HIBB-grown cells harbored Q. One can propose several evolutionary

explanations for this broadening of substrate specificity of the *B. henselae* salvage proteins. In one, the

enzyme and transporter specificities would become more relaxed in *B. henselae* as they are no longer under

selective pressure to maintain efficient preQ$_1$ selection, and we are observing an intermediate stage along

the evolutionary loss of the whole pathway. An argument against that hypothesis is that if $preQ_1$ synthesis

genes and *queA*, *queG* and *queH* genes seem to have been lost very early in the clade, *tgt* is often the last

maintained and we did not find any examples where *tgt* was lost with the other genes maintained, suggesting

a fitness advantage of maintaining the *tgt* gene. In addition, both *B. henselae* QPTR and bTGT encoding

genes are expressed and the proteins detected *in vitro* [see Table S5 of (36)]. The other possibility is that

we are observing a transition of a $preQ_1$ salvage to a q salvage that is working poorly in human cells but

could be efficient in an environment with more q/Q. Could the insect vector provide such an environment?

Answering these questions will require further studies, including additional quantitative data on $q/preQ_1$

levels in different environments and tRNA modification profiles along the pathogen's life cycle. In

summary, the study sheds light on the diverse and adaptable nature of queuosine metabolism in various

bacteria, particularly in intracellular pathogens. The unique characteristics of Q salvage observed in *B.*

henselae raise intriguing questions about its role in different host/vector environments. Further

investigations are warranted to unravel the complexities of Q salvage and its implications to *Bartonella*'s

virulence.

Materials and methods

Comparative genomics and bioinformatics

The BLAST tools (37) resources at the National Center for Biotechnology Information (NCBI) and BV-

BRC (38) were routinely used. Multiple sequence alignments were built using MUSCLE v5. 1 (39) and

visualized and edited with Jalview2 (40). Protein sequences were retrieved from the NCBI using the

following accession numbers: Ct YhhQ, NP_219643.1; Ct TGT, NP_219697.1; Cd ECF-A,

YP_001086568.1; Cd ECF-A, YP_001086569.1; Cd ECF-T, YP_001086570.1; Cd Queuosine hydrolase,

YP_001088185.1; Cd ECF-S, YP_001088186.1; Cd Queuine lyase, YP_001088187.1; Ec YhhQ, WP

001100469.1; Ec bTGT; Bh YhhQ, WP_011181356.; Bh bTGT, WP_011180873.1. Protein IDs used for

this study are listed in Table S6. Complete genomes of Alphaproteobacteria with good quality were

retrieved from BV-BRC (38).

For metabolic reconstruction analyses in each taxonomic rank, all protein members were retrieved

from the InterPro database (41) using the following IPR family ID: FolE1, IPR001474; FolE2, IPR003801;

QueD, IPR007115; QueE, IPR024924; QueC, IPR018317; QueF, IPR00029500; TGT, IPR004803; QueA,

IPR003699; QueG, IPR004453; QueH, IPR003828. A universal single-copy small ribosomal protein uS2,

(IPR001865) was used to estimate the number of organisms in each rank. The number of proteins per

taxonomic rank was computed and the criteria used for filtering the groups ranks encoding just TGT were

the following: 1) the number of TGT proteins was no less than 10 so we were not polluted with small

taxonomic sample size; 2) the number of each of the QueDECFAGH proteins was no more than a fifth of

the number of TGT proteins. To analyze the conserved residues in the substrate-binding pockets, the

sequences of TGT proteins from select taxonomic groups were retrieved from UniProt and aligned using

MUSCLE v5.1 (39). The conserved residues were visualized using weblogo3 (42).

The structure of TGT proteins (*Bartonella henselae* str. Houston-1 bTGT, A0A0R4J8M4;

Anaplasma phagocytophilum bTGT, S6G6J1; *Nakamurella multipartita* bTGT, C8X7A7) were modeled

using SWISS-MODEL (40) using Alpha Fold structure (*Bartonella fuyuanensis* bTGT (ID

A0A840DZ06_9HYPH), *Anaplasma phagocytophilum* str ApMUC09 TGT (ID A0A0F3NAG3_ANAPH),

and *Nakamurella multipartita* bTGT (ID C8X7A7_NAKMY) as templates respectively (43). The cartoon

representation of protein structure was produced by PyMol (version 2.5) (44) and colored by domain (red,

N-terminus; light blue, C-terminus).

Sequence Similarity Networks (SSNs)

The Enzyme Function Initiative (EFI) suite of web tools was used to generate the SSN (45). Visualization

of SSNs was carried out using Cytoscape 3.10.1ape (46). 7,625 PF02592 family sequences were retrieved

from UniProt using the family option with fraction of 3 and submitted to EFI. The initial SSN was generated

with an alignment score cutoff set such that each connection (edge) represents a sequence identity of above

approximately 40%. The obtained SSN was first colored according to the configurations for salvaging

preQ$_1$, preQ$_0$, queuine, and Queuosine *de novo* synthesis. Then more stringent SSNs were created by

increasing the alignment score cutoff in small increments (usually by 5). This process was repeated until

most clusters were homogeneous in their colors. The UniProt IDs were associated with the genome ID

including GenBank/EMBL, RefSeq nucleotide, BV-BRC genome ID, Ensembl genome ID, using

homemade scripts (scripts available upon request). The UniProt IDs of PF02592 family sequences in the

SSN are listed in Table S7 as well as corresponding presence of Q pathway genes. The connection between

UniProt IDs and genome information was performed by querying UniProt ID mapping file using homemade

scripts (scripts available upon request).

Phylogenetic investigations and Q gene presence/absence distribution pattern analysis

For phylogenetic analysis of species, the 20 ribosomal protein data (L2, L3, L4, L5, L6, L9, L10, L14, L16,

L18, L22, L24, S2, S3, S4, S5, S8, S10, S17, and S19) were retrieved from BV-BRC and aligned

independently using MUSCLE v5.1 (39). Alignments were trimmed using BMGE v1.12 using default

settings (47) and then concatenated. To build the phylogenetic tree of TGT proteins, the sequences of TGT

proteins from each taxonomic rank were retrieved (ID are listed in Table S6) and subject to alignment with

MUSCLE (39) and trimming with BMGE (47). The maximum likelihood tree was built by FastTree (48)

using the LG-cat model with a hundred bootstraps. Trees were visualized using iTOL (49). To view the

gene clustering near *tgt* in different bacterial genomes, GizmoGene (http://www.gizmogene.com) was used

to extract gene regions and subjected to visualization using Gene Graphics (50). The genomic regions are

listed in Table S4.

Strains, media, and growth conditions

Strains and plasmids used in this study are listed in Table S9. LB medium (tryptone 10 grams/liter, yeast

extract 5 grams/liter, sodium chloride 5 grams/liter) was routinely used for *E. coli* growth at 37°C. The

medium was solidified using 15 g/L of agar. As needed, kanamycin (50 g/mL), ampicillin (100 g/mL), and

chloramphenicol (25 g/mL) were added. In the presence of exogenous Q precursors as previously described

(51), cells were cultured in M9-defined medium containing 1% glycerol (Thermo Fisher Scientific,

Waltham, MA, USA) for the purpose of eliminating background Q-tRNA. After cells reached an optical

density at 600 nm (OD_{600nm}) of 0.1-0.2, 0.2% arabinose was added to induce the expression of genes under

the pBAD promoter. After cells reached an OD_{600nm} of 0.2, DMSO, $preQ_0$, $preQ_1$, q, or Q were added. The

transport reaction was stopped at time points of 30 or 60 min after supplementing with DMSO or different

Q precursors by placing samples on melting ice and then centrifuging, followed by tRNA extraction. Q was

purchased from Epitoire (Singapore), q from Santa Cruz Biotechnology, $preQ_1$ and $preQ_0$ from Sigma-

Aldrich.

$\quad$ *B. henselae* Houston I was obtained from the American Type Culture Collection (ATCC 49882)

and cultivated as previously described (52) on HIBB agar plates [Bacto heart infusion agar (Becton,

Dickinson, Sparks, MD) supplemented with 4% defibrinated sheep blood and 2% sheep serum (Quad Five,

Ryegate, MT) by volume] for 4 days at 37°C, 5% CO_2 and 100% relative humidity. When required, $preQ_1$

was added to a final concentration of 100 nM. Following harvest into ice-cold heart infusion broth, tRNA

was collected from the bacterial cells.

Construction of *E. coli* strains and plasmids

B. henselae yhhQ gene (Bh *yhhQ*) was chemically synthesized (without optimization) in pTWIST–Kan

vector. XbaI and HindIII restrictions sites were added at the 5′ and 3′ ends, respectively (Twist Bioscience

HQ) (**Table S10**). Bh *yhhQ* DNA sequence was amplified using two primers pairs (F_Bh

yhhQ_XbaI_PBAD33 and R_Bh yhhQ_HindIII_PBAD33) by PCR with the addition of restrictions

sites XbaI and HindIII at their 5′ and 3′ ends, respectively. Bh *yhhQ* was cloned into the XbaI and HindIII

sites of pBAD33. *B. henselae tgt* (Bh *tgt*) was amplified by PCR from *B. henselae* genomic DNA using

the KpnI-RBS-TGTBh-F_PBAD24 and TGTBh-SbfI-R _PBAD24 primers and cloned into the KpnI and

SbfI sites of pBAD24. The UniProt IDs for Bh *yhhQ* and Bh *tgt* are A0A0H3M726_BARHE and

A0A0R4J8M4_BARHE, respectively. *E. coli* transformations were performed using the $CaCl_2$ chemical

transformation procedure (53). Transformants were selected on LB agar supplemented with ampicillin or

chloramphenicol (100 µg/mL). The clones were validated through sequencing and PCR analyses using

primers designed specifically for Bh *yhhQ* and Bh *tgt* genes. All primers used in this study are listed in

Table S10.

tRNA extraction and migration

Cells were harvested by centrifugation at 16,000 x g for 2 minutes at 4°C. Immediately after pelleting, the

cells were resuspended in 1 mL of Trizol (Thermo Fisher Scientific, Waltham, MA, USA). According to

the manufacturer's instructions, small RNA was extracted with the PureLink™ miRNA Isolation kit

(Thermo Fisher Scientific, Waltham, MA, USA). 50 µL of RNase-free water were used to elute the purified

RNAs. Quantification of prepared tRNA was performed using a Nanodrop 1000 spectrophotometer. We

loaded 150 ng of tRNAs per well on a denaturing 8 M urea, 8% polyacrylamide gel containing 0.5% 3-

(Acrylamido) phenylboronic acid (Sigma-Aldrich) after resuspending in a 2X RNA Loading Dye (NEB).

Migration was performed in a mixture of 1X TAE at 4°C. With a wet transfer apparatus in 1X TAE at 150

mA at 4°C for 90 minutes, tRNAs were transferred onto a Biodyne B precut nylon membrane (Thermo

Scientific). The membrane was UV irradiated in a UV crosslinker (Fisher FB-UVXL-1000) at a preset UV

energy dosage of 120 mJ/cm2. The North2South Chemiluminescent Hybridization and Detection Kit

(Thermo) was used to detect tRNA^{Asp}. As the DIG Easy Hyb (Roche) drastically reduces the background

noise, it was used as the initial membrane-blocking buffer instead of the North2South kit's membrane-

blocking buffer. Hybridization was done at 60°C, using the specific biotinylated primer for tRNA Asp GUC

(14) (5' biotin-CCCTGCGTGACAGGCAGG 3' for *E. coli* added to a final concentration of 50 ng/mL.

The blot was visualized by the iBright™ Imaging Systems.

tRNA profiling by mass spectrometry

tRNA for each sample (1.8 µg) was hydrolyzed in a 30 µL digestion cocktail containing 2.49 U benzonase,

3 U CIAP (calf intestinal alkaline phosphatase), 0.07 U PDE I (phosphodiesterase I), 0.1 mM deferoxamine,

0.1 mM BHT (butylated hydroxytoluene), 3 ng coformycin, 25 nM 15N-dA (internal standard [^{15}N]$_5$-

deoxyadenosine), 2.5 mM MgCl$_2$ and 5 mM Tris-HCL buffer pH 8.0. The digestion mixture was incubated

at 37 °C for 6 h. After digestion, all samples were analyzed by chromatography-coupled triple-quadrupole

mass spectrometry (LC-MS/MS). For each sample, 600 ng of hydrolysate was injected for two technical

replicates. Using synthetic standards, HPLC retention times of RNA modifications were confirmed on a

Waters Acuity BEH C18 column (50×2.1 mm inner diameter, 1.7 µm particle size) coupled to an Agilent

1290 HPLC system and an Agilent 6495 triple-quad mass spectrometer. The Agilent sample vial insert was

used. The HPLC system was operated at 25 °C and a flow rate of 0.3 mL/min in a gradient Table S11 with

Buffer A (0.02% formic acid in double distilled water) and Buffer B (0.02% formic acid in 70%

acetonitrile). The HPLC column was coupled to the mass spectrometer with an electrospray ionization

source in positive mode with the following parameters: Dry gas temperature, 200 °C; gas flow, 11 L/min;

nebulizer, 20 psi; sheath gas temperature, 300 °C; sheath gas flow, 12 L/min; capillary voltage, 3000 V;

nozzle voltage, 0 V. Multiple reaction monitoring (MRM) mode was used for detection of product ions

derived from the precursor ions for all the RNA modifications with instrument parameters including the

collision energy (CE) optimized for maximal sensitivity for the modification. Based on synthetic standards

(Biosynth) with optimized collision energies, the following transitions and retention times (except k^2C,

which we do not have standard) were monitored: $cmnm^5s^2U$, m/z 348 → 141, 2.36 min; k^2C, m/z 372.1 →

240.1; $m^{2,2}G$, m/z 312 → 180, 9.70 min; $preQ_1$, m/z 312 → 163, 2.15 min; Q, m/z 410 → 163, 5.53 min.

Signal intensities for each ribonucleoside were normalized by dividing by the sum of the UV signal

intensities of the four canonical ribonucleosides recorded with an in-line UV spectrophotometer at 260 nm.

Conflicts of Interest

No conflicts of interest.

Funding Information

This work was funded by the National Institutes of Health (project numbers GM070641, ES026856, and

ES024615) and the National Research Foundation of Singapore through the Singapore-MIT Alliance for

Research and Technology Antimicrobial Resistance Interdisciplinary Research Group.

Acknowledgments

We are thankful to Dr Brian Greene for communicating unpublished results.

References

1. El Yacoubi B, Bailly M, de Crécy-Lagard V. Biosynthesis and function of posttranscriptional modifications of transfer RNAs. Annu Rev Genet. 2012 Dec 15;46(1):69–95.

2. Antoine L, Bahena-Ceron R, Bunwaree HD, Gobry M, Loegler V, Romby P, et al. RNA modifications in pathogenic bacteria: Impact on host adaptation and virulence. Vol. 12, Genes. MDPI AG; 2021. p. 1125.

3. Fleming BA, Blango MG, Rousek AA, Kincannon WM, Tran A, Lewis AJ, et al. A tRNA modifying enzyme as a tunable regulatory nexus for bacterial stress responses and virulence. Nucleic Acids Res. 2022 Jul 22;50(13):7570–90.

4. Hutinet G, Swarjo MA, de Crécy-Lagard V. Deazaguanine derivatives, examples of crosstalk between RNA and DNA modification pathways. Vol. 14, RNA Biology. Taylor and Francis Inc.; 2017. p. 1175–84.

5. Fergus C, Barnes D, Alqasem MA, Kelly VP. The queuine micronutrient: Charting a course from microbe to man. Vol. 7, Nutrients. MDPI AG; 2015. p. 2897–929.

6. Zallot R, Brochier-Armanet C, Gaston KW, Forouhar F, Limbach PA, Hunt JF, et al. Plant, animal, and fungal micronutrient queuosine is salvaged by members of the DUF2419 protein family. ACS Chem Biol. 2014 Aug 15;9(8):1812–25.

7. Chevance FFV, Le Guyon S, Hughes KT. The effects of codon context on in vivo translation speed. PLoS Genet. 2014;10(6):e1004392.

8. Whittle CA, Kulkarni A, Chung N, Extavour CG. Adaptation of codon and amino acid use for translational functions in highly expressed cricket genes. BMC Genomics. 2021 Dec 1;22(1):234.

9.   Pollo-Oliveira L, Davis NK, Hossain I, Ho P, Yuan Y, Salguero García P, et al. The absence of the

queuosine tRNA modification leads to pleiotropic phenotypes revealing perturbations of metal and

oxidative stress homeostasis in *Escherichia coli* K12. Metallomics. 2022 Sep 24;14(9):mfac065.

10.   Durand JM, Dagberg B, Uhlin BE, Björk GR. Transfer RNA modification, temperature and DNA

superhelicity have a common target in the regulatory network of the virulence of *Shigella flexneri*:

the expression of the *virF* gene. Mol Microbiol. 2000;35(4):924–35.

11.   Díaz-Rullo J, González-Pastor JE. tRNA queuosine modification is involved in biofilm formation

and virulence in bacteria. Nucleic Acids Res. 2023 Aug 28;51(18):9821–37.

12.   Fruchard L, Babosan A, Carvalho A, Lang M, Li B, Duchateau M, et al. Queuosine modification of

tRNA-Tyrosine elicits translational reprogramming and enhances growth of *Vibrio cholerae* with

aminoglycosides. bioRxiv]. 2022 Jan 1;2022.09.26.509455.

13.   Zallot R, Ross R, Chen WH, Bruner SD, Limbach PA, de Crécy-Lagard V. Identification of a novel

epoxyqueuosine reductase family by comparative genomics. ACS Chem Biol. 2017 Feb

8;12(3):844–51.

14.   Zallot R, Yuan Y, De Crécy-Lagard V. The *Escherichia coli* COG1738 member YhhQ is involved

in 7-cyanodeazaguanine (preQ$_0$) transport. Biomolecules. 2017 Mar 1;7(1):12.

15.   Yuan Y, Zallot R, Grove TL, Payan DJ, Martin-Verstraete I, Šepić S, et al. Discovery of novel

bacterial queuine salvage enzymes and pathways in human pathogens. Proc Natl Acad Sci U S A.

2019 Sep 17;116(38):19126–35.

16.   Guillén N. Pathogenicity and virulence of *Entamoeba histolytica*, the agent of amoebiasis. Vol. 14,

Virulence. 2023. p. 2158656.

17.   Dixit S, Kessler AC, Henderson J, Pan X, Zhao R, D'Almeida GS, et al. Dynamic queuosine changes

in tRNA couple nutrient levels to codon choice in *Trypanosoma brucei*. Nucleic Acids Res. 2021

Dec 16;49(22):12986–99.

18.   Jin X, Gou Y, Xin Y, Li J, Sun J, Li T, et al. Advancements in understanding the molecular and

immune mechanisms of Bartonella pathogenicity. Front Microbiol. 2023 Jul 27;14:1196700.

19.   Hansmann Y, DeMartino S, Piémont Y, Meyer N, Mariet P, Heller R, et al. Diagnosis of cat scratch
disease with detection of *Bartonella henselae* by PCR: A study of patients with lymph node
enlargement. J Clin Microbiol. 2005 Aug;43(8):3800–6.

20.   La Scola B, Liang Z, Zeaiter Z, Houpikian P, Grimont PAD, Raoult D. Genotypic characteristics of
two serotypes of *Bartonella henselae*. J Clin Microbiol. 2002;40(6):2002–8.

21.   Zeaiter Z, Fournier PE, Raoult D. Genomic variation of *Bartonella henselae* strains detected in
lymph nodes of patients with cat scratch disease. J Clin Microbiol. 2002;40(3):1023–30.

22.   Johannsson S, Neumann P, Ficner R. Crystal structure of the human tRNA guanine transglycosylase
catalytic subunit QTRT1. Biomolecules. 2018 Sep 1;8(3):81.

23.   Stengl B, Reuter K, Klebe G. Mechanism and substrate specificity of tRNA-Guanine
transglycosylases (TGTs): tRNA-modifying enzymes from the three different kingdoms of life share
a common catalytic mechanism. Chem Bio Chem. 2005;6(11):1926–39.

24.   Slany RK, Bosl M, Kersten H. Transfer and isomerization of the ribose moiety of AdoMet during
the biosynthesis of queuosine tRNAs, a new unique reaction catalyzed by the QueA protein from
Escherichia coli. Biochimie. 1994;76(5):389–93.

25.   Boccaletto P, MacHnicka MA, Purta E, Pitkowski P, Baginski B, Wirecki TK, et al. MODOMICS:
A database of RNA modification pathways. 2017 update. Nucleic Acids Res. 2018;46(D1):D303–
7.

26.   Ferla MP, Thrash JC, Giovannoni SJ, Patrick WM. New rRNA gene-based phylogenies of the
Alphaproteobacteria provide perspective on major groups, mitochondrial ancestry and phylogenetic
instability. PLoS One. 2013 Dec 11;8(12):e83383.

27.   Cermakian N, Cedergren R. Modified nucleosides always were: an evolutionary model. In: Grosjean
H, Benne R, editors. Modification and Editing of RNA. Washington, D. C.: ASM Press; 1998. p.
535–41.

28.   Garzón MJ, Reyes-Prieto M, Gil R. The minimal translation machinery: what we can learn from
naturally and experimentally reduced genomes. Front Microbiol. 2022;13:858983.

29.   Yan F, Xiang S, Shi L, Zhu X. Synthesis of queuine by colonic gut microbiome via cross-feeding.

Food Frontiers1. 2023 Sep 15;1–14.

30.   Grosjean H, Breton M, Sirand-Pugnet P, Tardy F, Thiaucourt F, Citti C, et al. Predicting the minimal

translation apparatus: lessons from the reductive evolution of Mollicutes. PLoS Genet. 2014;10(5).

31.   de Crécy-Lagard V, Marck C, Grosjean H. Decoding in Candidatus *Riesia pediculicola*, close to a

minimal tRNA modification set? Trends Cell Mol Biol. 2012; 7:11–34.

32.   Murray GGR, Charlesworth J, Miller EL, Casey MJ, Lloyd CT, Gottschalk M, et al. Genome

reduction Is associated with bacterial pathogenicity across different scales of temporal and

ecological divergence. Mol Biol Evol. 2021;38(4):1570–9.

33.   McClure EE, Chávez ASO, Shaw DK, Carlyon JA, Ganta RR, Noh SM, et al. Engineering of

obligate intracellular bacteria: Progress, challenges and paradigms. Vol. 15, Nature Reviews

Microbiology. 2017. p. 544–58.

34.   Loterio RK, Zamboni DS, Newton HJ. Keeping the host alive - Lessons from obligate intracellular

bacterial pathogens. Vol. 79, Pathogens and Disease. 2021. p. ftab052.

35.   Alsmark CM, Frank AC, Karlberg EO, Legault BA, Ardell DH, Canbäck B, et al. The louse-borne

human pathogen *Bartonella quintana* is a genomic derivative of the zoonotic agent *Bartonella*

henselae. Proc Natl Acad Sci U S A. 2004;101(26):9716–21.

36.   Omasits U, Quebatte M, Stekhoven DJ, Fortes C, Roschitzki B, Robinson MD, et al. Directed

shotgun proteomics guided by saturated RNA-seq identifies a complete expressed prokaryotic

proteome. Genome Res. 2013 Nov;23(11):1916–27.

37.   Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, Miller W, et al. Gapped BLAST and PSI-

BLAST: a new generation of protein database search programs. Nucleic Acids Res.

1997;25(17):3389–402.

38.   Olson RD, Assaf R, Brettin T, Conrad N, Cucinell C, Davis JJ, et al. Introducing the Bacterial and

Viral Bioinformatics Resource Center (BV-BRC): a resource combining PATRIC, IRD and ViPR.

Nucleic Acids Res. 2023 Jan 6;51(D1):D678–89.

39.   Edgar RC. Muscle5: High-accuracy alignment ensembles enable unbiased assessments of sequence
homology and phylogeny. Nat Commun. 2022 Nov 15;13(1):6968.

40.   Waterhouse AM, Procter JB, Martin DMA, Clamp M, Barton GJ. Jalview Version 2-A multiple
sequence alignment editor and analysis workbench. Bioinformatics. 2009;25(9):1189–91.

41.   Paysan-Lafosse T, Blum M, Chuguransky S, Grego T, Pinto BL, Salazar GA, et al. InterPro in 2022.
Nucleic Acids Res. 2023 Jan 6;51(D1):D418–27.

42.   Crooks GE, Hon G, Chandonia JM, Brenner SE. WebLogo: A Sequence Logo Generator. Genome
Res 2004;14(6):1188–90.

43.   Varadi M, Anyango S, Deshpande M, Nair S, Natassia C, Yordanova G, et al. AlphaFold Protein
Structure Database: Massively expanding the structural coverage of protein-sequence space with
high-accuracy models. Nucleic Acids Res. 2022 Jan 7;50(D1):D439–44.

44.   Zhu K, Day T, Warshaviak D, Murrett C, Friesner R, Pearlman D. Antibody structure determination
using a combination of homology modeling, energy-based refinement, and loop prediction. Proteins:
Structure, Function and Bioinformatics. 2014;82(8):1646–55.

45.   Gerlt JA, Bouvier JT, Davidson DB, Imker HJ, Sadkhin B, Slater DR, et al. Enzyme function
initiative-enzyme similarity tool (EFI-EST): A web tool for generating protein sequence similarity
networks. Vol. 1854, Biochimica et Biophysica Acta - Proteins and Proteomics. Elsevier B.V.; 2015.
p. 1019–37.

46.   Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, et al. Cytoscape: A software
Environment for integrated models of biomolecular interaction networks. Genome Res. 2003
Nov;13(11):2498–504.

47.   Criscuolo A, Gribaldo S. BMGE (Block Mapping and Gathering with Entropy): a new software for
selection of phylogenetic informative regions from multiple sequence alignments. BMC Evol Biol.
2010;10:210.

48.   Price MN, Dehal PS, Arkin AP. FastTree 2 – approximately maximum-likelihood trees for large
alignments. PLoS One. 2010 Mar 10;5(3):e9490.

49.   Letunic I, Bork P. Interactive tree of life (iTOL) v5: An online tool for phylogenetic tree display

532   and annotation. Nucleic Acids Res. 2021;49(W1):W293–6.

533   50.   Harrison KJ, de Crécy-Lagard V, Zallot R. Gene Graphics: A genomic neighborhood data

534   visualization web application. Bioinformatics. 2018;34(8):1406–8.

535   51.   Kelley LA, Mezulis S, Yates CM, Wass MN, Sternberg MJE. The Phyre2 web portal for protein

536   modeling, prediction and analysis. Nat Protoc. 2015 Jun 30;10(6):845–58.

537   52.   Battisti JM, Minnick MF. Laboratory maintenance of *Bartonella quintana*. Curr Protoc Microbiol.

538   2008 Aug;10(1):3C.1.1-3C.1.13.

539   53.   Green R, Rogers EJ. Transformation of chemically competent *E. coli*. In: Methods in Enzymology.

540   Academic Press Inc.; 2013. p. 329–36.

Figure legends

Fig 1. Known or predicted Q synthesis pathways.

(A) $preQ_0$/$preQ_1$ synthesis and salvage pathways in *E. coli*; (B) q salvage pathway in *C. trachomatis*; (C)

$preQ_1$, q and Q salvage pathways in *C. difficile*. The ECF transporters include 4 subunits: S, the substrate-

specific transmembrane component (QueT); T, the energy-coupling module; A and A', a pair of ABC

ATPase. (D) Possible q and Q salvage pathways in *B. henselae* Houston 1.

Figure 2. Protein sequence similarity network (SSN) of 7,625 QPTR (PF02592 family) proteins.

Each node in the network represents a QPTR protein. An edge (represented as a line) is drawn between two

nodes with a BLAST *E*-value cutoff of better than 10^{-70} (alignment score of 70). The nodes are colored

based on the presence/absence of the other Q synthesis genes in the corresponding genome. Cases with

inconclusive Q pathway gene distribution are colored in light gray. QPTR proteins without genome

information are colored in dark gray. QPTR homologs from *E. coli* and *C. trachomatis* are indicated by

yellow and blue arrows, respectively. The solitary TGTs from genomes harboring only *tgt* that cluster with

TGTs in complete Q pathways are circled in blue.

Figure 3. Comparison of the phylogenetic tree of taxonomic ranks of bacteria that harbor *tgt* only

and the maximum likelihood tree of their TGT proteins.

The maximum likelihood tree of marker proteins of taxonomic ranks of bacteria that harbor *tgt* only without

other Q proteins (left). The maximum likelihood tree of TGT proteins from organisms of each

corresponding rank (right). Human and *Pyrococcus horikoshii* tRNA-guanine (15) transglycosylases were

used as the outgroup for each tree, respectively. TGT of *E. coli* was used for comparison. Bootstraps above

0.5 are shown under the branch. A multiple sequence alignment of consensus residues in the TGT proteins

in the moiety of 7-substitue of deazapurine is shown (middle). The residues are highlighted by the percent

identity. The dashed lines connect each organism and their TGT sequence. The branch of TGTs from

Bartonella, Pelagibacter, E. coli, and humans is highlighted in red.

Figure 4. Bh TGT and Bh YhhQ salvage preQ$_1$ in *E. coli*.

Detection of Q-tRNAAsp GUC by the APB assay. Q-modified tRNAs that migrate slower are indicated by

an arrow. tRNAs were extracted from WT and mutant strains expressing different Q salvage genes. The

strains used are denoted in the first line. The genes and corresponding vectors are indicated in the second

line. Plasmid and strain information are given in Table S9. Cells were grown in minimal media in the

presence of exogenous preQ$_0$, preQ$_1$, q or Q as noted. DMSO was used as control when no deazapurine

was supplemented.

Figure 5. Quantification of preQ1 and Q in *B. henselae*.

LC-MS/MS analysis of ribonucleosides containing preQ1 and Q was performed on 600 ng of hydrolyzed small RNA, with signal intensities normalized to the sum of UV absorbances of the canonical ribonucleosides, using an inline UV detector, to correct for differences in amounts of injected RNA. Data represent mean ± SD for three biological replicates of *B. henselae* BH-WT+preQ1 (BH-WT+preQ1) and mean ± deviation about the mean for two biological replicates of *B. henselae* BH-WT (BH-WT); technical duplicates were performed for each biological replicate. All modifications were validated with standards.

Figure 6. Phylogenetic analysis of Queuosine biosynthesis genes in *Bartonellaceae*. The clade of *Bartonella spp.* (*Bartonellaceae*; Alphaproteobacteria; left). The branches of *Pseudochrobactrum, Brucella*, and *Ochrobactrum* were collapsed. The highlighted node 1 corresponds to the node in the tree in Figure S5. The presence of Q biosynthesis proteins is indicated by the circles with TGTs highlighted in red. The comparative view of the corresponding truncated and full-length *tgt* gene variants in different *Bartonella* species in the tree (right). Red, *tgt*; yellow, *queA*; blue, genes encoding a transporter-like protein; black, hypothetical genes; gray, other genes. The fragmented *tgt* genes in *B. quintana* strains are boxed. Gene IDs are provided in supplementary Table S4.

Figure 7. The prevalence of Q pathway proteins in selected taxonomic ranks in Uniprot database.

(A) A cladogram of the order Hyphomicrobiales. (B) A cladogram of the order Mycobacteriales. (C) A cladogram of the class Spirochaetia. (D) A cladogram of the order Rickettsiales. Each pie chart represents the percentage of organisms in each taxonomic rank that contains the corresponding Q pathway protein. The size of each triangle correlates with the number of genomes in each taxonomic unit. The sequences of TGTs' substrate binding sites from ranks that encoded only *tgt* were analyzed and presented in Table. S2. Only taxonomic ranks that contain more than 15 TGTs are shown.

Figure 1

Figure 2

Genotype | Substrate of TGT
FolE1/2,QueD,QueE,QueC,QueF,TGT,QueA,QueG/H | preQ$_1$/preQ$_0$
QueF,TGT,QueA,QueG/H | preQ$_1$/preQ$_0$
TGT,QueA,QueG/H | preQ$_1$
TGT | q
Other than above | n.a.
Lack of genome information | n.a.

◁ E.Coli
◁ Chlamydia trachomatis

Figure 3

Figure 4

A

WT | Δtgt | *ΔqueD ΔqueF Δtgt*::Kan pBAD33::Ct *yhhQ*

pBAD24 | pBAD24::Bh *tgt*

10 | 100 | 0 | 10 | 100 | preQ₁ (nM)

← Q-tRNA
← No Q

B

ΔqueD ΔyhhQ::Kan

pBAD33 | pBAD33::Bh *yhhQ* | pBAD33 | Δtgt | ΔqueD | WT

| 0 | 10 | 100 | | 100 | | | |
100 | 0 | | | | | 10 | 100 | | 100 | preQ₀ (nM)
preQ₁ (nM)

← Q-tRNA
← No Q

C

ΔqueDΔqueFΔyhhQΔtgt::Kan | Δtgt | WT

pBAD33 | Ct *yhhQ* | Ec *yhhQ* | Ct *tgt* | Bh *yhhQ*
pBAD24 | Ct *tgt* | Ct *tgt* | Ec *yhhQ* | Bh *tgt*

q (nM) | 100 | 5000 | 1000 | 5000 | 1000 | 2000 | 1000 | 500

Q-tRNA →
No Q →

D

WT | Δtgt | *ΔqueDΔyhhQ*::Kan | *ΔqueDΔyhhQ*::Kan

pBAD24 | | | *queKqueLqueT* | *queKqueLqueT*
pBAD33 | | | Bh *yhhQ* | Cd ECF-TAA' *queT*

Q (nM) | | | 500 | 500 | 5000 | 5000 | 100 | DMSO

Q-tRNA →
No Q →

Figure 5

Figure 6

Tree scale: 0.1

Sale: ⊢—⊣ 1kB

Figure 7