Post-GWAS Prioritization of Genome-Phenome Associations in Sorghum

Debasmita Pal[1*], Kevin Schaper[2], Addie Thompson[3], Jessica Guo[4], Pankaj Jaiswal[5], Curtis

Lisle[6], Laurel Cooper[5], David LeBauer[4], Anne Thessen[2†], Arun Ross[1*†]

[1]Department of Computer Science and Engineering, Michigan State University, East Lansing,

Michigan, USA

[2]Translational & Integrated Sciences Laboratory, University of Colorado Anschutz Medical Campus,

Department of Biomedical Informatics, Aurora, Colorado, USA

[3]Department of Plant, Soil and Microbial Sciences, Michigan State University, East Lansing,

Michigan, USA

[4]Arizona Experiment Station, University of Arizona, Tucson, Arizona, USA

[5]Department of Botany and Plant Pathology, Oregon State University, Corvallis, Oregon, USA

[6]KnowledgeVis, LLC., Altamonte Springs, Florida, USA

[†]These authors share last authorship

* Correspondence: paldebas@msu.edu, rossarun@cse.msu.edu

Keywords: plant phenotype, maximum canopy height, maximum growth rate, *Sorghum bicolor*,

SNP, GWAS, pattern recognition, feature engineering

Abstract

This study presents an innovative approach for understanding the genetic underpinnings of two key

phenotypes in *Sorghum bicolor*: maximum canopy height and maximum growth rate. Genome-Wide

Association Studies (GWAS) are widely used to decipher the genetic basis of traits in organisms, but

the challenge lies in selecting an appropriate statistically significant threshold for analysis. Our goal

was to employ GWAS to pinpoint the genetic markers associated with the phenotypes of interest

using specific permissive-filtered threshold values that allows the inclusion of broader collections of

explanatory candidate genes. Then, we utilized a pattern recognition technique to prioritize a set of

informative genes, which hold potential for further investigation and could find applications in

Artificial Intelligence systems. Utilizing a subset of the Sorghum Bioenergy Association Panel

cultivated at the Maricopa Agricultural Center in Arizona, we sought to unveil patterns between

phenotypic similarity and genetic proximity among accessions in order to organize Single Nucleotide

Polymorphisms (SNPs) which are likely to be associated with the phenotypic trait. Additionally, we

explored the impact of this method by considering all SNPs versus focusing on SNPs classified

through the GWAS pre-filter. Experimental results indicated that our approach effectively prioritizes

SNPs and genes influencing the phenotype of interest. Moreover, this methodology holds promise in

the feature selection from genomic data for predicting complex phenotypic traits influenced by

numerous genes and environmental conditions and could pave the way for further research in this

field.

Author Summary

Analyzing the relationships of an organism's phenotypes with their genotypes and environments

stands as a critical aspect in plant biology. Over the past few decades, researchers have focused on

identifying impactful genes controlling the plant phenotypes in diverse environmental conditions, a

pursuit especially relevant in the face of global climate change. Applied field experiments and

40 quantitative genetics have identified numerous regions of the genome influencing complex traits.

41 However, the substantial time and cost required for physical experiments have prompted the use of

42 computational methods to facilitate this field of research. Our work aims to unravel the patterns

43 between phenotypic similarity and genetic proximity across a panel of sorghum accessions to

44 prioritize genes influencing the phenotype of interest. Our results suggest that this method is an

45 effective way to prioritize informative genes for further analysis, and for genomic features selection

46 that can be used by machine learning and deep learning algorithms for predicting complex

47 phenotypic traits.

48 1. Introduction

49 To mitigate the effects of climate change and population growth, we need to better understand the

50 dynamic interplay between an organism's genome, phenome, and environmental effects. Artificial

51 Intelligence (AI)-based systems using machine learning (ML) and deep learning (DL) algorithms

52 have shown great promise in capturing the complex and nonlinear response of natural systems and

53 genomics (Lin & Ngiam, 2023, Ghannam & Techtmann, 2021; Hsieh, 2022; Thessen, 2016), but

54 several barriers still exist (Danilevicz et al, 2022), including the absence of AI-ready data for most

55 biological applications. This is especially true for genomic data, which contains millions of potential

56 features that can quickly overwhelm an AI model. Therefore, genomic data needs filtering through an

57 automated pattern recognition process to identify the most distinguishing components for use in

58 training the ML/DL algorithms. For model organisms, this can be a simple process of selecting data

59 from genome regions that are known to be causal for the phenotype of interest, but it is challenging

60 for complex phenotypes, such as growth rate and biomass accumulation, which are regulated by

61 molecular interactions and biological pathways involving many genes, and greatly influenced by

62 environmental conditions.

Genome-wide association study (GWAS, Visscher et al., 2012) is a widely-used technique to deduce

the genetic basis of phenotypic traits in organisms. However, the challenge lies in selecting an

appropriate statistically significant threshold (p-value) for the analysis, considering the false

discovery rate (Hebbring, 2014). A stringent significance threshold for associations would possibly

miss causative variance present at lower frequencies in the population. Other methods to decrease

genomic feature set size, for example, correlation with the trait of interest, would be potentially

confounded with population structure. Though these features would potentially be highly predictive

of the phenotypes, they would likely not lend insight into underlying biology and gene function -

only to breeding history or source of accessions in the population. Thus, our approach was to use

certain permissive-filtered GWAS thresholds that retain many more features than what the GWAS

analysis itself would have identified as significantly associated with the phenotypic trait. This would

likely increase false positives as well as true positives and lead to subsets of potentially informative

SNPs, whose association with the phenotypes was presumably not confounded with the population

structure.

On top of the genetic markers identified by permissive-filtered GWAS thresholds (allowing more

explanatory candidate genes), we utilized a pattern recognition technique based on statistical

correlation coefficient to establish genome-phenome associations in the context of environmental

variation. The idea was to compare the resemblance of phenotypic similarity determined by the

normalized phenotypic trait measurements among accessions with their genetic relatedness. Through

this process, our objective was to prioritize Single Nucleotide Polymorphisms (SNPs) and genes,

which are likely to be correlated with the phenotype under study. Our algorithm also incorporated a

kind of ordering for the prioritized SNPs based on their impact on phenotypic traits, which alleviates

the hurdle of discovering SNPs with lower effect estimates through GWAS (Moore et al, 2010). We

considered the phenotypes, viz. maximum canopy height and maximum growth rate of *Sorghum*

bicolor, to examine the usefulness of our method. Empirical evidence in the form of ontology-based

gene functions, spatial and temporal expression, and similarity to known homologs were curated,

describing that our methodology shows a promising avenue to extract the genes on precedence that

may be informative for the studied phenotype. We hypothesize that these SNPs and genes could

potentially be considered for new plant breeding and genomics projects to empirically validate their

functional annotations.

Moreover, the prioritized SNPs and genes extracted by our approach could be utilized to improve the

performance of ML/DL algorithms. At the outset of this work, we envisaged to develop ML models

to predict the phenotype given information about genomes and environments. Our study focused on a

subset of the Sorghum Bioenergy Association Panel, grown at the Maricopa Agricultural Center in

Arizona, as part of the TERRA-REF project. Throughout the growing season, various phenotype

measurements were collected regularly by an automated field scanner system and field scientists. The

difficulty to train ML models based on this dataset was the size of an entire genome compared to the

availability of a limited number of phenotypic observations. In general, it is hugely expensive to

cultivate at such a large scale to obtain a comparable number of phenotypic measurements with

genetic makeup, necessitating some method of genomic feature selection. In GWAS, prefiltering

(e.g., on missingness, error probability, linkage disequilibrium, minor allele frequency, or other

metrics) could be a favorable way to reduce the dimensionality of the datasets (the number of SNPs),

however, they manifested mixed outcomes (Danilevicz et al, 2022, Rice and Lipka, 2019), and the

best approach for doing so is still a subject of research.

We primarily intended to leverage GWAS in our experiment as it has already shown its capability in

finding causal SNPs for complex phenotypic traits and posed our method as post-GWAS analysis

upon the selection of lower stringent thresholds. Our approach aims to establish the associations

between genotypes and phenotypes based on the statistical correlation in contrast to regression,

which was mostly used by genome-phenome association studies, e.g., GWAS and any BLUP-based

methods (Zhang et al., 2022). Regression analysis typically derives a specific equation to express the

relationship between independent and dependent variables, whereas correlation helps in determining

if at all a relationship exists between two independent distributions, and their outcomes might vary

depending on the patterns present in the dataset. Furthermore, we analyzed the relationships at the

granularity of every pair of accessions rather than at an accession level. Additionally, the use of

normalized phenotypic trait measurements introduced more resilience to outliers in our experiments

and made the extracted results more comparable across different environmental conditions. We also

explored the prospect of our method by investigating the difference in results with applying GWAS

pre-filter and without GWAS pre-filter.

Background:

The most intensive and comprehensive studies of genotypes, environments, and phenotypes are being

conducted in agricultural research, where plant breeding programs and seed and livestock companies

try to find genes or combinations of genes for desired phenotypes (Lobos & Hancock, 2015; Lopes et

al., 2015). The most important traits of interest in agricultural plant breeding, such as growth, plant

height and yield are generally determined by suites of genes, rather than a single gene, and are

influenced by environmental factors. For example, in soybean, Cao et al. (2019) found a number of

quantitative trait loci (QTL) implicated in controlling plant height with environmental and epistatic

interaction effects. This complicates the process of identifying informative areas of the genome and

leads to the development of novel approaches.

In sorghum, QTL analysis was used to identify genomic regions important for plant height (Brown et

al., 2008), early-season cold tolerance (Burow et al., 2011), and Ma2, a regulator of flowering time

(Casto et al., 2019), among others. Using a biocuration strategy, Mace et al. (2019) compiled a

comprehensive QTL Atlas (https://aussorgm.org.au/sorghum-qtl-atlas/) with the projected locations

of ~6000 QTL on the sorghum reference genome by data mining approximately 150 peer-reviewed

publications in sorghum from 1995 to the present. Another approach involved the development of

populations of sorghum plants with induced mutations, for example, the 6400 ethyl methane

sulfonate-mutagenized lines developed by Jiao et al. (2016), which was utilized to identify genes

involved in drought tolerance (Jiao et al., 2016) and spikelet fertility (Gladman et al., 2019).

With the advent of whole genome sequencing in sorghum (Mace et al., 2013; McCormick et al.,

2018; Boyles et al., 2019; Cooper et al., 2019), Single Nucleotide Polymorphisms (SNPs) and

insertions/deletions were identified and can be associated with variations in traits or phenotypes of

interest (Rafalski, 2002). This led to newer approaches such as association mapping (Zhu et al.,

2008) and GWAS (Visscher et al., 2012). In sorghum, GWAS has been utilized for determining the

genetic basis for such complex phenotypes as root system architecture (Zheng et al., 2020), grain

polyphenol content (Rhodes et al., 2014), grain size (Tao et al., 2020), disease responses (Ahn et al.,

2019), and grain yield (Boyles et al., 2019).

Association studies in specialized panels or populations led to advances in our understanding of plant

height (Brown et al., 2008), grain quality (Sukumaran et al., 2012), and bioenergy traits (Brenton et

al., 2016). Other methods based on whole genome analysis, including such approaches as genomic

selection (Meuwissen et al., 2001; Budhlakoti et al., 2022) and pan-genome analysis (Tao et al.,

2021, Naithani et al., 2023), were also utilized.

Further, to improve phenotyping for the genomic prediction model, Watanabe et al. (2017) suggested

a high-throughput phenotyping system for sorghum plant height, a complex quantitative trait by

using unmanned aerial vehicle (UAV) imagery. Researchers also attempted to apply ML approaches

for genomic selection that primarily rely on the availability of genomic data to predict phenotypic

traits (Tong and Nikoloski, 2021). Costa et al. (2022) proposed genomic prediction for traits

modulated by epistatic effect by applying ML and artificial neural networks. Their comparative study

indicated better performance with neural networks for complex quantitative traits. Further, Jubair and

Domaratzki (2023) presented a comprehensive overview of deep learning approaches in the genomic

selection process by incorporating weather data. Similarly, Adak et al. (2021), using their UAV-

based weekly observance of plant height phenotype for a growing season of the maize crop, observed

that the two SNP markers previously identified by GWAS methods show a greater effect on early-

season growth, depicting a more dynamic relationship, that is often not observed in classical

experiments. These approaches seek to identify important genes by accounting for the variability in

the environment.

2.   Materials and Methods

Our Approach:

To establish genome-phenome associations, we compared the phenotypic similarity of each pair of

accessions with their genetic relatedness in terms of matching alleles at a particular SNP location on

the chromosome. We utilized the differences among genetically diverse accessions in the normalized

phenotypic trait measurements to build a "target" similarity matrix describing the proximity of each

pair of accessions. This is based on the assumption that pairs of accessions showing small differences

in phenotype are most likely to be genetically similar. The normalization was obtained by fitting a

logistic growth curve model to the measurements. The next step was to construct a similarity matrix

corresponding to each SNP identified across the same set of accessions to discover which similarity

matrices are analogous to the "target" similarity matrix. This was achieved by feature engineering

using Pearson correlation coefficient, which helped in prioritizing a set of SNPs likely to be

informative about the phenotype under study. Figure 1 demonstrates the pipeline of our approach.

We utilized permissive-filtered GWAS thresholds at various p-value levels resulting in multiple sets

of potentially informative SNPs to be included in our experiment. We also compared these results

with that obtained considering all the variants across all chromosomes without applying GWAS pre-

filter. In the following, we have provided a detailed description of the datasets and the methods used

in our experiments. Our code and experimental results are available on GitHub

(https://github.com/genophenoenvo/phenotype-to-genotype).

Figure 1. Method Pipeline for Prioritizing SNPs. This pipeline begins with *normalized* phenotypic trait data and a set of GWAS-filtered SNPs in separate files and ends with lists of prioritized SNPs impacting the phenotypic traits.

Plant Material and Phenotype Dataset Description:

The multi-year TERRA Phenotyping Reference Platform (TERRA-REF) program (Burnette et al.,

2018; LeBauer et al., 2020) studied how the environment and other factors affected the growth of

agricultural products. An underlying goal of the program is to study how to maximize growth and, as

a result, maximize agricultural bioenergy output. Season 6 of the program was held in the summer of

2017. *Sorghum bicolor* was chosen as the target organism for this study because of its drought

tolerance, variety of food, fuel, and fiber applications, and potential improvement through targeted

breeding. A subset of 336 sorghum accessions from the Bioenergy Association Panel (BAP) (Brenton

et al., 2016) were planted in the study field in Arizona (Maricopa Agricultural Center). An automatic

large field scanner system observed these plants for phenotype measurements every day or two

throughout the growth season. In addition to comparing automated observations, a small number of

common phenotypes were also recorded by field scientists.

A suite of measurements, including photographs, canopy height, leaf characteristics, and soil

environmental factors, were captured regularly by automated systems to analyze each accession's

growth patterns (Burnette et al., 2018). Two replicate plots of each accession were planted within a

200m x 20m field. Data collected by TERRA-REF is available in the public domain, along with

experimental design, measurement methods, and other study metadata (LeBauer et al., 2020). Our

analysis used canopy height phenotype and whole-genome resequencing data. Plot-level canopy

height measurements were estimated from laser scans using an algorithm calibrated to hand

measurements. Canopy height measurements are available for most days during the growing season.

Genomics data are from whole-genome resequencing of the BAP genotypes included in the study.

To test the impact of a different environment, we used end-of-season height data from the same

accessions grown in Florence County, South Carolina (Brenton et al., 2016).

Phenotype Data Preparation and Normalization:

Automated measurements of canopy height over the growing season were fitted with logistic growth

curves to obtain the phenotypic traits of maximum canopy height and maximum growth rate relative

to growing degree days (gdd), defined as the accumulation of heat energy above 10°C following the

"Method 1" described by McMaster & Wilhelm (1997). This approach can yield phenotypic traits

that are less sensitive to outliers, more comparable across environmental conditions, and permit

greater biological inference.

First, we applied a cleaning algorithm that excluded accessions having a number of automatic

measurements per season of less than 35 or having the measurements on fewer than 40% of days,

which resulted in a total of 326 accessions. This preprocessing yielded a time series that could

robustly estimate the parameters of maximum canopy height and maximum growth rate.

Time series of canopy height and growing degree days (gdd) for each accession were modeled in a

Bayesian framework with the likelihood:

$$height_i \sim Normal(\mu_i, \sigma^2) \qquad (1)$$

223   where, μ is the expected value of height, σ^2 is the measurement error variance, and i indexes each

224   observation. The logistic model was defined as:

$$\mu_i = \frac{c}{(1+ae^{b*gdd})} \qquad (2)$$

226   with the logistic parameters a, b, and c and the covariate *gdd* associated with each observation.

227   However, we re-parameterized the model to obtain more biologically meaningful parameters by

228   defining:

$$Y_{max} = c \qquad (3)$$

$$Y_{min} = \frac{c}{(1+a)} \qquad (4)$$

$$R_{half} = -\frac{c*b}{4} \qquad (5)$$

232   Finally, we transformed the logistic parameters to the whole real line, which allowed for faster

233   mixing and improved model convergence:

$$c = e^{\theta_c} \qquad (6)$$

$$a = e^{\theta_a} - 1 \qquad (7)$$

$$236 \qquad b = -e^{\theta_b} \qquad\qquad (8)$$

All root nodes were given wide, relatively non-informative standard priors, including

Normal(0,1000) for all transformed logistic parameters (θ_a, θ_b, θ_c) and *Gamma(0.1, 0.1)* for the

observation precision $\left(\frac{1}{\sigma^2}\right)$, where *Normal()* and *Gamma()* refer to probability distributions.

We implemented the above model in JAGS 4.3.0 (Plummer, 2003) via R/rjags (Plummer et al. 2022;

R Core Team, 2021). Three parallel Markov Chain Monte Carlo (MCMC) sequences were assigned

with dispersed starting values. The models were run until convergence was achieved at a Gelman and

Rubin (1992) diagnostic parameter of < 1.2. An additional run of 10,000 iterations with the thinning

parameter as 10 yielded a total of 3,000 relatively independent posterior samples for each parameter

of interest. The posterior median of Y_{max} (maximum canopy height, cm) and R_{half} (maximum growth

rate, cm/*gdd*) of the accessions were considered to estimate the "target" similarity matrix for the

corresponding phenotypic traits and in the GWAS analysis (described below). Figure 2 shows the

logistic growth curve obtained for two sample accessions. The data and code for the logistic growth

curve model are publicly available (Guo & LeBauer, 2022).

Estimation of "Target" Similarity Matrix:

Based on the above-mentioned, *normalized* phenotypic measurements of the *Sorghum* accessions, we

computed the "target" similarity matrix for each of the phenotypes (maximum canopy height and

maximum growth rate) *separately*. We assumed that the accessions having close genetic proximity

with each other will exhibit a smaller difference in their phenotypic observations. Hence, this matrix

helped us estimate the potential correlation coefficient values among accessions. The rows and

columns of this matrix correspond to the accessions and the values indicate the "target" similarity

Figure 2. Example of logistic growth curves to illustrate phenotypes of two sorghum accessions. Using automated canopy heights for each accession, logistic growth curves were fitted and produced maximum canopy height (Y_{max}) and maximum growth rate (R_{half}). The cleaning algorithm applied ensured a sufficient number of observations to produce confident estimations of the parameters.

257 value (potential correlation coefficient), with a range of [0,1]. The higher the similarity value, the

258 greater is the similarity between the accessions, based on the phenotype under study.

259 The "target" similarity value between the accessions *'i'* and *'j'* was calculated as follows:

260  1) First, the difference (x_{ij}) in their phenotypic measurement was computed.

261  2) Min-max scaling was applied to the difference (x_{ij}), using the following equation, in order to

262   scale the difference in the range [0,1]:

263
$$x_{ij}{}^{scaled} = \frac{x_{ij} - x_{min}}{x_{max} - x_{min}} \qquad (9)$$

264 where, x_{min} and x_{max}, respectively, denote the minimum and maximum values obtained over all x_{ij}'s.

3) The scaled difference was subtracted from 1 ($x_{ij}' = 1 - x_{ij}^{scaled}$), so that a higher value indicates a

greater similarity between accessions.

Genomic Data Preparation and GWAS:

Throughout the project, we used the Sorghum BTX623 reference genome v3.1, and the analysis of

function (described in the "SNP Annotation and Function" subsection) was carried out on either

Phytozome (Goodstein et al., 2012), Gramene (Tello-Ruiz et al., 2021), Planteome (Cooper et al.,

2018) or the Gene Ontology (The Gene Ontology Consortium, 2021) online databases. Whole

genome sequencing data was obtained from the TERRA-REF project (LeBauer et al., 2020,

https://docs.terraref.org/protocols/genomic-data), where 384 sorghum accessions from the sorghum

Bioenergy Association Panel (BAP) were sequenced with ~25x coverage. Shotgun sequencing (127-

bp paired-end) was done using an Illumina X10 instrument at the HudsonAlpha Institute for

Biotechnology. After aligning the sequence reads (BTX623 reference genome v3.1 available from

Phytozome), GVCF files with variant calls were downloaded from CyVerse, merged, and filtered

using protocols described in the TERRA-REF documentation. SNPs were filtered using GATK

Variant Filtration, and accessions without phenotypic observations were removed, and only biallelic

sites were considered.

GWAS was performed using rMVP (Yin et al., 2021) with the FarmCPU algorithm (Liu et al., 2016)

for each of the normalized phenotypes (maximum canopy height and maximum growth rate) using a

kinship matrix and 1 principle component to account for relatedness and structure within the

population. The results were filtered with p-values of .0001, .0005, and .001 to create six different

lists of SNPs (Supplementary File). Additionally, these six lists were filtered against QTL regions

known to be associated with height and growth obtained from the Sorghum QTL Atlas (Mace et al.,

2019). After this step, we had 12 lists of SNPs, which were utilized to filter the GVCF file to obtain

12 VCF files, hereafter, referred to as experimental conditions (Table 1 - Total #SNPs).

Table 1. Experimental Summary and Resulting SNP Lists using Permissive-Filtered GWAS Thresholds

p-value	Phenotype	QTL Filtering	Total #SNPs[1]	Prioritized #SNPs[2]	Prioritized #SNPs directly map to known genes	Highest Correlation Coefficient[3]
p0001	Maximum Growth	w/ QTL	534	77	18	0.41
		No QTL	1,055	107	28	0.49
	Maximum Height	w/ QTL	4,630	286	69	0.61
		No QTL	6,003	323	84	0.63
p0005	Maximum Growth	w/ QTL	2,861	174	30	0.55
		No QTL	5,575	198	48	0.60
	Maximum Height	w/ QTL	15,888	538	119	0.67
		No QTL	20,899	537	133	0.68
p001	Maximum Growth	w/ QTL	6,106	230	47	0.59
		No QTL	11,698	326	82	0.64
	Maximum Height	w/ QTL	27,536	634	140	0.68
		No QTL	36,033	755	182	0.69

[1] Total number of SNPs identified through GWAS.
[2] Number of prioritized SNPs extracted.
[3] Pearson correlation coefficient of "target" similarity matrix with the summation of SNP similarity matrices corresponding to the prioritized SNPs.

Creation of SNP Similarity Matrices:

The above VCF files were processed separately to create a similarity matrix for each SNP, comparing

the allelic similarity of every pair of accessions at that location on the chromosome. The values of

these similarity matrices were computed with an addition of 0.5 for each matching allele between any

pair of accessions, along with reversed heterozygous SNPs receiving a value of 1.0. For instance, if a

single VCF file consists of 'N' number of SNPs across 'a' number of accessions, there would be a

total of 'N' SNP similarity matrices; each of these similarity matrices is a square matrix of

dimensions 'a x a', and their cell values encoded among three discrete values (0, 0.5, 1.0). Out of the

326 accessions obtained as part of the logistic growth model, we used 274 accessions that were

common to both Seasons 4 and 6 of the TERRA-REF project for generating the similarity matrices

(common set of accessions were considered for future work). Thus, each SNP similarity matrix has

dimensions of 274 x 274 in our experiment.

Prioritization of SNPs:

Next, we compared the corresponding SNP similarity matrices (described above) to the phenotype

similarity matrix ("target" similarity matrix) using the algorithm described below. The notion was to

find out the SNP similarity matrices wherein the variation between the accessions is similar to that in

the phenotypic measurements. Pearson correlation coefficient ($\rho_{(X,Y)}$) was used to measure the

relationship between two data matrices. It is defined by the covariance of two data matrices X and Y

divided by their standard deviations.

$$\rho_{(X,Y)} = \frac{E[(X-\mu_X)(Y-\mu_Y)]}{\sigma_X \sigma_Y} \qquad (10)$$

where, μ_X and μ_Y are, respectively, the mean and, σ_X and σ_Y are, respectively, the standard deviations

of the data matrices X and Y. We followed the feature selection technique, known as Sequential

Forward Selection (SFS) (Pudil et al, 1994) in which the important features are sequentially selected

using an evaluation criterion and appended to the empty candidate list until the addition of further

features does not reduce the criterion.

We prioritized a certain number of SNPs from a set of SNP similarity matrices, each corresponding

to a SNP, by using Pearson correlation coefficient (Table 1 - Prioritized #SNPs). To accomplish this,

we initiated the process by defining a "*maximum desired number of prioritized SNPs*" (e.g., 1000),

and a *null* "candidate" SNP list. In the first iteration, we selected a SNP similarity matrix from the set

of SNP similarity matrices that resulted in a maximum correlation coefficient with the "target"

similarity matrix and added the corresponding SNP to the candidate SNP list. Starting from the

second iteration, we chose the SNP similarity matrix from the remaining set (i.e., the corresponding

SNP not present in the candidate SNP list), which, when summed up with the similarity matrices

corresponding to SNPs already added to the candidate SNP list, yielded in the maximum correlation

coefficient with "target" similarity matrix. Then, the corresponding selected SNP was appended to

the candidate SNP list. We continued this process iteratively, seeking the next *cumulatively impactful*

SNP and appended it to the candidate SNP list sequentially until the number of SNPs in the candidate

SNP list reached the "*maximum desired number of prioritized SNPs*". At the end, we derived a

"prioritized" SNP list by selecting the first 'K' SNPs from the candidate SNP list, whose

corresponding similarity matrices, when summed up, results in the maximum correlation coefficient

with the "target" similarity matrix. In other words, our method generated a "final" SNP similarity

matrix (sum of similarity matrices corresponding to the prioritized SNPs 'K') based on the allelic

proximities among accessions maximizing the correlation coefficient (Table 1 - Highest Correlation

Coefficient) with the "target" similarity matrix. Our algorithm is described below:

Input:

● *Similarity matrices among 'a' unique accessions corresponding to 'N' no. of SNPs (S_j,*

335   *where $j = 1, 2, ..., N$, and dimension of each S_j is a x a, and the cell values of S_j are*

336   *among three discrete values [0.0, 0.5, 1.0])*

337   ● *"Target" similarity matrix among 'a' accessions ('T' of dimension a x a)*

338   ● *Maximum desired number of prioritized SNPs 'n'*

339   **Output:**

340   ● *Prioritized SNP list 'L_K', where 'K' is the no. of prioritized SNPs and K<=n*

341   **Initialize:**

342   ● *Candidate SNP list 'L_n' $=\varnothing$*

343   **Algorithm:**

344   ● *Step 1: Compute Pearson correlation coefficient of each S_j with T, for j = 1,2,...,N.*

345   ● *Step 2: Select S_{j*} with highest correlation coefficient found in Step 1, where $j^* \in$*

346   *{1,2,...,N} and append the corresponding SNP index j^* to L_n*

347   ● *Step 3: Sum the similarity matrices corresponding to the SNP indices present in L_n*

348   *and S_j together, and compute Pearson correlation coefficient with T, where $j \in$*

349   *{1,2,...,N} and $j \not\subset Ln$*

350   ● *Step 4: Select S_j from step 3 resulting in the highest correlation coefficient value and*

351   *append the corresponding SNP index 'j' to L_n*

352   ● *Step 5: Repeat steps 3 and 4 until the number of SNPs in L_n reaches the maximum*

353   *desired number of prioritized SNPs 'n'.*

354   ● *Step 6: Select the first K no. of SNPs from L_n , the summation of whose corresponding*

355   *similarity matrices resulted in the maximum correlation coefficient value with 'T' and*

356   *assign them to L_K*

We executed the above algorithm independently for each of the experimental conditions described in

Table 1. In cases where the total number of SNPs exceeded 1000, the *maximum desired number of*

prioritized SNPs (n) was set to 1000. We observed that this limit did not impact the prioritization of

informative SNPs as the highest correlation coefficient was consistently achieved at an index below

1000 for all experimental conditions. Figure 3 shows the trend of the correlation coefficient over the

iterations while adding SNPs to the candidate SNP list for the experiment based on maximum canopy

height with a *p*-value of .001 without QTL filtering. Further, our prioritized set of SNPs provides a

sort of ranking based on their influence on the phenotypic trait as they have been extracted through a

sequential algorithm. Hence, while doing analysis on these prioritized SNP sets (discussed later in

this paper), we used the term "Top k", which refers to the first 'k' SNPs in the prioritized SNP list.

Figure 3. Trend of correlation coefficient while adding SNPs to the candidate SNP list based on their similarity matrices considering the phenotype maximum height with GWAS pre-filtering at p-value of .001.

Prioritizing SNPs without GWAS Pre-filter:

Additionally, we conducted an experiment on the entire set of SNPs without GWAS pre-filter to

compare the results with that of the GWAS-filtered sets of SNPs for maximum canopy height. As we

were unable to process the whole GVCF file at once due to the larger volume and limited

computational resources, 10 VCF files containing SNPs (Table 2 - Total #SNPs) corresponding to

each of the chromosomes were created (Supplementary File). The goal was to first process each VCF

file individually to prioritize SNPs corresponding to each chromosome with our algorithm (described

above). During these executions, we set the *maximum desired number of prioritized SNPs (n)* as

1000, similar to the executions with GWAS-filtered SNP set. Next, we merged the prioritized lists of

SNPs from each chromosome together and executed the same algorithm again on the combined set of

SNPs to obtain a final set of prioritized SNPs for the analysis. While applying the algorithm to the

combined prioritized set of SNPs from all chromosomes, we iterated through the entire list as the

total number of SNPs was already reduced during chromosome-wise execution, i.e., in this case, the

maximum desired number of prioritized SNPs (n) was set to the total number of SNPs in the set.

Table 2. Experimental Summary and Resulting SNP Lists using all Variants without GWAS pre-filter

Chromosome No.	Phenotype	Total #SNPs[1]	Prioritized #SNPs[2]	Prioritized #SNPs directly map to known genes	Highest Correlation Coefficient[3]
01	Maximum Height	106,824	595	259	0.64
02	Maximum Height	157,148	753	256	0.64
03	Maximum Height	132,661	656	216	0.64
04	Maximum Height	126,273	690	208	0.64
05	Maximum Height	157,930	758	198	0.64
06	Maximum Height	157,808	649	121	0.64
07	Maximum Height	119,573	635	136	0.64
08	Maximum Height	124,899	644	153	0.63
09	Maximum Height	124,843	679	145	0.63
10	Maximum Height	128,622	587	171	0.63
"Combined prioritized" SNPs from all chromosomes	Maximum Height	6646	987	247	0.71

[1] Total number of SNPs.

[2] Number of prioritized SNPs extracted.

[3] Pearson correlation coefficient of "target" similarity matrix with the summation of SNP similarity matrices corresponding to the "prioritized" SNPs.

381 **SNP Annotation and Function:**

382   The prioritized SNPs extracted by the above process for each experimental condition were annotated

383   with overlapping or nearest known gene identifiers using the SnpEff package (Cingolani et al., 2012).

384   A list of unique gene identifiers for each phenotype, p-value, GWAS filter, and QTL filter was

Figure 4. Trend of correlation coefficient while prioritizing SNPs based on their similarity matrices considering the phenotype maximum height without GWAS pre-filter.

submitted for Gene Ontology (GO) Enrichment Analysis using PANTHER 17.0 (Thomas et al.,

2022) and the Sorghum bicolor reference list (Mi et al., 2019). The lists of genes to be analyzed can

be found in GitHub (https://github.com/genophenoenvo/phenotype-to-

genotype/tree/main/significant_SNPs_annotated). We did separate enrichment analyses using the

complete set of GO Biological Process complete, Molecular Function, and Cellular Component

annotation data sets. The specific test performed was Fisher's exact test with FDR correction.

The genes identified that neighbor the "Top 10" SNPs (from SnpEff) for both phenotypes with the

GWAS filtering (with and without QTL filtering) at p=.0001 were further explored for their

functional role by using Gramene and the EMBL-EBI Gene Expression Atlas. This was also done for

the "Top 10" SNPs for the height phenotype without the GWAS filtering. InterPro protein domain

annotations (Mitchell et al., 2019) were obtained by querying Gramene BioMart (Tello-Ruiz et al.,

2018, Tello-Ruiz et al., 2021). The gene IDs were also used to query the EMBL-EBI Gene

Expression Atlas (Papatheodorou et al., 2018) to explore their baseline gene expression. The baseline

expression data of genes showing a default minimum expression level of 0.1 transcripts per million

(TPM) were downloaded for experiments E-MTAB-3839 (Olson et al., 2014), E-MTAB-4400

(Davidson et al., 2012), E-GEOD-98817 (Kebrom et al., 2017). Expression heatmaps were generated

using ClustVis (Beta) (Metsalu et al., 2015). The Sorghum QTL data was queried from the Sorghum

QTL Atlas (Mace et al. 2019) and mapped on Gramene's sorghum genome browser. The SNPs were

also queried using Gramene's variant effect prediction (VEP) tool (Naithani et al. 2017; McLaren et

al. 2016) to find any overlapping genes or existing SNPs in that database.

Generalizability of Our Method:

The generalizability of this method was tested by examining the predictive value of these prioritized

SNPs on the same accessions grown in a different environment. As part of this experiment, we

created a "target" similarity matrix based on the phenotypic observations (maximum plant height)

collected in Florence County, South Carolina (Clemson University; Brenton et al. 2016) using the

process described above. Thereafter, the Pearson correlation coefficient was calculated between this

"target" similarity matrix with the "final" SNP similarity matrix (i.e., the summation of similarity

matrices corresponding to the prioritized SNPs extracted) obtained for different experimental

conditions based on the maximum canopy height of MAC Season 6 dataset.

3.   Results and Discussion

For both phenotypes, a smaller p-value resulted in fewer prioritized sets of SNPs, as expected (Table

1 - Prioritized #SNPs). Filtering by QTLs reduced the number of SNPs; however, it did not improve

the highest correlation coefficient achieved (Table 1 - Highest Correlation Coefficient). We found a

little overlap in prioritized SNPs between the two phenotypes and only at the two higher p values

(Figure 5). Further, most of the prioritized SNPs extracted without GWAS pre-filter were not

identified as informative compared to results obtained from GWAS pre-filter for maximum height

(Figure 5-D).

Figure 5. UpSet Plot showing the Overlap of Prioritized SNPs between the growth and height phenotypes. The intersection size is shown by vertical black bars (top). In the matrix, each column corresponds to a set (with the vertical black bar showing the size of the set) and each row (SNP list) corresponds to a possible intersection. The filled cells show which set is part of the intersection. Lists of prioritized SNPs for height with and without QTL filtering have high overlap, whereas the lists of prioritized SNPs for growth with and without QTL filtering have a moderate amount of overlap. At each p-value and for each phenotype, the QTL filtering added between 18% and 31% more unique SNPs. (Lex et al. 2014) Figure made with UpSetPlot v0.8.0 (https://github.com/jnothman/UpSetPlot)

(A). There is very little overlap of prioritized SNPs between the growth and height phenotype. At $p0001$ there is no overlap.

(B). At $p0005$ there are 7 SNPs in common between phenotypes.

(C). At $p001$ there are 11 SNPs in common.

(D). For the height phenotype, the majority of SNPs identified as important without the GWAS pre-filter had not been identified when using the GWAS pre-filter.

Few prioritized SNPs were mapped to known genes (Table 1 and Table 2), with the majority

mapping to intergenic regions. Enrichment analysis showed that 43 unique GO terms related to

biosynthetic processes, metabolic processes, and gene expression were over-represented in the

annotations of the genes associated with prioritized SNPs. The triterpenoid biosynthetic process

(GO:0016104) and triterpenoid metabolic process (GO:0006722) were the most over-enriched

(20.13-fold enrichment) across all the SNP lists and were located in the SNP lists for the growth

phenotype with GWAS filtering at p0.001 with and without QTL filtering. Nitrogen compound

metabolic process (GO:0006807) was one of the few under-enriched (0.72-fold enrichment) GO

annotations and was only present in SNP lists for the height phenotype at p0.0005 with and without

QTL filtering. The positive regulation of amide metabolic process (GO:0034250) was the only GO

term that was over-enriched without GWAS filtering, but not with GWAS filtering. There were more

overrepresented GO annotations for the height phenotype than for the growth phenotype - possibly

because we know more about genes for height. We did not see much difference between the

annotations enriched in prioritized SNP lists based on QTL filtering and without QTL filtering.

Of the "Top 40" prioritized SNPs extracted with GWAS pre-filter (both phenotypes with p0.0001),

all except two were in intergenic regions within 700 bp of a known transcript. One SNP,

S05_1688138, a deletion event (AATGTG→A) located on Chromosome-5 at 1688138 bp position,

was present in the intron of a gene, SORBI_3005G018800, that overlaps with QTLs known to impact

dry matter growth rate and plant height (Supplementary Figure 3). Another SNP, S06_45415791, lies

between two pre-miRNA (MIR2118) genes, ENSRNA04948199 and ENSRNA049481970, which

are conserved in plants and induce the phased small interfering RNAs (phasiRNAs) production

known to impact plant development and fertility (Supplementary Figure 4; Araki et al. 2020). A

closer look at the 85,709bp region around another intergenic SNP, S01_50208809, shows that it

overlaps with QTLs for phenotypes such as plant height, days to flowering, and tiller height. Gene

SORBI_3001G265700 flanks this SNP on the right and gene SORBI_3001G265600 on the left.

SORBI_3001G265700 encodes a serine-threonine-like kinase, and SORBI_3001G265600 encodes

an aldolase-1-epimerase like protein (SbA1ELP). The expression profile of the 51 genes that flank

and/or overlap these SNPs (according to SnpEff) showed baseline expression of 44 genes in plant

structures important for growth and height, such as the root, vegetative meristem, shoot, and stem

internodes (Supplementary Figure 5 (S5A-C); E-MTAB3839, E-MTAB-4400, and E-GEOD-98817).

Included in this list of 51 genes are the genes known to impact transcription factors, cytochrome

P450, sugar transporters, protein kinases, Leucine-rich Ser/Thr-kinase membrane proteins, and

proteins involved in regulating microtubule binding and assembly that are key to regulating mitosis

and growth during the cell cycle and thus height and growth. Most of these genes are expressed

across experiments (Supplementary Figure 5 (S5D)).

Of the "Top 10" prioritized SNPs obtained based on all variants without GWAS pre-filter, all but

three were in intergenic regions within 700 bp of a known transcript. The genes most closely

mapping to these prioritized SNPs were different from the genes identified with the GWAS pre-filter

(SORBI_3001G279900, SORBI_3002G119500, SORBI_3005G116400, SORBI_3001G262900,

SORBI_3002G153900, SORBI_3002G212200, SORBI_3006G040401, SORBI_3002G132000,

SORBI_3007G086800, SORBI_3002G119600, SORBI_3005G116500, SORBI_3007G115500,

SORBI_3001G262950, SORBI_3002G154000, SORBI_3006G040500, SORBI_3007G086900). All

these genes overlapped with known QTLs for plant height and most also overlapped with panicle

length and panicle emergence QTLs. These panicle traits may also regulate plant height in the

reproductive phase. The expression profile of these genes showed baseline expression in plant

structures important for growth and height, such as the root, vegetative meristem, shoot, and stem

internodes (Supplementary Figure 6 (S6A-C); E-MTAB-4400, E-MTAB-3839.and E-GEOD-98817).

Two of these genes (SORBI_3001G262950 and SORBI_3006G040401) did not show expression in

any of these studies. One gene (SORBI_3005G116400) showed expression in only one study (E-

MTAB-4400) where it was expressed in leaf tissues.

Priority SNPs were found near many of the genes known to affect height, but only without the

GWAS pre-filter and most of the prioritized SNPs for height did not map closely to any of the known

height genes in sorghum (Quinby and Karper, 1953; Figure 6A&B). With the GWAS pre-filter, the

only known height gene with a nearby SNP is *Ma1* (SOBIC_006G057866), which has three SNPs

nearby and is known to regulate flowering time in *Sorghum* (Quinby and Karper, 1945, Figure 6A). It

reflects the influence of flowering time on plant height. Without the GWAS pre-filter, prioritized

SNPs are identified near Ma5, Ma2, Ma1, Dw2, and Dw1 (Figure 6B). Recent work on *Sorghum*

grown in Ethiopia (Enyew et al. 2022) gives the seven most impactful SNPs on plant height, which

fall within known genes (Figure 6A&B). With the GWAS pre-filter, four of these are also near SNPs

we identified as significant for plant height (SOBIC.003G119600, SOBIC.006G111800,

SOBIC.010G085400, and SOBIC.010G066100), two are near known height genes (*DW1*,

SOBIC.009G223500 and *Ma5*, SOBIC.001G017500), and one is not near either

(SOBIC.006G235400) (Figure 6A). Without the GWAS pre-filter, an additional SNP identified in the

Ethiopian study is also near a SNP identified in this study (Figure 6B).

Based on the above data mining and literature review on known genes to support our findings, we

observed that our method is capable of prioritizing the informative genes influencing the trait of

interest leveraging GWAS, that can be utilized for the purpose of phenotype prediction. Further, the

experiment without GWAS pre-filter was able to locate certain SNPs, which are known to be

associated with the plant height phenotype. These SNPs were not retrieved after applying GWAS

with permissive-filtered thresholds. Though the experiment without GWAS pre-filter does not

explicitly account for the population structure, this analysis demonstrated the efficacy of our

approach in finding genes that may be predictive for the phenotype. Our method effectively reduced

the number of SNPs to be considered as input to ML/DL models, and population structure can be

Figure 6. Chromosomal location of prioritized SNPs with known height genes. Prioritized SNPs (red triangles) are scattered throughout the ten chromosomes. The known plant height-associated genes are labeled in blue vertical lines. The green vertical lines show the location of SNPs important for height in sorghum grown in Ethiopia (Enyew et al. 2022)

(A) The 286 SNPs identified for maximum height based on GWAS-filtered set of SNPs with QTL filtering (p-value of .0001)

(B) The 987 SNPs identified for maximum height considering all variants without the GWAS pre-filter.

accounted for later by incorporating certain techniques like clustering and principal component

analysis in the models (López-Cortés et al., 2020).

While the lists of prioritized SNPs explained about half of the variation in the phenotype in the

TERRA-REF *Sorghum* data set (Table 1 and Table 2), they had very little explanatory power for the

same phenotypes in the same accessions grown in a very different environment. There were 271

accessions in common between the Clemson and TERRA-REF datasets, and the correlation

coefficient is found to be approximately 0.07 and 0.06, respectively, for all the cases in Table 1 and

Table 2 with the Clemson "target" similarity matrix while tested with the prioritized SNPs extracted

based on MAC Season 6 dataset.

504   **4.   Conclusions**

This method is promising for prioritizing the SNPs and genes for further analysis in the domain of

functional genetics as well as identifying areas of the genome that are cumulatively informative for

predicting complex phenotypes using ML/DL methods. The highest correlation coefficient we found

was 0.69 with GWAS pre-filter (Table 1- Highest Correlation Coefficient) and 0.71 without GWAS

pre-filter (Table 2- Highest Correlation Coefficient), which are moderately strong and unsurprising

because of the influence of the environment on height and growth. We hypothesize that the total

number of SNPs that can determine height or growth is much larger than what was found in this

study, but depending on what environmental pressures the plant faces, different sub-groups of SNPs

will prove to be predictive. Our results from the South Carolina data set (Clemson) support this. In

addition, the partial overlap in important SNPs with the Ethiopia study could reflect the hot and arid

climate, which is more similar to Arizona than South Carolina despite using different accessions. A

logical next step is to find a data set from a similar environment using different accessions to see if

the significant SNPs in this study are similarly predictive.

In the absence of empirical data on gene function from sorghum, we can only conclude that this

method has the potential to identify specific areas of the genome that can help predict height and

growth phenotypes. We found gene homology and expression data supporting the importance of

some of the identified sorghum SNPs for the height and growth phenotypes. These SNPs are present

in the vicinity of gene homologs. For example, tobacco aldolase-1-epimerase (*NbA1ELP*) is known

to regulate pectin methylesterase (*NbPME*) antagonistically (Sheshukova et al. 2017). Lower

expression levels of tobacco *NbA1ELP* resulted in the dwarf plant phenotype. It gives us the

hypothesis that the sorghum ortholog of this tobacco gene (SORBI_3001G265600 *SbAiELP*) may be

associated with plant height and biomass yield in sorghum. Existing GO annotations do not include

these traits due to a lack of evidence from sorghum. However, this homologous sorghum gene is

expressed in embryo, shoot internode, root, emerging inflorescence, and stem internodes

(Supplementary Figure 5 (S5A-C)) suggesting that its expression in these tissues are likely

contributors for plant height and growth, that needs further validation by functional genomics studies.

Our original goal was to use biological knowledge graphs that link genes to phenotypes to help with

feature selection in genomic data. We quickly discovered that functional annotation of the *Sorghum*

genome is either insufficient or inferred from *Arabidopsis*, which tends to be grown in the laboratory

under standardized conditions. While genes are annotated as being useful for drought resistance or

salt tolerance, etc., there are no functional annotations of a genome that list a function in a specific

environmental context. This method could help add contextualized functional annotations to

genomes. These results have implications for using genotype-to-phenotype graphs in knowledge

guided machine learning (KGML) for phenotype prediction. Specifically, these graphs will need to

include genotype or SNP information (some biological graphs do include limited genotype

information) linked to the gene, and the functional annotations should include environmental context.

Functional annotations discovered in the laboratory can have limited use in the natural environment.

Additionally, we observed that the phenotypic dataset used in our experiment consists of accessions

having limited variation in their phenotypic measurements. Hence, our approach resulted in a "target"

similarity matrix where most of the accessions are highly correlated with each other considering the

phenotype under study. We hypothesize that the extraction of informative genes from phenotypes by

our approach would be more prominent if accessions with larger variations in their phenotypic traits

are selected.

Our method is currently limited in capturing linear relationships between genotypes and phenotypes,

and the environmental variations were guarded by normalizing the inputs itself rather than explicitly

incorporating in the procedure. In future, this algorithm could potentially be extended by using

weighted correlation metrics to capture non-linearity as well as incorporating environmental

variations explicitly in the procedure. In addition, we utilized a sequential algorithm for selecting the

significant SNPs, wherein an SNP once identified as informative, cannot be deleted from the

prioritized SNP list. In future, the process of prioritizing SNPs can be improved by employing other

feature selection techniques, viz., Sequential Floating Forward Selection (SFFS). We believe that the

work reported in this paper provides a novel path leveraging pattern recognition techniques for

prioritization of informative SNPs that can be further studied and exploited to predict the plant

phenotypes.

Acknowledgements

We thank the TERRA-REF project for sharing their high throughput phenomics and genomics data.

Funding

NSF Awards 1940059, 1940062, 1940330 Harnessing the Data Revolution

Author Contributions

Project PIs AR, AET, PJ, DL and AT conceived the idea. DL and JG collected, analyzed and

assembled the TERRA-REF project data. DP, KS, AET, JG, PJ and AT analyzed the data. LC helped

in project coordination and CL helped in data visualization. All authors contributed to combined data

analysis, writing, and editing the manuscript. All authors approved the manuscript.

Conflicts of Interest

Author Curtis Lisle is employed by KnowledgeVis LLC. The remaining authors declare that the

research was conducted in the absence of any commercial or financial relationships that could be

construed as a potential conflict of interest.

References

Adak A, Conrad C, Chen Y, Wilde SC, Murray SC, Anderson II SL,Subramanian NK (2021)

'Validation of functional polymorphisms affecting maize plant height by unoccupied aerial systems

discovers novel temporal phenotypes', G3 Genes|Genomes|Genetics, 11(6), doi:

10.1093/g3journal/jkab075

Ahn E, Hu Z, Perumal R, Prom LK, Odvody G, Upadhyaya HD, Magill C (2019) 'Genome wide

association analysis of sorghum mini core lines regarding anthracnose, downy mildew, and head

smut', PLOS ONE, 14: e0216671, doi:10.1371/journal.pone.0216671

Araki S, Le NT, Koizumi K et al. (2020) 'miR2118-dependent U-rich phasiRNA production in rice

anther wall development', Nat Commun, 11, 3115, doi:10.1038/s41467-020-16637-3

Blears MJ, De Grandis SA, Lee H, Trevors JT (1998) 'Amplified fragment length polymorphism

(AFLP): a review of the procedure and its applications', J Ind Microbiol Biotech 21:99–114,

doi:10.1038/sj.jim.2900537

Boyles RE, Brenton ZW, Kresovich S (2019) 'Genetic and genomic resources of sorghum to connect

genotype with phenotype in contrasting environments', The Plant Journal 97: 19–39,

doi:10.1111/tpj.14113

Brenton ZW, Cooper EA, Myers MT, Boyles, RE, Shakoor N, Zielinski KJ, Rauh BL, Bridges WC,

Morris GP, Kresovich S (2016) 'A Genomic Resource for the Development, Improvement, and

Exploitation of Sorghum for Bioenergy', Genetics 204: 21–33, doi:10.1534/genetics.115.183947

Brown PJ, Rooney WL, Franks C, Kresovich S (2008) 'Efficient Mapping of Plant Height

Quantitative Trait Loci in a Sorghum Association Population With Introgressed Dwarfing Genes',

Genetics, 180: 629–637, doi:10.1534/genetics.108.092239

Budhlakoti N, Kushwaha AK, Rai A, Chaturvedi KK, Kumar A, Pradhan AK, Kumar U, et al. (2022)

'Genomic Selection: A Tool for Accelerating the Efficiency of Molecular Breeding for Development

of Climate-Resilient Crops', Frontiers in Genetics, vol. 13, doi: 10.3389/fgene.2022.832153

Burnette M, Kooper R, Maloney JD, Rohde GS, Terstriep JA, Willis C, et al. (2018) 'TERRA-REF

Data Processing Infrastructure', Proceedings of the Practice and Experience on Advanced Research

Computing, USA, pp 1–7, doi: 10.1145/3219104.3219152

Burow G, Burke JJ, Xin Z, Franks CD (2011) 'Genetic dissection of early-season cold tolerance in

sorghum (Sorghum bicolor (L.) Moench)', Mol Breeding, 28:391-402, doi: 10.1007/s11032-010-

9491-4

Cao Y, Li S, Chen G, Wang Y, Bhat JA, Karikari B, Kong J, Gai J, Zhao T (2019) 'Deciphering the

Genetic Architecture of Plant Height in Soybean Using Two RIL Populations Sharing a Common

M8206 Parent', Plants, 8(10):373, doi: 10.3390/plants8100373

Casto AL, Mattison AJ, Olson SN, Thakran M, Rooney WL, Mullet JE (2019) 'Maturity2, a novel

regulator of flowering time in Sorghum bicolor, increases expression of SbPRR37 and SbCO in long

days delaying flowering', PLOS ONE, 14: e0212154, doi: 10.1371/journal.pone.0212154

Cingolani P, Platts A, Wang LL, Coon M, Nguyen T, Wang L, Land SJ, Lu X, Ruden DM (2012) 'A

program for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs

in the genome of Drosophila melanogaster strain w1118', iso-2; iso-3. Fly, 6(2), 80–92, doi:

10.4161/fly.19695

Cooper EA, Brenton ZW, Flinn BS, Jenkins J, Shu S, Flowers D, Luo F, Wang Y, Xia, P., et al.

(2019) 'A new reference genome for Sorghum bicolor reveals high levels of sequence similarity

between sweet and grain genotypes: implications for the genetics of sugar metabolism', BMC

Genomics, 20: 420, doi: 10.1186/s12864-019-5734-x

Cooper L, Meier A, Laporte M-A, Elser JL, Mungall C, Sinn BT, Cavaliere D, et al. (2018) 'The

Planteome database: an integrated resource for reference ontologies, plant genomics and phenomics',

Nucleic Acids Research, 46: D1168–D1180, doi: 10.1093/nar/gkx1152

Costa WG, Celeri MO, Barbosa IP, Silva GN, Azevedo CF, Borem A, Nascimento M, Cruz CD

(2022) 'Genomic prediction through machine learning and neural networks for traits with epistasis',

Computational and Structural Biotechnology Journal, 20:5490-5499, doi: 10.1016/j.csbj.2022.09.029

Davidson RM, Gowda M, Moghe G, Lin H, Vaillancourt B, Shiu S-H, Jiang N, Robin Buell C.

(2012) 'Comparative transcriptomics of three Poaceae species reveals patterns of gene expression

evolution', The Plant Journal, 71: 492-502, doi: 10.1111/j.1365-313X.2012.05005.x

Danilevicz MF, Gill M, Anderson R, Batley J, et al. (2022) Plant Genotype to Phenotype Prediction

Using Machine Learning. Front. Genet. 13:822173. doi: 10.3389/fgene.2022.822173

Endelman JB (2011). 'Ridge regression and other kernels for genomic selection with R package

rrBLUP', Plant Genome, 4, 250-255

Enyew M, Feyissa T, Carlsson AS, Tesfaye K, Hammenhag C, Seyoum A, Geleta M. (2022)

'Genome-wide analyses using multi-locus models revealed marker-trait associations for major

agronomic traits in Sorghum bicolor', Front. Plant Sci, p.3837, doi: 10.3389/fpls.2022.999692

Gelman A, Rubin DB (1992) 'Inference from iterative simulation using multiple sequences',

Statistical science, 7(4):457-472, doi: 10.1214/ss/1177011136

Ghannam RB, Techtmann S (2021) 'Machine learning applications in microbial ecology, human

microbiome studies, and environmental monitoring', Computational and Structural Biotechnology

Journal, 19: 1092–1107, doi: 10.1016/j.csbj.2021.01.028

Gladman N, Jiao Y, Lee YK, Zhang L, Chopra R, Regulski M, Burow G, Hayes C, et al. (2019)

'Fertility of Pedicellate Spikelets in Sorghum Is Controlled by a Jasmonic Acid Regulatory Module',

International Journal of Molecular Sciences, 20: 4951, doi: 10.3390/ijms20194951

Goodstein DM, Shu S, Howson R, Neupane R, et al. (2012) 'Phytozome: a comparative platform for

green plant genomics', Nucleic Acids Research, 40: D1178–D1186, doi: 10.1093/nar/gkr944

Guo, J., LeBauer, D. (2022). 'genophenoenvo/JAGS-logistic-growth: v0.1.0', doi:

10.5281/zenodo.7114677

Hebbring SJ (2014) 'The challenges, advantages and future of phenome-wide association studies',

Immunology, 141(2):157-65. doi: 10.1111/imm.12195

Hsieh WW (2022) 'Evolution of machine learning in environmental science—A perspective',

Environmental Data Science, 1: e3, doi: 10.1017/eds.2022.2

Jiao Y, Burke J, Chopra R, Burow G, et al. (2016) 'A Sorghum Mutant Resource as an Efficient

Platform for Gene Discovery in Grasses', The Plant Cell, 28: 1551–1562, doi: 10.1105/tpc.16.00373

Jubair S, Domaratzki M (2023) 'Crop genomic selection with deep learning and environmental data:

A survey', Front. Artif. Intell, 5:1040295, doi: 10.3389/frai.2022.1040295

Kearsey MJ, Farquhar AGL (1998) 'QTL analysis in plants; where are we now?' Heredity 80: 137–

142, doi: 10.1046/j.1365-2540.1998.00500.x

Kebrom TH, McKinley B, Mullet JE (2017) 'Dynamics of gene expression during development and

expansion of vegetative stem internodes of bioenergy sorghum', Biotechnol Biofuels 10, 159 (2017).

doi: 10.1186/s13068-017-0848-3

LeBauer D, Maxwell B, Demieville J, Fahlgren N, French A, Garnett R, Hu Z, Huynh K, Kooper R,

Li Z, et al. (2020) 'Data From: TERRA-REF, An open reference data set from high resolution

genomics, phenomics, and imaging sensors', Dryad, Dataset, doi: 10.5061/dryad.4b8gtht99

Lex A, Gehlenborg N, Strobelt H, Vuillemot R, Pfister H (2014) 'UpSet: Visualization of

Intersecting Sets', IEEE Transactions on Visualization and Computer Graphics (InfoVis), 20: 1983–

1992, doi: 10.1109/TVCG.2014.2346248

Lin J, Ngiam KY (2023) 'How data science and AI-based technologies impact genomics', Singapore

Med J.,64(1):59-66, doi: 10.4103/singaporemedj.SMJ-2021-438

Liu X, Huang M, Fan B, Buckler ES, Zhang Z (2016) 'Iterative Usage of Fixed and Random Effect

Models for Powerful and Efficient Genome-Wide Association Studies', PLoS Genet 12(2):

e1005767, doi: 10.1371/journal.pgen.1005767

Lobos GA, Hancock JF (2015) 'Breeding blueberries for a changing global environment: a review',

Frontiers in Plant Science, doi: 10.3389/fpls.2015.00782

Lopes MS, El-Basyoni I, Baenziger PS, Singh S, Royo C, Ozbek K, Aktas H, Ozer E, Ozdemir F, et

al. (2015) 'Exploiting genetic diversity from landraces in wheat breeding for adaptation to climate

change', Journal of Experimental Botany, 66: 3477–3486, doi: 10.1093/jxb/erv122

López-Cortés XA, Matamala F, Maldonado C, Mora-Poblete F, Scapim CA (2020) 'A Deep

Learning Approach to Population Structure Inference in Inbred Lines of Maize', Front. Genet., vol.

11, doi: 10.3389/fgene.2020.543459

Mace ES, Tai S, Gilding EK, Li Y, Prentis PJ, et al. (2013) 'Whole-genome sequencing reveals

untapped genetic potential in Africa's indigenous cereal crop sorghum', Nat Commun, 4: 2320, doi:

10.1038/ncomms3320

Mace E, Innes D, Hunt C, Wang X, Tao Y, Baxter J, et al. (2019) 'The Sorghum QTL Atlas: a

powerful tool for trait dissection, comparative genomics and crop improvement', TAG. Theoretical

and Applied Genetics, 132(3), 751–766, doi: 10.1007/s00122-018-3212-5

McCormick RF, Truong SK, Sreedasyam A, Jenkins J, Shu S, Sims D, et al (2018) 'The Sorghum

bicolor reference genome: improved assembly, gene annotations, a transcriptome atlas, and

signatures of genome organization' The Plant Journal, 93: 338–354, doi: 10.1111/tpj.13781

McLaren W, Gil L, Hunt SE, Riat HS, Ritchie GRS, Thormann A, Flicek P, Cunningham F (2016)

'The Ensembl Variant Effect Predictor', Genome Biology, 17: 122, doi: 10.1186/s13059-016-0974-4

McMaster GS, Wilhelm WW (1997) 'Growing degree-days: one equation, two interpretations',

Agricultural and Forest Meteorology, 87: 291–300, doi: 10.1016/S0168-1923(97)00027-0

Metsalu T, Vilo J (2015) 'ClustVis: a web tool for visualizing clustering of multivariate data using

Principal Component Analysis and heatmap', Nucleic Acids Research, Volume 43, Issue W1, W566–

W570, doi: 10.1093/nar/gkv468

Meuwissen THE, Hayes BJ, Goddard ME (2001) 'Prediction of Total Genetic Value Using Genome-

Wide Dense Marker Maps', Genetics, 157: 1819–1829, doi: 10.1093/genetics/157.4.1819

Mi H, Muruganujan A, Ebert D, Huang X, Thomas PD (2019) 'PANTHER version 14: more

genomes, a new PANTHER GO-slim and improvements in enrichment analysis tools', Nucleic Acids

Research, 47(D1), D419–D426, doi: 10.1093/nar/gky1038

Mitchell AL, Attwood TK, Babbitt PC, Blum M, Bork P, Bridge A, Brown SD, et al. (2019)

'InterPro in 2019: improving coverage, classification and access to protein sequence annotations',

Nucleic Acids Research, 47(D1), D351–D360, doi: 10.1093/nar/gky1100

Moore JH, Asselbergs FW, Williams SM (2010), 'Bioinformatics challenges for genome-wide

association studies', Bioinformatics, 26(4):445-55. doi: 10.1093/bioinformatics/btp713

Naithani S, Cecilia HD, Sahu SK, Jaiswal P (2023) "Exploring Pan-Genomes: An Overview of

Resources and Tools for Unraveling Structure, Function, and Evolution of Crop Genes and

Genomes", Biomolecules 13(9): 1403, doi: 10.3390/biom13091403

Naithani S, Geniza M, Jaiswal P (2017) 'Variant Effect Prediction Analysis Using Resources

Available at Gramene Database', Methods Mol Biol.,1533:279–297, doi: 10.1007/978-1-4939-6658-

5_17

709 Olson A, Klein RR, Dugas DV, Lu Z, Regulski M, Klein PE, Ware D (2014) 'Expanding and vetting

710 sorghum bicolor gene annotations through transcriptome and methylome sequencing', The Plant

711 Genome, 7(2), lantgenome2013.08.0025.

712 Papatheodorou I, Fonseca NA, Keays M, Tang YA, Barrera E, Bazant W, et al. (2018) 'Expression

713 Atlas: gene and protein expression across multiple studies and organisms', Nucleic Acids Research,

714 Volume 46, Issue D1, D246–D251, doi: 10.1093/nar/gkx1158

715 Plummer M (2003) 'JAGS: A Program for Analysis of Bayesian Graphical Models Using Gibbs

716 Sampling', Proceedings of the 3rd International Workshop on Distributed Statistical Computing,

717 Vienna, 1-10

718 Plummer M (2022) 'rjags: Bayesian Graphical Models using MCMC. R package version 4-13', url:

719 https://CRAN.R-project.org/package=rjags

720 Pudil P, Novovičová J, Kittler J (1994) 'Floating search methods in feature selection', Pattern

721 Recognition Letters, Vol. 15, Issue 11, pp. 1119-1125, doi: 10.1016/0167-8655(94)90127-9

722 Quinby J, Karper R (1945) 'The inheritance of three genes that influence time of floral initiation and

723 maturity date in milo', J Am Soc Agron, 37:916-936, doi:

724 10.2134/agronj1945.00021962003700110006x

725 Quinby J, Karper R (1953) 'Inheritance of height in sorghum. Inherit. height. Sorghum', Abstracts of

726 the Annual Meetings of the American Society of Agronomy, Dallas, Texas, pp. 98–99

727 R Core Team (2021) 'R: A language and environment for statistical computing. R Foundation for

728 Statistical Computing', Vienna, Austria. url: https://www.R-project.org/

Rafalski A (2002) 'Applications of single nucleotide polymorphisms in crop genetics', Current

Opinion in Plant Biology, 5:94-100, doi: 10.1016/s1369-5266(02)00240-6

Rhodes DH, Hoffmann L, Rooney WL, Ramu P, Morris GP, Kresovich S (2014) 'Genome-Wide

Association Study of Grain Polyphenol Concentrations in Global Sorghum [Sorghum bicolor (L.)

Moench] Germplasm', J Agric Food Chem, 62(45):10916–10927, doi: 10.1021/jf503651t

Rice B, Lipka AE (2019), Evaluation of RR-BLUP Genomic Selection Models that Incorporate Peak

Genome-Wide Association Study Signals in Maize and Sorghum. The Plant Genome, 12: 180052.

doi: 10.3835/plantgenome2018.07.0052

Shavrukov YN (2016) 'CAPS markers in plant biology', Russ J Genet Appl Res, 6: 279–287, doi:

10.1134/S2079059716030114

Sheshukova EV, Komarova TV, Pozdyshev DV, Ershova NM, Shindyapina AV, et al. (2017) 'The

Intergenic Interplay between Aldose 1-Epimerase-Like Protein and Pectin Methylesterase in Abiotic

and Biotic Stress Control', Front Plant Sci. 8:1646. doi: 10.3389/fpls.2017.01646

Sukumaran S, Xiang W, Bean SR, Pedersen JF, Kresovich S, Tuinstra MR, Tesso TT, Hamblin MT,

Yu J (2012) 'Association Mapping for Grain Quality in a Diverse Sorghum Collection', The Plant

Genome. doi: 10.3835/plantgenome2012.07.0016

Tanksley SD, Young ND, Paterson AH, Bonierbale MW (1989) 'RFLP Mapping in Plant Breeding:

New Tools for an Old Science', Nat Biotechnol, 7: 257–264, doi: 10.1038/nbt0389-257

Tao Y, Zhao X, Wang X, Hathorn A, Hunt C, Cruickshank AW, Oosterom EJ, Godwin ID, Mace ES,

Jordan DR (2020) 'Large-scale GWAS in sorghum reveals common genetic control of grain size

among cereals', Plant Biotechnology Journal, 18: 1093–1105, doi: 10.1111/pbi.13284

Tao Y, Luo H, Xu J, Cruickshank A, Zhao X, Teng F, Hathorn A, Wu X, Liu Y, Shatte T, et al.

(2021) 'Extensive variation within the pan-genome of cultivated and wild sorghum', Nat Plants, 7:

766–773, doi: 10.1038/s41477-021-00925-x

Tello-Ruiz MK, Naithani S, Stein JC, Gupta P, Campbell M, Olson A, Wei S, Preece J, et al. (2018),

'Gramene 2018: unifying comparative genomics and pathway resources for plant research. Nucleic

Acids Research, 46(D1), D1181–D1189, doi: 10.1093/nar/gkx1111

Tello-Ruiz MK, Naithani S, Gupta P, Olson A, Wei S, Preece J, Jiao Y, Wang B, et al. (2021)

'Gramene 2021: harnessing the power of comparative genomics and pathways for plant research',

Nucleic Acids Research, Volume 49, Issue D1, D1452–D1463, doi: 10.1093/nar/gkaa979

The Gene Ontology Consortium (2021) 'The Gene Ontology resource: enriching a GOld mine',

Nucleic Acids Research, 49: D325–D334, doi: 10.1093/nar/gkaa1113

Thessen A (2016) 'Adoption of Machine Learning Techniques in Ecology and Earth Science', One

Ecosystem, 1: e8621, doi: 10.3897/oneeco.1.e8621

Thomas PD, Ebert D, Muruganujan A, et al.(2022) 'PANTHER: Making genome-scale

phylogenetics accessible to all', Protein Science, 31: 8–22, doi: 10.1002/pro.4218

Tong H, Nikoloski Z (2021) 'Machine learning approaches for crop improvement: Leveraging

phenotypic and genotypic big data', Journal of Plant Physiology, vol. 257, doi:

10.1016/j.jplph.2020.153354.

Visscher PM, Brown MA, McCarthy MI, Yang J (2012) 'Five Years of GWAS Discovery', The

American Journal of Human Genetics, 90: 7–24, doi: 10.1016/j.ajhg.2011.11.029

Watanabe K, Guo W, Arai K, Takanashi H, Kajiya-Kanegae H, Kobayashi M, et al. (2017) 'High-

Throughput Phenotyping of Sorghum Plant Height Using an Unmanned Aerial Vehicle and Its

Application to Genomic Prediction Modeling', Front. Plant Sci., 8:421, doi: 10.3389/fpls.2017.00421

Yin L, Zhang H, Tang Z, Xu J, Yin D, Zhang Z, et al. (2021) 'rMVP: A Memory-efficient,

Visualization-enhanced, and Parallel-accelerated Tool for Genome-wide Association Study',

Genomics, Proteomics & Bioinformatics, 19(4):619-628, doi: 10.1016/j.gpb.2020.10.007.

Zhang Q, Zhang Q, Jensen J (2022), Association Studies and Genomic Prediction for Genetic

Improvements in Agriculture, Front. Plant Sci.,vol. 13, doi: 10.3389/fpls.2022.904230

Zheng Z, Hey S, Jubery T, Liu H, et al. (2020) 'Shared Genetic Control of Root System Architecture

between Zea mays and Sorghum bicolor', Plant Physiology, 182: 977–991, doi: 10.1104/pp.19.00752

Zhu C, Gore M, Buckler ES, Yu J (2008) 'Status and Prospects of Association Mapping in Plants',

The Plant Genome, doi: 10.3835/plantgenome2008.02.0089