1

2   Insect Detect: An open-source DIY camera trap for automated insect monitoring

5   Maximilian Sittinger[1*], Johannes Uhler[1], Maximilian Pink[1], Annette Herz[1]

7   [1] Institute for Biological Control, Julius Kühn Institute, Dossenheim, Germany

10   * Corresponding author

11   E-mail: maximilian.sittinger@julius-kuehn.de (MS)

Abstract

Insect monitoring is essential to design effective conservation strategies, which are

indispensable to mitigate worldwide declines and biodiversity loss. For this purpose, traditional

monitoring methods are widely established and can provide data with a high taxonomic

resolution. However, processing of captured insect samples is often time-consuming and

expensive, which limits the number of potential replicates. Automated monitoring methods can

facilitate data collection at a higher spatiotemporal resolution with a comparatively lower effort

and cost. Here, we present the Insect Detect DIY (do-it-yourself) camera trap for non-invasive

automated monitoring of flower-visiting insects, which is based on low-cost off-the-shelf

hardware components combined with open-source software. Custom trained deep learning

models detect and track insects landing on an artificial flower platform in real time on-device

and subsequently classify the cropped detections on a local computer. Field deployment of the

solar-powered camera trap confirmed its resistance to high temperatures and humidity, which

enables autonomous deployment during a whole season. On-device detection and tracking

reliably estimated insect activity/abundance after metadata post-processing. Our insect

classification model achieved a high top-1 accuracy on the test dataset and generalized well

on a real-world dataset with captured insect images. The camera trap design and open-source

software are highly customizable and can be adapted to different use cases. With custom

trained detection and classification models, as well as accessible software programming, many

possible applications surpassing our proposed deployment method can be realized.

Introduction

The worldwide decline in insect biomass, abundance and diversity has been reported in numerous studies in recent years [1–3]. To identify potential drivers of insect decline, quantify their impact and design effective conservation strategies, more long-term monitoring data across ecological gradients is needed, preferably at high spatiotemporal resolutions [4, 5]. An obstacle to the implementation of large-scale monitoring schemes are often financial restrictions. The estimated costs for a national pollinator monitoring scheme range from ~525 € per site and year for volunteer floral observations with crowd-sourced identification to group level, up to ~8130 € for a professional monitoring with pan traps, transect walks, floral observations and expert identification to species level [6]. Due to these high costs, as well as the time-consuming processing and identification of insect specimens, traditional monitoring methods are usually deployed with a low number of spatial and/or temporal replicates. This often generates snapshot data with restricted potential for analysis and interpretation.

Non-invasive automated monitoring methods, which can acquire data at a high spatiotemporal resolution, can complement traditional monitoring methods that often generate data with a comparatively higher taxonomic resolution. When integrated into frameworks based on artificial intelligence (AI) approaches for rapid automated data processing and information extraction, large amounts of high-quality data can be collected with comparatively lower time and effort required [7]. Standardizing these automated monitoring methods and providing easy accessibility and reproducibility could furthermore decentralize monitoring efforts and strengthen the integration of independent biodiversity observations, also by non-professionals (Citizen Science) [8].

A range of different sensors can be used for automated insect monitoring [9]. These include acoustic [10] and opto-electronic sensors [11–13], as well as cameras [14–20]. Several low-cost DIY camera trap systems use scheduled video or image recordings, which are analyzed in subsequent processing steps [14, 15]. Other systems utilize motion detection software as

trigger for the video or image capture [16, 18]. Similar to traditional camera traps used for the

monitoring of mammals, these systems often generate large amounts of video/image data,

which is most efficiently processed and analyzed by making use of machine learning (ML) and

especially deep learning (DL) algorithms [21–23] automatically extract information such as

species identity, abundance or behavior [24]. While automated insect classification with

specifically trained DL models does not yet reach the accuracy of taxonomic experts, results

are still of sufficient quality to perform routine identification tasks in a fraction of time with the

potential to significantly reduce human workload [25].

Small DL models with few parameters and relatively low computational costs can be run on

suitable devices directly in the field ("AI at the edge"), to enable real-time detection and/or

classification of objects the model was trained on. An existing camera trap for automated insect

monitoring combines scheduled time-lapse image recordings at 0.33 fps (frames per second)

with subsequent on-device insect detection and classification running in parallel, while tracking

is implemented during post-processing on a local computer [17]. As an alternative approach,

frames produced by the camera can also be used as direct input for an insect detection model

running with a high frame rate in real time on the camera trap hardware. In this way, the

appearance and detection of an insect can automatically trigger the image capture. This is

usually more robust to false triggers (e.g. moving background) compared to motion detection

and can drastically reduce the amount of data that has to be stored.

Following this approach, we present the design and proof of concept of a novel DIY camera

trap system for automated visual monitoring of flower-visiting insects. The system is based on

low-cost off-the-shelf hardware components and completely open-source software, capable of

on-device AI inference at the edge with custom trained models. Our goal was to develop a

camera trap that could be easily utilized in monitoring projects involving citizen scientists to

achieve a broader application potential. As such, it should be inexpensive, easy to assemble

and set up, and provide reliable results without the requirement of expert taxonomic knowledge

during data collection. Detailed step-by-step instructions to enable a simple reproduction and

optional customization of the camera trap can be found at the corresponding documentation

website [26].

The camera trap is resistant to high temperatures and humidity, as well as fully solar-powered,

which makes it possible to deploy autonomously during a whole season. Insects landing on a

platform with colored flower shapes are detected and tracked in real time at ~12.5 fps, while

an image of each detected insect is cropped from synchronized high-resolution frames and

saved together with relevant metadata to a microSD card every second. The insect images

can then be automatically classified with a custom trained model in a subsequent step on a

local computer. The on-device tracking capabilities were tested with a fast-flying hoverfly

species (*Episyrphus balteatus*) in a lab experiment, and different metadata post-processing

settings for a reliable activity/abundance estimation were compared. Five camera traps were

continuously deployed during four months in 2023 with a total recording time of 1919 hours.

Captured images and metadata were classified, post-processed and analyzed. The

generalization capability of the classification model was validated on a real-world dataset of

images captured during field deployment.

Materials and methods

Hardware

The camera trap system is based on three main hardware components: (1) OpenCV AI Kit

(OAK-1) (Luxonis, Littleton, USA), including a 12MP image sensor and a specific chip for on-

device AI inference with custom models; (2) Raspberry Pi Zero 2 W (RPi) (Raspberry Pi

Foundation, Cambridge, UK), a single-board computer that is used as host for the OAK-1 and

stores the captured data on a microSD card; (3) PiJuice Zero pHAT (Pi Supply, East Sussex,

UK), an expansion board for the RPi with integrated RTC (real-time clock) for power

management and scheduling of recording times.

Two rechargeable batteries (~91Wh) connected to a 9W 6V solar panel (Voltaic Systems, New

York, USA) are used as power supply for the system. All electronic components are installed

in a weatherproof enclosure, which can be mounted to a standard wooden or steel post (Fig

1B). Below the camera trap enclosure, a sheet (e.g. acrylic glass or lightweight foam board)

with colored flower shapes printed on top is attached to the same post, which acts as visual

attractant and landing platform for flower-visiting insects (Fig 1A). This standardizable platform

provides a homogeneous plane background and leads to a uniform posture of insects sitting

on it, which increases detection, tracking and classification accuracy with less data required

for model training. The cost for the full camera trap setup including all components is ~700 €,

a minimal setup can be built for ~530 €. A complete list of all required components and step-

by-step assembly instructions can be found at the documentation website [26].

Fig 1. Camera trap design. (A) Field deployment of the camera trap and flower platform on

wooden post. (B) Weatherproof camera trap enclosure with integrated hardware and

connected solar panel.

Software

Insect detection

All of the camera trap software and associated insect detection models are available at GitHub

[27]. The software includes Python scripts for livestreaming of camera frames together with

the detection model and/or object tracker output, and for capturing high-resolution frames

(1080p/4K/12MP) or videos (1080p/4K). All scripts use auto focus, auto exposure and auto

white balance for the OAK-1 camera by default. Together with the software, custom trained

YOLOv5n [28], YOLOv6n [29], YOLOv7-tiny [30] and YOLOv8n [31] object detection models

are provided, which can be run on the OAK-1 chip to enable real-time insect detection and

tracking. The respective model weights, pre-trained on the MS COCO dataset [32], were fine-

tuned on an image dataset, collected with a camera trap prototype in 2022 [33]. The dataset

is composed of 1,335 images, of which 1,225 contain at least one annotation (110 background

images). A total of 2,132 objects were annotated, including 664 wasps, 454 flies, 297 honey

bees, 297 other arthropods, 233 shadows of insects and 187 hoverflies. For detection model

training, all originally annotated classes were merged into one generic class ("insect"), the

original 4K resolution (3840x2160 pixel) of the images was downscaled and stretched to

320x320 pixel, and the dataset was split into train (1,069 images), validation (133 images) and

test (133 images) subsets with a ratio of 0.8/0.1/0.1.

The provided YOLO models are general insect detectors and can also detect insects not

included in the dataset and/or on other homogeneous backgrounds. Due to the downscaled

input image size of 320x320 pixel, the models can achieve a high on-device inference speed,

while still keeping good precision and recall values (Table 1). The YOLOv5n model achieved

the highest mAP (mean average precision) and recall on the dataset validation split and is

used as default model by the camera trap software, with an IoU (intersection over union) and

confidence threshold of 0.5 respectively. Google Colab notebooks are provided at GitHub [34]

to reproduce the model training and validation, or train detection models on custom datasets.

Table 1. Metrics of the YOLO insect detection models.

Model	Image size [pixel]	mAP^{val} (@0.5 IoU)	Precision^{val}	Recall^{val}	Speed OAK-1 [fps]	Parameters (million)
YOLOv5n	320	0.969	0.955	0.961	49	1.76
YOLOv6n	320	0.951	0.969	0.898	60	4.63
YOLOv7-tiny	320	0.957	0.947	0.942	52	6.01
YOLOv8n	320	0.944	0.922	0.899	39	3.01

167 All models were trained on a custom dataset with 1,335 images (1,069 in train split) to 300

168 epochs with batch size 32 and default hyperparameters. Metrics (mAP, Precision, Recall) are

169 shown on the dataset validation split (133 images) for the original PyTorch (.pt) models before

170　conversion to .blob format. Speed (fps) is shown for the converted .blob models running on the

171　OAK-1 connected to the RPi Zero 2 W.

Processing pipeline

The main Python script for continuous automated insect monitoring uses a specific processing

pipeline to run insect detection on downscaled LQ (low-quality) frames (320x320 pixel). This

increases inference speed of the model and thereby accuracy of the object tracker that uses

the coordinates from the model output as input for tracking insects. The object tracker is based

on the Kalman Filter and Hungarian algorithm, which can keep track of a moving object by

comparing the bounding box coordinates of the current frame with the object's trajectory on

previous frames. By assigning a unique ID to each insect landing on or flying above the flower

platform, multiple counting of the same individual can be avoided as long as it is present in the

frame. As the lack of visual features on LQ frames would significantly decrease classification

accuracy, all tracked detections are synchronized with HQ (high-quality) frames (1920x1080

or 3840x2160 pixel) in a script node on-device in real time (Fig 2). The detected and tracked

insects are then cropped from the synchronized HQ frames by utilizing the bounding box

coordinates (Fig 3). The cropped detections are saved as individual .jpg images to the microSD

card of the RPi together with relevant metadata including timestamp, confidence score and

tracking ID with an image capture frequency of ~1 s. The image capture frequency can be

adjusted optionally, without decreasing the overall pipeline and inference speed. Due to the

synchronization of the tracker/model output with the HQ frames, the whole pipeline runs at

~12.5 fps (1080p HQ frame resolution), which is sufficient to reliably track most insects (~3.4

fps for 4K HQ frame resolution).

Fig 2. Diagram of the processing pipeline. HQ frames (1080p or 4K) are downscaled to LQ

frames (320x320 pixel), while both HQ and LQ frames run in two parallel streams. The LQ

frames are used as input for the insect detection model. The object tracker uses the

coordinates from the model output to track detected insects and assign unique tracking IDs.

Detected insects are cropped from synchronized HQ frames in real time and saved to the

microSD card of the RPi together with relevant metadata.

Fig 3. LQ frame and synced HQ frame with cropped detection. Downscaled LQ frames

(320x320 pixel) are used as model input. Detected insects are cropped from synchronized HQ

frames (1080p or 4K) on-device (OAK-1).

The mean peak power consumption of the camera trap system is ~4.4 W, while constantly

detecting and tracking five insects, and saving the cropped detections to the RPi microSD card

every second (Fig 4). With a combined battery capacity of ~91Wh, recordings can be run for

~20 hours, even if no sunlight is available for the solar panel to recharge the batteries. The

PiJuice Zero pHAT is used for efficient power management. If the PiJuice battery charge level

drops below a specified threshold, the RPi will immediately shut down before starting a

recording. The respective duration of each recording interval is conditional on the current

charge level, which can prevent gaps in monitoring data during periods with less sunlight. The

scheduled recording times, charge level thresholds and recording durations can be easily

adjusted for each use case.

Fig 4. Power consumption of the camera trap system. Power consumption was measured

during a 5 min recording interval while constantly detecting and tracking five insects, and

saving the cropped detections together with metadata to the RPi microSD card every second.

The insect images are classified in a subsequent step, followed by post-processing of the

metadata. This two-stage approach makes it possible to achieve high detection and tracking

accuracy in real time on-device (OAK-1), combined with a high classification accuracy of the

insect images cropped from HQ frames on a local computer.

Insect classification

The data output from the Python script for automated insect monitoring includes insect images

together with associated .csv files containing metadata for further analysis. Classification of

the insect images is done in a subsequent step on a local computer, after the camera trap data

has been collected from the microSD card. A modified YOLOv5 [28] classification script is used

to classify all insect images captured with the camera trap and append the classification results

(top-3 classes and probabilities) as new columns to the merged metadata .csv files. The script

is available at GitHub [35] together with other modified YOLOv5 scripts, including classification

model training and validation.

For training of the insect classification model we used an EfficientNet-B0 model, pre-trained

on the ImageNet dataset [36]. The pre-trained weights were fine-tuned on a custom image

dataset, mainly containing cropped detections captured by six camera traps [37]. The dataset

is composed of 21,000 images, of which 18,597 images contain various arthropods (mainly

insects). We sorted all images to 27 different classes, including four classes without insects

("none_*"). These additional classes are used to filter out images of background, shadows, dirt

(e.g. leaves and bird droppings) and birds that can be captured by the camera trap. Most of

the images were collected with six different camera traps deployed in the field in 2022 and

2023. Additional images of some classes with insufficient occurrences in the field data

(including "ant", "bee_bombus", "beetle_cocci", "bug", "bug_grapho", "hfly_eristal",

"hfly_myathr", "hfly_syrphus") were captured with a lab setup of the camera trap hardware to

increase accuracy and generalization capability of the model. Detailed descriptions of all

classes can be found in S1 Table and example images are shown in Fig 5.

Fig 5. Example images of the 27 classes from the dataset for classification model

training. All images were captured automatically by the camera trap and are shown unedited,

but resized to the same dimension. Original pixel values are plotted on the x- and y-axis for

each image. Four example images are shown for the class "other", to account for its high

intraclass heterogeneity.

For model training, the dataset was split into train (14,686 images), validation (4,189 images)

and test (2,125 images) subsets with a ratio of 0.7/0.2/0.1. We compared the metrics of three

different model architectures supported by YOLOv5 classification model training and different

hyperparameter settings to find the combination with the highest accuracy on our dataset (S2

Table). The EfficientNet-B0 model [38] trained to 20 epochs with batch size 64 and images

scaled to 128x128 pixel achieved the best balance between accuracy and inference speed,

while generalizing better to a real-world dataset compared to the same model trained to only

ten epochs. With this model, high top-1 accuracies on the test dataset for all insect classes

could be achieved (Fig 6, Table 2). Only three classes ("beetle", "bug", "other") did not reach a

top-1 accuracy > 0.9, probably because of a high visual intraclass heterogeneity and not

enough images in the dataset.

Fig 6. Normalized confusion matrix for the EfficientNet-B0 insect classification model,

validated on the dataset test split. The model was trained on a custom dataset with 21,000

images (14,686 in train split), scaled to 128x128 pixel, to 20 epochs with batch size 64 and

default hyperparameters. Metrics are shown on the dataset test split (2,125 images) for the

converted model in ONNX format.

Table 2. Metrics of the EfficientNet-B0 insect classification model, validated on the dataset test split.

Class	Images^{test}	Top-1 accuracy^{test}	Precision^{test}	Recall^{test}	F1 score^{test}
all	2125	0.972	0.971	0.967	0.969
ant	111	1.0	0.991	1.0	0.996
bee	107	0.963	0.972	0.963	0.967
bee_apis	31	1.0	0.969	1.0	0.984
bee_bombus	127	1.0	0.992	1.0	0.996
beetle	52	0.885	0.92	0.885	0.902
beetle_cocci	78	0.987	1.0	0.987	0.994
beetle_oedem	21	0.905	0.905	0.905	0.905
bug	39	0.846	1.0	0.846	0.917
bug_grapho	19	1.0	1.0	1.0	1.0
fly	173	0.971	0.944	0.971	0.957
fly_empi	19	1.0	1.0	1.0	1.0
fly_sarco	33	0.909	0.938	0.909	0.923
fly_small	167	0.958	0.952	0.958	0.955
hfly_episyr	253	0.996	0.996	0.996	0.996
hfly_eristal	197	0.99	0.995	0.99	0.992
hfly_eupeo	137	0.985	0.993	0.985	0.989
hfly_myathr	60	1.0	1.0	1.0	1.0
hfly_sphaero	39	0.974	1.0	0.974	0.987
hfly_syrphus	50	0.98	1.0	0.98	0.99
lepi	24	1.0	0.96	1.0	0.98
none_bg	86	0.988	0.966	0.988	0.977
none_bird	8	1.0	1.0	1.0	1.0
none_dirt	85	0.976	0.902	0.976	0.938
none_shadow	66	0.924	0.953	0.924	0.938
other	79	0.861	0.883	0.861	0.872
scorpionfly	12	1.0	1.0	1.0	1.0
wasp	52	1.0	1.0	1.0	1.0

The model was trained on a custom dataset with 21,000 images (14,686 in train split), scaled

to 128x128 pixel, to 20 epochs with batch size 64 and default hyperparameters. Metrics are

shown on the dataset test split (2,125 images) for the converted model in ONNX format.

The insect classification model was exported to ONNX format, which enables faster inference

speed on standard CPUs (~10-20 ms per image), and is available at GitHub [34]. To reproduce

the model training and validation, or train classification models on custom datasets, Google

Colab notebooks are provided in the same repository [34].

Metadata post-processing

By running the classification script, all available metadata .csv files from the camera trap output

are merged and the top-3 classification results for each insect image are appended to new

columns. In most cases, the newly generated .csv file still contains multiple rows for each insect

with a unique tracking ID, with the number of rows depending on the image capture frequency

(default: ~1 s) and on the duration the insect was present in the frame. With multiple captured

images for each individual insect, classification of the images can result in different top-1

classes for the same tracking ID. The Python script for post-processing and analysis [34] is

used to generate the final metadata .csv file that can then be used for further data analysis.

The weighted mean classification probability is calculated to determine the top-1 class with the

overall highest probability for each tracking ID, by calculating the mean probability of each top-

1 class per tracking ID and multiplying it with the proportion of images classified to the

respective top-1 class to the total number of images per tracking ID:

$$weighted\ mean\ probability_{top1\ class} = mean(probability_{top1\ class}) \times \frac{\#\ images_{top1\ class}}{\#\ images_{tracking\ ID}}$$

Only the top-1 class with the highest weighted mean classification probability per tracking ID

is kept in the final metadata file, in which each row corresponds to an individual tracked insect.

To make the activity/abundance estimations more reliable, all individual insects (= unique

tracking IDs) with less than three or more than 1,800 images are excluded by default. These

values can be adjusted optionally, e.g. if a different image capture frequency is used and/or to

make the final data more robust to inaccurate tracking of fast-moving insects. This can result

in "jumping" tracking IDs, as the limited inference speed of the detection model is providing the

object tracker with bounding box coordinates in a frequency that is too low to keep track of the

fast-moving insect (S1B Fig). In this case, the track of the insect is lost and a new tracking ID

is assigned to the same individual. With the default image capture frequency of ~1 s, the insect

had to be tracked for at least slightly over two seconds to be kept in the final dataset. The

default upper limit of 1,800 images per tracking ID removes all IDs that were tracked for more

than ~30-40 min at the default image capture frequency. This maximum duration depends on

the number of simultaneous detections, which can slightly decrease the capture frequency

over time. From our experience, objects tracked for > 30 min are most often non-insect

detections, e.g. leaves fallen on the flower platform.

For estimation of the respective insect size, the absolute bounding box sizes in mm are

calculated by supplying the true frame width and height (e.g. flower platform dimensions).

Insect tracking evaluation

The accuracy of the object tracker was tested in a lab experiment. The camera trap hardware

was installed 40 cm above the big flower platform (50x28 cm), so that the platform was filling

out the whole camera frame, and both were placed in the middle of a 200x180x155 cm insect

cage. The script for continuous automated insect monitoring was run with 4K HQ frame

synchronization (~3.4 fps) for 15 min for each recording interval. 15 *Episyrphus balteatus*

hoverflies (reared under lab conditions) were released inside the cage and recorded for 15 min

for each of the ten replications. The flower platform was simultaneously filmed with a

smartphone camera during each recording interval. The videos were played back afterwards

with reduced speed and all frame/platform visits of the hoverflies were counted manually. The

captured metadata was post-processed with ten different settings for the minimum number of

images required per tracking ID to be kept in the final data output ("-min_tracks 1-10"). The

true platform visits from the video count were then compared with the number of unique

tracking IDs for each setting.

Insect classification validation

While the metrics of the custom trained EfficientNet-B0 classification model show a high accuracy on the dataset test split, the underlying dataset was curated and insect images were selected in such a way to include images where the respective class was clearly identifiable and also some images with more difficult cases, e.g. with only a part of the insect visible. For a more realistic measure of the classification accuracy and an estimation of the generalization capability of the model, we compiled a dataset with images captured during field deployment of five camera traps between August 11 and September 18, 2023. No images from this recording period were included in the dataset for classification model training. All captured 93,215 images from camera trap 1 were classified and subsequently manually verified, to ensure that all images were sorted to the correct class. As some classes were only present with a small number of images, more images from camera traps 2-5 were added to achieve a better class balance, resulting in a total dataset size of 97,671 images. All images from each class were added to the dataset, including false negative classifications (image of target class classified to wrong class) and false positive classifications (image of other class classified wrongly to target class). No images were removed to reflect a real-world use case, which means that many nearly identical images are present in the dataset (e.g. insects moving only slightly or none-insect detections such as leaves) and the overall class balance is biased towards classes with more captured images. For two classes ("bug_grapho", "fly_empi"), no images were captured during the selected recording period. To run the model validation, a dummy image for each of these classes was added to the dataset and thus results for both classes must be ignored.

Field deployment

Starting in mid-May 2023, five camera traps were deployed at five different sites in southwestern Germany. All sites were located within a range of 50 km and separated by at least 5 km, with an elevation ranging from 90-170 m.a.s.l. As study sites, extensively managed orchard meadows were chosen, where an insect monitoring with focus on hoverflies using traditional methods (Malaise traps, yellow pan traps) was also being conducted. While the area, age and tree density differed between the sites, all meadows were cut at least once per year.

All camera traps were positioned in full sunlight with the solar panel and flower platform facing to the south. Scheduled recording intervals with a respective duration of 40 min (if the current battery charge level was > 70%) were run four to seven times per day, resulting in a recording duration of 160 to 280 min for each camera trap per day. Recordings started at 6, 7, 8, 10, 11, 14, 16, 18 and 19 o'clock, with the start time differing per month and in some cases also per camera trap. Due to differences in the date of first deployment (camera trap 3, camera trap 5) and initial hardware problems (camera trap 1), the total recording times differed between all camera traps (Table 3). The respective duration of each recording interval is conditional on the current battery charge level, which additionally influenced the total recording time per camera trap, depending on the amount of sunlight available to charge the batteries. During field deployment, the artificial flower platform was changed at least once for each camera trap (except camera trap 5) to test different sizes, materials and designs (flower shapes and colors). Therefore, the results presented below should be interpreted with a certain degree of caution in relation to the insect monitoring data, and should be considered primarily as proof of concept for the camera trap system.

Table 3. Recording times of the five deployed camera traps and site details of the orchard meadows.

Camera trap ID	Total recording time [h]	Site ID	Coordinates	Area [ha]
Camtrap 1	362	Gross-Rohrheim	49.718741, 8.509769	1.4
Camtrap 2	456	Boehl-Iggelheim	49.379616, 8.330233	0.7
Camtrap 3	349	Malsch	49.236468, 8.670059	0.9
Camtrap 4	421	Bensheim	49.703971, 8.590160	1.7
Camtrap 5	330	Dossenheim	49.446449, 8.649950	5.4

Data visualization

383 Data analysis of the automatically post-processed metadata captured by five camera traps

384 deployed in the field and creation of plots presented in the following was conducted with R

385 version 4.3.1 [39] and the R packages tidyverse version 2.0.0 [40], patchwork version 1.1.3

386 [41] and viridis version 0.6.4 [42]. All required R scripts and associated data to reproduce the

387 plots are available at Zenodo [43].

Results

Insect tracking evaluation

With 10 replications of 15 min recording intervals, the object tracker accuracy was tested

together with cropping detected insects from synchronized 4K HQ frames (~3.4 fps) every

second. High activity with many frame/platform visits of the hoverflies during the 15 min

recording interval resulted in a decreased tracking accuracy, mainly due to hoverflies flying

fast and erratic and/or coming close to each other (S1 Fig). This behavior can lead to "jumping"

tracking IDs and consequently multiple counting of the same individual, which is reflected in

the number of unique tracking IDs if none are removed during post-processing with the setting

"-min_tracks 1" (Fig 7). By excluding tracking IDs with less than a specified number of images during post-processing, the activity/abundance estimation can get more robust. Even with a decreased pipeline speed of ~3.4 fps during synchronization with 4K HQ frames, an exclusion of tracking IDs with less than six images ("-min_tracks 6") led to a relatively precise estimation of hoverfly activity/abundance in the data from the lab experiment (Fig 7).

Fig 7. Evaluation of the insect tracking accuracy in a lab experiment. Data from 10 replications of 15 min recording intervals with 15 *E. balteatus* hoverflies placed in a cage with the camera trap and flower platform is shown. Linear regression lines illustrate the effect of 10 different tracking IDs filter methods for post-processing of the captured metadata of each recording interval. With "-min_tracks 1" no tracking IDs are excluded. Dashed line indicates optimal result. True frame visits were manually counted from video recordings of the flower platform.

Insect classification validation

To estimate the generalization capability of the custom trained EfficientNet-B0 insect classification model, a dataset with images captured between August 11 and September 18, 2023 was compiled. For some classes ("beetle", "bug", "other", "wasp") classification accuracy was very low (Fig 8, S3 Table). For the classes "beetle_cocci" and "beetle_oedem", some of the images were classified as "beetle", which is the correct taxonomic order and could therefore be analyzed and interpreted without drawing overly wrong conclusions. In cases of uncertainty and wrong model predictions, images seem to have a tendency to be classified to the class "other", which has a high intraclass heterogeneity in the training dataset. For our focus group of hoverflies ("hfly_*"), a high classification accuracy was achieved for all classes of species (-groups). Some images of *Syrphus* sp. hoverflies ("hfly_syrphus") were wrongly predicted as *Eupeodes* sp. ("hfly_eupeo"), which could be explained by the high visual similarity of both

genera. Overall, classes with a high visual intraclass heterogeneity, such as "beetle", "bug"

and "other" could not be classified with a sufficient accuracy in the real-world dataset. Classes

with less intraclass heterogeneity achieved a high classification accuracy in most cases.

Fig 8. Normalized confusion matrix for the EfficientNet-B0 insect classification model,

validated on a real-world image dataset. The model was trained on a custom dataset with

21,000 images (14,686 in train split), scaled to 128x128 pixel, to 20 epochs with batch size 64

and default hyperparameters. Metrics are shown on a real-world dataset (97,671 images) for

the converted model in ONNX format. All images were classified and subsequently verified

and sorted to the correct class in the case of a wrong classification by the model. A dummy

image was added for each of the classes "bug_grapho" and "fly_empi", as no images of both

were captured. Results for both classes must be ignored.

Field deployment

Five camera traps were deployed at five different sites in southwestern Germany, starting in

mid-May 2023. The capability of the camera traps to withstand high temperatures and humidity

was tested, as well as the performance of the solar panel and two connected batteries used

as power supply for the system. The captured insect images were classified with the custom

trained EfficientNet-B0 classification model, and metadata was post-processed with the default

settings ("-min_tracks 3" and "-max_tracks 1800") to exclude all tracking IDs with less than

three or more than 1,800 images.

Weather resistance

The maximum air temperature measured at the nearest weather station during deployment of the camera traps was 37.5°C in July. The maximum temperature measured during the camera trap recordings was 81°C for the OAK-1 CPU and 66°C for the RPi CPU (Fig 9). Both measurements still lie in the safe operating temperature range for these devices and a normal functionality of the camera trap can be expected if the air temperature does not exceed ~38°C.

Fig 9. Maximum air and OAK-1/RPi CPU temperatures per day. Weather data was taken from the nearest weather station (source: Deutscher Wetterdienst).

For camera trap 5, a humidity/temperature USB logger was placed inside the enclosure from July to mid-August. Even during days with a high mean air humidity, measured at the nearest weather station, the mean humidity inside the enclosure only increased slightly to a maximum of ~15% (Fig 10). For all five camera traps, no buildup of moisture during the recording time until mid-September could be noticed, even without exchanging the originally installed 50g Silica gel pack.

Fig 10. Mean humidity per day measured inside the enclosure and at nearest weather station. Weather data was taken from the nearest weather station (source: Deutscher Wetterdienst).

Two rechargeable batteries connected to a 9W solar panel are used as power supply for the camera trap system. During sunny days with a sunshine duration of ~8-12 h per day, the charge level of the PiJuice battery stayed relatively constant at > 80% for all five camera traps (Fig 11). A drop in the battery charge level can be noticed during end of July and end of August,

when the sunshine duration per day decreased to ~0-4 h for several days. Since the duration

of each recording interval is conditional on the current charge level, decreased recording

durations preserved battery charge even when less sunlight was available to recharge the

batteries. A fast recovery to charge levels > 80% restored the normal recording durations within

a few days, after the sunshine duration increased again.

Fig 11. Mean PiJuice battery charge level and sum of the sunshine duration per day. The

PiJuice battery was charged by a second battery, connected to a 9W solar panel. Weather

data was taken from the nearest weather station (source: Deutscher Wetterdienst).

Insect monitoring data

During a recording time of ~1919 h of all five camera traps combined, a total of ~2.34 million

images with 49,583 unique tracking IDs were captured between mid-May and mid-September

2023. After post-processing of the metadata with default settings, 23,900 tracking IDs with less

than three images (= tracked less than ~2 s) and 85 tracking IDs with more than 1,800 images

(= tracked longer than ~30-40 min) were removed from the final dataset, resulting in ~2.03

million images with 25,598 unique tracking IDs.

Out of the 25,598 unique tracking IDs, 8,677 were classified to one of the non-insect classes

("none_bg", "none_bird", "none_dirt", "none_shadow") (S2 Fig). Flies ("fly_small", "fly") were

the most frequently captured insects, followed by bees ("bee") and other arthropods ("other")

(Fig 12F). For hoverflies, 1,090 unique tracking IDs of *E. balteatus* ("hfly_episyr"), followed by

672 unique tracking IDs of *Eupeodes corollae* or *Scaeva pyrastri* ("hfly_eupeo") and 220

unique tracking IDs of the other four hoverfly classes were recorded by all camera traps.

Differences in the composition of the captured insects between the camera traps can be

observed, which could have been influenced by site conditions and/or by different flower

platforms used for testing purposes during field deployment.

Fig 12. Total number of unique tracking IDs for each predicted insect class. (A – E) Data from camera traps 1-5. (F) Merged data from all camera traps. Data of non-insect classes not shown. All tracking IDs with less than three or more than 1,800 images were removed.

When an insect leaves the frame/platform and re-enters it again, a new tracking ID is assigned with the risk to count the same individual multiple times, which could influence the activity/abundance estimations. To assess this risk, the minimum time difference to the previous five tracking IDs that were classified as the same insect class was calculated for each unique tracking ID. A total of 3,150 tracking IDs showed a time difference of less than five seconds to the previous tracking ID that was classified as the same class (Fig 13). This equals ~18.6% of the 16,921 total tracking IDs classified as insects in our present dataset, of which some might have been the same individuals re-entering the frame.

Fig 13. Time difference to previous tracking ID classified as the same class. Minimum time difference < 30 s of the previous five tracking IDs classified as the same class is shown. Data of non-insect classes not shown. All tracking IDs with less than three or more than 1,800 images were removed.

Data from the images classified as one of the hoverfly classes is presented in more detail in the following plots, as an example for a functional group of special interest. Overall, the number of captured hoverfly tracking IDs varied throughout May and June, but also the total recording time per day was comparatively low during this period (Fig 14F). A peak in late June/early July is followed by steady numbers throughout July. Less hoverflies were recorded in August, though the recording time per day was also decreased during the first half of the month, due to lower battery charge levels caused by less available sunshine. Significantly lower numbers

of hoverflies were captured by camera trap 5 compared to the other four camera traps (Fig 14E). While the maximum recording time per day was normally 280 min for each camera trap, an overlap of changed recording start times resulted in 320 min total recording time per day for camera trap 3 in June 28 (Fig 14C).

Fig 14. Total number of unique tracking IDs classified as hoverfly and recording time per day. (A – E) Data from camera traps 1-5. (F) Merged data from all camera traps. Grey lines/areas indicate the recording time per day. Dashed lines indicate a change of the flower platform. All tracking IDs with less than three or more than 1,800 images were removed.

With differences in the total recording time per day between days and camera traps, the activity, calculated as the number of unique tracking IDs per hour of active recording, can be a more adequate estimation of hoverfly activity/abundance. While the range of hoverfly activity was mostly similar between all camera traps, a peak in activity at the first day of deployment (May 16) can be observed for camera trap 1 (Fig 15A). In total, 41 unique tracking IDs of hoverflies were captured during a single 40 min recording interval at that day, which extrapolates to 62 tracking IDs per hour. This initially high activity could have been caused by a smaller group of hoverflies visiting the platform multiple times due to a higher attractivity induced by novelty. The highest hoverfly activity for a prolonged period was recorded by camera trap 3 in early July (Fig 15C). A decrease in hoverfly activity for all camera traps in late July/early August could have been caused by more rainfall during this time (Fig 15F).

Fig 15. Number of unique tracking IDs classified as hoverfly per hour and precipitation sum per day. (A – E) Data from camera traps 1-5. (F) Merged data from all camera traps. Shaded areas indicate days without recordings. All tracking IDs with less than three or more

than 1,800 images were removed. Weather data was taken from the nearest weather station

(source: Deutscher Wetterdienst).

The start of the recording intervals was scheduled at 6, 7, 8, 10, 11, 14, 16, 18 and 19 o'clock,

with the respective start times differing per month and in some cases also per camera trap.

This resulted in differences in the total recording time per hour, with most recordings available

from 8, 10, 16 and 18 o'clock, and only few recordings available from 14 o'clock (S3 Fig).

Overall, the highest hoverfly activity was measured at 8 and 10 o'clock, with approximately two

unique hoverfly tracking IDs captured per hour on average (Fig 16F). About one hoverfly was

captured per hour at 7 and 16 o'clock. A similarity in high activity during hours before noon

compared to the other times of day can be seen also for bees (S3 Fig) and flies (S4 Fig).

Fig 16. Estimated hoverfly activity (number of unique tracking IDs per hour) per time of

day. (A – E) Data from camera traps 1-5. (F) Merged data from all camera traps. Shaded areas

indicate hours without recordings. All tracking IDs with less than three or more than 1,800

images were removed.

Discussion

The Insect Detect DIY camera trap system and associated software can be considered as an

open-source development platform for anyone interested in automated insect monitoring. Due

to its resistance to high temperatures and humidity, as well as a low power consumption of

~4.4 W combined with energy supplied by a solar panel, the camera trap system can be

autonomously deployed during a whole season. We specifically target non-professionals as

potential user group for deployment of the camera trap, such as citizen scientists involved in

large-scale monitoring projects. Detailed step-by-step instructions for hardware assembly and

software setup are provided at the corresponding documentation website [26] and allow

everyone a full reproduction of the system with additional tips on software programming. The

insect detection and classification models can be easily retrained on custom datasets without

special hardware requirements by using the provided Google Colab notebooks [34].

The artificial flower platform, which is used as attractant for flower-visiting insects and

background for the captured images, increases detection and tracking accuracy and can be

standardized similar to yellow pan traps to compare insect populations at different sites.

Ongoing efforts are made to test and compare alternative materials, shapes and colors. An

updated platform design could further increase the visual attraction for a wider range of insect

taxa or for specific target groups. By omitting the flower platform, the camera trap system can

be easily adapted to different use cases, e.g. to monitor insects visiting real flowers, ground-

dwelling arthropods (platform on ground) or nocturnal insects (vertical setup with light source).

Processing pipeline

We implemented a two-stage approach that combines reliable on-device insect detection and

tracking in real time at ~12.5 fps on low-resolution frames in the first stage, with accurate

classification of the insect images cropped from synchronized high-resolution frames in the

second stage. By using our proposed processing pipeline running on the camera trap

hardware, it is sufficient to only store the cropped insect detections (~5-10 KB per image at

1080p resolution). This can save a significant amount of space on the microSD card, compared

to storing the full frame (~0.5-1 MB per image at 1080p resolution). The separation of both

steps also simplifies dataset management and encourages retraining of the classification

model, as sorting the cropped insect images into class folders is much easier and faster

compared to bounding box annotations required to train a new detection model. Cases with a

low detection confidence and images that were classified to the wrong class or with a low

probability (e.g. new insect species) should be identified, relabeled/resorted and added to a

new dataset version to retrain the detection and/or classification model incrementally. Through

this iterative active learning loop, the accuracy of the deployed models can be increased over

time and will adapt to specific use cases and/or environments [44].

603 Insect detection and tracking

While the provided insect detection models will generalize well for different homogeneous

backgrounds and can also detect insect taxa not included in the training dataset, accuracy will

drop significantly when using them with complex and dynamic backgrounds, such as natural

vegetation. By utilizing datasets with annotated images that include these complex

backgrounds, models can also be trained to detect and classify insects under more challenging

conditions [45–47]. Additionally, other techniques can be used to increase the detection and

potentially also tracking accuracy of small insects in complex scenes, e.g. by using motion-

informed enhancement of the images prior to insect detection [46].

The accuracy of the implemented object tracker depends on the inference speed of the

detection model. While most insects landing and walking on the platform are tracked with a

high precision, fast and/or erratically moving insects, as well as insects coming very close to

each other, can result in "jumping" tracking IDs and multiple counting of the same individuals.

We show that the selection of different post-processing settings regarding the exclusion of

tracking IDs with less than a specified number of images can result in more precise activity

and abundance estimations. Further experiments with other insect species under natural

conditions are necessary to validate the activity/abundance estimation by using the number of

unique tracking IDs. Implementation of additional post-processing options, e.g. by

incorporating temporal (timestamp) and spatial (insect coordinates) information from the

captured metadata could furthermore increase the accuracy of the tracking results. More

sophisticated tracking methods are available in other systems, e.g. a video-based processing

pipeline that is able to detect, track and classify insects in heterogeneous environments of

natural vegetation and enables the automated analysis of pollination events [48].

Insect classification

When using our provided insect classification model for images of insect species that were not

included in the training dataset, wrong classification results must be expected. New

approaches incorporate the taxonomic hierarchy into the classification process to identify the

lowest taxonomic rank for which a reliable classification result is achieved [49, 50]. Thereby,

also completely new insect species that were not included in the training dataset can be

correctly classified to higher taxonomic ranks (e.g. family or order), if they are morphologically

similar to other species in the respective rank, which were included in the dataset.

When only images are used for automated classification, it is currently not possible to identify

many insects to species level, as often microscopic structures have to be examined to

distinguish closely related species. New approaches could fuse data generated by different

sensors, such as images produced by an image sensor and wing beat frequency data

generated by an opto-electronic sensor, to be able to identify insect species that could not be

distinguished from similar species with images only [51].

Insect monitoring data

In ecological insect surveys, traditional monitoring methods (e.g. Malaise traps or yellow pan

traps) are usually deployed to lethally capture insect specimens that are then identified by

trained experts to species level. Due to the high effort and cost required, these methods are

often deployed in very limited timeframes and interpretation of the resulting data is restricted

by its low temporal resolution. In contrast, data captured with automated methods is available

at a significantly higher temporal resolution of up to several hours for each day and covers the

whole season. Although the taxonomic resolution is currently still low for the automated

classification of many insect taxa, combining data from both automated and traditional

methods would benefit analyses significantly and widen the scope of possible interpretations

[7]. At the same time, data on estimated insect abundance extracted from the camera trap

metadata could be systematically compared with abundances acquired from traditional

methods to calculate a conversion factor to allow for a more direct comparison of insect data

captured with both methods.

The insect monitoring data presented in this paper can be seen as a proof of concept for

automated monitoring with our proposed camera trap system. In this example, we mainly focus

on hoverflies as target species, as this group provides important ecosystem services, such as

pollination and natural pest control [52]. With camera trap data on the estimated daily activity

of different hoverfly species, phenological information on different temporal scales can be

extracted. This information could be used to e.g. study the impact of climate change on the

phenology of specific species [53]. Activity changes during the day can also be investigated,

e.g. to determine appropriate times for pesticide application with reduced impact on beneficial

insects [54].

By changing the trap design and/or by adding olfactory attractants, such as pheromone

dispensers, the camera trap system could be adapted to a more targeted monitoring of specific

pest or invasive insect species [55, 56]. This also opens the opportunity to simultaneously

monitor beneficial and pest insect species with the potential to significantly reduce pesticide

use by facilitating data-informed decision making.

671 Future perspectives

With rapid improvements in AI-based software and hardware capabilities, a broad application

of these technologies for the next generation of biodiversity monitoring tools can be expected

in the near future [7, 9, 19]. However, this also means that it is currently difficult to maintain a

methodological continuity over longer periods, which is crucial for standardized long-term

monitoring of insects. Storing the raw data generated by automated monitoring devices

together with detailed metadata [57], including in-depth descriptions of the deployed trap

design, enables reproduction and reprocessing of this data and will allow a comparison to

updated technologies in the future [58].

Contrary to traditional methods, automated camera traps generate a permanent visual record

of insects in a given location and time that can be reanalyzed with enhanced software

capabilities in the future. As it is crucial to collect as many observations as possible to identify

long-term trends and causes of insect population changes, future researchers will be grateful

for every available record from the past that is our present now [59].

Acknowledgements
686

687 We would like to thank all caretakers and owners of the orchard meadows for letting us deploy

688 the camera traps during the whole season. Thanks a lot to everybody in the automated insect

689 monitoring community for constant feedback and support.

References
691

1. Hallmann CA, Sorg M, Jongejans E, Siepel H, Hofland N, Schwan H, et al. More than

75 percent decline over 27 years in total flying insect biomass in protected areas. PLoS

ONE. 2017; 12(10): e0185809. doi: 10.1371/journal.pone.0185809

2. Seibold S, Gossner MM, Simons NK, Blüthgen N, Müller J, Ambarlı D, et al. Arthropod

decline in grasslands and forests is associated with landscape-level drivers. Nature.

2019; 574: 671–674. doi: 10.1038/s41586-019-1684-3

3.   Wagner DL. Insect Declines in the Anthropocene. Annu Rev Entomol. 2020; 65: 457–

480. doi: 10.1146/annurev-ento-011019-025151

4.   Samways MJ, Barton PS, Birkhofer K, Chichorro F, Deacon C, Fartmann T, et al.

Solutions for humanity on how to conserve insects. Biological Conservation. 2020; 242:

108427. doi: 10.1016/j.biocon.2020.108427

5.   van Klink R, Bowler DE, Gongalsky KB, Swengel AB, Gentile A, Chase JM. Meta-

analysis reveals declines in terrestrial but increases in freshwater insect abundances.

Science. 2020; 368: 417–420. doi: 10.1126/science.aax9931

6.   Breeze TD, Bailey AP, Balcombe KG, Brereton T, Comont R, Edwards M, et al.

Pollinator monitoring more than pays for itself. J Appl Ecol. 2021; 58: 44–57. doi:

10.1111/1365-2664.13755

7.   Besson M, Alison J, Bjerge K, Gorochowski TE, Høye TT, Jucker T, et al. Towards the

fully automated monitoring of ecological communities. Ecology Letters. 2022; 25: 2753–

2775. doi: 10.1111/ele.14123

8.   Kühl HS, Bowler DE, Bösch L, Bruelheide H, Dauber J, Eichenberg David, et al.

Effective Biodiversity Monitoring Needs a Culture of Integration. One Earth. 2020; 3:

462–474. doi: 10.1016/j.oneear.2020.09.010

9.   van Klink R, August T, Bas Y, Bodesheim P, Bonn A, Fossøy F, et al. Emerging

technologies revolutionise insect ecology and monitoring. Trends in Ecology &

Evolution. 2022; 37(10): 872-885. doi: 10.1016/j.tree.2022.06.001

10. Kawakita S, Ichikawa K. Automated classification of bees and hornet using acoustic

analysis of their flight sounds. Apidologie. 2019; 50: 71–79. doi: 10.1007/s13592-018-

0619-6

11. Potamitis I, Rigakis I, Fysarakis K. Insect Biometrics: Optoacoustic Signal Processing
and Its Applications to Remote Monitoring of McPhail Type Traps. PLoS ONE. 2015;
10(11): e0140474. doi: 10.1371/journal.pone.0140474

12. Rydhmer K, Bick E, Still L, Strand A, Luciano R, Helmreich S, et al. Automating insect
monitoring using unsupervised near-infrared sensors. Sci Rep. 2022; 12: 2603. doi:
10.1038/s41598-022-06439-6

13. Parmezan ARS, Souza VMA, Seth A, Žliobaitė I, Batista GEAPA. Hierarchical
classification of pollinating flying insects under changing environments. Ecological
Informatics. 2022; 70: 101751. doi: 10.1016/j.ecoinf.2022.101751

14. Droissart V, Azandi L, Onguene ER, Savignac M, Smith TB, Deblauwe V. PICT: A
low-cost, modular, open-source camera trap system to study plant-insect interactions.
Methods Ecol Evol. 2021; 12: 1389–1396. doi: 10.1111/2041-210X.13618

15. Geissmann Q, Abram PK, Wu D, Haney CH, Carrillo J. Sticky Pi is a high-frequency
smart trap that enables the study of insect circadian activity under natural conditions.
PLoS Biol. 2022; 20(7): e3001689. doi: 10.1371/journal.pbio.3001689

16. Bjerge K, Nielsen JB, Sepstrup MV, Helsing-Nielsen F, Høye TT. An Automated Light
Trap to Monitor Moths (Lepidoptera) Using Computer Vision-Based Tracking and Deep
Learning. Sensors. 2021; 21: 343. doi: 10.3390/s21020343

17. Bjerge K, Mann HMR, Høye TT. Real-time insect tracking and monitoring with computer
vision and deep learning. Remote Sens Ecol Conserv. 2021; rse2.245. doi:
10.1002/rse2.245

18. Pegoraro L, Hidalgo O, Leitch IJ, Pellicer J, Barlow SE. Automated video monitoring of
insect pollinators in the field. Emerging Topics in Life Sciences. 2020; 4: 87–97. doi:
10.1042/ETLS20190074

19.   Høye TT, Ärje J, Bjerge K, Hansen OLP, Iosifidis A, Leese F, et al. Deep learning and

746   computer vision will transform entomology. Proc Natl Acad Sci USA. 2021; 118:

747   e2002545117. doi: 10.1073/pnas.2002545117

748   20.   Darras KFA, Balle M, Xu W, Yan Y, Zakka VG, Toledo-Hernández M, et al. Eyes on

749   nature: Embedded vision cameras for multidisciplinary biodiversity monitoring. BioRxiv

750   [Preprint]. 2023 bioRxiv 2023.07.26.550656 [posted 2023 Jul 29; cited 2023 Nov 21].

751   Available from: https://www.biorxiv.org/content/10.1101/2023.07.26.550656v1 doi:

752   10.1101/2023.07.26.550656

753   21.   Pichler M, Hartig F. Machine learning and deep learning—A review for ecologists.

754   Methods Ecol Evol. 2023; 14: 994-1016. doi: 10.1111/2041-210X.14061

755   22.   Wäldchen J, Mäder P. Machine learning for image based species identification.

756   Methods Ecol Evol. 2018; 9: 2216–2225. doi: 10.1111/2041-210X.13075

757   23.   Borowiec ML, Dikow RB, Frandsen PB, McKeeken A, Valentini G, White AE. Deep

758   learning as a tool for ecology and evolution. Methods Ecol Evol. 2022; 13: 1640–1660.

759   doi: 10.1111/2041-210X.13901

760   24.   Tuia D, Kellenberger B, Beery S, Costelloe BR, Zuffi S, Risse B, et al. Perspectives in

761   machine learning for wildlife conservation. Nat Commun. 2022; 13: 792. doi:

762   10.1038/s41467-022-27980-y

763   25.   Ärje J, Raitoharju J, Iosifidis A, Tirronen V, Meissner K, Gabbouj M, et al. Human

764   experts vs. machines in taxa recognition. Signal Processing: Image Communication.

765   2020; 87: 115917. doi: 10.1016/j.image.2020.115917

766   26.   Sittinger M. Insect Detect Docs - Documentation website for the Insect Detect DIY

767   camera trap system. 2023 [cited 2023 Nov 21]. Available from:

768   https://maxsitt.github.io/insect-detect-docs/

27.   Sittinger M. Insect Detect - Software for automated insect monitoring with a DIY camera

trap system. 2023 [cited 2023 Nov 21]. Available from: https://github.com/maxsitt/insect-

detect

28.   Jocher G. YOLOv5 by Ultralytics (Version 7.0). 2022 [cited 2023 Nov 21]. Available

from: https://github.com/ultralytics/yolov5

29.   Li C, Li L, Jiang H, Weng K, Geng Y, Li L, et al. YOLOv6: A Single-Stage Object

Detection Framework for Industrial Applications. arXiv:2209.02976v1. 2022 [cited 2023

Nov 21]. Available from: http://arxiv.org/abs/2209.02976 doi:

10.48550/arXiv.2209.02976

30.   Wang C-Y, Bochkovskiy A, Liao H-YM. YOLOv7: Trainable bag-of-freebies sets new

state-of-the-art for real-time object detectors. arXiv:2207.02696v1. 2022 [cited 2023

Nov 21]. Available from: http://arxiv.org/abs/2207.02696 doi:

10.48550/arXiv.2207.02696

31.   Jocher G, Chaurasia A, Qiu J. YOLO by Ultralytics (Version 8.0.0). 2023 [cited 2023

Nov 21]. Available from: https://github.com/ultralytics/ultralytics

32.   Lin T-Y, Maire M, Belongie S, Hays J, Perona P, Ramanan D, et al. Microsoft COCO:

Common Objects in Context. In: Fleet D, Pajdla T, Schiele B, Tuytelaars T, editors.

Computer Vision – ECCV 2014. Lecture Notes in Computer Science, vol 8693.

Springer, Cham. 2014. pp. 740–755. doi: 10.1007/978-3-319-10602-1_48

33.   Sittinger M. Image dataset for training of an insect detection model for the Insect Detect

DIY camera trap. 2023 [cited 2023 Nov 21]. Database: Zenodo [Internet]. Available

from: https://zenodo.org/records/7725941 doi: 10.5281/zenodo.7725941

34.   Sittinger M. Insect Detect ML - Software for classification of images and analysis of

metadata from a DIY camera trap system. 2023 [cited 2023 Nov 21]. Available from:

https://github.com/maxsitt/insect-detect-ml

35. Sittinger M. Custom YOLOv5 fork for the Insect Detect DIY camera trap. 2023 [cited

2023 Nov 21]. Available from: https://github.com/maxsitt/yolov5

36. Deng J, Dong W, Socher R, Li L-J, Li K, Fei-Fei L. ImageNet: A large-scale hierarchical

image database. 2009 IEEE Conference on Computer Vision and Pattern Recognition.

Miami, FL: IEEE; 2009. pp. 248–255. doi: 10.1109/CVPR.2009.5206848

37. Sittinger M, Uhler J, Pink M. Insect Detect - insect classification dataset v2. 2023 [cited

2023 Nov 21]. Database: Zenodo [Internet]. Available from:

https://zenodo.org/records/8325384 doi: 10.5281/zenodo.8325384

38. Tan M, Le QV. EfficientNet: Rethinking Model Scaling for Convolutional Neural

Networks. arXiv:1905.11946v5. 2020 [cited 2023 Nov 21]. Available from:

http://arxiv.org/abs/1905.11946 doi: 10.48550/arXiv.1905.11946

39. R Core Team. R: A Language and Environment for Statistical Computing. 2023 [cited

2023 Nov 21]. R Foundation for Statistical Computing, Vienna, Austria. Available from:

https://www.R-project.org/

40. Wickham H, Averick M, Bryan J, Chang W, McGowan L, François R, et al. Welcome to

the Tidyverse. JOSS. 2019; 4(43): 1686. doi: 10.21105/joss.01686

41. Pedersen T. patchwork: The Composer of Plots. 2023 [cited 2023 Nov 21]. Available

from: https://CRAN.R-project.org/package=patchwork

42. Garnier S, Ross N, Rudis R, Camargo AP, Sciaini M, Scherer C. viridis(Lite) -

Colorblind-Friendly Color Maps for R. 2023 [cited 2023 Nov 21]. Available from:

https://CRAN.R-project.org/package=viridis

43. Sittinger, M. R scripts and data for the paper "Insect Detect: An open-source DIY

camera trap for automated insect monitoring". 2023 [cited 2023 Nov 21]. Database:

Zenodo [Internet]. Available from: https://zenodo.org/records/10171524 doi:

10.5281/zenodo.10171524

44.   Bodesheim P, Blunk J, Körschens M, Brust C-A, Käding C, Denzler J. Pre-trained

models are not enough: active and lifelong learning is important for long-term visual

monitoring of mammals in biodiversity research—Individual identification and attribute

prediction with image features from deep neural networks and decoupled decision

models applied to elephants and great apes. Mamm Biol. 2022; 102: 875-897. doi:

10.1007/s42991-022-00224-8

45.   Bjerge K, Alison J, Dyrmann M, Frigaard CE, Mann HMR, Høye TT. Accurate detection

and identification of insects from camera trap images with deep learning. Fang W-T,

editor. PLOS Sustain Transform. 2023; 2: e0000051. doi: 10.1371/journal.pstr.0000051

46.   Bjerge K, Frigaard CE, Karstoft H. Object Detection of Small Insects in Time-Lapse

Camera Recordings. Sensors. 2023; 23: 7242. doi: 10.3390/s23167242

47.   Stark T, Ştefan V, Wurm M, Spanier R, Taubenböck H, Knight TM. YOLO object

detection models can locate and classify broad groups of flower-visiting arthropods in

images. Sci Rep. 2023; 13: 16364. doi: 10.1038/s41598-023-43482-3

48.   Ratnayake MN, Amarathunga DC, Zaman A, Dyer AG, Dorin A. Spatial Monitoring and

Insect Behavioural Analysis Using Computer Vision for Precision Pollination. Int J

Comput Vis. 2022; 131: 591-606. doi: 10.1007/s11263-022-01715-4

49.   Bjerge K, Geissmann Q, Alison J, Mann HMR, Høye TT, Dyrmann M, et al. Hierarchical

classification of insects with multitask learning and anomaly detection. Ecological

Informatics. 2023; 77: 102278. doi: 10.1016/j.ecoinf.2023.102278

50.   Badirli S, Picard CJ, Mohler G, Richert F, Akata Z, Dundar M. Classifying the unknown:

Insect identification with deep hierarchical Bayesian learning. Methods Ecol Evol. 2023;

14: 1515–1530. doi: 10.1111/2041-210X.14104

51.   Tschaikner M, Brandt D, Schmidt H, Bießmann F, Chiaburu T, Schrimpf I, et al.

843   Multisensor data fusion for automatized insect monitoring (KInsecta). In: Neale CM,

844   Maltese A, editors. Proc. SPIE 12727, Remote Sensing for Agriculture, Ecosystems,

845   and Hydrology XXV, 1272702. 2023. doi: 10.1117/12.2679927

846   52.   Rodríguez-Gasol N, Alins G, Veronesi ER, Wratten S. The ecology of predatory

847   hoverflies as ecosystem-service providers in agricultural systems. Biological Control.

848   2020; 151: 104405. doi: 10.1016/j.biocontrol.2020.104405

849   53.   Forrest JR. Complex responses of insect phenology to climate change. Current Opinion

850   in Insect Science. 2016; 17: 49–54. doi: 10.1016/j.cois.2016.07.002

851   54.   Karbassioon A, Stanley DA. Exploring relationships between time of day and pollinator

852   activity in the context of pesticide use. Basic and Applied Ecology. 2023; 72: 74–81. doi:

853   10.1016/j.baae.2023.06.001

854   55.   Preti M, Verheggen F, Angeli S. Insect pest monitoring with camera-equipped traps:

855   strengths and limitations. J Pest Sci. 2021; 94: 203–217. doi: 10.1007/s10340-020-

856   01309-4

857   56.   Teixeira AC, Ribeiro J, Morais R, Sousa JJ, Cunha A. A Systematic Review on

858   Automatic Insect Detection Using Deep Learning. Agriculture. 2023; 13: 713. doi:

859   10.3390/agriculture13030713

860   57.   Reyserhove L, Norton B, Desmet P. Best Practices for Managing and Publishing

861   Camera Trap Data. Community review draft. 2023 [cited 2023 Nov 21]. Available from:

862   https://docs.gbif-uat.org/camera-trap-guide/en/ doi: 10.35035/doc-0qzp-2x37

863   58.   van Klink R. Delivering on a promise: Futureproofing automated insect monitoring

864   methods. EcoEvoRxiv [Preprint]. 2023 [posted 2023 Sep 01; revised 2023 Oct 26; cited

865   2023 Nov 21]. Available from: https://ecoevorxiv.org/repository/view/5888/ doi:

866   10.32942/X2PW3X

867   59.   Kitzes J, Schricker L. The Necessity, Promise and Challenge of Automated Biodiversity

868   Surveys. Envir Conserv. 2019; 46: 247–250. doi: 10.1017/S0376892919000146

Supporting information

S1 Table. Description of the 27 classes from the image dataset that was used to train the insect classification model. The images were sorted to the respective class by considering taxonomic and visual distinctions. In some cases, clear taxonomical separations are difficult from images only and the decision to sort an image to the respective class was based more on visual distinction.

S2 Table. Comparison of different classification model architectures and hyperparameter settings supported by YOLOv5 classification model training. All models were trained on a custom dataset with 21,000 images (14,686 in train split) and default hyperparameters. Metrics are shown on the dataset validation split (4,189 images) and dataset test split (2,125 images) for the converted models in ONNX format.

S3 Table. Metrics of the EfficientNet-B0 insect classification model, validated on a real-world dataset. The model was trained on a custom dataset with 21,000 images (14,686 in train split), scaled to 128x128 pixel, to 20 epochs with batch size 64 and default hyperparameters. Metrics are shown on a real-world dataset (97,671 images) for the converted model in ONNX format. All images were classified and subsequently verified and sorted to the correct class in the case of a wrong classification by the model. A dummy image was added for each of the classes "bug_grapho" and "fly_empi", as no images of both were captured during the deployment period. Results for both classes must be ignored.

S1 Fig. Examples for limitations in the detection and tracking accuracy. (A) The same tracking ID is assigned to insects coming close to each other. (B) A fast-moving insect is not correctly tracked, with the risk of a new tracking ID being assigned to the same individual.

S2 Fig. Total number of unique tracking IDs for each predicted class. Merged data from all five camera traps deployed from mid-May to mid-September 2023. All tracking IDs with less than three or more than 1,800 images were removed.

S3 Fig. Total recording time per time of day. Merged data from all five camera traps deployed from mid-May to mid-September 2023.

S4 Fig. Estimated bee activity (number of unique tracking IDs per hour) per time of day. Merged data from all five camera traps deployed from mid-May to mid-September 2023. Shaded areas indicate hours without recordings. All tracking IDs with less than three or more than 1,800 images were removed.

S5 Fig. Estimated fly activity (number of unique tracking IDs per hour) per time of day. Merged data from all five camera traps deployed from mid-May to mid-September 2023. Shaded areas indicate hours without recordings. All tracking IDs with less than three or more than 1,800 images were removed.

Fig1

OAK-1

Image sensor
12MP resolution

CROP

4K resolution

DOWNSCALE
(optional)

HQ frame
1920x1080 px
/ 3840x2160 px

STRETCH + DOWNSCALE

LQ frame
320x320 px

Detection model

DETECTIONS

Object tracker

Script node
sync HQ frame with
detections & tracklets

DETECTIONS

TRACKLETS

RPi Zero 2 W

cropped detections
from HQ frame
+ metadata

microSD card

Fig2

Fig3

Fig4

Fig5

Fig6

Fig7

Fig8

Fig9

Fig10

Fig11

Fig12

Fig13

Fig14

Fig15

Fig16