1 **Expanding the toolbox: Novel class IIb microcins show activity against Gram-negative**
2 **ESKAPE and plant pathogens**
3

Benedikt M. Mortzfeld[1,2,*], Shakti K. Bhattarai[1,2], Vanni Bucci[1,2,3]
[1]Department of Microbiology, University of Massachusetts Medical School, Worcester, MA, USA
[2]Program in Microbiome Dynamics, University of Massachusetts Medical School, Worcester, MA,
USA,
[3]Program in Systems Biology, University of Massachusetts Medical School, Worcester, MA, USA
*Corresponding author

10
11 **Short title: Novel class IIb microcins show activity against pathogens**
12

Correspondence to:
Benedikt M. Mortzfeld benedikt.mortzfeld@umassmed.edu

Competing Interests
The authors of this manuscript have the following competing interests: V.B. receives support from a
sponsored research agreement from Vedanta Biosciences, Inc. The authors B.M.M. and S.K.B have
declared that no competing interests exist.

Abstract

Interspecies interactions involving direct competition via bacteriocin production play a vital role in shaping ecological dynamics within microbial ecosystems. For instance, the ribosomally-produced siderophore bacteriocins, known as class IIb microcins, affect the colonization of host-associated pathogenic *Enterobacteriaceae* species. Notably, to date, only five of these antimicrobials have been identified, all derived from specific *Escherichia coli* and *Klebsiella pneumoniae* strains. We hypothesized that class IIb microcin production extends beyond these specific compounds and organisms. With a customized informatics-driven approach, screening bacterial genomes in public databases with BLAST and manual curation, we have discovered twelve previously unknown class IIb microcins in seven additional *Enterobacteriaceae* species, encompassing phytopathogens and environmental isolates. We introduce three novel clades of microcins (MccW, MccX, and MccZ), while also identifying eight new variants of the five known class IIb microcins. To validate their antimicrobial potential, we heterologously expressed these microcins in *E. coli* and demonstrated efficacy against a variety of bacterial isolates, including plant pathogens from the genera *Brenneria*, *Gibbsiella,* and *Rahnella*. Two newly discovered microcins exhibit activity against Gram-negative ESKAPE pathogens, *i.e. Acinetobacter baumannii* or *Pseudomonas aeruginosa,* providing the first evidence that class IIb microcins can target bacteria outside of the *Enterobacteriaceae* family. This study underscores that class IIb microcin genes are more prevalent in the microbial world than previously recognized and that synthetic hybrid microcins can be a viable tool to target clinically relevant drug-resistant pathogens. Our findings hold significant promise for the development of innovative engineered live biotherapeutic products tailored to combat these resilient bacteria.

Introduction

A large body of theoretical and experimental work has shown that dynamics of microbiomes are shaped by the network of interbacterial interactions[1–4]. These cooperative and competitive interactions are often achieved *via* the secretion of cross-feeding metabolites[5], antimicrobial peptides[6] and bacterially produced small molecules[7, 8] and are crucial to ecological properties including stability and ability to respond to external perturbations[9, 10]. Among the competitive interactions, bacteriocin production is proposed to be a prominent mediator of microbiome dynamics[11] and, specifically, several reports including ours have shown that a bacteriocin subclass, Class IIb microcins, mediates *Enterobacteriaceae* dynamics *in vivo*[12–14].

Class IIb microcins are ribosomally synthesized bacteriocins between 5 kDa to 10 kDa in size with activity against closely related strains or species[13, 15–18]. Unlike all other microcins, they carry a serine-rich C-terminal motif for a posttranslational modification with a siderophore, here an enterobactin or an enterobactin derivative, before they are secreted into the extracellular space[19]. Siderophores are iron-chelating molecules commonly employed by various bacteria to scavenge free iron to compete with other bacteria, particularly in resource-scarce environments such as the gastrointestinal tract[16, 17, 20] and are often associated with increased pathogenicity or virulence[21–23]. The iron chelating moiety of these posttranslationally modified antimicrobial peptides is recognized by high-affinity receptors and functions as a Trojan Horse key to susceptible bacteria as it triggers import into the periplasmic space, where the peptide inhibits the molecular target of susceptible bacteria[20, 24–26]. Because of these features, delivery of class IIb microcins by wildtype and engineered probiotics has been recently proposed as a strategy to combat drug-resistant enteric bacteria[12, 13, 17, 27], which is in line with a growing body of work from the past decade that explores siderophore conjugation, including with enterobactin, to specifically deliver antibiotics and other small molecules to drug-resistant Gram-negative pathogens[28–30].

To date only five class IIb microcins have been described and only four have been characterized in terms of their antimicrobial activity. Specifically, the class IIb microcins MccE492 and MccG492 (uncharacterized) are solely present in *Klebsiella pneumoniae* (*Kp*), whereas MccH47 is specific for *Escherichia coli (Ec)*[16]. Additionally, truncated versions of *mciA* (MccI47) and *mcmM* (MccM) are present in *Kp* RYC492, whereas they are intact in the *Ec* CA46 genome[16]. Interestingly, while the genes encoding for microcin posttranslational modifications are highly conserved between *Ec* and *Kp*, suggesting a conserved pathway for microcin maturation, the toxin and corresponding immunity genes are significantly more variable (**Fig S1**). We hypothesized that class IIb microcin production extends beyond these specific compounds and organisms and identified a total of twelve novel class IIb microcins in seven additional *Enterobacteriaceae* species. Utilizing heterologous expression of these compounds in our *E. coli* system optimized for enterobactin conjugation, we show potent antimicrobial activity by the encoded toxins against a library of bacteria, including Gram-negative ESKAPE and plant pathogens. This demonstrates that class IIb microcin genes are more prevalent in the microbial world than previously recognized and that synthetic hybrid microcins can be a viable tool to target clinically relevant drug-resistant pathogens.

Materials and Methods

Bioinformatic class IIb microcin identification

We developed a pipeline that by leveraging BLAST[31] enabled us to mine publicly available

genome databases for novel, previously undescribed class IIb microcins. We included *mchCDEF* and

mcmL for *Ec* as well as *mceCDGHIJ* for *Kp* for posttranslational modification and export, expecting

more reliable hits for longer and functionally conserved proteins in close proximity to class IIb

microcin and immunity genes. Thus, we first ran *tblastn*[31] against RefSeq[32], to screen for all

genes related to biosynthesis pathways, known microcin genes, as well as immunity gene

sequences exhibiting homology to the microcin gene clusters found in *Ec* CA46 and *Kp* RYC492.

Homology to the microcin gene clusters were guided by BLAST parameters sseqid (Genome ID) ,

pident (percentage of identical positions) along with sstart (start of alignment in Genome) and send

(end of alignment position in Genome). Resulting hits were concatenated by genome ID and

assessed for their proximity to one another in the genome. These gene clusters should, at best,

contain all the known genes required for toxin maturation, including *mchCDEF* and *mcmL*[16, 17].

In addition to genomic hits to the known microcins, small ORFs of 50 to 150 amino acids in size

close to the biosynthesis genes were screened and annotated manually using the criteria described

below as well as their domains were predicted using SMART[33]. The ORFs were meticulously

examined and assessed against established class IIb microcin criteria known from *Ec* H47, *Ec* I47,

Ec M, *Kp* E492, and *Kp* G492: (i) a serine-rich C-terminus culminating in a final serine, (ii) the

presence of fewer than two cysteine residues, (iii) a signal peptide within the initial 15 amino acids

ending with GG or GA, and (iv) close proximity (≤200 bp) to an ORF featuring a predicted

transmembrane domain, typically encoding an immunity peptide. The identified genes were

included in the pipeline's input to expand the scope of gene detection. We repeated this process

iteratively through the pipeline until no additional genes were added to the output. Subsequently,

blastp was used to assess microcin similarity shown in Table 1 and Table S1.

Phylogenetic analyses

For the native full length coding sequence of the microcin and immunity genes a codon-based

sequence alignment was generated using the MUSCLE algorithm[34]. For phylogeny of all

microcins, the nucleotide sequences without the respective signal peptides were codon-aligned.

Subsequently, we determined the best fit substitution models for maximum likelihood phylogenetic

analyses, resulting in the General Time Reversible model with discrete gamma distribution

(GTR+G) and the Hasegawa-Kishino-Yano model with discrete gamma distribution (HKY+G),

respectively. A bootstrap test with 1000 replicates for maximum likelihood and random seed was

conducted for all trees. Alignment, model testing, and tree building was performed in MEGA11[35].

antiSMASH analyses

To test if similar results of class IIb microcin identification could be obtained with automated

bioinformatic tools, we ran antiSMASH 7.0[36], a widely used tool for microbial genome mining and

biosynthetic gene cluster detection. As input, we utilized the seven genomes from the newly

identified class IIb microcins: (i) *Bg* CP014137, (ii) *Gq* CP014136, (iii) *Ko* CP033844, (iv) *Ps*

CP034363, (v) *Ro* CP008886, (vi) *Se* CP030220, (vii) *Sf* CP033055. As a positive control for the well-

established microcins *Kp* E492 and *Kp* G492 as well as *Ec* H47 and *Ec* M we used the accession

numbers CP127839 (*Kp* RYC492) and CP148105 (*Ec* Nissle 1917), respectively. Notably, using the

"loose" setting, in none of the cases a class IIb microcin biosynthesis gene cluster was detected, nor

were any microcin genes identified. This was the case for both, the novel microcins and the original,

well-annotated, microcins.

Plasmids and heterologous class IIb microcin expression

ORFs of identified microcin and immunity genes were codon optimized for frequent *Ec* codon usage

without creating repetitive sequences and synthesized by Integrated DNA Technologies (Coralville,

IA) with 18 bp of native 5' upstream sequence and 20 bp of native 3' downstream sequence,

respectively. Using Gibson Assembly[37], the genes were cloned into our previously established *Ec*

class IIb microcin expression system that results in mature class IIb microcins posttranslationally

modified with an MGE[13, 17]. Briefly, the antimicrobial and the immunity genes are co-expressed

under the control of an arabinose-inducible pBad/araC promoter in a high copy plasmid with a

pUC-derived origin of replication. All assemblies were verified using whole plasmid sequencing.

DNA files for all used plasmids can be found as supplementary material.

Static inhibition assays

Cultures of strains with confirmed plasmid assemblies were spread in LB agar plates containing

100 µg/ml ampicillin. In addition to a pUC19 control without microcin expression, single colonies

for each microcin were picked with a sterile pipet tip and all placed into the same solid LB agar

plate containing 100 µg/ml ampicillin for plasmid retention, 0.2 mM 2,2-dipyridyl to create iron-

limited conditions during the growth phase, and 0.4% L-arabinose for induction of gene expression.

Plates were incubated at 37°C for up to 72 h, before they were overlaid with the target bacterial

isolates. Note that testing all microcin-expressing stains on the same plate allowed us to confidently
assess differential inhibitory activity between all 17 tested microcins. For the overlay, the microcin-
producing bacteria in the stabs were inactivated using chloroform vapors and ten minutes under
ultraviolet light. Then, target bacteria were diluted 1:2000 from overnight culture in LB media
containing 100 µg/ml ampicillin and 0.2 mM 2,2-dipyridyl. *Ec* and *S. flexneri* strains were diluted
1:200 to acquire dense bacterial lawns. Finally, 0.5 ml of molten agar was added to the liquid media
and the resulting soft agar was spread on the plate with the inactivated bacteria and incubated for
16 h at 37°C. The pUC19 control strain was unable to create any zone of inhibition against any of
the tested target bacteria.

Relative MIC dilution factors

For enrichment of microcin *Se* G492, an MBP-microcin fusion protein was expressed from pHMT-
SeG492 in *E. coli* BL21 cells as previously described[13, 17]. Harvested cells were resuspended in
column buffer (200 mM NaCl, 20 mM Tris-HCl, pH 7.5), lysed by sonication, and passed through a
high flow amylose resin (*New England Biolabs, Ipswich, MA*) as recommended by the manufacturer.
The protein was eluted with 10 mM maltose, cleaved with Tobacco etch virus (TEV) protease, and
further processed as previously reported[13, 17]. The relative MIC assays were conducted using
sterile 96-well round bottom microplates. The plates were prepared as follows: the first row
contained 20 µl of 2x LB with 0.4 mM 2,2'-dipyridyl and 20 µl of *Se* G492 containing solution in
amylose resin elution buffer (200 mM NaCl, 20 mM Tris-HCl, 10 mM maltose, pH 7.5). All other
wells were filled with 20 µl of 1x LB, 0.2 mM 2,2'-dipyridyl, and 0.5× amylose resin elution buffer,
and a two-fold serial dilution was performed across the plate. The target bacteria were grown
overnight in LB at 200 rpm and 37°C and were added to a final dilution of 10^{-4} into the wells. The
plates were then incubated in the dark at 37°C with gentle agitation. Relative MICs were
determined as the lowest concentration at which no growth was observed after 24 hours. All
reported values represent the median of at least three biological replicates.

Results

With the hypothesis that class IIb microcin production is a common trait among
Enterobacteriaceae, we anticipated that the genes encoding for the antimicrobial and immunity
would exhibit a high degree of dissimilarity to already known peptides as target specificity may
result in accelerated adaptive coevolution[38]. Therefore, in addition to the known microcin and
immunity genes from MccE492, MccG492, MccH47, MccI47, and MccM, in our informatic approach

we included the genes that are necessary for mature class IIb microcin biosynthesis, extending our search to longer sequences for more reliable Basic Local Alignment Search Tool (BLAST) results[31]. Moreover, we hypothesized that the amino acid sequences of genes responsible for posttranslational modification and microcin export would be less prone to evolutionary changes, thereby maintaining the functional integrity of the gene cluster[16, 39]. We then assessed their proximity in the respective genome location, because microcin genes are typically flanked by genes essential for toxin maturation[16]. Further, we manually assessed and annotated small open reading frames (ORFs) upstream and downstream of the maturation genes, allowing us to also identify novel class IIb microcins without significant sequence similarity to the known antimicrobials, enabling the discovery of compounds with new molecular targets or modes of action (see Methods).

Our informatics-driven analysis identified twelve promising class IIb microcin candidates from seven gene clusters with high similarity to *Ec* CA46 and *Kp* RYC492 in seven species across the *Enterobacteriaceae* family (**Fig 1A, Fig S2**): (i) *Brenneria goodwinii* (*Bg*; 2; GenBank: CP014137), (ii) *Gibbsiella quercinecans* (*Gq*; 1; CP014136), (iii) *Klebsiella oxytoca* (*Ko*; 1; CP033844), (iv) *Pantoea sp.* (*Ps*; 1; CP034363), (v) *Raoultella ornithinolytica* (*Ro*; 4; CP008886), (vi) *Salmonella enterica* (*Se*; 2; CP030220), (vii) *Serratia fonticola* (*Sf*; 1; CP033055). Although it has traditionally been a defining characteristic of class IIb microcins that all required genes are encoded within the chromosome[12], the gene cluster we discovered for *Se* is situated on a 159 kbp plasmid. Phylogenetic sequence analysis of both the antimicrobial and immunity peptide genes revealed the presence of eight different clades represented in both trees, respectively (**Fig 1B,C**). Regarding the well-established class IIb microcins MccH47, MccI47, MccM, MccG492, and MccE492, we identified novel members for each group, supported by nucleotide sequence similarity, amino acid identity, the closest *blastp* match, and domain predictions (**Table 1, Fig S3**). It is important to note that application of established tools for secondary metabolite identification (*e.g.*, antiSMASH 7.0)[36] to these genomes did not yield identification of any of the old or novel microcins providing support of the relevance of our approach. In order to then ensure that these novel microcins are unique and not part of any other microcin class, we performed phylogenetic analysis for all known microcin genes from the classes I, IIa, and IIb and show distinct clustering for all newly described sequences (**Fig S4**). In light of this discovery, we propose a new nomenclature for class IIb microcins that includes the species initials in which they were identified (*e.g.*, *Ec*, *Kp*), the closest relative already characterized class IIb microcin (G492, E492, H47, I47 or M), as well as the identifiers 'A' for antimicrobial or 'I' for immunity gene.

Based on this, the novel G492 relative found in *Salmonella enterica* will be called *Se* G492
with the antimicrobial peptide identified as *Se* G492A and the immunity peptide identified as *Se*
G492I. It is worth highlighting that in the case of the G492 group, all its members have the
immunity gene located downstream of the antimicrobial gene, whereas for the other clades, this
arrangement is reversed. In addition to uncovering eight novel variants of the five previously
characterized microcins, we have identified four additional microcins through manual curation of
ORFs in proximity to the microcin maturation genes. These novel microcins, which we name
microcin W (MccW), microcin X (MccX), and microcin Z (MccZ), seem to belong to three entirely
new clades based on nucleotide similarity (**Fig 1 B,C**). The two members of the microcin X group,
found in *B. goodwinii* (*Bg* X) and *R. ornithinolytica* (*Ro* X), only show significant similarity between
one another, but not to any of the other antimicrobial or immunity peptides. This holds true for the
nucleotide similarity (**Fig 1 B,C**) as well as amino acid identity and the closest *blastp* hits (**Fig. 1D,**
Table 1, Table S1). Similarly, MccW from *Gibbsiella quercinecans* (*Gq* W) does not show any
sequence similarity to either the known or novel antimicrobial or immunity peptides in terms of
nucleotide similarity, amino acid identity, the respective *blastp* hits, or phylogenetic localization
(**Fig 1,Table 1, Table S1**). Lastly, MccZ from *R. ornithinolytica* (Ro Z) shows insignificant amino
acid similarity with *Ec* MA (*mcmA*) for the antimicrobial, whereas the immunity peptide does not
have any match among the known or the novel microcins (**Fig 1 C, Table 1, Table S1**). Crucially,
the identification of MccX, and MccZ within the same gene clusters as representatives of the E492
(*Bg* E492), H47 (*Ro* H47), and I47 (*Ro* I47) groups strongly implies that they are functional
components of a microcin gene cluster.
To test the newly identified microcins for antimicrobial activity we used our previously
established *Ec* overexpression system[13, 17]. All antimicrobial and immunity peptides were codon
optimized, synthesized, and cloned into an inducible high copy vector (see Methods). Thus, we
extracted the novel microcins out of their native genomic context of siderophore biosynthesis and
transferred them into a heterologous expression background optimized for microcin-
monoglycosylated enterobactin (MGE) linkage[13, 17]. This allowed us to create hybrid compounds
that could be efficiently tested for antimicrobial activity in an *E. coli* background. Through static
plate inhibition assays involving live-producing cells[13, 17, 27], we successfully validated robust
antimicrobial activity of eleven out of the twelve newly discovered microcins (**Fig 2A**). Notably,
antimicrobial activity was only observed in iron-depleted media (**Fig 2B,C**). The hybrid microcins
exhibit a range of specificities, with some inhibiting targets narrowly (*e.g., Ps* G492AI), while others
exert a broader effect against multiple bacteria (*e.g., Se* G492AI). Moreover, our study also provides

the first evidence of inhibitory activity by *Kp* G492, a microcin whose existence and function have only been proposed in the scientific literature based on genetic sequence[16].

To this date class IIb microcins have been only shown to be very selective and only active against different species within the *Enterobacteriaceae* family[13, 16, 17, 27]. While the activity for the novel microcins varies, we here report, for the first time, antimicrobial activity outside of the *Enterobacteriaceae* family utilizing hybrid antimicrobial peptides. We demonstrate that microcins *Ps* G492 and *Se* G492 have activity against Gram-negative multidrug-resistant ESKAPE pathogens with both being capable of inhibiting *Acinetobacter baumannii* (BAA 1790), and with microcin *Se* G492 alone also showing activity against *Pseudomonas aeruginosa* (PA14) (**Fig 2**). Specifically, compared to *K. pneumoniae* (BAA 1705) *Se* G492 is 256-times more effective against *A. baumannii* (BAA 1790), 128-times more effective against *E. coli* (BAA 196), and 8-times more effective against *P. aeruginosa* (PA14) (**Fig. 2B**).

Discussion

With a comprehensive analysis of publicly available bacterial genomes, we unraveled twelve previously undiscovered class IIb microcins. Among these findings, we identified three novel microcin clades, specifically MccW, MccX, and MccZ. Through heterologous expression, we showed antimicrobial activity for all but one novel microcins and are the first to demonstrate activity for the known class IIb microcin *Kp* G492. Hence, this research demonstrates that class IIb microcin genes exhibit a higher prevalence in *Enterobacteriaceae* genomes than previously reported. As a result, their impact on ecological community dynamics in natural environments, including the growth of *Pseudomonadales* species, might be broader than previously thought. For antimicrobial activity testing, microcin and immunity genes were overexpressed recombinantly in our *Ec*-derived expression system optimized for microcin-MGE production[13, 17]. The common process of posttranslational modification with the siderophore consolidated the import mechanism of the hybrid microcins towards enterobactin, the most characteristic siderophore of the *Enterobacteriaceae* family. This allowed us to test the target-specific antimicrobial activity of the microcins irrespective of siderophore production in the native genomic background. However, it is important to note that class IIb microcin activity is dependent on active import through siderophore receptors and consequently some of these microcins might display different activity spectrums when tested in their native genomic background of siderophore biosynthesis. Furthermore, static plate inhibition assays exhibit lower sensitivity compared to purification approaches with quantitative minimum inhibitory concentration (MIC) assays. Thus, the activity

spectrums of the hybrid microcins could encompass a wider range than what has been described in
this study when tested as a purified product. However, historically the microcin literature proves
that ideal approaches for purification and MIC testing can vary between the antimicrobials[13, 16,
17, 25].

We were able to expand the origins of class IIb microcins from the enteric bacteria *Ec* and
Kp to other members of the *Enterobacteriaceae* family, including well-known phytopathogens[40–
42]. Specifically, *B. goodwinii* and *G. quercinecans* are associated with Acute Oak Decline (AOD) and
are frequently isolated together[43] and the two strains containing microcin genes were isolated
within the same research project. Notably, these bacteria grow synergistically[44], while
upregulating iron transporters during co-culture[45], hinting at class IIb microcin-related
competition. We were able to show activity of the overexpressed hybrid microcins against human-
derived enteric isolates, however, their native spectrum might have evolved to target more
frequently encountered strains from the genus *Brenneria* or *Gibbsiella*. Further, we demonstrated
activity of several class IIb microcins against the three tree pathogen genera *Brenneria*, *Gibbsiella*,
as well as *Rahnella*[44, 46–49]. Thus, treatment with potent microcins, purified or produced in live
bacteria, could present a viable option to target bacteria-caused plant diseases.

In health care settings the burden by Gram-negative ESKAPE pathogens and multidrug-
resistant *Enterobacteriaceae* weighs heavily on modern medicine and novel antimicrobials are
needed to develop new treatment options[50, 51]. In addition to enteric pathogens and
pathobionts, bacteria outside of the *Enterobacteriaceae* family have also been shown to scavenge
for and to import enterobactin, including *P. aeruginosa* and *A. baumannii* [52, 53]. Therefore,
different siderophore conjugates could be a viable option to target these pathogens as well or to
finetune the desired target range[28–30]. Antimicrobial peptides and particularly microcins are
promising candidates for selective eradication of enteric pathogens and have been demonstrated to
potently reduce pathogen colonization *in vivo*, when produced by a live probiotic[12, 13]. Here we
present the most comprehensive library of class IIb microcins created so far, that is suited for
heterologous expression and *in vivo* application for the development of novel live biotherapeutic
products against drug-resistant enteric bacteria and Gram-negative ESKAPE pathogens.

In this study, we challenge the prevailing notion that class IIb microcin production is limited
to *Ec* and *Kp*. Through comprehensive genomic analysis of publicly available bacterial genomes,
coupled with heterologous overexpression, we unveiled a set undiscovered class IIb microcins
across *Enterobacteriaceae* species. Our findings not only expand the known repertoire of class IIb
microcins but also hold significant implications for synthetic hybrid compounds. We demonstrate

that these newly identified class IIb microcins exert remarkable inhibitory effects on ESKAPE

pathogen species when expressed in a system for enterobactin-derived conjugation. This discovery

underscores their potential as agents against a broader spectrum of pathogens, including those

affecting humans and plants, thus opening new avenues for antimicrobial research and

applications.

Acknowledgements

The author would like to thank the administrative staff in the Microbiology Department at the

University of Massachusetts Chan Medical School, especially Annette Bohigian, Amy Parker, Dhruti

Desai, Marie Berardi, Richard Fish, and Tracey Rae, for their support.

Competing Interests

The authors of this manuscript have the following competing interests: V.B. receives support from a

sponsored research agreement from Vedanta Biosciences, Inc. The authors B.M.M. and S.K.B have

declared that no competing interests exist.

Data Availability Statement

The genomes are accessible with the following GenBank numbers: *Brenneria goodwinii*

(CP014137), *Gibbsiella quercinecans* (CP014136), *Klebsiella oxytoca* (CP033844), *Pantoea sp.*

(CP034363), R*aoultella ornithinolytica* (CP008886), *Salmonella enterica* (CP030220), *Serratia*

fonticola (CP033055). All information is included in the manuscript or supporting files. All plasmid

sequences as well as annotation files to produce Figs 1A, S1, S2, and S3 are available as

supplementary material.

Funding Disclosure

This work was supported by the CDMRP PRMP W81XWH2020013 to V.B., by the NIH NIA

1R01AG075283-01A1 to V.B., and by the Deutsche Forschungsgemeinschaft (DFG) project

457837076 to B.M.M. The funders had no role in study design, data collection and analysis,

decision to publish, or preparation of the manuscript.

Author Contributions

B.M.M, S.K.B, and V.B conceptualized the study. S.K.B. performed bioinformatic data mining. B.M.M

and V.B designed the experiments. B.M.M performed plasmid design, verification, and *in vitro*

activity testing. B.M.M, S.K.B, and V.B wrote the manuscript.

References

References

1. Stein RR, Bucci V, Toussaint NC, Buffie CG, Rätsch G, Pamer EG, et al. Ecological
Modeling from Time-Series Inference: Insight into Dynamics and Stability of Intestinal
Microbiota. *PLoS Comput Biol* 2013; **9**: e1003388.

2. Bucci V, Xavier JB. Towards Predictive Models of the Human Gut Microbiome. *J Mol
Biol* 2014; **426**: 3907–3916.

3. Coyte KZ, Schluter J, Foster KR. The ecology of the microbiome: Networks, competition,
and stability. *Science (1979)* 2015; **350**: 663–666.

4. Hromada S, Venturelli OS. Gut microbiota interspecies interactions shape the response of
Clostridioides difficile to clinically relevant antibiotics. *PLoS Biol* 2023; **21**: e3002100.

5. Culp EJ, Goodman AL. Cross-feeding in the gut microbiome: Ecology and mechanisms.
Cell Host Microbe 2023; **31**: 485–499.

6. Heilbronner S, Krismer B, Brötz-Oesterhelt H, Peschel A. The microbiome-shaping roles
of bacteriocins. *Nature Reviews Microbiology 2021 19:11* 2021; **19**: 726–739.

7. Hibbing ME, Fuqua C, Parsek MR, Peterson SB. Bacterial competition: surviving and
thriving in the microbial jungle. *Nature Reviews Microbiology 2009 8:1* 2009; **8**: 15–25.

8. Donia MS, Fischbach MA. Small molecules from the human microbiota. *Science (1979)*
2015; **349**.

9. Coyte KZ, Rakoff-Nahoum S. Understanding Competition and Cooperation within the
Mammalian Gut Microbiome. *Current Biology* 2019; **29**: R538–R544.

10. Coyte KZ, Rao C, Rakoff-Nahoum S, Foster KR. Ecological rules for the assembly of
microbiome communities. *PLoS Biol* 2021; **19**: e3001116.

11. Niehus R, Oliveira NM, Li A, Fletcher AG, Foster KR. The evolution of strategy in
bacterial warfare via the regulation of bacteriocins and antibiotics. *Elife* 2021; **10**.

12. Sassone-Corsi M, Nuccio S-P, Liu H, Hernandez D, Vu CT, Takahashi AA, et al.
Microcins mediate competition among Enterobacteriaceae in the inflamed gut. *Nature*
2016; **540**: 280–283.

13. Mortzfeld BM, Palmer JD, Bhattarai SK, Dupre HL, Mercado-Lubio R, Silby MW, et al.
Microcin MccI47 selectively inhibits enteric bacteria and reduces carbapenem-resistant
Klebsiella pneumoniae colonization in vivo when administered via an engineered live
biotherapeutic. *Gut Microbes* 2022; **14**.

14. Cherrak Y, Salazar MA, Yilmaz K, Kreuzer M, Hardt WD. Commensal E. coli limits
Salmonella gut invasion during inflammation by producing toxin-bound siderophores in a
tonB-dependent manner. *PLoS Biol* 2024; **22**: e3002616.

15. de Lorenzo V. Isolation and characterization of microcin E 492 from Klebsiella
pneumoniae. *Arch Microbiol* 1984; **139**: 72–75.

16. Vassiliadis G, Destoumieux-Garzón D, Lombard C, Rebuffat S, Peduzzi J. Isolation and
characterization of two members of the siderophore-microcin family, microcins M and
H47. *Antimicrob Agents Chemother* 2010; **54**: 288–297.

17. Palmer JD, Mortzfeld BM, Piattelli E, Silby MW, McCormick BA, Bucci V. Microcin
H47: A Class IIb Microcin with Potent Activity Against Multidrug Resistant
Enterobacteriaceae. *ACS Infect Dis* 2020; **6**: 672–679.

18. Baquero F, Lanza VF, Baquero MR, del Campo R, Bravo-Vázquez DA. Microcins in
Enterobacteriaceae: Peptide Antimicrobials in the Eco-Active Intestinal Chemosphere.
Front Microbiol . 2019. Frontiers Media S.A. , **10**: 2261

19. Azpiroz MF, Rodríguez E, Laviña M. The structure, function, and origin of the microcin
H47 ATP-binding cassette exporter indicate its relatedness to that of colicin V. *Antimicrob*
Agents Chemother 2001; **45**: 969–972.
20. Patzer SI, Baquero MR, Bravo D, Moreno F, Hantke K. The colicin G, H and X
determinants encode microcins M and H47, which might utilize the catecholate
siderophore receptors FepA, Cir, Fiu and IroN. *Microbiology (N Y)* 2003; **149**: 2557–
2570.
21. Miethke M, Marahiel MA. Siderophore-Based Iron Acquisition and Pathogen Control.
Microbiology and Molecular Biology Reviews 2007; **71**: 413–451.
22. Wilson BR, Bogdan AR, Miyazawa M, Hashimoto K, Tsuji Y. Siderophores in Iron
Metabolism: From Mechanism to Therapy Potential. 2016.
23. Khasheii B, Mahmoodi P, Mohammadzadeh A. Siderophores: Importance in bacterial
pathogenesis and applications in medicine and industry. *Microbiol Res* 2021; **250**:
126790.
24. Bieler S, Silva F, Soto C, Belin D. Bactericidal activity of both secreted and nonsecreted
microcin E492 requires the mannose permease. *J Bacteriol* 2006; **188**: 7049–7061.
25. Destoumieux-Garzón D, Thomas X, Santamaria M, Goulard C, Barthélémy M, Boscher
B, et al. Microcin E492 antibacterial activity: evidence for a TonB-dependent inner
membrane permeabilization on Escherichia coli. *Mol Microbiol* 2003; **49**: 1031–1041.
26. Rodríguez E, Laviña M. The Proton Channel Is the Minimal Structure of ATP Synthase
Necessary and Sufficient for Microcin H47 Antibiotic Action. *Antimicrob Agents*
Chemother 2003; **47**: 181–187.
27. Palmer JD, Piattelli E, McCormick BA, Silby MW, Brigham CJ, Bucci V. Engineered
Probiotic for the Inhibition of Salmonella via Tetrathionate-Induced Production of
Microcin H47. *ACS Infect Dis* 2018; **4**: 39–45.
28. Page MGP. The Role of Iron and Siderophores in Infection, and the Development of
Siderophore Antibiotics. *Clinical Infectious Diseases* 2019; **69**: S529–S537.
29. Negash KH, Norris JKS, Hodgkinson JT. Siderophore–Antibiotic Conjugate Design: New
Drugs for Bad Bugs? *Molecules* 2019; **24**.
30. Rayner B, Verderosa AD, Ferro V, Blaskovich MAT. Siderophore conjugates to combat
antibiotic-resistant bacteria. *RSC Med Chem* 2023; **14**: 800–822.
31. Boratyn GM, Schäffer AA, Agarwala R, Altschul SF, Lipman DJ, Madden TL. Domain
enhanced lookup time accelerated BLAST. *Biol Direct* 2012; **7**: 12.
32. O'Leary NA, Wright MW, Brister JR, Ciufo S, Haddad D, McVeigh R, et al. Reference
sequence (RefSeq) database at NCBI: current status, taxonomic expansion, and functional
annotation. *Nucleic Acids Res* 2016; **44**: D733–D745.
33. Letunic I, Khedkar S, Bork P. SMART: Recent updates, new developments and status in
2020. *Nucleic Acids Res* 2021; **49**: D458–D460.
34. Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high
throughput. *Nucleic Acids Res* 2004; **32**: 1792–1797.
35. Tamura K, Stecher G, Kumar S. MEGA11: Molecular Evolutionary Genetics Analysis
Version 11. *Mol Biol Evol* 2021; **38**: 3022–3027.
36. Blin K, Shaw S, Augustijn HE, Reitz ZL, Biermann F, Alanjary M, et al. antiSMASH 7.0:
new and improved predictions for detection, regulation, chemical structures and
visualisation. *Nucleic Acids Res* 2023; **51**: W46–W50.

37.   Gibson DG, Young L, Chuang RY, Venter JC, Hutchison CA, Smith HO. Enzymatic
442   assembly of DNA molecules up to several hundred kilobases. *Nature Methods 2009 6:5*
443   2009; **6**: 343–345.
444   38.   Paterson S, Vogwill T, Buckling A, Benmayor R, Spiers AJ, Thomson NR, et al.
445   Antagonistic coevolution accelerates molecular evolution. *Nature 2010 464:7286* 2010;
446   **464**: 275–278.
447   39.   Lynch M, Ackerman MS, Gout JF, Long H, Sung W, Thomas WK, et al. Genetic drift,
448   selection and the evolution of the mutation rate. *Nat Rev Genet* . 2016. Nature Publishing
449   Group. , **17**: 704–714
450   40.   Denman S, Brady C, Kirk S, Cleenwerck I, Venter S, Coutinho T, et al. Brenneria
451   goodwinii sp. nov., associated with acute oak decline in the UK. *Int J Syst Evol Microbiol*
452   2012; **62**: 2451–2456.
453   41.   Brady C, Denman S, Kirk S, Venter S, Rodríguez-Palenzuela P, Coutinho T. Description
454   of Gibbsiella quercinecans gen. nov., sp. nov., associated with Acute Oak Decline. *Syst*
455   *Appl Microbiol* 2010; **33**: 444–450.
456   42.   Allahverdipour T, Shahryari F, Charkhabi NF. First report of walnut bacterial canker
457   caused by Gibbsiella quercinecans and Brenneria roseae subsp. roseae in Iran. *New Dis*
458   *Rep* 2020; **41**: 12–12.
459   43.   Denman S, Doonan J, Ransom-Jones E, Broberg M, Plummer S, Kirk S, et al.
460   Microbiome and infectivity studies reveal complex polyspecies tree disease in Acute Oak
461   Decline. *The ISME Journal 2018 12:2* 2017; **12**: 386–399.
462   44.   Brady C, Orsi M, Doonan JM, Denman S, Arnold D. Brenneria goodwinii growth in vitro
463   is improved by competitive interactions with other bacterial species associated with Acute
464   Oak Decline. *Curr Res Microb Sci* 2022; **3**: 100102.
465   45.   McDonald JE, McDonald J, Koskella B, Julian Marchesi P, Doonan JM, Broberg M, et al.
466   Host–microbiota–insect interactions drive emergent virulence in a complex tree disease.
467   *Proceedings of the Royal Society B* 2020; **287**.
468   46.   Brady C, Hunter G, Kirk S, Arnold D, Denman S. Gibbsiella greigii sp. nov., a novel
469   species associated with oak decline in the USA. *Syst Appl Microbiol* 2014; **37**: 417–422.
470   47.   Hajialigol M, Falahi Charkhabi N, Shahryari F, Sarikhani S. Association of Rahnella
471   victoriana, Enterobacter hormaechei subsp. hoffmannii and Citrobacter braakii with
472   walnut decline. *Scientific Reports 2023 13:1* 2023; **13**: 1–14.
473   48.   Moradi-Amirabad Y, Khodakaramian G. Brenneria alni, causal agent bark canker of
474   Alnus subcordata. *Journal of Phytopathology* 2020; **168**: 516–523.
475   49.   Poret-Peterson AT, McClean AE, Chen L, Kluepfel DA. Complete Genome Sequences of
476   Brenneria rubrifaciens Strain 6D370 and Brenneria nigrifluens Strain ATCC 13028,
477   Causative Agents of Bark Cankers in Walnut. *Microbiol Resour Announc* 2019; **8**.
478   50.   World Health Organization. Antimicrobial resistance. Global report on surveillance.
479   *World Health Organization* . 2014.
480   51.   World Health Organization. Global priority list of antibiotic-resistant bacteria to guide
481   research, discovery, and development of new antibiotics. *World Health Organization* .
482   2017.
483   52.   Moynié L, Milenkovic S, Mislin GLA, Gasser V, Malloci G, Baco E, et al. The complex
484   of ferric-enterobactin with its transporter from Pseudomonas aeruginosa suggests a two-
485   site model. *Nature Communications 2019 10:1* 2019; **10**: 1–14.

486   53.   Subashchandrabose S, Smith S, DeOrnellas V, Crepin S, Kole M, Zahdeh C, et al.
487   Acinetobacter baumannii Genes Required for Bacterial Survival during Bloodstream
488   Infection. *mSphere* 2016; **1**.
489   54.   Sullivan MJ, Petty NK, Beatson SA. Easyfig: a genome comparison visualizer.
490   *Bioinformatics* 2011; **27**: 1009–1010.
491
492

Figure Legends

Fig 1: Novel class IIb microcins are found in numerous *Enterobacteriaceae* genomes. (A) Sequence alignments of the newly identified microcin and immunity genes with the gene clusters of *Ec* CA46 and *Kp* RYC492 using Easyfig[54]. Antimicrobial (A) and immunity (I) genes in the center are represented by darker and lighter shades, respectively. X=mchX, I=mchI, B=mchB, E=mceE, L=mceL, M=mceM. (B,C) Phylogenetic trees of antimicrobial and corresponding immunity genes using codon-aligned nucleotide sequences with General Time Reversible model with discrete gamma distribution (GTR+G) and the Hasegawa-Kishino-Yano model with discrete gamma distribution (HKY+G), respectively. (D) MUSCLE[34] alignment of the amino acid sequence of the signal peptide sequence as well as the C-terminus of the antimicrobial peptides.

Fig 2: Novel class IIb microcins are effective inhibitors of *Enterobacteriaceae* and Gram-negative ESKAPE pathogens. (A) Heatmap summarizing the inhibitory potential of known and novel class IIb microcins against a library of *Enterobacteriaceae*, *Pseudomonadales*, and Gram-positive bacteria, including multidrug-resistant isolates (red) as determined by static inhibition assays with live producing bacteria. *=activity determined through microcin purification and minimum inhibitory concentration assays[13, 17]. (B) Relative minimum inhibitory concentrations for *Se* G492 against different bacterial species. Note that *Se* G492 is 256-times more potent against *A. baumannii* (BAA 1790) compared to *K. pneumoniae* (BAA 1705). (C) Static inhibition assays comparing *Kp* E492, *Kp* G492, *Ec* H47, *Ec* I47, *Ec* M, and *Se* G492 activity from single colony production against multidrug-resistant *A. baumannii* (BAA 1790) and *P. aeruginosa* (PA14). Note that iron-limited conditions (DP) are required for antimicrobial activity, confirming action of class IIb microcins. L-ara=L-arabinose, DP=2,2-dipyridyl, scale bars: 1 cm.

513 **Table 1: *Blastp* results and closest matches to the known class IIb microcins MccE492, MccG492, MccH47, MccI47, or MccM.** Red color indicates no significant
514 match found.

Microcin Name	Species	Accession no.	Antimicrobial Gene Name	Antimicrobial closest match	Identical/Total Length	E-value	Immunity Gene Name	Immunity closest match	Identical/Total Length	E-value
Bg E492	*Brenneria goodwinii*	CP014137	*Bg* E492A	*Kp* E492A (mceA)	35/106	2.00E-14	*Bg* E492I	*Kp* E492I (mceB)	14/94	6.00E-04
Bg X	*Brenneria goodwinii*	CP014137	*Bg* XA	*Kp* E492A (mceA)	21/98	6.90E-02	*Bg* XI	*Ec* MI (mcmI)	10/78	2.60E+00
Gq W	*Gibbsiella quercinecans*	CP014136	*Gq* WA	*Ec* I47A (mciA)	7/103	8.10E+00	*Gq* WI	*Ec* MI (mcmI)	8/74	3.30E+02
Ko H47	*Klebsiella oxytoxa*	CP033844	*Ko* H47A	*Ec* H47A (mchB)	32/77	1.00E-07	*Ko* H47I	*Ec* H47I (mchI)	19/69	2.00E-14
Ps G492	*Pantoea sp.*	CP034363	*Ps* G492A	*Kp* G492A (mceL)	43/89	6.00E-09	*Ps* G492I	*Kp* G492I (mceM)	37/85	2.00E-20
Ro Z	*Raoultella ornithinolytica*	CP008886	*Ro* ZA	*Ec* MA (mcmA)	21/64	1.00E-03	*Ro* ZI	*Ec* MI (mcmI)	5/74	2.00E+01
Ro H47	*Raoultella ornithinolytica*	CP008886	Ro H47A	*Ec* H47A (mchB)	31/79	2.00E-07	*Ro* H47I	*Ec* H47I (mchI)	21/69	4.00E-14
Ro I47	*Raoultella ornithinolytica*	CP008886	*Ro* I47A	*Ec* I47A (mciA)	37/79	4.00E-26	*Ro* I47I	*Ec* I47I (mciI)	64/135	1.00E-43
Ro X	*Raoultella ornithinolytica*	CP008886	*Ro* XA	*Ec* MA (mcmA)	18/99	4.40E-01	*Ro* XI	*Ec* MI (mcmI)	9/61	1.00E-02
Sf H47	*Serratia fonticola*	CP033055	*Sf* H47A	*Ec* H47A (mchB)	38/87	8.00E-11	*Sf* H47I	*Ec* H47I (mchI)	48/68	4.00E-22
Se G492	*Salmonella enterica*	CP030220	*Se* G492A	*Kp* G492A (mceL)	56/85	1.00E-11	*Se* G492I	*Kp* G492I (mceM)	38/84	1.00E-23
Se M	*Salmonella enterica*	CP030220	*Se* MA	*Ec* MA (mcmA)	34/90	1.00E-17	*Se* MI	*Ec* MI (mcmI)	38/71	6.00E-04