Expanding the toolbox: Novel class IIb microcins show activity against gram-negative ESKAPE

and plant pathogens

Benedikt M. Mortzfeld[1,2,*], Shakti K. Bhattarai[1,2], Vanni Bucci[1,2,3]

[1]Department of Microbiology and Physiological Systems, University of Massachusetts Medical School,

Worcester, MA, USA

[2]Program in Microbiome Dynamics, University of Massachusetts Medical School, Worcester, MA, USA,

[3]Program in Systems Biology, University of Massachusetts Medical School, Worcester, MA, USA

[*]Corresponding author

Short title: Novel class IIb microcins show activity against pathogens

Correspondence to:

Benedikt M. Mortzfeld benedikt.mortzfeld@umassmed.edu

Abstract

Interspecies interactions involving direct competition via bacteriocin production play a vital role in shaping ecological dynamics within microbial ecosystems. For instance, the ribosomally-produced siderophore bacteriocins, known as class IIb microcins, have been observed to affect the colonization of pathogenic *Enterobacteriaceae* species within hosts. Notably, to date, only five of these antimicrobials have been identified and all were derived from specific *Escherichia coli* and *Klebsiella pneumoniae* strains. We hypothesized that class IIb microcin production extends beyond these specific compounds and organisms. By employing an informatics-driven approach, screening bacterial genomes in publicly accessible databases, we have discovered a total of twelve previously unknown class IIb microcins. Our investigation unveiled that these microcins are harbored within a diverse array of *Enterobacteriaceae* species, encompassing phytopathogens and environmental isolates. We introduce three novel clades of microcins (MccW, MccX, and MccZ), while also identifying eight new variants of the five previously known ones. To validate their antimicrobial potential, we heterologously expressed these microcins, along with their immunity peptides, in *E. coli* and unequivocally demonstrated their efficacy against a variety of bacterial isolates, including plant pathogens like *Gibbsiella* species *and Rahnella victoriana*. Remarkably, two of these newly discovered class IIb microcins exhibit activity against gram-negative ESKAPE pathogens, such as *Acinetobacter baumannii* or *Pseudomonas aeruginosa* providing the first evidence that class IIb microcins can target bacteria outside of the *Enterobacteriaceae* family. Our findings hold significant promise for the development of innovative live biotherapeutic products tailored to combat these resilient bacteria and underscore the notion that class IIb microcins are more prevalent and more broad-spectrum in the natural microbial world than previously recognized.

Introduction

A large body of theoretical and experimental work has shown that dynamics of microbiomes are shaped by the network of interbacterial interactions(1–4). These cooperative and competitive interactions are often achieved *via* the secretion of cross-feeding metabolites(5), antimicrobial peptides(6) and bacterially produced small molecules(7,8) and are crucial to ecological properties including stability and ability to respond to external perturbations(9,10). Among the competitive interactions, bacteriocin production is proposed to be a prominent mediator of microbiomes dynamics(11) and, specifically, several reports including ours have shown that a bacteriocins subclass, Class IIb microcins, mediates *Enterobacteriaceae* dynamics *in vivo*(12,13).

Class IIb microcins are ribosomally synthesized bacteriocins between 5 kDa to 10 kDa in size

with activity against closely related strains or species(13–17). Unlike all other microcins, they carry

a serine-rich C-terminal motif for a posttranslational modification with a siderophore, here an

enterobactin or an enterobactin derivative, before they are secreted into the extracellular space(18).

Siderophores are iron-chelating molecules commonly employed by various bacteria to scavenge free

iron to outgrow other bacteria, particularly in resource-scarce environments such as the

gastrointestinal tract(15,16,19) and are often associated with increased pathogenicity or

virulence(20–22). The iron chelating moiety of these siderophore antimicrobial peptides (sAMPs) is

recognized by high-affinity receptors and functions as a 'Trojan Horse' key to susceptible bacteria as

it triggers import into the periplasmic space, where the peptide inhibits the molecular target of

susceptible bacteria(19,23–25). Because of these features, delivery of class IIb microcins by wildtype

and engineered probiotics has been recently proposed as a strategy to combat drug-resistant enteric

bacteria(12,13,16,26).

To date only five class IIb microcins have been described and only four have been

characterized in terms of their antimicrobial activity. Specifically, the sAMPs MccE492 and MccG492

(uncharacterized) are solely present in *Klebsiella pneumoniae* (*Kp*), whereas MccH47 is specific for

Escherichia coli (Ec)(15). Additionally, truncated versions of *mciA* (MccI47) and *mcmM* (MccM) are

present in *Kp* RYC492, whereas they are intact in the *Ec* CA46 genome(15). Interestingly, while the

genes encoding for microcins posttranslational modifications are highly conserved between *Ec* and

Kp, suggesting a conserved pathway for microcin maturation, the toxin and corresponding immunity

genes are significantly more variable (Figure S1). We hypothesized that class IIb microcin production

extends beyond these specific compounds and organisms and identified a total of twelve novel class

IIb microcins in seven additional *Enterobacteriaceae* species. In inhibition assays utilizing

heterologous expression of these compounds, we demonstrate potent activity against a library of

bacteria, including gram-negative ESKAPE and plant pathogens, .proving that class IIb microcins are

more prevalent and more broad-spectrum in the natural microbial world than previously recognized.

76   Results

With the hypothesis that class IIb microcin production is a common trait among *Enterobacteriaceae*,

we anticipated that the genes encoding for the antimicrobial and immunity would exhibit a high

degree of dissimilarity to already known peptides as target specificity may result in accelerated

adaptive coevolution(27). Therefore, in addition to the known microcin and immunity genes from

MccE492, MccG492, MccH47, MccI47, and MccM, in our informatic approach we included the genes

that are necessary for mature class IIb microcin biosynthesis, extending our search to longer
sequences for more reliable Basic Local Alignment Search Tool (BLAST) results(28). Moreover, we
hypothesized that the amino acid sequences of genes responsible for posttranslational modification
and microcin export would be less prone to evolutionary changes, thereby maintaining the functional
integrity of the gene cluster(15,29). We then assessed their proximity in the respective genome
location, because microcin genes are typically flanked by genes essential for toxin maturation(15).
Further, we assessed small open reading frames upstream and downstream of the maturation genes,
allowing us to identify novel class IIb microcins without significant sequence similarity to the known
antimicrobials, enabling the discovery of compounds with new molecular targets or modes of action
(see Methods).

Our informatics-driven analysis identified twelve promising class IIb microcin candidates
from seven gene clusters with high similarity to *Ec* CA46 and *Kp* RYC492 in seven species across the
Enterobacteriaceae family (**Figure 1A, Figure S2**): (i) *Brenneria goodwinii* (*Bg*; 2; GenBank:
CP014137), (ii) *Gibbsiella quercinecans* (*Gq*; 1; CP014136), (iii) *Klebsiella oxytoca* (*Ko*; 1; CP033844),
(iv) *Pantoea sp*. (*Ps*; 1; CP034363), (v) *Raoultella ornithinolytica* (*Ro*; 4; CP008886), (vi) *Salmonella*
enterica (*Se*; 2; CP030220), (vii) *Serratia fonticola* (*Sf*; 1; CP033055). Remarkably, although it has
traditionally been a defining characteristic of class IIb microcins that all required genes are encoded
within the chromosome(12), the gene cluster we discovered for *Se* is situated on a 159 kbp plasmid.
Phylogenetic sequence analysis of both the antimicrobial and immunity peptide genes revealed the
presence of eight different clades represented in both trees, respectively (**Figure 1B,C**). Regarding
the well-established class IIb microcins MccH47, MccI47, MccM, MccG492, and MccE492, we
identified novel members for each group, supported by sequence similarity, amino acid identity, the
closest BLAST match, and domain predictions (**Table 1, Figure S3**). In light of this discovery, we
propose a new nomenclature for class IIb microcins that includes the species initials in which they
were identified (*e.g*., *Ec*, *Kp*), the closest relative already characterized class IIb microcin (G492, E492,
H47, I47 or M), as well as the identifiers 'A' for antimicrobial or 'I' for immunity gene.

Figure 1: Novel class IIb microcins are found in numerous *Enterobacteriaceae* genomes. (A) Sequence alignments of the newly identified microcin and immunity genes with the gene clusters of *Ec* CA46 and *Kp* RYC492 using Easyfig(30). Antimicrobial (A) and immunity (I) genes in the center are represented by darker and lighter shades, respectively. X=mchX, I=mchI, B=mchB, E=mceE, L=mceL, M=mceM. (B,C) Phylogenetic trees of antimicrobial and corresponding immunity genes using codon-aligned nucleotide sequences with General Time Reversible model with discrete gamma distribution (GTR+G) and the Hasegawa-Kishino-Yano model with discrete gamma distribution (HKY+G), respectively. (D) MUSCLE(31) alignment of the amino acid sequence of the signal peptide sequence as well as the C-terminus of the antimicrobial peptides.

Thus, the novel G492 relative found in *Salmonella enterica* will be called *Se* G492 with the
antimicrobial peptide identified as *Se* G492A and the immunity peptide identified as *Se* G492I. It is
worth highlighting that in the case of the G492 group, all its members have the immunity gene located
downstream of the antimicrobial gene, whereas for the other clades, this arrangement is reversed. In
addition to uncovering eight novel variants of the five previously characterized microcins, we have
identified four additional microcins that seem to belong to three entirely new clades. We name these
microcins microcin W (MccW), microcin X (MccX), and microcin Z (MccZ). The two members of the
microcin X group, found in *B. goodwinii* (*Bg* X) and *R. ornithinolytica* (Ro X), only show significant
similarity between one another, but not to any of the other antimicrobial or immunity peptides
(**Figure 1 B,C**, **Table 1, Table S1**). Similarly, MccW from *Gibbsiella quercinecans* (*Gq* W) does not
show any sequence similarity to either the known or novel antimicrobial or immunity peptides in
terms of sequence similarity, amino acid identity, the respective BLAST hits (**Table 1, Table S1**), or
phylogenetic localization (**Figure 1 B,C**). Lastly, MccZ from *R. ornithinolytica* (Ro Z) shows
insignificant similarity with *Ec* MA (*mcmA*) for the antimicrobial, whereas the immunity peptide does
not have any match among the known or the novel microcins (**Table 1, Table S1**). Crucially, the
identification of MccX, and MccZ within the same gene clusters as representatives of the E492 (*Bg*
E492), H47 (*Ro* H47), and I47 (*Ro* I47) groups strongly implies that they are functional components
of a gene cluster.

To test the newly identified microcins for antimicrobial activity in our previously established
Ec overexpression system(13,16), all antimicrobial and immunity peptides were codon optimized,
synthesized, and cloned into an inducible high copy vector (see Methods). Through static plate
inhibition assays involving live-producing cells(13,16,26), we successfully validated the robust
antimicrobial activity of eleven out of the twelve newly discovered microcins (Figure 2A). These
microcins exhibit a range of specificities, with some inhibiting targets narrowly (*e.g., Ps* G492AI),
while others exert a broader effect against multiple bacteria (*e.g., Se* G492AI). Notably, our study also
provides the first evidence of inhibitory activity by *Kp* G492, a microcin whose existence and function
have only been proposed in the scientific literature based on genetic sequence(15).

Figure 2: Novel sAMPs are effective inhibitors of *Enterobacteriaceae* and gram-negative ESKAPE pathogens. (A) Heatmap summarizing the inhibitory potential of known and novel class IIb microcins against a library of *Enterobacteriaceae*, *Pseudomonadales*, and gram-positive bacteria, including multidrug-resistant isolates (red) as determined by static inhibition assays with live producing bacteria. *=activity determined through microcin purification and minimum inhibitory concentration assays(13,16). (B) Static inhibition assays comparing *Kp* E492, *Kp* G492, *Ec* H47, *Ec* I47, *Ec* M, and *Se* G492 activity from single colony production against multidrug-resistant *A. baumannii* (BAA 1790) and *P. aeruginosa* (PA14). Note that iron-limited conditions (DP) are required for antimicrobial activity, confirming action of class IIb microcins. L-ara=L-arabinose, DP=2,2-dipyridyl, scale bars: 1 cm.

To this date class IIb microcins have been only shown to be very selective and only active
against different species within the *Enterobacteriaceae* family(13,15,16,26). While the activity for the
novel sAMPs varies, we here report, for the first time, antimicrobial activity outside of the
Enterobacteriaceae family. Specifically, we found that microcins *Ps* G492 and *Se* G492 have activity
against gram-negative multidrug-resistant ESKAPE pathogens with both being capable of inhibiting
Acinetobacter baumannii (BAA 1790), and with microcin *Se* G492 alone also showing activity against
Pseudomonas aeruginosa (PA14) (**Figure 2**).

Discussion
With a comprehensive analysis of publicly available bacterial genomes, we unraveled twelve
previously undiscovered class IIb microcins. Among these findings, we identified three novel
microcin clades, specifically MccW, MccX, and MccZ. Through heterologous expression, we showed
antimicrobial activity for all but one novel microcins and are the first to demonstrate activity for the
known class IIb microcin *Kp* G492. Hence, this research demonstrates that class IIb microcins exhibit
a higher prevalence in natural settings than previously anticipated. As a result, their impact on
ecological community dynamics in natural environments, including the growth of *Pseudomonadales*
species, is likely broader than previously thought. Antimicrobial and immunity genes were
overexpressed and tested in our *Ec*-derived expression system, which might have led to degrees of
incompatibility between the antimicrobial peptides and genes for microcin maturation.
Consequently, the process of posttranslational modification with the siderophore (*i.e.*,
monoglycosylated enterobactin) might be less efficient, and the export mediated by the native signal
peptide less targeted, potentially leading to reduced observed activity in the static plate assays.
Furthermore, it is important to note that these assays exhibit lower sensitivity compared to
purification approaches with minimum inhibitory concentration (MIC) assays. Thus, the activity
spectrums are likely to encompass a wider range than what has been described in this study.
However, historically the microcin literature proves that ideal approaches for purification and MIC
testing can vary between the antimicrobials(13,15,16,24).

Remarkably, we were able to expand the origins of class IIb microcins from the enteric
bacteria *Ec* and *Kp* to other members of the *Enterobacteriaceae* family, including well-known
phytopathogens(32–34). Specifically, *B. goodwinii* and *G. quercinecans* are associated with Acute Oak
Decline (AOD) and are frequently isolated together(35). Similarly, the two strains containing
microcin genes were isolated within the same research project. Notably, these bacteria grow
synergistically(36), while upregulating iron transporters during co-culture(37), hinting at class IIb

microcin-related competition. We were able to show activity of the overexpressed microcins against
human-derived enteric isolates, however, their antimicrobial spectrum might have evolved to target
more frequently encountered strains from the genus *Brenneria* or *Gibbsiella*. Further, we
demonstrated activity of several class IIb microcins against the tree pathogens *G. quercinecans, G.*
greigii as well as *Rahnella victoriana*(36,38,39)*. Thus, treatment with potent microcins, purified or
produced in live bacteria, could present a viable option to target bacteria-caused plant diseases.

In health care settings the burden by gram-negative ESKAPE pathogens and multidrug-
resistant *Enterobacteriaceae* weighs heavily on modern medicine and novel antimicrobials are
needed to develop new treatment options(40,41). Antimicrobial peptides and particularly microcins
are promising candidates for selective eradication of enteric pathogens and have been demonstrated
to potently reduce pathogen colonization *in vivo*, when produced by a live probiotic(12,13). Here we
present the most comprehensive library of class IIb microcins created so far, that is suited for
heterologous expression and *in vivo* application for the development of novel live biotherapeutic
products against drug-resistant enteric bacteria and gram-negative ESKAPE pathogens.

In this study, we challenge the prevailing notion that class IIb microcin production is limited
to *Ec* and *Kp*. Through comprehensive genomic analysis of publicly available bacterial genomes,
coupled with heterologous overexpression, we unveiled a set undiscovered class IIb microcins across
Enterobacteriaceae species. Our findings not only expand the known repertoire of class IIb microcins
but also hold significant implications. We demonstrate that these newly identified class IIb microcins
exert remarkable inhibitory effects on ESKAPE pathogen species not affected by any of the current
ones. This discovery underscores their potential as agents against a broader spectrum of pathogens,
including those affecting humans and plants, thus opening new avenues for antimicrobial research
and applications.

211
Materials and Methods
Bioinformatic class IIb microcin identification
We developed a pipeline that by leveraging BLAST(28) enabled us to mine publicly available genome
databases for novel, previously undescribed sAMPs. We included *mchCDEF* and *mcmL* for *Ec* as well
as *mceCDGHIJ* for *Kp* for posttranslational modification and export, expecting more reliable hits for
longer and functionally conserved proteins in close proximity to class IIb microcin and immunity
genes. Thus, we first ran *tblastn*(28) against RefSeq(42), to screen for all genes related to
biosynthesis pathways, known microcin genes, as well as immunity gene sequences exhibiting
homology to the microcin gene clusters found in *Ec* CA46 and *Kp* RYC492. Homology to the microcin

gene clusters were guided by BLAST parameters sseqid (Genome ID) , pident (percentage of identical
positions) along with sstart (start of alignment in Genome) and send (end of alignment position in
Genome). Resulting hits were concatenated by genome ID and assessed for their proximity to one
another in the genome. These gene clusters should, at best, contain all the known genes required for
toxin maturation, including *mchCDEF* and *mcmL*(15,16). In addition to genomic hits to the known
microcins, small ORFs of 50 to 150 amino acids in size close to the biosynthesis genes were annotated
and their domains were predicted using SMART(43). The ORFs were meticulously examined and
assessed against established class IIb microcin criteria: (i) a serine-rich C-terminus culminating in a
final serine, (ii) the presence of fewer than one cysteine residue, (iii) a signal peptide within the initial
15 amino acids ending with GG or GA, and (iv) close proximity (≤200 bp) to an ORF featuring a
predicted transmembrane domain, typically encoding an immunity peptide. The identified genes
were included in the pipeline's input to expand the scope of gene detection. We repeated this process
iteratively through the pipeline until no additional genes were added to the output. Subsequently,
blastp was used to assess microcin similarity shown in Table 1 and Table S1.

Phylogenetic analyses
For the native full length coding sequence of the microcin and immunity genes a codon-based
sequence alignment was generated using the MUSCLE algorithm(31). Then, we determined the best
fit substitution models for maximum likelihood phylogenetic analyses, resulting in the General Time
Reversible model with discrete gamma distribution (GTR+G) and the Hasegawa-Kishino-Yano model
with discrete gamma distribution (HKY+G), respectively. A bootstrap test with 1000 replicates for
maximum likelihood and random seed was conducted for both trees. Alignment, model testing, and
tree building was performed in MEGA11(44).

Plasmids and heterologous class IIb microcin expression
ORFs of identified microcin and immunity genes were codon optimized for frequent *Ec* codon usage
without creating repetitive sequences and synthesized by Integrated DNA Technologies (Coralville,
IA) with 18 bp of native 5' upstream sequence and 20 bp of native 3' downstream sequence,
respectively. Using Gibson Assembly(45), the genes were cloned into our previously established *Ec*
class IIb microcin expression system that results in mature class IIb microcins posttranslationally
modified with a monoglycosylated enterobactin(13,16). Briefly, the antimicrobial and the immunity
genes are co-expressed under the control of an arabinose-inducible pBad/araC promoter in a high

copy plasmid with a pUC-derived origin of replication. All assemblies were verified using whole

plasmid sequencing. DNA files for all used plasmids can be found at:

Static inhibition assays

Cultures of strains with confirmed plasmid assemblies were spread in LB agar plates containing 100

µg/ml ampicillin. Single colonies for each microcin were picked with a sterile pipet tip and inserted

into solid LB agar containing 100 µg/ml ampicillin for plasmid retention, 0.2 mM 2,2-dipyridyl to

create iron limited conditions during the growth phase, and 0.4% L-arabinose for induction of gene

expression. Plates were incubated at 37°C for up to 72 h, before they were overlaid with the target

bacterial isolates. For the overlay, the microcin-producing bacteria in the stabs were inactivated

using chloroform vapors and ten minutes under ultraviolet light. Then, target bacteria were diluted

1:2000 from overnight culture in LB media containing 100 µg/ml ampicillin and 0.2 mM 2,2-

dipyridyl. *Ec* and *S. flexneri* strains were diluted 1:200 to acquire dense bacterial lawns. Finally, 0.5

ml of molten agar was added to the liquid media and the resulting soft agar was spread on the plate

with the inactivated bacteria and incubated for 16 h at 37°C.

Availability of Data and Materials

The genomes are accessible with the following GenBank numbers: *Brenneria goodwinii* (GenBank: CP014137), *Gibbsiella quercinecans* (CP014136), *Klebsiella oxytoca* (CP033844), *Pantoea sp.* (CP034363), R*aoultella ornithinolytica* (CP008886), *Salmonella enterica* (CP030220), *Serratia fonticola* (CP033055). All all information is included in the manuscript or supporting files. All plasmid sequences as well as annotation files to produce Figures 1A, S1, S2, and S3 are available as supplementary material.

Funding Disclosure

This work was supported partially supported by the CDMRP PRMP W81XWH2020013 to V.B., by the NIH NIA 1R01AG075283-01A1 to V.B., and by the Deutsche Forschungsgemeinschaft (DFG) project 457837076 to B.M.M.

Competing Interests

V.B. receives support from a sponsored research agreement from Vedanta Biosciences, Inc.

Acknowledgements

The author would like to thank the administrative staff in the Microbiology and Physiological Systems Department at the University of Massachusetts Chan Medical School, especially Annette Bohigian, Amy Parker, Dhruti Desai, Marie Berardi, Richard Fish, and Tracey Rae, for their support.

Author Contributions

B.M.M, S.K.B, and V.B conceptualized the study. S.K.B. performed bioinformatic data mining. B.M.M and V.B designed the experiments. B.M.M performed plasmid design, verification, and *in vitro* activity testing. B.M.M, S.K.B, and V.B wrote the manuscript.

296 **Table 1: *Blastp* results and closest matches to the known class IIb microcins MccE492, MccG492, MccH47, MccI47, or MccM.** Red color indicates no significant
297 match found.

Microcin Name	Species	Accession no.	Antimicrobial Gene Name	Antimicrobial closest match	Identical/Total Length	E-value	Immunity Gene Name	Immunity closest match	Identical/Total Length	E-value
Bg E492	*Brenneria goodwinii*	CP014137	*Bg* E492A	*Kp* E492A (mceA)	35/106	2.0E-14	*Bg* E492I	*Kp* E492I (mceB)	14/94	6.0E-04
Bg X	*Brenneria goodwinii*	CP014137	*Bg* XA	*Kp* E492A (mceA)	21/98	6.9E-02	*Bg* XI	*Ec* MI (mcmI)	10/78	2.6E+00
Gq W	*Gibbsiella quercinecans*	CP014136	*Gq* WA	*Ec* I47A (mciA)	7/103	8.1E+00	*Gq* WI	*Ec* MI (mcmI)	8/74	3.3E+02
Ko H47	*Klebsiella oxytoxa*	CP033844	*Ko* H47A	*Ec* H47A (mchB)	32/77	1.0E-07	*Ko* H47I	*Ec* H47I (mchI)	19/69	2.0E-14
Ps G492	*Pantoea sp.*	CP034363	*Ps* G492A	*Kp* G492A (mceL)	43/89	6.0E-09	*Ps* G492I	*Kp* G492I (mceM)	37/85	2.0E-20
Ro Z	*Raoultella ornithinolytica*	CP008886	*Ro* ZA	*Ec* MA (mcmA)	21/64	1.0E-03	*Ro* ZI	*Ec* MI (mcmI)	5/74	2.0E+01
Ro H47	*Raoultella ornithinolytica*	CP008886	Ro H47A	*Ec* H47A (mchB)	31/79	2.0E-07	*Ro* H47I	*Ec* H47I (mchI)	21/69	4.0E-14
Ro I47	*Raoultella ornithinolytica*	CP008886	*Ro* I47A	*Ec* I47A (mciA)	37/79	4.0E-26	*Ro* I47I	*Ec* I47I (mciI)	64/135	1.0E-43
Ro X	*Raoultella ornithinolytica*	CP008886	*Ro* XA	*Ec* MA (mcmA)	18/99	4.4E-01	*Ro* XI	*Ec* MI (mcmI)	9/61	1.0E-02
Sf H47	*Serratia fonticola*	CP033055	*Sf* H47A	*Ec* H47A (mchB)	38/87	8.0E-11	*Sf* H47I	*Ec* H47I (mchI)	48/68	4.0E-22
Se G492	*Salmonella enterica*	CP030220	*Se* G492A	*Kp* G492A (mceL)	56/85	1.0E-11	*Se* G492I	*Kp* G492I (mceM)	38/84	1.0E-23
Se M	*Salmonella enterica*	CP030220	*Se* MA	*Ec* MA (mcmA)	34/90	1.0E-17	*Se* MI	*Ec* MI (mcmI)	38/71	6.0E-04

References

1.   Stein RR, Bucci V, Toussaint NC, Buffie CG, Rätsch G, Pamer EG, et al. Ecological
Modeling from Time-Series Inference: Insight into Dynamics and Stability of Intestinal
Microbiota. PLoS Comput Biol [Internet]. 2013 [cited 2023 Sep 12];9(12):e1003388.
Available from:
https://journals.plos.org/ploscompbiol/article?id=10.1371/journal.pcbi.1003388
2.   Bucci V, Xavier JB. Towards Predictive Models of the Human Gut Microbiome. J Mol
Biol. 2014 Nov 25;426(23):3907–16.
3.   Coyte KZ, Schluter J, Foster KR. The ecology of the microbiome: Networks, competition,
and stability. Science (1979) [Internet]. 2015 Nov 6 [cited 2023 Sep 12];350(6261):663–6.
Available from: https://www.science.org/doi/10.1126/science.aad2602
4.   Hromada S, Venturelli OS. Gut microbiota interspecies interactions shape the response of
Clostridioides difficile to clinically relevant antibiotics. PLoS Biol [Internet]. 2023 May 1
[cited 2023 Sep 12];21(5):e3002100. Available from:
https://journals.plos.org/plosbiology/article?id=10.1371/journal.pbio.3002100
5.   Culp EJ, Goodman AL. Cross-feeding in the gut microbiome: Ecology and mechanisms.
Cell Host Microbe. 2023 Apr 12;31(4):485–99.
6.   Heilbronner S, Krismer B, Brötz-Oesterhelt H, Peschel A. The microbiome-shaping roles
of bacteriocins. Nature Reviews Microbiology 2021 19:11 [Internet]. 2021 Jun 1 [cited
2023 Sep 12];19(11):726–39. Available from: https://www.nature.com/articles/s41579-
021-00569-w
7.   Hibbing ME, Fuqua C, Parsek MR, Peterson SB. Bacterial competition: surviving and
thriving in the microbial jungle. Nature Reviews Microbiology 2009 8:1 [Internet]. 2009
Nov 30 [cited 2023 Sep 12];8(1):15–25. Available from:
https://www.nature.com/articles/nrmicro2259
8.   Donia MS, Fischbach MA. Small molecules from the human microbiota. Science (1979)
[Internet]. 2015 Jul 24 [cited 2023 Sep 12];349(6246). Available from:
https://www.science.org/doi/10.1126/science.1254766
9.   Coyte KZ, Rakoff-Nahoum S. Understanding Competition and Cooperation within the
Mammalian Gut Microbiome. Current Biology. 2019 Jun 3;29(11):R538–44.
10. Coyte KZ, Rao C, Rakoff-Nahoum S, Foster KR. Ecological rules for the assembly of
microbiome communities. PLoS Biol [Internet]. 2021 Feb 19 [cited 2023 Sep
12];19(2):e3001116. Available from:
https://journals.plos.org/plosbiology/article?id=10.1371/journal.pbio.3001116
11. Niehus R, Oliveira NM, Li A, Fletcher AG, Foster KR. The evolution of strategy in
bacterial warfare via the regulation of bacteriocins and antibiotics. Elife. 2021 Sep 1;10.
12. Sassone-Corsi M, Nuccio SP, Liu H, Hernandez D, Vu CT, Takahashi AA, et al.
Microcins mediate competition among Enterobacteriaceae in the inflamed gut. Nature.
2016 Dec 31;540(7632):280–3.
13. Mortzfeld BM, Palmer JD, Bhattarai SK, Dupre HL, Mercado-Lubio R, Silby MW, et al.
Microcin MccI47 selectively inhibits enteric bacteria and reduces carbapenem-resistant
Klebsiella pneumoniae colonization in vivo when administered via an engineered live
biotherapeutic. Gut Microbes [Internet]. 2022 Dec 31 [cited 2022 Oct 4];14(1). Available
from: https://www.tandfonline.com/doi/full/10.1080/19490976.2022.2127633

14.   de Lorenzo V. Isolation and characterization of microcin E 492 from Klebsiella
345   pneumoniae. Arch Microbiol. 1984 Sep;139(1):72–5.
346   15.   Vassiliadis G, Destoumieux-Garzón D, Lombard C, Rebuffat S, Peduzzi J. Isolation and
347   characterization of two members of the siderophore-microcin family, microcins M and
348   H47. Antimicrob Agents Chemother. 2010 Jan 1;54(1):288–97.
349   16.   Palmer JD, Mortzfeld BM, Piattelli E, Silby MW, McCormick BA, Bucci V. Microcin
350   H47: A Class IIb Microcin with Potent Activity Against Multidrug Resistant
351   Enterobacteriaceae. ACS Infect Dis. 2020 Apr 10;6(4):672–9.
352   17.   Baquero F, Lanza VF, Baquero MR, del Campo R, Bravo-Vázquez DA. Microcins in
353   Enterobacteriaceae: Peptide Antimicrobials in the Eco-Active Intestinal Chemosphere.
354   Vol. 10, Frontiers in Microbiology. Frontiers Media S.A.; 2019. p. 2261.
355   18.   Azpiroz MF, Rodríguez E, Laviña M. The structure, function, and origin of the microcin
356   H47 ATP-binding cassette exporter indicate its relatedness to that of colicin V.
357   Antimicrob Agents Chemother [Internet]. 2001 [cited 2023 Sep 13];45(3):969–72.
358   Available from: https://journals.asm.org/doi/10.1128/aac.45.3.969-972.2001
359   19.   Patzer SI, Baquero MR, Bravo D, Moreno F, Hantke K. The colicin G, H and X
360   determinants encode microcins M and H47, which might utilize the catecholate
361   siderophore receptors FepA, Cir, Fiu and IroN. Microbiology (N Y). 2003 Sep
362   1;149(9):2557–70.
363   20.   Miethke M, Marahiel MA. Siderophore-Based Iron Acquisition and Pathogen Control.
364   Microbiology and Molecular Biology Reviews [Internet]. 2007 Sep [cited 2023 Sep
365   6];71(3):413–51. Available from: https://journals.asm.org/doi/10.1128/mmbr.00012-07
366   21.   Wilson BR, Bogdan AR, Miyazawa M, Hashimoto K, Tsuji Y. Siderophores in Iron
367   Metabolism: From Mechanism to Therapy Potential. 2016 [cited 2023 Sep 6]; Available
368   from: http://dx.doi.org/10.1016/j.molmed.2016.10.005
369   22.   Khasheii B, Mahmoodi P, Mohammadzadeh A. Siderophores: Importance in bacterial
370   pathogenesis and applications in medicine and industry. Microbiol Res. 2021 Sep
371   1;250:126790.
372   23.   Bieler S, Silva F, Soto C, Belin D. Bactericidal activity of both secreted and nonsecreted
373   microcin E492 requires the mannose permease. J Bacteriol. 2006 Oct 1;188(20):7049–61.
374   24.   Destoumieux-Garzón D, Thomas X, Santamaria M, Goulard C, Barthélémy M, Boscher
375   B, et al. Microcin E492 antibacterial activity: evidence for a TonB-dependent inner
376   membrane permeabilization on Escherichia coli. Mol Microbiol [Internet]. 2003 Aug 1
377   [cited 2023 Sep 4];49(4):1031–41. Available from:
378   https://onlinelibrary.wiley.com/doi/full/10.1046/j.1365-2958.2003.03610.x
379   25.   Rodríguez E, Laviña M. The Proton Channel Is the Minimal Structure of ATP Synthase
380   Necessary and Sufficient for Microcin H47 Antibiotic Action. Antimicrob Agents
381   Chemother. 2003 Jan 1;47(1):181–7.
382   26.   Palmer JD, Piattelli E, McCormick BA, Silby MW, Brigham CJ, Bucci V. Engineered
383   Probiotic for the Inhibition of Salmonella via Tetrathionate-Induced Production of
384   Microcin H47. ACS Infect Dis. 2018 Jan 12;4(1):39–45.
385   27.   Paterson S, Vogwill T, Buckling A, Benmayor R, Spiers AJ, Thomson NR, et al.
386   Antagonistic coevolution accelerates molecular evolution. Nature 2010 464:7286
387   [Internet]. 2010 Feb 24 [cited 2023 Sep 12];464(7286):275–8. Available from:
388   https://www.nature.com/articles/nature08798

389   28.   Boratyn GM, Schäffer AA, Agarwala R, Altschul SF, Lipman DJ, Madden TL. Domain
390   enhanced lookup time accelerated BLAST. Biol Direct [Internet]. 2012 Apr 17 [cited 2022
391   Jun 10];7(1):12. Available from:
392   http://biologydirect.biomedcentral.com/articles/10.1186/1745-6150-7-12

393   29.   Lynch M, Ackerman MS, Gout JF, Long H, Sung W, Thomas WK, et al. Genetic drift,
394   selection and the evolution of the mutation rate. Vol. 17, Nature Reviews Genetics. Nature
395   Publishing Group; 2016. p. 704–14.

396   30.   Sullivan MJ, Petty NK, Beatson SA. Easyfig: a genome comparison visualizer.
397   Bioinformatics. 2011 Apr 1;27(7):1009–10.

398   31.   Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high
399   throughput. Nucleic Acids Res [Internet]. 2004 Mar 1 [cited 2023 Sep 13];32(5):1792–7.
400   Available from: https://dx.doi.org/10.1093/nar/gkh340

401   32.   Denman S, Brady C, Kirk S, Cleenwerck I, Venter S, Coutinho T, et al. Brenneria
402   goodwinii sp. nov., associated with acute oak decline in the UK. Int J Syst Evol Microbiol
403   [Internet]. 2012 Oct 1 [cited 2023 Sep 4];62(10):2451–6. Available from:
404   https://www.microbiologyresearch.org/content/journal/ijsem/10.1099/ijs.0.037879-0

405   33.   Brady C, Denman S, Kirk S, Venter S, Rodríguez-Palenzuela P, Coutinho T. Description
406   of Gibbsiella quercinecans gen. nov., sp. nov., associated with Acute Oak Decline. Syst
407   Appl Microbiol. 2010 Dec 1;33(8):444–50.

408   34.   Allahverdipour T, Shahryari F, Charkhabi NF. First report of walnut bacterial canker
409   caused by Gibbsiella quercinecans and Brenneria roseae subsp. roseae in Iran. New Dis
410   Rep [Internet]. 2020 Jan 1 [cited 2023 Sep 4];41(1):12–12. Available from:
411   https://onlinelibrary.wiley.com/doi/full/10.5197/j.2044-0588.2020.041.012

412   35.   Denman S, Doonan J, Ransom-Jones E, Broberg M, Plummer S, Kirk S, et al.
413   Microbiome and infectivity studies reveal complex polyspecies tree disease in Acute Oak
414   Decline. The ISME Journal 2018 12:2 [Internet]. 2017 Oct 13 [cited 2023 Sep
415   4];12(2):386–99. Available from: https://www.nature.com/articles/ismej2017170

416   36.   Brady C, Orsi M, Doonan JM, Denman S, Arnold D. Brenneria goodwinii growth in vitro
417   is improved by competitive interactions with other bacterial species associated with Acute
418   Oak Decline. Curr Res Microb Sci. 2022 Jan 1;3:100102.

419   37.   McDonald JE, McDonald J, Koskella B, Julian Marchesi P, Doonan JM, Broberg M, et al.
420   Host–microbiota–insect interactions drive emergent virulence in a complex tree disease.
421   Proceedings of the Royal Society B [Internet]. 2020 Aug 26 [cited 2023 Sep 4];287(1933).
422   Available from: https://royalsocietypublishing.org/doi/10.1098/rspb.2020.0956

423   38.   Brady C, Hunter G, Kirk S, Arnold D, Denman S. Gibbsiella greigii sp. nov., a novel
424   species associated with oak decline in the USA. Syst Appl Microbiol. 2014 Sep
425   1;37(6):417–22.

426   39.   Hajialigol M, Falahi Charkhabi N, Shahryari F, Sarikhani S. Association of Rahnella
427   victoriana, Enterobacter hormaechei subsp. hoffmannii and Citrobacter braakii with
428   walnut decline. Scientific Reports 2023 13:1 [Internet]. 2023 Jul 12 [cited 2023 Sep
429   26];13(1):1–14. Available from: https://www.nature.com/articles/s41598-023-38427-9

430   40.   World Health Organization. Antimicrobial resistance. Global report on surveillance.
431   World Health Organization. 2014.

432   41.   World Health Organization. Global priority list of antibiotic-resistant bacteria to guide
433   research, discovery, and development of new antibiotics. World Health Organization.
434   2017.

435   42.   O'Leary NA, Wright MW, Brister JR, Ciufo S, Haddad D, McVeigh R, et al. Reference
436   sequence (RefSeq) database at NCBI: current status, taxonomic expansion, and functional
437   annotation. Nucleic Acids Res [Internet]. 2016 Jan 4 [cited 2023 Sep 13];44(D1):D733–
438   45. Available from: https://dx.doi.org/10.1093/nar/gkv1189
439   43.   Letunic I, Khedkar S, Bork P. SMART: Recent updates, new developments and status in
440   2020. Nucleic Acids Res [Internet]. 2021 Jan 8 [cited 2022 Jun 2];49(D1):D458–60.
441   Available from: https://academic.oup.com/nar/article/49/D1/D458/5940513
442   44.   Tamura K, Stecher G, Kumar S. MEGA11: Molecular Evolutionary Genetics Analysis
443   Version 11. Mol Biol Evol [Internet]. 2021 Jun 25 [cited 2023 Sep 13];38(7):3022–7.
444   Available from: https://dx.doi.org/10.1093/molbev/msab120
445   45.   Gibson DG, Young L, Chuang RY, Venter JC, Hutchison CA, Smith HO. Enzymatic
446   assembly of DNA molecules up to several hundred kilobases. Nature Methods 2009 6:5
447   [Internet]. 2009 Apr 12 [cited 2023 Sep 12];6(5):343–5. Available from:
448   https://www.nature.com/articles/nmeth.1318
449