Non-autonomous cell redox-pairs dictate niche homeostasis in multi-lineage stem populations

Olivia Majhi[1], Aishwarya Chhatre[1,2], Tanvi Chaudhary[1] and Devanjan Sinha[1#]

[1]Department of Zoology, Institute of Science, Banaras Hindu University, Varanasi, Uttar Pradesh, India

[2]Present address: Biological Research Centre, Institute of Genetics, Eötvös Loránd Research Network (ELKH), Szeged, Hungary

[#]To whom correspondence should be addressed:

Address for correspondence:
Dr. Devanjan Sinha,
Department of Zoology,
Institute of Science,
Banaras Hindu University,
Varanasi – 221005,
Uttar Pradesh, India
India
Email: devanjan@bhu.ac.in

Abstract

Reactive oxygen species (ROS), predominantly derived from mitochondrial respiratory complexes, have evolved as key molecules influencing cell fate decisions like maintenance and differentiation. These redox-dependent events are mainly considered to be cell intrinsic in nature, on contrary our observations indicate involvement of these oxygen-derived entities as intercellular communicating agents. In multi-lineage *Drosophila* germline, neighbouring Germline Stem Cells (GSCs) and Cystic Stem Cells (CySCs) maintain differential redox thresholds where CySCs by virtue of their higher redox-state regulate physiological ROS levels of germline. Disruption of the intercellular redox equilibrium between the two adjoining stem cell populations results in deregulated niche architecture and loss of GSCs, which was mainly attributed to loss of contact-based receptions and uncontrolled CySC proliferation due to ROS-mediated activation of self-renewing signals. Our observations hint towards the crucial role of intercellular redox gradients originating from somatic progenitors, CySCs in niche stability where they function not only as a source of their own maintenance cues but also serve as non-autonomous redox moderators of germline immortality. Our findings underscore the complexity of niche homeostasis and predicate the importance of intercellular redox communication in understanding stem cell microenvironments.

Introduction

Studies from the past few decades have shown the apparent role of ROS in influencing various biological processes[1-3]. Generally, the sources of ROS are compartmentalized in co-occurrence with their targets mediating important signalling events[4]. Mitochondria, a major source of oxidant species[5,6], localize dynamically towards nucleus, effecting oxidation induced reshaping of gene expression profiles[7]. Localized ROS production are contributed by NADH oxidases distributed across different cellular regions[8]. Intracellular hydrogen peroxide gradients maintained by thioredoxin system allow intercompartmental exchanges among endoplasmic reticulum (ER), mitochondria and peroxisomes[9-11].

The process of redox relays is conserved and plays a fundamental role in self-renewal and differentiation of stem cell populations[12]. Stem cells whether embryonic or adult maintain a low redox profile, characterized by subdued mitochondrial respiration[13-17]. Levels of ROS are tightly regulated and elevated amounts promote early differentiation and atypical stem cell behaviour[18]. However, leading evidences suggest that many transcription factors require oxidative environment for maintenance of pluripotent states. For instance, physiological ROS levels play a crucial role in genome maintenance of embryonic stem cells[13]. Multipotent hematopoietic progenitors and neural stem cells require relatively high baseline redox for their maintenance[14,16,19]. Suppression of Nox system or mitochondrial ROS compromises the maintenance of these self-renewing populations and promote their differentiation or death[18]. These evidences point towards essentiality of a perfectly tuned redox state for balancing the pluripotent and differentiated states. But how the stem cells harmonize their redox potential by possessing a restrained oxidant system is not very clear.

We addressed this fundamental question in multi-stem cell lineage-based niche architecture in *Drosophila* testis. The testicular stem cell niche is composed of a central cluster of somatic cells called the hub which contacts eight to eleven GSCs arranged in a round array[20,21]. A pair of cystic stem cells (CySCs) originating the somatic lineage cyst cells, enclose each GSC and make their independent connections with the hub and GSCs via adherens junctions[22-26]. The hub cells secrete Unpaired family of cytokines that defines niche occupancy and self-renewal of both GSCs and CySCs through Jak-Stat signalling[27,28]. CySCs also repress GSC

differentiation by suppressing transcription of bag-of-marbles (Bam)[29,30]. Concerted

asymmetric division of both GSC and CySC is essential for proper cyst formation[31]. A

developing cyst contains a dividing and differentiating gonialblast encircled by cyst

cells that provide nourishment to developing spermatids[32,33]. However, the

maintenance of GSCs in the spatially controlled microenvironment is still not very

clear.

We found the existence of a perfectly balanced redox differential between GSC and

CySC essential for niche homeostasis. CySCs by virtue of their higher redox threshold

and clustering of mitochondria at GSC-CySC interface, generate an intercellular redox

gradient that maintain physiological ROS levels in GSCs. Alterations in CySC redox

state affected self-renewal and differentiation propensities of the GSCs. Intercellular

redox-imbalance induced niche disequilibrium through precocious differentiation of

GSCs and deregulated CySC proliferation, due to activation of pro-proliferative

transductions and loss of cell-cell contact signalling. Our results in multi-stem cell

lineage-based germline system indicate a nuanced division of labour where a higher

redox state in CySCs not only maintains their self-renewing processes but also non-

autonomously foster GSCs maintenance.

Results

Redox state of GSC is maintained by asymmetric distribution of CySC
mitochondria

The apical region of the Drosophila testis incorporates a cluster of differentiated cells

the hub surrounded by GSCs in a rosette arrangement with each GSC being enclosed

by two CySCs (Fig. 1A). The mitochondria morphology and distribution in this cluster

was checked by immunostaining ATP5A subunit of mitochondrial ATPase. We

observed fragmented mitochondria in Vasa[+] GSCs as compared to a fused

organization in Traffic jam (Tj)[+] CySCs (Fig. 1B#, C, D). Distribution profiles of

mitochondria showed its higher propensity to overlap with Tj, indicating CySCs to be

a mitochondria dense region compared to GSC (Fig. 1F); and its mitochondria to be

predominantly clustered near the GSC-CySC boundary, illustrated by intensity profile

(Fig. S1A''') and X-Z and Y-Z sectioning (Fig. S1B-B'). The ATP5A labelling

overlapped with *TFAM-GFP*, a mitochondrial transcription factor[34], further confirming

the patterning observed between these two stem cell populations (Fig. S1C').

Mitochondria localization near the GSC-CySC boundary, together with its dispersion

pattern was in accordance with the intensity variance of ROS reporter line *gstD1-GFP*

[35] at the niche (Fig. 1H and S1D-D'). The redox gradient intensifies at the Tj-Vasa

boundary and then gradually diffuses into Vasa$^+$ GSCs, suggesting an establishment

of GSC redox state through CySCs (Fig. 1Q; *left panel*). A GSC intrinsic oxidative

environment should have resulted in a constant continuum of *gstD1*-GFP intensity

across the Vasa-stained structure. We confirmed the greater ROS gradient at the

GSC-CySC interface by 2D-Plot profile of Vasa, Tj and *gstD1*-GFP across the different

cell types (Fig. S1E).

To affirm the role of CySC in influencing GSC redox state, we asked if disrupting

dismutases in CySCs would influence the redox state of GSCs. We depleted the

expression of superoxide dismutase 1 (Sod1) in CySCs and early differentiating cyst

cells using *Tj-Gal4* driver (*Tj>Sod1i*). Higher CySC redox state resulted in

corresponding increase in GSC ROS profile, as indicated by a greater degree gstD1-

GFP intensity and DHE fluorescence at the Vasa-gstD1GFP (V∩G) intersection (Fig.

1K, P, R and S1F), suggesting non-autonomous maintenance of GSC redox gradient

by CySCs (Fig. 1Q; *right panel*). The surge in GSC ROS levels upon lowering of

antioxidant defences in CySCs, was additionally confirmed through enhanced

Vasa:*gstD1*-GFP colocalization intensities in *Sod1i* populations (MCC^{mean} = 0.482)

with respect of controls (MCC^{mean} = 0.276) (Fig. 1S). These observations hinted

towards a potential role of CySCs and its GSC polarized mitochondria in elevating the

cellular ROS gradient at the interface between two stem cell populations.

Balanced CySC redox profile is crucial for its proliferation and GSC

maintenance

Along with the shift in the redox gradient, we observed that high ROS in *Tj>Sod1i* had

a striking effect on the increase in DAPI$^+$-nuclei at the tip of testis (Fig. 1G and L). This

overcrowding was attributed to an increase in number and positional shift in Tj$^+$ CySCs

and early differentiating cyst cells (Fig. 2A', B' and E). The enhancement in the number

of CySCs was also reflected by ~2-fold change in Zfh1$^+$ cells (Fig. 2F). Vasa$^+$ cells

flanking the hub however, substantially reduced in number (Fig. 2C''-D'' and G), which

was validated through lower detectable levels of Vasa even when its expression per

cell remained unchanged (Fig. S2O). The observations were further verified using

Sod1i localized in different chromosome to avoid any chromosome or balancer-based

biasness (Fig. S1G-L). However, the phenotypic effect of higher ROS was limited to

its origin in CySCs only because ablation of GSC redox status did not cause any

marked change in niche composition. *Nos-Gal4* driven knock-down of Sod1 in GSCs

did not result in significant change in CySC number (Fig. S2A', B' and E) but effected

a considerable reduction on Vasa$^+$ cells (Fig. S2C', D' and F), aligning with previous

observations[17]. Although, Sod1 is the predominant enzyme which also localizes in the

intermembrane space of mitochondria, we observed a similar result upon depleting

the matrix localized Sod2 in both GSCs and CySCs (Fig. S2G-M), indicating

involvement of mitochondrial ROS for the observed cellular phenotypes.

To confirm active proliferation of cells and rule out the possibility of arrested growth,

we utilized fly- Fluorescent Ubiquitination-based Cell Cycle Indicator (FUCCI)

system[36] which comprises of two reporter constructs marking G1/S transition (green),

S (red) and G2/early mitosis (yellow) (Fig. 2H). *Tj>Sod1i* testes presented a ~2-fold

increase in cell number across all stages of cell-cycle (Fig 2I). The enhanced number

of dividing cells was contributed mainly by progressive division of CySCs, showing >2-

fold difference in accumulation of Zfh1$^+$ nuclei in S and G2/M phases (Fig. 2J, S2PV-

VII and QV-VII). The number of Zfh1 labelled cells in G1/S transition was also

substantially more (Fig. S2PIII and QIII), indicating dysregulated redox ensued

increased mitotic index of CySCs. Downregulation of Sod1 in GSCs through *Nos-Gal4*

demonstrated no significant change in G1/S, S and G2/M phase numbers, confirming

our previous observation that altering ROS levels in GSCs does not have any non-

autonomous effect on CySCs (Fig. S2N).

CySC induced ROS couples precocious differentiation of stem cell lineages

Since, in *Sod1i* testis the number of Tj$^+$ CySCs and early differentiating cells[37] were

elevated and more than CySC-specific Zfh1$^+$ cells (Fig. 2I and J), we sought to label

these cells with another differentiation marker Eya (Eyes absent). We found that Eya$^+$

cells clustered towards the hub along with a significant increase in their number in

Sod1 depleted CySCs (Fig. 3A, B and D). The supposed premature expression of Eya

resulted in a population of differentiating cyst precursor cells co-expressing Eya-Zfh1

(Fig. 3A'', B'' and D). This population was found to be deviating from the normal

partitioning of wild type representative populations (Fig. 3C). On contrary, the same

cells would not show a similar phenotypical arrangement when Sod1i depletion was

driven in the germline lineage by *Nos-Gal4* (Fig. S3A-D). These observations hint

towards an obvious change in redox state of GSCs affecting their number, even when

generated non-autonomously from CySCs.

Together with reduction in Vasa$^+$ GSCs shown earlier (Fig. 2G), we also detected

gonialblast expressing differentiation-promoting factor Bam to be much closer to the

hub in *Tj>Sod1i* testes (Fig. 3E'' and F''). The mean distance between the hub and

Bam$^+$ cells was found to be shortened by ~7 microns (Fig 3G). The relative

advancement of spermatogonial differentiation towards the hub was also ascertained

by tracking the transformation of GSC-specific spectrosome into branched fusomes

connecting the multi-cell gonialblast (Fig. 1A). The branching fusomes were found

more proximal to hub than controls (Fig. 3H, I'' and J''), along with parallel reduction

in number of spectrosomes, implying a decline in early-stage germ cells (Fig. S3F).

This loss could possibly be attributed to a higher oxidizing environment in GSCs and

loss of intercellular contacts, particularly with hub cells (Fig. 2C# and D#). The

observation corroborated with decreased expression pattern of E-cadherin in *Tj>Sod1i*

particularly near the hub (Fig. 3K-M and S3G) and could be one of the reasons for

GSCs dissociation and differentiation (Fig. 2D#), due to absence of self-renewing

signals. Indeed, disengagement of GSCs from the hub caused reduction in Socs36E

and Ptp61F transcripts that act as drivers of E-cadherin expression and are induced

by maintenance factor STAT originating from the hub (Fig. 3N and O). These results

suggest that disruption of intercellular redox gradient effected premature differentiation

of stem cells in the niche.

Disrupted niche architecture compromises GSC-CySC communication

GSCs and CySCs intercommunicate through redox dependent factors such as EGFR,

PI3K/Tor to support CySC maintenance[38]. EGF ligands secreted by GSCs maintain

differences in cell polarity and segregates self-renewing from differentiating

populations[39]. *Tj>Sod1i* cells showed lower levels of cell-polarity marker Dlg (Fig. 4A",

B" and S4A), that expanded more into the differentiated zone (Fig. S4B). Since, Dlg

expression is dependent on EGFR[40], impairment in its reception was reflected in net

reduction in pErk levels across sections in *Sod1i* testis (Fig. S4C). Due to loss of

contact, this reduction was mainly limited to regions surrounding the hub, in CySC

zone (Fig. 4C"-D" and E). Differentiated cyst cells presented relatively higher levels of

pErk expression (Fig. 4E, S4D'), particularly in *Tj>Sod1i* tissues where the pErk area

was significantly extended, mapping with that of Dlg (Fig. 4F) and located

comparatively closure to the hub (Fig. S4D), supporting its redox-dependent

enhancement to promote differentiation. The deregulated proliferation of cystic cells

reconciled with higher levels of its self-renewing inducers PI3K effector, p4E-BP (Fig.

4G', H' and I) and Hedgehog (Hh) (Fig. 4M' and N') [41] in *Tj>Sod1i* testis. The noticeably

high p4E-BP was associated with Cyclin-D accumulation (Fig. 4J). Similarly, depletion

of Sod1 in CySC resulted in elevated expression of Hh receptor Patched (Ptc) (Fig.

4O) which extended to regions farther from the hub, overlapping with expanded Tj^+

cells (Fig. 4M-N). Since, Ptc itself is a transcriptional Hh target, its expression in CySCs

suggest active Hh signalling[42], further supported by higher levels and similar pattern

of Hh transcriptional effector Cubitus interruptus (Ci) in *Sod1i* testis (Fig. 4P and S4E'''-

F'''). Since, Ptc, Ci and a GPCR-like signal transducer Smoothened (Smo) are all

transcriptionally driven by Hh, and Hh itself is regulated by higher levels of Ci[43,44], we

expectedly found higher levels of transcripts for these pathway genes (Fig. S4G-J),

indicating ROS-mediated upregulation of Hh activity. The overt proliferation of CySCs

under high ROS conditions was suppressed by reducing gene dosage either by single

copy of *PI3K^{DN}* (dominant-negative allele) or Hh-RNAi (Fig. 4L and Fig. S4K-N) which

was also associated with an equivalent maintenance of GSC population (Fig. 4K and

S4O-R). Together these observations suggest that intercellular redox gradients

balance the stem-cell populations in the niche through CySC maintenance factors.

239 Elevating CySC antioxidant defence promotes GSC self-renewal

To further substantiate the role of ROS in coordinating the multi-lineage stem

populations, we strengthened the cellular defences against oxidants by

overexpressing Sod1 in CySCs and monitored the relative GSC/CySC populations.
Indeed, we observed an increase in the number of Vasa$^+$ cells (Fig. 5A and S5A''-B''),
together with a slight reduction in Tj$^+$ cells (Fig. 5B and S5E''-F''). Any morphological
anomalies associated with cell-cell adhesion or abnormal cellular dispersion, was not
observed (Fig. S5A-B and E-F). Enhancing the levels of Sod1 in GSCs promoted the
growth of GSC-like cells (Fig. S5C'', D'' and I) but did not have any prominent effect
on CySCs (Fig. S5G'', H'' and J). The increment in Vasa$^+$ GSCs under CySC-induced
low redox conditions was also represented by parallel increase in spectrosome
number (Fig. 5C-E). The uptick in GSC number due to scavenging of ROS in CySC,
might be a result of delayed differentiation as indicated by displacement of Bam$^+$ zone
away from hub; thus, confirming non-cell-autonomous role of CySC ROS in
maintaining GSC fate (Fig. 5F-H). The data suggest that reducing redox profile of
CySCs differentially affected their own self-renewing propensies along with GSC
maintenance and optimum redox state of both the stem-cell populations is controlled
by CySCs.

Discussion:

ROS is a crucial mediator of stem cell maintenance where these species act as second
messengers to induce different post-translational protein modifications thereby,
affecting cell fate[45,46]. Effectuation of ROS is mainly considered to be autocrine in
nature where generation of these oxidative species is compartmentalized with their
site of action. Recent studies have reported non-autonomous generation of ROS by
NOX enzymes or upon stimulation by growth factors, to play a critical role in
regenerative growth[47-49]. However, in either of the cases, the target cell serves as the
source and consequent response. Our observations in multi-lineage *Drosophila* stem
cell niche suggested a previously unaccounted transmission of redox signals inter-
cellularly that act as an organizer of niche homoeostasis. The higher ROS threshold
of CySCs and resultant CySC-GSC oxidative gradient maintained the physiological
redox state of GSC. Reduced ROS in GSC was accompanied with its proliferation[50],
phenocopied when Sod1 was overexpressed in its neighbours (Fig. 6).

The GSCs are attached to the hub via adherens junction to receive the maintenance signals like, Upd that trigger the JAK-STAT pathway[27,28], suppression of which activates Bam, acting as a switch from transit amplifying cells to spermatocyte differentiation[51,52]. E-cadherin levels ensure only undifferentiated stem cells to persist within the niche, while differentiating GSCs lose the competition for niche occupancy[25]. Elevated ROS in CySC possibly affected STAT phosphorylation through S-glutathionylation[53,54], causing reduced presence of dependent transcripts, Socs36E and E-cadherin thereby, resulting in detachment of GSCs from the neighbours as under E-cadherin depleted conditions[55-58]. This resulted in compromised GSC-CySC paracrine receptions such as EGFR which play a crucial role in maintenance of cell-polarity that segregates self-renewing CySC populations from the ones receiving differentiation cues [39,59,60], leading to accumulation of cells co-expressing both stemness and differentiation markers. In contrast to GSCs, ROS imbalance in CySCs induced accelerated differentiation due to deregulation of PI3K/Tor and Hedgehog pathways. Although, redox modulation of PI3K pathway components has been previously demonstrated[61], in this study we found Hedgehog to be probably susceptible to redox regulation.

However, we do not rule out the possibilities of hub cells playing an equivalent role in GSC maintenance, given their parallel effect in suppressing GSC differentiation[28]. We had tried using E-132 driver to express *Sod1i* lines in hub cells but were challenged by substantial adverse effect on fly viability. We also avoided usage of Gal80 dependent clonal populations to maintain a homogenous genetic background and prevent false readouts due to diffusion of ROS signals in unaltered neighbourhood.

The proposed concept of intercellular ROS communication can be of importance in deciphering the biochemical adaptability and plasticity of different niches influencing stem cell fate, ensuring niche size and architecture to prevent stem cell loss and aging. This has been observed in neural stem cells where inflammation-induced quiescence recovers the regenerating capacity of aging brain[62]. In addition to metabolic variations, dependence of the germline on somatic neighbours for its redox state might be one the reasons behind its presumed immortal nature and renitence to aging[63]. The same can also be applied to cancer stem cells which maintain niche occupancy and resist oxidative stress induced apoptosis probably, by receiving redox cues from the

304 environment. However, further work is required to elucidate the myriad of driver
305 mechanisms intersecting at the realm of redox regulation that extend beyond the
306 present system into broader translational areas.

Materials and Methods

Fly strains

Fly stocks were maintained and crosses were set at 25 °C on normal corn meal and
yeast medium unless otherwise indicated. All fly stocks (BL-24493) *UAS SOD1 RNAi*,
(BL-32909) *UAS SOD1 RNAi*, (BL-29389) *UAS SOD1 RNAi*, (BL-32983) *UAS SOD2*
RNAi, (BL-24754) *UAS SOD1*, (BL- 32489) *UAS hhRNAi*, (BL-8288) *UAS PI3K^{DN}*, (BL-
25751) *UAS Dcr-nos*, (BL-55122) *UAS-FUCCI*, were obtained from Bloomington
Drosophila Stock Centre. *Nos-Gal4*, *Tj-Gal4*, *Tj-Gal4-BamGFP*, *gstD1-GFP*, *TFAM-*
GFP were kind gift from U. Nongthomba, P. Majumder, K. Ray, B. C. Mandal, Hong
Xu labs respectively. For optimal Gal4 activity parent crosses were shifted to 29 °C till
the eclosion of adults. 3-5 days old males were taken for all the experiments. The
control lines for all experiments are the corresponding Gal4-line crossed with wild type
OregonR⁺.

Dissection and immunostaining

Anesthetized flies were dissected in 1X Phosphate Buffer Saline (PBS). All incubations
were carried out at room temperature (25 °C) unless otherwise mentioned. Testes
were fixed in 4% paraformaldehyde for 30 minutes, followed by multiple washes with
0.1% PBTX (PBS + TritonX 100). Post-blocking in 0.5% Bovine Serum Albumin, testis
was incubated overnight at 4°C for primary antisera, followed by washing in 0.1%
PBTX 3 times 15 minutes each before incubating with secondary antibody. The tissues
were counterstained with DAPI (4,6-diamidino-2-phenylindole) for 20 minutes,
followed by three washes in 0.1% PBTX 10 minutes each. For PTC staining, a modified
protocol utilizing PIPES-EGTA buffer was used as described[64]. The dpErk was
labelled by dissecting fly testes in Schneider's media, followed by fixation in 4% PFA
for 30 min, 3x wash in 0.1% PBTX, blocked and incubated in primary antibody
overnight; each step supplemented with phosphatase inhibitor cocktail 2 (1:100,

Sigma, cat#P5726). Samples were mounted in DABCO anti-fade medium prior imaging.

The following primary antibodies were used in the experiments - anti-FasIII (7G10) (1:120, DSHB (Developmental Studies Hybridoma Bank)), anti-Eya (1:50, DSHB), anti-DEcad (DCAD2) (1:50, DSHB), anti-α-Spectrin (3A9) (1:20, DSHB), anti-Ptc (1:100,DSHB) , anti-Ci (1:50, DSHB), anti-βtubulin (1:300, DSHB), anti-pERK (1:100, Cell Signaling (4370)), ATP5A (1:700, Abcam (ab14748)), anti-Tj (1:5000), anti-Vasa (1:4000), anti-Zfh1(1:2000). The primary labelling was detected using appropriate Alexa-Flour tagged secondary antibody (ThermoFisher Scientific).

To assess superoxide levels, testes were dissected in Schneider's medium (SM) and immediately incubated in 30 µM Dihydroxyethidium (DHE) at 25 °C in dark. Testes were washed thrice with SM for 7 minutes each before quantifying the emitted fluorescence using SpectraMax iD5 (Molecular Devices).

Quantitative Reverse transcription–PCR

For semi-quantitative RT-PCR and qRT-PCR analyses, total RNA was isolated from testes of 3-5 days old male flies using Qiagen RNA extraction kit. RNA pellets were resuspended in nuclease-free water and quantity of RNA was spectrophotometrically estimated. First strand cDNA was synthesized from 1-2 µg of total RNA as described earlier. The prepared cDNAs were subjected to real time PCR using forward and reverse primer pairs as listed below, using 5 µl qPCR Master Mix (SYBR Green, ThermoFisher Scientific), 2 pmol/µl of each primer per reaction for 10 µl final volume in ABI 7500 Real time PCR machine. The fold change in expression was calculated through $2^{-\Delta\Delta Ct}$ method. The primers used are listed in Supplementary Table S1.

Imaging and image analysis

Confocal imaging was carried out using Zeiss LSM 900 confocal microscope, using appropriate dichroics and filters. Images were further analyzed and processed for brightness and contrast adjustments using ImageJ (Fiji). Mean intensity measurements were carried out using standard Fiji plug-ins, 2D surface plot quantification or Zeiss Zen software as indicated. Quantification of cell number/nuclei

in single or co-stained populations (Tj, Vasa, Zfh1, Eya, FUCCI) were performed from individual slices of z-stacks using semi-automated Cell Counter plugin in Fiji for all the mentioned genotypes. The relative distance from the two reference points in the images was estimated through Inter-edge Distance ImageJ Macro v2.0 from Github. All images were assembled using Adobe Photoshop version 21.2.1.

Immunoblot Analysis

Protein was extracted from the dissected testes using RIPA buffer as described previously and quantified using Bradford reagent (Biorad). Equivalent concentration of lysate was denatured using 1X Laemelli Buffer with 1 M DTT at 95°C, separated in 10% SDS-PAGE and transferred onto PVDF membranes. Membranes were blocked with commercial blocking solution in TBST base before sequential primary antibody incubation with anti-Vasa and anti-β-Tubulin (DSHB, E7). Secondary detection was performed using HRP-tag anti-mouse antibody (GE Amersham).

Statistical analyses

All data points represent at least three biological replicates and the sample sizes carrying adequate statistical power are mentioned in the figure legends. Statistical significance for each experiment were calculated using two tailed Student's t-test unless otherwise mentioned, using GraphPad Prism 8 software. The P values were calculated through pairwise comparison of the data with wild type or driver alone and driven RNAi lines. Significance values for sample sizes mentioned in figure legends were represented as $*P < 0.01$, $**P < 0.001$, or $***P < 0.0001$.

Author contribution

D.S. and O.M conceived the study and designed the experiments. O.M, A.C and T.C performed the experiments. D.S and O.M analyzed the data. D.S, O.M. and A.C. wrote the manuscript.

Acknowledgements

We thank P. Majumder, U. Nongthomba, K. Ray, G. Ratnaparkhi, B. C. Mandal, Hong

Xu lab and S. C. Lakhotia for kindly sharing some of the transgenic fly lines. Christian

Bökel for sharing experimental protocols. Anti-Tj antibody was a generous gift from

Prof. D. Godt, University of Toronto. Anti-Vasa and Anti-Zfh1 were kindly gifted by

Prof. R. Lehmann, New York University. We also thank Dr. Bama Charan Mandal for

generously sharing his Confocal microscope imaging system and Sudeshna

Majumder for preliminary data. We acknowledge BDSC for fly stocks and DSHB for

antibodies. This work is supported by Innovative Young Biotechnologist Award by

Department of Biotechnology (BT/12/IYBA/2019/01), Early Career Research Award

by Science and Engineering Research Board (ECR/2018/000009), BHU-Institute of

Eminence grant (R/Dev/IoE/Incentive/2021-22/32452). The authors also acknowledge

financial support from DST-INSPIRE fellowship (to O.M.), UGC-CAS doctoral

fellowship (to T.C.).

Declaration of interest

The authors declare no competing interests

Data availability

The data that support the findings of this study are available within the main text and

its Supplementary Information file. The lead contact will share all raw data associated

with this paper upon request. This study did not utilize or generate any unique datasets

or codes.

References

1.   Schieber, M., and Chandel, N.S. (2014). ROS function in redox signaling and oxidative
stress. Curr Biol *24*, R453-462. 10.1016/j.cub.2014.03.034.
2.   Holmstrom, K.M., and Finkel, T. (2014). Cellular mechanisms and physiological
consequences of redox-dependent signalling. Nat Rev Mol Cell Biol *15*, 411-421.
10.1038/nrm3801.
3.   D'Autreaux, B., and Toledano, M.B. (2007). ROS as signalling molecules: mechanisms
that generate specificity in ROS homeostasis. Nat Rev Mol Cell Biol *8*, 813-824.
10.1038/nrm2256.
4.   Maryanovich, M., and Gross, A. (2013). A ROS rheostat for cell fate regulation. Trends
Cell Biol *23*, 129-134. 10.1016/j.tcb.2012.09.007.
5.   Murphy, M.P. (2009). How mitochondria produce reactive oxygen species. Biochem J
417, 1-13. 10.1042/BJ20081386.

6.   Hamanaka, R.B., and Chandel, N.S. (2010). Mitochondrial reactive oxygen species
433   regulate cellular signaling and dictate biological outcomes. Trends Biochem Sci *35*,
434   505-513. 10.1016/j.tibs.2010.04.002.
435   7.   Al-Mehdi, A.B., Pastukh, V.M., Swiger, B.M., Reed, D.J., Patel, M.R., Bardwell, G.C.,
436   Pastukh, V.V., Alexeyev, M.F., and Gillespie, M.N. (2012). Perinuclear mitochondrial
437   clustering creates an oxidant-rich nuclear domain required for hypoxia-induced
438   transcription. Sci Signal *5*, ra47. 10.1126/scisignal.2002712.
439   8.   Bedard, K., and Krause, K.H. (2007). The NOX family of ROS-generating NADPH
440   oxidases: physiology   and   pathophysiology.   Physiol   Rev   *87*,   245-313.
441   10.1152/physrev.00044.2005.
442   9.   Appenzeller-Herzog, C., Banhegyi, G., Bogeski, I., Davies, K.J., Delaunay-Moisan, A.,
443   Forman, H.J., Gorlach, A., Kietzmann, T., Laurindo, F., Margittai, E., et al. (2016).
444   Transit of H2O2 across the endoplasmic reticulum membrane is not sluggish. Free
445   Radic Biol Med *94*, 157-160. 10.1016/j.freeradbiomed.2016.02.030.
446   10.   Mishina, N.M., Bogdanova, Y.A., Ermakova, Y.G., Panova, A.S., Kotova, D.A., Bilan,
447   D.S., Steinhorn, B., Arner, E.S.J., Michel, T., and Belousov, V.V. (2019). Which
448   Antioxidant System Shapes Intracellular H(2)O(2) Gradients? Antioxid Redox Signal
449   *31*, 664-670. 10.1089/ars.2018.7697.
450   11.   Yoboue, E.D., Sitia, R., and Simmen, T. (2018). Redox crosstalk at endoplasmic
451   reticulum (ER) membrane contact sites (MCS) uses toxic waste to deliver messages.
452   Cell Death Dis *9*, 331. 10.1038/s41419-017-0033-4.
453   12.   Rehman, J. (2010). Empowering self-renewal and differentiation: the role of
454   mitochondria in stem cells. J Mol Med (Berl) *88*, 981-986. 10.1007/s00109-010-0678-
455   2.
456   13.   Li, T.S., and Marban, E. (2010). Physiological levels of reactive oxygen species are
457   required to maintain genomic stability in stem cells. Stem Cells *28*, 1178-1185.
458   10.1002/stem.438.
459   14.   Le Belle, J.E., Orozco, N.M., Paucar, A.A., Saxe, J.P., Mottahedeh, J., Pyle, A.D., Wu,
460   H., and Kornblum, H.I. (2011). Proliferative neural stem cells have high endogenous
461   ROS levels that regulate self-renewal and neurogenesis in a PI3K/Akt-dependant
462   manner. Cell Stem Cell *8*, 59-71. 10.1016/j.stem.2010.11.028.
463   15.   Chuikov, S., Levi, B.P., Smith, M.L., and Morrison, S.J. (2010). Prdm16 promotes stem
464   cell maintenance in multiple tissues, partly by regulating oxidative stress. Nat Cell Biol
465   *12*, 999-1006. 10.1038/ncb2101.
466   16.   Owusu-Ansah, E., and Banerjee, U. (2009). Reactive oxygen species prime
467   Drosophila haematopoietic progenitors for differentiation. Nature *461*, 537-541.
468   10.1038/nature08313.
469   17.   Tan, S.W.S., Lee, Q.Y., Wong, B.S.E., Cai, Y., and Baeg, G.H. (2017). Redox
470   Homeostasis Plays Important Roles in the Maintenance of the Drosophila Testis
471   Germline Stem Cells. Stem Cell Reports *9*, 342-354. 10.1016/j.stemcr.2017.05.034.
472   18.   Zhou, G., Meng, S., Li, Y., Ghebre, Y.T., and Cooke, J.P. (2016). Optimal ROS
473   Signaling   Is   Critical   for   Nuclear   Reprogramming.   Cell   Rep   *15*,   919-925.
474   10.1016/j.celrep.2016.03.084.
475   19.   Sinenko, S.A., Shim, J., and Banerjee, U. (2011). Oxidative stress in the
476   haematopoietic niche regulates the cellular immune response in Drosophila. EMBO
477   Rep *13*, 83-89. 10.1038/embor.2011.223.
478   20.   Davies, E.L., and Fuller, M.T. (2008). Regulation of self-renewal and differentiation in
479   adult stem cell lineages: lessons from the Drosophila male germ line. Cold Spring Harb
480   Symp Quant Biol *73*, 137-145. 10.1101/sqb.2008.73.063.
481   21.   Hime, G.R., Brill, J.A., and Fuller, M.T. (1996). Assembly of ring canals in the male
482   germ line from structural components of the contractile ring. J Cell Sci *109 (Pt 12)*,
483   2779-2788. 10.1242/jcs.109.12.2779.

484   22.   Gonczy, P., and DiNardo, S. (1996). The germ line regulates somatic cyst cell
485   proliferation and fate during Drosophila spermatogenesis. Development *122*, 2437-
486   2447. 10.1242/dev.122.8.2437.

487   23.   Fabrizio, J.J., Boyle, M., and DiNardo, S. (2003). A somatic role for eyes absent (eya)
488   and sine oculis (so) in Drosophila spermatocyte development. Dev Biol *258*, 117-128.
489   10.1016/s0012-1606(03)00127-1.

490   24.   Hardy, R.W., Tokuyasu, K.T., Lindsley, D.L., and Garavito, M. (1979). The germinal
491   proliferation center in the testis of Drosophila melanogaster. J Ultrastruct Res *69*, 180-
492   190. 10.1016/s0022-5320(79)90108-4.

493   25.   Chen, S., Lewallen, M., and Xie, T. (2013). Adhesion in the stem cell niche: biological
494   roles and regulation. Development *140*, 255-265. 10.1242/dev.083139.

495   26.   Inaba, M., Yuan, H., Salzmann, V., Fuller, M.T., and Yamashita, Y.M. (2010). E-
496   cadherin is required for centrosome and spindle orientation in Drosophila male
497   germline stem cells. PLoS One *5*, e12473. 10.1371/journal.pone.0012473.

498   27.   Kiger, A.A., Jones, D.L., Schulz, C., Rogers, M.B., and Fuller, M.T. (2001). Stem cell
499   self-renewal specified by JAK-STAT activation in response to a support cell cue.
500   Science *294*, 2542-2545. 10.1126/science.1066707.

501   28.   Tulina, N., and Matunis, E. (2001). Control of stem cell self-renewal in Drosophila
502   spermatogenesis by JAK-STAT signaling. Science *294*, 2546-2549.
503   10.1126/science.1066700.

504   29.   Song, X., Wong, M.D., Kawase, E., Xi, R., Ding, B.C., McCarthy, J.J., and Xie, T.
505   (2004). Bmp signals from niche cells directly repress transcription of a differentiation-
506   promoting gene, bag of marbles, in germline stem cells in the Drosophila ovary.
507   Development *131*, 1353-1364. 10.1242/dev.01026.

508   30.   Kawase, E., Wong, M.D., Ding, B.C., and Xie, T. (2004). Gbb/Bmp signaling is
509   essential for maintaining germline stem cells and for repressing bam transcription in
510   the Drosophila testis. Development *131*, 1365-1375. 10.1242/dev.01025.

511   31.   Losick, V.P., Morris, L.X., Fox, D.T., and Spradling, A. (2011). Drosophila stem cell
512   niches: a decade of discovery suggests a unified view of stem cell regulation. Dev Cell
513   *21*, 159-171. 10.1016/j.devcel.2011.06.018.

514   32.   Jemc, J.C. (2011). Somatic gonadal cells: the supporting cast for the germline.
515   Genesis *49*, 753-775. 10.1002/dvg.20784.

516   33.   Zoller, R., and Schulz, C. (2012). The Drosophila cyst stem cell lineage: Partners
517   behind the scenes? Spermatogenesis *2*, 145-157. 10.4161/spmg.21380.

518   34.   Larsson, N.G., Wang, J., Wilhelmsson, H., Oldfors, A., Rustin, P., Lewandoski, M.,
519   Barsh, G.S., and Clayton, D.A. (1998). Mitochondrial transcription factor A is necessary
520   for mtDNA maintenance and embryogenesis in mice. Nat Genet *18*, 231-236.
521   10.1038/ng0398-231.

522   35.   Sykiotis, G.P., and Bohmann, D. (2008). Keap1/Nrf2 signaling regulates oxidative
523   stress tolerance and lifespan in Drosophila. Dev Cell *14*, 76-85.
524   10.1016/j.devcel.2007.12.002.

525   36.   Zielke, N., Korzelius, J., van Straaten, M., Bender, K., Schuhknecht, G.F.P., Dutta, D.,
526   Xiang, J., and Edgar, B.A. (2014). Fly-FUCCI: A versatile tool for studying cell
527   proliferation in complex tissues. Cell Rep *7*, 588-598. 10.1016/j.celrep.2014.03.020.

528   37.   Li, M.A., Alls, J.D., Avancini, R.M., Koo, K., and Godt, D. (2003). The large Maf factor
529   Traffic Jam controls gonad morphogenesis in Drosophila. Nat Cell Biol *5*, 994-1000.
530   10.1038/ncb1058.

531   38.   Amoyel, M., Hillion, K.H., Margolis, S.R., and Bach, E.A. (2016). Somatic stem cell
532   differentiation is regulated by PI3K/Tor signaling in response to local cues.
533   Development *143*, 3914-3925. 10.1242/dev.139782.

534   39.   Castanieto, A., Johnston, M.J., and Nystul, T.G. (2014). EGFR signaling promotes self-
535   renewal through the establishment of cell polarity in Drosophila follicle stem cells. Elife
536   *3*. 10.7554/eLife.04437.

537   40.   Papagiannouli, F., Berry, C.W., and Fuller, M.T. (2019). The Dlg Module and Clathrin-
538   Mediated Endocytosis Regulate EGFR Signaling and Cyst Cell-Germline Coordination
539   in the Drosophila Testis. Stem Cell Reports *12*, 1024-1040.
540   10.1016/j.stemcr.2019.03.008.
541   41.   Michel, M., Kupinski, A.P., Raabe, I., and Bokel, C. (2012). Hh signalling is essential
542   for somatic stem cell maintenance in the Drosophila testis niche. Development *139*,
543   2663-2669. 10.1242/dev.075242.
544   42.   Chen, Y., and Struhl, G. (1996). Dual roles for patched in sequestering and transducing
545   Hedgehog. Cell *87*, 553-563. 10.1016/s0092-8674(00)81374-4.
546   43.   Chen, Y., and Struhl, G. (1998). In vivo evidence that Patched and Smoothened
547   constitute distinct binding and transducing components of a Hedgehog receptor
548   complex. Development *125*, 4943-4948. 10.1242/dev.125.24.4943.
549   44.   Jiang, J., and Hui, C.C. (2008). Hedgehog signaling in development and cancer. Dev
550   Cell *15*, 801-812. 10.1016/j.devcel.2008.11.010.
551   45.   Sinenko, S.A., Starkova, T.Y., Kuzmin, A.A., and Tomilin, A.N. (2021). Physiological
552   Signaling Functions of Reactive Oxygen Species in Stem Cells: From Flies to Man.
553   Front Cell Dev Biol *9*, 714370. 10.3389/fcell.2021.714370.
554   46.   Liang, R., and Ghaffari, S. (2014). Stem cells, redox signaling, and stem cell aging.
555   Antioxid Redox Signal *20*, 1902-1916. 10.1089/ars.2013.5300.
556   47.   Fu, X.J., Peng, Y.B., Hu, Y.P., Shi, Y.Z., Yao, M., and Zhang, X. (2014). NADPH
557   oxidase 1 and its derived reactive oxygen species mediated tissue injury and repair.
558   Oxid Med Cell Longev *2014*, 282854. 10.1155/2014/282854.
559   48.   Taylor-Fishwick, D.A. (2013). NOX, NOX Who is There? The Contribution of NADPH
560   Oxidase One to Beta Cell Dysfunction. Front Endocrinol (Lausanne) *4*, 40.
561   10.3389/fendo.2013.00040.
562   49.   Santabarbara-Ruiz, P., Esteban-Collado, J., Perez, L., Viola, G., Abril, J.F., Milan, M.,
563   Corominas, M., and Serras, F. (2019). Ask1 and Akt act synergistically to promote
564   ROS-dependent regeneration in Drosophila. PLoS Genet *15*, e1007926.
565   10.1371/journal.pgen.1007926.
566   50.   Pan, L., Chen, S., Weng, C., Call, G., Zhu, D., Tang, H., Zhang, N., and Xie, T. (2007).
567   Stem cell aging is controlled both intrinsically and extrinsically in the Drosophila ovary.
568   Cell Stem Cell *1*, 458-469. 10.1016/j.stem.2007.09.010.
569   51.   Bausek, N. (2013). JAK-STAT signaling in stem cells and their niches in Drosophila.
570   JAKSTAT *2*, e25686. 10.4161/jkst.25686.
571   52.   Gonczy, P., Matunis, E., and DiNardo, S. (1997). bag-of-marbles and benign gonial
572   cell neoplasm act in the germline to restrict proliferation during Drosophila
573   spermatogenesis. Development *124*, 4361-4371. 10.1242/dev.124.21.4361.
574   53.   Butturini, E., Darra, E., Chiavegato, G., Cellini, B., Cozzolino, F., Monti, M., Pucci, P.,
575   Dell'Orco, D., and Mariotto, S. (2014). S-Glutathionylation at Cys328 and Cys542
576   impairs STAT3 phosphorylation. ACS Chem Biol *9*, 1885-1893. 10.1021/cb500407d.
577   54.   Xiong, Y., Uys, J.D., Tew, K.D., and Townsend, D.M. (2011). S-glutathionylation: from
578   molecular mechanisms to health outcomes. Antioxid Redox Signal *15*, 233-270.
579   10.1089/ars.2010.3540.
580   55.   DeGennaro, M., Hurd, T.R., Siekhaus, D.E., Biteau, B., Jasper, H., and Lehmann, R.
581   (2011). Peroxiredoxin stabilization of DE-cadherin promotes primordial germ cell
582   adhesion. Dev Cell *20*, 233-243. 10.1016/j.devcel.2010.12.007.
583   56.   Karsten, P., Hader, S., and Zeidler, M.P. (2002). Cloning and expression of Drosophila
584   SOCS36E and its potential regulation by the JAK/STAT pathway. Mech Dev *117*, 343-
585   346. 10.1016/s0925-4773(02)00216-2.
586   57.   Richmond, C.A., Rickner, H., Shah, M.S., Ediger, T., Deary, L., Zhou, F., Tovaglieri,
587   A., Carlone, D.L., and Breault, D.T. (2018). JAK/STAT-1 Signaling Is Required for
588   Reserve Intestinal Stem Cell Activation during Intestinal Regeneration Following Acute
589   Inflammation. Stem Cell Reports *10*, 17-26. 10.1016/j.stemcr.2017.11.015.

590   58.   Leatherman, J.L., and Dinardo, S. (2010). Germline self-renewal requires cyst stem
591   cells and stat regulates niche adhesion in Drosophila testes. Nat Cell Biol *12*, 806-811.
592   10.1038/ncb2086.
593   59.   Kiger, A.A., White-Cooper, H., and Fuller, M.T. (2000). Somatic support cells restrict
594   germline stem cell self-renewal and promote differentiation. Nature *407*, 750-754.
595   10.1038/35037606.
596   60.   Schulz, C., Wood, C.G., Jones, D.L., Tazuke, S.I., and Fuller, M.T. (2002). Signaling
597   from germ cells mediated by the rhomboid homolog stet organizes encapsulation by
598   somatic support cells. Development *129*, 4523-4534. 10.1242/dev.129.19.4523.
599   61.   Koundouros, N., and Poulogiannis, G. (2018). Phosphoinositide 3-Kinase/Akt
600   Signaling and Redox Metabolism in Cancer. Front Oncol *8*, 160.
601   10.3389/fonc.2018.00160.
602   62.   Kalamakis, G., Brune, D., Ravichandran, S., Bolz, J., Fan, W., Ziebell, F., Stiehl, T.,
603   Catala-Martinez, F., Kupke, J., Zhao, S., et al. (2019). Quiescence Modulates Stem
604   Cell Maintenance and Regenerative Capacity in the Aging Brain. Cell *176*, 1407-1419
605   e1414. 10.1016/j.cell.2019.01.040.
606   63.   Snoeck, H.W. (2015). Can Metabolic Mechanisms of Stem Cell Maintenance Explain
607   Aging and the Immortal Germline? Cell Stem Cell *16*, 582-584.
608   10.1016/j.stem.2015.04.021.
609   64.   Michel, M., Raabe, I., Kupinski, A.P., Perez-Palencia, R., and Bokel, C. (2011). Local
610   BMP receptor activation at adherens junctions in the Drosophila germline stem cell
611   niche. Nat Commun *2*, 415. 10.1038/ncomms1426.

Figure Legends

Figure 1. Mitochondrial Distribution and Redox Profile of GSCs and CySCs in Adult *Drosophila* Testes

(A) Schematic representation of adult *Drosophila* testicular niche showing arrangement of different stem cell populations (GSC – Germinal Stem Cell, CySC – Cystic Stem Cell). (B-E) Differential distribution of mitochondria labelled with ATP5A (green) in Vasa[+] GSCs and Tj[+] CySCs of wild-type fly testis; B# shows digitally zoomed image of the dotted area. (F) Relative mitochondria density profile of GSCs and CySCs denoted through cooccurrence of ATP5A with Tj or Vasa. A.U. - arbitrary units. (G-P) Redox profiling of testicular stem cell niche using *gstD1*-GFP as intrinsic ROS reporter in control (G-K) and *Tj-Gal4* driven *Sod1RNAi (Sod1i)* testis (L-P). (Q) Gaussian distribution curve for the variations in *gstD1*-GFP fluorescence at CySC-GSC intersection region (inset) in *Tj-Gal4* control and *Sod1i* testis. (R) Quantification of mean *gstD1*-GFP fluorescence intensity (MFI) upon Sod1 knockdown represented as fold change over control. Data denotes mean ± s.e.m, n = 25 (S). Colocalization intensity between Vasa and *gstD1*-GFP expressed through Manders' co-localization

630   coefficient (MCC) among control and *Sod1i* cells (G). Controls are the indicated driver
631   line crossed with *Oregon R⁺*. Scale bar - 10 µm. See also Figure S1.

Figure 2. Redox disequilibrium in Cyst stem cell lineage deregulates early CySCs and decreases the GSC number

(A-B) Distribution of Tj⁺ CySC/early cyst cell around the hub (FasIII) in control and
driven *Sod1RNAi* testis. (C-D) Represents the rosette arrangement of Vasa⁺ GSCs
flanking the hub. The digitally magnified region around the hub (dotted line) as seen
in control (C#) and *Sod1RNAi* (D#). (E-G) Mean number of Tj⁺ cells, Zfh1⁺ early CySCs
and GSCs in control and *Sod1RNAi* lines shown as mean ± s.e.m, n = 10. (H)
Construct design of the FUCCI reporter line for tracking the cell cycle stages in
Drosophila tissues, containing degrons of Cyclin B and E2F1 proteins fused with RFP
or GFP. G1, S and G2/M is represented by GFP⁺, RFP⁺ and dual labelled cells
respectively. (I-J) Comparative changes in the total cell population (I) or Zfh1⁺ cells (J)
present in G1, S and G2/M phase at the niche zone between control and *Tj>Sod1i*.
Data points denotes mean ± s.e.m, n = 10. Scale bar - 10 µm. See also Figure S2.

Figure 3. ROS imbalance in CySCs promotes differentiation of both GSCs and CySCs

(A-D) Number of early CySCs (Zfh1⁺) (A'-B'), late differentiating cyst cells (Eya⁺) (A-
B) and Zfh1, Eya co-expressing population were imaged (A''-B''), represented
schematically (C) and quantified with respect to control and *Sod1RNAi* lines (D). Data
points represent mean ± s.e.m, n = 10. (E-F) The effect of Sod1 depletion in CySCs
on the differentiation status of GSCs as observed using *Bam-GFP* reporter line. (G)
The relative distance of the differentiation initiation zone (Bam⁺) from the hub (red) in
control and *Sod1i* testis shown as mean ± s.e.m, n = 10. (H-J) The shape and size of
the spectrosomes marked with α-spectrin (green) were imaged (I''-J'') and quantified
for their distance from the hub (marked with asterisk) (H), n = 10. Branched fusome
marks differentiating populations (I'-J'). (K-L) Comparative staining of CySC-GSC
contacts through adherens junction using E-cadherin (monochrome/green). Dotted
area near the hub has been expanded as inset to show loss of E-cadherin network.

(M) 3D surface intensity histogram for E-cadherin expression near the hub. (N-O) Fold

change in expression of Stat dependent transcripts Socs36E (N) and Ptp61F (O)

among control and *Sod1i* niche. Data is shown as mean ± s.e.m, n = 3. Scale bar - 10

µm. See also Figure S3.

Figure 4. Altered CySC redox state affects niche maintenance signals.

(A-B) The expression of cell polarity marker Dlg (red) in control and *Sod1i* stem-cell

niche (A''-B''), (C-D) Representative image showing pErk distribution parallelly with Tj$^+$

cells and its expression pattern through Fire LUT (C''-D''). Scale bar - 10 µm. (E)

Segregated pErk expression patterns in CySCs near the hub and differentiating cyst

cells (CC), mean ± s.e.m, n = 200. (F) Estimation of the differentiation zone through

quantification of high pErk expressing regions. (G-I) Activation of PI3K/Tor in somatic

lineage was adjudged through reporter p4E-BP in *Tj>Sod1i* (G'-H') and levels of the

active form was quantified (I), MFI – mean fluorescence intensity shown as mean ±

s.e.m, n = 40. (J) Induction of Cyclin D in *Sod1i* represented as fold change over

controls by mean ± s.e.m, n = 3. (K-L) The number of Vasa$^+$ (K) and Tj$^+$ (L) cells across

control, *Sod1i*, *Sod1i/PI3K^{DN}* (dominant negative allele) and *Sod1i/Hhi* rescue samples

depicted as mean ± s.e.m, n = 10. (M'-N') Representative image showing distribution

of Patched (Ptc) in Tj$^+$ cells in control and *Tj>Sod1i*. (O-P) Quantification of Ptc (O)

and Hh effector Ci (P) expression through fluorescence intensity as mean ± s.e.m, n

(Ptc) = 90, n (Ci) = 200. See also Figure S4.

Figure 5. Low CySC ROS sustains GSC maintenance

(A-B) Bar graphs illustrating variations in Vasa$^+$ GSC (A) and Tj$^+$ CySCs (B) numbers

upon overexpressing Sod1 (*Tj>Sod^{OE}*), represented as mean ± s.e.m, n = 10. (C-E)

The spectrosomes labelled with α-spectrin (α-spec) were imaged (C'-D') and their

number was quantified (E). (F-H) The initiation of gonialblast differentiation was

determined by measuring the distance of Bam$^+$ zone from hub (red), n = 10 (F), (G''-

H'') shows distribution of Bam$^+$ reporter expressing cells. Scale bar - 10 µm. See also

Figure S5.

Figure 6. Proposed role of intercellular redox gradients in GSC maintenance and differentiation

Mitochondrial abundance in GSC-CySC boundary maintains redox differential between two stem cell population. Disequilibrated intercellular redox gradient enhances precocious GSC differentiation and enhances CySC proliferation, thereby disrupting the homeostatic balance between the two stem populations.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Normal Condition

High-ROS Condition

E-cadherin

Hub

Hub

GSC

CySC

GSC

↓Sod1

Redox Gradient

↓Sod1

Redox Disequilibrium

GSCs ↓

Maintained Niche Architecture

Hh ↑
EGFR ↓
PI3K/Tor↑

No change in CySCs/CCs & Niche Architecture

CySCs and CCs ↑↑   GSCs ↓↓

Early Differentiation