SURVEY AND SUMMARY

Unlocking the Full Potential of Nanopore Sequencing: Tips, Tricks, and Advanced Data Analysis Techniques

Daria Meyer [1,2,*], **Winfried Goettsch** [1,3,*], **Jannes Spannenberg** [1,*], **Bettina Stieber** [2], **Sebastian Krautwurst** [1,4], **Martin Hölzer** [5], **Christian Brandt** [6], **Jörg Linde** [7], **Christian Höner zu Siederdissen** [1], **Akash Srivastava** [1,3], **Milena Zarkovic** [1,3], **Damian Wollny** [1,3,5,#], **and Manja Marz** [1,3,4,8−11,#]

[1] RNA Bioinformatics and High-Throughput Analysis, Friedrich Schiller University, Leutragraben 1, Jena, 07743, Germany
[2] Oncgnostics GmbH, Löbstedter Str. 41, Jena, 07749, Germany
[3] Leibniz Institute on Aging - Fritz Lipmann Institute, Beutenbergstraße 11, Jena, 07745, Germany
[4] European Virus Bioinformatics Center, Friedrich Schiller University, Leutragraben 1, Jena, 07743, Germany
[5] Genome Competence Center (MF1), Robert Koch Institute, Berlin 13353, Germany
[6] Institute of Infectious Diseases and Infection Control, Jena-University Hospital, Friedrich Schiller University, 07747 Jena, Germany
[7] Institute of Bacterial Infections and Zoonoses (IBIZ), Friedrich-Löffler-Institute, Naumburger Str. 96 a, 07743 Jena, Germany
[8] Max Planck Institute for Evolutionary Anthropology, Deutscher Pl. 6, Leipzig, 04103, Germany
[9] German Center for Integrative Biodiversity Research (iDiv), Puschstraße 4, Leipzig, 04103, Germany
[10] Michael Stifel Center Jena, Friedrich Schiller University, Ernst-Abbe-Platz 2, Jena, 07743, Germany
[11] Cluster of Excellence "Balance of the Microverse", Friedrich Schiller University, Fürstengraben 1, Jena, 07743, Germany
[*] These authors contributed equally.
[#] These authors contributed equally.

ABSTRACT

Nucleic acid sequencing is the process of identifying the sequence of DNA or RNA, with DNA used for genomes and RNA for transcriptomes. Deciphering this information has the potential to greatly advance our understanding of genomic features and cellular functions. In comparison to other available sequencing methods, nanopore sequencing stands out due to its unique advantages of processing long nucleic acid strands in real time, within a small portable device, enabling the rapid analysis of samples in diverse settings. Evolving over the past decade, nanopore sequencing remains in a state of ongoing development and refinement, resulting in persistent challenges in protocols and technology. This article employs an interdisciplinary approach, evaluating experimental and computational methods to address critical gaps in our understanding in order to maximize the information gain from this advancing technology. Here we present both overview and analysis of all aspects of nanopore sequencing by providing statistically supported insights. Thus, we aim to provide fresh perspectives on nanopore sequencing and give comprehensive guidelines for the diverse challenges that frequently impede optimal experimental outcomes.

INTRODUCTION

Nanopore sequencing is a transformative technology in genomics, offering the unique ability to sequence DNA or RNA molecules in their native form. Nanopore sequencing can generate long sequencing reads by measuring disturbances in the ion current as biological molecules such as DNA and RNA pass through a nanopore (1, 2). During or after sequencing, the raw DNA/RNA signal can be transformed into nucleotide sequences. This capability provides invaluable insights into

genetic variation and molecular modifications. Since the debut of the MinION device by Oxford Nanopore Technologies (ONT) in 2014, nanopore sequencing has experienced a surge in popularity and has become a cornerstone in genomic research. For DNA sequencing (2), it is attractive in diagnostics (3) and for metagenomic assemblies (4, 5, 6, 7), enabling the study of microbial communities and their functions. In direct RNA sequencing, it supports diverse tasks such as *de novo* transcriptome assembly, isoform expression quantification (8), and the direct detection of RNA modifications (9). These applications highlight the technology's versatility and its ability to address a wide range of biological questions.

One of the key advantages of nanopore sequencing is its ability to generate data rapidly, facilitated by quick library preparation and real-time data acquisition during sequencing (10). Additionally, nanopore sequencing can directly sequence native RNA molecules without the need for reverse transcription or amplification (8). Its capacity for long-read sequencing, where fragments up to two megabases in length can be read in a single pass (11), further enhances its utility by providing comprehensive genomic and transcriptomic insights. Furthermore, due to their portability and affordability, devices such as the MinION (12) are particularly valuable for applications such as monitoring virus outbreaks, where fast and on-site sequencing is essential.

Nanopore sequencing offers many advantages but comes with technical challenges. When first introduced, the technology was limited by lower accuracy ($\sim$60%) and throughput, though these shortcoming have improved significantly over time (13). With the introduction of ONT's MinION device in 2014 accuracy improved to 92% (14), with current rates reaching 99% (15). While in 2021 nanopore sequencing continued to exhibit a higher error rate compared to Illumina sequencing (2), in 2024 sequencing of nearly complete bacterial genomes without short-read or reference polishing became possible (16, 17). Nonetheless, sequencing accuracy remains a key consideration for certain high-precision applications, particularly in direct RNA sequencing (18).

Another challenge lies in the customization of nanopore wet lab workflows. While numerous reviews provide detailed overviews of tools for standard nanopore sequencing applications (10, 19, 20, 21), the field's relative novelty has given rise to a broad range of non-standard approaches aimed at maximizing sample utility. The versatility of nanopore sequencing lends itself to customization to meet specific requirements, including the use of various flow cells, library preparation kits, sequencing buffers, and an extensive set of computational analysis tools. The sheer number of options can be daunting for newcomers. Optimizing protocols, designing libraries, and selecting parameters require careful consideration. Compounding this issue is the prevalence of unverified claims in the field, as many methods lack experimental validation, making them challenging to integrate into standardized workflows. This is particularly problematic

when information is available solely through the Nanopore Community[1] and lacks proper citation.

This perspective work integrates insights gained from over 300 nanopore sequencing runs, which span a variety of species, sequencing devices, preparation methods, and sequencing protocols. These data-driven findings provide valuable guidance for addressing common challenges in nanopore sequencing. For instance, we show that flow cell performance is influenced more by the sample type than by factors such as read length, flow cell age, or the number of active pores at the start of a sequencing run. Techniques such as flow cell washing and adaptive sampling (22) have been demonstrated to enhance output and improve sequencing yield. Additionally, we offer practical advice for optimizing library preparation, particularly for obtaining long reads and adjusting loading amounts when working with small sample sizes. We also address the complexities of data analysis, providing guidelines for constructing customized bioinformatics pipelines that align with specific experimental goals. Effective methods for calling RNA and DNA modifications, as well as strategies for normalizing raw signal data, are also discussed. By bridging the gap between experimental data and best practices, this review equips researchers with actionable strategies for optimizing nanopore sequencing experiments.

BACKGROUND ON NANOPORE HARDWARE

At the heart of nanopore sequencing is a protein channel (nanopore) through which DNA or RNA molecules can pass (Fig. 1). These nanopores are sensitive to electrical changes as molecules move through them (23). Flow cells are filled with an electrolyte solution that establishes in absence of DNA or RNA a relatively constant electrical current through the nanopore. The sensor continuously records this current as a raw electrical signal in picoamperes (pA) (24, 25, 26). During library preparation of DNA/RNA, a motor protein is attached to the nucleotide strand, guiding the DNA or RNA to the nanopore and then unwinding it while controlling the pace at which the strand passes through, ensuring accurate sequencing (2). As each DNA or RNA molecule passes through a nanopore, it disrupts the electrical current in a unique way based on its sequence at a time (9 nt in the nanopore serve for the signal in flowcell R10 and 5 nt for the discontinued flowcell R9) (14, 27). The recorded electrical signal of the sensor, often referred to as the "squiggle", transmits the data for downstream processing (10).

Each flow cell contains thousands of nanopores, which are grouped in channels of 4 pores. A standard MinION flow cell has 2 048 pores organized in 512 channels (2). Flow cells are the standard, full-sized sequencing units used in devices like MinION, GridION, and PromethION, which generate tens to hundreds of gigabases (Gb) of data (28).

Flow cells differ in the design of the nanopores, which impacts their performance mainly in terms of sequencing accuracy (29). In this work, we use R9 and R10 flow cells using single-constriction pores which were designed to be

---

[1]https://community.nanoporetech.com

robust, reliable, and capable of handling a wide range of sequencing tasks.

The R10 flow cells feature double-constriction pores, where the nucleotide strand passes through two constriction points instead of one. The double-constriction design enhances the resolution of basecalling and can achieve higher raw sequencing accuracy (30).

A flow cell check can be performed manually in the `MinKNOW` software to assess the number of nanopores available in the flow cell before a sequencing run. At the start of each sequencing run an initial flow cell check is performed providing the starting number of available pores (31, 32). Each pore can only process a limited number of molecules before it becomes inactive. Therefore the number of available pores decreases over time (30, 31). Usually even a new flow cell never has the maximum number of pores available. A warranty is provided by ONT that allows flow cells to be returned if they do not have enough available pores within three months of purchase[2].

The channels in the flow cell are multiplexed (mux), meaning that only a subset of nanopores can be actively used for sequencing at any given time. A mux scan identifies the most efficient nanopores on a flow cell by testing multiple groups of nanopores (mux groups) in different channels (33, 34). One pore per channel is selected every 1.5 h to continue sequencing within the channel, whereby the current pore can also be re-selected directly. The interval of 1.5 h is the default, but the time can be set in the `MinKNOW` software at the start of each sequencing run. This process helps to optimize the sequencing run by ensuring only high-quality pores are actively sequencing and increase the overall lifetime of the flow cell.

For more details we refer to the nanopore documentation[3].

DATA CHARACTERIZATION AND DESCRIPTION

This perspective draws its conclusions from an extensive and diverse dataset, encompassing over 300 nanopore sequencing runs Among these, a specific focus is placed on about 241 sequencing runs conducted using R9 and R10 MinION and GridION flow cells. This dataset offers robust insights into sequencing performance across various conditions. The study also acknowledges instances of sequencing failures and runs conducted on earlier flow cell types (e.g., flongles or R8 flow cells), although these are not included in the accompanying statistical analyses.

The 241 sequencing runs we focus on (see Tab. S1) include 207 DNA sequencing runs and 34 direct RNA sequencing runs. These runs span a wide range of organisms, including 33 virus, 45 bacteria, 5 protist, 2 plant, 47 insect, 17 mouse, 38 human, 51 metagenomic, and 3 synthetic samples. The amount of input material loaded onto the flow cells ranged from 42 ng to 1440 ng.

The output of sequencing runs was assessed based on several metrics: (1) **EB12** As the sequencing yield, defined as the total amount of sequenced bases, depends on the runtime

Figure 1. Nanopore sequencing on a MinION. A MinION flow cell contains 512 **channels** with 4 nanopores in each channel, resulting in 2 048 nanopores available for sequencing DNA or RNA. Each sensor well (90 μm deep) holds a single nanopore and a platinum electrode (at the base of the well). The electrode is controlled via the **PCB** (Printed Circuit Board) and measures the ion flow as **ionic current**. During sequencing, a **motor protein** guides the DNA/RNA through the nanopore following the electric gradient. The DNA/RNA disrupts the current by partially blocking the ion flow. This can be measured in a resulting **squiggle**.

of the flow cell, we decided to compare the total number of bases produced after 12 h We chose 12 h because some runs included in this study were stopped after that time to wash the flowcell. Referred to as EB12 in this study and estimated by `MinKNOW`, EB12 ranged from $7 \cdot 10^6$ to $28 \cdot 10^9$ nucleotides. For washed flow cells, only the data from the first run was included in the analysis of EB12. (2) The **median read length** calculated after basecalling varied significantly, ranging from 419 to 23 746 bases. We chose to calculate the median instead of the mean, as it is robust against outliers. (3) Another key metric was the **pore half time** of the flow cells, defined as the duration of sequencing until only half of the initially active pores remained active. This value ranged from 1 to 59 hours, highlighting variations in flow cell durability and performance across experiments. Due to artifacts observed in some flow cells, we manually re-evaluated the pore half time of 22 flow cells, as shown in Tab. S3. (4) Other flow cell parameters evaluated in this study include the number of active channels, active pores, and the age of the flow cells. The number of **active channels**, as reported in ONT run reports, ranged from 93 to 512 out of a possible 512 channels. (5) Similarly, the number of **active pores** varied between 749 and 1 984 out of a total of 2 048 available pores. As the number of active pores can increase shortly after starting sequencing, we used the maximum number of active pores during the first 10 minutes of each run. (6) The **flow cell age** was another important variable considered. This was calculated from the time of their arrival at the laboratory, as the production dates were not accessible. We analyzed flow cells that were used between 2 days and 249 days after their arrival, with ONT's

---

[2]https://nanoporetech.com/document/flow-cell-check
[3]https://nanoporetech.com/document/hardware

recommended maximum shelf life being 90 days when stored at 2–8°C. (7) Finally, we elaborated how the amount of loaded library (in ng) affects sequencing yield.

The following analyses provide deeper insights into flow cell performance and durability and have been conducted using custom scripts.

CHALLENGES IN LIBRARY PREPARATION FOR ONT SEQUENCING

Library preparation for ONT sequencing involves multiple steps. For instance, DNA library preparation using the Ligation Sequencing DNA V14 kit (SQK-LSK114) entails DNA repair and end-preparation, followed by adapter ligation and clean-up, before priming and loading the prepared library onto the flow cell. An alternative, if short preparation times are a main focus, is the Rapid Sequencing kit V14 (SQK-RAD114) which reduces the preparation time to as little as 10 min at the cost of sequencing accuracy. For DNA, multiplexing is possible (35) using the Native Barcoding Kit 96 V14 (SQK-NBD114.96).

Similarly, library preparation for direct RNA sequencing (SQK-RNA004) begins with adapter ligation of the reverse transcription (RTA), followed by a RT reaction to stabilize the single RNA strand, and concludes with the ligation of the RNA Ligation Adapter (RLA), priming, and flow cell loading. Each stage of these protocols requires multiple handling and washing steps such as pipetting, mixing, and centrifugation, which introduce challenges in maintaining nucleic acid length and achieving consistent yields.

Obtaining long reads

To obtain the longest possible DNA or RNA reads, it is crucial to minimize any shearing of nucleic acids.

To avoid shearing during library preparation, it is important to reduce the suction force when pipetting. This can be minimized by using cut-tips and pipetting slowly, as recommended by Prall et al. (36). Additionally, vortexing is considered too harsh for nucleic acids, as it often results in shorter fragments, so it is preferable to use gentle tapping for mixing instead (37).

Many DNA/RNA isolation kits, particularly those using beat-beating- or column-based methods, are known to shear and fragment nucleic acids (38, 39). We recommend the classic phenol-chloroform extraction method, as it preserves longer fragments. Recently, there has been a growing interest in high molecular weight DNA extraction techniques. Innovations in this area include the surface topography of silica lamellae and new solid-phase methods introduced by different brands, which aim to improve yield and ease of handling. These kits are often more user-friendly but typically come at a higher reagent cost (40, 41).

In our analysis of DNA and RNA sequencing reads, we mainly used slow and gentle phenol-chloroform extraction to avoid nucleic acid shearing and to maximize read length. We achieved notably long reads in both DNA and RNA samples. The longest DNA read obtained was 1.4 Mb, which came from a metagenomic sample (sample ID: 117.1) and was classified as *Homo sapiens* using `kraken2` (42), with a GC content of 41.4 %. While nanopore sequencing is generally more robust in handling GC-rich regions than PCR-dependent platforms, high GC content can still affect read quality, base-calling accuracy, and sequencing coverage (43, 44). For RNA, the longest read was 175 kb from sample ID 103.1, which classified as bacterial with a GC content of 68.6 %.

Adjustment of loading amount

Loading dependent on pore availability. While it may seem intuitive to increase the amount of DNA/RNA loaded to enhance data yield, there is a threshold beyond which additional loading may not yield proportional benefits[4]. This is particularly true due to saturation effects caused by a limited number of pores on the flow cell. The quantity of data generated is directly proportional to the initial number of active pores (Fig. 2A) and channels (Fig. 2B). When working with flow cells that initially have fewer pores available, it is advisable to load less sample material. Our experiments demonstrated that for flow cells with at least 1 500 active pores, a DNA load of 350 ng produced reasonable sequencing output. For flow cells with approximately 1 200 active pores, we made experience with 300 ng of DNA, and for those with around 800 active pores, we loaded about 250 ng, as detailed in Tab. S1. However, the amount of loaded library does not correle (R^2 - 0.08) with the resulting EB12 for DNA (see Fig. 2C). A relatively weak trend towards higher EB12 is observed when loading more starting material for RNA (R^2 - 0.20).

Note that R10 flow cells exhibit similar behavior in this regard as the discontinued R9 flow cells.

Instead, having extra input material available is advantageous for repeated washing of the flow cell, as it helps recover nanopores and extend overall sequencing yield. Specifically, when planning to wash a flow cell, we recommend splitting the library for different sequencing runs in advance, see details in Sec. Washing and reusing flow cells. **Minimal DNA loads for flow cells** For optimal results in DNA sequencing, ONT recommends to start the library preparation with 1000 ng (equivalent to 100–200 fmol) of total DNA (SQK-LSK114 DNA library preparation kit[5]). Following library preparation, approx. 50 % of the DNA is typically available for loading into the flow cell.

For samples with limited DNA quantities that cannot be amplified, the following methods have yielded successful results: Although ONT's instruction recommend a minimum of 100 ng[6], studies show that without special treatment, 1 ng of DNA (prior to library preparation) yielded 6,118 reads with an N50 of 3,907 bases (45). This result indicates good quality; however, reduced input correlates with decreased output. Carrier sequencing allows for the sequencing of very small amounts of target DNA by mixing it with a larger quantity of non-target "carrier" DNA, ensuring adequate representation without amplification. This technique successfully detected as little as 0.2 ng of target *Bacillus subtilis* DNA combined with 1 000 ng of carrier Lambda DNA, producing high-quality data without amplification (46).

---

[4]https://nanoporetech.com/document/chemistry-technical-document
[5]https://community.nanoporetech.com/attachments/10325/download
[6]https://nanoporetech.com/document/requirements/lig-seq-input-load

Figure 2. (A) The number of active pores (maximum number of active pores during the first 10 minutes) at the beginning of the sequencing run seem to influence EB12. Note the generally lower number of active pores for RNA runs. **(B)** The number of active channels at the beginning of the sequencing run has a weaker influence on EB12. Although a high yield is only achievable when many channels are active, starting with many active channels does not automatically guarantee a high EB12. R^2 values were calculated for second degree polynomial regression. For DNA a correlation can be seen between estimated bases after 12 h and amount of pores and channels at the sequencing start, respectively. For RNA, a relatively weak trend can be seen only for the amount of active pores. **(C)** For DNA, the amount of estimated bases after 12 h is independent of the amount of starting material loaded onto the flow cell (R^2 = 0.08). For RNA, a relatively weak trend shows increased amount of estimated bases after 12 h with increased amount of starting material loaded onto the flow cell (R^2 = 0.20).

The 10X Genomics Chromium Controller is an advanced microfluidics-based platform that enables high-throughput single-cell and spatial analysis of DNA, RNA, and proteins. DNA extracted from the Chromium Controller can be sequenced on nanopore sequencers enhancing the analysis of low-input samples (47). The technology enables sequencing of genomic DNA quantities as low as 50 pg.

In our hands, when loading as low as 30 ng DNA onto a flow cell (435 ng before library preparation, ID 67.6) we successfully got 648 Mb data. In another run we loaded 50 ng onto the flow cell (ID 129.2), which resulted in even 1.5 Gb data output. Simon *et al.* reported the successful use of 50 ng in metagenomic samples with eight species (48).

Minimal RNA load for flow cells. RNA sequencing requires poly-adenylated 3' ends (poly-A tails) that serve as consistent binding sites for ligating adapter molecules to create a continuous RNA-adapter complex that can pass through the nanopore, see Sec. Adapter ligation. Ribosomal RNA (rRNA), which lacks poly-A tails, makes up the majority of cellular RNA; for instance, *E. coli* cells contain around 85 % rRNA (49). Ribosomal RNA is naturally excluded when using a poly-A-based sequencing approach. Usually, this is intended, given that rRNA sequencing is often undesired. For optimal RNA sequencing, ONT recommends to use 500 ng of total RNA or 50 ng of poly-A-tailed RNA (which is in agreement with the ratio of rRNAs in the cell).

Many important RNA molecules, including certain mRNAs (e.g., histone-coding) and most polymerase III transcripts (like ncRNAs, non-canonical miRNAs, and lncRNAs), lack poly-A tails (50, 51). To sequence all RNAs except rRNAs, it is recommended to ligate poly-A tails to the sample and perform an rRNA depletion step prior to poly-A ligation (52). A method for total RNA sequencing with only 10 ng input material has been reported (53). In our hands, 42 ng RNA (ID 152.1) produced 157 Mb of data.

Using external control sequences

The use of external control sequences is a critical aspect of controlling library preparation and benchmarking the sequencing process in nanopore workflows. Both, the DNA

Figure 3. Structure of the ONT DNA spike-in: The positive control spike-in for DNA sequencing is a 3.6 kb amplicon, which maps to the 3' end of the Lambda phage genome.

ligation sequencing kit (SQK-LSK114) and the direct RNA sequencing kit (SQK-RNA004) contain positive control strands (spike-ins). The DNA Control Strand (DCS, Fig. 3) is a 3.6 kb standard amplicon that maps to the 3' end of the Lambda virus genome. It is provided at a concentration of 10 ng/μL.[7]

The RNA Calibrate Strand (RCS) is derived from the *Saccharomyces* genome, specifically the Enolase II gene (ENO2, YHR174W). It is supplied at a concentration of 15 ng/μL[8]. Both DCS and RCS are utilized as positive controls for the library preparation and sequencing process; and therefore are not used for normalization of sequencing results. In our experiments, we detected approx. 1 % of DCS/RCS sequences in our samples.

Adapter ligation

Adapter ligation, the final library preparation step for nanopore sequencing, attaches adapters with motor proteins to sample molecules. These adapters guide DNA or RNA into the nanopore and regulate sequencing speed, making ligation efficiency crucial for optimal output. The concentration of the adapters in the mixes is proprietary; however, it is standardized for reactions involving approx. 100-200 fmol (approx. 1 μg

---

[7]according to ONT Live support asking "What is DCS?", https://nanoporetech.com, 12.12.2024, see Fig. S1

[8]according to ONT Live support asking "What is RCS?", https://nanoporetech.com, 12.12.2024, see Fig. S2

of DNA[9]), 50 ng for poly(A)-tailed RNA, and 500 ng for total RNA. Although the impact of RNA modifications on adapter ligation efficiency is not yet well investigated, these modifications remain an essential factor to consider for RNA samples (8).

New England Biolabs (NEB) generally recommends maintaining a 10:1 adapter-to-sample ratio to ensure efficient ligation without excessive free adapters[10]. According to ONT, high adapter-to-sample proportions generally do not hinder sequencing as long as the majority of pores are actively sequencing[11]

Choosing between LFB/SFB buffer

The buffer choice in the final library prep step determines DNA fragment retention. Long Fragment Buffer (LFB) enriches fragments ≥3 kb for long-read sequencing, while Short Fragment Buffer (SFB) retains a broad size range, affecting sequencing yield and flow cell performance. Our observations indicated that using SFB during the library preparation resulted in more sequencing output (EB12) when compared to the use of LFB (Fig. 4), which also applies to the estimated number of sequenced bases after 24 h and 36 h (see Fig. S3). These differences are evident when analyzing both flow cell types, R9 and R10, together. However, when analyzing R10 flow cells only, the difference between the buffer types was not as pronounced, Fig. S8. In the case of R9 data only, a slightly larger difference emerged between the outcomes of LFB and SFB applications, Fig. S9. These findings suggest that buffer choice may have a variable impact depending on the flow cell generation, potentially due to differences in pore design and chemistry.

Figure 4. When combining R9 and R10 data, there are significant differences between the used buffers. As expected, RNA runs show significantly different amount of estimated bases after 12 h, but also the usage of LFB or SFB results in significantly different amount of sequenced bases after 12 h. For separate plots see Supplement Fig. S9 and Fig. S8. The finding remains unchanged for estimated bases after 24 h and 36 h, see Fig. S3.

---

[9]Note: 1 μg DNA of 8 kb fragments are 200 fmol; 5 kb are about 300 fmol; 50 kb are 30 fmol

[10]https://international.neb.com/protocols/2012/11/01/protocol-for-use-with-end-user-supplied-primers-and-adaptors-e6000

[11]https://nanoporetech.com/document/experiment-companion-minknow "Chapter 43. Troubleshooting your run from the pore activity plots"

Amplicon sequencing

Amplicon sequencing is one common strategy to enrich low-concentrated nucleic acid target material from complex samples for subsequent sequencing. Amplicon sequencing enriches specific DNA regions by amplifying them via PCR with specific primers. The resulting PCR products, or "amplicons," are then sequenced to provide high-resolution information on the targeted regions. A key example of amplicon sequencing is its use in viruses, such as SARS-CoV-2 during COVID-19, for cost-effective sequencing and genomic surveillance. (54). ONT has specific advantages in sequencing long amplicons without additional fragmentation, *e.g.* to retain information of co-occurring mutations on single amplicons supporting virus variant deconvolution and lineage assignment (55).

In nanopore sequencing, amplicons should be mixed in equal nanomolar concentrations (56, 57, 58). When working with amplicons of the same size, mass concentration (nanograms per microliter) can be used instead of molar concentration. Due to the specific target enrichment, amplicon sequencing can ensure sufficient sequencing depth to reliably detect genetic mutations, supporting detailed genomic analysis in fields requiring accurate variant detection.

Amplicon sequencing can be an effective approach to explore entire virus genomes with low viral titers. While a lot of work was done on SARS-CoV-2, amplicon designs are increasingly being explored for other viruses, such as PPV. High-quality assemblies require overlapping amplicons, as accurate strain identification is crucial for diverse pathogens (see Fig. 5A). In some protocols, however, such as the widely used `ARTIC` protocol[12] for SARS-CoV-2 sequencing, smaller amplicons of 392 bp are employed, with an overlap of 90 bp. In our study of SARS-CoV-2, we achieved high-quality results using longer amplicons of approx. 1200 bp, with an overlap of 117 bp between consecutive fragments (59). We demonstrate amplicon sequencing for Plum pox virus (PPV), which exhibits substantial genetic diversity across strains (Fig. 5A). We gained a uniform distribution of coverage across the entire genome (see Fig. 5B), demonstrating the effectiveness of well-designed amplicons for comprehensive genomic representation. Monitoring amplicon coverage is essential, as the aforementioned diversity may result in poor amplicon performance due to insufficient primer binding in mutated regions.

Direct RNA Sequencing

The direct RNA sequencing protocol provides an efficient approach for sequencing RNA molecules directly without the need for converting them into complementary DNA (cDNA). This method allows for sequencing RNA in its native form, preserving RNA modifications, without amplification, and potentially capturing full-length RNA transcripts. An optional library preparation step, binding the RNA strand to a synthesized, complementary DNA strand, can increase throughput and improve read quality by unfolding complex RNA structures, which aids in producing more consistent

---

[12]https://github.com/artic-network/artic-ncov2019

Figure 5. A Genetic diversity for each position in the plum pox virus (PPV) genome across 109 PPV isolates of all known PPV strains reduced in the P3 gene region and enhanced in the genetic region encoding for the capsid protein (CP). **B** Coverage comparison of three sequencing methods: direct RNA sequencing using nanopore (green), Illumina shot-gun RNA-Seq (blue) and nanopore-based PCR amplicon cDNA sequencing (pink) at the example of PPV. A characteristic feature of amplicon sequencing is the clear visibility of higher coverage regions due to the overlapping amplicon design and specific enrichment

sequencing data. However, this approach has downsides: (i) Prolonged incubation of RNA at high temperatures with divalent cations during cDNA synthesis can cause RNA degradation; (ii) Common reverse transcriptases may cleave RNA strands. This can reduce the availability of full-length RNA and impacting the efficiency and accuracy of direct RNA sequencing.

A distinctive feature of ONT's direct RNA sequencing is its ability to detect RNA modifications. However, the bioinformatics analysis of these modifications is still underdeveloped. In this work, we focus intensively on this aspect, see Sec. Hidden treassures in raw data.

TECHNICAL CHALLENGES IN NANOPORE SEQUENCING

Maximizing sequencing yield

Sequencing yield, also referred to as throughput, represents the total amount of data generated during a sequencing run. According to Wang *et al.* (2), the expected sequencing yield of a flow cell primarily depends on three factors: the number of active nanopores, the DNA/RNA translocation speed through the nanopore, and the running time. If the translocation speed is too fast, the system may struggle to differentiate between nucleotides, resulting in inaccurate sequencing. Conversely, if the translocation speed is too slow, throughput is reduced. The translocation speed is largely dependent on the motor protein used. For DNA, it typically operates at around 400-450 nt per second, while for RNA, the speed is slower, at approx. 70 nt per second (2). This speed can be monitored in real-time if live basecalling is enabled, although it can only be

controlled indirectly. For example, increasing the temperature may accelerate the translocation speed, whereas overloading the flow cell might lead to a reduction in speed.

To analyze sequencing yield we explore over 300 ONT sequencing experiments across a variety of species, see Sec. Data characterization and description, Tab. S1. We observed significant variation in sequencing yield and investigate potential reasons for this yield disparity (Fig. 2, Fig. 6). We first investigate whether EB12 is influenced by the pore half time, which refers to the duration after which half of the nanopores in a sequencing run become inactive. A strong correlation is observed between pore half time and EB12 across all sample types except bacteria (Fig. 6A), consistent for both R9 and R10 flow cells This correlation remains significant even when normalizing EB12 for the number of starting pores, as shown in Fig. S4. As expected, a slower degradation rate of the flow cell leads to a higher sequencing yield.

Next, we analyze the impact of sample type on EB12. DNA sequencing yields in our data are generally higher than the ones for RNA, as shown in Fig. 6B, likely due to increased translocation speed and higher number of starting pores. Although DNA mouse samples have a slightly higher EB12 than other sample types with p-values of 0.04 or below (except p = 0.067 for viruses), see Fig. S5, it appears that after 12 hours of sequencing, the sequencing yield is roughly similar, regardless of the sample type.

Contrary, the pore half time seems to vary significantly depending on the type of sample being sequenced (Fig. 6C, with a median pairwise p-value of 0.011, see Fig. S6). For DNA samples, for instance, we found that insect and protist samples tend to have a much shorter pore half time than mouse, metagenomic, or bacterial samples. Furthermore, flow cells used for bacterial, metagenomic, and human samples exhibited high variability in pore half time. The flow cell pore half time appears to be influenced not only by RNA/DNA characteristics but also by the sample type. We assume these differences might originate from factors such as the quality and purity of RNA/DNA, the extraction method, the sequencing kit used or the cell type composition. Further research is needed to investigate those factors.

Despite variations in pore half time, the distribution of EB12 remains relatively consistent across different sample types (Fig. 6B). Notably, insects and protists, which exhibit shorter pore half time, do not stand out as outliers regarding EB12. When comparing R9 to R10, as shown in Fig. S10, there are no significant differences across all sample types. However, a significant reduction ($p < 2.4 \cdot 10^{-7}$)in both EB12 and pore half time is observed for R10 when analyzing metagenomic samples specifically. Taken together, after 12 hours, the sequencing output remains relatively consistent regardless of sample type, pore count and flow cell type, but the sample type seems to influences the degradation rate of the flow cell.

The median read length had no impact on EB12, see Fig. 6D, nor the amount of estimated bases after 24 h and 36 h, see Fig. S7. Most runs had a median read length below 3 kb, with EB12 values ranging from 0 to nearly 8 Gb and showing a fairly even distribution. Runs with a median read length exceeding 3 kb tended to have higher EB12 values,

Figure 6. Maximizing sequencing yield. Sequencing yield from R9 (dots) and R10 (triangle) for DNA (orange) and RNA (blue) samples from a variety of species. Differences between RNA and DNA are evident in all plots, with a lower sequencing yield for RNA in general. Note, that the lower RNA yield is dependent on the slower sequencing speed (2) and the generally low number of active starting pores during RNA runs. **(A)** Correlation (R^2) between pore half time and the estimated number of bases after 12 hours of sequencing (EB12) is observed across most sample types. No significant correlation is evident for bacteria due to high variability. Regarding R9 and R10 data separately, the correlation for bacteria slightly increases (R9: $R^2 = 0.39$, R10: $R^2 = 0.11$), but still remains weaker than for the other sample types. For the other sample types the correlation does not change when separating R9 and R10 flow cell data. **(B)** The EB12 yield is consistent across all sample types and flow cell type. **(C)** In contrast to EB12, the distribution of pore half time differs strongly between the sample types. Insect samples have a lower pore half time than other samples. Metagenomic and bacteria samples show a high variance. Data shown in B and C indicate differences between R9 and R10 data for metagenomic samples, see Fig. S10. **(D)** The median read length does not influence EB12 (nor the amount of estimated bases after 24 h and 36 h, see Fig. S7) for DNA. For RNA, a weak positive correlation is visible, which is slightly stronger after 24 h and 36 h, see Fig. S7. But note that there are only a few data points for RNA after 24 h and 36 h, and the amount of starting pores for RNA was generally lower. **(E)** The age of the flow cell (number of days from arrival to usage) does not affect EB12, although our data for R10 flow cells are limited.

highlighting a clear distinction between RNA and DNA. R10 median read lengths aligned with those of R9 data.

We then turned our attention to the properties of the flow cells themselves. The age of the flow cell (the time between delivery and use) did not significantly affect EB12. Surprisingly, flow cells over 200 days old (Fig. 6E) performed similar to younger ones, although exceeding the ONT warranty period of three months by a factor of two. The oldest flow cell we tested was 249 days old, performing normally (1080 pores, 3.1 Gb output), see Tab. S1, ID 172.1.

In summary, a high EB12 is only achievable when many channels/pores are active at the beginning of the sequencing run (Fig. 2A,B). However, starting with many active channels does not automatically guarantee a high EB12. On the other hand, as expected, the number of active pores at the beginning of a sequencing run influence EB12, particularly when sequencing DNA (Fig. 2A).

Washing and reusing flow cells

Flow cells naturally degrade with use as active pores decrease, often due to blocking during sequencing. Blocked pores shift from "single pore" to "unavailable" in `MinKNOW`. While ONT acknowledges pore blocking, reasons for its variability remain

Figure 7. Influence of washing the flow cell on the amount of active pores. **(A)** A flowcell which has been paused (blue bars) two times for washing. **(B)** A flowcell which has been stopped (blue bars) five times for washing.

unclear. Pores can be unblocked using voltage reversal during mux scans or by washing the flow cell.

Washing is performed with ONT's "Flow Cell Wash Kit" (EXP-WSH003, EXP-WSH004), which uses DNase I to digest and remove loaded genetic material, freeing blocked pores. The cost of washing is minimal, approx. €16 per wash step, as the Wash Kit EXP-WSH004 costs €95 for six wash steps[13]. The sequencing run can be paused or stopped before washing: pausing allows continuation in the same file, while stopping generates a new file for subsequent sequencing. After washing, the flow cell can be reused for sequencing a new sample, but the previous run must be stopped rather than paused. Washed flow cells can also be stored for future use. However, they will not perform as well as new ones; the number of active pores will be lower than initially but higher than before washing (see Fig. 7).

Washing can also be employed during a single sequencing run to increase sequencing yield by reactivating blocked pores. We identified three key factors to consider before sequencing: (i) potential washing steps, (ii) optimal washing timing, and (iii) available sample material. We investigated the impact of washing flow cells on pore recovery by analyzing 34 flow cells that had undergone 1-6 washes, totaling 58 washing steps, see Tab. S1. Examplarily, Fig. 7 shows two such washing steps. Statistics of these two flow cells are displayed in Tab. S4. In Fig. 7A, human DNA sample (ID 111.1) was washed twice during sequencing (at 10 h and 30 h), pausing the run (blue bars). Active pores decreased over time (red arrows) but increased after each wash (blue arrows). In Fig. 7B, the cricket DNA sample (ID 124.1–124.6) was washed five times during sequencing. A mean loss of 84.2 % of active pores could be compensated by revoking on average 85.6 % of those pores. ONT recommends washing and reusing a flow cell up to four times. Our findings show that flow cells can be washed at least five times and still produces data.

A key factor for high sequencing yield is determining the appropriate time point for washing, as yield depends on the number of active pores, Fig. 2A. It seems that pores do not block faster after washing. Therefore, we recommend monitoring cumulative output in MinKNOW: when the curve flattens (Fig. 7A, green line), we advise washing.

To ensure sufficient material for reloading, we recommend splitting the sample before starting the sequencing run if washing is planned to increase yield. For instance, when planning two wash steps, we recommend allocating 50 %, 30 %, and 20 % of the total library for the initial run and subsequent washes, respectively. This decreasing allocation accounts for the reduction in active pores after each wash. Adding excess material beyond the capacity of available pores will not further increase sequencing yield (see Fig. 2C and Fig. 7)

Adaptive sampling

Adaptive sampling allows for real-time selection or rejection of DNA fragments based on their sequence content, enabling targeted enrichment or depletion of specific genomic regions, organisms, or sequence types during a sequencing run[14] (60). For example, adaptive sampling can enrich gene panels or exomes (61), eliminate unwanted sequences such as highly repetitive regions or contaminants, and target specific organisms or deplete host in metagenomic studies (22, 62, 63). This approach enhances coverage of targeted genes or regions without requiring complex library preparation steps like amplification or hybridization probes.

As DNA molecules pass through a nanopore, the device generates real-time sequence data (Fig. 8A). This requires a GPU and a fast basecalling model for adaptive sampling. The system quickly compares each fragment's partial sequence (approx. 400 bases, corresponds to one second of data) to a reference genome or user-defined targets. When enriching for regions of interest, the sequencing continues only if the sequence matches the region of interest; otherwise, the DNA fragment is ejected, conserving flow cell resources and improving yield for desired regions. The depletion mode works vice versa.

As the acceptance or rejection of a read is based solely on its first chunk, strands may be rejected, if the region of interest is not located at the beginning of the read. To address this, the target region of interest should be extended by adding a "buffer" on both sites in the configuration file to accommodate sequencing in either direction, as sequencing can initiate from either DNA strand (Fig. 8B). Thus, for adaptive sampling, buffer describes an amount of nucleotides, by which the region

---

[13]23 October 2024

[14]https://nanoporetech.com/document/adaptive-sampling

Figure 8. **(A)** The basic principle of adaptive sampling: For each read, the first ∼400 bases are mapped to a given reference, and sequencing continues only if they match a region of interest (22). About 450 bases are sequenced per second for R9 nanopore flow cells (2). Rejecting a read takes about 0.5 seconds, with additional time required to capture the next read. **(B)** In order not to miss the reads where the region of interest (green region) is not at the beginning of the read, the target region should be defined with an extended buffer (e.g. 2000 nt) on both sides (pink regions) in the configuration .bed file. **(C)** Adaptive sampling was used to enrich over 30,000 CpG islands (CGIs) to increase their coverage. To increase enrichment, each CGI region of interested was extended by a 500 nt buffer region (blue dots) and by a 2000 nt buffer region (pink stars) on both sides.

of interest is extended in the bed file (a text file format used to store genomic regions).

Using adaptive sampling may yield a higher number of short reads (reducing N50 and median read length), and necessitating re-basecalling with a higher accuracy model post-sequencing to enhance data quality.

We have applied adaptive sampling to enrich CpG islands in the human genome, see Tab. S5. We targeted approx. 30 000 CpG islands, with an average length of 777 nt (IDs 153.1–178.1). We applied adaptive sampling without adding a buffer region, with a 500 nt buffer region, and a 2 000 nt buffer region on both sides of the target regions. We achieved up to 17 X mean sequencing depth on CpG islands, representing a fourfold increase compared to the mean sequencing depth of the whole genome, when applying 2 000 nt buffer. According to ONT, the theoretical maximum increase in sequencing depth achievable through adaptive sampling is approximately 5–10-fold[15]. In summary, adaptive sampling can effectively enhance sequencing depth in specific genomic regions or target species, albeit with increased time requirements for post-processing, such as re-basecalling and filtering out short, unwanted reads.

CHALLENGES IN ANALYZING ONT DATA

Assembling an ONT pipeline

Bioinformatics analysis of ONT sequencing data is based on modular pipelines, that can be assembled and utilized effectively, even by users with limited computational

Figure 9. The standard pipeline for nanopore sequencing data involves basecalling of the raw data, followed by quality control and filtering. The reads can then be analyzed with specialized long-read tools for mapping, polished assembly, or other workflows.

backgrounds (Fig. 9). Instead of relying solely on standard, one-size-fits-all pipelines, it is critical to carefully select tools that best align with specific experimental goals and data characteristics.

A well-constructed ONT analysis pipeline includes several steps, which can be addressed by different tools:

Basecalling is the transformation of raw signal data, or 'squiggles,' into the original nucleotide sequences using neural networks. Typically, the basecalling tools have a pre-processing step, such as removing the adaptors and normalizing individual signals. The following three tools are all provided by ONT. Currently, Dorado[16] is the state-of-the-art software for basecalling on ONT platforms, that supports the older fast5 and the new pod5 format, which is a

---

more compact format enabling faster and more efficient file access. `Dorado` includes basecalling models for all ONT kits, predicts modified bases, and built-in demultiplexing capabilities for barcode detection from ONT barcoding kits. `Bonito`[17] is an alternative tool that allows users to train their own neural networks for basecalling. `Remora`[18] adds an additional layer, enabling nucleotide modification calling by training neural networks along the standard bases. Alternative tools for nucleotide modifications prediction are discussed in detail below. Basecalling results are typically saved in `fastq` format for subsequent analysis steps.

After basecalling, **quality control** (QC) is essential to evaluate the reliability of the sequencing data and to identify any potential issues in the dataset, such as degradation, contamination, and read length distribution. One of the most comprehensive QC tools for ONT data is `pycoQC`[19] (64), which provides an interactive `html` overview that includes information on read quality, read length, sequence coverage, active channels, and quality over sequencing time. Other QC tools, such as `NanoPlot`[20] (65) and `nanoQC`[21], provide a restricted functionality, but with a different representation of the data.

Following QC, **filtering** is often necessary, especially for experiments requiring specific read quality or length thresholds. Filtering tools help to remove low-quality reads, adapters, barcodes, or reads outside the desired range, increasing the overall quality and relevance of the dataset. `chopper`[22] from the `NanoPack` suite (65) or `Filtlong`[23] are widely used for filtering ONT reads. `Dorado` also provides parameters to trim adapters and barcodes.

Once high-quality reads are obtained, subsequent analyses such as mapping and assembly can be performed using specialized tools designed for long-read data. For read **mapping**, `minimap2`[24] (66) is widely used. `Winnowmap`[25] (67) is an alternative mapping tool that has been specifically optimized for complex genomes. It has demonstrated superior performance when mapping both simulated PacBio and ONT data for human genomes (67). For assembly, `Flye` is one of the most widely used tools, demonstrating good performance in terms of contig length and error rate in bacteria (68). It includes polishing and can be told how many iterations of polishing should be performed. Alternative tools, such as `Shasta`[26] (69), have been shown to perform faster, while `NextDenovo`[27] (70) can generate longer contigs, particularly in the case of mollusks (71). **Error correction** (polishing) can be subsequently utilized to refine the assembly. `Medaka`[28] is a ONT-specific neural networks, trained on typical ONT errors to correct these

errors effectively. `Medaka` has proven particularly effective in reducing deletion errors (68). Additionally, `Racon`[29] (72) is a consensus caller using high quality reads from technologies such as Illumina to reduce single nucleotide variants and insertion errors, enhancing the overall accuracy of the assembled data.

Hidden treassures in raw data

Beyond the standard pipeline of ONT tools, the raw data from ONT sequencing contains a hidden treasure of information. Analyzing this data requires significant self-implementation and detailed knowledge. Here, a demonstration of the fundamentals necessary to fully exploit the potential of ONT by using DNA and RNA modification detection as an example is given. This description will pave the way to complement existing tools for nucleotide modification analysis, which are currently designed for only a small fraction of modifications. To achieve this, we will focus on processing the raw data, which can be divided into five distinct phases: (1) Accessing raw data: in this initial step, the raw signal data from the nanopore sequencer is imported into the computational analysis. This is particularly important because it serves as the foundation for all subsequent analyses. (2) Improving Signal Segmentation: is applied to refine the process of identifying individual signals corresponding to nucleotides. Improving segmentation helps enhance the accuracy of the subsequent basecalling step by ensuring that the software can effectively distinguish between different signal components. (3) DNA Modification Detection: Once the signal data is well-segmented, this step focuses on identifying specific modifications to DNA (e.g., methylation). Detection of DNA modifications is important for understanding epigenetic regulation and gene expression, providing insights into cellular functions and biological processes. (4) RNA Modification Detection: Similar to DNA modification detection, this step involves identifying modifications in RNA sequences. RNA modifications can influence gene expression and stability, making their detection crucial for studying transcriptomics and understanding post-transcriptional regulation. (5) Differential Signal Detection: Alternatively to predicting modification types directly, the differences in signal patterns between conditions or samples can be analyzed to reveal a general change of the nucleotide.

Preprocessing raw data. ONT stores the 'squiggles' as integers (73) to save storage space, which are converted back to pA, see SEqu. **1**. Normalization of raw ONT signals is essential for reliable comparisons within and across sequencing experiments. Signal variations caused by differences in flow cells, nanopores, and sensors can introduce biases (Fig. S11), making normalization (Fig. 10A) critical for accurate and reproducible analyses (SEqu. **2**). Outliers in ONT signals, as in Fig. 10B, often caused by faulty sensor measurements, can distort downstream analyses. To ensure data quality, these outliers should be identified and filtered prior to further signal processing. The ONT signal, comprising a continuous series of data points, may occasionally include multiple adjacent sequencing reads, as shown in Fig. 10C.

[17]https://github.com/nanoporetech/bonito
[18]https://github.com/nanoporetech/remora
[19]https://github.com/a-slide/pycoQC
[20]https://github.com/wdecoster/NanoPlot
[21]https://github.com/wdecoster/nanoQC
[22]https://github.com/wdecoster/chopper
[23]https://github.com/rrwick/Filtlong
[24]https://github.com/lh3/minimap2
[25]https://github.com/marbl/Winnowmap
[26]https://github.com/paoloshasta/shasta
[27]https://github.com/Nextomics/NextDenovo
[28]https://github.com/nanoporetech/medaka

[29]https://github.com/isovic/racon

Figure 10. Overview of Signal Data Preprocessing Steps. **A** To standardize the data, the unnormalized time series (left) are converted (arrow) into the corresponding normalized signal (right, mostly ranging from -4 to 4). **B** Detected outliers (pink circles) indicate points that deviate significantly from the expected signal pattern, likely due to sensor errors. **C** A single signal can contain more than one read. Here indicated with signal containing three recurring components for 'Read 1' and 'Read 2' specific to RNA sequencing: 'adapter,' 'poly A,' and 'transcript.'

Figure 12. Modified nucleotides have their own characteristic signal. Machine learning models can detect this signal directly or by comparing it to the unmodified counterpart. The red and blue lines represent the mean signal of 200 reads respectively, while the vertical bulges indicate the signal distribution per base following a normal distribution. Modifications can lead to error patterns in the reads that can be detected by tools.

Figure 11. Overview of Signal Processing and Modification Calling. The raw ONT signal can be segmented and corrected for basecalling errors by resquigglers using the raw signal and basecalls. Some resquiggling tools also require a reference sequence and a mapping.

The reason for this phenomenon is unresolved. To ensure accurate and reliable downstream analyses, such signals must be split into individual reads using the coordinates provided by `Dorado`.

The python package `read5`[30] is a wrapper that unifies access to `fast5`, `pod5`, and `slow5` data formats and provides functions for signal conversion (SEqu. **1**) and normalization (SEqu. **2**). Each of these formats has its own API using different function names which often requires to script format-specific code to handle inconsistencies.

Improving signal segmentation. The analysis of raw data plays a crucial role for detecting chemical nucleotide modifications. This is particularly relevant for the vast amount of around 170 RNA modifications for which no basecalling models currently exist. These modifications also produce characteristic signals when translocating through the nanopore. Analyzing raw signals for modification detection requires improved signal segmentation, known as 'resquiggling.' This process involves realigning the raw current signal to the nucleotides, see Fig. 11. This segmentation is crucial because the motor protein that moves the nucleotides through the pore does not operate at a constant speed. The time a nucleotide spends in the pore ('dwell time') varies depending on the specific nucleotides (74).

Several tools have been developed to facilitate the resquiggling process and require generally (i) the raw ONT signal (`fast5`, `slow5` or `pod5`) and (ii) the basecalls generated by the basecalling process (`fastq`). Optionally, (iii) a reference sequence (`fasta`) and (iv) the alignments of the basecalls to the reference (`bam`) can be added. The most widely used tool `f5c` (75) is a GPU-accelerated, multi-threaded re-implementation of `Nanopolish Eventalign`. It offers significant reductions in runtime while maintaining the known output formats. Notably, `f5c` also supports newer pore chemistries, such as R10 and RP4. For instance, a sequencing run of a 32,000 bp Coronavirus genome (ID 13.1) with 1.3 Gb across 857,000 reads produced a 181.5 GB output file using `Nanopolish Eventalign`, which has been reduced to 84 GB with `f5c`. `Tombo` has not been supported by ONT since 2020 and is compatible only with older single-read `fast5` formats. Despite its limitations, `Tombo` remains in use today for specific applications.

Future developments in resquiggling tools are expected to further enhance compatibility with emerging nanopore technologies and data formats.

---

[30] https://github.com/rnajena/read5

Detecting DNA modifications. The direct detection of DNA modifications without amplification or pre-treatment of the DNA is an exciting feature of nanopore sequencing, providing insights into epigenetic regulation and genomic variability. DNA modifications alter the electrical signal as the DNA passes through the nanopore resulting in a differing picoampere signal for k-mers (for R9 k=5; for R10 k=9) containing modified bases compared to those consisting of only unmodified bases, see Fig. 12. Currently, 17 types of DNA modifications have been identified in the genomes of bacteria and eukaryotes (76), with the most well-characterized being methylation of cytosine 5-methylcytosine, 5mC (77); 5-hydroxymethylcytosine, 5hmC; and adenine N6-methyladenine, 6mA (78).

Several computational tools have been developed to identify modifications from nanopore data, with a primary focus on 5mC and 6mA, using different computational approaches, in particular Hidden Markov Models, statistical tests, or neural networks, see Tab. S2. Input data for the methylation calling tools is usually the raw or basecalled `fast5/pod5` files.

DNA modification calling was traditionally performed after basecalling using tools like `Nanopolish` (79) or ONT's `Megalodon`[31]. Currently, ONT's `Dorado`[32] detects CpG methylations directly during basecalling if specified, and provides greater accuracy than tools like `Nanopolish`(80), which has difficulty when nearby CpG sites differ in methylation (79). The performance of modification-detection tools varies by slight over-prediction (e.g. `Nanopolish`) or under-prediction (e.g. `DeepMod`) as shown in several reviews (21, 80, 81, 82, 83) Validation remains essential, although tools like `Nanopolish`, `Megalodon`, `DeepSignal`, and `Dorado` already show high correlation with the gold standard, whole-genome bisulfite sequencing (WGBS) (80, 81). Meta approaches improve modification detection accuracy by combining the outputs of multiple tools. For example `METEORE` combines results from up to six tools using a random forest model, significantly improving accuracy but at the cost of increased runtime (21).

Detecting RNA modifications. The detection of RNA modifications using nanopore sequencing is less developed compared to DNA modifications, possibly because RNA is known to contain over 170 distinct modifications (84). RNA modifications are of utmost importance for RNA function and regulation.

Computational tools exist only for a handful of RNA modifications, see Tab. S2. Beside these downstream analysis tools, ONT's `Dorado` directly enables modification detection by specifying specific basecalling models. Users can further customize the process by manually disabling additional filters as needed. Excitingly, basecallers like `Bonito` and `Remora`, allow researchers to train custom models for detection of modifications of any kind. These models can be created from scratch or fine-tuned from pre-trained versions, offering flexibility for different datasets and research objectives. The models can perform binary classification to predict a certain

modification, or directly call modified bases alongside the standard nucleotide sequence.

Detecting differential signals. Comparing raw ONT signals between different samples provides a powerful approach for detecting differential signals, which can point to mutations, modifications and isotopic labeling (85). The goal of this approach is not necessarily to identify the exact type of modification but rather to detect the presence of a modification. The signal may differ in its measured pA range, including mean, variance, skewness, kurtosis, and dwell time. The calculation of these measures can be applied to full-length nanopore reads. To minimize normalization biases, comparative analyses should ideally be performed using the same flow cell. When comparisons across multiple flow cells are unavoidable, further normalization steps are needed.

On the other hand, for example, in the case of modifications such as deuterium in the DNA/RNA backbone (85), normalization can inadvertently obscure biologically meaningful information. This is particularly true when the signal-to-noise ratio—where the signal represents the modification and the noise stems from instrumental bias—is low. To address these challenges, carefully controlled experimental designs are essential to ensure that observed differences in the nanopore signal reflect genuine modifications rather than artifacts of the sequencing process.

CONCLUSION

In summary, this study has yielded critical insights into refining nanopore sequencing using the MinION platform, with far-reaching implications for research and application. We displayed suggestions for the user of nanopore sequencing in the three categories (1) library preparation tips; (2) technical tips; and (3) analyzing ONT data, see a summary in Fig. 13. We showed general statistics for the number of sequenced bases after 12 h on both R9 and R10 flow cells. EB12 is influenced by the number of pores and channels at the start of the run for DNA, as well as the pore half time, which surprisingly varies depending on the sample type. In contrast, EB12 shows no correlation with the amount of input material, read length, or flow cell age. Additionally, we showed, that the flow cell half-life depends for DNA samples on the buffer used (LFB vs. SFB) and that flow cells can perform well after they have been expired. Meanwhile, we could also verify most of the observations on R10.4 flow cells. However, for the current time point the sample size is too small to make statistically valid statements.

With this publication, correlations and assertions from the community have been statistically substantiated and are now citable for future work.

Here, we aim to describe an additional observation: within the community, it is often posited claimed that theoretically, one could sequence reads of unlimited length if the DNA fragments were arbitrarily long. We extracted fragments of 500,000 nt from algae *Chlamydomonas euryale*, *Chlamydomonas reinhardtii*; however, we encountered significant challenges during the sequencing process. Despite trying various methods, we consistently observed good sequencing performance only in the initial minutes before the sequencing had to be aborted. Regarding this observation, we

---

[31]https://github.com/nanoporetech/megalodon
[32]https://github.com/nanoporetech/dorado

TIPS & TRICKS for ONT sequencing

LIBRARY PREPARATION TIPS

Obtaining long reads:
> do not vortex your samples
> use cut tips and pipette as slowly as possible
> use phenol-chloroform extraction

Optimizing loading amount and dealing with small sample sizes:
> aim for 1,000 ng of DNA or 500 ng of RNA before library preparation
> balance the DNA/RNA load on the flow cell based on active pore count
> once all nanopores are occupied, additional input material has minimal impact on sequencing yield
> avoid loading less than 50 ng of DNA, as yields drop significantly
> for small amounts of DNA, use carrier sequencing with a non-target "carrier" DNA
> when ligating poly-A tails to all RNA species (if interested in sequencing a broader range of RNA), use a depletion method to remove ribosomal RNA (rRNA)

External control sequences:
> incorporate spike-ins for valuable assurance of successful library preparation
> for well understood, standardized workflows, exclude spike-ins to increase output

Adapter ligation:
> for very high molecular weight DNA, use fewer adapters as there are fewer fragment ends to ligate
> RNA modifications may influence the adapter ligation efficiency

Choosing the buffer:
> use Long Fragment Buffer (LFB) when focusing on long-read sequencing
> Short Fragment Buffer (SFB) typically yields higher sequencing output than LFB
> for R10 flow cells, the difference between LFB and SFB is less pronounced

Amplicon sequencing:
> carefully design primers for effective assembly and complete genome coverage
> for high-quality assembly, use 1000 bp amplicons with an overlap of 200 bp
> for pathogens with diverse strains, use the correct strain for primer design

TECHNICAL TIPS

Flow cell quality:
> Ensure a high number of active pores (e.g., by washing) to maximize sequencing yield.
> Flow cells older than 200 days still function well.

Washing:
> good results can be achieved even after five washing steps
> If you do not have enough sample material do not use washing steps in between
> If you have enough sample material, split the library into decreasing portions to load after each washing step (e.g. 50%, 30% and 20%)

Adaptive sampling:
> Use adaptive sampling for enriching or depleting specific genomic regions or organisms
> Extend the region of interest in the configuration file with a buffer region on both sides

ANALYZING ONT DATA

Pipelines:
> build modular pipelines tailored to your specific experimental goals

Working with raw data:
> normalization is essential to eliminate biases, ensuring reliable comparisons
> normalize each read individually

Modification detection:
> use combined tools for high-confidence calls when computational resources allow
> validate results with complementary methods to ensure robust conclusions
> RNA modifications are vastly more diverse than DNA modifications
> custom models can be trained from scratch or fine-tuned from pre-existing ones
> resquiggling is particularly vital for RNA modification detection for accurately assigning signal segments to specific nucleotides or nucleotide groups

Detecting Differential Signals:
> use summary statistics at different levels of granularity to identify differences in nanopore signals
> be cautious of confounding variables, which can overshadow biological signals
> use consistent sample preparation and sequencing protocols to minimize noise

Figure 13. TIPS & TRICKS mentioned within the manuscript are summarized here for a better overview.

currently have a few potential explanations: (1) the molarity of the adaptors with respect to the high molecular weight fragments might have been incorrect; or (2) algae may contain a high amount of polysaccharides that could influence pore activity. As seen for this example many of the observations can still not be entirely explained and call systematic analysis.

ACKNOWLEDGEMENTS

We thank Franziska Hufsky for her valuable assistance in improving the writing of this manuscript and Emanuel Barth for his valuable input on statistical questions. We thank Carolin Dippmann for her experimental work and insightful discussions that contributed to the development of this study and Franziska Aron for her support and help in the lab. We thank Marie Lataretu for her contribution to the ONT DNA spike-in and Adrian Viehweger for his work on figures. We thank Patrick Bohn for discussion about different flow cell types. We thank the Sequencing Core Facility of the Genome Competence Center (MF1) of the Robert Koch Institute (RKI), Germany for providing excellent sequencing services and André Götze from the RKI MF1 DigiLab for outstanding support in obtaining technical metadata information of the associated runs. We thank Matthew Huska and Christian Blumenscheit from MF1 for R10.4.1 data obtained in the context of an RKI special research funds project about adaptive sequencing of a Zymo Mock community. Competing interests: DM and BS are employees of oncgnostics GmbH, a company that aims to commercialize DNA methylation markers. This work was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany´s Excellence Strategy – EXC 2051 – Project-ID 390713860, the German DFG Collaborative Research Centre AquaDiva (CRC 1076 AquaDiva), the German state of Thuringia via the Thüringer Aufbaubank (2021 FGI 0009), the Carl-Zeiss-Stiftung within the program Scientific Breakthroughs in Artificial Intelligence (project "Interactive Inference"), and the Ministry for Economics, Sciences and Digital Society of Thuringia (TMWWDG), under the framework of the Landesprogramm ProDigital (DigLeben-5575/10-9).

Conflict of interest statement. None declared.

REFERENCES

1. Laszlo, A. H., Derrington, I. M., Ross, B. C., Brinkerhoff, H., Adey, A., Nova, I. C., Craig, J. M., Langford, K. W., Samson, J. M., Daza, R., et al. (2014) Decoding long nanopore sequencing reads of natural DNA. *Nature biotechnology,* **32**(8), 829–833.
2. Wang, Y., Zhao, Y., Bollas, A., Wang, Y., and Au, K. F. (2021) Nanopore sequencing technology, bioinformatics and applications. *Nature biotechnology,* **39**(11), 1348–1365.

3. Petersen, L. M., Martin, I. W., Moschetti, W. E., Kershaw, C. M., and Tsongalis, G. J. (2019) Third-generation sequencing in the clinical laboratory: exploring the advantages and challenges of nanopore sequencing. *Journal of clinical microbiology,* **58**(1), e01315–19.

4. Nicholls, S. M., Quick, J. C., Tang, S., and Loman, N. J. (2019) Ultra-deep, long-read nanopore sequencing of mock microbial community standards. *Gigascience,* **8**(5), giz043.

5. Overholt, W. A., Hölzer, M., Geesink, P., Diezel, C., Marz, M., and Küsel, K. (2020) Inclusion of Oxford Nanopore long reads improves all microbial and viral metagenome-assembled genomes from a complex aquifer system. *Environmental microbiology,* **22**(9), 4000–4013.

6. Liu, L., Yang, Y., Deng, Y., and Zhang, T. (2022) Nanopore long-read-only metagenomics enables complete and high-quality genome reconstruction from mock and complex metagenomes. *Microbiome,* **10**(1), 209.

7. Chen, Z., Grim, C. J., Ramachandran, P., and Meng, J. (2024) Advancing metagenome-assembled genome-based pathogen identification: unraveling the power of long-read assembly algorithms in Oxford Nanopore sequencing. *Microbiology Spectrum,* pp. e00117–24.

8. Oikonomopoulos, S., Bayega, A., Fahiminiya, S., Djambazian, H., Berube, P., and Ragoussis, J. (2020) Methodologies for transcript profiling using long-read technologies. *Frontiers in genetics,* **11**, 606.

9. Jain, M., Abu-Shumays, R., Olsen, H. E., and Akeson, M. (2022) Advances in nanopore direct RNA sequencing. *Nature Methods,* **19**(10), 1160–1164.

10. Rang, F. J., Kloosterman, W. P., and de Ridder, J. (2018) From squiggle to basepair: computational approaches for improving nanopore sequencing read accuracy. *Genome biology,* **19**(1), 90.

11. Loose, M., Rakyan, V., Holmes, N., and Payne, A. (2019) Whale watching with BulkVis: A graphical viewer for Oxford Nanopore bulk fast5 files. *Bioinformatics,* **35**(13).

12. Faria, N. R., Sabino, E. C., Nunes, M. R., Alcantara, L. C. J., Loman, N. J., and Pybus, O. G. (2016) Mobile real-time surveillance of Zika virus in Brazil. *Genome medicine,* **8**(1), 1–4.

13. Amarasinghe, S. L., Su, S., Dong, X., Zappia, L., Ritchie, M. E., and Gouil, Q. (2020) Opportunities and challenges in long-read sequencing data analysis. *Genome biology,* **21**(1), 1–16.

14. Jain, M., Olsen, H. E., Paten, B., and Akeson, M. (2016) The Oxford Nanopore MinION: delivery of nanopore sequencing to the genomics community. *Genome biology,* **17**, 1–11.

15. Ferreira, M. R., Carratto, T. M. T., Recalde, T. S. F., Bonadio, R. S., Jain, M., de Oliveira, S. F., Castelli, E. C., and Mendes-Junior, C. T. (2024) Advances in forensic genetics: Exploring the potential of long read sequencing. *Forensic Science International: Genetics,* p. 103156.

16. Sereika, M., Kirkegaard, R. H., Karst, S. M., Michaelsen, T. Y., Sørensen, E. A., Wollenberg, R. D., and Albertsen, M. (2022) Oxford Nanopore R10. 4 long-read sequencing enables the generation of near-finished bacterial genomes from pure cultures and metagenomes without short-read or reference polishing. *Nature methods,* **19**(7), 823–826.

17. Zhao, W., Zeng, W., Pang, B., Luo, M., Peng, Y., Xu, J., Kan, B., Li, Z., and Lu, X. (2023) Oxford nanopore long-read sequencing enables the generation of complete bacterial and plasmid genomes without short-read sequencing. *Frontiers in Microbiology,* **14**, 1179966.

18. Liu-Wei, W., van der Toorn, W., Bohn, P., Hölzer, M., Smyth, R. P., and von Kleist, M. (2024) Sequencing accuracy and systematic errors of nanopore direct RNA sequencing. *BMC genomics,* **25**(1), 528.

19. Magi, A., Semeraro, R., Mingrino, A., Giusti, B., and D'aurizio, R. (2018) Nanopore sequencing data analysis: state of the art, applications and challenges. *Briefings in bioinformatics,* **19**(6), 1256–1272.

20. Leggett, R. M. and Clark, M. D. (2017) A world of opportunities with nanopore sequencing. *Journal of experimental botany,* **68**(20), 5419–5429.

21. Yuen, Z. W.-S., Srivastava, A., Daniel, R., McNevin, D., Jack, C., and Eyras, E. (2021) Systematic benchmarking of tools for CpG methylation detection from nanopore sequencing. *Nature communications,* **12**(1), 3438.

22. Martin, S., Heavens, D., Lan, Y., Horsfield, S., Clark, M. D., and Leggett, R. M. (2022) Nanopore adaptive sampling: a tool for enrichment of low abundance species in metagenomic samples. *Genome biology,* **23**(1), 1–27.

23. Timp, W., Mirsaidov, U. M., Wang, D., Comer, J., Aksimentiev, A., and Timp, G. (2010) Nanopore sequencing: electrical measurements of the code of life. *IEEE transactions on nanotechnology,* **9**(3), 281–294.

24. Diederichs, T., Pugh, G., Dorey, A., Xing, Y., Burns, J. R., Hung Nguyen, Q., Tornow, M., Tampé, R., and Howorka, S. (2019) Synthetic protein-conductive membrane nanopores built with DNA. *Nature communications,* **10**(1), 5018.

25. Howorka, S. and Siwy, Z. (2009) Nanopore analytics: sensing of single molecules. *Chemical Society Reviews,* **38**(8), 2360–2384.

26. Lindsay, S. (2016) The promises and challenges of solid-state sequencing. *Nature nanotechnology,* **11**(2), 109–111.

27. MacKenzie, M. and Argyropoulos, C. (2023) An introduction to nanopore sequencing: past, present, and future considerations. *Micromachines,* **14**(2), 459.

28. Logsdon, G. A., Vollger, M. R., and Eichler, E. E. (2020) Long-read human genome sequencing and its applications. *Nature Reviews Genetics,* **21**(10), 597–614.

29. Sanderson, N. D., Kapel, N., Rodger, G., Webster, H., Lipworth, S., Street, T. L., Peto, T., Crook, D., and Stoesser, N. (2023) Comparison of R9. 4.1/Kit10 and R10/Kit12 Oxford Nanopore flowcells and chemistries in bacterial genome reconstruction. *Microbial genomics,* **9**(1), 000910.

30. Ni, Y., Liu, X., Simeneh, Z. M., Yang, M., and Li, R. (2023) Benchmarking of Nanopore R10. 4 and R9. 4.1 flow cells in single-cell whole-genome amplification and whole-genome shotgun sequencing. *Computational and Structural Biotechnology Journal,* **21**, 2352–2364.

31. Castro-Wallace, S. L., Chiu, C. Y., John, K. K., Stahl, S. E., Rubins, K. H., McIntyre, A. B., Dworkin, J. P., Lupisella, M. L., Smith, D. J., Botkin, D. J., et al. (2017) Nanopore DNA sequencing and genome assembly on the International Space Station. *Scientific reports,* **7**(1), 18022.

32. McIntyre, A. B., Rizzardi, L., Yu, A. M., Alexander, N., Rosen, G. L., Botkin, D. J., Stahl, S. E., John, K. K., Castro-Wallace, S. L., McGrath, K., et al. (2016) Nanopore sequencing in microgravity. *npj Microgravity,* **2**(1), 1–9.

33. Kovaka, S., Fan, Y., Ni, B., Timp, W., and Schatz, M. C. (2021) Targeted nanopore sequencing by real-time mapping of raw electrical signal with UNCALLED. *Nature biotechnology,* **39**(4), 431–441.

34. Technologies, O. N. MinKNOW: Mux scan and active channel selection. (2024) https://community.nanoporetech.com/nanopore_learning/lessons/mux-scan-and-acs [Accessed: Dezember, 21, 2024].

35. Koch, C., Reilly-O'Donnell, B., Gutierrez, R., Lucarelli, C., Ng, F. S., Gorelik, J., Ivanov, A. P., and Edel, J. B. (2023) Nanopore sequencing of DNA-barcoded probes for highly multiplexed detection of microRNA, proteins and small biomarkers. *Nature Nanotechnology,* **18**(12), 1483–1491.

36. Prall, T. M., Neumann, E. K., Karl, J. A., Shortreed, C. G., Baker, D. A., Bussan, H. E., Wiseman, R. W., and O'Connor, D. H. (2021) Consistent ultra-long DNA sequencing with automated slow pipetting. *BMC genomics,* **22**(1), 1–12.

37. 'Giron' Koetsier, P. A. and Cantor, E. J. (2021) A simple approach for effective shearing and reliable concentration measurement of ultra-high-molecular-weight DNA. *BioTechniques,* (2), 439–444.

38. Jones, A., Torkel, C., Stanley, D., Nasim, J., Borevitz, J., and Schwessinger, B. (2021) High-molecular weight DNA extraction, clean-up and size selection for long-read sequencing. *PLoS One,* **16**(7), e0253830.

39. Maghini, D. G., Moss, E. L., Vance, S. E., and Bhatt, A. S. (2021) Improved high-molecular-weight DNA extraction, nanopore sequencing and metagenomic assembly from the human gut microbiome. *Nature protocols,* **16**(1), 458–471.

40. Zhang, Y., Zhang, Y., Burke, J. M., Gleitsman, K., Friedrich, S. M., Liu, K. J., and Wang, T.-H. (2016) A Simple Thermoplastic Substrate Containing Hierarchical Silica Lamellae for High-Molecular-Weight DNA Extraction. *Advanced Materials,* **28**(48), 10630–10636.

41. Jaudou, S., Tran, M.-L., Vorimore, F., Fach, P., and Delannoy, S. (2022) Evaluation of high molecular weight DNA extraction methods for long-read sequencing of Shiga toxin-producing Escherichia coli. *Plos one,* **17**(7), e0270751.

42. Wood, D. E., Lu, J., and Langmead, B. (2019) Improved metagenomic analysis with Kraken 2. *Genome biology,* **20**, 1–13.

43. Browne, P. D., Nielsen, T. K., Kot, W., Aggerholm, A., Gilbert, M. T. P., Puetz, L., Rasmussen, M., Zervas, A., and Hansen, L. H. (2020) GC bias affects genomic and metagenomic reconstructions, underrepresenting GC-poor organisms. *GigaScience,* **9**(2), giaa008.

44. Delahaye, C. and Nicolas, J. (2021) Sequencing DNA with nanopores: Troubles and biases. *PloS one,* **16**(10), e0257521.

45. Heavens, D., Chooneea, D., Giolai, M., Cuber, P., Aanstad, P., Martin, S., Alston, M., Misra, R., Clark, M. D., and Leggett, R. M. (October, 2021) How low can you go? Driving down the DNA input requirements for nanopore sequencing. *biorxiv,*.

46. Mojarro, A., Hachey, J., Ruvkun, G., Zuber, M. T., and Carr, C. E. (March, 2018) CarrierSeq: a sequence analysis workflow for low-input nanopore sequencing. *BMC Bioinformatics,* **19**(1), 108.

47. Arakawa, K. (2023) Ultralow-input genome library preparation for nanopore sequencing with droplet MDA. *Methods Mol. Biol.,* **2632**, 91–100.

48. Simon, S. A., Schmidt, K., Griesdorn, L., Soares, A. R., Bornemann, T. L., and Probst, A. J. (2023) Dancing the Nanopore limbo–Nanopore metagenomics from small DNA quantities for bacterial genome reconstruction. *BMC genomics,* **24**(1), 727.

49. Scott, M., Gunderson, C. W., Mateescu, E. M., Zhang, Z., and Hwa, T. (November, 2010) Interdependence of cell growth and gene expression: origins and consequences. *Science,* **330**(6007), 1099–1102.

50. Marzluff, W. F., Wagner, E. J., and Duronio, R. J. (November, 2008) Metabolism and regulation of canonical histone mRNAs: life without a poly(A) tail. *Nat. Rev. Genet.,* **9**(11), 843–854.

51. Turowski, T. W. and Boguta, M. (May, 2021) Specific features of RNA polymerases I and III: Structure and assembly. *Front. Mol. Biosci.,* **8**, 680090.

52. O'Neil, D., Glowatz, H., and Schlumpberger, M. (Jul, 2013) Ribosomal RNA depletion for efficient use of RNA-seq capacity. *Curr Protoc Mol Biol,* **Chapter 4**, Unit 4.19.

53. Fang, Y., Changavi, A., Yang, M., Sun, L., Zhang, A., Sun, D., Sun, Z., Zhang, B., and Xu, M. Q. (Jan, 2022) Nanopore Whole Transcriptome Analysis and Pathogen Surveillance by a Novel Solid-Phase Catalysis Approach. *Adv Sci (Weinh),* **9**(3), e2103373.

54. Freed, N. E., Vlková, M., Faisal, M. B., and Silander, O. K. (2020) Rapid and inexpensive whole-genome sequencing of SARS-CoV-2 using 1200 bp tiled amplicons and Oxford Nanopore Rapid Barcoding. *Biology Methods and Protocols,* **5**(1), bpaa014.

55. Whitford, W., Hawkins, V., Moodley, K. S., Grant, M. J., Lehnert, K., Snell, R. G., and Jacobsen, J. C. (2022) Proof of concept for multiplex amplicon sequencing for mutation identification using the MinION nanopore sequencer. *Scientific reports,* **12**(1), 8572.

56. McCrone, J. T., Woods, R. J., Martin, E. T., Malosh, R. E., Monto, A. S., and Lauring, A. S. (May, 2018) Stochastic processes constrain the within and between host evolution of influenza virus. *Elife,* **7**.

57. Gilbert, K. B., Fahlgren, N., Kasschau, K. D., Chapman, E. J., Carrington, J. C., and Carbonell, A. (November, 2014) Preparation of multiplexed small RNA libraries from plants. *Bio Protoc.,* **4**(21).

58. Taylor, B. C., Lejzerowicz, F., Poirel, M., Shaffer, J. P., Jiang, L., Aksenov, A., Litwin, N., Humphrey, G., Martino, C., Miller-Montgomery, S., Dorrestein, P. C., Veiga, P., Song, S. J., McDonald, D., Derrien, M., and Knight, R. (March, 2020) Consumption of fermented foods is associated with systematic differences in the gut microbiome and metabolome. *mSystems,* **5**(2).

59. Brandt, C., Krautwurst, S., Spott, R., Lohde, M., Jundzill, M., Marquet, M., and Hölzer, M. (July, 2021) PoreCov-an easy to use, fast, and robust workflow for SARS-CoV-2 genome reconstruction via nanopore sequencing. *Front. Genet.,* **12**, 711437.

60. Payne, A., Holmes, N., Clarke, T., Munro, R., Debebe, B. J., and Loose, M. (2021) Readfish enables targeted nanopore sequencing of gigabase-sized genomes. *Nature biotechnology,* **39**(4), 442–450.

61. Filser, M., Schwartz, M., Merchadou, K., Hamza, A., Villy, M.-C., Decees, A., Frouin, E., Girard, E., Caputo, S. M., Renault, V., et al. (2023) Adaptive nanopore sequencing to determine pathogenicity of BRCA1 exonic duplication. *Journal of Medical Genetics,* **60**(12), 1206–1209.

62. De Meulenaere, K., Cuypers, W. L., Gauglitz, J. M., Guetens, P., Rosanas-Urgell, A., Laukens, K., and Cuypers, B. (2024) Selective whole-genome sequencing of Plasmodium parasites directly from blood samples by Nanopore adaptive sampling. *Mbio,* **15**(1), e01967–23.

63. Marquet, M., Zöllkau, J., Pastuschek, J., Viehweger, A., Schleußner, E., Makarewicz, O., Pletz, M. W., Ehricht, R., and Brandt, C. (2022) Evaluation of microbiome enrichment and host DNA depletion in human vaginal samples using Oxford Nanopore's adaptive sequencing. *Scientific reports,* **12**(1), 4000.

64. Leger, A. and Leonardi, T. (February, 2019) pycoQC, interactive quality control for Oxford Nanopore Sequencing. *Journal of Open Source Software,* **4**(34), 1236.

65. De Coster, W. and Rademakers, R. (05, 2023) NanoPack2: population-scale evaluation of long-read sequencing data. *Bioinformatics,* **39**(5) btad311.

66. Li, H. (May, 2018) Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics,* **34**(18), 3094–3100.

67. Jain, C., Rhie, A., Zhang, H., Chu, C., Walenz, B. P., Koren, S., and Phillippy, A. M. (July, 2020) Weighted minimizer sampling improves long read mapping. *Bioinformatics,* **36**(Supplement_1), i111–i118.

68. Boostrom, I., Portal, E. A. R., Spiller, O. B., Walsh, T. R., and Sands, K. (March, 2022) Comparing Long-Read Assemblers to Explore the Potential of a Sustainable Low-Cost, Low-Infrastructure Approach to Sequence Antimicrobial Resistant Bacteria With Oxford Nanopore Sequencing. *Frontiers in Microbiology,* **13**.

69. Shafin, K., Pesout, T., Lorig-Roach, R., Haukness, M., Olsen, H. E., Bosworth, C., Armstrong, J., Tigyi, K., Maurer, N., Koren, S., Sedlazeck, F. J., Marschall, T., Mayes, S., Costa, V., Zook, J. M., Liu, K. J., Kilburn, D., Sorensen, M., Munson, K. M., Vollger, M. R., Monlong, J., Garrison, E., Eichler, E. E., Salama, S., Haussler, D., Green, R. E., Akeson, M., Phillippy, A., Miga, K. H., Carnevali, P., Jain, M., and Paten, B. (May, 2020) Nanopore sequencing and the Shasta toolkit enable efficient de novo assembly of eleven human genomes. *Nature Biotechnology,* **38**(9), 1044–1053.

70. Hu, J., Wang, Z., Sun, Z., Hu, B., Ayoola, A. O., Liang, F., Li, J., Sandoval, J. R., Cooper, D. N., Ye, K., Ruan, J., Xiao, C.-L., Wang, D.-P., Wu, D.-D., and Wang, S. (March, 2023) An efficient error correction and accurate assembly tool for noisy long reads. *biorxiv,*.

71. Sun, J., Li, R., Chen, C., Sigwart, J. D., and Kocot, K. M. (April, 2021) Benchmarking Oxford Nanopore read assemblers for high-quality molluscan genomes. *Philosophical Transactions of the Royal Society B: Biological Sciences,* **376**(1825), 20200160.

72. Vaser, R., Sović, I., Nagarajan, N., and Šikić, M. (January, 2017) Fast and accurate de novo genome assembly from long uncorrected reads. *Genome Research,* **27**(5), 737–746.

73. Gholami, A., Kim, S., Dong, Z., Yao, Z., Mahoney, M. W., and Keutzer, K. (2021) A Survey of Quantization Methods for Efficient Neural Network Inference. *arXiv,*.

74. Fleming, A. M., Mathewson, N. J., Manage, S. A. H., and Burrows, C. J. (September, 2021) Nanopore Dwell Time Analysis Permits Sequencing and Conformational Assignment of Pseudouridine in SARS-CoV-2. *ACS Central Science,* **7**(10), 1707–1717.

75. Gamaarachchi, H., Lam, C. W., Jayatilaka, G., Samarakoon, H., Simpson, J. T., Smith, M. A., and Parameswaran, S. (2020) GPU accelerated adaptive banded event alignment for rapid comparative nanopore signal analysis. *BMC bioinformatics,* **21**(1), 1–13.

76. Raiber, E.-A., Hardisty, R., van Delft, P., and Balasubramanian, S. (2017) Mapping and elucidating the function of modified bases in DNA. *Nature Reviews Chemistry,* **1**(9), 0069.

77. Klungland, A. and Robertson, A. B. (2017) Oxidized C5-methyl cytosine bases in DNA: 5-Hydroxymethylcytosine; 5-formylcytosine; and 5-carboxycytosine. *Free Radical Biology and Medicine,* **107**, 62–68.

78. Heyn, H. and Esteller, M. (2015) An adenine code for DNA: a second life for N6-methyladenine. *Cell,* **161**(4), 710–713.

79. Simpson, J. T., Workman, R. E., Zuzarte, P., David, M., Dursi, L., and Timp, W. (2017) Detecting DNA cytosine methylation using nanopore sequencing. *Nature methods,* **14**(4), 407–410.

80. Sigurpalsdottir, B. D., Stefansson, O. A., Holley, G., Beyter, D., Zink, F., Hardarson, M., Sverrisson, S., Kristinsdottir, N., Magnusdottir, D. N., Magnusson, O., et al. (2024) A comparison of methods for detecting DNA methylation from long-read sequencing of human genomes. *Genome Biology,* **25**(1), 69.

81. Liu, Y., Rosikiewicz, W., Pan, Z., Jillette, N., Wang, P., Taghbalout, A., Foox, J., Mason, C., Carroll, M., Cheng, A., et al. (2021) DNA methylation-calling tools for Oxford Nanopore sequencing: a survey and human epigenome-wide evaluation. *Genome biology,* **22**(1), 1–33.

82. Yao, B., Hsu, C., Goldner, G., Michaeli, Y., Ebenstein, Y., and Listgarten, J. (2021) Nanopore callers for epigenetics from limited supervised data. *bioRxiv,* pp. 2021–06.

83. Akbari, V., Garant, J.-M., O'Neill, K., Pandoh, P., Moore, R., Marra, M. A., Hirst, M., and Jones, S. J. (2021) Megabase-scale methylation phasing using nanopore long reads and NanoMethPhase. *Genome biology,* **22**(1), 1–21.

84. Chen, H., Yao, J., Bao, R., Dong, Y., Zhang, T., Du, Y., Wang, G., Ni, D., Xun, Z., Niu, X., Ye, Y., and Li, H.-B. (February, 2021) Cross-talk of four types of RNA modification writers defines

tumor microenvironment and pharmacogenomic landscape in colorectal cancer. *Molecular Cancer,* **20**(1).

85. Höner zu Siederdissen, C., Spangenberg, J., Bisdorf, K., Krautwurst, S., Srivastava, A., Marz, M., and Taubert, M. (December, 2024) Nanopore sequencing enables novel detection of deuterium incorporation in DNA. *Computational and Structural Biotechnology Journal,* **23**, 3584–3594.

86. Stoiber, M., Quick, J., Egan, R., Eun Lee, J., Celniker, S., Neely, R. K., Loman, N., Pennacchio, L. A., and Brown, J. (2016) De novo identification of DNA modifications enabled by genome-guided nanopore signal processing. *BioRxiv,* p. 094672.

87. Rand, A. C., Jain, M., Eizenga, J. M., Musselman-Brown, A., Olsen, H. E., Akeson, M., and Paten, B. (2017) Mapping DNA methylation with high-throughput nanopore sequencing. *Nature methods,* **14**(4), 411–413.

88. Liu, Q., Georgieva, D. C., Egli, D., and Wang, K. (2019) NanoMod: a computational tool to detect DNA modifications using Nanopore long-read sequencing data. *BMC genomics,* **20**(1), 31–42.

89. McIntyre, A. B., Alexander, N., Grigorev, K., Bezdan, D., Sichtig, H., Chiu, C. Y., and Mason, C. E. (2019) Single-molecule sequencing detection of N 6-methyladenine in microbial reference materials. *Nature communications,* **10**(1), 579.

90. Ni, P., Huang, N., Zhang, Z., Wang, D.-P., Liang, F., Miao, Y., Xiao, C.-L., Luo, F., and Wang, J. (2019) DeepSignal: detecting DNA methylation state from Nanopore sequencing reads using deep-learning. *Bioinformatics,* **35**(22), 4586–4595.

91. Liu, Q., Fang, L., Yu, G., Wang, D., Xiao, C.-L., and Wang, K. (2019) Detection of DNA base modifications by deep recurrent neural network on Oxford Nanopore sequencing data. *Nature communications,* **10**(1), 2449.

92. Ahsan, M. U., Gouru, A., Chan, J., Zhou, W., and Wang, K. (2024) A signal processing and deep learning framework for methylation detection using Oxford Nanopore sequencing. *Nature Communications,* **15**(1), 1448.

93. Zhang, Y.-z., Yamaguchi, K., Hatakeyama, S., Furukawa, Y., Miyano, S., Yamaguchi, R., and Imoto, S. (2021) On the application of BERT models for nanopore methylation detection. In *2021 IEEE International Conference on Bioinformatics and Biomedicine (BIBM)* IEEE pp. 320–327.

94. Bonet, J., Chen, M., Dabad, M., Heath, S., Gonzalez-Perez, A., Lopez-Bigas, N., and Lagergren, J. (2022) DeepMP: a deep learning tool to detect DNA base modifications on Nanopore sequencing data. *Bioinformatics,* **38**(5), 1235–1243.

95. Lorenz, D. A., Sathe, S., Einstein, J. M., and Yeo, G. W. (2020) Direct RNA sequencing enables m6A detection in endogenous transcript isoforms at base-specific resolution. *Rna,* **26**(1), 19–28.

96. Liu, H., Begik, O., and Novoa, E. M. (2021) EpiNano: detection of m 6 A RNA modifications using oxford nanopore direct RNA sequencing. *RNA Modifications: Methods and Protocols,* pp. 31–52.

97. Ueda, H. (2020) nanoDoc: RNA modification detection using Nanopore raw reads with Deep One-Class Classification. *bioRxiv,* pp. 2020–09.

98. Maier, K. C., Gressel, S., Cramer, P., and Schwalb, B. (September, 2020) Native molecule sequencing by nano-ID reveals synthesis and stability of RNA isoforms. *Genome Research,* **30**(9), 1332–1344.

99. Price, A. M., Hayer, K. E., McIntyre, A. B. R., Gokhale, N. S., Abebe, J. S., Della Fera, A. N., Mason, C. E., Horner, S. M., Wilson, A. C., Depledge, D. P., and Weitzman, M. D. (November, 2020) Direct RNA sequencing reveals m6A modifications on adenovirus RNA are necessary for efficient splicing. *Nature Communications,* **11**(1).

100. Pratanwanich, P. N., Yao, F., Chen, Y., Koh, C. W., Wan, Y. K., Hendra, C., Poon, P., Goh, Y. T., Yap, P. M., Chooi, J. Y., et al. (2021) Identification of differential RNA modifications from nanopore direct RNA sequencing with xPore. *Nature biotechnology,* **39**(11), 1394–1402.

101. Gao, Y., Liu, X., Wu, B., Wang, H., Xi, F., Kohnen, M. V., Reddy, A. S., and Gu, L. (2021) Quantitative profiling of N 6-methyladenosine at single-base resolution in stem-differentiating xylem of Populus trichocarpa using Nanopore direct RNA sequencing. *Genome biology,* **22**, 1–17.

102. Begik, O., Lucas, M. C., Pryszcz, L. P., Ramirez, J. M., Medina, R., Milenkovic, I., Cruciani, S., Liu, H., Vieira, H. G. S., Sas-Chen, A., et al. (2021) Quantitative profiling of pseudouridylation dynamics in native RNAs with nanopore sequencing. *Nature biotechnology,* **39**(10), 1278–1291.

103. Jenjaroenpun, P., Wongsurawat, T., Wadley, T. D., Wassenaar, T. M., Liu, J., Dai, Q., Wanchai, V., Akel, N. S., Jamshidi-Parsian, A., Franco, A. T., et al. (2021) Decoding the epitranscriptional landscape from native RNA

sequences. *Nucleic acids research,* **49**(2), e7–e7.

104. Leger, A., Amaral, P. P., Pandolfini, L., Capitanchik, C., Capraro, F., Miano, V., Migliori, V., Toolan-Kerr, P., Sideri, T., Enright, A. J., et al. (2021) RNA modifications detection by comparative Nanopore direct RNA sequencing. *Nature communications,* **12**(1), 7198.

105. Hendra, C., Pratanwanich, P. N., Wan, Y. K., Goh, W. S., Thiery, A., and Göke, J. (2022) Detection of m6A from direct RNA sequencing using a multiple instance learning framework. *Nature Methods,* **19**(12), 1590–1598.

106. Hassan, D., Acevedo, D., Daulatabad, S. V., Mir, Q., and Janga, S. C. (July, 2022) Penguin: A tool for predicting pseudouridine sites in direct RNA nanopore sequencing data. *Methods,* **203**, 478–487.

107. Qin, H., Ou, L., Gao, J., Chen, L., Wang, J.-W., Hao, P., and Li, X. (January, 2022) DENA: training an authentic neural network model using Nanopore sequencing data of Arabidopsis transcripts for detection and quantification of N6-methyladenosine on RNA. *Genome Biology,* **23**(1).

108. Spangenberg, J., zu Siederdissen, C. H., Žarković, M., Triebel, S., Rose, R., Christophersen, C. M., Paltzow, L., Hegab, M. M., Wansorra, A., Srivastava, A., Krumbholz, A., and Marz, M. (March, 2023) Magnipore: Prediction of differential single nucleotide changes in the Oxford Nanopore Technologies sequencing signal of SARS-CoV-2 samples. *bioRxiv,*.

109. Chan, A., Naarmann-de Vries, I. S., Scheitl, C. P. M., Höbartner, C., and Dieterich, C. (April, 2024) Detecting m6A at single-molecular resolution via direct RNA sequencing and realistic training data. *Nature Communications,* **15**(1).

SUPPLEMENTARY INFORMATION

Table S1: **Sample Summary** – overview of all runs used for the analysis of Fig. 2, Fig. 4, Fig. 6 and Fig. S4–S10. RNA or DNA sequencing; fc version – R9 or R10 flow cells; device – MinION or GridION device; #cha – amount of active channels at the begin of the sequencing run; #por – amount of active pores at the beginning of the sequencing run; fc age – time passed since delivery of the Flow cell in days; sample type – from which kind of species the sequenced material originated in nine categories: virus, bacteria, protist, metagenomic, insect, mouse, human; buffer – LFB, SFB or RNA buffer; ng load – amount of loaded library (DNA/RNA) in ng; EB 12 – amount of sequenced bases after 12 h, estimated by sequencing software MinKNOW; mean rl – mean estimated read length of all sequenced reads in bases; pore hl – calculated pore half life time of the flow cell in hours (see main text).

ID	RNA/ DNA	fc version	device	#cha	#por	fc age	sample type	buffer	ng load	Estimated bases	EB12	mean rl	pore hl	fc id
1.1	DNA	R10	GridIon	510	NA	NA	virus	SFB	NA	8161	4721	2036	998	FAU91805
2.1	DNA	R10	GridIon	480	NA	NA	virus	SFB	NA	4745	4745	2934	1630	FAX97604
3.1	DNA	R10	GridIon	467	1766	NA	virus	SFB	NA	6041	3977	2137	643	FAY00760
4.1	DNA	R10	GridIon	502	1860	NA	virus	SFB	NA	8139	5074	2640	1012	FAY01081
5.1	DNA	R10	GridIon	501	1869	NA	virus	SFB	NA	9339	5444	3776	1397	FAY01249
6.1	DNA	R10	GridIon	504	NA	NA	virus	SFB	NA	9932	5699	4022	1510	FAY01385
7.1	DNA	R10	MinIon	293	1368	172	virus	SFB	60	177	153	419	51	FAO88286
8.1	RNA	R9	MinIon	295	749	174	virus	RNA	140	523	423	1530	900	FAM95379
9.1	RNA	R9	MinIon	487	1333	56	virus	RNA	210	1571	923	1700	NA	FAO07649
10.2	RNA	R9	MinIon	489	1385	49	virus	RNA	90	1495	1031	1454	NA	FAO10418
11.1	RNA	R9	MinIon	501	1282	10	virus	RNA	74	1324	591	1072	1638	FAO53185
12.1	RNA	R9	MinIon	506	1372	112	virus	RNA	308	1247	885	1046	797	FAO83518
13.1	RNA	R9	MinIon	473	1201	127	virus	RNA	186	1326	869	1577	810	FAO86549
14.1	RNA	R9	MinIon	470	1201	102	virus	RNA	304	118	117	1206	79	FAO86555
15.1	RNA	R9	MinIon	505	1380	111	virus	RNA	392	1028	801	1123	637	FAO86826
16.1	RNA	R9	MinIon	510	1342	34	virus	RNA	256	483	462	982	269	FAO86858
17.1	RNA	R9	MinIon	500	1241	111	virus	RNA	348	136	136	1112	113	FAO86895
18.1	RNA	R9	MinIon	500	1308	110	virus	RNA	266	626	580	1099	414	FAO86907
19.1	RNA	R9	MinIon	498	1298	110	virus	RNA	420	296	296	1140	212	FAO86991
20.1	RNA	R9	MinIon	442	1142	93	virus	RNA	224	284	283	1354	225	FAO86996
21.1	RNA	R9	MinIon	495	1228	111	virus	RNA	360	314	312	1133	215	FAO87031
22.1	RNA	R9	MinIon	482	1249	127	virus	RNA	276	295	294	1353	232	FAO87106
23.1	RNA	R9	MinIon	432	1002	9	virus	RNA	167	1400	650	1677	1348	FAO88388
24.1	DNA	R9	MinIon	512	1960	3	virus	SFB	575	13251	6708	765	1439	FAO92019
25.1	RNA	R9	MinIon	491	1327	110	virus	RNA	346	600	556	1111	352	FAO92042
26.1	DNA	R9	MinIon	511	1907	6	virus	SFB	516	9748	6396	866	958	FAO92112
27.1	RNA	R9	MinIon	497	1307	110	virus	RNA	356	768	657	1123	526	FAO92116
28.1	RNA	R9	MinIon	502	1365	111	virus	RNA	254	1141	847	975	795	FAO92254
29.1	RNA	R9	MinIon	322	893	93	virus	RNA	236	317	316	1151	368	FAO92303
30.1	RNA	R9	MinIon	350	1210	167	virus	RNA	268	1708	767	1666	1373	FAP02359
31.1	RNA	R9	MinIon	93	152	51	virus	RNA	580	46	44	1256	166	FAQ01609
31.2	RNA	R9	MinIon	NA	814	NA	virus	RNA	580	NA	487	1246	557	FAQ01609
32.1	RNA	R9	MinIon	420	965	147	virus	RNA	52	7	7	605	53	FAQ69187
33.1	DNA	R10	GridIon	510	1609	NA	bacteria	LFB	NA	22320	4567	10233	3562	FAU91295
34.1	DNA	R10	GridIon	511	1556	NA	bacteria	LFB	NA	10173	3988	7108	2071	FAU92937
35.1	DNA	R10	GridIon	502	1491	NA	bacteria	LFB	NA	12166	3944	7911	1930	FAW21704
36.1	DNA	R10	GridIon	510	1460	NA	bacteria	LFB	NA	12767	3779	1553	2366	FAW22654
37.1	DNA	R10	GridIon	510	1578	NA	bacteria	LFB	NA	14943	4433	3423	2026	FAW22671
38.1	DNA	R10	GridIon	348	1019	NA	bacteria	LFB	NA	6458	2713	1629	1675	FAW23628
39.1	DNA	R10	GridIon	498	1567	NA	bacteria	LFB	NA	6065	3604	6144	719	FAW45145
40.1	DNA	R10	GridIon	510	1566	NA	bacteria	LFB	NA	14424	4395	8707	2285	FAW45147
41.1	DNA	R10	GridIon	478	1501	NA	bacteria	LFB	NA	7288	3437	9545	1050	FAW45167
42.1	DNA	R10	GridIon	413	1262	NA	bacteria	LFB	NA	2715	2177	1855	456	FAW98004
43.1	DNA	R10	GridIon	503	NA	NA	bacteria	LFB	NA	3563	NA	NA	NA	FAW98056
44.1	DNA	R10	GridIon	499	1899	NA	bacteria	LFB	NA	13610	6244	3383	1169	FAW99941
45.1	DNA	R10	GridIon	495	NA	NA	bacteria	LFB	NA	4855	NA	NA	NA	FAX00084
46.1	DNA	R10	GridIon	484	1484	NA	bacteria	LFB	NA	7281	3929	2441	851	FAX00224
47.1	DNA	R10	GridIon	487	1812	105	bacteria	LFB	1216	11256	4949	3599	934	FAX12900
48.1	DNA	R10	GridIon	221	813	112	bacteria	LFB	403	4209	2002	3186	657	FAX13587
49.1	DNA	R10	GridIon	511	NA	98	bacteria	LFB	552	14912	5538	1931	NA	FAX14134
50.1	DNA	R10	GridIon	506	NA	105	bacteria	LFB	324	10673	4891	2555	NA	FAX18203
51.1	DNA	R10	GridIon	484	1844	NA	bacteria	LFB	NA	21675	5881	3121	2151	FAX70780
52.1	DNA	R10	MinIon	235	999	186	bacteria	SFB	139.2	1239	770	1169	1531	FAO85473

ID	RNA/ DNA	fc version	device	#cha	#por	fc age	sample type	buffer	ng load	Estimated bases	EB12	mean rl	pore hl	fc id
53.1	DNA	R10	MinIon	348	NA	36	bacteria	LFB	NA	2625	NA	NA	NA	FAS37108
xx.1	DNA	R10	MinIon	428	1116	35	bacteria	LFB	1008	1242	1041	6008	NA	FAS37108
54.1	DNA	R10	MinIon	132	NA	31	bacteria	LFB	298	997	707	9448	NA	FAS38416
xx.1	DNA	R10	MinIon	417	NA	30	bacteria	LFB	276	4825	3281	9328	NA	FAS38416
55.1	DNA	R10	MinIon	229	NA	1	bacteria	LFB	1440	651	443	2796	NA	FAV74660
56.1	DNA	R10	MinIon	500	NA	7	bacteria	LFB	157	11585	4816	3574	NA	FAW66139
57.1	DNA	R10	MinIon	468	NA	2	bacteria	LFB	NA	9767	4480	5356	NA	FAW73005
xx.1	DNA	R10	MinIon	NA	NA	1	bacteria	LFB	NA	NA	NA	NA	NA	FAW73005
58.1	DNA	R10	MinIon	226	NA	1	bacteria	LFB	199	4678	NA	NA	NA	FAW75095
59.1	DNA	R10	MinIon	487	NA	1	bacteria	LFB	226	0	NA	NA	NA	FAX18284
60.1	DNA	R10	MinIon	473	NA	1	bacteria	LFB	154	12573	4714	4304	NA	FAX30985
61.1	DNA	R9	GridIon	484	1447	NA	bacteria	LFB	NA	9421	3623	7487	1791	FAU91863
62.1	DNA	R9	GridIon	511	1398	NA	bacteria	LFB	NA	6066	3190	13070	2225	FAV24719
63.1	DNA	R9	GridIon	494	1512	NA	bacteria	LFB	NA	18854	6177	9601	2187	FAV30532
64.1	DNA	R9	MinIon	139	482	101	bacteria	LFB	200	100	NA	5093	18	FAO16410
65.1	DNA	R9	MinIon	499	1439	4	bacteria	SFB	238	18063	6136	6785	2038	FAO53260
66.1	DNA	R9	MinIon	416	1021	114	bacteria	SFB	352	11310	4237	3302	2384	FAO87074
67.4	DNA	R9	MinIon	449	1541	37	bacteria	LFB	200	654	NA	4932	NA	FAO92001
68.1	DNA	R9	MinIon	416	1161	178	bacteria	SFB	600	7394	4172	2023	1108	FAP00827
69.1	DNA	R9	MinIon	449	1659	121	bacteria	SFB	775	2507	NA	18442	NA	FAQ01004
69.2	DNA	R9	MinIon	355	1393	122	bacteria	SFB	444	5911	3731	12254	2	FAQ01004
70.1	DNA	R9	MinIon	494	1620	44	bacteria	SFB	1428	8812	4452	1262	1652	FAQ58659
71.1	DNA	R9	MinIon	454	1692	58	bacteria	SFB	504	18406	5950	20443	2303	FAQ58677
72.1	DNA	R9	MinIon	493	1312	41	bacteria	SFB	479.6	25403	6853	2023	2787	FAQ69194
73.1	DNA	R9	MinIon	480	1468	87	bacteria	LFB	700	23019	6807	23746	2554	FAR96927
74.1	DNA	R9	MinIon	406	1398	14	protist	LFB	600	4265	3028	1592	446	FAQ01569
75.1	DNA	R9	MinIon	485	1692	80	protist	SFB	310	4620	3452	1988	608	FAQ01665
76.1	DNA	R9	MinIon	508	1881	16	protist	SFB	528	10925	5862	1852	1169	FAQ01773
77.1	DNA	R9	MinIon	499	1785	78	protist	SFB	470	4948	3880	1867	557	FAQ01847
77.2	DNA	R9	MinIon	371	1329	79	protist	SFB	160	2085	1749	2093	258	FAQ01847
78.1	DNA	R10	GridIon	503	1719	NA	metagenomic	LFB	NA	5046	2434	5637	821	FAW10206
79.1	DNA	R10	GridIon	501	1723	NA	metagenomic	LFB	NA	5681	2805	5293	932	FAW13592
80.1	DNA	R10	GridIon	457	1556	NA	metagenomic	LFB	NA	5468	2379	6421	1184	FAX33641
81.1	DNA	R10	GridIon	500	1733	NA	metagenomic	LFB	NA	7313	3103	5179	1068	FAX57960
82.1	DNA	R10	GridIon	507	1657	NA	metagenomic	LFB	NA	5308	2229	7428	1191	FAX59252
83.1	DNA	R10	MinIon	450	1657	156	metagenomic	SFB	600	7086	3361	1648	1199	FAO79888
84.1	DNA	R9	MinIon	NA	1953	NA	metagenomic	SFB	447	NA	7190	1355	2371	FAK75664
85.1	DNA	R9	MinIon	NA	1921	NA	metagenomic	SFB	471.6	NA	6634	1246	2878	FAL78178
85.2	DNA	R9	MinIon	NA	19	NA	metagenomic	SFB	471.6	NA	NA	836	NA	FAL78178
86.1	DNA	R9	MinIon	NA	186	NA	metagenomic	SFB	280	NA	555	2355	NA	FAL78306
86.2	DNA	R9	MinIon	NA	1915	NA	metagenomic	SFB	280	NA	7182	3067	3253	FAL78306
87.1	DNA	R9	MinIon	507	1933	154	metagenomic	SFB	252	17853	7006	2561	2339	FAL80840
87.2	DNA	R9	MinIon	396	1452	155	metagenomic	SFB	252	10310	5187	2378	1001	FAL80840
88.1	DNA	R9	MinIon	512	NA	167	metagenomic	SFB	256	19674	7000	2717	3098	FAL81216
88.2	DNA	R9	MinIon	378	1279	168	metagenomic	SFB	256	9050	4769	2434	1110	FAL81216
89.1	DNA	R9	MinIon	506	1793	147	metagenomic	SFB	301	6030	4453	1833	729	FAL83332
89.2	DNA	R9	MinIon	457	1617	148	metagenomic	SFB	301	2492	2204	1124	271	FAL83332
90.1	DNA	R9	MinIon	510	1966	179	metagenomic	SFB	1300	27326	6719	2119	2953	FAL83351
91.1	DNA	R9	MinIon	479	1813	177	metagenomic	SFB	431	13234	6063	1166	1671	FAL84179
92.1	DNA	R9	MinIon	485	1802	171	metagenomic	SFB	379	19643	5541	2215	2231	FAL85078
93.1	DNA	R9	MinIon	508	NA	150	metagenomic	SFB	360	14580	7302	4228	2041	FAL91762
93.2	DNA	R9	MinIon	420	1588	151	metagenomic	SFB	360	13564	5661	2288	1945	FAL91762
94.1	DNA	R9	MinIon	507	1891	181	metagenomic	SFB	400	17020	5792	1623	2302	FAM93261
95.1	DNA	R9	MinIon	376	1365	174	metagenomic	SFB	410	8949	4597	2509	1247	FAM95301
96.1	DNA	R9	MinIon	503	1843	190	metagenomic	SFB	609	25293	6351	1795	2686	FAM95308
97.1	DNA	R9	MinIon	508	1945	181	metagenomic	SFB	458	26777	7184	3221	2585	FAM95383
98.1	DNA	R9	MinIon	271	1005	174	metagenomic	SFB	300	2850	2272	2305	606	FAM96150
99.1	DNA	R9	MinIon	503	1896	178	metagenomic	SFB	564	17602	5838	1871	2520	FAM96240
100.1	DNA	R9	MinIon	465	1691	171	metagenomic	SFB	253	8206	5247	3313	1363	FAM96327
100.2	DNA	R9	MinIon	353	1293	172	metagenomic	SFB	253	6930	4180	1905	1291	FAM96327
101.1	DNA	R9	MinIon	507	1876	190	metagenomic	SFB	552	28264	6597	1462	3230	FAM96767
102.1	DNA	R9	MinIon	481	1780	54	metagenomic	SFB	460	12460	5585	1483	1076	FAO10159
103.1	DNA	R9	MinIon	499	1771	39	metagenomic	SFB	372	2181	1580	1713	812	FAO10214

ID	RNA/ DNA	fc version	device	#cha	#por	fc age	sample type	buffer	ng load	Estimated bases	EB12	mean rl	pore hl	fc id
103.2	DNA	R9	MinIon	240	796	45	metagenomic	SFB	460	3263	2633	1585	850	FAO10214
104.1	DNA	R9	MinIon	507	1594	40	metagenomic	SFB	960	4117	2129	922	1157	FAO10253
105.1	DNA	R9	MinIon	491	1602	41	metagenomic	SFB	408	2037	1597	1785	NA	FAO10336
106.1	DNA	R9	MinIon	498	1742	39	metagenomic	SFB	393	4078	2218	1444	1008	FAO10413
10.5	DNA	R9	MinIon	337	1157	55	metagenomic	SFB	564	3713	3571	1340	644	FAO10418
107.1	DNA	R9	MinIon	506	1948	42	metagenomic	SFB	634	23235	7055	1672	2661	FAO12115
108.1	DNA	R9	MinIon	510	1955	42	metagenomic	SFB	538	18522	6564	1740	2241	FAO12140
109.1	DNA	R9	MinIon	501	1881	42	metagenomic	SFB	600	18284	6171	1912	2292	FAO12169
110.1	DNA	R9	MinIon	484	NA	20	metagenomic	SFB	672	10957	5227	2273	1750	FAO16442
111.1	DNA	R9	MinIon	512	1853	48	metagenomic	SFB	240	19079	6678	1685	2428	FAO83526
112.1	DNA	R9	MinIon	509	1842	34	metagenomic	SFB	474	24501	6387	2150	3460	FAO86805
113.1	DNA	R9	MinIon	463	1637	87	metagenomic	SFB	189.6	3831	1777	2757	968	FAP46095
114.1	DNA	R9	MinIon	488	1659	87	metagenomic	SFB	158.4	15254	5286	1071	1976	FAP49364
115.1	DNA	R9	MinIon	500	1526	66	metagenomic	SFB	286.8	10070	3581	2934	1208	FAQ01716
116.1	DNA	R9	MinIon	479	1728	52	metagenomic	SFB	470.4	6142	2766	2240	834	FAQ01897
117.1	DNA	R9	MinIon	484	1680	52	metagenomic	SFB	477.6	7060	3049	1966	1301	FAQ10186
118.1	DNA	R9	MinIon	495	1850	36	metagenomic	SFB	674.4	12282	5140	1137	1761	FAQ37458
119.1	DNA	R9	MinIon	496	1817	37	metagenomic	SFB	574	11248	4596	1583	2062	FAQ69372
120.1	DNA	R9	MinIon	499	1904	151	insect	LFB	103	2509	NA	13240	264	FAL78077
120.2	DNA	R9	MinIon	481	1707	165	insect	SFB	113	5427	4328	5385	576	FAL78077
121.1	DNA	R9	MinIon	509	1984	150	insect	LFB	232	5127	4677	10627	438	FAL83572
122.1	DNA	R9	MinIon	380	1456	20	insect	SFB	255	2097	1977	5473	221	FAO10283
10.1	DNA	R9	MinIon	508	NA	49	insect	LFB	144	2393	NA	20800	386	FAO10418
10.3	DNA	R9	MinIon	390	1405	54	insect	LFB	144	1651	NA	21548	273	FAO10418
10.4	DNA	R9	MinIon	354	1257	55	insect	SFB	255	1084	NA	5500	162	FAO10418
123.1	DNA	R9	MinIon	462	1666	7	insect	SFB	226	7190	5361	4274	729	FAO10430
123.2	DNA	R9	MinIon	373	1394	21	insect	SFB	255	1473	1441	10126	165	FAO10430
124.1	DNA	R9	MinIon	497	1843	69	insect	LFB	200	2766	2638	16530	263	FAO12153
124.2	DNA	R9	MinIon	447	1699	74	insect	LFB	200	2253	2205	16834	244	FAO12153
124.3	DNA	R9	MinIon	388	1512	75	insect	LFB	160	1771	NA	15219	229	FAO12153
124.4	DNA	R9	MinIon	358	1433	76	insect	LFB	160	1413	NA	16750	226	FAO12153
124.5	DNA	R9	MinIon	355	1340	88	insect	LFB	150	1716	NA	19073	249	FAO12153
124.6	DNA	R9	MinIon	230	951	89	insect	LFB	100	1003	NA	14562	258	FAO12153
125.1	DNA	R9	MinIon	436	1540	60	insect	LFB	300	3879	3655	16927	361	FAO12159
125.2	DNA	R9	MinIon	410	1489	61	insect	LFB	200	2721	2671	18638	272	FAO12159
125.3	DNA	R9	MinIon	416	1591	61	insect	LFB	200	3097	NA	13451	547	FAO12159
125.4	DNA	R9	MinIon	364	1366	62	insect	LFB	200	2271	NA	14192	338	FAO12159
125.5	DNA	R9	MinIon	321	1196	67	insect	LFB	200	1319	NA	16557	175	FAO12159
125.6	DNA	R9	MinIon	264	1086	68	insect	LFB	100	719	NA	16788	151	FAO12159
126.1	DNA	R9	MinIon	510	1935	49	insect	SFB	200	9855	6485	4358	1016	FAO86712
126.2	DNA	R9	MinIon	476	1811	50	insect	SFB	100	6855	5140	4640	724	FAO86712
126.3	DNA	R9	MinIon	419	1570	51	insect	SFB	67	4758	4070	4859	545	FAO86712
126.4	DNA	R9	MinIon	307	1381	62	insect	LFB	70	1701	1587	7019	264	FAO86712
127.1	DNA	R9	MinIon	445	1709	8	insect	LFB	200	4467	4156	18700	443	FAO86834
127.2	DNA	R9	MinIon	468	1785	9	insect	LFB	170	3610	3305	18665	336	FAO86834
128.1	DNA	R9	MinIon	383	1424	6	insect	LFB	200	1543	1510	19752	113	FAO88446
129.1	DNA	R9	MinIon	487	1796	55	insect	LFB	190	4379	3694	6359	357	FAO88464
129.2	DNA	R9	MinIon	279	1046	57	insect	LFB	50	1507	NA	6788	311	FAO88464
130.1	DNA	R9	MinIon	511	1944	44	insect	SFB	200	9475	6638	4590	922	FAO91712
130.2	DNA	R9	MinIon	464	1753	48	insect	SFB	100	6429	4884	4961	692	FAO91712
130.3	DNA	R9	MinIon	425	NA	51	insect	SFB	67	2592	2592	4828	525	FAO91712
67.1	DNA	R9	MinIon	499	1907	8	insect	LFB	200	4891	4586	18872	455	FAO92001
67.2	DNA	R9	MinIon	485	1852	9	insect	LFB	170	4284	3895	19414	417	FAO92001
67.3	DNA	R9	MinIon	473	1776	13	insect	LFB	140	3997	3559	18674	368	FAO92001
67.5	DNA	R9	MinIon	427	1508	44	insect	SFB	78	4948	3728	3529	607	FAO92001
67.6	DNA	R9	MinIon	111	758	51	insect	LFB	30	648	578	14885	408	FAO92001
131.1	DNA	R9	MinIon	512	1959	55	insect	LFB	190	7742	6090	6314	803	FAO92100
131.2	DNA	R9	MinIon	498	1864	57	insect	LFB	100	3275	NA	6876	NA	FAO92100
131.3	DNA	R9	MinIon	469	1772	57	insect	LFB	85	5929	4737	4932	894	FAO92100
131.4	DNA	R9	MinIon	380	1421	58	insect	LFB	60	3028	2746	6700	370	FAO92100
131.5	DNA	R9	MinIon	223	874	62	insect	LFB	70	807	778	6584	264	FAO92100
132.1	DNA	R9	MinIon	465	1812	57	insect	LFB	100	4724	4089	7116	498	FAO92220
132.2	DNA	R9	MinIon	359	1379	58	insect	LFB	60	2643	2390	6769	334	FAO92220

ID	RNA/ DNA	fc version	device	#cha	#por	fc age	sample type	buffer	ng load	Estimated bases	EB12	mean rl	pore hl	fc id
133.1	DNA	R9	MinIon	506	1951	58	insect	LFB	190	6564	5533	6795	688	FAO92291
133.2	DNA	R9	MinIon	433	1628	62	insect	LFB	85	3866	3453	7461	426	FAO92291
134.1	DNA	R9	MinIon	488	1725	72	plant	SFB	350	8694	3799	2288	620	FAR96897
135.1	DNA	R9	MinIon	473	1615	72	plant	SFB	330	9350	3929	4101	532	FAR96905
136.1	RNA	R9	MinIon	511	1399	128	mouse	RNA	1252	1407	910	1058	915	FAO83512
137.1	RNA	R9	MinIon	509	1383	128	mouse	RNA	1372	832	705	746	638	FAO92305
138.1	RNA	R9	MinIon	494	1275	12	mouse	RNA	1092	1206	680	733	1094	FAP46096
139.1	RNA	R9	MinIon	474	1263	12	mouse	RNA	1252	1174	708	772	1176	FAP46400
140.1	RNA	R9	MinIon	501	1353	12	mouse	RNA	1220	1940	888	1049	1390	FAP46452
141.1	RNA	R9	MinIon	486	1320	12	mouse	RNA	1044	1337	800	1091	827	FAP49780
142.1	DNA	R9	MinIon	494	NA	86	mouse	SFB	385	10671	6305	10446	1935	FAQ58673
142.2	DNA	R9	MinIon	NA	1236	NA	mouse	SFB	385	NA	2280	9714	370	FAQ58673
143.1	DNA	R9	MinIon	479	1827	63	mouse	SFB	300	6680	4769	8821	599	FAQ58674
144.1	DNA	R9	MinIon	447	1714	57	mouse	SFB	300	5561	4246	9428	612	FAQ61173
145.1	DNA	R9	MinIon	499	NA	83	mouse	SFB	245	11137	6360	10034	1800	FAQ61176
145.2	DNA	R9	MinIon	NA	1205	NA	mouse	SFB	245	NA	2779	9350	510	FAQ61176
146.1	DNA	R9	MinIon	508	1949	63	mouse	SFB	282	21067	6967	9473	1993	FAQ69208
147.1	DNA	R9	MinIon	508	1875	83	mouse	SFB	245	19740	6786	9155	1855	FAQ69274
148.1	DNA	R9	MinIon	499	1909	83	mouse	SFB	385	19034	6723	9377	1832	FAQ69338
149.1	DNA	R9	MinIon	507	1895	49	mouse	SFB	270	24681	7030	6962	2518	FAQ69370
150.1	DNA	R9	MinIon	490	1829	57	mouse	SFB	280	22628	6920	8963	2237	FAQ69578
151.1	DNA	R9	MinIon	509	1964	170	human	SFB	444	18354	6760	1550	2234	FAL84151
151.2	DNA	R9	MinIon	213	728	172	human	SFB	444	2207	1506	1485	737	FAL84151
152.1	RNA	R9	MinIon	505	1360	47	human	RNA	42	157	117	475	431	FAO10420
153.1	DNA	R9	MinIon	502	NA	68	human	SFB	538	12217	6859	2414	1762	FAP46521
154.1	DNA	R9	MinIon	489	NA	43	human	SFB	451	10568	6294	4487	1612	FAP49434
154.2	DNA	R9	MinIon	448	1708	44	human	SFB	451	13249	5635	4181	1369	FAP49434
155.1	DNA	R9	MinIon	NA	1919	NA	human	SFB	688.8	NA	6763	4626	2319	FAP49725
156.1	DNA	R9	MinIon	501	1928	189	human	LFB	400	12833	NA	5179	1386	FAQ01752
157.1	DNA	R9	MinIon	467	1430	57	human	SFB	382	7352	4147	515	712	FAQ09810
158.1	DNA	R9	MinIon	480	1356	49	human	SFB	821	6529	4541	454	899	FAQ10598
158.2	DNA	R9	MinIon	187	536	50	human	SFB	547	2326	1799	467	791	FAQ10598
158.3	DNA	R9	MinIon	76	239	51	human	SFB	274	299	227	854	NA	FAQ10598
159.1	DNA	R9	MinIon	488	1878	162	human	SFB	300	13259	5853	2776	1275	FAQ69371
160.1	DNA	R9	MinIon	476	NA	92	human	SFB	400	15275	5655	2746	1314	FAR29925
161.1	DNA	R9	MinIon	NA	1690	238	human	LFB	405	NA	4401	2257	544	FAR33296
161.2	DNA	R9	MinIon	293	866	NA	human	LFB	405	2509	1905	927	350	FAR33296
162.1	DNA	R9	MinIon	509	NA	123	human	LFB	394	19595	NA	NA	1491	FAR33573
163.1	DNA	R9	MinIon	456	NA	130	human	LFB	305	8441	NA	643	759	FAR38117
164.1	DNA	R9	MinIon	505	NA	27	human	LFB	365	9027	NA	NA	935	FAR39219
165.1	DNA	R9	MinIon	498	NA	37	human	LFB	321	7285	NA	765	750	FAR39290
166.1	DNA	R9	MinIon	508	NA	74	human	LFB	453	6073	NA	944	834	FAR39317
167.1	DNA	R9	MinIon	377	1197	242	human	LFB	340	3463	2744	677	356	FAR39502
168.1	DNA	R9	MinIon	490	NA	91	human	SFB	400	17476	5708	4673	1212	FAR93058
169.1	DNA	R9	MinIon	383	993	247	human	LFB	340	2266	1726	848.	221	FAR96341
170.1	DNA	R9	MinIon	481	1833	92	human	SFB	400	15910	5763	5055	1078	FAR96408
171.1	DNA	R9	MinIon	483	1777	91	human	SFB	400	10105	4151	2782	1001	FAR96434
172.1	DNA	R9	MinIon	384	1080	249	human	LFB	425	3106	2509	1029	397	FAR96514
173.1	DNA	R9	MinIon	510	1882	37	human	SFB	300	7049	5165	1412	718	FAR96669
173.2	DNA	R9	MinIon	300	1079	38	human	SFB	200	5071	2886	1414	726	FAR96669
174.1	DNA	R9	MinIon	479	1294	238	human	LFB	389	5313	3079	849	446	FAR96893
175.1	DNA	R9	MinIon	438	1308	171	human	LFB	405	3491	2671	947	630	FAS60674
176.1	DNA	R9	MinIon	509	NA	90	human	LFB	450	20857	6391	587	2090	FAV38963
176.2	DNA	R9	MinIon	NA	728	91	human	LFB	411	NA	1729	1164	584	FAV38963
177.1	DNA	R9	MinIon	NA	NA	69	human	LFB	400	NA	NA	NA	NA	FAV38989
177.2	DNA	R9	MinIon	452	1445	NA	human	LFB	330	8250	4531	907	984	FAV38989
178.1	DNA	R9	MinIon	511	NA	60	human	LFB	400	14634	5579	622	1222	FAV39381
179.1	DNA	R9	MinIon	NA	1437	NA	human	LFB	412	NA	6287	679	1644	FAV87040
180.1	DNA	R9	MinIon	NA	1371	NA	human	LFB	414.96	NA	5864	569	1207	FAW80737
9.2	RNA	R9	MinIon	270	694	83	synthetic	RNA	230	505	362	893	801	FAO07649
124.7	RNA	R9	MinIon	189	484	103	synthetic	RNA	207	201	183	1002	469	FAO12153
125.7	RNA	R9	MinIon	200	522	103	synthetic	RNA	212	195	179	1038	418	FAO12159

Figure S1. Screenshot from the Nanopore Community about DNA control sequence (DCS)

ACCESS RAW ONT DATA

The following equation is used to convert the squiggles stored in the raw data to the pA signal:

$$pA\_signal = (signal + offset) * range / digitization \qquad (1)$$

where `signal` represents the raw integer value, `offset` is the shift in the pA signal, `range` is the range of the pA signal, and `digitization` is the resolution of the conversion used to convert the continuous pA signal into discrete integer values. All parameters are provided by ONT in the raw data files.

NORMALIZATION

Normalization corrects for biases, ensuring that signals from different experiments or sequencing channels are directly comparable. It can be performed on each read individually using statistical measures such as the mean and standard deviation or the median and median absolute deviation (MAD). The median and MAD are more robust to outliers, which are common in sequencing data, see Fig. 10. ONT tools, including their basecallers, typically employ normalization during data processing. The standard formula for signal normalization, based on the median and MAD, is as follows:

$$norm\_signal = \frac{pA\_signal - median(pA\_signal)}{mad(pA\_signal)} \qquad (2)$$

This process ensures that the signal distributions across reads become uniform, eliminating biases (see Fig. **??**B).

Table S2. Overview of methylation calling tools (adapted from Liu *et al.*), sorted by publication date. 5mC – 5-methylcytosine in DNA; m5C – 5-methylcytosine in RNA; 6mA – N6-methyladenosine in DNA; m6A – N6-methyladenosine in RNA; 5hmC – 5-hydroxymethylcytosine; psU – pseudouridine; m1A – N1-methyladenosine; 5moU – 5-methoxyuridine; m7G – N7-methylguanosine; Ino – inosine; f5C – 5-formylcytidine;

comp. approach	tool	DNA / RNA	modification type	year	citation
hidden Markov models	Nanopolish	DNA	5mC	2017	(79)
statistical	Tombo	DNA / RNA	4mC, 5mC, m5C, 6mA	2017	(86)
statistical	SignalAlign	DNA	5mC, 5hmC, 6mA	2017	(87)
neural networks	Dorado	DNA	5mC, 6mA		ONT[1]
statistical	NanoMod	DNA	5mC	2019	(88)
neural networks	mCaller	DNA	6mA	2019	(89)
neural networks	DeepSignal	DNA	5mC, 6mA	2019	(90)
neural networks	DeepMod	DNA	5mC, 6mA	2019	(91)
neural network	DeepMod2	DNA	5mC	2024	(92)
neural networks	Megalodon	DNA	5mC, 6mA		ONT[2]
hidden Markov models	f5c	DNA	5mC	2020	(75)
neural networks	methBERT	DNA	5mC, 6mA	2021	(93)
meta approaches	METEORE	DNA	5mC, 6mA	2021	(21)
neural networks	DeepMP	DNA	5mC, 6mA	2021	(94)
statistical	–	DNA	D2O	2024	(85)
machine learning	MINES	RNA	m6A	2019	(95)
error profile & machine learning	EpiNano	RNA	m6A, psU	2019	(96)
neural networks	nanoDoc	RNA	m6A	2020	(97)
neural networks	nano-ID	RNA	e5U	2020	(98)
statistical	DRUMMER	RNA	m6A	2020	(99)
statistical	xPore	RNA	m6A	2021	(100)
machine learning	Nanom6A	RNA	m6A	2021	(101)
statistical	Yanocomp	RNA	m6A	2021	(?)
machine learning	nanoRMS	RNA	psU	2021	(102)
error profile	ELIGOS	RNA	m6A, m1A, 5moU, psU, m7G, Ino, hm5C, f5C	2021	(103)
statistical	Nanocompore	RNA	m6A	2021	(104)
neural networks	m6Anet	RNA	m6A	2022	(105)
machine learning	Penguin	RNA	psU	2022	(106)
neural networks	DENA	RNA	m6A	2022	(107)
statistical	Magnipore	RNA	-	2023	(108)
neural networks	mAFIA	RNA	m6A	2024	(109)

[1] https://github.com/nanoporetech/dorado
[2] https://github.com/nanoporetech/megalodon

Figure S2. Screenshot from the Nanopore Community about RNA control sequence (RCS)

Figure S3. T-test shows higher estimated number of sequenced bases using SFB compared to LFB buffer after 24 h (**A**) and 36 h (**B**); addition to Fig. 4.

Figure S4. Correlation (R^2) between pore half life and the estimated number of bases after 12 hours of sequencing (EB12) is observed across most sample types. The EB12 values were divided by number of starting pores, to corrected for any potential bias. The resulting plots reflect the same statements as Fig. 6A.

Figure S5. T-test shows higher EB12 in mouse compared to other sample types, addition to Fig. 6B.

Figure S6. T-test shows significant differences in pore half time between sample types; addition to Fig. 6C.

Figure S7. Distribution of estimated number of sequenced bases after 24 h (**A**) and 36 h (**B**) for DNA (red) and RNA (blue) samples; addition to Fig. 6D.

Table S3. Re-evaluated hand picked pore half time for Different Run IDs

ID	Half Life
1.1	998
2.1	1630
6.1	1510
10.1	386
88.1	3098
93.1	2041
110.1	1750
130.3	525
142.1	1935
145.1	1800
153.1	1762
154.1	1612
160.1	1314
162.1	1491
163.1	759
164.1	935
165.1	750
166.1	834
168.1	1212
178.1	1222
176.1	1971
176.1	2090

Figure S9. T-test shows significant differences between the buffers used for R9 data.

Figure S8. T-test shows no significant differences between the buffers used for R10 data. This might be due to the low number of data points.

Table S4. Sequencing statistics for the samples used in Fig. 7. cum. yield (Gb) – cumulative amount of estimated bases for all sequencing runs on this flow cell. done

	ID 156	ID 124
#pores	1 928	1 843
#channels	501	497
# wash steps	2	6
cum. yield (Gb)	12.83	11.13
sample type	human	insect
fc version	R9	R9
device	MinION	MinION
RNA/ DNA	DNA	DNA

Table S5. Samples used for adaptive sampling in Fig. 8. # por – amount of active pores at sequencing start; yield – cumulative sequencing yield in Gb; depth – mean sequencing depth on CpG islands; wash – number of times the flow cell was washed in between; buffer – extension length of the CpG islands

id	# por	yield	depth	wash	buffer
NA	NA	10.7	3.01	0	0
NA	1519	19.7	6.10	0	0
155	1504	16.7	4.75	0	0
154	1305	23.3	6.67	1	0
153	1379	11.4	3.28	0	1
153	822	2.58	1.39	0	1
NA	1001	2.59	0.67	0	500
158	1192	8.65	2.13	2	500
157	1269	8.72	4.76	1	500
156	1250	13.49	3.85	2	0
164	1405	9.68	8.79	1	2000
165	1433	7.74	8.88	2	2000
166	1548	6.39	7.60	2	2000
162	1620	21.26	17.79	1	2000
163	969	9.00	7.24	1	2000
174	1352	6.46	5.40	1	2000
167	693	3.77	3.42	0	2000
169	749	2.60	2.13	0	2000
172	787	3.51	3.08	0	2000
161	1063	5.45	1.52	1	0

Figure S10. For the data shown in Fig. 6 B,C a t-test reveals no significant differences in estimated bases after 12 hours and pore half time between R9 and R10 data. However, when separating the data based on sample types, both EB12 and pore half time show significant differences for the metagenomic samples.

Figure S11. Read mean distribution (top) and read standard deviation (bottom) before (left) and after (right) normalization of six randomly picked channels from the sequencing runs 5.1 and 12.1. Biases caused by pores, flow cells, or sensors vanish after normalization.