Deciphering the actin structure-dependent preferential cooperative binding of cofilin

Kien Xuan Ngo[1,#,*], Huong T Vu[2,#], Kenichi Umeda[1], Noriyuki Kodera[1], Taro Q.P. Uyeda[3]

[1]Nano Life Science Institute (WPI-NanoLSI), Kanazawa University, Kanazawa, Ishikawa 920-1192, Japan.
[2]Centre for Mechanochemical Cell Biology, Warwick Medical School, Coventry CV4 7AL, UK.
[3]Department of Physics, Faculty of Advanced Science and Engineering, Waseda University, Shinjuku, Tokyo 169-8555, Japan.

# These authors equally contributed.
* Corresponding authors
Email: ngoxuankien@staff.kanazawa-u.ac.jp

Abstract
The mechanism underlying the preferential and cooperative binding of cofilin and the expansion of clusters toward the pointed-end side of actin filaments remains poorly understood. To address this, we conducted a principal component analysis based on available filamentous actin (F-actin) and C-actin (cofilins were excluded from cofilactin) structures and compared to monomeric G-actin. The results strongly suggest that C-actin, rather than F-ADP-actin, represented the favourable structure for binding preference of cofilin. High-speed atomic force microscopy explored that the shortened bare half helix adjacent to the cofilin clusters on the pointed end side included fewer actin protomers than normal helices. The mean axial distance (MAD) between two adjacent actin protomers along the same long-pitch strand within shortened bare half helices was longer (5.0–6.3 nm) than the MAD within typical helices (4.3–5.6 nm). The inhibition of torsional motion during helical twisting, achieved through stronger attachment to the lipid membrane, led to more pronounced inhibition of cofilin binding and cluster formation than the presence of inorganic phosphate (Pi) in solution. F-ADP-actin exhibited more naturally supertwisted half helices than F-ADP.Pi-actin, explaining how Pi inhibits cofilin binding to F-actin with variable helical twists. We propose that protomers within the shorter bare helical twists, either influenced by thermal fluctuation or induced allosterically by cofilin clusters, exhibit characteristics of C-actin like structures with an elongated MAD, leading to preferential and cooperative binding of cofilin.
Keywords: actin, cofilin, actin binding proteins, high-speed atomic force microscopy, principal component analysis.

Introduction
　　　　Cofilin is a member of the actin-depolymerizing factor/cofilin (ADF/cofilin) family that is present in all eukaryotes. There are two major isoforms of cofilin in mammals: cofilin 1 and cofilin 2, predominantly found in non-muscle and muscle tissues, respectively (Bamburg and Bernstein, 2010). Cofilin is widely recognized as a key regulator of actin filament dynamics, particularly in nonequilibrium assembly and disassembly processes. Cofilin promotes the depolymerization and severing of actin filaments in a concentration-dependent manner (Andrianantoandro and Pollard, 2006) and collaborates or competes with other actin binding

proteins (ABPs) in this process (Bibeau et al., 2021; Grintsevich et al., 2016; Lappalainen and
Drubin, 1997; Ngo et al., 2016).

The structure of an actin filament has traditionally been represented as a simplistic atomic
model derived from the G-actin structure in the Holmes model (Holmes et al., 1990) without
considering the conformational changes occurring within each protomer during actin
polymerization (Kasai et al., 1962; Rich and Estes, 1976). The rigid-body rotation of the G-actin
crystal structure used in the Holmes model prompted the development of accurate models that
consider structural changes within protomers, such as the X-ray diffraction Lorenz model
(Lorenz et al., 1993) and Oda model (Oda et al., 2009).

The actin protomer consists of two domains, the inner domain (ID) and the outer domain
(OD), with relatively limited contact between them. The polypeptide chain traverses between
these domains twice: at the loop centered around residue Lys336 and at the linker helix Gln137-
Ser145, which functions as the axis of a hinge between the domains. As a result, two clefts are
formed between these domains (Dominguez and Holmes, 2011). The upper cleft, located
between subdomain (SD2) and SD4, serves as a binding site for nucleotides and divalent ions
like Mg^{2+}. On the other hand, the lower cleft, located between SD1 and SD3, is characterized by
residues Tyr143, Ala144, Gly146, Thr148, Gly168, Ile341, Ile345, Leu346, Leu349, Thr351, and
Met355, which are predominantly hydrophobic. This cleft serves as the primary binding site for
most ABPs including cofilin and is also responsible for critical longitudinal contacts between
actin protomers within the filament (Dominguez, 2004; Fujii et al., 2010; Oda et al., 2009).

The actin filament is a double-stranded helical structure formed by the crossover of two
long-pitch helices; the distance between the crossover points is known as the half helical pitch
(HHP), which is approximately 36 nm (Fujii et al., 2010; Holmes et al., 1990; Lorenz et al.,
1993; Oda et al., 2009). A canonical half helix is composed of 13 actin protomers (6.5 protomer
pairs). Thus, the mean axial distance (MAD) between two adjacent protomers along the same
long-pitch strand is 5.5 nm (36 nm/(13/2)) (Egelman, 1994). However, long actin filaments
exhibit variable helical twists (Egelman et al., 1982). Recent cryo-EM studies have demonstrated
the atomic structures of short actin segments with cofilin (referred to as cofilactin) (Galkin et al.,
2011; Huehn et al., 2020; Tanaka et al., 2018). When cofilin binds to actin in a 1:1 ratio to form
a cofilactin segment, the helices became supertwisted, and the HHP is decreased by
approximately 25% (McGough et al., 1997), reducing the number of actin protomers to ~11 in
each half helix (5.5 protomer pairs) (PDB 3J0S) (Galkin et al., 2011). In a simple model,
assuming that the MAD remains unchanged in both bare actin and cofilactin segments, the
difference in the number of actin protomers impacts the formation of short and long HHPs.
Nonetheless, whether the alterations in MAD dynamics are comparable between normal and
supertwisted half helices remains uncertain. Therefore, additional research is needed to
investigate this particular issue.

In this study, we utilized several techniques, including principal component analysis
(PCA) and high-speed atomic force microscopy (HS-AFM). We aimed to understand how the
allosteric conformational changes induced by cofilin clusters in neighboring bare actin filaments
contribute to the preferential binding of additional cofilin molecules toward the pointed-end (PE)
side, as reported in our previous study (Ngo et al., 2015). Therefore, we generated pseudo AFM
images representing long bare actin filaments (referred to as F-actin structures) and cofilactin
filaments without cofilin molecules (referred to as C-actin structures) using available protein data
bank (PDB) structures such as 6VAU (bare actin filament from a partially cofilin-decorated
sample) and 6VAO (human cofilin-1 decorated actin filament) (Huehn et al., 2020). We

measured the HHPs and the number of actin protomers per HHP in pseudo-AFM images and
used them as the reference values for the data obtained from our real HS-AFM analyses. In
principle, we classify that filament regions with supertwisted half helices and a reduced number
of protomer pairs per HHP, as detected in our AFM measurements, are indicative of C-actin like
structure. We aimed to address several issues: (**i**) Does the number of actin protomers per HHP
vary between typical F-actin and C-actin like structure regions? (**ii**) Does the MAD change, and
do these changes affect cofilin binding within the unbound actin regions adjacent to cofilin
clusters on the PE and barbed-end (BE) sides? Are these changes quantitatively similar or
distinct? (**iii**) Can we evaluate the impact of Pi on torsional flexibility during helical twisting of
actin filaments to elucidate how they influence the preferential cooperative binding and
expansion of cofilin clusters?
An innovative and previously unexplored facet of this study is the visualization of the
actin protomer's structure within a bare half helix adjacent to small cofilin clusters on the PE and
BE sides using HS-AFM. Our findings elucidate the dynamic structural changes in half helices
and the number of actin protomers per HHP in normal F-actin. Our results deviated from the
conventional ensemble-averaged helical twists usually observed in EM and X-ray diffraction,
depicting a rigid half helix. Moreover, we show that shortened bare half helices adjacent to the
cofilin clusters on the PE side contain less actin protomers than conventional half helices. The
MAD within the shortened bare HHP (C-actin like structure) is longer than the MAD within the
canonical HHP (F-actin structure). This elongated MAD can be interpreted as resulting from the
allosteric transmission of structural changes originating in the cofilactin region, extending
beyond the boundary between cofilactin and bare actin regions. This extended influence affects
the structure of the shortened bare half helix on the PE side. Additionally, our findings suggest
that the cooperative binding of cofilin is significantly impacted by the torsional flexibility of
actin filaments. The inhibition of torsional motion during helical twisting, achieved via physical
perturbation, inhibited cofilin binding and the expansion of larger clusters more than Pi binding.
We also verify our previous hypothesis, which supports cofilin's initial binding to the
favorable structure of actin protomers because F-actin, without ADF/cofilin proteins, could exist
in the 162° twist state observed for cofilin-decorated F-actin (Galkin et al., 2011, 2001; Ngo et
al., 2016, 2015). Our hypothesis focuses on the evidence of decreasing the number of protomers
within the bare half helix adjacent to the cofilin cluster on the PE side. This reduction leads to a
shorter HHP but paradoxically results in a longer MAD in the C-actin like structure area. The
protomers within a shortened bare half helix near cofilin clusters are converted to C-actin like
structures via our proposed F-actin to C-actin transition pathway, thereby facilitating the
expansion of clusters on the PE side. If cofilin molecules fail to promptly attach to protomers
within the shortened bare region, the protomers resembling the C-actin structure in that region
can quickly change the closed nucleotide binding clefts to the open state and transition to a G-
actin state. This transition (C-actin to monomer G-actin like structure) ultimately results in
severing of the filament at or near the boundary of cofilactin and bare actin regions. Furthermore,
this study suggests that Pi binding mainly inhibits the initial binding of cofilin by reducing the
fraction of supertwisted half helices in F-ADP.Pi-actin, whereas the removal of Pi enhances the
initial binding of cofilin by elevating the fraction of supertwisted half helices in F-ADP-actin.
Nevertheless, Pi binding only slightly inhibits the propagation of the supertwisted helical
structure within cofilin clusters to bare half helices on the PE side in F-ADP.Pi-actin when
compared to F-ADP-actin. Together, this clarifies that the initial interactions of cofilin with F-

actin and the subsequent expansion of clusters are closely linked to the variability in helical
twists.
139
Results
Principal component analysis suggests C-actin like structure as the optimal actin structure
for cofilin binding
To investigate the relationship between actin structure and cofilin binding, we explored
the collective changes in actin conformations as they transitioned from G-actin to F-actin and
then to C-actin. We applied PCA to identify the principal components (PCs), as depicted in
Figure 1. Our analysis included an array of actin chains sourced from 41 available PDB
structures of filamentous actin protomers bound to cofilins, profilin, DNAse I, and various
nucleotides (ATP, AMPPNP, ADP.Pi, ADP, and Apo). We then compared these findings with 5
additional PDB structures featuring free G-actin and twinfilin-1-bound G-actin (**Table**
supplement 1).
We performed PCA with the same dataset using different alignments for subdomains 1
(SD1), 2 (SD2), 3 (SD3), and 4 (SD4), the outer domain (OD, SD1 + SD2), the inner domain
(ID, SD3 + SD4), and the entire actin structure. In all cases, we observed clustering of different
groups of actin chains. Interestingly, aligning only the IDs of all actin chains yielded the most
meaningful PCs. Importantly, PCA can elucidate the conformational transition between C-actin
and G-actin structures, and vice versa, through PC1, while PC2 delineates the pathway between
F-actin and C-actin structures (**Figure 1A**). To visualize the conformational changes associated
with each PC, we back-calculated and depicted the PCs in **Figure 1B and 1C**, respectively.
These illustrations revealed two outer domain motion modes. The first mode (PC1, accounting
for 45% of the structural variance) involved rotations of the OD around the hinge region to seal
the nucleotide binding cleft by reducing the SD2-SD4 distance (as indicated by the yellow
arrowhead in **Figure 1B and Video 1**), depicting the transformation between G-actin and C-
actin. Structures with higher PC1 values such as F-actin and C-actin are associated with
conformations featuring a closer nucleotide binding cleft. Conversely, structures with lower PC1
values, including tilted-actin and G-actin, are related to conformations featuring a more open
nucleotide binding cleft. The second prominent mode (PC2, accounting for 28% of the structural
variance) included the twisting of the OD during the transition between F-actin and C-actin
(**Figure 1C and Video 1**). A higher PC2 value indicates that the OD is more flattened or
untwisted, as seen in the tilted-actin and F-actin. On the other hand, a lower PC2 value suggests
that the OD is more twisted, as observed in G-actin and C-actin.
Our PCA results indicated that the structure of an actin filament with singly bound cofilin
– 6UBY (orange box in **Figure 1A**) was in the transition pathway between the F-actin and C-
actin conformations, particularly along PC2. This cluster also includes structures of two
protomers positioned at the BE side near a cofilactin cluster (including chains E&H combined as
E and F&G combined as F in PDB: 6UC4 (**Table supplement 1**)). Consistent with previous
studies (Galkin et al., 2011; Tanaka et al., 2018), our analysis suggests that the attachment of a
single cofilin to an actin filament induces a slight twisting of the OD relative to the ID (~5°).
Furthermore, this effect was cumulative, with more extensive cofilin binding leading to larger
OD twisting angles (~10°), thus representing the C-actin state. Throughout this transition, the
nucleotide binding cleft remains closed as observed through the minimal variations in PC1
values. This suggests that cofilin prefers to bind to the C-actin structure and the transitional actin
states featuring structures resembling C-actin within the actin filaments. However, we also

predicted that cofilins do not preferentially bind to the tilted structural state of actin, as tilted
actin structures exhibited opened nucleotide-binding clefts, but without the twisting of the OD
(Galkin et al., 2015) (indicated by lower PC1 and higher PC2 values, as represented by the
magenta box in **Figure 1A**). Additionally, these tilted actin structures did not align with the
conformational transition pathway between F-actin and C-actin.
Noticeably, PCA analysis revealed higher structural flexibility in F-ADP-actin (red dots),
exploring a larger space than F-ADP-Pi-actin structures (orange dots) within the F-actin cluster
(inset in **Figure 1A**). This included the F-ADP-actin structure (3MFP) (Fujii et al., 2010), which
notably exhibited a PC2 value significantly approaching that of C-actin structure (**Figure 1A**).
However, upon comparison of structures simultaneously resolved in (Chou and Pollard, 2019), a
slightly smaller PC2 value for the F-ADP-actin (6DJO) was observed in contrast to F-AMPPNP-
actin (6DJM) and F-ADP.Pi-actin (6DJN) (inset of **Figure 1A**). Significantly, the magnified
views in the F-actin cluster highlight structural resemblance between F-ADP-actin and F-
ADP.Pi-actin in recent studies (Oosterheert et al., 2022; Reynolds et al., 2022). The F-ADP-actin
structure (8D13) and F-ADP.Pi-actin structure (8D14) exhibit remarkably similar PC2 values
(**Figure supplement 1A**). Furthermore, the PC2 results demonstrate notable similarities between
Mg^{2+}-F-ADP-actin (8A2T) and Mg^{2+}-F-ADP.Pi-actin (8A2S), and Mg^{2+}-F-ADP.BeF$_3$-actin
(8A2R) (**Figure supplement 1B**).
In order to better understand the effect of the collective PC2 movements of individual
ODs on the global half helices of filaments, we compared the side views and outer views of F-
actin and C-actin protomers in their filamentous structures (**Figure 2A-C**). The twisting
movement of OD that we found in **Figure 1C** is best represented again from the side views with
the rotation axes (marked as red stars in blue boxes of **Figure 2A-B**). We identified that cofilin
clusters rotate the ODs approximately 10°, burying the SD2 including D-loops deeper into the
core of the filament. As a result, the twisting angles between two adjacent protomers in the same
strand appears to increase from approximately 26° to around 36° if view from the outer side
(black boxes of **Figure 2A-B**), as reported before (Galkin et al., 2011; Tanaka et al., 2018). Such
change in the twisting angles from the outer views can be represented as the change of the OD
lines on 2D surfaces of imaginary cylinders of the filaments, unwrapped along imaginary lines
that an AFM tip would scan through (**Figure 2D-E**). Such diagrams depict the varying twisting
angles of OD domains within F-actin and C-actin (PC2 mode), leading to the formation of HHPs
with different numbers of detected protomer pairs. Hence, collective movements of individual
ODs that decrease PC2 values would lead to a globally shorter HHPs and fewer number of
protomer pairs per HHP.
217
The half helices within long actin filaments exhibit heterogeneity in their lengths and the
number of protomers per half helical pitch when observed through HS-AFM
In our actual AFM measurements, we determined the half helices, the number of
protomer pairs per HHP, and MAD as illustrated in the pseudo-AFM images representing F-
actin, C-actin, and cofilactin structures (**Figure supplement 2, Table supplement 2 and**
Methods). To investigate the presence of random variable helical twists in long actin filaments
in solution, we employed HS-AFM to examine the dynamic structures and distribution of actin
protomers within multiple half helices, with high spatial (1 nm in the xy plane and 0.1 nm in the
z-axis) and temporal (2 frames per second) resolution. As depicted in **Figures 3 and 4**, **Table**
supplement 3, and **Video 2**, we successfully imaged and quantified the number of actin
protomers within many half helices, allowing us to precisely measure the distribution of the

number of protomer pairs per HHP (**Figure 3A**). Subsequently, we established a correlation
between the HHP and the number of protomer pairs per HHP. We also calculated the HHPs
assuming a constant MAD of 5.5 nm and compared them with actual data (**Figure 3B**). We
analyzed the linear regression analysis results and noted that the slope derived from our actual
AFM measurements was smaller than that computed assuming that the MAD remained
unchanged.
To address this discrepancy, we analyzed our measurements, revealing a range of
protomer pairs within each HHP, varying from 4.5 to 8.5 with distinct proportions (**Figure 3C,
Table supplement 3**). The most commonly observed numbers of protomer pairs per HHP were
6.5 and 7.5, accounting for 60.9% (n = 1355 HHPs) and 28.0% (n = 624 HHPs) of the samples,
respectively. Conversely, the least frequently observed numbers of protomer pairs per HHP were
4.5, 5.5 and 8.5, accounting for 0.2% (n = 5 HHPs), 8.2% (n = 182 HHPs) and 2.7% (n = 60
HHPs) of the samples, respectively. Subsequently, we computed the MAD by dividing the length
of each measured HHP by the corresponding number of protomer pairs per HHP, thereby
establishing a correlation between these variables (**Figure 3C**). The MAD values (mean ± SD)
for HHPs containing 6.5 and 7.5 were 5.1 ± 0.2 nm (n = 8807.5 pairs) and 5.0 ± 0.3 nm (n =
4680 pairs), respectively. Corresponding values for HHPs with 5.5 and 8.5 protomer pairs were
5.3 ± 0.3 nm (n = 1001 pairs) and 4.6 ± 0.3 nm (n = 510 pairs) (**Table supplement 3**). Notably,
we observed that HHPs with fewer protomer pairs exhibited elongated MADs. Our analysis also
indicates an extremely uncommon occurrence, with approximately 0.2% of the actin protomers
within the F-actin structure (4.5 protomer pairs per HPP) demonstrating elongated MADs (6.0 -
6.6 nm).
We next conducted a temporal autocorrelation function (ACF) analysis aimed at
estimating the time constants governing the decaying self-correlation in the axial distance (AD)
between two adjacent actin protomers along the same long-pitch strand (**Figure 3D-F**). The core
objective of this analysis was to determine the temporal extent of AD similarity and elucidate the
dynamic behaviors and conformational changes in the distance between adjacent actin protomers
over time. The results revealed that self-correlation of the ADs within actin filaments,
immobilized on lipid membrane, rapidly decreased within a few hundred milliseconds and then
gradually decayed within a few seconds (**Figure 3E-F**). These results could potentially account
for the fluctuation observed in HHPs, the number of protomer pairs per HHP, and the MADs
seen within the same actin filaments over time (**Figure 4A-B**).
To assess the degree of spatial clustering in the structure of actin protomers, we utilized a
spatial ACF. This analysis offered a quantitative evaluation of the consistency in the amplitude
variations among actin protomers over different distances within long actin filaments, observed
across several time points. The height amplitude of protomers, used to measure the periodicity of
the distance between protomers and the slopes in heights between them, reflects the surface
corrugation formed by protomers within the filaments. The spatial ACF analysis provides an
indirect measure of the degree to which height values near each other in space are similar to one
another in actin filaments. If protomers can change the shape and/or rotate around the axis
parallel to the filament axis, the individual protomers may have slightly different thickness
and/or orientations, leading to variations in the heights of protomers and, consequently,
variations in the periodicity of distances and the slopes in heights between protomers (depicted in
Figure supplement 3, Methods). As seen in **Figure 4C**, many spatial ACF curves (as depicted
by the envelope curves) consistently exhibited a decreasing then increasing pattern for almost
every half helix, suggesting the randomness in the protomer distance periodicity, with the steep

downward and upward slopes in heights between protomers observed across each half helix
segment.
277
The shortened bare HHP adjacent to the cofilin clusters on the PE side includes fewer actin
protomers than the normal HHP, resulting in MAD elongation
Using HS-AFM, we aimed to simultaneously capture the helical structures and
distribution of actin protomers in the bare F-actin, cofilactin, and hybrid cofilactin/bare actin
segments within the same imaging area (see Methods). As shown in **Figure supplement 4**, we
successfully distinguished cofilactin and bare F-actin structures based on a height difference of
approximately 1-1.6 nm and a length difference in the half helices of approximately 25%. A bare
F-actin segment consisting of two consecutive half helices (HHP1, HHP2) exhibited the
expected normal lengths of 36 nm (6.5 pairs) and 36.5 nm (6.5 pairs) (**Figure supplement 4A**).
In contrast, a cofilactin segment with two consecutive half helices (HHP3, HHP4) displayed
shorter lengths of 28.6 nm (4.5 pairs) and 29.8 nm (4.5 pairs) (**Figure supplement 4B**). Next, we
presented a typical segment of the hybrid bare actin/cofilactin structure containing two
consecutive half helices (HHP5, HHP6), both showing comparably shortened lengths of 27.9 nm
(4.5 pairs) and 30.5 nm (4.5 pairs) (**Figure supplement 4C**). Based on our previous study (Ngo
et al., 2015), we judged that the bare half helix with the shorter HHP adjacent to the cofilin
cluster is on the PE side and that the normal HHP is located on the BE side. These typical data
suggest that, similar to cofilactin segments, shortened bare half helices adjacent to cofilin
clusters on the PE side include fewer actin protomers than normal half helices. We next analyzed
the MAD variation in this hybrid filament over time and made interesting observations (**Figure**
supplement 5). We discovered that MAD5 in bare actin HHP5 was elongated similarly to that in
cofilactin HHP6 and HHP7. Notably, MAD5 (5.7 ± 0.5 nm, n = 13) was extended and closely
resembled MAD6 (5.9 ± 0.3 nm) and MAD7 (6.1 ± 0.3 nm). This finding suggests that the MAD
elongation in the cofilactin region was transmitted beyond the cofilactin/bare actin boundary to
extend MAD5 in the shortened bare half helix.
We then conducted similar analyses on the hybrid cofilactin/bare actin filaments (**Figures**
5–6, **Video 3**). First, we observed the minimal impact of the lipid membrane on the half helices
(HHP1, HHP2), number of protomer pairs per HHP, and MADs (MAD1, MAD2) in normal actin
filaments (**Figure 5A**). We examined the bare half helices (HHP3–HHP6) and the number of
protomer pairs within each HHP on either side of a small cofilin cluster and found that the
shortened bare half helix on the PE side includes fewer actin protomers than the normal half
helix (**Figure 5B, 5C**). For example, the cofilin cluster (estimated as 2 to 4 molecules) shortened
one adjacent bare half helix (HHP6) on the PE side (25.5 nm, 4.5 pairs), while the half helix
(HHP5) on the BE side was similar to the normal half helix (37.9 nm, 7.5 pairs) (**Figure 5C**). In
this case, the MAD6 calculated within the shortened bare half helix (HHP6) was 5.7 nm and was
larger than the MAD5 value of 5.1 nm calculated within the normal half helix (HHP5) (**Figure**
5C). Despite a complete decoration requiring 9 to 11 cofilin molecules in a half helix, the
observation that 2 to 4 molecules can initiate the supertwisting of a half helix, achieved by
reducing the number of protomer pairs per HHP, strongly implies the presence of allosteric
transitioning in the supertwisting structures from the bound to the unbound region on the PE
side. We anticipate that the unbound region may contain approximately 1 to 2 pairs (2 to 4
protomers) resembling C-actin structures. When analyzing the fluctuation in the MADs over time
in this hybrid filament, we observed that MAD6 (5.4 ± 0.3 nm, n = 12) was significantly greater

than MAD5 (5.0 ± 0.2 nm, n = 12) (**Figure 6A, 6B**). Our analysis suggests that the MADs are longer in the shorter bare half helices than in the normal half helices (**Table supplement 4**).

We also analyzed a spatial ACF to demonstrate the extent of structural similarity between actin protomers between the bound and unbound cofilin regions. In contrast to the control actin filament (**Figure 4C**), the presence of small cofilin clusters led to a more pronounced linear decline in the spatial ACF for each half helix (**Figure 6C**). The results suggest that the periodicity of the protomer distances and the slope between the decorated and undecorated cofilin regions become more regular and are flatter than those in normal actin filaments. This result further suggests that the structural similarity among actin protomers in the cofilactin region extends beyond the half helix, encompassing neighboring bare actin regions.

The transmission of the supertwisted helical structure within the cofilin cluster to the adjacent bare half helices is essential in facilitating the preferential cooperative binding of cofilin

The cooperative binding of cofilin to actin filaments has been previously reported (Bobkov et al., 2006; McGough et al., 1997; Ngo et al., 2016, 2015). However, the role of cooperative conformational changes in actin filaments in driving the preferential binding of cofilin remains unclear, as indicated in several notable studies (Bibeau et al., 2021; Bobkov et al., 2006; Galkin et al., 2011; Hayakawa et al., 2014; Huehn et al., 2018, 2020; McCullough et al., 2008; Narita, 2020; Ngo et al., 2015; Suarez et al., 2011; Tanaka et al., 2018).

We observed the cooperative binding of cofilin to actin filaments when they were loosely absorbed onto a lipid membrane in an F-buffer with cofilin, ATP, and inorganic phosphate (Pi) (**Figure 7A**, **Video 4**). We also examined the peak heights and lengths of the half helices in the cofilactin and bare actin regions during the unidirectional growth of the cofilin clusters (**Figure 7A-C**). Consistent with our previous findings (Ngo et al., 2015), the cofilin cluster binding led to an approximately 1–2 nm increase in the peak height and an approximately 25% decrease in the HHPs in the cofilactin region. Then, we analyzed the relationship between the time needed to shorten the bare half helix beyond the cofilactin/bare actin boundary and to form a newly matured cofilin cluster within the shortened bare half helix. Finally, we determined the lag time for the growth of the cofilin cluster within the individual shortened bare half helices adjacent to the cofilin cluster.

The analysis suggested that even with 10 mM Pi, the cofilin cluster could supertwist the half helices in the actin filaments. Notably, the supertwisted half helices within the cofilactin region propagated beyond the cofilactin/bare actin boundary, shortening the adjacent bare half helix before the saturated cofilin cluster grew (**Figure 7A, 7B**, **Table supplement 5**). For example, when the HHP between P2 and P3 was shortened at 2 seconds, P2 was bound to a cofilin cluster, but P3 remained unbound. A lag time of approximately $t = 3$ s was needed for the saturated cofilin cluster to be recruited and grew in P3 at 5 seconds. This pattern was also observed when the HHPs between P3-P4, P4-P5, and P5-P6 shortened before the cofilin clusters expanded. These findings suggest that the shortening of the bare half helix beyond the cofilactin/bare actin boundary drives the actin structure-dependent preferential cooperative binding of cofilin clusters. The cooperative conformational changes during helical twisting may provide a mechanism for understanding the preferential binding of cofilin clusters toward the PE side, as reported in previous studies (Bibeau et al., 2021; Ngo et al., 2015).

As negative control experiments, we immobilized actin filaments onto lipid membranes (DPPC/DPTAP=75/25 wt%) to physically hinder the changes with helical twisting. We then

added a high concentration of cofilin (2000 nM) and observed its cooperative binding with actin filaments in the presence of ATP (**Figure 8A, Figure supplement 6A, Video 5**). However, the binding and expansion of cofilin clusters on these actin filaments were severely inhibited during long incubation times. Instead of cooperatively forming large clusters, cofilin formed only a few small clusters that transiently associated and dissociated, even at cofilin concentrations much higher than k_d (k_d = ~5–10 nM) (Andrianantoandro and Pollard, 2006). The histograms of the peak heights and HHPs were analyzed and showed single peaks at 8.2 ± 0.8 nm (n = 1930) and 34.3 ± 2.6 nm (n = 1556), respectively, which were similar to those of normal actin filaments (**Table supplement 6**). This result strongly suggests that the cooperative binding of cofilin is greatly influenced by cooperative conformational changes during helical twisting.

Examining the inhibitory effect of Pi on the binding and expansion of cofilin clusters

Next, we examined the nucleotide effect on the cooperative binding of cofilin to actin filaments (**Figure 8B-D, Figure supplement 6B-D, Video 6-8**). We analyzed the time-dependent changes in the peak heights and HHPs of representative actin filaments under each condition to track the growth of the cofilin clusters. We made actin filaments incubated with different nucleotides (ADP and ATP + Pi), leading to the formation of F-ADP-actin and F-ADP.Pi-actin, and subsequently mixed with cofilin. We consistently observed the preferential binding of cofilin to shortened bare HHPs toward the PE side. However, we noted that in rare instances, cofilin clusters also grew on both sides in the regular bare half helices when ATP or ADP was present. The histograms of the peak heights and HHPs followed Gaussian distributions corresponding to the cofilactin and normal bare actin regions (**Figure 8B-C, Figure supplement 6B-C**). The growth rates of the cofilin clusters within the shortened bare half helices adjacent to the cofilin clusters in both F-ADP.Pi-actin and F-ADP-actin were not significantly different, as shown in **Figure 9A-B**. Currently, we have no experimental data to discuss about whether and when bound Pi is released during this process. However, three major processes could be considered once a cofilin cluster with a critical size has formed, the allosteric transmission of the supertwisted structure of the half helices within the cofilactin region to an adjacent bare half helix on the PE side results in C-actin like protomers. First, Pi is released accompanying (preceding) supertwisting structural change in the bare region near cofilin clusters on the PE side. Because most structural studies suggest that cofilin binding promotes Pi release, leading to the formation of C-actin with ADP binding (**Figure 1A**), we anticipate that protomers within the shortened bare half helix adjacent to cofilin clusters in F-ADP.Pi-actin rapidly converts to F-ADP-actin and C-actin like structures, thereby facilitating the expansion of cofilin clusters toward the states of actin bound to ADP. Second, Pi remains bound to protomers within the supertwisted bare section. The successive interactions with cofilin molecules promote Pi release. Third, similar to process described in second one, but in this case, subsequent interactions with cofilin molecules do not induce the release of Pi. Consequently, numerous actin protomers within the newly formed cofilactin region remain bound to ADP.Pi. When we observed neighboring bare half helices on both sides of a cofilin cluster in F-ADP-actin and F-ADP-Pi-actin within one imaging field, the HHP was always shorter on one side, while the HHP was essentially normal on the other side in both F-ADP-actin and F-ADP.Pi-actin. These results suggest that bound Pi has a minor effect, hindering the propagation of the supertwisted helical structure within cofilin clusters to the neighboring bare half helices on the PE side.

To examine how bound Pi inhibits cofilin binding (Blanchoin and Pollard, 1999; Carlier et al., 1997; Carlier and Pantaloni, 1988), we assessed the torsional flexibility by measuring the

HHPs in control actin filaments incubated with ADP and ATP+Pi, or ATP + phalloidin (**Figure**
9C-D). In all cases, the histograms followed typical HHP distributions. The mean HHPs were
nearly identical. However, the variances of the HHPs in F-ADP.Pi-actin and phalloidin-stabilized
F-actin were significantly smaller than that in F-ADP-actin. Consequently, the proportion of
naturally supertwisted half helices with HHPs shorter than 30 nm was 5.8% for F-ADP-actin but
only 1.1% and 0.2% for F-ADP.Pi-actin and phalloidin-stabilized F-actin, respectively. Our
finding provides the structural support for earlier research that the nucleotide state of actin could
influence F-actin's flexibility and its ability to undergo rearrangements that mediate ABP binding
(Blanchoin and Pollard, 1999; Suarez et al., 2011; Zimmermann et al., 2015). This is in line with
measurements of micrometer-scale persistence length, indicating that F-ADP.Pi-actin is more
rigid than F-ADP-actin (Isambert et al., 1995), and with the mechanical control of twisting and
bending of actin's nucleotide-bound states (De La Cruz et al., 2010). Assuming that initial cofilin
binding occurs at sites with naturally supertwisted helical structures (Galkin et al., 2001), we
suggest that bound Pi in F-ADP.Pi-actin mainly inhibits initial cofilin binding by reducing the
fraction of supertwisted helical structures in comparison to F-ADP-actin.

Discussion
We investigated the dynamic conformational changes in half helices of F-actin and C-
actin like structure segments by observing the fluctuations in the number of actin protomers per
HHP and MAD. The results provide robust support for a prior study (Egelman et al., 1982),
which reported variable helical twisting within actin filaments. Nonetheless, the Cryo-EM and
X-ray diffraction findings depict the rigid and inflexible helical twists of actin filaments (HHP =
36 nm). Certainly, HS-AFM captures the dynamic structural changes of proteins during
functional processes in solution, offering a trade-off with lower spatial resolution but superior
temporal resolution compared to Cryo-EM (Ando, 2012; Ando et al., 2008). We leverage the
advantage of this technique to capture the torsional flexibility during helical twisting of actin
filaments and elucidate their adaptable structure as a variable helical twist, which plays crucial
roles in the binding and cluster expansion of cofilin. While it was previously believed that the
interaction between the upper and lower clefts establishes the structural basis for how nucleotide-
dependent conformation changes modulate the binding affinities for ABPs including cofilin
(Dominguez and Holmes, 2011), recent structural analyses have revealed a remarkable similarity
in the average structure of actin protomers bound with ADP.BeF$_3$, ADP.Pi, and ADP in the
presence of Mg^{2+} (Oosterheert et al., 2022; Reynolds et al., 2022) (**Figure supplement 1**),
significantly challenging the hypothesis regarding the modulation of ABP binding affinities
through nucleotide-dependent conformational changes. So, why is the flexibility in torsion
during helical twisting crucial for the preferential cooperative binding of cofilin to actin
filaments toward the PE side? Unfortunately, we currently lack the atomic structures of actin
protomers adjacent to cofilin clusters on the PE side to fully address this question.
Nonetheless, our PCA results reveal that key longitudinal interactions between the D-
loop of SD2 positioned in the pocket of the next protomer's SD1 contribute to keeping the
nucleotide binding cleft closed and the OD untwisted in the F-actin (**Figures 1, 2**). When cofilin
binds and sits on the sides of SD1 and SD2 of two adjacent actin protomers (**Figure 2B, C**), the
OD (SD1, SD2) in bound protomers becomes more twisted (a decrease in PC2), but without
affecting the closing of the nucleotide binding cleft (nearly unchanging PC1). This
transformation ultimately leads to the conformational changes from F-actin to C-actin (**Figure**
1). The transition from the C-actin to breaking the filament into monomer G-actin structure

additionally requires the opening of the nucleotide binding cleft (decreasing PC1). However, this
process is normally hindered in the presence of cofilin, as the bridging between two protomers
can compensate for all the losses in the disrupted interactions between the SD2 D-loop and SD1
pocket (**Figure 2B, C**). Strikingly, the HS-AFM results suggest that cofilin clusters can convert
bare protomers within the shortened bare half helices on the PE side into C-actin like structures.
Unless additional cofilin molecules quickly bind to the protomers with C-actin like structures on
the PE side, the disrupted interactions would readily lead to the nucleotide binding cleft
transitioning from the closed to open states, resembling C-actin to G-actin transition (**Figure 1**).
Our proposed hypothesis regarding the shift from C-actin like structure to G-actin aligns with
earlier studies suggesting that the binding sites of cofilin encompass a region that overlaps with
G-actin structure (Carlier et al., 1997; Mannherz et al., 2007; Narita, 2020; Tanaka et al., 2018).
Additionally, the protomers within the shortened bare half helix near a cofilin cluster on the PE
side of F-ADP.Pi-actin may be quickly converted to F-ADP-actin and C-actin like structure
through F-actin to C-actin transition. Together, this explains the preferential binding of cofilin to
ADP-actin on the PE side and why filament breakages are more commonly observed at or near
the boundary of cofilactin and bare actin on the PE side, rather than where actin protomers are
fully bound with cofilin (Bibeau et al., 2021; Ngo et al., 2015; Suarez et al., 2011). However, our
finding regarding the structure partially contradicts the existing cryo-EM structure, which
suggested that conformational changes induced by cofilin clusters, shifting from F-actin to C-
actin, are restricted to the protomers directly in contact with bound cofilin. According to this
model, cofilin tunes adjacent protomers, resembling F-ADP-actin structure on the PE side, to
expand the clusters (Huehn et al., 2020; Suarez et al., 2011). Nonetheless, their hypothesis is
insufficient to explain the PCA results supporting the structural evidence obtained from our
AFM measurements that a cofilin cluster not only has the capacity to convert F-actin to C-actin
but also allosterically expands the C-actin structure from the bound to unbound region on the PE
side. Thus, additional cofilin molecules preferentially bind to actin protomers resembling C-actin
within the shortened bare zone than F-ADP-actin structure near a cofilin cluster on the PE side
(**Figures 1, 5-9**). Even though HS-AFM did not allow for the direct observation of the different
nucleotide states bound to actin, these unidirectional propagation of the supertwisted structure in
cofilin clusters to the neighboring bare zone on the PE side and preferential and cooperative
bindings of cofilin to those bare sections were consistently and unequivocally observed in the
presence of ADP and ADP.Pi (Ngo et al., 2015). Further studies are needed to identify the source
of this discrepancy between HS-AFM and cryo-EM observations.
Our aim is to understand the underlying mechanism driving the preferential and
cooperative attachment of cofilin. This mechanism is largely influenced by the preferred helical
twisting structure within the actin filament, as opposed to an alternate theory that suggests cofilin
binding depends on the nucleotide state of actin (**Figure 10**). While both hypotheses might
explain cofilin binding along the actin filament, they propose different mechanisms underlying
cofilin's cooperative behavior and binding preference.
Our hypothesis (**Figure 10E-H**) aligns with the results in the cofilactin region, where
actin protomers are fully decorated by cofilin (**Figure 10A, 10C, 10D**). However, it is partially
incongruent with the structures of actin protomers near cofilin clusters on the PE side, as
depicted in **Figure 10D**. Additionally, we propose two potential hypotheses that require further
verification in the near future to explain the preferential cooperative bindings and cluster
expansion of cofilin to undecorated C-actin like structure within the shortened bare half helix on
the PE side (**Figure 10E-F**): (i) Because of the disrupted connections between the BE-

protomer's SD2 D-loop and PE-protomer's SD1 pocket in the area resembling the C-actin
structure, where cofilins are not bound, the SD2 could change between untwisted and twisted
states (F-actin to C-actin transition) and the nucleotide-binding cleft could shift between the
closed and open states (C-actin to G-actin transition). These transitions might prompt the rapid
transformation of protomers within the shortened bare half helix adjacent to the cofilin clusters
on the PE side of F-ADP.Pi-actin to F-ADP-actin and C-actin like structure. This phenomenon
also explains the binding preference of cofilin for ADP-actin on the PE side; (ii) In the canonical
actin protofilament, the SD2 D-loop of BE-protomer is inserted between the tips of SD1 and SD3
of PE-protomer (**Figure 2A**). However, the SD2 D-loop of BE protomer does not directly
interact with the tips of SD1 and SD3. Instead, it is deeply buried within the core of the cofilactin
protofilament, representing the C-actin structure (**Figure 2B-C**). Conversely, in the region
representing the C-actin like structure, where cofilins are unbound, the SD2 D-loops might
exhibit various conformations, potentially allowing flexible interaction with the tips of SD1 and
SD3 including that of the neighboring protomers on the PE side. This interaction could lead to an
extended MAD in C-actin like structure area with optimal conformational changes in the targets-
binding cleft (hydrophobic - lower cleft) between SD1 and SD3 of the protomers on the PE side,
thereby facilitating the preferential binding and expansion of cofilin clusters (**Figure 10E-F**).
Furthermore, we observed a significant correlation between the nucleotide-bound states
of actin and the proportion of naturally supertwisted half helices in F-ADP-actin and F-ADP.Pi-
actin (**Figure 9**). We suggest that the bound Pi reduces the fraction of naturally supertwisted half
helices and hinders the initial binding of cofilin and/or initial growth to a critical size of cluster
set at 2 to 4 molecules (**Figure 5-6**) that can induce allosteric conformational changes to the
neighboring bare zone on the PE side for cooperative bindings of cofilin (**Figures 9C-D, 10H**).
This discovery aligns with previous research, which showed that achieving the most effective
cooperative binding of cofilin also requires a critical size for a cofilin cluster set at 2 molecules
(Huehn et al., 2020). However, Pi only minimally impedes the transition of the supertwisted
helical structure within the cofilin cluster to the bare actin zone in F-ADP.Pi-actin compared to
F-ADP-actin. This transition is pivotal in driving the preferential cooperative binding and
expansion of the cofilin cluster (**Figures 7, 8**). In terms of structure, the PCA results also
strongly agree with previous research indicating that ADP.BeF$_3$, ADP.Pi, and ADP F-actin
structures (**Figure supplement 1**) display a striking resemblance (Oosterheert et al., 2022;
Reynolds et al., 2022). Additionally, the reduction in torsional flexibility during helical twisting
resulted in a significant hindrance to the cooperative binding and expansion of cofilin clusters
(**Figure 8A**). Therefore, we suggest that the reduction in torsional flexibility associated with
variable helical twists, rather than the large conformational change in F-actin's protomers linked
to ATP hydrolysis, accounts for the inhibition of cofilin binding by Pi (**Figure 10H**).
Nonetheless, it remains plausible that the structural flexibility exhibited by ADP-bound actin
protomers could result in subtle variations in the conformations of the DNase binding loop (D-
loop) G46-M47-G48-N49, as suggested in (Chou and Pollard, 2019). We suggest that the
absence of bound Pi possibly increases the torsional flexibilities during helical twisting of ADP-
bound actin filaments in contrast to their ADP.Pi-bound counterparts.
Actin structure-dependent preferential binding of actin-binding proteins (ABPs) have
been investigated at the molecular and cellular levels (Hayakawa et al., 2022; Jegou and Romet-
Lemonne, 2019; Jégou and Romet-Lemonne, 2016; Ngo et al., 2016; Tokuraku et al., 2020,
2009). Notably, Harris and colleagues reported biased localization of tandem calponin homology
domains (CH1-CH2) that can be mutated to selectively bind different actin conformations at the

leading or trailing edge of motile cells (Harris et al., 2020). The untwisting of the long-pitch
helix of the actin filament induced by diaphanous-related formin mDia1 inhibits cofilin binding
(Mizuno et al., 2018). Stretching actin filaments within cells enhances their binding affinities for
the myosin II motor domain (Uyeda et al., 2011). The importance of cooperative and allosteric
conformational changes in actin filaments warrants further investigation of the actin structure-
dependent preferential cooperative binding and function of other ABPs to clarify the role of
structural polymorphisms and physiological implications of actin filaments in working with
different ABPs in cells.
558

Methods
Proteins
Rabbit skeletal actin and human cofilin 1, without a His-tag, were prepared following the
same protocol as described in our previous report (Ngo et al., 2015). The purified proteins were
rapidly frozen in liquid nitrogen and stored at -80°C until use. Prior to experimentation, the
proteins were thawed on ice and subjected to ultracentrifugation (typically at 80,000 rpm for 10
minutes at 4°C) to eliminate any undesired proteins present in the pellet. The remaining high-
quality proteins in the supernatant were collected, kept on ice, and utilized for HS-AFM
experiments.
568

Principal component analysis
Principal component analysis (PCA) is a commonly used statistical method for detecting
correlations in large datasets by reducing the dimensionality of the data. In the analysis of PDB
structures, PCA can reveal critical information regarding collective changes in the original
conformations. The process involves aligning the original data in three-dimensional space to
remove trivial movements, calculating the displacement covariance matrix between common
atoms, and performing eigendecomposition. This results in a small number of new variables
called principal components (PCs), which capture most of the original data's information.
Clustering is a byproduct of PCA, as the original dataset can be projected onto the identified
collective components and naturally grouped in the new PC dimensional space.
To ensure consistent and accurate PCA results, we conducted several preprocessing steps.
After cleaning the selected PDBs, we specifically extracted the alpha carbons of all reported
protein chains for further analysis. The sequences of these chains were aligned using the AlignIO
module from the Biopython package (Cock et al., 2009), and we identified the residues that were
commonly present across the PDBs. Specifically, we found that residues 8 to 41, 54 to 63, 66 to
74, 76 to 234, and 236 to 373 in the ACTs_RABIT gene (P68135) were common among the 46
chosen PDBs. Importantly, these residue numbers differ from the standard residue IDs used in
PDBs because the initial two residues present in genes are typically cleaved in the final actin
chains.
We then performed structural alignment only for residues in the IDs, following the
definition reported in (Kabsch et al., 1990). After aligning the structures, we used the entire set
of commonly solved residues to establish the covariance matrix, which was subjected to
eigendecomposition to obtain different eigenvectors (PCs). As shown in **Figure 1 and Figure**
supplement 1, we utilized the two eigenvectors with the largest values, denoted as PC1 and PC2.
In this study, the significance of PCA lies in its capacity to unveil the conformational transition
between G-actin and C-actin, and vice versa, through PC1, while PC2 delineates the pathway
between F-actin and C-actin, and vice versa.

Analyses of pseudo AFM images of F-actin and C-actin structures constructed from existing PDB structures

To illustrate how cofilin clusters impact the helical twisting of long actin filaments, leading to the shortening of HHPs and less protomers per HHP, resembling the C-actin structure as measured by AFM, we built long actin filaments at different states along the F-actin to C-actin transitions by using 8 PDB chains (**Table supplement 2**). In **Figure supplement 2**, we show pseudo-AFM images featuring long filaments representing F-actin (constructed from 6VAU), C-actin (with cofilins excluded) and cofilactin (constructed from 6VAO) structures.

To construct the long filament featuring F-actin, we performed a series of steps. First, we repeatedly rotated and translated two actin chains from the selected PDBs to generate the coordinates of 62 consecutive protomers. Since not all of the selected PDBs had reported BIOMT matrices, we used transformation matrices that minimized the root mean square deviation (RMSD) between the two chains. To construct long filaments featuring C-actin and cofilactin, a similar process was followed, but it involved two chains of actin protomers and two chains of cofilins. The lists of chains and matrices utilized to build the long filaments are presented in **Table supplement 2**.

The generated long filaments were then aligned along the x-axis and pseudo-imaged using Afmize software (https://github.com/ToruNiina/afmize) (Niina et al., 2021, 2020). A probe radius of 2.0 nm with an angle of 5.0° was set, and the image resolution was configured to be 1.0 nm in the xy plane and 0.64 Å on the z-axis, with a noise level of 0.1 nm. To analyze the structures, we calculated longitudinal section profiles of the HHPs by determining the average height within ± 3 nm around the centerline (red and blue lines in **Figure supplement 2A**). The offset values of the height profile were set to zero by referring to virtual mica surface, where filament was placed in the pseudo-AFM images.

In our AFM analyses, we determined the HHPs by measuring the distances between major peaks observed in longitudinal section profiles along the filament axis (red markers in **Figure supplement 2A-B, 2D-E, and 2G-H**). These major peaks likely represent two distinct actin protofilaments that are successively probed by the AFM tips (red markers in **Figure supplement 2C, 2F, and 2I**). Thus, the number of actin protomers within each HHP is odd, leading to a noninteger number of actin pairs. For example, in canonical actin filaments, each HHP is typically composed of 13 actin protomers (equivalent to 6.5 protomer pairs). To account for this, we determined the number of protomer pairs within each HHP by counting the major and minor peaks (noted by the red and green markers within each HHP subtracted by 0.5) in the same longitudinal-section profiles. This adjustment was applied for both the pseudo and real AFM images of the filaments representing F-actin and C-actin. Our pseudo-AFM image displayed the challenge of accurately detecting actin pairs within filaments representing the cofilactin structure (**Figure supplement 2G-I**) due to the presence of cofilin decorations and the tip-sample dilation effect (Ando, 2022).

High-speed atomic force microscopy

We used a laboratory-built high-speed atomic microscope (HS-AFM), as described previously (Ando et al., 2013). HS-AFM imaging in the tapping mode was carried out in solution with small cantilevers (BL-AC10DS-A2, Olympus), and the spring constant, resonant frequency in water, and quality factor in water were ~0.1 N/m, ~500 kHz, and ~1.5, respectively. An additional tip was grown on the original cantilever tip in gas with sublimable ferrocene powder

by electron beam deposition (EBD) using scanning electron microscopy (ZEISS Supra 40
VP/Gemini column, Zeiss, Jena, Germany). Typically, the EBD tip was grown under vacuum (1-
5×10^{-6} Torr), with an aperture size of 10 µm and an electron beam voltage of 20 kV for 30 s.
The EBD ferrocene tip was sharpened using a radio frequency plasma etcher (Tergeo Plasma
Cleaner, Pie Scientific, Union City, CA) under an argon gas atmosphere (typically at 180 mTorr
and 20 W for 90 s). For HS-AFM imaging, the free-oscillation peak-to-peak amplitude of the
cantilever (A_0) was set at ~1.6-1.8 nm, and the feedback amplitude set-point was set at ~0.9 A_0.
HS-AFM imaging was performed as described in detail elsewhere (Ando et al., 2013; Kodera et
al., 2021; Ngo et al., 2015), except for the use of a recently developed only tracing imaging
(OTI) mode (Fukuda and Ando, 2021).

HS-AFM imaging
The lipid compositions used in this study predominantly consisted of neutral 1,2-
dipalmitoyl-*sn*-glycero-3-phosphocholine (DPPC, Avanti) and positively charged 1,2-
dipalmitoyl-3-trimethylammonium-propane (DPTAP, Avanti). Liposomes were prepared using
DPPC/DPTAP (90/10, wt%), DPPC/DPTAP (75/25, wt%), and DPPC/DPTAP (50/50, wt%)
ratios, following our established sample preparation protocol (Ngo et al., 2015). These
liposomes, along with mica-supported lipid bilayers, were created accordingly.
Actin filaments were generated in F1 buffer containing 40 mM KCl, 20 mM PIPES-KOH
(pH 6.8), 1 mM MgCl$_2$, 0.5 mM EGTA, 0.5 mM DTT, and 1 mM ATP, following the procedure
described in our previous work (Ngo et al., 2015). To obtain actin filaments bound with different
nucleotides (ATP, ADP, ADP.Pi) and investigate the cooperative binding of cofilin to these
filaments, a similar protocol was employed (Ngo et al., 2015). Briefly, actin filaments (5–10 µM)
were made in F1 buffer and gently immobilized on a positively charged lipid membrane
composed of DPPC/DPTAP (90/10 wt%). The actin filaments on the lipid surface were washed
with 20 µl of imaging F1 buffer additionally including either 0.5 mM ATP, 1 mM ADP/5 U/ml
hexokinase/10 mM glucose, or 1 mM ATP + 10 mM Pi (K$_2$HPO$_4$/KH$_2$PO$_4$ (pH 6.8) buffer),
resulting in F-ATP-actin, F-ADP-actin, or F-ADP.Pi-actin, respectively. Subsequently, cofilin,
prepared in the imaging F1 buffer with the desired concentration and nucleotides, was introduced
into the imaging chamber to observe the cooperative binding of cofilin along the actin filaments.
To compare the helical twisting structures, the number of protomers per HHP, and MADs
in the normal F-actin structure with that in bare actin segments on both sides of the cofilin
clusters in the hybrid cofilactin/bare actin filaments at high spatiotemporal resolution in the same
imaging field, we immobilized these filaments sequentially on the same lipid membrane
(referred to **Figures supplement 4-5**). First, actin filaments were prepared by incubating G-actin
in F1 buffer at a concentration of 20 µM for approximately 1 hour at 22°C, resulting in F-actin.
Second, cofilactin and hybrid cofilactin/bare actin filaments were formed by mixing F-actin (10
µM) with cofilin (1 µM) in a small tube in F1 buffer. The final molar ratio was 10/1 (actin to
cofilin). The mixture was incubated for 10 minutes at 22°C to allow the binding, cluster
expansion and actin filament severing to reach equilibrium. Third, a 2 µl solution containing
cofilactin and hybrid cofilactin/bare actin filaments was introduced onto a lipid membrane
composed of DPPC/DPTAP (50/50 wt%) and incubated for 10 minutes at 22°C to immobilize
the filaments. Fourth, the surface was washed with 20 µl of F1 buffer, and then a 2 µl solution
containing normal actin filaments (10 µM) was introduced onto the same lipid membrane. This
sample was also incubated for 10 minutes at 22°C. Fifth, the surface was washed again with 20
µl of imaging F1 buffer prior to HS-AFM imaging.

Analysis of the growth rate of the cofilin cluster within the shortened bare half helices adjacent to the cofilin cluster in F-ADP.Pi-actin and F-ADP-actin

We measured the growth rates of the cofilin clusters within shortened bare half helices adjacent to the cofilin clusters on the PE side in F-ADP.Pi-actin and F-ADP-actin. As noted in the section of Results, when we were able to observe bare sections on both sides of a cofilin cluster, one of them had shorter HHP while the other was close to normal HHP, and based on our previous report (Ngo et al., 2015), we judged the bare section with shorter HHP is on the PE side of an actin filament. It is important to clarify that the collected data did not include the growth rates of cofilin within the normal half helices on the PE or BE sides, in order to exclude the random binding events of cofilin to actin filaments under various nucleotide conditions. By employing HS-AFM, we were able to monitor changes in the length of a half helix adjacent to a cofilin cluster and the peak height at individual crossover points. The lag time measurements were performed at the precise moment when the bare half helix adjacent to the cofilin cluster on the PE side began to shorten and continued until a newly saturated cluster emerged within this neighboring bare half helix. This determination was made by analyzing the increase in the peak heights (P1) before and after forming a newly matured cofilin cluster adjacent to a preformed cofilin cluster (P2) (as depicted in **Figure 9A**). The data presented in **Figures 7, 8C** were also included in this analysis. Each fully decorated-supertwisted half helix consists of 11 cofilin molecules (referred to **Figure supplement 1I**). To calculate the mean growth rate of the cofilin cluster within a shortened bare half helix adjacent to a cofilin cluster, we can divide 11 cofilin molecules by the lag time in seconds.

We assessed torsional flexibility of actin filaments by measuring the HHPs in control actin filaments incubated with 1 mM ADP and 1mM ATP + 10 mM Pi, or 1 mM ATP + phalloidin saturated to actin, resulting in F-ADP-actin, F-ADP.Pi-actin, and phalloidin-stabilized F-actin, respectively, using our previous method (Ngo et al., 2015). Because, we did not examine the nucleotide-bound actin states when actin filaments were incubated with 1 mM ATP and phalloidin excess, we simply labelled them as phalloidin-stabilized F-actin and used as the control. The fraction of naturally supertwisted half-helices with HHPs less than 30 nm was determined by computing the areas under the curves derived from single Gaussian fitting curves (referred to **Figure 9C-D**).

HS-AFM data analysis and processing

The AFM data were analyzed using custom software, including Kodec, UMEX Viewer, and Umex Viewer for Drift Analysis, as described in previous studies (Hayakawa et al., 2022; Ngo et al., 2015). To measure the half helical pitch (HHP) and the number of actin protomer pairs per HHP, we utilized the custom software Umex Viewer for Drift Analysis, as described in (Hayakawa et al., 2022). Additionally, AFM images and videos were processed using Fiji-ImageJ (NIH, USA) and After Effects (Adobe, USA).

To quantitatively analyze the HHPs and count the number of protomer pairs per HHP, we employed Umex Viewer for Drift Analysis, which enabled the semiautomatic determination and measurement of the distance between the highest "cross-over" points between two long-pitch strands. This involved creating a topographical line profile along the actin filaments, similar to previous studies conducted by (Amyot et al., 2022; Hayakawa et al., 2022). Specifically, one HHP was measured between the two highest cross-over points, while one helical pitch (HP) was measured between the three highest cross-over points. In brief, we drew a longitudinal section

profile line along the actin filament, covering a length of 1-4 consecutive HHPs. The offset
values were set to zero with reference to the height of the lowest position along the longitudinal
section lines. Before analysis, we corrected the nonlinearity of the XY piezos by applying
nonlinear image scaling and reduced image noise using a Gaussian smoothing filter with a
standard deviation of 0.76 nm. The profile was obtained by averaging the signal within a 5-8 nm
bandpass filter along the filament. To minimize the influence of noise, we set minimum
threshold pitch values of 3 nm for the actin protomers and 20 nm for the HHPs. Typically, we
selected 30-61 actin filaments and measured the HHPs for approximately 20 consecutive time
frames. To minimize measurement errors in the HHP and the number of protomer pairs per HHP,
we specifically chose straight actin segments with clearly discernible protomers and drew line
profiles (with an average width of 3%) to measure 1-4 consecutive HHPs. This method was
consistently applied to analyze normal bare actin, cofilactin, and hybrid cofilactin/bare actin
filaments. We distinguished normal F-actin and cofilactin structures based on differences in the
peak height and HHP, as described in our previous study (Ngo et al., 2015).
To calculate the number of protomer pairs per HHP and the mean axial distance (MAD)
between two adjacent protomers along the same long-pitch strand, we semiautomatically
detected and counted the number of protomer pairs per HHP, as mentioned in other parts and
shown in **Figure supplement 2**. The MADs within an HHP were determined by dividing the
HHP by the actual counted number of protomer pairs per HHP.
To enhance the characteristics of actin protomers in the different types of filaments (bare
F-actin, cofilactin and C-actin like structure in the hybrid cofilactin/bare actin segments), we
converted the raw AFM images into Laplacian-filtered images. This was achieved by applying a
high-pass filter known as the Laplacian of Gaussian (LoG) filter, which can be found at the
following link:
(https://homepages.inf.ed.ac.uk/rbf/HIPR2/log.htm). The Laplacian filter used in this process is a
type of high-pass filter, characterized by the kernel $\begin{bmatrix} -1 & -1 & -1 \\ -1 & 8 & -1 \\ -1 & -1 & -1 \end{bmatrix}$. Prior to the application of
the Laplacian filter, we employed a Gaussian filter with a typical sigma value of 1.27 pixel/nm.
This preliminary step helped reduce sensitivity to noise and enhanced the resulting Laplacian-
filtered images, resulting in improved visualization of the actin protomers.
763
Temporal autocorrelation analysis
The temporal autocorrelation analysis method can be found at the following link:
https://real-statistics.com/time-series-analysis/stochastic-processes/autocorrelation-function/.
In summary, the analysis involves the following steps: (**i**) Data processing and collection:
We directly measured and collected time series data for the axial distance (AD) between two
adjacent actin protomers along the same long-pitch strand in different actin filaments over time,
as shown in **Figure 3D**, using a custom semiautomatic analysis software called Umex View for
Drift Analysis. First, we converted raw AFM images containing actin filaments into Laplacian-
filtered images by applying a Laplacian of Gaussian (LoG) filter (typically with sigma ~ 1.27
pixel/nm) to eliminate low-frequency and high-frequency noise signals. Second, we selected a
short segment of the actin filament (~4 HHPs) located at the center of the AFM image and used a
segmented line running along the axis of the actin filament to measure the distance between actin
protomer pairs, as depicted in **Figure 3D**. To facilitate the semiautomatic analysis and determine
the peak-to-peak distance between individual ADs, we applied a bandpass filter (5-8 nm) to
remove noise, setting the low-pass frequency (LPF) cutoff at 3 nm and the high-pass frequency

(HPF) cutoff at 6 nm for all analyses. Typically, the variation in the AD within each actin
filament over time was assessed using a series of AFM images (including 258 images captured at
a rate of 2 frames per second (fps)). (**ii**) Calculation of the autocorrelation coefficients: Each
autocovariance value was divided by the autocovariance at lag 0 to obtain the autocorrelation
coefficients. These coefficients represent the strength and direction of the linear relationship
between observations at different time lags (**Equation 1**). (**iii**) Plotting the autocorrelation
function (ACF): The autocorrelation coefficients were visualized by plotting the autocorrelation
function (ACF). The ACF plot displays the autocorrelation coefficients on the y-axis and the
time lags on the x-axis, as depicted in **Figure 3E**. (**iv**) Estimation of the time constants: The ACF
plot was fitted with a second-order exponential decay function to extract the time constants (t_1,
t_2), as depicted in **Figure 3E-F**.
The *ACF* (k) of a time series of values (y_i) at lag k is defined as
$ACF(k) = [\frac{1}{n}\sum_{i=1}^{n-k}(y_i - \bar{y})(y_{i+k} - \bar{y})]/[\sum_{i=1}^{n}(y_i - \bar{y})^2]$ (**Equation 1**)
In our analysis, a time series of values (y_i) of the axial distance between two adjacent
actin pairs at lag k ($k \geq 0$) was measured for each actin filament in 258 consecutive AFM images
($n = 258$) collected at a frame rate of 2 fps. The mean value, $\bar{y}$, was calculated.
795
Spatial autocorrelation analysis
To evaluate the degree of spatial clustering in the actin protomer structural data, we
conducted spatial autocorrelation analysis, as depicted in **Figure supplement 3**. Initially, we
generated a line profile by utilizing cubic spline interpolation along the ridgeline of an actin
filament (**Figure supplement 3a**). The actin filament includes a large helical twist and protomer
structures with periodicities of 36 nm and 6–7 nm, respectively. The helical twist exhibits a
typical height of 3 nm (**Figure supplement 3b**), which is three times larger than the 0.5–1.5 nm
height of the protomer (**Figure supplement 3c**). To explore the spatial self-correlation among
protomers, we applied a fast Fourier transform (FFT) and utilized a bandpass filter (BPF). The
filter isolated heights within the 5 to 8 nm range in the Fourier domain, corresponding to the
actin protomer height along the filament (depicted in **Figure supplement 3c**). Then, we derived
the autocorrelation function (ACF) of the protomer heights by obtaining the complex conjugate
(magnitude) and then performed an inverse FFT. Additionally, we applied the Hilbert transform
in the Fourier domain by doubling the positive frequency data, zeroing the negative frequency
data, and applying an inverse FFT. This process produced an envelope curve. As a result, the
envelop curves of the height profiles with and without the BPF exhibit alternating positive and
negative values at half helical pitch and protomer periodicities, respectively (**Figure supplement
5d-e**). In contrast, the envelope curve consistently shows positive values; thus, this curve is
suitable for evaluating spatial autocorrelations. In the main text, we compared the envelope
curves representing the spatial ACF results, measured at multiple time points and different
conditions.
817

References
Amyot R, Marchesi A, Franz CM, Casuso I, Flechsig H. 2022. Simulation atomic force
microscopy for atomic reconstruction of biomolecular structures from resolution-limited
experimental images. *PLoS Comput Biol* **18**:1–15. doi:10.1371/journal.pcbi.1009970
Ando T. 2022. High-Speed Atomic Force Microscopy in Biology: Directly Watching Dynamics
of Biomolecules in Action. Springer Berlin, Heidelberg. doi:10.1007/978-3-662-64785-1
Ando T. 2012. High-speed atomic force microscopy coming of age. *Nanotechnology* **23**:1–27.

825   doi:10.1088/0957-4484/23/6/062001

826   Ando T, Uchihashi T, Kodera N. 2013. High-speed AFM and applications to biomolecular
827   systems. *Annu Rev Biophys* **42**:393–414. doi:10.1146/annurev-biophys-083012-130324

828   Ando T, Uchihashi T, Kodera N, Yamamoto D, Miyagi A, Taniguchi M, Yamashita H. 2008.
829   High-speed AFM and nano-visualization of biomolecular processes. *Pflugers Arch Eur J*
830   *Physiol* **456**:211–225. doi:10.1007/s00424-007-0406-0

831   Andrianantoandro E, Pollard TD. 2006. Mechanism of Actin Filament Turnover by Severing and
832   Nucleation at Different Concentrations of ADF/Cofilin. *Mol Cell* **24**:13–23.
833   doi:10.1016/j.molcel.2006.08.006

834   Bamburg JR, Bernstein BW. 2010. Roles of ADF/cofilin in actin polymerization and beyond.
835   *F1000 Biol Rep* **2**:1–7. doi:10.3410/B2-62

836   Bibeau JP, Gray S, De La Cruz EM. 2021. Clusters of a Few Bound Cofilins Sever Actin
837   Filaments. *J Mol Biol* **433**:166833. doi:10.1016/j.jmb.2021.166833

838   Blanchoin L, Pollard TD. 1999. Mechanism of interaction of Acanthamoeba actophorin
839   (ADF/Cofilin) with actin filaments. *J Biol Chem* **274**:15538–15546.
840   doi:10.1074/jbc.274.22.15538

841   Bobkov A a, Muhlrad A, Pavlov D a, Kokabi K, Yilmaz A, Reisler E. 2006. Cooperative effects
842   of cofilin (ADF) on actin structure suggest allosteric mechanism of cofilin function. *J Mol*
843   *Biol* **356**:325–34. doi:10.1016/j.jmb.2005.11.072

844   Carlier MF, Laurent V, Santolini J, Melki R, Didry D, Xia GX, Hong Y, Chua NH, Pantaloni D.
845   1997. Actin depolymerizing factor (ADF/cofilin) enhances the rate of filament turnover:
846   Implication in actin-based motility. *J Cell Biol* **136**:1307–1322. doi:10.1083/jcb.136.6.1307

847   Carlier MF, Pantaloni D. 1988. Binding of phosphate to F-ADP-actin and role of F-ADP-P(i)-
848   actin in ATP-actin polymerization. *J Biol Chem* **263**:817–825. doi:10.1016/s0021-
849   9258(19)35428-6

850   Chou SZ, Pollard TD. 2019. Mechanism of actin polymerization revealed by cryo-EM structures
851   of actin filaments with three different bound nucleotides. *Proc Natl Acad Sci U S A*
852   **116**:4265–4274. doi:10.1073/pnas.1807028115

853   Cock PJA, Antao T, Chang JT, Chapman BA, Cox CJ, Dalke A, Friedberg I, Hamelryck T,
854   Kauff F, Wilczynski B, De Hoon MJL. 2009. Biopython: Freely available Python tools for
855   computational molecular biology and bioinformatics. *Bioinformatics* **25**:1422–1423.
856   doi:10.1093/bioinformatics/btp163

857   De La Cruz EM, Roland J, McCullough BR, Blanchoin L, Martiel J-L. 2010. Origin of twist-
858   bend coupling in actin filaments. *Biophys J* **99**:1852–60. doi:10.1016/j.bpj.2010.07.009

859   Dominguez R. 2004. Actin-binding proteins - A unifying hypothesis. *Trends Biochem Sci*
860   **29**:572–578. doi:10.1016/j.tibs.2004.09.004

861   Dominguez R, Holmes KC. 2011. Actin Structure and Function. *Annu Rev Biophys* **40**:1–20.
862   doi:10.1146/annurev-biophys-042910-155359.

863   Egelman EH. 1994. Actin Filament Structure: The ghost of ribbons past. *Curr Biol* **4**:79–81.
864   doi:10.1016/S0960-9822(00)00020-8

865   Egelman EH, Francis N, DeRosier DJ. 1982. F-actin is a helix with a random variable twist.
866   *Nature* **298**:131–135. doi:10.1038/298131a0

867   Fujii T, Iwane AH, Yanagida T, Namba K. 2010. Direct visualization of secondary structures of
868   F-actin by electron cryomicroscopy. *Nature* **467**:724–728. doi:10.1038/nature09372

869   Fukuda S, Ando T. 2021. Faster high-speed atomic force microscopy for imaging of
870   biomolecular processes. *Rev Sci Instrum* **92**. doi:10.1063/5.0032948

Galkin VE, Orlova a, Lukoyanova N, Wriggers W, Egelman EH. 2001. Actin depolymerizing
factor stabilizes an existing state of F-actin and can change the tilt of F-actin subunits. *J
Cell Biol* **153**:75–86.
Galkin VE, Orlova A, Kudryashov DS, Solodukhin A, Reisler E, Schröder GF, Egelman EH.
2011. Remodeling of actin filaments by ADF/cofilin proteins. *Proc Natl Acad Sci U S A*
108:20568–20572. doi:10.1073/pnas.1110109108
Galkin VE, Orlova A, Vos MR, Schröder GF, Egelman EH. 2015. Near-Atomic Resolution for
One State of F-Actin. *Structure* **23**:173–182. doi:10.1016/j.str.2014.11.006
Grintsevich EE, Yesilyurt HG, Rich SK, Hung R-J, Terman JR, Reisler E. 2016. F-actin
dismantling through a redox-driven synergy between Mical and cofilin. *Nat Cell Biol*
18:876–85. doi:10.1038/ncb3390
Harris AR, Jreij P, Belardi B, Joffe AM, Bausch AR, Fletcher DA. 2020. Biased localization of
actin binding proteins by actin filament conformation. *Nat Commun* **11**:1–13.
doi:10.1038/s41467-020-19768-9
Hayakawa K, Sakakibara S, Sokabe M, Tatsumi H. 2014. Single-molecule imaging and kinetic
analysis of cooperative cofilin-actin filament interactions. *Proc Natl Acad Sci U S A*
111:9810–9815. doi:10.1073/pnas.1321451111
Hayakawa Y, Takaine M, Ngo KX, Imai T, Yamada M, Behjat AB, Umeda K, Hirose K,
Yurtsever A, Kodera N, Tokuraku K, Numata O, Fukuma T, Ando T, Nakano K, Uyeda
TQP. 2022. Actin binding domain of Rng2 sparsely bound on F-actin strongly inhibits actin
movement on myosin II. *Life Sci Alliance* **6**:e202201469. doi:10.26508/lsa.202201469
Holmes KC, Popp D, Gebhard W, Kabsch W. 1990. Atomic model of the actin filament. *Nature*
347:44–49. doi:10.1038/347044a0
Huehn A, Cao W, Austin Elam W, Liu X, De La Cruz EM, Sindelar C V. 2018. The actin
filament twist changes abruptly at boundaries between bare and cofilin-decorated segments.
J Biol Chem **293**:5377–5383. doi:10.1074/jbc.AC118.001843
Huehn AR, Bibeau JP, Schramm AC, Cao W, De La Cruz EM, Sindelar C V. 2020. Structures of
cofilin-induced structural changes reveal local and asymmetric perturbations of actin
filaments. *Proc Natl Acad Sci U S A* **117**:1478–1484. doi:10.1073/pnas.1915987117
Isambert H, Venier P, Maggs AC, Fattoum A, Kassab R, Pantaloni D, Carlier MF. 1995.
Flexibility of actin filaments derived from thermal fluctuations. Effect of bound nucleotide,
phalloidin, and muscle regulatory proteins. *J Biol Chem* **270**:11437–11444.
doi:10.1074/jbc.270.19.11437
Jegou A, Romet-Lemonne G. 2019. The many implications of actin filament helicity. *Semin Cell
Dev Biol* **102**:65–72. doi:10.1016/j.semcdb.2019.10.018
Jégou A, Romet-Lemonne G. 2016. Single Filaments to Reveal the Multiple Flavors of Actin.
Biophys J **110**:2138–2146. doi:10.1016/j.bpj.2016.04.025
Kabsch W, Mannherz HG, Suck D, Pai EF, Holmes KC. 1990. Atomic structure of the actin:
DNase I complex. *Nature* **347**:37–44. doi:10.1038/347037a0
Kasai M, Asakura S, Oosawa F. 1962. The cooperative nature of G-F transformation of actin.
Biochim Biophys Acta **57**:22–31.
Kodera N, Noshiro D, Dora SK, Mori T, Habchi J, Blocquel D, Gruet A, Dosnon M, Salladini E,
Bignon C, Fujioka Y, Oda T, Noda NN, Sato M, Lotti M, Mizuguchi M, Longhi S, Ando T.
2021. Structural and dynamics analysis of intrinsically disordered proteins by high-speed
atomic force microscopy. *Nat Nanotechnol* **16**:181–189. doi:10.1038/s41565-020-00798-9
Lappalainen P, Drubin DG. 1997. Cofilin promotes rapid actin filament turnover in vivo. *Nature*

388:78–82. doi:10.1038/40418

Lorenz M, Popp D, Holmes KC. 1993. Refinement of the F-actin model against x-ray fiber
diffraction data by the use of a directed mutation algorithm. *J Mol Biol* **234**:826–836.

Mannherz HG, Ballweber E, Galla M, Villard S, Granier C, Steegborn C, Schmidtmann A,
Jaquet K, Pope B, Weeds AG. 2007. Mapping the ADF/Cofilin Binding Site on Monomeric
Actin by Competitive Cross-linking and Peptide Array: Evidence for a Second Binding Site
on Monomeric Actin. *J Mol Biol* **366**:745–755. doi:10.1016/j.jmb.2006.11.100

McCullough BR, Blanchoin L, Martiel JL, De La Cruz EM. 2008. Cofilin increases the bending
flexibility of actin filaments: implications for severing and cell mechanics. *J Mol Biol*
381:550–558. doi:10.1016/j.jmb.2008.05.055

McGough A, Pope B, Chiu W, Weeds A. 1997. Cofilin changes the twist of F-actin: implications
for actin filament dynamics and cellular function. *J Cell Biol* **138**:771–781.

Mizuno H, Tanaka K, Yamashiro S, Narita A, Watanabe N. 2018. Helical rotation of the
diaphanous-related formin mDia1 generates actin filaments resistant to cofilin. *Proc Natl
Acad Sci U S A* **115**:E5000–E5007. doi:10.1073/pnas.1803415115

Narita A. 2020. ADF/cofilin regulation from a structural viewpoint. *J Muscle Res Cell Motil*
41:141–151. doi:10.1007/s10974-019-09546-6

Ngo KX, Kodera N, Katayama E, Ando T, Uyeda TQ. 2015. Cofilin-induced unidirectional
cooperative conformational changes in actin filaments revealed by high-speed atomic force
microscopy. *Elife* **4**:1–22. doi:10.7554/eLife.04806

Ngo KX, Umeki N, Kijima ST, Kodera N, Ueno H, Furutani-Umezu N, Nakajima J, Noguchi
TQP, Nagasaki A, Tokuraku K, Uyeda TQP. 2016. Allosteric regulation by cooperative
conformational changes of actin filaments drives mutually exclusive binding with cofilin
and myosin. *Sci Rep* **6**:1–11. doi:10.1038/srep35449

Niina T, Fuchigami S, Takada S. 2020. Flexible Fitting of Biomolecular Structures to Atomic
Force Microscopy Images via Biased Molecular Simulations. *J Chem Theory Comput*
16:1349–1358. doi:10.1021/acs.jctc.9b00991

Niina T, Matsunaga Y, Takada S. 2021. Rigid-body fitting to atomic force microscopy images
for inferring probe shape and biomolecular structure. *PLoS Comput Biol* **17**:1–24.
doi:10.1371/journal.pcbi.1009215

Oda T, Iwasa M, Aihara T, Maéda Y, Narita A. 2009. The nature of the globular- to fibrous-actin
transition. *Nature* **457**:441–445. doi:10.1038/nature07685

Oosterheert W, Klink BU, Belyy A, Pospich S, Raunser S. 2022. Structural basis of actin
filament assembly and aging. *Nature* **611**:374–379. doi:10.1038/s41586-022-05241-8

Reynolds MJ, Hachicho C, Carl AG, Gong R, Alushin GM. 2022. Bending forces and nucleotide
state jointly regulate F-actin structure. *Nature*. doi:10.1038/s41586-022-05366-w

Rich SA, Estes JE. 1976. Detection of conformational changes in actin by proteolytic digestion:
Evidence for a new monomeric species. *J Mol Biol* **104**:777–792. doi:10.1016/0022-
2836(76)90181-9

Suarez C, Roland J, Boujemaa-Paterski R, Kang H, McCullough BR, Reymann A-C, Guérin C,
Martiel J-L, De la Cruz EM, Blanchoin L. 2011. Cofilin tunes the nucleotide state of actin
filaments and severs at bare and decorated segment boundaries. *Curr Biol* **21**:862–8.
doi:10.1016/j.cub.2011.03.064

Tanaka K, Takeda S, Mitsuoka K, Oda T, Kimura-Sakiyama C, Maéda Y, Narita A. 2018.
Structural basis for cofilin binding and actin filament disassembly. *Nat Commun* **9**:1–3.
doi:10.1038/s41467-018-04290-w

Tokuraku K, Kuragano M, Uyeda TQP. 2020. Long-range and directional allostery of actin
filaments plays important roles in various cellular activities. *Int J Mol Sci* **21**.
doi:10.3390/ijms21093209
Tokuraku K, Kurogi R, Toya R, Uyeda TQP. 2009. Novel mode of cooperative binding between
myosin and Mg2+-actin filaments in the presence of low concentrations of ATP. *J Mol Biol*
386:149–162. doi:10.1016/j.jmb.2008.12.008
Uyeda TQP, Iwadate Y, Umeki N, Nagasaki A, Yumura S. 2011. Stretching actin filaments
within cells enhances their affinity for the myosin ii motor domain. *PLoS One* **6**:e26200.
doi:10.1371/journal.pone.0026200
Zimmermann D, Santos A, Kovar DR, Rock RS. 2015. Actin Age Orchestrates Myosin-5 and
Myosin-6 Runlengths. *Curr Biol* **25**:2057–2062. doi:10.1016/j.cub.2015.06.033.

Acknowledgments
The authors acknowledge the facility and financial support of WPI Nano Life Science Institute
(WPI NanoLSI), Kanazawa University and the instrumental support of Toshio Ando. The
authors acknowledge Huan Tran from Georgia Institute of Technology, and Damien Hall and
Thao Phuong Ngo from WPI-NanoLSI, Kanazawa University for their help in the data analyses.
The authors acknowledge for the KAKENHI (Japan Society for the Promotion of Science)
supporting K.X.N. (19K06581, 23K05713, 23H02452-01) and T.Q.P.U. (#23H02452) and N.K.
(20H00327); and CREST, Japan Science and Technology Agency (JPMJCR1762 to N.K.).

Author contributions
K.X.N., H.T.V., N.K., T.Q.P.U. designed the concepts and experiments. K.X.N., H.T.V., N.K.
performed experiments, K.X.N., H.T.V., K.U. analysed the data. K.X.N., H.T.V., T.Q.P.U. wrote
the manuscript. All authors revised the manuscript.
988

989 **Conflicts of interest**
990 The authors declare no conflicts of interest. The funders have no role in interpreting the data.

Figure 1 The significant movements of the OD during F-actin to C-actin to G-actin transitions, as revealed by PCA.

(**A**) PCA of actin chains reported in 46 PDB structures (referred to **Table supplement 1**) reveals collective transition pathway between different actin conformations: G-actin (soluble monomer – blue box), F-actin (filamentous protomer reported with no cofilin bound – green box), C-actin (filamentous protomer reported within cofilactin after removal of cofilin – red box), tilted actin (reported in 3J8J and 3J8K – magenta box) and single-cofilin-bound actin, where one cofilin molecule bound to an actin protomer within an actin filament (reported in 6UBY – orange box). Protomers at the most barbed end side of a cofilin cluster in 6UC4 also belong to the latter cluster. Notably, a twinfilin-1 bound structure of soluble actin molecule (3DAW) (Paavilainen et al., 2008), previously classified as C-actin (Tanaka et al., 2018), was found between the G-actin and C-actin clusters. PDB entries are coloured with nucleotide molecules present, and structures lacking reported nucleotides are labelled "Apo". The inset shows a magnified region of the F-actin cluster (corresponding to the black box in the main graph).

(**B**) The PC1, represented by blue arrows, induces a rotational motion of the OD around hinge region, causing the nucleotide binding cleft between SD2 and SD4 (depicted as a yellow arrowhead) to close during the transition from G-actin (cyan chain) to C-actin (pink chain).

(**C**) The PC2, indicated by red arrows, results in an untwisting motion of the SD2, leading to the flattening of OD during the transformation from C-actin (pink chain) to F-actin (green chain), but without affecting the closing of nucleotide binding cleft (nearly unchanging PC1). The rotational axes in panels B and C were denoted by blue and red asterisks, respectively.

Figure 2 The variations in half helices and the number of protomer pairs per HHP in F-actin, C-actin, and cofilactin structures in relationship with PC2 analysis.

(**A-C**) Side-views (represented by blue boxes, similar to the right panel of Figure 1C) and outer-views (represented by black boxes) of F-actin, C-actin protomers without and with cofilins along long filaments are depicted. The white translucent cylinders represent the central axes of the filaments. In panel B, a yellow arrowhead highlights a portion of SD2 including D-loop that is buried deeper into the C-actin filament as a result of the reduction of PC2 (represented by the red arrow, corresponding to the red arrow in the right panel of Figure 1C).

(**D-E**) Diagrams of 2D surfaces (unwrapped from imaginary cylinders along the filaments and cut open along imaginary lines scanned by the AFM tip) illustrate the different twisting angles between adjacent protomers in the same strand appears to increase from approximately 26° to around 36° if view from the outer side, resulting in HHPs with varying numbers of detected protomer pairs (marked between red lines). The green and pink objects represent outer-views of OD domains of the focused chains, along with cofilins shown in black. The opposite chains are depicted in grey.

1089
Figure 3 Exploring the dynamics of helical structures of long actin filaments using HS-AFM.
(**A**) The provided image shows actin filaments with varying numbers of protomer pairs in one
half helix. To measure the HHPs and the number of protomer pairs per HHP, a longitudinal
section profile of two consecutive half helices was taken as depicted, following the previous
method (Hayakawa et al., 2022). Actin filaments were gently immobilized on lipid membrane
composed of DPPC/DPTAP (90/10 wt%) and imaged in F1 buffer. The offset values were set to
zero with reference to the height of the lowest position along the longitudinal section lines. The
scale bar represents 25 nm. For the detail, see Methods.
(**B**) The correlation between the counted protomer pairs from image A and the HHP probed by
HS-AFM is shown. Black symbols represent the experimental data, with the linear fit (y = 10.71
+ 3.43x, R = 0.9903). The calculated data (red symbols) also demonstrate a correlation between
the number of protomer pairs per HHP and the HHP, which is calculated as 5.5 nm multiplied by
the number of counted protomer pairs per HHP.
(**C**) The distribution of MADs is shown for half helices composed of different number of
counted protomer pairs. The overall MAD (mean ± SD) is 5.1 ± 0.3 nm (n = 15021).
(**D**) The axial distance (AD) between two adjacent actin protomers along the same long-pitch
strand in individual bare actin filaments were directly measured as the peak-to-peak distance in
four consecutive HHPs. Each protomer pair within the filament was assigned a number ranging

from 1 to 27. The ADs were measured in a time series, with the imaging rate of 0.5 seconds per frame and a total of 258 frames. These AD values for 27 protomer pairs within the actin filament in a time series of 258 frames were utilized to determine the autocorrelation coefficients. The still AFM image was shown after processing with Laplacian of Gaussian filter.

(**E**) An example of the ACF plot displays the autocorrelation coefficients on the y-axis against the time lags on the x-axis. The plotted data was fitted using a second-order exponential decay function with the time constants of (t_1, t_2), represented by the red curve.

(**F**) A histogram was constructed using the time constants obtained from many similar ACF plots and fittings obtained from different actin filaments ($n = 61$), as showcased in (D, E). The number of data points was 122. These time constants are associated with the overall trend of time decay in autocorrelation of axial distance between two adjacent protomers in the same protofilament.

1154
Figure 4 The variation of helical twists measured by HHPs, the number of protomer pairs per
HHP, and the MADs in long actin filaments through HS-AFM. Actin filaments were gently
immobilized on lipid membrane composed of DPPC/DPTAP (90/10 wt%) and imaged in F1
buffer.
(**A**) The graph demonstrates the changes in HHPs, the number of protomer pairs per HHP, and
the MADs over time. The statistical analysis using two-population *t-test* at a significance level of
$p \leq 0.05$ revealed that the difference between MAD1 (5.1 ± 0.3 nm, n = 32) and MAD2 ($5.2 \pm$

0.2 nm, n = 32) is not statistically significant (p = 0.2365). However, MAD1 and MAD3 (5.2 ±

0.3 nm, n = 32) exhibit a slight difference (p = 0.02895).

(**B**) Representative still images obtained from AFM are presented. Arrows indicate the peaks

(P1, P2, P3, P4) identified in the raw AFM images. HHP1, HHP2, and HHP3 were measured

between P1-P2, P2-P3, and P3-P4, respectively. Scale bars: 25 nm.

(**C**) The spatial ACF, plotted for each actin filament (F1, F2, F3, F4, F5), visually demonstrates

the extent of structural similarity between actin protomers' height amplitudes over distance. The

spatial ACF analysis provides an indirect measure of the degree to which height values near each

other in space are similar to one another in actin filaments (see Methods). This analysis was

conducted at different time points (the imaging rate of 0.5 s per frame and a total of 50 frames

each filament (F1-F5)). The still AFM image is shown after processing with Laplacian of

Gaussian filter. Bar: 25 nm.

1174

1175

1176

1177

1178

1179

1180

1181

1182

1183

(A) Bare actin filaments

(B) Hybrid cofilactin/bare actin

(C) Hybrid cofilactin/bare actin

Figure 5 Asymmetric helical twisting on both sides of a small cofilin cluster. Red arrowheads indicate small area of cofilin clusters (in boxes). Scale bars: 25 nm.
(**A**) A still AFM image shows normal actin filaments, along with a longitudinal section profile representing two consecutive half helices 1 and 2 (denoted HHP1, HHP2).

(**B**) A still AFM image displays a hybrid segment consisting of cofilactin and bare actin areas,
with a longitudinal section profile representing two consecutive half helices. A short filament
was observed where a small cofilin cluster cut filament at cofilactin/bare actin boundary on the
BE side, followed by two consecutive bare half helices 3 and 4 (denoted HHP3, HHP4) on the
PE side. The shortened bare half helix 3 and normal bare half helix 4 are presented. The hybrid
cofilactin/bare actin filaments, priorly made in solution, were strongly immobilized on lipid
membrane composed of DPPC/DPTAP (50/50 wt%) and imaged in F1 buffer. See Methods.
(**C**) A still AFM image depicts a hybrid segment comprising cofilactin and bare actin areas,
along with a longitudinal section profile representing two consecutive half helices 5 and 6
(denoted HHP5, HHP6). In this case, a short filament contained a small cofilin cluster (estimated
as 2 to 4 molecules) in the middle, configured by a normal bare half helix 5 on the BE side and a
shortened bare half helix 6 on the PE side. We estimated the size of a cofilin cluster by
measuring its height and width within the white box around P8, distinguishing it with the control
bare section lacking cofilin bindings (P7, P9). Despite the invisibility of cofilin molecules below,
we made the assumption that cofilin bound uniformly to two strands at a molar ratio of 1:1,
enabling us to account for their presence.
1206

Figure 6 The variation of half helices, the number of protomer pairs per HHP, and the MADs in a hybrid cofilactin/bare actin filament probed by HS-AFM.

(**A**) Representative still images obtained from AFM in **Figure 5C** are displayed. Arrows indicate the peaks (P7, P8, P9) identified in the raw AFM images. A normal bare half helix 5 (denoted HHP5) and a shortened bare half helix 6 (denoted HHP6) were measured between P7-P8 and P8-P9, respectively. Cyan and white arrows indicate the peak positions within cofilin

clusters and bare actin peaks, respectively. Boxes denote the cofilin clusters. Scale bars are 25
nm.
(**B**) The graph depicts the analysis of half helices 5 and 6, the number of protomer pairs in each
half helix (Protomer Pairs (5) and Protomer Pairs (6)), and the MAD5 and MAD6 over time.
Two-populations *t-test* was used to validate the statistical difference between MAD5 (5.0 ± 0.2
nm) and MAD6 (5.4 ± 0.3 nm). The sample size was 12.
(**C**) The spatial ACF was plotted for a normal bare half helix 5 and a shorter bare half helix 6 in a
hybrid cofilactin/bare actin filament across multiple time points. This plot visually demonstrates
the extent of structural similarity between actin protomers over distance from decorated to
undecorated cofilin region. The spatial ACF analysis provides an indirect measure of the degree
to which height values near each other in space are similar to one another in actin filaments. This
analysis was conducted at different time points (the imaging rate of 0.5 s per frame and a total of
50 frames). The peak positions of the cofilactin regions (depicted in boxes) and the bare actin
regions are indicated by cyan and white arrows, respectively. The still AFM image in A is shown
after processing with Laplacian of Gaussian filter.
1229
1230
1231
1232

1233

Figure 7 Propagation of supertwisted half helices within the cofilactin region beyond the
boundary of cofilactin and bare actin, leading to the pre-shortening of a bare half helix and
subsequent recruitment of cofilin cluster growth.

(**A**) Still AFM images show the time-dependent growth of cofilin clusters beyond the boundary.
The arrow indicates the direction of cofilin cluster growth. The peaks of cofilactin and bare actin
segments are marked by yellow and white arrowheads, respectively. Peak heights (P1-P6) were
determined in five consecutive half helices. Actin filaments were loosely immobilized on a lipid
membrane composed of DPPC and positively charged DPTAP (90/10 wt%) and imaged in an F1
buffer containing 300 nM cofilin, 1 mM ATP and 10 mM Pi. Scale bars: 25 nm.

(**B**) Examination of changes in peak heights and HHPs in the cofilactin and bare actin regions
over time. Peak heights and HHPs were measured as described previously (Ngo et al., 2015).

(**C**) Histograms of peak heights (8.6 ± 0.4 nm, 10.3 ± 0.5 nm, n = 726) and HHPs (25.3 ± 2.6 nm,
36.9 ± 3.6 nm, n = 605) were obtained from the data in (B). The significant difference between
the peaks in double Gaussian fitting was statistically examined using an *F-test*. The green and
red curves denote the fractions of F-actin and cofilactin structures, respectively.

1249

Figure 8 Cooperative binding of cofilin is heavily dependent on the dynamic and cooperative conformational changes involving helical twisting within actin filaments.

(**A**) The cooperative binding of cofilin to actin filaments was significantly inhibited when the torsional motion during helical twisting within the filaments was suppressed. The filaments were strongly attached onto a highly positively charged lipid membrane containing DPPC/DPTAP (75/25 wt%) and imaged in F1 buffer containing 2000 nM cofilin and 0.5 mM ATP.

(**B, C, D**) The normal cooperative bindings of cofilin to actin filaments were observed when the filaments were loosely immobilized on weakly positively charged lipid membrane comprising DPPC/DPTAP (90/10 wt%) and imaged in F1 buffer under different conditions: 300 nM cofilin + 0.5 mM ATP (B), 300 nM cofilin + 1 mM ADP (C), and 75 nM cofilin + 1 mM ATP + 10 mM Pi (D). The changes in peak heights and HHPs of a representative actin filament under each condition were analyzed over time. Histograms were constructed by analyzing the changes in peak heights and HHPs of all actin filaments in the observed field during the growth of the cofilin cluster at various time intervals after adding cofilin. The statistical significance of the peaks in the histograms was examined using an *F-test*. The green and red curves denote the fractions of F-actin and cofilactin structures, respectively. Time labels in the still AFM images indicate the elapsed time after adding cofilin to the imaging buffer. Peaks of bare actin and cofilactin are denoted by white and yellow arrowheads, respectively. Scale bars: 25 nm.

1271

Figure 9 Comparison of the growth rate of cofilin clusters within the shortened bare half helices that resemble the C-actin structure adjacent to cofilin clusters on the PE side, for both F-ADP-actin and F-ADP.Pi-actin.

(**A**) In this cartoon representation, hybrid cofilactin/bare actin filaments are depicted, consisting of four half helices. Two of these half helices are fully decorated with cofilin clusters in the middle, followed by a shortened bare half helix and a normal long bare half helix on the PE and BE sides, respectively. P1-P5 were labels indicating peak heights at individual crossover points. To make a meaningful comparison, the lag time measurements were initiated exclusively when a bare half helix between P1 and P2 experienced shortening, and they continued until a newly matured cluster emerged in P1 within the shortened bare half helix (P1-P2) adjacent to the preformed cofilin cluster (P2). Each fully decorated supertwisted half helix contains 11 cofilin molecules (referred to **Figure supplement 1I**). To determine the mean growth rate of the cofilin cluster within a shortened bare half helix adjacent to the cofilin cluster, 11, the number of cofilin molecules in one half helix, was divided by the lag time in seconds.

(**B**) This part presents a comparison of the growth rate of cofilin clusters within shortened bare half helices that resemble the C-actin structure adjacent to cofilin clusters on the PE side. This comparison was made when actin filaments were incubated with the same cofilin concentration (i.e., 300 nM) in the F1 buffer containing 1 mM ADP/5 U/ml hexokinase/10 mM glucose or 1

mM ATP + 10 mM Pi, resulting in filamentous protomers representing F-ADP-actin or F-
ADP.Pi-actin structures. The growth rates (mean ± SD molecules/HHP/s) were measured for 22
shortened bare half helices in F-ADP-actin (10.0 ± 8.3) and 28 ones in F-ADP.Pi-actin (8.3 ±
7.7). Using the two-population *t-test*, the results showed that the difference was not significant (p
= 0.47846) at the $p \leq 0.05$ level.
(**C, D**) Histograms illustrate the distribution of the HHP (mean ± SD) in control actin filaments
for three conditions: (i) phalloidin-stabilized F-actin (37.0 ± 2.3 nm), (ii) F-ADP-actin (36.9 ±
3.8 nm), and (iii) F-ADP.Pi-actin (37.0 ± 2.8 nm), with each condition having 2487 data points.
For comparison, single Gaussian fits in D were integrated and displayed in C. The inset within C
exhibited fractions of the naturally supertwisted half helices with HHPs shorter than 30 nm.
Utilizing one-way ANOVA with a significance level of $p \leq 0.05$, there were significant
differences observed between (i) and (ii) (p = 0.00837, F = 6.95754) as well as between (ii) and
(iii) (p = 0.04080, F = 4.18633), primarily attributable to the statistical difference in the variance
of HHP between condition (ii) vs. (i) and (iii). The difference observed between (i) and (iii) was
not significant (p = 0.58272, F = 0.30189).

1336

Figure 10 The hypothesis regarding the actin structure-dependent preferential cooperative binding of cofilin. For simplicity, a single actin protofilament comprising 6 protomers labelled from -2 to 8 is shown, both without and with cofilin. The black and red arrows represent stable bonds and unstable bonds between domains of two adjacent actin protomers in the same strand, respectively. The broken red arrows indicate completely disrupted bonds between domains of two adjacent actin molecules. Additionally, broken red and purple lines show positions sensitive to being severed, with the purple lines indicating a stronger severing activity.

(**A**) The conformational transition between G-actin, F-actin (bound with ADP.Pi, ADP), and C-actin structures. In the lower model of C-actin, the binding of a single cofilin molecule to an actin filament induces a small movement of SD2, causing a rotational shift of approximately 5° in the OD relative to the ID. In the upper one, when the cofilin clusters bind, they induce a larger rotation of approximately 10° in the OD (Galkin et al., 2011; Tanaka et al., 2018).

(**B**) A single protofilament of an actin filament. Upon polymerization, OD rotates relative to the ID to close nucleotide binding cleft and untwist the OD to flatten actin molecule. The orange color corresponds to OD (SD1+SD2), while green corresponds to ID (SD3+SD4) (Oda et al., 2009).

(**C**) A single protofilament of an actin filament bound to a single cofilin (Huehn et al., 2020; Tanaka et al., 2018).

(D) A single protofilament of an actin filament bound to cofilin clusters. Actin conformational changes induced by cofilin clusters are local and limited to protomers directly contacting bound cofilin. The cofilin bindings accelerate a rapid transition from F-ATP-actin to F-ADP.Pi-actin and F-ADP-actin states of protomers on the PE side. Cofilin tunes adjacent protomers, resembling F-ADP-actin structure on the PE side, to expand the clusters The cooperative binding of cofilin requires a critical size of a cofilin cluster set at 2. Notably, the severing of the actin filament is predominantly observed at the junction between cofilactin and bare actin (Huehn et al., 2020; Suarez et al., 2011; Tanaka et al., 2018).

(E-F) A single protofilament of an actin filament bound to cofilin clusters in the authors' previous study (Ngo et al., 2015) and in the current study. Our hypothesis supports that actin conformational changes induced by cofilin clusters are not local and not limited to protomers directly contacting bound cofilin (C-actin) on the PE side. Cofilin clusters have the ability to shorten a bare half helix on the PE side by reducing the number of protomers. Additionally, cofilin cluster not only has the capacity to convert F-actin to C-actin but also allosterically expands the C-actin structure from the bound to unbound region on the PE side (**E**). The MAD_c in the undecorated C-actin like structure region are elongated compared to MAD_f in F-actin region. The shortening of half helices and the elongated MAD_c play a pivotal role in facilitating the preferential cooperative binding of cofilin to actin protomers via F-actin to C-actin transition pathway. If cofilin does not promptly attach to protomers in shortened bare half helix region (as depicted in E-F), the protomers within C-actin like structure regions can readily open the nucleotide binding clefts and transition back to G-actin state via C-actin to G-actin transition, ultimately severing the filament at or near boundaries (broken lines).

(G) HS-AFM measurements of MAD values measured in half helices of filaments representing F-actin (MAD_f) and undecorated C-actin like structure (MAD_c) adjacent to cofilin cluster on the PE side.

(H) A hypothesis describing how the function of cofilin is regulated by the torsional flexibility during helical twisting of actin filaments. The F-ADP-actin has increased the fractions of naturally supertwisted helical structures and the torsional flexibility during helical twisting, thereby creating some proposed C-actin like structures to facilitate the initial binding and cluster expansion of cofilin, compared to F-ADP.Pi-actin. To emphasize the inhibitory effect of Pi on the initial binding of cofilin, we depict a simplified model of F-ADP.Pi-actin, featuring very infrequent structures resembling C-actin. As a result, the initial binding of cofilin and growing of critical cluster set at 2 to 4 cofilin molecules are more challenging compared to F-ADP-actin.

1401
1402
Figure supplement 1 A closer examination of PC1 and PC2 in the F-actin clusters.
Within these clusters of F-actin, similar to the inset of **Figure 1A**, there is a larger presence of C-
actin like structures in F-ADP-actin compared to F-ADP.Pi-actin. However, the magnified views
of the F-actin cluster also highlight structural resemblance between F-ADP-actin and F-ADP.Pi-
actin, as recently elucidated by (Reynolds et al., 2022) (**A**) and (Oosterheert et al., 2022) (**B**).
Significantly, the F-ADP-actin structure (8D13) and F-ADP.Pi-actin structure (8D14) exhibit
remarkably similar PC2 values. Furthermore, the PC2 results demonstrate notable similarities
between Mg^{2+}-F-ADP-actin (8A2T) and Mg^{2+}-F-ADP.Pi-actin (8A2S), as well as Mg^{2+}-F-
ADP.BeF_3-actin (8A2R).

Figure supplement 2 Pseudo-AFM images featuring long filaments representing F-actin and C-actin structures. It is noted that both filamentous structures share the same length and contain an equal number of actin protomers, specifically 62 protomers. However, the filament representing C-actin exhibits a greater number of half helices than F-actin. This difference can be attributed to the decrease in the number of actin protomers per HHP in C-actin structure.

(**A**) A pseudo-AFM image featuring a long filament representing F-actin structure constructed using the 6VAU. The scale bar represents 25 nm.

(**B**) Longitudinal section profiles of the middle half helices in panel A were used to identify the HHPs (distances between major peaks marked in red, reported in nm) and the numbers of actin protomer pairs per HHP (as counted major and small peaks marked in red and green within each HHP subtracted by 0.5). The offset values of the height profile were set to zero by referring to virtual mica surface, where filament was placed in the pseudo-AFM images. The MADs were computed and shown in brackets. The HHP, the number of protomers per HHP, and MAD were shown accordingly.

(**C**) A side-view of the long actin filament imaged in panel A. To enhance visibility of protomer pairs, one of the chains was depicted as translucent.

(**D-F**) Panels D-F follow a similar format as panels A-C but display the results obtained for a long filament representing C-actin structure (cofilins are removed from initial cofilactin structure reported in 6VAO). The HHP, the number of protomers per HHP, and axial distance (AD) were shown in E accordingly.

(**G-I**) Panels G-I also followed a similar format as panels A-C, but showcased the results obtained for a long cofilactin filament (constructed from 6VAO). Accurately detecting actin pairs in this case was challenging due to cofilin decoration and the tip-sample dilation effect, as explained by (Ando, 2022).

1466
Figure supplement 3 A schematic overview of the spatial autocorrelation analysis. In panel (**a**),
an HS-AFM image is depicted, with a line profile drawn along an actin filament. Panels (**b**) and
(**c**) show the height profiles before and after applying the bandpass filter (BPF), respectively. The
spatial autocorrelation function (ACF) and envelope curve before and after BPF processing and
after applying the Hilbert transform are displayed in panels (**d**) and (**e**), respectively. Panels (**f**)
and (**g**) illustrate the surface topography of the actin protomers' heights along filaments, which
were used in making the spatial ACF (as depicted by the envelope curve). The green curves in (**f**)
represent the regular periodicity of distance and flatter slopes in the heights between protomers,
while the red curves in (**g**) depict the irregular periodicity of distance and steeper slopes in the
heights between protomers. In theory, the envelope curve (green) versus distance in (**f**) should
linearly decrease. In contrast, the envelope curve (red) versus distance in (**g**) shows a more
pronounced decrease. During the actual analysis, the selected space intervals (ΔD) for measuring
the degree to which height values near each other in space are similar to one another in F-actin
and hybrid cofilactin/bare actin segments were 0.4 nm and 0.3 nm, respectively. The
measurement process involved shifting these space intervals from left to right and up to down,

with increasing interval values from the initial position (0 ΔD). P1-P7 denote the peak positions
of individual protomers.

1527
1528 **Figure supplement 4** The propagation of the supertwisted half helices induced by cofilin
1529 clusters results in the shortening of a bare half helix beyond the cofilactin/bare actin boundary.

The HHPs and peak heights were determined using methods similar to our previous reports (Hayakawa et al., 2022; Ngo et al., 2015). Scale bars: 25 nm. Arrowheads indicated the peaks of cofilin clusters. h_0 and h_1 represented the peak heights in two consecutive HHPs of bare actin and cofilactin segments. The offset values were set to zero with reference to the height of the lowest position along the longitudinal section lines. The height and width differences were utilized to determine the bare actin and cofilactin segments. The bare actin, cofilactin, and hybrid cofilactin/bare actin filaments, priorly made in solution, were strongly immobilized on lipid membrane composed of DPPC/DPTAP (50/50 wt%) and imaged in F1 buffer. See Methods.

(**A**) A still AFM image presents a normal actin filament, accompanied by a longitudinal section profile across two consecutive half helices to measure the HHP1 and HHP2 and the number of actin protomer pairs per HHP accordingly. The MADs were computed from AFM analysis and shown.

(**B**) A still AFM image depicts a cofilactin filament, accompanied by a longitudinal section profile across two consecutive half helices to measure the HHP3 and HHP4 and the numbers of actin protomer pairs per HHP accordingly. The MADs were calculated from our AFM analysis and shown.

(**C**) A still AFM image showcases a hybrid bare actin/cofilactin filament, accompanied by a longitudinal section profile across two consecutive half helices to measure the shortened bare HHP5 and shortened cofilactin HHP6 and the number of actin protomers per HHP. The MADs were calculated from our AFM analysis and shown.

1576

Figure supplement 5 The analysis of the variation in helical twisting by measuring HHPs, the number of protomer pairs per HHP, and MADs in a hybrid bare actin/cofilactin segment using HS-AFM.

(**A**) Representative still AFM image of a hybrid bare actin/cofilactin segment in **Figure supplement 4C** is provided. The peaks (P7, P8, P9, P10) in the raw AFM image are indicated by white and cyan arrows for the bare actin and cofilactin regions, respectively. HHP5, HHP6, and HHP7 and the number of protomer pairs within each HHP were measured between P7-P8, P8-P9, and P9-P10, respectively and MAD5, MAD6, and MAD7 were computed accordingly. Scale bars: 25 nm.

(**B**) The variation of HHPs, the number of protomer pairs per HHP, and MADs in shortened bare and cofilactin regions were examined over time.

(**C**) A comparison between MAD5 (5.7 ± 0.5 nm) observed in the shortened bare area and MAD6 (5.9 ± 0.3 nm) along with MAD7 (6.1 ± 0.3 nm) within the cofilactin region was presented. A two-population *t-test* with a sample size of n = 13 was used.

(D) The relationship between the HHP and the number of protomer pairs per HHP measured for
C-actin like structures within the shortened bare half helix adjacent to cofilin cluster, as explored
through HS-AFM, was illustrated using black symbols. The collected data was subjected to
fitting with a linear equation (y = 10.99 + 3.21x, R = 0.9975) in the region of the shortened bare
half helix situated close to the boundary of cofilactin and bare actin. Calculated data, depicted
with red symbols, presented the correlation between the number of protomer pairs and the HHP
that was calculated as 5.5 nm multiplied by the number of protomer pairs within each HHP.
(E) The histograms depict the distribution of MADs for C-actin like structures within the
shortened bare half helices adjacent to cofilin clusters consisting of different number of protomer
pairs.

(A) + 2000 nM cofilin + 0.5 mM ATP/ filaments were immobilized on 25% DPTAP membrane

(B) + 300 nM cofilin + 0.5 mM ATP/ filaments were immobilized on 10% DPTAP membrane

(C) + 300 nM cofilin + 1 mM ADP/ filaments were immobilized on 10% DPTAP membrane

(D) + 75 nM cofilin + 1 mM ATP + 10 mM Pi/ filaments were immobilized on 10% DPTAP membrane

Figure supplement 6 Cooperative binding of cofilin is strongly dependent on the cooperative conformational changes involving helical twisting in actin filaments. Arrowheads denote cofilin clusters. The time labels indicate the elapsed time after adding cofilin to the imaging buffer. Peaks of bare actin and cofilactin are denoted by white and yellow arrowheads, respectively. (**A**) When the torsional motion of actin filaments was suppressed, the cooperative binding of cofilin to the filaments was greatly inhibited. The filaments were strongly attached onto a highly positively charged lipid membrane containing DPPC/DPTAP (75/25 wt%) and imaged in F1 buffer containing 2000 nM cofilin and 0.5 mM ATP. Scale bars: 25 nm. (**B, C, D**) The cooperative binding of cofilin to actin filaments was examined under different conditions: incubation in F1-buffer containing 300 nM cofilin + 0.5 mM ATP (B), 300 nM cofilin + 1 mM ADP (C), and 75 nM cofilin + 1 mM ATP + 10 mM Pi (D). Scale bars: 25 nm.

Table supplement 1 List of 46 protein data bank (PDB) structures that were utilized in the PCA,
as depicted in **Figure 1**. Due to the presence of varying structures among different chains in
some PDB structures, each chain was considered as an individual structure. Notably, in the case
of 6UC4, chains E and H were combined and renamed as chain E (E&H = E), while chains F and
G were combined and renamed as chain F (F&G = F). Similarly, in the case of 6UBY, chains G
and H were combined and renamed as chain G (G&H = G).
1663

PDB ID	List of actin chains	Method	Resolution (Å)	Nucleotide	Protein bound	First author	Year
1ATN	A	X-RAY	2.8	ATP	DNASE I	KABSCH	1990
1J6Z	A	X-RAY	1.54	ADP-RHO		OTTERBEIN	2001
2BTF	A	X-RAY	2.55	ATP	PROFILIN	SCHUTT	1993
2ZWH	A	X-RAY	3.3	ADP		ODA	2009
3DAW	A	X-RAY	2.55	ATP	TWINFILIN-1	PAAVILAINEN	2008
3HBT	A	X-RAY	2.7	ATP		WANG	2010
3J0S	A, B, C, D, E, F, G, H, I, J, K, L	EM	9		COFILIN-2	GALKIN	2011
3J8I	D, E, F, G, H	EM	4.7	ADP		GALKIN	2015
3J8J	A, B, C, D, E, F, G, H, I, J, K	EM	12			GALKIN	2015
3J8K	A, B, C, D, E, F, G, H, I, J	EM	12			GALKIN	2015
3MFP	A	EM	6.6	ADP		FUJII	2010
4A7N	A, B, C, D, E	EM	8.9	ADP		BEHRMANN	2012
5ONV	A, B, C, D, E	EM	4.1	ADP		MERINO	2018
5OOC	A, B, C, D, E	EM	3.6	ADP JASP		MERINO	2018
5OOD	A, B, C, D, E	EM	3.7	ADP-Pi JASP		MERINO	2018
5OOE	A, B, C, D, E	EM	3.6	ANP		MERINO	2018
5OOF	A, B, C, D, E	EM	3.4	ADP-BeF$_x$		MERINO	2018
5YU8	A, B, C, D, E	EM	3.8	ADP	COFILIN-2	TANAKA	2018
6BNO	A, B, C, D, E, F, G, H	EM	5.5	ADP		GUREL	2017
6BNU	A, B, C, D, E, F, G, H	EM	7.5			GUREL	2017
6DJM	A, B, C, D	EM	3.1	AMPPNP		CHOU	2019
6DJN	A, B, C, D	EM	3.1	ADP-Pi		CHOU	2019
6DJO	A, B, C, D	EM	3.6	ADP		CHOU	2019
6FHL	A, B, C, D, E	EM	3.3	ADP-Pi		MERINO	2018
6KLL	A, B, C, D	EM	3	ADP		ODA	2020
6KLN	A, B, C, D	EM	3.4	ADP		ODA	2020
6UBY	A, B, C, D, E, F, G	EM	7.5	ADP	COFILIN-1	HUEHN	2020
6UC0	A, B, C, D, E, F, G	EM	7.5	ADP	COFILIN-1	HUEHN	2020
6UC4	A, B, C, D, E, F, J, K, L	EM	9.2	ADP	COFILIN-1	HUEHN	2020
6VAO	D, A, B, C, E	EM	3.4	ADP	COFILIN-1	HUEHN	2020
6VAU	B, A, C, D, E	EM	3.5	ADP		HUEHN	2020
7Q8B	A, B, C, D, E	EM	3.3	ADP-Pi		KOTILA	2022
7Q8C	A, B, C, D, E	EM	2.72	ADP		KOTILA	2022
7Q8S	A, C, D, E, F	EM	3.4	ADP	ADF/COFILIN	KOTILA	2022
8A2R	C, A, B, D, E	EM	2.17	ADP-BeF$_3$		OOSTERHEERT	2022
8A2S	C, A, B, D, E	EM	2.22	ADP-Pi		OOSTERHEERT	2022
8A2T	C, A, B, D, E	EM	2.24	ADP		OOSTERHEERT	2022

8A2U	C, A, B, D, E	EM	2.21	ADP-BeF$_3$		OOSTERHEERT	2022
8A2Y	C, A, B, D, E	EM	2.15	ADP-Pi		OOSTERHEERT	2022
8A2Z	C, A, B, D, E	EM	2.15	ADP		OOSTERHEERT	2022
8D13	A, B, C	EM	2.43	ADP		REYNOLDS	2022
8D14	A, B, C	EM	2.51	ADP-Pi		REYNOLDS	2022
8D15	A, B, C, D, E, F, G	EM	3.61	ADP		REYNOLDS	2022
8D16	A, B, C, D, E, F, G	EM	3.71	ADP-Pi		REYNOLDS	2022
8D17	A, B, C, D, E, F, G	EM	3.69	ADP		REYNOLDS	2022
8D18	A, B, C, D, E, F, G	EM	3.66	ADP		REYNOLDS	2022

Table supplement 2 The lists of chains and matrices utilized to build the long filaments

PDB ID	Chains	RMSD (Å)	Transformation matrix as Euler angles		
6VAU	BA	0.000497	[166.588953	-2.1209466e-01	-2.62674783e-02]
8A2S	ED	0.000704	[-166.93535314	-10.63461097	-5.38436149]
8A2T	AB	0.000704	[-167.39029871	-13.65820025	-6.3200023]
3MFP	AB	0.000398	[166.656064	2.13617418e-05	7.68141187e-05]
6UC4	EF	0.619515	[164.58920564	-0.207997407	-0.80958005]
6VAO	CA	0.000699	[162.515987	-4.24094412e-02	-1.82361154e-01]
5YU8	ED	0.000561	[-162.100031	-4.00777476e-05	4.03030046e-05]
3J0S	KL	0.001134	[-162.12014856	-0.17228856	-0.18642638]
*For cofilactin filament					
6VAO*	CAJF	0.000699	[162.515987	-4.24094412e-02	-1.82361154e-01]

Table supplement 3 Variation during helical twisting measured by HHPs, the number of protomer pairs per HHP, and the MAD_f observed for protomers representing F-actin structures within long bare actin filaments, as probed by HS-AFM. The total number of HHPs observed was 2226. The total number of protomer pairs used for estimating MAD_f was calculated by multiplying the number of protomer pairs per HHP by the number of HHPs.

The number of protomer pairs per HHP	HHPs (nm)	MAD_f (nm)	The number of HHPs (proportion)	The total number of protomer pairs
4.5	28.6 ± 1.4	6.3 ± 0.3	5 (0.2%)	22.5
5.5	29.3 ± 1.8	5.3 ± 0.3	182 (8.2%)	1001
6.5	33.1 ± 1.3	5.1 ± 0.2	1355 (60.9%)	8807.5
7.5	37.2 ± 2.0	5.0 ± 0.3	624 (28.0%)	4680
8.5	39.1 ± 2.4	4.6 ± 0.3	60 (2.7%)	510

Table supplement 4 Variation during helical twisting measured by HHPs, the number of protomer pairs per HHP, and the MAD_c observed for protomers resembling C-actin structures within shortened bare half helices adjacent to the cofilactin/bare actin boundary, as probed by HS-AFM. The total number of HHPs observed was 34. The total number of protomer pairs used for estimating MAD_c was calculated by multiplying the number of protomer pairs by the number of HHPs.

The number of protomer pairs per HHP	HHP (nm)	MAD_c (nm)	The number of HHPs (proportion)	The total number of protomer pairs
3.5	22.2 ± 0.2	6.3 ± 0.1	2 (5.9%)	7
4.5	26.1 ± 1.4	5.8 ± 0.3	21 (61.8%)	94.5
5.5	28.5 ± 0.9	5.2 ± 0.2	9 (32.4%)	49.5

1809	**Table supplement 5** The correlation between the time it takes to shorten a bare HHP beyond the
1810	cofilactin/bare actin boundary and to form a newly matured cofilin cluster in a pre-shortened half
1811	helix. The lag time (*t*) represents the duration required for a newly saturated cofilin cluster to
1812	bind and decorate onto a bare half helix adjacent to the preformed cluster after this bare half
1813	helix has been shortened by approximately 25%. This table is related to **Figure 7**.
1814	

The time required to form a newly matured cofilin cluster in a pre-shortened half helix		The duration needed to shorten a bare half helix beyond the cofilactin/bare actin boundary				
		P1 – P2	**P2 – P3**	**P3 – P4**	**P4 – P5**	**P5 – P6**
P1	0 s					
P2	0 s	0 s				
P3	5.0 s		2.0 s (*t* = 3.0 s)			
P4	36 s			30.5 s (*t* = 5.5 s)		
P5	54.5 s				54 s (*t* = 0.5 s)	
P6	58.5 s					58.5 s (*t* = ~0 s)

	Peak Height (nm)		Half Helical Pitch (nm)		Frequency
	Peak 1 (proportion) [F-test, p]	Peak 2 (proportion) [F-test, p]	Peak 1 (proportion) [F-test, p]	Peak 2 (proportion) [F-test, p]	
(A) 2000 nM cofilin, 0.5 mM ATP/ on DPPC/DPTAP (75/25 wt%)	8.2 ± 0.8 (100%)	ND	34.3 ± 2.6 (100%)	ND	Height: 1930 HHP: 1556
(B) 300 nM cofilin, 0.5 mM ATP/ on DPPC/DPTAP (90/10 wt%)	9.2 ± 0.4 (47.0%) [0.0204]	10.5 ± 0.7 (53.0%) [0.0204]	28.5 ± 3.0 (55.1%) [0.0135]	37.4 ± 3.6 (44.9%) [0.0135]	Height: 2212 HHP: 1780
(C) 300 nM cofilin, 1 mM ADP/ on DPPC/DPTAP (90/10 wt%)	8.4 ± 0.5 (54.8%) [0.0081]	9.9 ± 0.7 (45.2%) [0.0081]	30.1 ± 3.8 (70.3%) [0.0195]	37.8 ± 2.9 (29.7%) [0.0195]	Height: 3183 HHP: 2779
(D) 75 nM cofilin, 1 mM ATP, 10 mM Pi/ on DPPC/DPTAP (90/10 wt%)	8.3 ± 0.7 (55.3%) [1.48×10^{-6}]	11.0 ± 1.0 (44.7%) [1.48×10^{-6}]	30.3 ± 2.9 (25.2%) [0.0002]	38.7 ± 3.6 (74.8%) [0.0002]	Height: 2316 HHP: 1865

Legend of videos

Videos can be downloaded here:

Video 1 Possible transition pathways from G-actin to C-actin (constructed by +PC1) and C-actin to F-actin (constructed by +PC2), and vice versa, constructed by -PC1 and -PC2, respectively. The transition from G-actin to F-actin ultimately requires both +PC1 + PC2. Related to **Figure 1**.

Video 2 HS-AFM imaging to investigate the dynamic structures during helical twisting by measuring the variations of HHPs, the number of actin protomers per HHP, and MADs in F-actin structures within long actin filaments. Our results provided strong evidence supporting the variable helical twisting within actin filaments. However, the results contrast the existing rigid and inflexible depiction of helical twists in cryo-EM and X-ray diffraction structures. Actin filaments were gently immobilized on lipid membrane composed of DPPC/DPTAP (90/10 wt%). Bars: 25 nm. Imaging rate: 2 frames per second (fps). Video plays at 5 fps. Related to **Figure 3**.

Video 3 HS-AFM imaging to investigate the dynamic structures during helical twisting by measuring the variations of HHPs, the number of actin protomers per HHP, and MADs in the normal and shorter half helices on the BE and PE sides of a cofilin cluster. Bars: 25 nm. Imaging rate: 2 fps. Video plays at 5 fps. Related to **Figure 5**.

Video 4 HS-AFM imaging to demonstrate the preferentially cooperative binding of cofilin toward a pre-shortened bare half helix resembling C-actin structures along actin filaments. Even with 10 mM Pi, the cofilin cluster could supertwist the half helices in the actin filaments. Bars: 25 nm. Imaging rate: 2 fps. Video plays at 5 fps. Related to **Figure 7**.

Video 5 Cooperative binding of cofilins along actin filaments was greatly inhibited when actin filaments were strongly immobilized on lipid membrane composed of DPPC/DPTAP (75/25 wt%). The strong immobilization hindered the torsional motion of actin filaments during helical twisting, resulting in suppressing cofilin binding and cluster expansion. Actin filaments were imaged in an F1 buffer containing 2000 nM cofilin and 0.5 mM ATP. Time label denotes the elapsed time after addition of cofilin. Bars: 25 nm. Imaging rate: 2 fps. Video plays at 10 fps. Related to **Figure 8A**.

Video 6 Cooperative binding of cofilins along actin filaments was normal when actin filaments were loosely immobilized on lipid membrane composed of DPPC/DPTAP (90/10 wt%). Actin filaments were imaged in an F1 buffer containing 300 nM cofilin and 0.5 mM ATP. Time label denotes the elapsed time after addition of cofilin. Bars: 25 nm. Imaging rate: 2 fps. Video plays at 10 fps. Related to **Figure 8B**.

Video 7 Cooperative binding of cofilins along actin filaments was normal when actin filaments were loosely immobilized on lipid membrane composed of DPPC/DPTAP (90/10 wt%). Actin filaments were imaged in an F1 buffer containing 300 nM cofilin and 1 mM ADP (see Methods). Time label denotes the elapsed time after addition of cofilin. Bars: 25 nm. Imaging rate: 2 fps. Video plays at 10 fps. Related to **Figure 8C**.

1906

1907 **Video 8** Growth of cofilin cluster along actin filaments was normal when actin filaments were
1908 loosely immobilized on lipid membrane composed of DPPC/DPTAP (90/10 wt%). Even with 10
1909 mM Pi, the cofilin cluster could supertwist the half helices in the actin filaments. Actin filaments
1910 were imaged in an F1 buffer containing 75 nM cofilin and 1 mM ATP+10 mM Pi (See
1911 Methods). Time label denotes the elapsed time after addition of cofilin. Bars: 25 nm. Imaging
1912 rate: 2 fps. Video plays at 10 fps. Related to **Figure 8D**.
1913