Deciphering the actin structure-dependent preferential cooperative binding of cofilin

Kien Xuan Ngo[1,#,*], Huong T Vu[2,#], Kenichi Umeda[1], Minh-Nhat Trinh[3], Noriyuki Kodera[1], Taro Q.P. Uyeda[4]

[1]Nano Life Science Institute (WPI-NanoLSI), Kanazawa University, Kanazawa, Ishikawa 920-1192, Japan.
[2]Centre for Mechanochemical Cell Biology, Warwick Medical School, Coventry CV4 7AL, UK.
[3]School of Electrical and Electronic Engineering, Hanoi University of Science and Technology, Hanoi, Vietnam.
[4]Department of Physics, Faculty of Advanced Science and Engineering, Waseda University, Shinjuku, Tokyo 169-8555, Japan.

# These authors equally contributed.
* Corresponding authors
Email: ngoxuankien@staff.kanazawa-u.ac.jp

Abstract

The mechanism underlying the preferential and cooperative binding of cofilin and the expansion of clusters toward the pointed-end side of actin filaments remains poorly understood. To address this, we conducted a principal component analysis based on available filamentous actin (F-actin) and C-actin (cofilins were excluded from cofilactin) structures and compared to monomeric G-actin. The results strongly suggest that C-actin, rather than F-ADP-actin, represented the favourable structure for binding preference of cofilin. High-speed atomic force microscopy explored that the shortened bare half helix adjacent to the cofilin clusters on the pointed end side included fewer actin protomers than normal helices. The mean axial distance (MAD) between two adjacent actin protomers along the same long-pitch strand within shortened bare half helices was longer (5.0–6.3 nm) than the MAD within typical helices (4.3–5.6 nm). The inhibition of torsional motion during helical twisting, achieved through stronger attachment to the lipid membrane, led to more pronounced inhibition of cofilin binding and cluster formation than the presence of inorganic phosphate (Pi) in solution. F-ADP-actin exhibited more naturally supertwisted half helices than F-ADP.Pi-actin, explaining how Pi inhibits cofilin binding to F-actin with variable helical twists. We propose that protomers within the shorter bare helical twists, either influenced by thermal fluctuation or induced allosterically by cofilin clusters, exhibit characteristics of C-actin-like structures with an elongated MAD, leading to preferential and cooperative binding of cofilin.

Keywords: actin, cofilin, actin binding proteins, high-speed atomic force microscopy, principal component analysis.

Introduction
Cofilin is a member of the actin-depolymerizing factor/cofilin (ADF/cofilin) family that is present in all eukaryotes. There are two major isoforms of cofilin in mammals: cofilin 1 and cofilin 2, predominantly found in non-muscle and muscle tissues, respectively (Bamburg and Bernstein, 2010). Cofilin is widely recognized as a key regulator of actin filament dynamics, particularly in nonequilibrium assembly and disassembly processes. Cofilin promotes the depolymerization and severing of actin filaments in a concentration-dependent manner

(Andrianantoandro and Pollard, 2006) and collaborates or competes with other actin binding
proteins (ABPs) in this process (Bibeau et al., 2021; Grintsevich et al., 2016; Lappalainen and
Drubin, 1997; Ngo et al., 2016).
The structure of an actin filament has traditionally been represented as a simplistic atomic
model derived from the G-actin structure in the Holmes model (Holmes et al., 1990) without
considering the conformational changes occurring within each protomer during actin
polymerization (Kasai et al., 1962; Rich and Estes, 1976). The rigid-body rotation of the G-actin
crystal structure used in the Holmes model prompted the development of accurate models that
consider structural changes within protomers, such as the X-ray diffraction Lorenz model
(Lorenz et al., 1993) and Oda model (Oda et al., 2009).
The actin protomer consists of two domains, the inner domain (ID) and the outer domain
(OD), with relatively limited contact between them. The polypeptide chain traverses between
these domains twice: at the loop centered around residue Lys336 and at the linker helix Gln137-
Ser145, which functions as the axis of a hinge between the domains. As a result, two clefts are
formed between these domains (Dominguez and Holmes, 2011). The upper cleft, located
between subdomain (SD2) and SD4, serves as a binding site for nucleotides and divalent ions
like Mg^{2+}. On the other hand, the lower cleft, located between SD1 and SD3, is characterized by
residues Tyr143, Ala144, Gly146, Thr148, Gly168, Ile341, Ile345, Leu346, Leu349, Thr351, and
Met355, which are predominantly hydrophobic. This cleft serves as the primary binding site for
most ABPs including cofilin and is also responsible for critical longitudinal contacts between
actin protomers within the filament (Dominguez, 2004; Fujii et al., 2010; Oda et al., 2009).
The actin filament is a double-stranded helical structure formed by the crossover of two
long-pitch helices; the distance between the crossover points is known as the half helical pitch
(HHP), which is approximately 36 nm (Fujii et al., 2010; Holmes et al., 1990; Lorenz et al.,
1993; Oda et al., 2009). A canonical half helix is composed of 13 actin protomers (6.5 protomer
pairs). Thus, the mean axial distance (MAD) between two adjacent protomers along the same
long-pitch strand is 5.5 nm (36 nm/(13/2)) (Egelman, 1994). However, long actin filaments
exhibit variable helical twists (Egelman et al., 1982). Recent cryo-EM studies have demonstrated
the atomic structures of short actin segments with cofilin (referred to as cofilactin) (Galkin et al.,
2011; Huehn et al., 2020; Tanaka et al., 2018). When cofilin binds to actin in a 1:1 ratio to form
a cofilactin segment, the helices became supertwisted, and the HHP is decreased by
approximately 25% (McGough et al., 1997), reducing the number of actin protomers to ~11 in
each half helix (5.5 protomer pairs) (PDB 3J0S) (Galkin et al., 2011). In a simple model,
assuming that the MAD remains unchanged in both bare actin and cofilactin segments, the
difference in the number of actin protomers impacts the formation of short and long HHPs.
Nonetheless, whether the alterations in MAD dynamics are comparable between normal and
supertwisted half helices remains uncertain. Therefore, additional research is needed to
investigate this particular issue.
In this study, we utilized several techniques, including principal component analysis
(PCA) and high-speed atomic force microscopy (HS-AFM). We aimed to understand how the
allosteric conformational changes induced by cofilin clusters in neighboring bare actin filaments
contribute to the preferential binding of additional cofilin molecules toward the pointed-end (PE)
side, as reported in our previous study (Ngo et al., 2015). Therefore, we generated pseudo AFM
images representing long bare actin filaments (referred to as F-actin structures) and cofilactin
filaments without cofilin molecules (referred to as C-actin structures) using available protein data
bank (PDB) structures such as 6VAU (bare actin filament from a partially cofilin-decorated

sample) and 6VAO (human cofilin-1 decorated actin filament) (Huehn et al., 2020). We
measured the HHPs and the number of actin protomers per HHP in pseudo-AFM images and
used them as the reference values for the data obtained from our real HS-AFM analyses. In
principle, we classify that filament regions with supertwisted half helices and a reduced number
of protomer pairs per HHP, as detected in our AFM measurements, are indicative of C-actin-like
structure. We aimed to address several issues: (**i**) Does the number of actin protomers per HHP
vary between typical F-actin and C-actin-like structure regions? (**ii**) Does the MAD change, and
do these changes affect cofilin binding within the unbound actin regions adjacent to cofilin
clusters on the PE and barbed-end (BE) sides? Are these changes quantitatively similar or
distinct? (**iii**) Can we evaluate the impact of Pi on torsional flexibility during helical twisting of
actin filaments to elucidate how they influence the preferential cooperative binding and
expansion of cofilin clusters?

An innovative and previously unexplored facet of this study is the visualization of the
actin protomer's structure within a bare half helix adjacent to small cofilin clusters on the PE and
BE sides using HS-AFM. Our findings elucidate the dynamic structural changes in half helices
and the number of actin protomers per HHP in normal F-actin. Our results deviated from the
conventional ensemble-averaged helical twists usually observed in EM and X-ray diffraction,
depicting a rigid half helix. Moreover, we show that shortened bare half helices adjacent to the
cofilin clusters on the PE side contain less actin protomers than conventional half helices. The
MAD within the shortened bare HHP (C-actin-like structure) is longer than the MAD within the
canonical HHP (F-actin structure). This elongated MAD can be interpreted as resulting from the
allosteric transmission of structural changes originating in the cofilactin region, extending
beyond the boundary between cofilactin and bare actin regions. This extended influence affects
the structure of the shortened bare half helix on the PE side. Additionally, our findings suggest
that the cooperative binding of cofilin is significantly impacted by the torsional flexibility of
actin filaments. The inhibition of torsional motion during helical twisting, achieved via physical
perturbation, inhibited cofilin binding and the expansion of larger clusters more than Pi binding.

We also verify our previous hypothesis, which supports cofilin's initial binding to the
favorable structure of actin protomers because F-actin, without ADF/cofilin proteins, could exist
in the 162° twist state observed for cofilin-decorated F-actin (Galkin et al., 2011, 2001; Ngo et
al., 2016, 2015). Our hypothesis focuses on the evidence of decreasing the number of protomers
within the bare half helix adjacent to the cofilin cluster on the PE side. This reduction leads to a
shorter HHP but paradoxically results in a longer MAD in the C-actin-like structure area. The
protomers within a shortened bare half helix near cofilin clusters are converted to C-actin-like
structures via our proposed F-actin to C-actin transition pathway, thereby facilitating the
expansion of clusters on the PE side. If cofilin molecules fail to promptly attach to protomers
within the shortened bare region, the protomers resembling the C-actin structure in that region
can quickly change the closed nucleotide binding clefts to the open state and transition to a G-
actin state. This transition (C-actin-like to monomer G-actin-like structure) ultimately results in
severing of the filament at or near the boundary of cofilactin and bare actin regions. Furthermore,
this study suggests that Pi binding mainly inhibits the initial binding of cofilin by reducing the
fraction of supertwisted half helices in F-ADP.Pi-actin, whereas the removal of Pi enhances the
initial binding of cofilin by elevating the fraction of supertwisted half helices in F-ADP-actin.
Nevertheless, Pi binding only slightly inhibits the propagation of the supertwisted helical
structure within cofilin clusters to bare half helices on the PE side in F-ADP.Pi-actin when
compared to F-ADP-actin. Together, this clarifies that the initial interactions of cofilin with F-

actin and the subsequent expansion of clusters are closely linked to the variability in helical
twists.
141
Results
Principal component analysis suggests C-actin-like structure as the optimal actin structure
for cofilin binding
To investigate the relationship between actin structure and cofilin binding, we explored
the collective changes in actin conformations as they transitioned from G-actin to F-actin and
then to C-actin. We applied PCA to identify the principal components (PCs), as depicted in
Figure 1. Our analysis included an array of actin chains sourced from 41 available PDB
structures of filamentous actin protomers bound to cofilin, profilin, DNAse I, and various
nucleotides (ATP, AMPPNP, ADP.Pi, ADP, and Apo). We then compared these findings with 5
additional PDB structures featuring free G-actin and twinfilin-1-bound G-actin (**Table**
supplement 1).
We performed PCA with the same dataset using different alignments for subdomains 1
(SD1), 2 (SD2), 3 (SD3), and 4 (SD4), the outer domain (OD, SD1 + SD2), the inner domain
(ID, SD3 + SD4), and the entire actin structure. In all cases, we observed clustering of different
groups of actin chains. Interestingly, aligning only the IDs of all actin chains yielded the most
meaningful PCs. Importantly, PCA can elucidate the conformational transition between C-actin
and G-actin structures, and vice versa, through PC1, while PC2 delineates the pathway between
F-actin and C-actin structures (**Figure 1A**). To visualize the conformational changes associated
with each PC, we back-calculated and depicted the PCs in **Figure 1B and 1C**, respectively.
These illustrations revealed two outer domain motion modes. The first mode (PC1, accounting
for 45% of the structural variance) involved rotations of the OD around the hinge region to seal
the nucleotide binding cleft by reducing the SD2-SD4 distance (as indicated by the yellow
arrowhead in **Figure 1B and Video 1**), depicting the transformation between G-actin and C-
actin. Structures with higher PC1 values such as F-actin and C-actin are associated with
conformations featuring a closer nucleotide binding cleft. Conversely, structures with lower PC1
values, including tilted-actin and G-actin, are related to conformations featuring a more open
nucleotide binding cleft. The second prominent mode (PC2, accounting for 28% of the structural
variance) included the twisting of the OD during the transition between F-actin and C-actin
(**Figure 1C and Video 1**). A higher PC2 value indicates that the OD is more flattened or
untwisted, as seen in the tilted-actin and F-actin. On the other hand, a lower PC2 value suggests
that the OD is more twisted, as observed in G-actin and C-actin.
Our PCA results indicated that the structure of an actin filament with singly bound cofilin
– 6UBY (orange box in **Figure 1A**) was in the transition pathway between the F-actin and C-
actin conformations, particularly along PC2. This cluster also includes structures of two
protomers positioned at the BE side near a cofilactin cluster (including chains E&H combined as
E and F&G combined as F in PDB: 6UC4 (**Table supplement 1**)). Consistent with previous
studies (Galkin et al., 2011; Tanaka et al., 2018), our analysis suggests that the attachment of a
single cofilin to an actin filament induces a slight twisting of the OD relative to the ID (~5°).
Furthermore, this effect was cumulative, with more extensive cofilin binding leading to larger
OD twisting angles (~10°), thus representing the C-actin state. Throughout this transition, the
nucleotide binding cleft remains closed as observed through the minimal variations in PC1
values. This suggests that cofilin prefers to bind to the C-actin structure and the transitional actin
states featuring structures resembling C-actin within the actin filaments. However, we also

predicted that cofilins do not preferentially bind to the tilted structural state of actin, as tilted
actin structures exhibited opened nucleotide-binding clefts, but without the twisting of the OD
(Galkin et al., 2015) (indicated by lower PC1 and higher PC2 values, as represented by the
magenta box in **Figure 1A**). Additionally, these tilted actin structures did not align with the
conformational transition pathway between F-actin and C-actin.
Noticeably, PCA analysis revealed higher structural flexibility in F-ADP-actin (red dots),
exploring a larger space than F-ADP-Pi-actin structures (orange dots) within the F-actin cluster
(inset in **Figure 1A**). This included the F-ADP-actin structure (3MFP) (Fujii et al., 2010), which
notably exhibited a PC2 value significantly approaching that of C-actin structure (**Figure 1A**).
However, upon comparison of structures simultaneously resolved in (Chou and Pollard, 2019), a
slightly smaller PC2 value for the F-ADP-actin (6DJO) was observed in contrast to F-AMPPNP-
actin (6DJM) and F-ADP.Pi-actin (6DJN) (inset in **Figure 1A**). Significantly, the magnified
views in the F-actin cluster highlight structural resemblance between F-ADP-actin and F-
ADP.Pi-actin in recent studies (Oosterheert et al., 2022; Reynolds et al., 2022). The F-ADP-actin
structure (8D13) and F-ADP.Pi-actin structure (8D14) exhibit remarkably similar PC2 values
(**Figure supplement 1A**). Furthermore, the PC2 results demonstrate notable similarities between
Mg^{2+}-F-ADP-actin (8A2T) and Mg^{2+}-F-ADP.Pi-actin (8A2S), and Mg^{2+}-F-ADP.BeF$_3$-actin
(8A2R) (**Figure supplement 1B**). Altogether, we suggest that the conformation adopted by actin
in the cofilactin region, different from actin-nucleotide-solely dependent state, is the most
favorable one for binding cofilin.
In order to better understand the effect of the collective PC2 movements of individual
ODs on the global half helices of filaments, we compared the side views and outer views of F-
actin and C-actin protomers in their filamentous structures (**Figure 2A-C**). The twisting
movement of OD that we found in **Figure 1C** is best represented again from the side views with
the rotation axes (marked as red stars in blue boxes of **Figure 2A-B**). We identified that cofilin
clusters rotate the ODs approximately 10°, burying the SD2 including D-loops deeper into the
core of the filament. As a result, the twisting angles between two adjacent protomers in the same
strand appears to increase from approximately 26° to around 36° if viewed from the outer side
(black boxes of **Figure 2A-B**), as reported before (Galkin et al., 2011; Tanaka et al., 2018). Such
change in the twisting angles from the outer views can be represented as the change of the OD
lines on 2D surfaces of imaginary cylinders of the filaments, unwrapped along imaginary lines
that an AFM tip would scan through (**Figure 2D-E**). Such diagrams depict the varying twisting
angles of OD domains within F-actin and C-actin (PC2 mode), leading to the formation of HHPs
with different numbers of detected protomer pairs. Hence, collective movements of individual
ODs that decrease PC2 values would lead to a globally shorter HHPs and fewer number of
protomer pairs per HHP. Thus, this clearly demonstrates that the global shortened HHP is
directly related to the OD twists of the local actin protomers (**Figure 2B, C, E**), which represent
the C-actin structure.
223
The half helices within long actin filaments exhibit heterogeneity in their lengths and the
number of protomers per half helical pitch when observed through HS-AFM
In our actual AFM measurements, we determined the half helices, the number of
protomer pairs per HHP, and MAD as illustrated in the pseudo-AFM images representing F-
actin, C-actin, and cofilactin structures (**Figure supplement 2, Table supplement 2,** and
Methods). To investigate the presence of random variable helical twists in long actin filaments
in solution, we employed HS-AFM to examine the dynamic structures and distribution of actin

protomers within multiple half helices, with high spatial (1 nm in the xy plane and 0.1 nm in the
z-axis) and temporal (2 frames per second) resolution. As depicted in **Figures 3 and 4**, **Table**
supplement 3, and **Video 2**, we successfully imaged and quantified the number of actin
protomers within many half helices, allowing us to precisely measure the distribution of the
number of protomer pairs per HHP (**Figure 3A**). Subsequently, we established a correlation
between the HHP and the number of protomer pairs per HHP. We also calculated the HHPs
assuming a constant MAD of 5.5 nm and compared them with actual data (**Figure 3B**). We
analyzed the linear regression analysis results and noted that the slope derived from our actual
AFM measurements was smaller than that computed assuming that the MAD remained
unchanged.
To address this discrepancy, we analyzed our measurements, revealing a range of
protomer pairs within each HHP, varying from 4.5 to 8.5 with distinct proportions (**Figure 3C,**
Table supplement 3). The most commonly observed numbers of protomer pairs per HHP were
6.5 and 7.5, accounting for 60.9% (n = 1355 HHPs) and 28.0% (n = 624 HHPs) of the samples,
respectively. Conversely, the least frequently observed numbers of protomer pairs per HHP were
4.5, 5.5 and 8.5, accounting for 0.2% (n = 5 HHPs), 8.2% (n = 182 HHPs) and 2.7% (n = 60
HHPs) of the samples, respectively. Subsequently, we computed the MAD by dividing the length
of each measured HHP by the corresponding number of protomer pairs per HHP, thereby
establishing a correlation between these variables (**Figure 3C**). The MAD values (mean ± SD)
for HHPs containing 6.5 and 7.5 were 5.1 ± 0.2 nm (n = 8807.5 pairs) and 5.0 ± 0.3 nm (n =
4680 pairs), respectively. Corresponding values for HHPs with 5.5 and 8.5 protomer pairs were
5.3 ± 0.3 nm (n = 1001 pairs) and 4.6 ± 0.3 nm (n = 510 pairs) (**Table supplement 3**). Notably,
we observed that HHPs with fewer protomer pairs exhibited elongated MADs. Our analysis also
indicates an extremely uncommon occurrence, with approximately 0.2% of the actin protomers
within the F-actin structure (4.5 protomer pairs per HPP) demonstrating elongated MADs (6.0 -
6.6 nm).
We next conducted a temporal autocorrelation function (ACF) analysis aimed at
estimating the time constants governing the decaying self-correlation in the axial distance (AD)
between two adjacent actin protomers along the same long-pitch strand (**Figure 3D-F, Figure**
supplement 3, and **Video 3**). The core objective of this analysis was to determine the temporal
extent of AD similarity and elucidate the dynamic behaviors and conformational changes in the
distance between adjacent actin protomers over time. The results revealed that self-correlation of
the ADs within actin filaments, immobilized on lipid membrane, rapidly decreased within a few
hundred milliseconds and then gradually decayed within a few seconds (**Figure 3E-F, Figure**
supplement 3). The t_1 and t_2 values correspond to the first and second time constant decays of
AD between two adjacent actin protomers within F-ADP-actin and F-ADP.Pi-actin segments. At
the current imaging rate, these time constants remain very similar between actin filaments bound
with ADP or ADP.Pi (**Figure supplement 3, Table supplement 5**). These results potentially
explain the intrinsic dynamics and fluctuations observed in HHPs, the number of protomer pairs
per HHP, and the MADs within the actin filaments over time.
To assess the degree of spatial clustering in the structure of actin protomers, we utilized a
spatial ACF analysis. This analysis provides an indirect measure of the degree to which height
values near each other in space are similar to one another in actin filaments. If protomers can
change the shape and/or rotate around the axis parallel to the filament axis, the individual
protomers may have slightly different thickness and/or orientations, leading to variations in the
heights of protomers and, consequently, variations in the slopes in heights between protomers

(depicted in **Figure supplement 4**, **Methods**). As seen in **Figure 4C and Figure supplement**
5A, the spatial ACF curves were effectively modelled using an exponential decaying sinusoidal
function. The decay constant ($D1$) values (mean ± SD) were determined to be 27.2 ± 6.7 nm.
This result provided an indirect measure of the degree to which height values near each other in
space are similar to one another in canonical actin filaments. Because our AFM images provide
detailed surface topography along a longitudinal section profile with outstanding spatial
resolution in height (z-axis: ~1 Å), we suggest that the similarity in height among protomers in
F-actin persists within a region encompassing approximately 5 pairs (~10 actin protomers).
However, the limited lateral resolution (x- and y-axis) in our AFM images, restricted to 1–3 nm,
prevented us from delving further into the structural details of actin protomers within this
distance decay.
288
The shortened bare HHP adjacent to the cofilin clusters on the PE side includes fewer actin
protomers than the normal HHP, resulting in MAD elongation
Using HS-AFM, we aimed to simultaneously capture the helical structures and
distribution of actin protomers in the bare F-actin, cofilactin, and hybrid cofilactin/bare actin
segments within the same imaging area (see Methods). As shown in **Figure supplement 6**, we
successfully distinguished cofilactin and bare F-actin structures based on a height difference of
approximately 1-1.6 nm and a length difference in the half helices of approximately 25%. A bare
F-actin segment consisting of two consecutive half helices (HHP1, HHP2) exhibited the
expected normal lengths of 36 nm (6.5 pairs) and 36.5 nm (6.5 pairs) (**Figure supplement 6A**).
In contrast, a cofilactin segment with two consecutive half helices (HHP3, HHP4) displayed
shorter lengths of 28.6 nm (4.5 pairs) and 29.8 nm (4.5 pairs) (**Figure supplement 6B**). Next, we
presented a typical segment of the hybrid bare actin/cofilactin structure containing two
consecutive half helices (HHP5, HHP6), both showing comparably shortened lengths of 27.9 nm
(4.5 pairs) and 30.5 nm (4.5 pairs) (**Figure supplement 6C**). Based on our previous study (Ngo
et al., 2015), we judged that the bare half helix with the shorter HHP adjacent to the cofilin
cluster is on the PE side and that the normal HHP is located on the BE side. These typical data
suggest that, similar to cofilactin segments, shortened bare half helices adjacent to cofilin
clusters on the PE side include fewer actin protomers than normal half helices. We next analyzed
the MAD variation in this hybrid filament over time and made interesting observations (**Figure**
supplement 7). We discovered that MAD5 in bare actin HHP5 was elongated similarly to that in
cofilactin HHP6 and HHP7. Notably, MAD5 (5.7 ± 0.5 nm, n = 13) was extended and closely
resembled MAD6 (5.9 ± 0.3 nm) and MAD7 (6.1 ± 0.3 nm). This finding suggests that the MAD
elongation in the cofilactin region was transmitted beyond the cofilactin/bare actin boundary to
extend MAD5 in the shortened bare half helix.
We then conducted similar analyses on the hybrid cofilactin/bare actin filaments (**Figures**
5–6, **Video 4**). First, we observed the minimal impact of the lipid membrane on the half helices
(HHP1, HHP2), number of protomer pairs per HHP, and MADs (MAD1, MAD2) in normal actin
filaments (**Figure 5A**). We examined the bare half helices (HHP3–HHP6) and the number of
protomer pairs within each HHP on either side of a small cofilin cluster and found that the
shortened bare half helix on the PE side includes fewer actin protomers than the normal half
helix (**Figure 5B, 5C**). For example, the cofilin cluster (estimated as 2 to 4 molecules) shortened
one adjacent bare half helix (HHP6) on the PE side (25.5 nm, 4.5 pairs), while the half helix
(HHP5) on the BE side was similar to the normal half helix (37.9 nm, 7.5 pairs) (**Figure 5C**). In
this case, the MAD6 calculated within the shortened bare half helix (HHP6) was 5.7 nm and was

larger than the MAD5 value of 5.1 nm calculated within the normal half helix (HHP5) (**Figure**
5C). Despite a complete decoration requiring 9 to 11 cofilin molecules in a half helix, the
observation that 2 to 4 molecules can initiate the supertwisting of a half helix, achieved by
reducing the number of protomer pairs per HHP, strongly implies the presence of allosteric
transitioning in the supertwisting structures from the bound to the unbound region on the PE
side. We anticipate that the unbound region may contain approximately 1 to 2 pairs (2 to 4
protomers) resembling C-actin structures. When analyzing the fluctuation in the MADs over
time in this hybrid filament, we observed that MAD6 (5.4 ± 0.3 nm, n = 12) was significantly
greater than MAD5 (5.0 ± 0.2 nm, n = 12) (**Figure 6A, 6B**). Our analysis suggests that the
MADs are longer in the shorter bare half helices than in the normal half helices (**Table**
supplement 4). Furthermore, we analyzed the fluctuation of the MAD in the F-ADP-actin
segments on both sides of large cofilin clusters over time (**Figure supplement 8, Video 5**). Our
analysis suggests that the MADs fluctuate over time and skew towards longer ranges in a
supertwisted bare half helix on the PE side compared to an untwisted or normal half helix on the
BE side.
We also analyzed a spatial ACF to indirectly measure of the degree to which height
values near each other in space are similar to one another, specifically focusing on regions from
bound to unbound cofilin. In contrast to the normal actin filament (**Figure 4C, Figure**
supplement 5A), the presence of small cofilin clusters led to a more pronounced linear decline
in the spatial ACF curves for each half helix (**Figure 6C, Figure supplement 5B**). The
distribution of the decay constant (*D1*) values was analyzed, yielding the mean $\pm$ SD values for
each half helix on the PE and BE side of 12.7 ± 7.0 nm and 12.6 ± 6.6 nm, respectively. We
propose that the similarity in height among protomers in the bound cofilin region equally extends
to the unbound region on the PE and BE sides, encompassing approximately 1 to 2 pairs (~2 to 4
actin protomers). Nevertheless, the restricted lateral resolution in our AFM images hindered a
detailed exploration of the structural intricacies of unbound actin protomers on the PE or BE
side. We propose the future utilization of Cryo-EM to examine the structural details of bare
protomers adjacent to a cofilin cluster within this hybrid cofilactin/bare actin filament, which is
attached similarly to a supported lipid membrane.
352
The transmission of the supertwisted helical structure within the cofilin cluster to the
adjacent bare half helices is essential in facilitating the preferential cooperative binding of
cofilin
The cooperative binding of cofilin to actin filaments has been previously reported
(Bobkov et al., 2006; McGough et al., 1997; Ngo et al., 2016, 2015). However, the role of
cooperative conformational changes in actin filaments in driving the preferential binding of
cofilin toward the PE side remains unclear, as indicated in several notable studies (Bibeau et al.,
2021; Bobkov et al., 2006; Galkin et al., 2011; Hayakawa et al., 2014; Huehn et al., 2018, 2020;
McCullough et al., 2008; Narita, 2020; Ngo et al., 2015; Suarez et al., 2011; Tanaka et al., 2018;
Wioland et al., 2017).
We observed the cooperative binding of cofilin to actin filaments when they were loosely
absorbed onto a lipid membrane in an F-buffer with cofilin, ATP, and inorganic phosphate (Pi)
(**Figure 7A, Video 6**). We also examined the peak heights and lengths of the half helices in the
cofilactin and bare actin regions during the unidirectional growth of the cofilin clusters (**Figure**
7A-C). Consistent with our previous findings (Ngo et al., 2015), the cofilin cluster binding led to
an approximately 1–2 nm increase in the peak height and an approximately 25% decrease in the

HHPs in the cofilactin region. Then, we analyzed the relationship between the time needed to
shorten the bare half helix beyond the cofilactin/bare actin boundary and to form a newly
matured cofilin cluster within the shortened bare half helix. Finally, we determined the lag time
for the growth of the cofilin cluster within the individual shortened bare half helices adjacent to
the cofilin cluster.
The analysis suggested that even with 10 mM Pi, the cofilin cluster could supertwist the
half helices in the actin filaments. Notably, the supertwisted half helices within the cofilactin
region propagated beyond the cofilactin/bare actin boundary, shortening the adjacent bare half
helix before the saturated cofilin cluster grew (**Figure 7A, 7B**, **Table supplement 6**). For
example, when the HHP between P2 and P3 was shortened at 2 seconds, P2 was bound to a
cofilin cluster, but P3 remained unbound. A lag time of approximately $t = 3$ s was needed for the
saturated cofilin cluster to be recruited and grew in P3 at 5 seconds. This pattern was also
observed when the HHPs between P3-P4, P4-P5, and P5-P6 shortened before the cofilin clusters
expanded. These findings suggest that the shortening of the bare half helix beyond the
cofilactin/bare actin boundary drives the actin structure-dependent preferential cooperative
binding of cofilin clusters. The cooperative conformational changes during helical twisting may
provide a mechanism for understanding the preferential binding of cofilin clusters toward the PE
side, as reported in previous studies (Bibeau et al., 2021; Ngo et al., 2015).
As negative control experiments, we immobilized actin filaments onto lipid membranes
(DPPC/DPTAP=75/25 wt%) to physically hinder the changes with helical twisting. We then
added a high concentration of cofilin (2000 nM) and observed its cooperative binding with actin
filaments in the presence of ATP (**Figure 8A, Video 7**). However, the binding and expansion of
cofilin clusters on these actin filaments were severely inhibited during long incubation times.
Instead of cooperatively forming large clusters, cofilin formed only a few small clusters that
transiently associated and dissociated, even at cofilin concentrations much higher than k_d ($k_d =$
~5–10 nM) (Andrianantoandro and Pollard, 2006). The histograms of the peak heights and HHPs
were analyzed and showed single peaks at 8.2 ± 0.8 nm (n = 1930) and 34.3 ± 2.6 nm (n = 1556),
respectively, which were similar to those of normal actin filaments (**Table supplement 7**). The
decay constant ($D1$) values were determined to be 26.8 ± 6.3 nm (**Figure supplement 9**),
suggesting that the resemblance in height among protomers in sparsely decorated cofilin-actin
filaments extends within a region covering approximately 5 pairs (~10 actin protomers). These
values were found to be comparable to those (27.2 ± 6.7 nm) observed in normal actin filaments
(**Figure 4C**). Together, our result strongly suggests that the cooperative binding of cofilin is
greatly influenced by cooperative conformational changes during helical twisting.
403
Examining the inhibitory effect of Pi on the binding and expansion of cofilin clusters
Next, we examined the nucleotide effect on the cooperative binding of cofilin to actin
filaments (**Figure 8B-D, Video 8-10**). We analyzed the time-dependent changes in the peak
heights and HHPs of representative actin filaments under each condition to track the growth of
the cofilin clusters. We made actin filaments incubated with different nucleotides (ADP and ATP
+ Pi), leading to the formation of F-ADP-actin and F-ADP.Pi-actin, and subsequently mixed
with cofilin. We consistently observed the preferential binding of cofilin to shortened bare HHPs
toward the PE side. However, we noted that in rare instances, cofilin clusters also grew on both
sides in the regular bare half helices when only ADP was present. The histograms of the peak
heights and HHPs followed Gaussian distributions corresponding to the cofilactin and normal
bare actin regions (**Figure 8B-C, Table supplement 7**). The growth rates of the cofilin clusters

within the shortened bare half helices adjacent to the cofilin clusters in both F-ADP.Pi-actin and F-ADP-actin were not significantly different, as shown in **Figure 9A-B**. Currently, we have no experimental data to discuss about whether and when bound Pi is released during this process. However, three major processes could be considered once a cofilin cluster with a critical size has formed, the allosteric transmission of the supertwisted structure of the half helices within the cofilactin region to an adjacent bare half helix on the PE side results in C-actin-like protomers. First, Pi is released accompanying (preceding) supertwisting structural change in the bare region near cofilin clusters on the PE side. Because most structural studies suggest that cofilin binding promotes Pi release, leading to the formation of C-actin with ADP binding (**Figure 1A**), we anticipate that protomers within the shortened bare half helix adjacent to cofilin clusters in F-ADP.Pi-actin rapidly converts to F-ADP-actin and C-actin-like structures, thereby facilitating the expansion of cofilin clusters toward the states of actin bound to ADP. Second, Pi remains bound to protomers within the supertwisted bare section. The successive interactions with cofilin molecules promote Pi release. Third, similar to the process described in the second one, but in this case, subsequent interactions with cofilin molecules do not induce the release of Pi. Consequently, numerous actin protomers within the newly formed cofilactin region remain bound to ADP.Pi. When we observed neighboring bare half helices on both sides of a cofilin cluster in F-ADP-actin and F-ADP-Pi-actin within one imaging field, the HHP was always shorter on one side, while the HHP was essentially normal on the other side in both F-ADP-actin and F-ADP.Pi-actin. These results suggest that bound Pi has a minor effect, hindering the propagation of the supertwisted helical structure within cofilin clusters to the neighboring bare half helices on the PE side.

To examine how bound Pi inhibits cofilin binding (Blanchoin and Pollard, 1999; Carlier et al., 1997; Carlier and Pantaloni, 1988), we assessed the torsional flexibility by measuring the HHPs in control actin filaments incubated with ADP and ATP+Pi, or ATP + phalloidin (**Figure 9C-D**). In all cases, the histograms followed typical HHP distributions. The mean HHPs were nearly identical. However, the variances of the HHPs in F-ADP.Pi-actin and phalloidin-stabilized F-actin were significantly smaller than that in F-ADP-actin. Consequently, the proportion of naturally supertwisted half helices with HHPs shorter than 30 nm was 5.8% for F-ADP-actin but only 1.1% and 0.2% for F-ADP.Pi-actin and phalloidin-stabilized F-actin, respectively. Our finding provides the structural support for earlier research that the nucleotide state of actin could influence F-actin's flexibility and its ability to undergo rearrangements that mediate ABP binding (Blanchoin and Pollard, 1999; Okura et al., 2023; Suarez et al., 2011; Zimmermann et al., 2015). This is in line with measurements of micrometer-scale persistence length, indicating that F-ADP.Pi-actin is more rigid than F-ADP-actin (Isambert et al., 1995), and with the mechanical control of twisting and bending of actin's nucleotide-bound states (De La Cruz et al., 2010). Assuming that initial cofilin binding occurs at sites with naturally supertwisted helical structures (Galkin et al., 2001), we suggest that bound Pi in F-ADP.Pi-actin mainly inhibits initial cofilin binding by reducing the fraction of supertwisted helical structures in comparison to F-ADP-actin.

Discussion

We investigated the dynamic conformational changes in half helices of F-actin and C-actin-like structure segments by observing the fluctuations in the number of actin protomers per HHP and MAD. The results provide robust support for a prior study (Egelman et al., 1982), which reported variable helical twisting within actin filaments. Nonetheless, the Cryo-EM and X-ray diffraction findings depict the rigid and inflexible helical twists of actin filaments (HHP =

36 nm). Certainly, HS-AFM captures the dynamic structural changes of proteins during
functional processes in solution, offering a trade-off with lower spatial resolution but superior
temporal resolution compared to Cryo-EM (Ando, 2012; Ando et al., 2023, 2008). We leverage
the advantage of this technique to capture the torsional flexibility during helical twisting of actin
filaments and elucidate their adaptable structure as a variable helical twist, which plays crucial
roles in the binding and cluster expansion of cofilin. While it was previously believed that the
interaction between the upper and lower clefts establishes the structural basis for how nucleotide-
dependent conformation changes in F-actin structure modulate the binding affinities for ABPs
including cofilin (Dominguez and Holmes, 2011), recent structural analyses have revealed a
remarkable similarity in the average structure of actin protomers bound with $ADP.BeF_3$, ADP.Pi,
and ADP in the presence of Mg^{2+} (Oosterheert et al., 2022; Reynolds et al., 2022) (**Figure
supplement 1**), significantly challenging the hypothesis regarding the modulation of ABP
binding affinities through nucleotide-dependent conformational changes. So, why is the
flexibility in torsion during helical twisting crucial for the preferential cooperative binding of
cofilin to actin filaments toward the PE side? Unfortunately, we currently lack the atomic
structures of actin protomers adjacent to cofilin clusters on the PE side to fully address this
question.
Nonetheless, our PCA results reveal that key longitudinal interactions between the D-
loop of SD2 positioned in the pocket of the next protomer's SD1 contribute to keeping the
nucleotide binding cleft closed and the OD untwisted in the F-actin (**Figures 1, 2**). When cofilin
binds and sits on the sides of SD1 and SD2 of two adjacent actin protomers (**Figure 2B, C**), the
OD (SD1, SD2) in bound protomers becomes more twisted (a decrease in PC2), but without
significantly affecting the closing of the nucleotide binding cleft (nearly unchanging PC1). This
transformation ultimately leads to the conformational changes from F-actin to C-actin (**Figure
1**). The transition from the C-actin to breaking the filament into monomer G-actin structure
additionally requires the opening of the nucleotide binding cleft (decreasing PC1). However, this
process is normally hindered in the presence of cofilin, as the bridging between two protomers
can compensate for all the losses in the disrupted interactions between the SD2 D-loop and SD1
pocket (**Figure 2B, C**). Strikingly, the HS-AFM results suggest that cofilin clusters can convert
bare protomers within the shortened bare half helices on the PE side into C-actin-like structures.
Unless additional cofilin molecules quickly bind to the protomers with C-actin-like structures on
the PE side, the disrupted interactions would readily lead to the nucleotide binding cleft
transitioning from the closed to open states, resembling C-actin to G-actin transition (**Figure 1**).
Our proposed hypothesis regarding the shift from C-actin-like structure to G-actin aligns with
earlier studies suggesting that the binding sites of cofilin encompass a region that overlaps with
G-actin structure (Carlier et al., 1997; Mannherz et al., 2007; Narita, 2020; Tanaka et al., 2018).
The dissociation of the D-loop from an adjacent actin subunit directly causes the transition to the
G-form of actin, which is regarded as the most stable configuration for the actin molecule (Oda
et al., 2019). Additionally, the protomers within the shortened bare half helix near a cofilin
cluster on the PE side of F-ADP.Pi-actin may be quickly converted to F-ADP-actin and C-actin-
like structure through F-actin to C-actin transition. Together, this explains the preferential
binding of cofilin to ADP-actin on the PE side and why filament breakages are more commonly
observed at or near the boundary of cofilactin and bare actin on the PE side, rather than where
actin protomers are fully bound with cofilin (Bibeau et al., 2021; Ngo et al., 2015; Suarez et al.,
2011). However, our finding regarding the structure partially contradicts the existing Cryo-EM

structure, which suggested that conformational changes induced by cofilin clusters, shifting from
F-actin to C-actin, are restricted to the protomers directly in contact with bound cofilin.
According to this model, cofilin tunes adjacent protomers, resembling F-ADP-actin structure on
the PE side, to expand the clusters (Huehn et al., 2020; Suarez et al., 2011). Also, cofilin binding
may be facilitated by a shortening of the helix pitch, independent of any transition to the C-form
of actin (Narita, 2020). Nonetheless, their hypotheses are insufficient to explain the PCA results
supporting the structural evidence obtained from our AFM measurements that a cofilin cluster
not only has the capacity to convert F-actin to C-actin but also allosterically expands the C-actin
structure from the bound to unbound region on the PE side. Thus, additional cofilin molecules
preferentially bind to actin protomers resembling C-actin within the shortened bare zone than F-
ADP-actin structure near a cofilin cluster on the PE side (**Figures 1, 5-9**). Even though HS-AFM
did not allow for the direct observation of the different nucleotide states of actin, these
unidirectional propagation of the supertwisted structure in cofilin clusters to the neighboring bare
zone on the PE side and preferential and cooperative bindings of cofilin to those bare sections
were consistently and unequivocally observed in the presence of ADP and ADP.Pi (Ngo et al.,
2015). Further studies are needed to identify the source of this discrepancy between HS-AFM
and Cryo-EM observations.
Our aim is to understand the underlying mechanism driving the preferential and
cooperative attachment of cofilin to actin filament. This mechanism is largely influenced by the
preferred helical twisting structure within the actin filament, as opposed to an alternate theory
that suggests cofilin binding depends on the nucleotide state of actin (**Figure 10**). While both
hypotheses might explain biased cofilin binding along the actin filament, they propose different
mechanisms underlying cofilin's cooperative behavior and binding preference.
Our hypothesis (**Figure 10E-H**) aligns with the results in the cofilactin region, where
actin protomers are fully decorated by cofilin (**Figure 10A, 10C, 10D**). However, it is partially
incongruent with the structures of actin protomers near cofilin clusters on the PE side, as
depicted in **Figure 10D**. Additionally, we propose two potential hypotheses that require further
verification in the near future to explain the preferential cooperative bindings and cluster
expansion of cofilin to undecorated C-actin-like structure within the shortened bare half helix on
the PE side (**Figure 10E-F**): (i) Because of the disrupted connections between the BE-
protomer's SD2 D-loop and PE-protomer's SD1 pocket in the area resembling the C-actin
structure, where cofilins are not bound, the SD2 could change between untwisted and twisted
states (F-actin to C-actin transition) and the nucleotide-binding cleft could shift between the
closed and open states (C-actin to G-actin transition). These transitions might prompt the rapid
transformation of protomers within the shortened bare half helix adjacent to the cofilin clusters
on the PE side of F-ADP.Pi-actin to F-ADP-actin and C-actin-like structure. This phenomenon
also explains the binding preference of cofilin for ADP-actin on the PE side; (ii) In the canonical
actin protofilament, the SD2 D-loop of BE-protomer is inserted between the tips of SD1 and SD3
of PE-protomer (**Figure 2A**). However, the SD2 D-loop of BE protomer does not directly
interact with the tips of SD1 and SD3. Instead, it is deeply buried within the core of the cofilactin
protofilament, representing the C-actin structure (**Figure 2B-C**). Conversely, in the region
representing the C-actin-like structure, where cofilins are unbound, the SD2 D-loops in twisted
OD might exhibit various dynamic conformations, potentially allowing flexible interaction with
the tips of SD1 and SD3 including that of the neighboring protomers on the PE side. The
collective movements of individual ODs that decrease PC2 values would lead to a shorter HHP
and fewer number of protomer pairs per HHP (**Figure 2**). These changes cause an extended

MAD in C-actin-like structure area with optimal conformational changes in lower cleft between
SD1 and SD3 of the protomers on the PE side, thereby facilitating the preferential binding and
expansion of cofilin clusters (**Figure 10E-F**).
In this study, we experimentally demonstrated that helical twisting, the number of
protomers per half helix, and MADs are intrinsically dynamic and fluctuate around their mean
values over time. Our findings support the variability (Egelman et al., 1982) and irregularity
(Fineberg et al., 2024) in helical twists and dynamics of actin filaments both *in vitro* and *in vivo*
(Ivanova et al., 2024), challenging the traditional view, based on Cryo-EM and X-ray diffraction,
that these parameters remain unchanged. Furthermore, we observed a significant correlation
between the nucleotide-bound states of actin and the proportion of naturally supertwisted half
helices in F-ADP-actin and F-ADP.Pi-actin (**Figure 9**). We suggest that the bound Pi reduces the
fraction of naturally supertwisted half helices and hinders the initial binding of cofilin and/or
initial growth to a critical size of cluster set at 2 to 4 molecules (**Figure 5-6**) that can induce
allosteric conformational changes to the neighboring bare zone on the PE side for cooperative
bindings of cofilin (**Figures 9C-D, 10H**). This discovery aligns with previous research, which
showed that achieving the most effective cooperative binding of cofilin also requires a critical
size for a cofilin cluster set at 2 molecules (Huehn et al., 2020). However, Pi only minimally
impedes the propagation of the supertwisted helical structure within the cofilin cluster to the bare
actin zone in F-ADP.Pi-actin compared to F-ADP-actin. This transition is pivotal in driving the
preferential cooperative binding and expansion of the cofilin cluster (**Figures 7, 8**). In terms of
structure, the PCA results also strongly agree with previous research indicating that ADP.BeF$_3$,
ADP.Pi, and ADP F-actin structures (**Figure supplement 1**) display a striking resemblance
(Oosterheert et al., 2022; Reynolds et al., 2022). Additionally, the reduction in torsional
flexibility during helical twisting resulted in a significant hindrance to the cooperative binding
and expansion of cofilin clusters (**Figure 8A**). Therefore, we suggest that the reduction in
torsional flexibility associated with variable helical twists, rather than the large conformational
change in F-actin's protomers linked to ATP hydrolysis, accounts for the inhibition of cofilin
binding by Pi (**Figure 10H**). Nonetheless, it remains plausible that the structural flexibility
exhibited by ADP-bound actin protomers could result in subtle variations in the conformations of
the DNase binding loop (D-loop) G46-M47-G48-N49, as suggested in (Chou and Pollard, 2019).
The crystal structure of the F-form revealed that Pi in ADP.Pi connects the two large domains of
the actin molecule, thereby stabilizing the F-form. When Pi is released, this connection is
significantly weakened (Kanematsu et al., 2022). We suggest that the absence of bound Pi
possibly increases the torsional flexibilities during helical twisting of ADP-bound actin filaments
in contrast to their ADP.Pi-bound counterparts.
Actin structure-dependent preferential binding of actin-binding proteins (ABPs) have
been investigated at the molecular and cellular levels (Hayakawa et al., 2022; Jegou and Romet-
Lemonne, 2019; Jégou and Romet-Lemonne, 2016; Ngo et al., 2016; Tokuraku et al., 2020,
2009). Notably, Harris and colleagues reported biased localization of tandem calponin homology
domains (CH1-CH2) that can be mutated to selectively bind different actin conformations at the
leading or trailing edge of motile cells (Harris et al., 2020). The untwisting of the long-pitch
helix of the actin filament induced by diaphanous-related formin mDia1 inhibits cofilin binding
(Mizuno et al., 2018). Stretching actin filaments within cells enhances their binding affinities for
the myosin II motor domain (Uyeda et al., 2011). The importance of dynamic, cooperative, and
allosteric conformational changes in actin filaments warrants further investigation of the actin
structure-dependent preferential cooperative binding and function of other ABPs to clarify the

role of dynamics, structural polymorphisms, and physiological implications of actin filaments in
working with different ABPs in cells.
600
Methods
Proteins
Rabbit skeletal actin and human cofilin 1, without a His-tag, were prepared following the
same protocol as described in our previous report (Ngo et al., 2015). The purified proteins were
rapidly frozen in liquid nitrogen and stored at -80°C until use. Prior to experimentation, the
proteins were thawed on ice and subjected to ultracentrifugation (typically at 80,000 rpm for 10
minutes at 4°C) to eliminate any undesired proteins present in the pellet. The remaining high-
quality proteins in the supernatant were collected, kept on ice, and utilized for HS-AFM
experiments.
610
Principal component analysis
Actin molecules have been categorized into four groups based on the orientations of the
core of the outer domain (OD): F-form in naked F-actin, C-form in cofilactin, O-form in profilin-
actin, and G-form in presumably free actin, as determined by Hierarchical Cluster Analysis
(HCA) (Oda et al., 2019). However, HCA serves only for clustering and lacks the capability to
clearly visualize or map back to the original structural data to derive further meaningful insights.
In contrast, Principal Component Analysis (PCA) is typically used to reduce the dimensionality
of large datasets, retaining essential information. Through this dimension reduction, data clusters
naturally form, and critical information can be mapped back to the original data, revealing
collective movements between clusters. PCA has previously been applied to analyze actin
structures (Xue et al., 2014) using a small set of PDBs.
The process involves aligning the original data in three-dimensional space to remove
trivial movements, calculating the displacement covariance matrix between common atoms, and
performing eigendecomposition. This results in a small number of new variables called principal
components (PCs), which capture most of the original data's information. Clustering is a
byproduct of PCA, as the original dataset can be projected onto the identified collective
components and naturally grouped in the new PC dimensional space.
To ensure consistent and accurate PCA results, we conducted several preprocessing steps.
After cleaning the selected PDBs, we specifically extracted the alpha carbons of all reported
protein chains for further analysis. The sequences of these chains were aligned using the AlignIO
module from the Biopython package (Cock et al., 2009), and we identified the residues that were
commonly present across the PDBs. Specifically, we found that residues 8 to 41, 54 to 63, 66 to
74, 76 to 234, and 236 to 373 in the ACTs_RABIT gene (P68135) were common among the 46
chosen PDBs. Importantly, these residue numbers differ from the standard residue IDs used in
PDBs because the initial two residues present in genes are typically cleaved in the final actin
chains.
We then performed structural alignment only for residues in the IDs, following the
definition reported in (Kabsch et al., 1990). After aligning the structures, we used the entire set
of commonly solved residues to establish the covariance matrix, which was subjected to
eigendecomposition to obtain different eigenvectors (PCs). As shown in **Figure 1 and Figure**
supplement 1, we utilized the two eigenvectors with the largest values, denoted as PC1 and PC2.
In this study, the significance of PCA lies in its capacity to unveil the conformational transition

between G-actin and C-actin, and vice versa, through PC1, while PC2 delineates the pathway
between F-actin and C-actin, and vice versa.
645
Analyses of pseudo AFM images of F-actin and C-actin structures constructed from
existing PDB structures
To illustrate how cofilin clusters impact the helical twisting of long actin filaments,
leading to the shortening of HHPs and less protomers per HHP, resembling the C-actin structure
as measured by AFM, we built long actin filaments at different states along the F-actin to C-actin
transitions by using 8 PDB chains (**Table supplement 2**). In **Figure supplement 2**, we show
pseudo-AFM images featuring long filaments representing F-actin (constructed from 6VAU), C-
actin (with cofilins excluded) and cofilactin (constructed from 6VAO) structures.
To construct the long filament featuring F-actin, we performed a series of steps. First, we
repeatedly rotated and translated two actin chains from the selected PDBs to generate the
coordinates of 62 consecutive protomers. Since not all of the selected PDBs had reported
BIOMT matrices, we used transformation matrices that minimized the root mean square
deviation (RMSD) between the two chains. To construct long filaments featuring C-actin and
cofilactin, a similar process was followed, but it involved two chains of actin protomers and two
chains of cofilins. The lists of chains and matrices utilized to build the long filaments are
presented in **Table supplement 2**. Zoom-in views of these long filaments are shown in **Figure 2**.
The generated long filaments were then aligned along the x-axis and pseudo-imaged
using Afmize software (https://github.com/ToruNiina/afmize) (Niina et al., 2021, 2020). A probe
radius of 2.0 nm with an angle of 5.0° was set, and the image resolution was configured to be 1.0
nm in the xy plane and 0.64 Å on the z-axis, with a noise level of 0.1 nm. To analyze the
structures, we calculated longitudinal section profiles of the HHPs by determining the average
height within ± 3 nm around the center line (red and blue lines in **Figure supplement 2A**). The
offset values of the height profile were set to zero by referring to virtual mica surface, where
filament was placed in the pseudo-AFM images.
In our AFM analyses, we determined the HHPs by measuring the distances between
major peaks observed in longitudinal section profiles along the filament axis (red markers in
Figure supplement 2A-B, 2D-E, and 2G-H). These major peaks likely represent two distinct
actin protofilaments that are successively probed by the AFM tips (red markers in **Figure**
supplement 2C, 2F, and 2I). Thus, the number of actin protomers within each HHP is odd,
leading to a noninteger number of actin pairs. For example, in canonical actin filaments, each
HHP is typically composed of 13 actin protomers (equivalent to 6.5 protomer pairs). To account
for this, we determined the number of protomer pairs within each HHP by counting the major
and minor peaks (noted by the red and green markers within each HHP subtracted by 0.5) in the
same longitudinal section profiles. This adjustment was applied for both the pseudo and
experimental AFM images of the filaments representing F-actin and C-actin. Our pseudo-AFM
image displayed the challenge of accurately detecting actin pairs within filaments representing
the cofilactin structure (**Figure supplement 2G-I**) due to the presence of cofilin decorations and
the tip-sample dilation effect (Ando, 2022).
684
High-speed atomic force microscopy
We used a laboratory-built high-speed atomic force microscope (HS-AFM), as described
previously (Ando et al., 2013). HS-AFM imaging in the tapping mode was carried out in solution
with small cantilevers (BL-AC10DS-A2, Olympus), and the spring constant, resonant frequency

in water, and quality factor in water were ~0.1 N/m, ~500 kHz, and ~1.5, respectively. An
additional tip was grown on the original cantilever tip in gas with sublimable ferrocene powder
by electron beam deposition (EBD) using scanning electron microscopy (ZEISS Supra 40
VP/Gemini column, Zeiss, Jena, Germany). Typically, the EBD tip was grown under vacuum (1-
5×10^{-6} Torr), with an aperture size of 10 μm and an electron beam voltage of 20 kV for 30 s.
The EBD ferrocene tip was sharpened using a radio frequency plasma etcher (Tergeo Plasma
Cleaner, Pie Scientific, Union City, CA) under an argon gas atmosphere (typically at 180 mTorr
and 20 W for 90 s). For HS-AFM imaging, the free-oscillation peak-to-peak amplitude of the
cantilever (A_0) was set at ~1.6-1.8 nm, and the feedback amplitude set-point was set at ~0.9 A_0.
HS-AFM imaging was performed as described in detail elsewhere (Ando et al., 2013; Kodera et
al., 2021; Ngo et al., 2015), except for the use of a recently developed only tracing imaging
(OTI) mode (Fukuda and Ando, 2021).
701
HS-AFM imaging
The lipid compositions used in this study predominantly consisted of neutral 1,2-
dipalmitoyl-*sn*-glycero-3-phosphocholine (DPPC, Avanti) and positively charged 1,2-
dipalmitoyl-3-trimethylammonium-propane (DPTAP, Avanti). Liposomes were prepared using
DPPC/DPTAP (90/10, wt%), DPPC/DPTAP (75/25, wt%), and DPPC/DPTAP (50/50, wt%)
ratios, following our established sample preparation protocol (Ngo et al., 2015). These
liposomes, along with mica-supported lipid bilayers, were created accordingly.
Actin filaments were generated in F1 buffer containing 40 mM KCl, 20 mM PIPES-KOH
(pH 6.8), 1 mM $MgCl_2$, 0.5 mM EGTA, and 0.5 mM DTT in the presence of 0.5-1 mM ATP,
following the procedure described in our previous work (Ngo et al., 2015). To obtain actin
filaments bound with different nucleotides (ATP, ADP, ADP.Pi) and investigate the cooperative
binding of cofilin to these filaments, a similar protocol was employed (Ngo et al., 2015). Briefly,
actin filaments (5–10 μM) were made in F1 buffer containing 0.5-1 mM ATP and gently
immobilized on a positively charged lipid membrane composed of DPPC/DPTAP (90/10 wt%).
The actin filaments on the lipid surface were washed with 20 μl of imaging F1 buffer
additionally including either 0.5 mM ATP, 1 mM ADP/5 U/ml hexokinase/10 mM glucose, or 1
mM ATP + 10 mM Pi (K_2HPO_4/KH_2PO_4 (pH 6.8) buffer), resulting in F-ATP-actin, F-ADP-
actin, or F-ADP.Pi-actin, respectively. Subsequently, cofilin, prepared in the imaging F1 buffer
with the desired concentration and nucleotides, was introduced into the imaging chamber to
observe the cooperative binding of cofilin along the actin filaments.
To compare the helical twisting structures, the number of protomers per HHP, and MADs
in the normal F-actin structure with that in bare actin segments on both sides of the cofilin
clusters in the hybrid cofilactin/bare actin filaments at high spatiotemporal resolution in the same
imaging field, we immobilized these filaments sequentially on the same lipid membrane
(referred to **Figures supplement 6-7**). First, actin filaments were prepared by incubating G-actin
in F1 buffer and 0.5 mM ATP at a concentration of 20 μM for approximately 1 hour at 22°C,
resulting in F-actin. Second, cofilactin and hybrid cofilactin/bare actin filaments were formed by
mixing F-actin (10 μM) with cofilin (1 μM) in a small tube in F1 buffer and 0.5 mM ATP. The
final molar ratio was 10/1 (actin to cofilin). The mixture was incubated for 10 minutes at 22°C to
allow the binding, cluster expansion and actin filament severing to reach equilibrium. Third, a 2
μl solution containing cofilactin and hybrid cofilactin/bare actin filaments was introduced onto a
lipid membrane composed of DPPC/DPTAP (50/50 wt%) and incubated for 10 minutes at 22°C
to immobilize the filaments. Fourth, the surface was washed with 20 μl of imaging F1 buffer

containing 0.5 mM ATP, and then a 2 μl solution containing normal actin filaments (10 μM) was introduced onto the same lipid membrane. This sample was also incubated for 10 minutes at 22°C. Fifth, the surface was washed again with 20 μl of imaging F1 buffer containing 0.5 mM ATP prior to HS-AFM imaging.

Analysis of the growth rate of the cofilin cluster within the shortened bare half helices adjacent to the cofilin cluster in F-ADP.Pi-actin and F-ADP-actin

We measured the growth rates of the cofilin clusters within shortened bare half helices adjacent to the cofilin clusters on the PE side in F-ADP.Pi-actin and F-ADP-actin. As noted in the section of Results, when we were able to observe bare sections on both sides of a cofilin cluster, one of them had shorter HHP while the other was close to normal HHP, and based on our previous report (Ngo et al., 2015), we judged the bare section with shorter HHP is on the PE side of an actin filament. It is important to clarify that the collected data did not include the growth rates of cofilin within the normal half helices on the PE or BE sides, in order to exclude the random binding events of cofilin to actin filaments under various nucleotide conditions. By employing HS-AFM, we were able to monitor changes in the length of a half helix adjacent to a cofilin cluster and the peak height at individual crossover points. The lag time measurements were performed at the precise moment when the bare half helix adjacent to the cofilin cluster on the PE side began to shorten and continued until a newly saturated cluster emerged within this neighboring bare half helix. This determination was made by analyzing the increase in the peak heights (P1) before and after forming a newly matured cofilin cluster adjacent to a preformed cofilin cluster (P2) (as depicted in **Figure 9A**). The data presented in **Figures 7, 8C** were also included in this analysis. Each fully decorated-supertwisted half helix consists of 11 cofilin molecules (referred to **Figure supplement 1I**). To calculate the mean growth rate of the cofilin cluster within a shortened bare half helix adjacent to a cofilin cluster, we divided 11 cofilin molecules by the lag time in seconds.

We assessed torsional flexibility of actin filaments by measuring the HHPs in control actin filaments incubated with 1 mM ADP, 1mM ATP + 10 mM Pi, or 1 mM ATP + phalloidin saturated to actin, resulting in F-ADP-actin, F-ADP.Pi-actin, or phalloidin-stabilized F-actin, respectively, using our previous method (Ngo et al., 2015). Because, we did not examine the nucleotide-bound actin states when actin filaments were incubated with 1 mM ATP and phalloidin excess, we simply labelled them as phalloidin-stabilized F-actin and used as the control. The fraction of naturally supertwisted half-helices with HHPs less than 30 nm was determined by computing the areas under the curves derived from single Gaussian fitting curves (referred to **Figure 9C-D**).

HS-AFM data analysis and processing

The AFM data were analyzed using custom software, including Kodec, UMEX Viewer, and Umex Viewer for Drift Analysis, as described in previous studies (Hayakawa et al., 2022; Ngo et al., 2015). To measure the half helical pitch (HHP) and the number of actin protomer pairs per HHP, we utilized the custom software Umex Viewer for Drift Analysis, as described in (Hayakawa et al., 2022). Additionally, AFM images and videos were processed using Fiji-ImageJ (NIH, USA) and After Effects (Adobe, USA).

To quantitatively analyze the HHPs and count the number of protomer pairs per HHP, we employed Umex Viewer for Drift Analysis, which enabled the semiautomatic determination and measurement of the distance between the highest "cross-over" points between two long-pitch

strands. This involved creating a topographical line profile along the actin filaments, similar to
previous studies conducted by (Amyot et al., 2022; Hayakawa et al., 2022). Specifically, one
HHP was measured between the two highest cross-over points, while one helical pitch (HP) was
measured between the three highest cross-over points. In brief, we drew a longitudinal section
profile line along the actin filament, covering a length of 1-4 consecutive HHPs. The offset
values were set to zero with reference to the height of the lowest position along the longitudinal
section lines. Before analysis, we corrected the nonlinearity of the XY piezos by applying
nonlinear image scaling and reduced image noise using a Gaussian smoothing filter with a
standard deviation of 0.76 nm. The profile was obtained by averaging the signal within a 5-8 nm
bandpass filter along the filament. To ensure correct detection of axial distance (AD) between
two adjacent protomers within the same long-pitch strand, we set a range of 3–6.6 nm for minor
pitches. To facilitate correct detection of the HHPs between two crossover points, we set a range
of 15–25 nm for minimum major pitches for cofilactin and bare actin segments. These allow the
software to automatically penalize incorrect peaks. In some rare cases, the analyst also made the
decisions to add missing peaks or to remove incorrect peaks caused by noise. Typically, we
selected 30-61 actin filaments and measured the HHPs for approximately 20 consecutive time
frames. To minimize measurement errors in the HHP and the number of protomer pairs per HHP,
we specifically chose straight actin segments with clearly discernible protomers and drew line
profiles (with an average width of 3%) to measure 1-4 consecutive HHPs. This method was
consistently applied to analyze normal bare actin, cofilactin, and hybrid cofilactin/bare actin
filaments. We distinguished normal F-actin and cofilactin structures based on differences in the
peak height and HHP, as described in our previous study (Ngo et al., 2015).
To calculate the number of protomer pairs per HHP and the mean axial distance (MAD)
between two adjacent protomers along the same long-pitch strand, we semiautomatically
detected AD and counted the number of protomer pairs per HHP, as mentioned in other parts and
shown in **Figure supplement 2**. The MADs within an HHP were determined by dividing the
HHP by the actual counted number of protomer pairs per HHP.
To enhance the characteristics of actin protomers in the different types of filaments (bare
F-actin, cofilactin and C-actin-like structure in the hybrid cofilactin/bare actin segments), we
converted the raw AFM images into Laplacian-filtered images. This was achieved by applying a
high-pass filter known as the Laplacian of Gaussian (LoG) filter, which can be found at the
following link:
(https://homepages.inf.ed.ac.uk/rbf/HIPR2/log.htm). The Laplacian filter used in this process is a
type of high-pass filter, characterized by the kernel $\begin{bmatrix} -1 & -1 & -1 \\ -1 & 8 & -1 \\ -1 & -1 & -1 \end{bmatrix}$. Prior to the application of
the Laplacian filter, we employed a Gaussian filter with a typical sigma value of 1.27 pixel/nm.
This preliminary step helped reduce sensitivity to noise and enhanced the resulting Laplacian-
filtered images, resulting in improved visualization of the actin protomers.
818
Temporal autocorrelation analysis
The temporal autocorrelation analysis method can be found at the following link:
https://real-statistics.com/time-series-analysis/stochastic-processes/autocorrelation-function/.
In summary, the analysis involves the following steps: (**i**) Data processing and collection:
We directly measured and collected time series data for the axial distance (AD) between two
adjacent actin protomers along the same long-pitch strand in different actin filaments over time,
as shown in **Figure 3D**, using a custom semiautomatic analysis software called Umex View for

Drift Analysis. First, we converted raw AFM images containing actin filaments into Laplacian-
filtered images by applying a Laplacian of Gaussian (LoG) filter (typically with sigma ~ 1.27
pixel/nm) to eliminate low-frequency and high-frequency noise signals. Second, we selected a
short segment of the actin filament (~4 HHPs) located at the center of the AFM image and used a
segmented line running along the axis of the actin filament to measure the distance between actin
protomer pairs, as depicted in **Figure 3D**. To facilitate the semiautomatic analysis and determine
the peak-to-peak distance between individual ADs, we applied a bandpass filter to remove noise
by setting the low-pass frequency (LPF) cutoff at 3 nm and the high-pass frequency (HPF) cutoff
at 6 nm for all analyses. To ensure correct detection of AD between two adjacent protomers
within the same long-pitch strand, we set a range of 3–6.6 nm for minor pitches, allowing the
software to penalize incorrect peaks. Typically, the variation in the AD within each actin
filament over time was assessed using a series of AFM images (including 258 images captured at
a rate of 2 frames per second (fps)). (**ii**) Calculation of the autocorrelation coefficients: Each
autocovariance value was divided by the autocovariance at lag 0 to obtain the autocorrelation
coefficients. These coefficients represent the strength and direction of the linear relationship
between observations at different time lags (**Equation 1**). (**iii**) Plotting the autocorrelation
function (ACF): The autocorrelation coefficients were visualized by plotting the autocorrelation
function (ACF). The ACF plot displays the autocorrelation coefficients on the y-axis and the
time lags on the x-axis, as depicted in **Figure 3E**. (**iv**) Estimation of the time constants: The ACF
plot was fitted with a second-order exponential decay function to extract the time constants (t_1,
t_2), as depicted in **Figure 3E-F**.
The *ACF* (k) of a time series of values (y_i) at lag k is defined as
$$ACF(k) = [\frac{1}{n} \sum_{i=1}^{n-k}(y_i - \bar{y})(y_{i+k} - \bar{y})]/[\sum_{i=1}^{n}(y_i - \bar{y})^2] \text{ (\textbf{Equation 1})}$$
In our analysis, a time series of values (y_i) of the axial distance between two adjacent
actin pairs at lag k ($k \geq 0$) was measured for each actin filament in 258 consecutive AFM images
($n = 258$) collected at a frame rate of 2 fps, or 5 fps and 10 fps (**Figure supplement 3**). The
mean value, $\bar{y}$, was calculated.
853
Spatial autocorrelation analysis
To evaluate the degree of spatial clustering in the actin protomer structural data, we
conducted spatial autocorrelation analysis, as depicted in **Figure supplement 4**. Initially, we
generated a line profile by utilizing cubic spline interpolation along the ridgeline of an actin
filament (**Figure supplement 4A**). The actin filament includes a large helical twist and protomer
structures with periodicities of 36 nm and 6–7 nm, respectively. The helical twist exhibits a
typical height of 3 nm (**Figure supplement 4B**), which is three times larger than the 0.5–1.5 nm
height of the protomer (**Figure supplement 4C**). To explore the spatial self-correlation among
protomers, we applied a fast Fourier transform (FFT) and utilized a low-pass filter (LPF) and
high-pass filter (HPF). The filters isolated heights using LPF and HPF within a range of 5-8 nm
in the Fourier domain, corresponding to the actin protomer height along the filament (depicted in
Figure supplement 4C). To ensure correct detection of AD between two adjacent protomers
within the same long-pitch strand, we set a range of 3-6.6 nm for minor pitches, allowing the
software to penalize incorrect peaks. Then, we derived the autocorrelation function (ACF) of the
protomer heights by obtaining the complex conjugate (magnitude) and then performed an inverse
FFT. Additionally, we applied the Hilbert transform in the Fourier domain by doubling the
positive frequency data, zeroing the negative frequency data, and applying an inverse FFT. This
process produced an envelope curve. As a result, the envelop curves of the height profiles with

and without the BPF exhibit alternating positive and negative values at half helical pitch and protomer periodicities, respectively (**Figure supplement 4D-E**). In contrast, the envelope curve consistently shows positive values; thus, this curve is suitable for evaluating spatial autocorrelations. In the main text, we compared the envelope curves representing the spatial ACF results, measured at multiple time points and different conditions.

By visualizing the spatial ACF curves with distance (d) derived from F-actin and hybrid cofilactin/bare actin segments (**Figure 4C, 6C and Figure supplement 5A-B**), along with sparsely decorated cofilin-actin filaments strongly attached on lipid surface (**Figure supplement 7**), we predicted that the curves could be modelled using an exponential decaying sinusoidal function. This function, characterized by two primary components (exponential decay and sinusoid frequencies), enables us to derive parameters for the decay constant ($D1$) and the frequencies of the sinusoidal oscillation ($D2, D3, ...D9$), taking into account the phase angle of the sinusoidal waves ($\gamma1, \gamma2, \gamma3, \gamma4$). Based on previous studies (Ju and Rappaport, 2021; Zhang et al., 2008), we successfully formulated and applied a novel exponential decaying sinusoidal function, denoted as $\rho(d)$, to fit the spatial ACF curves (**Equation 2**).

$\rho(d) = exp(-d/D1) [D2cos (d/D3 + \gamma1) + D4sin (d/D5 + \gamma2) + D6cos (d/D7 + \gamma3) + D8sin (d/D9 +\gamma4)]$ (**Equation 2**)

For fitting, we utilized the Nadam optimizer (Dozat, 2016) and employed gradient descent in our approach. The mean square error served as our selected loss function for fine-tuning parameters. This optimization framework not only aided in the model's convergence but also improved its capacity to precisely capture the inherent patterns in the data. Consequently, it enhanced the effectiveness of our fitting solution. Finally, the $D1$ values, providing an indirect measure of the degree to which height values near each other in space are similar to one another in actin filaments, were shown.

References

Amyot R, Marchesi A, Franz CM, Casuso I, Flechsig H. 2022. Simulation atomic force microscopy for atomic reconstruction of biomolecular structures from resolution-limited experimental images. *PLoS Comput Biol* **18**:1–15. doi:10.1371/journal.pcbi.1009970

Ando T. 2022. High-Speed Atomic Force Microscopy in Biology: Directly Watching Dynamics of Biomolecules in Action. Springer Berlin, Heidelberg. doi:10.1007/978-3-662-64785-1

Ando T. 2012. High-speed atomic force microscopy coming of age. *Nanotechnology* **23**:1–27. doi:10.1088/0957-4484/23/6/062001

Ando T, Fukuda S, Ngo KX, Flechsig H. 2023. High-speed atomic force microscopy for filming protein molecules in dynamic action. *Annu Rev Biophys* **53**:19–39. doi:10.1146/annurev-biophys-030722-113353

Ando T, Uchihashi T, Kodera N. 2013. High-speed AFM and applications to biomolecular systems. *Annu Rev Biophys* **42**:393–414. doi:10.1146/annurev-biophys-083012-130324

Ando T, Uchihashi T, Kodera N, Yamamoto D, Miyagi A, Taniguchi M, Yamashita H. 2008. High-speed AFM and nano-visualization of biomolecular processes. *Pflugers Arch Eur J Physiol* **456**:211–225. doi:10.1007/s00424-007-0406-0

Andrianantoandro E, Pollard TD. 2006. Mechanism of Actin Filament Turnover by Severing and Nucleation at Different Concentrations of ADF/Cofilin. *Mol Cell* **24**:13–23.

doi:10.1016/j.molcel.2006.08.006

Bamburg JR, Bernstein BW. 2010. Roles of ADF/cofilin in actin polymerization and beyond.
F1000 Biol Rep **2**:1–7. doi:10.3410/B2-62

Bibeau JP, Gray S, De La Cruz EM. 2021. Clusters of a Few Bound Cofilins Sever Actin
Filaments. *J Mol Biol* **433**:166833. doi:10.1016/j.jmb.2021.166833

Blanchoin L, Pollard TD. 1999. Mechanism of interaction of Acanthamoeba actophorin
(ADF/Cofilin) with actin filaments. *J Biol Chem* **274**:15538–15546.
doi:10.1074/jbc.274.22.15538

Bobkov A a, Muhlrad A, Pavlov D a, Kokabi K, Yilmaz A, Reisler E. 2006. Cooperative effects
of cofilin (ADF) on actin structure suggest allosteric mechanism of cofilin function. *J Mol*
Biol **356**:325–34. doi:10.1016/j.jmb.2005.11.072

Carlier MF, Laurent V, Santolini J, Melki R, Didry D, Xia GX, Hong Y, Chua NH, Pantaloni D.
1997. Actin depolymerizing factor (ADF/cofilin) enhances the rate of filament turnover:
Implication in actin-based motility. *J Cell Biol* **136**:1307–1322. doi:10.1083/jcb.136.6.1307

Carlier MF, Pantaloni D. 1988. Binding of phosphate to F-ADP-actin and role of F-ADP-P(i)-
actin in ATP-actin polymerization. *J Biol Chem* **263**:817–825. doi:10.1016/s0021-
9258(19)35428-6

Chou SZ, Pollard TD. 2019. Mechanism of actin polymerization revealed by cryo-EM structures
of actin filaments with three different bound nucleotides. *Proc Natl Acad Sci U S A*
116:4265–4274. doi:10.1073/pnas.1807028115

Cock PJA, Antao T, Chang JT, Chapman BA, Cox CJ, Dalke A, Friedberg I, Hamelryck T,
Kauff F, Wilczynski B, De Hoon MJL. 2009. Biopython: Freely available Python tools for
computational molecular biology and bioinformatics. *Bioinformatics* **25**:1422–1423.
doi:10.1093/bioinformatics/btp163

De La Cruz EM, Roland J, McCullough BR, Blanchoin L, Martiel J-L. 2010. Origin of twist-
bend coupling in actin filaments. *Biophys J* **99**:1852–60. doi:10.1016/j.bpj.2010.07.009

Dominguez R. 2004. Actin-binding proteins - A unifying hypothesis. *Trends Biochem Sci*
29:572–578. doi:10.1016/j.tibs.2004.09.004

Dominguez R, Holmes KC. 2011. Actin Structure and Function. *Annu Rev Biophys* **40**:1–20.
doi:10.1146/annurev-biophys-042910-155359.

Dozat T. 2016. Incorporating Nesterov Momentum into Adam. *ICLR Work* 1–4.

Egelman EH. 1994. Actin Filament Structure: The ghost of ribbons past. *Curr Biol* **4**:79–81.
doi:10.1016/S0960-9822(00)00020-8

Egelman EH, Francis N, DeRosier DJ. 1982. F-actin is a helix with a random variable twist.
Nature **298**:131–135. doi:10.1038/298131a0

Fineberg A, Takagi Y, Thirumurugan K, Andrecka J, Billington N, Young G, Cole D, Burgess
SA. 2024. Myosin-5 varies its step length to carry cargo straight along the irregular F-actin
track. *Proc Natl Acad Sci* **121**:1–12. doi:10.1073/pnas.2401625121

Fujii T, Iwane AH, Yanagida T, Namba K. 2010. Direct visualization of secondary structures of
F-actin by electron cryomicroscopy. *Nature* **467**:724–728. doi:10.1038/nature09372

Fukuda S, Ando T. 2021. Faster high-speed atomic force microscopy for imaging of
biomolecular processes. *Rev Sci Instrum* **92**. doi:10.1063/5.0032948

Galkin VE, Orlova a, Lukoyanova N, Wriggers W, Egelman EH. 2001. Actin depolymerizing
factor stabilizes an existing state of F-actin and can change the tilt of F-actin subunits. *J
Cell Biol* **153**:75–86.

Galkin VE, Orlova A, Kudryashov DS, Solodukhin A, Reisler E, Schröder GF, Egelman EH.
2011. Remodeling of actin filaments by ADF/cofilin proteins. *Proc Natl Acad Sci U S A*
108:20568–20572. doi:10.1073/pnas.1110109108

Galkin VE, Orlova A, Vos MR, Schröder GF, Egelman EH. 2015. Near-Atomic Resolution for
One State of F-Actin. *Structure* **23**:173–182. doi:10.1016/j.str.2014.11.006

Grintsevich EE, Yesilyurt HG, Rich SK, Hung R-J, Terman JR, Reisler E. 2016. F-actin
dismantling through a redox-driven synergy between Mical and cofilin. *Nat Cell Biol*
18:876–85. doi:10.1038/ncb3390

Harris AR, Jreij P, Belardi B, Joffe AM, Bausch AR, Fletcher DA. 2020. Biased localization of
actin binding proteins by actin filament conformation. *Nat Commun* **11**:1–13.
doi:10.1038/s41467-020-19768-9

Hayakawa K, Sakakibara S, Sokabe M, Tatsumi H. 2014. Single-molecule imaging and kinetic
analysis of cooperative cofilin-actin filament interactions. *Proc Natl Acad Sci U S A*
111:9810–9815. doi:10.1073/pnas.1321451111

Hayakawa Y, Takaine M, Ngo KX, Imai T, Yamada M, Behjat AB, Umeda K, Hirose K,
Yurtsever A, Kodera N, Tokuraku K, Numata O, Fukuma T, Ando T, Nakano K, Uyeda
TQP. 2022. Actin binding domain of Rng2 sparsely bound on F-actin strongly inhibits actin
movement on myosin II. *Life Sci Alliance* **6**:e202201469. doi:10.26508/lsa.202201469

Holmes KC, Popp D, Gebhard W, Kabsch W. 1990. Atomic model of the actin filament. *Nature*
347:44–49. doi:10.1038/347044a0

Huehn A, Cao W, Austin Elam W, Liu X, De La Cruz EM, Sindelar C V. 2018. The actin
filament twist changes abruptly at boundaries between bare and cofilin-decorated segments.
J Biol Chem **293**:5377–5383. doi:10.1074/jbc.AC118.001843

Huehn AR, Bibeau JP, Schramm AC, Cao W, De La Cruz EM, Sindelar C V. 2020. Structures of
cofilin-induced structural changes reveal local and asymmetric perturbations of actin
filaments. *Proc Natl Acad Sci U S A* **117**:1478–1484. doi:10.1073/pnas.1915987117

Isambert H, Venier P, Maggs AC, Fattoum A, Kassab R, Pantaloni D, Carlier MF. 1995.
Flexibility of actin filaments derived from thermal fluctuations. Effect of bound nucleotide,
phalloidin, and muscle regulatory proteins. *J Biol Chem* **270**:11437–11444.
doi:10.1074/jbc.270.19.11437

Ivanova JR, Benk AS, Schaefer J V., Dreier B, Hermann LO, Plückthun A, Missirlis D, Spatz JP.
2024. Designed Ankyrin Repeat Proteins as Actin Labels of Distinct Cytoskeletal Structures

992   in Living Cells. *ACS Nano*. doi:10.1021/acsnano.3c12265

993 Jegou A, Romet-Lemonne G. 2019. The many implications of actin filament helicity. *Semin Cell*
994   *Dev Biol* **102**:65–72. doi:10.1016/j.semcdb.2019.10.018

995 Jégou A, Romet-Lemonne G. 2016. Single Filaments to Reveal the Multiple Flavors of Actin.
996   *Biophys J* **110**:2138–2146. doi:10.1016/j.bpj.2016.04.025

997 Ju S, Rappaport TS. 2021. 140 GHz Urban Microcell Propagation Measurements for Spatial
998   Consistency Modeling. *IEEE Int Conf Commun* 1–6. doi:10.1109/ICC42927.2021.9500677

999 Kabsch W, Mannherz HG, Suck D, Pai EF, Holmes KC. 1990. Atomic structure of the actin:
1000   DNase I complex. *Nature* **347**:37–44. doi:10.1038/347037a0

1001 Kanematsu Y, Narita A, Oda T, Koike R, Ota M, Takano Y, Moritsugu K, Fujiwara I, Tanaka K,
1002   Komatsu H, Nagae T, Watanabe N, Iwasa M, Maéda Y, Takeda S. 2022. Structures and
1003   mechanisms of actin ATP hydrolysis. *Proc Natl Acad Sci U S A* **119**:1–12.
1004   doi:10.1073/pnas.2122641119

1005 Kasai M, Asakura S, Oosawa F. 1962. The cooperative nature of G-F transformation of actin.
1006   *Biochim Biophys Acta* **57**:22–31.

1007 Kodera N, Noshiro D, Dora SK, Mori T, Habchi J, Blocquel D, Gruet A, Dosnon M, Salladini E,
1008   Bignon C, Fujioka Y, Oda T, Noda NN, Sato M, Lotti M, Mizuguchi M, Longhi S, Ando T.
1009   2021. Structural and dynamics analysis of intrinsically disordered proteins by high-speed
1010   atomic force microscopy. *Nat Nanotechnol* **16**:181–189. doi:10.1038/s41565-020-00798-9

1011 Lane TJ. 2023. Protein structure prediction has reached the single-structure frontier. *Nat Methods*
1012   **20**:170–173. doi:10.1038/s41592-022-01760-4

1013 Lappalainen P, Drubin DG. 1997. Cofilin promotes rapid actin filament turnover in vivo. *Nature*
1014   **388**:78–82. doi:10.1038/40418

1015 Lorenz M, Popp D, Holmes KC. 1993. Refinement of the F-actin model against x-ray fiber
1016   diffraction data by the use of a directed mutation algorithm. *J Mol Biol* **234**:826–836.

1017 Mannherz HG, Ballweber E, Galla M, Villard S, Granier C, Steegborn C, Schmidtmann A,
1018   Jaquet K, Pope B, Weeds AG. 2007. Mapping the ADF/Cofilin Binding Site on Monomeric
1019   Actin by Competitive Cross-linking and Peptide Array: Evidence for a Second Binding Site
1020   on Monomeric Actin. *J Mol Biol* **366**:745–755. doi:10.1016/j.jmb.2006.11.100

1021 McCullough BR, Blanchoin L, Martiel JL, De La Cruz EM. 2008. Cofilin increases the bending
1022   flexibility of actin filaments: implications for severing and cell mechanics. *J Mol Biol*
1023   **381**:550–558. doi:10.1016/j.jmb.2008.05.055

1024 McGough A, Pope B, Chiu W, Weeds A. 1997. Cofilin changes the twist of F-actin: implications
1025   for actin filament dynamics and cellular function. *J Cell Biol* **138**:771–781.

1026 Miller MD, Phillips GN. 2021. Moving beyond static snapshots: Protein dynamics and the
1027   Protein Data Bank. *J Biol Chem* **296**:100749. doi:10.1016/j.jbc.2021.100749

1028 Mizuno H, Tanaka K, Yamashiro S, Narita A, Watanabe N. 2018. Helical rotation of the
1029   diaphanous-related formin mDia1 generates actin filaments resistant to cofilin. *Proc Natl*

Acad Sci U S A **115**:E5000–E5007. doi:10.1073/pnas.1803415115

Narita A. 2020. ADF/cofilin regulation from a structural viewpoint. *J Muscle Res Cell Motil*
41:141–151. doi:10.1007/s10974-019-09546-6

Ngo KX, Kodera N, Katayama E, Ando T, Uyeda TQ. 2015. Cofilin-induced unidirectional
cooperative conformational changes in actin filaments revealed by high-speed atomic force
microscopy. *Elife* **4**:1–22. doi:10.7554/eLife.04806

Ngo KX, Umeki N, Kijima ST, Kodera N, Ueno H, Furutani-Umezu N, Nakajima J, Noguchi
TQP, Nagasaki A, Tokuraku K, Uyeda TQP. 2016. Allosteric regulation by cooperative
conformational changes of actin filaments drives mutually exclusive binding with cofilin
and myosin. *Sci Rep* **6**:1–11. doi:10.1038/srep35449

Niina T, Fuchigami S, Takada S. 2020. Flexible Fitting of Biomolecular Structures to Atomic
Force Microscopy Images via Biased Molecular Simulations. *J Chem Theory Comput*
16:1349–1358. doi:10.1021/acs.jctc.9b00991

Niina T, Matsunaga Y, Takada S. 2021. Rigid-body fitting to atomic force microscopy images
for inferring probe shape and biomolecular structure. *PLoS Comput Biol* **17**:1–24.
doi:10.1371/journal.pcbi.1009215

Oda T, Iwasa M, Aihara T, Maéda Y, Narita A. 2009. The nature of the globular- to fibrous-actin
transition. *Nature* **457**:441–445. doi:10.1038/nature07685

Oda T, Takeda S, Narita A, Maéda Y. 2019. Structural Polymorphism of Actin. *J Mol Biol*
431:3217–3228. doi:10.1016/j.jmb.2019.05.048

Okura K, Matsumoto T, Narita A, Tatsumi H. 2023. Mechanical Stress Decreases the Amplitude
of Twisting and Bending Fluctuations of Actin Filaments. *J Mol Biol* **435**:168295.
doi:10.1016/j.jmb.2023.168295

Oosterheert W, Klink BU, Belyy A, Pospich S, Raunser S. 2022. Structural basis of actin
filament assembly and aging. *Nature* **611**:374–379. doi:10.1038/s41586-022-05241-8

Paavilainen VO, Oksanen E, Goldman A, Lappalainen P. 2008. Structure of the actin-
depolymerizing factor homology domain in complex with actin. *J Cell Biol* **182**:51–59.
doi:10.1083/jcb.200803100

Reynolds MJ, Hachicho C, Carl AG, Gong R, Alushin GM. 2022. Bending forces and nucleotide
state jointly regulate F-actin structure. *Nature*. doi:10.1038/s41586-022-05366-w

Rich SA, Estes JE. 1976. Detection of conformational changes in actin by proteolytic digestion:
Evidence for a new monomeric species. *J Mol Biol* **104**:777–792. doi:10.1016/0022-
2836(76)90181-9

Suarez C, Roland J, Boujemaa-Paterski R, Kang H, McCullough BR, Reymann A-C, Guérin C,
Martiel J-L, De la Cruz EM, Blanchoin L. 2011. Cofilin tunes the nucleotide state of actin
filaments and severs at bare and decorated segment boundaries. *Curr Biol* **21**:862–8.
doi:10.1016/j.cub.2011.03.064

Tanaka K, Takeda S, Mitsuoka K, Oda T, Kimura-Sakiyama C, Maéda Y, Narita A. 2018.

Structural basis for cofilin binding and actin filament disassembly. *Nat Commun* **9**:1–3.
1069   doi:10.1038/s41467-018-04290-w

1070   Tokuraku K, Kuragano M, Uyeda TQP. 2020. Long-range and directional allostery of actin
1071   filaments plays important roles in various cellular activities. *Int J Mol Sci* **21**.
1072   doi:10.3390/ijms21093209

1073   Tokuraku K, Kurogi R, Toya R, Uyeda TQP. 2009. Novel mode of cooperative binding between
1074   myosin and Mg2+-actin filaments in the presence of low concentrations of ATP. *J Mol Biol*
1075   **386**:149–162. doi:10.1016/j.jmb.2008.12.008

1076   Uyeda TQP, Iwadate Y, Umeki N, Nagasaki A, Yumura S. 2011. Stretching actin filaments
1077   within cells enhances their affinity for the myosin ii motor domain. *PLoS One* **6**:e26200.
1078   doi:10.1371/journal.pone.0026200

1079   Wioland H, Guichard B, Senju Y, Myram S, Lappalainen P, Jégou A, Romet-Lemonne G. 2017.
1080   ADF/Cofilin Accelerates Actin Dynamics by Severing Filaments and Promoting Their
1081   Depolymerization at Both Ends. *Curr Biol* **27**:1956-1967.e7. doi:10.1016/j.cub.2017.05.048

1082   Xue B, Leyrat C, Grimes JM, Robinson RC. 2014. Structural basis of thymosin-β4/profilin
1083   exchange leading to actin filament polymerization. *Proc Natl Acad Sci U S A* **111**:E4596–
1084   E4605. doi:10.1073/pnas.1412271111

1085   Zhang Y, Zhang J, Dongt D, Nie X, Liu G, Zhang P. 2008. A novel spatial autocorrelation model
1086   of shadow fading in urban macro environments. *IEEE GLOBECOM 2008 - 2008 IEEE*
1087   *Glob Telecommun Conf New Orleans, LA, USA* 1–5. doi:10.1109/GLOCOM.2008.ECP.801

1088   Zimmermann D, Santos A, Kovar DR, Rock RS. 2015. Actin Age Orchestrates Myosin-5 and
1089   Myosin-6 Runlengths. *Curr Biol* **25**:2057–2062. doi:10.1016/j.cub.2015.06.033.

1090

Acknowledgments
The authors express their gratitude for the facility and financial support provided by the WPI
Nano Life Science Institute (WPI NanoLSI), Kanazawa University, as well as the instrumental
assistance provided by Toshio Ando. Special thanks are extended to Huan Tran from Georgia
Institute of Technology, and Damien Hall and Thao Phuong Ngo from WPI-NanoLSI, Kanazawa
University, for their valuable contribution to the data analyses. The authors also acknowledge the
financial support from KAKENHI (Japan Society for the Promotion of Science) for K.X.N.
(19K06581, 23K05713, 23H02452-01) and T.Q.P.U. (#23H02452) and N.K. (20H00327); and
CREST, Japan Science and Technology Agency (JPMJCR1762 to N.K.).

1100

1101   **Author contributions**
1102   K.X.N., H.T.V., N.K., T.Q.P.U. designed the concepts and experiments. K.X.N., H.T.V., N.K.
1103   performed experiments, K.X.N., H.T.V., K.U., M.N.T. analysed the data. K.X.N., H.T.V.,
1104   T.Q.P.U. wrote the manuscript. All authors revised the manuscript.

1105

1106   **Conflicts of interest**
1107   The authors declare no conflicts of interest. The funders have no role in interpreting the data.

1108
1109

(A)

(B) front view   side view

(C)

1110

Figure 1 The significant movements of the OD during F-actin to C-actin to G-actin transitions,
as revealed by PCA.
(**A**) PCA of actin chains reported in 46 PDB structures (referred to **Table supplement 1**) reveals
collective transition pathway between different actin conformations: G-actin (soluble monomer –
blue box), F-actin (filamentous protomer reported with no cofilin bound – green box), C-actin
(filamentous protomer reported within cofilactin after removal of cofilin – red box), tilted actin
(reported in 3J8J and 3J8K – magenta box) and single-cofilin-bound actin, where one cofilin
molecule bound to an actin protomer within an actin filament (reported in 6UBY – orange box).
Protomers at the most barbed end side of a cofilin cluster in 6UC4 also belong to the latter
cluster. Notably, a twinfilin-1 bound structure of soluble actin molecule (3DAW) (Paavilainen et
al., 2008), previously classified as C-actin (Tanaka et al., 2018), was found between the G-actin
and C-actin clusters. PDB entries are coloured with nucleotide molecules present, and structures
lacking reported nucleotides are labelled "Apo". The inset shows a magnified region of the F-
actin's PCA cluster (corresponding to the black box in the main graph).
(**B**) The PC1, represented by blue arrows, induces a rotational motion of the OD around hinge
region, causing the nucleotide binding cleft between SD2 and SD4 (depicted as a yellow
arrowhead) to close during the transition from G-actin (cyan chain) to C-actin (pink chain).
(**C**) The PC2, indicated by red arrows, results in an untwisting motion of the SD2, leading to the
flattening of OD during the transformation from C-actin (pink chain) to F-actin (green chain),
but without affecting the closing of nucleotide binding cleft (nearly unchanging PC1). The
rotational axes in panels B and C were denoted by blue and red asterisks, respectively.

1157
Figure 2 The variations in half helices and the number of protomer pairs per HHP in F-actin, C-
actin, and cofilactin structures in relationship with PC2 analysis.
(**A-C**) Side-views (represented by blue boxes, similar to the right panel of Figure 1C) and outer-
views (represented by black boxes) of F-actin, C-actin protomers without and with cofilins along
long filaments are depicted. The white translucent cylinders represent the central axes of the
filaments. In panel B, a yellow arrowhead highlights a portion of SD2 including D-loop that is
buried deeper into the C-actin filament as a result of the reduction of PC2 (represented by the red
arrow, corresponding to the red arrow in the right panel of Figure 1C).
(**D-E**) Diagrams of 2D surfaces (unwrapped from imaginary cylinders along the filaments and
cut open along imaginary lines scanned by the AFM tip) illustrate the different twisting angles
between adjacent protomers in the same strand appears to increase from approximately 26° to
around 36° if view from the outer side, resulting in HHPs with varying numbers of detected
protomer pairs (marked between red lines). This demonstrates that the global shortened pitch is
directly related to the OD twist of the local actin protomers. The green and pink objects represent
outer-views of OD domains of the focused chains, along with cofilins shown in black. The
opposite chains are depicted in grey.

1190
Figure 3 Exploring the dynamics of helical structures of long actin filaments using HS-AFM.
(**A**) The provided image shows actin filaments with varying numbers of protomer pairs in one
half helix. To measure the HHPs and the number of protomer pairs per HHP, a longitudinal
section profile of two consecutive half helices was taken as depicted, following the previous
method (Hayakawa et al., 2022). Actin filaments were gently immobilized on lipid membrane
composed of 1,2-dipalmitoyl-sn-glycero-3-phosphocholine (DPPC) and 1,2-dipalmitoyl-3-
trimethylammonium-propane (DPTAP) (90/10 wt%) and imaged in F1 buffer containing 0.5 mM
ATP. The offset values were set to zero with reference to the height of the lowest position along
the longitudinal section lines. The scale bar represents 25 nm. For the detail, see Methods.
(**B**) The correlation between the counted protomer pairs from image A and the HHP probed by
HS-AFM is shown. Black symbols represent the experimental data, with the linear fit (y = 10.71
+ 3.43x, *R* = 0.9903). The calculated data (red symbols) also demonstrate a correlation between
the number of protomer pairs per HHP and the HHP, which is calculated as 5.5 nm multiplied by
the number of counted protomer pairs per HHP.
(**C**) The distribution of MADs is shown for half helices composed of different number of
counted protomer pairs. The overall MAD (mean ± SD) is 5.1 ± 0.3 nm (*n* = 15021).
(**D**) The axial distance (AD) between two adjacent actin protomers along the same long-pitch
strand in individual bare actin filaments were directly measured as the peak-to-peak distance in

four consecutive HHPs. Each protomer pair within the filament was assigned a number ranging
from 1 to 27. The ADs were measured in a time series, with the imaging rate of 0.5 seconds per
frame and a total of 258 frames. These AD values for 27 protomer pairs within the actin filament
in a time series of 258 frames were utilized to determine the autocorrelation coefficients. The
still AFM image was shown after processing with Laplacian of Gaussian filter.
(**E**) An example of the ACF plot displays the autocorrelation coefficients on the y-axis against
the time lags on the x-axis. The plotted data was fitted using a second-order exponential decay
function with the time constants of (t_1, t_2), represented by the red curve.
(**F**) A histogram was constructed using the time constants obtained from many similar ACF plots
and fittings obtained from different actin filaments ($n = 61$), as showcased in (D, E). The number
of data points was 122. These time constants are associated with the overall trend of time decay
in autocorrelation of axial distance between two adjacent protomers in the same protofilament.

Figure 4 The variation of helical twists measured by HHPs, the number of protomer pairs per HHP, and the MADs in long actin filaments through HS-AFM. Actin filaments were gently immobilized on lipid membrane composed of 1,2-dipalmitoyl-sn-glycero-3-phosphocholine (DPPC) and 1,2-dipalmitoyl-3-trimethylammonium-propane (DPTAP) (90/10 wt%) and imaged in F1 buffer containing 0.5 mM ATP.

(**A**) The graph demonstrates the changes in HHPs, the number of protomer pairs per HHP, and the MADs over time. The statistical analysis using two-population *t-test* at a significance level of $p \leq 0.05$ revealed that the difference between MAD1 (5.1 ± 0.3 nm, n = 32) and MAD2 (5.2 ± 0.2 nm, n = 32) is not statistically significant ($p = 0.2365$). However, MAD1 and MAD3 (5.2 ± 0.3 nm, n = 32) exhibit a slight difference ($p = 0.02895$).

(**B**) Representative still images obtained from AFM are presented. Arrows indicate the peaks (P1, P2, P3, P4) identified in the raw AFM images. HHP1, HHP2, and HHP3 were measured between P1-P2, P2-P3, and P3-P4, respectively. Scale bars: 25 nm.

(**C**) The spatial ACF curves were produced for a canonical actin filament (F1) at various time
points (i.e., Data 1, 2, 3). The curves were fitted using an exponential decaying sinusoidal
function (**Equation 2**), and the resulting decay constant (*D1*) values from individual curves were
presented. The still AFM image is shown after processing with Laplacian of Gaussian filter. Bar:
25 nm.
(**D**) The *D1* values were determined through a similar fitting of spatial ACF curves, as illustrated
in (**C**). This examination was conducted at different time points (with an imaging rate of 0.5
seconds per frame and a total of 50 frames for each canonical filament (F1-F5) using the data, as
shown in **Figure Supplement 5A**). Subsequently, the distribution of *D1* values was analyzed,
yielding the mean ± SD value of 27.2 ± 6.7 nm, based on a sample size of 248.

Figure 5 Asymmetric helical twisting on both sides of a small cofilin cluster. Red arrowheads indicate small area of cofilin clusters (in boxes). Scale bars: 25 nm. Based on our previous study (Ngo et al., 2015), we judged that the bare half helix with the shorter HHP adjacent to the cofilin cluster is on the PE side and that the normal HHP is located on the BE side.

(**A**) A still AFM image shows normal actin filaments, along with a longitudinal section profile
representing two consecutive half helices 1 and 2 (denoted HHP1, HHP2).
(**B**) A still AFM image displays a hybrid segment consisting of cofilactin and bare actin areas,
with a longitudinal section profile representing two consecutive half helices. A short filament
was observed where a small cofilin cluster cut filament at cofilactin/bare actin boundary on the
BE side, followed by two consecutive bare half helices 3 and 4 (denoted HHP3, HHP4) on the
PE side. The shortened bare half helix 3 and normal bare half helix 4 are presented. The hybrid
cofilactin/bare actin filaments, priorly made in solution, were strongly immobilized on lipid
membrane composed of 1,2-dipalmitoyl-sn-glycero-3-phosphocholine (DPPC) and 1,2-
dipalmitoyl-3-trimethylammonium-propane (DPTAP) (50/50 wt%) and imaged in F1 buffer
containing 0.5 mM ATP (see Methods).
(**C**) A still AFM image depicts a hybrid segment comprising cofilactin and bare actin areas,
along with a longitudinal section profile representing two consecutive half helices 5 and 6
(denoted HHP5, HHP6). In this case, a short filament contained a small cofilin cluster (estimated
as 2 to 4 molecules) in the middle, configured by a normal bare half helix 5 on the BE side and a
shortened bare half helix 6 on the PE side. We estimated the size of a cofilin cluster by
measuring its height and width within the white box around P8, distinguishing it with the control
bare section lacking cofilin bindings (P7, P9). Despite the invisibility of cofilin molecules below,
we made the assumption that cofilin bound uniformly to two strands at a molar ratio of 1:1,
enabling us to account for their presence.

1366
Figure 6 The variation of half helices, the number of protomer pairs per HHP, and the MADs in
a hybrid cofilactin/bare actin filament probed by HS-AFM. Based on our previous study (Ngo et
al., 2015), we judged that the bare half helix with the shorter HHP adjacent to the cofilin cluster
is on the PE side and that the normal HHP is located on the BE side.
(**A**) Representative still images obtained from AFM in **Figure 5C** are displayed. Arrows
indicate the peaks (P7, P8, P9) identified in the raw AFM images. A normal bare half helix 5
(denoted HHP5) and a shortened bare half helix 6 (denoted HHP6) were measured between P7-

P8 and P8-P9, respectively. Cyan and white arrows indicate the peak positions within cofilin
clusters and bare actin peaks, respectively. Boxes denote the cofilin clusters. Scale bars are 25
nm.
(**B**) The graph depicts the analysis of half helices 5 and 6, the number of protomer pairs in each
half helix (Protomer Pairs (5) and Protomer Pairs (6)), and the MAD5 and MAD6 over time.
Two-populations *t-test* was used to validate the statistical difference between MAD5 (5.0 ± 0.2
nm) and MAD6 (5.4 ± 0.3 nm). The sample size was 12.
(**C**) The spatial ACF curves were generated for a normal half helix 5 and a shorter half helix 6
near a cofilin cluster on either the PE or BE side in a hybrid cofilactin/bare actin filament at
various time points (i.e., Data 1, 2, 3). The curves were fitted using an exponential decaying
sinusoidal function (**Equation 2**), and the resulting decay constant ($D1$) values from individual
curves were presented. This examination was conducted at different time points (with an imaging
rate of 0.5 seconds per frame and a total of 50 frames for each half helix on the PE or BE side
using the data, as shown in **Figure Supplement 5B**). The $D1$ values were determined through a
similar fitting of spatial ACF curves. Subsequently, the distribution of $D1$ values was analyzed,
yielding the mean $\pm$ SD values for each half helix on the PE and BE side of 12.7 ± 7.0 nm and
12.6 ± 6.6 nm, respectively. The sample size of each case was 50. The statistical analysis using
two-population *t-test* at a significance level of $p \leq 0.05$ revealed that the difference between two
mean values was not significant. The still AFM images are shown after processing with
Laplacian of Gaussian filter. Scale bars: 25 nm.

1420
Figure 7 Propagation of supertwisted half helices within the cofilactin region beyond the
boundary of cofilactin and bare actin, leading to the pre-shortening of a bare half helix and
subsequent recruitment of cofilin cluster growth.
(**A**) Still AFM images show the time-dependent growth of cofilin clusters beyond the boundary.
The arrow indicates the direction of cofilin cluster growth. The peaks of cofilactin and bare actin
segments are marked by yellow and white arrowheads, respectively. Peak heights (P1-P6) were
determined in five consecutive half helices. Actin filaments were loosely immobilized on a lipid
membrane composed of 1,2-dipalmitoyl-sn-glycero-3-phosphocholine (DPPC) and 1,2-
dipalmitoyl-3-trimethylammonium-propane (DPTAP) (90/10 wt%) and imaged in an F1 buffer
containing 300 nM cofilin, 1 mM ATP and 10 mM Pi. Scale bars: 25 nm.
(**B**) Examination of changes in peak heights and HHPs in the cofilactin and bare actin regions
over time. Peak heights and HHPs were measured as described previously (Ngo et al., 2015).
(**C**) Histograms of peak heights (8.6 ± 0.4 nm, 10.3 ± 0.5 nm, n = 726) and HHPs (25.3 ± 2.6 nm,
36.9 ± 3.6 nm, n = 605) were obtained from the data in (B). The significant difference between
the peaks in double Gaussian fitting was statistically examined using an *F-test*. The green and
red curves denote the fractions of F-actin and cofilactin structures, respectively.

Figure 8 Cooperative binding of cofilin is heavily dependent on the dynamic and cooperative conformational changes involving helical twisting within actin filaments.

(**A**) The cooperative binding of cofilin to actin filaments was significantly inhibited when the torsional motion during helical twisting within the filaments was suppressed. The filaments were strongly attached onto a highly positively charged lipid membrane containing 1,2-dipalmitoyl-sn-glycero-3-phosphocholine (DPPC) and 1,2-dipalmitoyl-3-trimethylammonium-propane (DPTAP) (75/25 wt%) and imaged in F1 buffer containing 2000 nM cofilin and 0.5 mM ATP.

(**B, C, D**) The normal cooperative bindings of cofilin to actin filaments were observed when the filaments were loosely immobilized on weakly positively charged lipid membrane comprising DPPC/DPTAP (90/10 wt%) and imaged in F1 buffer under different conditions: 300 nM cofilin + 0.5 mM ATP (B), 300 nM cofilin + 1 mM ADP (C), and 75 nM cofilin + 1 mM ATP + 10 mM Pi (D). The changes in peak heights and HHPs of a representative actin filament under each condition were analyzed over time. Histograms were constructed by analyzing the changes in peak heights and HHPs of all actin filaments in the observed field during the growth of the cofilin cluster at various time intervals after adding cofilin. The statistical significance of the peaks in the histograms was examined using an *F-test*. The green and red curves denote the fractions of F-actin and cofilactin structures, respectively. Time labels in the still AFM images indicate the elapsed time after adding cofilin to the imaging buffer. Peaks of bare actin and cofilactin are denoted by white and yellow arrowheads, respectively. Scale bars: 25 nm.

1484
Figure 9 Comparison of the growth rate of cofilin clusters within the shortened bare half helices
that resemble the C-actin structure adjacent to cofilin clusters on the PE side, for both F-ADP-
actin and F-ADP.Pi-actin.
(**A**) In this cartoon representation, hybrid cofilactin/bare actin filaments are depicted, consisting
of four half helices. Two of these half helices are fully decorated with cofilin clusters in the
middle, followed by a shortened bare half helix and a normal long bare half helix on the PE and
BE sides, respectively. P1-P5 were labels indicating peak heights at individual crossover points.
To make a meaningful comparison, the lag time measurements were initiated exclusively when a
bare half helix between P1 and P2 experienced shortening, and they continued until a newly
matured cluster emerged in P1 within the shortened bare half helix (P1-P2) adjacent to the
preformed cofilin cluster (P2). Each fully decorated supertwisted half helix contains 11 cofilin
molecules (referred to **Figure supplement 2F-I**). To determine the mean growth rate of the
cofilin cluster within a shortened bare half helix adjacent to the cofilin cluster, 11, the number of
cofilin molecules in one half helix, was divided by the lag time in seconds.
(**B**) This part presents a comparison of the growth rate of cofilin clusters within shortened bare
half helices that resemble the C-actin structure adjacent to cofilin clusters on the PE side. This
comparison was made when actin filaments were incubated with the same cofilin concentration
(i.e., 300 nM) in the F1 buffer containing 1 mM ADP + 5 U/ml hexokinase + 10 mM glucose or

1 mM ATP + 10 mM Pi, resulting in filamentous protomers representing F-ADP-actin or F-
ADP.Pi-actin structures. The growth rates (mean ± SD) were measured for 22 shortened bare
half helices in F-ADP-actin (10.0 ± 8.3 molecules/HHP/s) and 28 shortened bare half helices in
F-ADP.Pi-actin (8.3 ± 7.7 molecules/HHP/s). Using the two-population *t-test*, the results showed
that the difference was not significant ($p = 0.47846$) at the $p \leq 0.05$ level.
(**C, D**) Histograms illustrate the distribution of the HHP (mean ± SD) in control actin filaments
for three conditions: (i) phalloidin-stabilized F-actin (37.0 ± 2.3 nm), (ii) F-ADP-actin ($36.9 \pm$
3.8 nm), and (iii) F-ADP.Pi-actin (37.0 ± 2.8 nm), with each condition having 2487 data points.
For comparison, single Gaussian fits in D were integrated and displayed in C. The inset within C
exhibited fractions of the naturally supertwisted half helices with HHPs shorter than 30 nm.
Utilizing one-way ANOVA with a significance level of $p \leq 0.05$, there were significant
differences observed between (i) and (ii) ($p = 0.00837$, $F = 6.95754$) as well as between (ii) and
(iii) ($p = 0.04080$, $F = 4.18633$), primarily attributable to the statistical difference in the variance
of HHP between condition (ii) vs. (i) and (iii). The difference observed between (i) and (iii) was
not significant ($p = 0.58272$, $F = 0.30189$).

Figure 10 The hypothesis regarding the actin structure-dependent preferential cooperative binding of cofilin. For simplicity, a single actin protofilament comprising 6 protomers labelled from -2 to 8 is shown, both without and with cofilin. The black and red arrows represent stable bonds and unstable bonds between domains of two adjacent actin protomers in the same strand, respectively. The broken red arrows indicate completely disrupted bonds between domains of two adjacent actin molecules. Additionally, broken red and purple lines show positions sensitive to being severed, with the purple lines indicating a stronger severing activity.

(**A**) The conformational transition between G-actin, F-actin (bound with ADP.Pi, ADP), and C-actin structures. In the lower model of C-actin, the binding of a single cofilin molecule to an actin filament induces a small movement of SD2, causing a rotational shift of approximately 5° in the OD relative to the ID. In the upper one, when the cofilin clusters bind, they induce a larger rotation of approximately 10° in the OD (Galkin et al., 2011; Tanaka et al., 2018).

(**B**) A single protofilament of an actin filament. Upon polymerization, OD rotates relative to the ID to close nucleotide binding cleft and untwist the OD to flatten actin molecule. The orange color corresponds to OD (SD1+SD2), while green corresponds to ID (SD3+SD4) (Oda et al., 2009).

(**C**) A single protofilament of an actin filament bound to a single cofilin (Huehn et al., 2020; Tanaka et al., 2018).

(D) A single protofilament of an actin filament bound to cofilin clusters. Actin conformational changes induced by cofilin clusters are local and limited to protomers directly contacting bound cofilin. The cofilin bindings accelerate a rapid transition from F-ATP-actin to F-ADP.Pi-actin and F-ADP-actin states of protomers on the PE side. Cofilin tunes adjacent protomers, resembling F-ADP-actin structure on the PE side, to expand the clusters The cooperative binding of cofilin requires a critical size of a cofilin cluster set at 2. Notably, the severing of the actin filament is predominantly observed at the junction between cofilactin and bare actin (Huehn et al., 2020; Suarez et al., 2011; Tanaka et al., 2018).

(E-F) A single protofilament of an actin filament bound to cofilin clusters in the authors' previous study (Ngo et al., 2015) and in the current study. Our hypothesis supports that actin conformational changes induced by cofilin clusters are not local and not limited to protomers directly contacting bound cofilin (C-actin) on the PE side. Cofilin clusters have the ability to shorten a half helix by reducing the number of protomers. Additionally, cofilin cluster not only has the capacity to convert F-actin to C-actin within cofilactin region but also allosterically expands the C-actin structure from the bound to unbound region on the PE side (**E**). The MAD_c in the undecorated C-actin-like structure region are elongated compared to MAD_f in F-actin region. The shortening of half helices and the elongated MAD_c play a pivotal role in facilitating the preferential cooperative binding of cofilin to actin protomers via F-actin to C-actin transition pathway. If cofilin does not promptly attach to protomers in shortened bare half helix region (as depicted in E-F), the protomers within C-actin-like structure regions can readily open the nucleotide binding clefts and transition back to G-actin state via C-actin to G-actin transition, ultimately severing the filament at or near boundaries (broken lines).

(G) HS-AFM measurements of MAD values measured in half helices of filaments representing F-actin (MAD_f) and undecorated C-actin-like structure (MAD_c) adjacent to cofilin cluster on the PE side.

(H) A hypothesis describing how the function of cofilin is regulated by the torsional flexibility during helical twisting of actin filaments. The F-ADP-actin has increased the fractions of naturally supertwisted helical structures and the torsional flexibility during helical twisting, thereby creating some proposed C-actin-like structures to facilitate the initial binding and cluster expansion of cofilin, compared to F-ADP.Pi-actin. To emphasize the inhibitory effect of Pi on the initial binding of cofilin, we depict a simplified model of F-ADP.Pi-actin, featuring very infrequent structures resembling C-actin. As a result, the initial binding of cofilin and growing of critical cluster set at 2 to 4 cofilin molecules are more challenging compared to F-ADP-actin.

Figure supplement 1 A closer examination of PC1 and PC2 in the PCA clusters of F-actin
structures.

Within these PCA clusters of F-actin, similar to the inset of **Figure 1A**, there is a larger presence
of C-actin-like structures in F-ADP-actin compared to F-ADP.Pi-actin. However, the magnified
views of the F-actin's PCA clusters also highlight structural resemblance between F-ADP-actin
and F-ADP.Pi-actin, as recently elucidated by (Reynolds et al., 2022) (**A**) and (Oosterheert et al.,
2022) (**B**). Significantly, the F-ADP-actin structure (8D13) and F-ADP.Pi-actin structure (8D14)
exhibit remarkably similar PC2 values. Furthermore, the PC2 results demonstrate notable
similarities between Mg^{2+}-F-ADP-actin (8A2T) and Mg^{2+}-F-ADP.Pi-actin (8A2S), as well as
Mg^{2+}-F-ADP.BeF_3-actin (8A2R). These diverse PDB structures may reflect the intrinsic
dynamics and polymorphism in actin structure. Regardless of whether a structure is derived from
crystallography, Cryo-EM, or NMR data, it is more accurate to refer to it as a structure—one of
many possible conformations (Lane, 2023; Miller and Phillips, 2021).

1645

Figure supplement 2 Pseudo-AFM images featuring long filaments representing F-actin and C-
actin structures. It is noted that both filamentous structures share the same length and contain an
equal number of actin protomers, specifically 62 protomers. However, the filament representing
C-actin exhibits a greater number of half helices than F-actin. This difference can be attributed to
the decrease in the number of actin protomers per HHP in C-actin structure.
(**A**) A pseudo-AFM image featuring a long filament representing F-actin structure constructed
using the 6VAU. The scale bar represents 25 nm.
(**B**) Longitudinal section profiles of the middle half helices in panel A were used to identify the
HHPs (distances between major peaks marked in red, reported in nm) and the numbers of actin
protomer pairs per HHP (as counted major and small peaks marked in red and green within each
HHP subtracted by 0.5). The offset values of the height profile were set to zero by referring to
virtual mica surface, where filament was placed in the pseudo-AFM images. The MADs were
computed and shown in brackets. The HHP, the number of protomers per HHP, and MAD were
shown accordingly.
(**C**) A side-view of the long actin filament imaged in panel A. To enhance visibility of protomer
pairs, one of the chains was depicted as translucent.
(**D-F**) Panels D-F follow a similar format as panels A-C but display the results obtained for a
long filament representing C-actin structure (cofilins are removed from initial cofilactin structure
reported in 6VAO). The HHP, the number of protomers per HHP, and axial distance (AD) were
shown in E accordingly.
(**G-I**) Panels G-I also followed a similar format as panels A-C, but showcased the results
obtained for a long cofilactin filament (constructed from 6VAO). Accurately detecting actin
pairs in this case was challenging due to cofilin decoration and the tip-sample dilation effect, as
explained by (Ando, 2022).

1692
Figure supplement 3 Exploring the dynamics of axial distance (AD) between two adjacent
protomers within the same actin filaments. Actin filaments were gently immobilized on a lipid
membrane composed of 1,2-dipalmitoyl-sn-glycero-3-phosphocholine (DPPC) and 1,2-
dipalmitoyl-3-trimethylammonium-propane (DPTAP) (90/10 wt%). The still AFM images are
shown after processing with Laplacian of Gaussian filter. Scale bars: 25 nm.
(**A, B**) Longitudinal section profiles and distributions of AD between two adjacent protomers
within actin segments bound with ADP (F-ADP-actin), captured at high temporal and spatial
resolutions using a Quantum AC-10-Super-Sharp high density carbon tip (Nanotools, Germany).
F-ADP-actin segments were prepared as described in the Methods section and imaged in F1
buffer containing 1 mM ADP, 5 U/ml hexokinase and 10 mM glucose.
(**C**) Longitudinal section profiles and distributions of AD between two adjacent protomers within
actin segments bound with ADP.Pi (F-ADP.Pi-actin), captured at high temporal and spatial
resolutions using the same cantilever. F-ADP.Pi-actin segments were prepared as described in
the Methods section and imaged in F1 buffer containing 1 mM ATP and 10 mM Pi
(K_2HPO_4/KH_2PO_4, pH 6.8 buffer).

(**D**) The examples of the ACF plots display the autocorrelation coefficients on the y-axis against
the time lags on the x-axis. The plotted data was fitted using a second-order exponential decay
function with the time constants of (t_1, t_2), represented by the red curve. The t_1 and t_2 values are
associated with first and second time decay of AD between two adjacent actin protomers within
F-ADP-actin and F-ADP.Pi-actin segments, recorded experimentally at high temporal resolution
using HS-AFM. A two-population *t-test* is employed for statistical analysis to evaluate the
significant differences (*p*-values) between the two populations (mean ± SD). We implemented
running frame averages (3 frames) in the ACF analyses. See **Table supplement 5**.

1754

Figure supplement 4 A schematic overview of the spatial ACF analysis.

(**A**) An HS-AFM image is depicted, with a line profile drawn along an actin filament.

(**B, C**) The height profiles before and after applying the bandpass filter (BPF), respectively.

(**D, E**) The spatial ACF or envelope curves before and after BPF processing and after applying

the Hilbert transform are displayed, respectively.

(**F, G**) Illustrating the surface topography of the actin protomers' heights along filaments, which

were used in making the spatial ACF curve (as depicted by the envelope curve). The green

waves in (**F**) represent flatter slopes in the heights between protomers, while the red waves in

(**G**) depict the steeper slopes in the heights between protomers. In theory, the envelope curve

(green) versus distance in (**F**) should linearly decrease. In contrast, the envelope curve (red)

versus distance in (**G**) shows a more pronounced decrease. During the actual analysis, the

selected space intervals (ΔD) for measuring the degree to which height values near each other in

space are similar to one another in F-actin and hybrid cofilactin/bare actin segments were 0.4 nm

and 0.3 nm, respectively. The measurement process involved shifting these space intervals from

left to right and up to down, with increasing interval values from the initial position (0 ΔD). P1-

P7 denote the peak positions of individual protomers.

1771
Figure supplement 5 Analysis of spatial ACF curves
(**A**) Spatial ACF curves were depicted for canonical actin filaments. This analysis was conducted
at different time points (the imaging rate of 0.5 s per frame and a total of 50 frames for each
filament (F1-F5)). The still AFM image is shown after processing with Laplacian of Gaussian
filter. Scale bar: 25 nm. Related to **Figure 4C-D**.
(**B**) Spatial ACF curves were depicted for a hybrid cofilactin/bare actin filament, featuring a bare
half helix near a cofilin cluster on either the PE or BE side. This analysis was conducted at
different time points (the imaging rate of 0.5 s per frame and a total of 50 frames for each half
helix on the PE or BE side). Based on our previous study (Ngo et al., 2015), we judged that the
bare half helix with the shorter HHP adjacent to the cofilin cluster is on the PE side and that the

normal HHP is located on the BE side. The still AFM images are shown after processing with
Laplacian of Gaussian filter. Scale bars: 25 nm. Related to **Figure 6C**.

1828
1829 **Figure supplement 6** The propagation of the supertwisted half helices induced by cofilin
1830 clusters results in the shortening of a bare half helix beyond the cofilactin/bare actin boundary.

The HHPs and peak heights were determined using methods similar to our previous reports (Hayakawa et al., 2022; Ngo et al., 2015). Scale bars: 25 nm. Arrowheads indicated the peaks of cofilin clusters. h_0 and h_1 represented the peak heights in two consecutive HHPs of bare actin and cofilactin segments. The offset values were set to zero with reference to the height of the lowest position along the longitudinal section lines. The height and width differences were utilized to determine the bare actin and cofilactin segments. The bare actin, cofilactin, and hybrid cofilactin/bare actin filaments, priorly made in solution, were strongly immobilized on lipid membrane composed of DPPC/DPTAP (50/50 wt%) and imaged in F1 buffer containing 0.5 mM ATP. See Methods.

(**A**) A still AFM image presents a normal actin filament, accompanied by a longitudinal section profile across two consecutive half helices to measure the HHP1 and HHP2 and the number of actin protomer pairs per HHP accordingly. The MADs were computed from AFM analysis and shown.

(**B**) A still AFM image depicts a cofilactin filament, accompanied by a longitudinal section profile across two consecutive half helices to measure the HHP3 and HHP4 and the numbers of actin protomer pairs per HHP accordingly. The MADs were calculated from our AFM analysis and shown.

(**C**) A still AFM image showcases a hybrid bare actin/cofilactin filament, accompanied by a longitudinal section profile across two consecutive half helices to measure the shortened bare HHP5 and shortened cofilactin HHP6 and the number of actin protomers per HHP. The MADs were calculated from our AFM analysis and shown. It should be emphasized that only MADs obtained in the bare regions without cofilin decoration were used to make Histograms in **Figure supplement 7**.

1877

Figure supplement 7 The analysis of the variation in helical twisting by measuring HHPs, the
number of protomer pairs per HHP, and MADs in a hybrid bare actin/cofilactin segment using
HS-AFM.

(**A**) Representative still AFM image of a hybrid bare actin/cofilactin segment in **Figure
supplement 6C** is provided. The peaks (P7, P8, P9, P10) in the raw AFM image are indicated by
white and cyan arrows for the bare actin and cofilactin regions, respectively. HHP5, HHP6, and
HHP7 and the number of protomer pairs within each HHP were measured between P7-P8, P8-
P9, and P9-P10, respectively and MAD5, MAD6, and MAD7 were computed accordingly. Scale
bars: 25 nm.

(**B**) The variation of HHPs, the number of protomer pairs per HHP, and MADs in shortened bare
and cofilactin regions were examined over time.

(**C**) A comparison between MAD5 (5.7 ± 0.5 nm) observed in the shortened bare area and
MAD6 (5.9 ± 0.3 nm) along with MAD7 (6.1 ± 0.3 nm) within the cofilactin region was
presented. A two-population *t-test* with a sample size of n = 13 was used.

(D) The relationship between the HHP and the number of protomer pairs per HHP measured for
C-actin-like structures within the shortened bare half helix adjacent to cofilin cluster, as explored
through HS-AFM, was illustrated using black symbols. The collected data was subjected to
fitting with a linear equation ($y = 10.99 + 3.21x$, $R = 0.9975$) in the region of the shortened bare
half helix situated close to the boundary of cofilactin and bare actin. Calculated data, depicted
with red symbols, presented the correlation between the number of protomer pairs and the HHP
that was calculated as 5.5 nm multiplied by the number of protomer pairs within each HHP.
(E) The histograms depict the distribution of MADs for C-actin-like structures within the
shortened bare half helices adjacent to cofilin clusters consisting of different number of protomer
pairs.

1938

Figure Supplement 8 The fluctuation of the mean axial distance (MAD) between two adjacent
actin protomers along the same long-pitch strand of F-ADP-actin segments as a function of time.
(**A**) The longitudinal section profiles and distributions of axial distance (AD) within barely
supertwisted and untwisted half helices neighboring cofilin clusters on the PE and BE sides,
respectively. White and cyan arrows indicate the peak positions (P1-P5) within bare actin and
cofilin clusters, respectively. Notably, P4 was completely decorated with cofilin clusters,
resulting in a supertwisted HHP measured between P3 and P4 (Video 5). This supertwisted
structure persists even after cofilin clusters are removed during imaging. This structure,
containing fewer protomer pairs, closely resembles the bare HHP measured between P1 and P2
on the PE side, as opposed to the normal HHP between P4 and P5 on the BE side. The number of
protomer pairs per half helix determined by longitudinal section profiles is presented and
calculated by subtracting 0.5 from the total number of peaks (see Methods). F-ADP-actin
segments were prepared as described in the Methods section and imaged in F1 buffer containing
1 mM ADP, 5 U/ml hexokinase, 10 mM glucose, and 300 nM cofilin. Actin filaments were
gently immobilized on a lipid membrane composed of 1,2-dipalmitoyl-sn-glycero-3-
phosphocholine (DPPC) and 1,2-dipalmitoyl-3-trimethylammonium-propane (DPTAP) (90/10
wt%) and captured at high temporal and spatial resolutions using a Quantum AC-10-Super-Sharp
high-density carbon tip. There was minimal nonspecific binding of cofilin to this lipid membrane
surface. Based on our previous study (Ngo et al., 2015), we judged that the bare half helix with

the shorter HHP adjacent to the cofilin cluster is on the PE side and that the normal HHP is
located on the BE side. Scale bar: 25 nm.
(**B, C**) The MADs were calculated as described in the Methods to follow the time course of the
variation in MADs in barely supertwisted (5.3 ± 0.3, $N = 55$) on the PE side, untwisted ($5.0 \pm$
0.3, $N = 57$) on the BE side, and normally twisted (4.8 ± 0.3, $N = 129$) half helices, for
comparison between them. A two-population *t-test* was employed for statistical analysis to
evaluate significant differences (*p*-values) between the two populations (mean $\pm$ SD).

2004
Figure supplement 9 The spatial ACF curve analysis.
(**A**) The spatial ACF curves were produced for sparsely decorated cofilin-actin filaments (F1, F2,
F3), strongly attached on lipid surface, at various time points (i.e., Data 1, 2, 3). The curves were
fitted using an exponential decaying sinusoidal function (**Equation 2**). The still AFM image in
Figure 8A is shown after processing with Laplacian of Gaussian filter. Bar: 25 nm.
(**B**) The decay constant (*D1*) values were determined through a similar fitting of spatial ACF
curves, as illustrated in **Figure 4C**. This examination was conducted at different time points
(with an imaging rate of 0.5 seconds per frame and a total of 50 frames). Subsequently, the
distribution of *D1* values was analyzed, yielding the mean $\pm$ SD value of 26.8 ± 6.3 nm, based on
a sample size of 147.

Table supplement 1 List of 46 protein data bank (PDB) structures that were utilized in the PCA,
as depicted in **Figure 1**. Due to the presence of varying structures among different chains in
some PDB structures, each chain was considered as an individual structure. Notably, in the case
of 6UC4, chains E and H were combined and renamed as chain E (E&H = E), while chains F and
G were combined and renamed as chain F (F&G = F). Similarly, in the case of 6UBY, chains G
and H were combined and renamed as chain G (G&H = G).
2021

PDB ID	List of actin chains	Method	Resolution (Å)	Nucleotide	Protein bound	First author	Year
1ATN	A	X-RAY	2.8	ATP	DNASE I	KABSCH	1990
1J6Z	A	X-RAY	1.54	ADP-RHO		OTTERBEIN	2001
2BTF	A	X-RAY	2.55	ATP	PROFILIN	SCHUTT	1993
2ZWH	A	X-RAY	3.3	ADP		ODA	2009
3DAW	A	X-RAY	2.55	ATP	TWINFILIN-1	PAAVILAINEN	2008
3HBT	A	X-RAY	2.7	ATP		WANG	2010
3J0S	A, B, C, D, E, F, G, H, I, J, K, L	EM	9		COFILIN-2	GALKIN	2011
3J8I	D, E, F, G, H	EM	4.7	ADP		GALKIN	2015
3J8J	A, B, C, D, E, F, G, H, I, J, K	EM	12			GALKIN	2015
3J8K	A, B, C, D, E, F, G, H, I, J	EM	12			GALKIN	2015
3MFP	A	EM	6.6	ADP		FUJII	2010

4A7N	A, B, C, D, E	EM	8.9	ADP		BEHRMANN	2012
5ONV	A, B, C, D, E	EM	4.1	ADP		MERINO	2018
5OOC	A, B, C, D, E	EM	3.6	ADP JASP		MERINO	2018
5OOD	A, B, C, D, E	EM	3.7	ADP-Pi JASP		MERINO	2018
5OOE	A, B, C, D, E	EM	3.6	ANP		MERINO	2018
5OOF	A, B, C, D, E	EM	3.4	ADP-BeF_x		MERINO	2018
5YU8	A, B, C, D, E	EM	3.8	ADP	COFILIN-2	TANAKA	2018
6BNO	A, B, C, D, E, F, G, H	EM	5.5	ADP		GUREL	2017
6BNU	A, B, C, D, E, F, G, H	EM	7.5			GUREL	2017
6DJM	A, B, C, D	EM	3.1	AMPPNP		CHOU	2019
6DJN	A, B, C, D	EM	3.1	ADP-Pi		CHOU	2019
6DJO	A, B, C, D	EM	3.6	ADP		CHOU	2019
6FHL	A, B, C, D, E	EM	3.3	ADP-Pi		MERINO	2018
6KLL	A, B, C, D	EM	3	ADP		ODA	2020
6KLN	A, B, C, D	EM	3.4	ADP		ODA	2020
6UBY	A, B, C, D, E, F, G	EM	7.5	ADP	COFILIN-1	HUEHN	2020

6UC0	A, B, C, D, E, F, G	EM	7.5	ADP	COFILIN-1	HUEHN	2020
6UC4	A, B, C, D, E, F, J, K, L	EM	9.2	ADP	COFILIN-1	HUEHN	2020
6VAO	D, A, B, C, E	EM	3.4	ADP	COFILIN-1	HUEHN	2020
6VAU	B, A, C, D, E	EM	3.5	ADP		HUEHN	2020
7Q8B	A, B, C, D, E	EM	3.3	ADP-Pi		KOTILA	2022
7Q8C	A, B, C, D, E	EM	2.72	ADP		KOTILA	2022
7Q8S	A, C, D, E, F	EM	3.4	ADP	ADF/COFILIN	KOTILA	2022
8A2R	C, A, B, D, E	EM	2.17	ADP-BeF$_3$		OOSTERHEERT	2022
8A2S	C, A, B, D, E	EM	2.22	ADP-Pi		OOSTERHEERT	2022
8A2T	C, A, B, D, E	EM	2.24	ADP		OOSTERHEERT	2022
8A2U	C, A, B, D, E	EM	2.21	ADP-BeF$_3$		OOSTERHEERT	2022
8A2Y	C, A, B, D, E	EM	2.15	ADP-Pi		OOSTERHEERT	2022
8A2Z	C, A, B, D, E	EM	2.15	ADP		OOSTERHEERT	2022
8D13	A, B, C	EM	2.43	ADP		REYNOLDS	2022
8D14	A, B, C	EM	2.51	ADP-Pi		REYNOLDS	2022

8D15	A, B, C, D, E, F, G	EM	3.61	ADP		REYNOLDS	2022
8D16	A, B, C, D, E, F, G	EM	3.71	ADP-Pi		REYNOLDS	2022
8D17	A, B, C, D, E, F, G	EM	3.69	ADP		REYNOLDS	2022
8D18	A, B, C, D, E, F, G	EM	3.66	ADP		REYNOLDS	2022

Table supplement 2 The lists of chains and matrices utilized to build the long filaments

PDB ID	Chains	RMSD (Å)	Transformation matrix as Euler angles		
6VAU	BA	0.000497	[166.588953	-2.1209466e-01	-2.62674783e-02]
8A2S	ED	0.000704	[-166.93535314	-10.63461097	-5.38436149]
8A2T	AB	0.000704	[-167.39029871	-13.65820025	-6.3200023]
3MFP	AB	0.000398	[166.656064	2.13617418e-05	7.68141187e-05]
6UC4	EF	0.619515	[164.58920564	-0.207997407	-0.80958005]
6VAO	CA	0.000699	[162.515987	-4.24094412e-02	-1.82361154e-01]
5YU8	ED	0.000561	[-162.100031	-4.00777476e-05	4.03030046e-05]
3J0S	KL	0.001134	[-162.12014856	-0.17228856	-0.18642638]
*For cofilactin filament					
6VAO*	CAJF	0.000699	[162.515987	-4.24094412e-02	-1.82361154e-01]

Table supplement 3 Variation during helical twisting measured by HHPs, the number of
protomer pairs per HHP, and the MAD_f observed for protomers representing F-actin structures
within long bare actin filaments, as probed by HS-AFM. The total number of HHPs observed
was 2226. The total number of protomer pairs used for estimating MAD_f was calculated by
multiplying the number of protomer pairs per HHP by the number of HHPs.

The number of protomer pairs per HHP	HHPs (nm)	MAD_f (nm)	The number of HHPs (proportion)	The total number of protomer pairs
4.5	28.6 ± 1.4	6.3 ± 0.3	5 (0.2%)	22.5
5.5	29.3 ± 1.8	5.3 ± 0.3	182 (8.2%)	1001
6.5	33.1 ± 1.3	5.1 ± 0.2	1355 (60.9%)	8807.5
7.5	37.2 ± 2.0	5.0 ± 0.3	624 (28.0%)	4680
8.5	39.1 ± 2.4	4.6 ± 0.3	60 (2.7%)	510

Table supplement 4 Variation during helical twisting measured by HHPs, the number of
protomer pairs per HHP, and the MAD_c observed for protomers resembling C-actin structures
within shortened bare half helices adjacent to the cofilactin/bare actin boundary, as probed by
HS-AFM. The total number of HHPs observed was 34. The total number of protomer pairs used
for estimating MAD_c was calculated by multiplying the number of protomer pairs by the number
of HHPs.
2129

The number of protomer pairs per HHP	HHP (nm)	MAD_c (nm)	The number of HHPs (proportion)	The total number of protomer pairs
3.5	22.2 ± 0.2	6.3 ± 0.1	2 (5.9%)	7
4.5	26.1 ± 1.4	5.8 ± 0.3	21 (61.8%)	94.5
5.5	28.5 ± 0.9	5.2 ± 0.2	9 (32.4%)	49.5

2161 **Table supplement 5** Quantifying time constants (t_1, t_2) obtained from fitting individual ACF
2162 curves. The t_1 and t_2 values are associated with first and second time constants of axial distance
2163 (AD) between two adjacent actin protomers within F-ADP-actin and F-ADP.Pi-actin segments.
2164 These time constants reflect the time decay in the autocorrelation of AD, recorded
2165 experimentally at high temporal resolution using HS-AFM. Related to **Figure supplement 3**.
2166

Experimental conditions	t_1 (s)	t_2 (s)
F-ADP-actin	0.4 ± 0.2	0.5 ± 0.7
(200 ms/frame)	(N = 82)	(N = 80)
F-ADP-actin	0.3 ± 0.3	1.2 ± 1.8
(100 ms/frame)	(N = 78)	(N = 78)
F-ADP.Pi-actin	0.2 ± 0.3	0.5 ± 1.1
(100 ms/frame)	(N = 152)	(N = 141)

2200 **Table supplement 6** The correlation between the time it takes to shorten a bare HHP beyond the
2201 cofilactin/bare actin boundary and to form a newly matured cofilin cluster in a pre-shortened half
2202 helix. The lag time (t) represents the duration required for a newly saturated cofilin cluster to
2203 bind and decorate onto a bare half helix adjacent to the preformed cluster after this bare half
2204 helix has been shortened by approximately 25%. This table is related to **Figure 7**.
2205

The time required to form a newly matured cofilin cluster in a pre-shortened half helix		The duration needed to shorten a bare half helix beyond the cofilactin/bare actin boundary				
		P1 – P2	P2 – P3	P3 – P4	P4 – P5	P5 – P6
P1	0 s					
P2	0 s	0 s				
P3	5.0 s		2.0 s ($t = 3.0$ s)			
P4	36 s			30.5 s ($t = 5.5$ s)		
P5	54.5 s				54 s ($t = 0.5$ s)	
P6	58.5 s					58.5 s ($t = \sim 0$ s)

Table supplement 7 A summary of peak heights and HHPs obtained from the histograms in **Figure 8**. Single and double Gaussian fitting methods were applied to the data. The significant difference between the peaks in the Gaussian fitting was statistically evaluated and indicated using an *F-test*. "ND" indicates cases where the information was not determined.

	Peak Height (nm)		Half Helical Pitch (nm)		Frequency
	Peak 1 (proportion) [*F-test, p*]	Peak 2 (proportion) [*F-test, p*]	Peak 1 (proportion) [*F-test, p*]	Peak 2 (proportion) [*F-test, p*]	
(A) 2000 nM cofilin, 0.5 mM ATP/ on DPPC/DPTAP (75/25 wt%)	8.2 ± 0.8 (100%)	ND	34.3 ± 2.6 (100%)	ND	Height: 1930 HHP: 1556
(B) 300 nM cofilin, 0.5 mM ATP/ on DPPC/DPTAP (90/10 wt%)	9.2 ± 0.4 (47.0%) [0.0204]	10.5 ± 0.7 (53.0%) [0.0204]	28.5 ± 3.0 (55.1%) [0.0135]	37.4 ± 3.6 (44.9%) [0.0135]	Height: 2212 HHP: 1780
(C) 300 nM cofilin, 1 mM ADP/ on DPPC/DPTAP (90/10 wt%)	8.4 ± 0.5 (54.8%) [0.0081]	9.9 ± 0.7 (45.2%) [0.0081]	30.1 ± 3.8 (70.3%) [0.0195]	37.8 ± 2.9 (29.7%) [0.0195]	Height: 3183 HHP: 2779
(D) 75 nM cofilin, 1 mM ATP, 10 mM Pi/ on DPPC/DPTAP (90/10 wt%)	8.3 ± 0.7 (55.3%) [1.48×10^{-6}]	11.0 ± 1.0 (44.7%) [1.48×10^{-6}]	30.3 ± 2.9 (25.2%) [0.0002]	38.7 ± 3.6 (74.8%) [0.0002]	Height: 2316 HHP: 1865

Legend of videos

Video 1 Possible transition pathways from G-actin to C-actin (constructed by +PC1) and C-actin to F-actin (constructed by +PC2), and vice versa, constructed by -PC1 and -PC2, respectively. The transition from G-actin to F-actin ultimately requires both + PC1 + PC2. Related to **Figure 1**.

Video 2 HS-AFM imaging to investigate the dynamic structures during helical twisting by measuring the variations of HHPs, the number of actin protomers per HHP, and MADs in F-actin structures within long actin filaments. Our results provided strong evidence supporting the variable helical twisting and dynamics in MADs within actin filaments. However, the results contrast the existing rigid and inflexible depiction of helical twists in Cryo-EM and X-ray diffraction structures. Actin filaments were gently immobilized on lipid membrane composed of 1,2-dipalmitoyl-sn-glycero-3-phosphocholine (DPPC) and 1,2-dipalmitoyl-3-trimethylammonium-propane (DPTAP) (90/10 wt%). Bars: 25 nm. Imaging rate: 2 frames per second (fps). Video plays at 5 fps. Related to **Figure 3**.

Video 3 HS-AFM imaging to investigate the dynamic structures during helical twisting by measuring the variability of MADs in F-ADP-actin and F-ADP.Pi-actin structures within long actin filaments. Our findings provided strong evidence supporting the dynamics in MADs within these actin filaments, though the current temporal resolution could not distinguish a significant difference between F-ADP-actin and F-ADP.Pi-actin, even at high imaging rates. Actin filaments were gently immobilized on lipid membrane composed of 1,2-dipalmitoyl-sn-glycero-3-phosphocholine (DPPC) and 1,2-dipalmitoyl-3-trimethylammonium-propane (DPTAP) (90/10 wt%) and imaged in F1 buffer containing either 1 mM ADP + 5 U/ml hexokinase + 10 mM glucose or 1 mM ATP + 10 mM Pi. Bars: 25 nm. Imaging was conducted at rates of 5 and 10 fps for F-ADP-actin and at 10 fps for F-ADP.Pi-actin. The video is played at 5 fps. Related to **Figure supplement 3**.

Video 4 HS-AFM imaging to investigate the dynamic structures during helical twisting by measuring the variations of HHPs, the number of actin protomers per HHP, and MADs in the normal and shorter half helices neighboring a cofilin cluster on the BE and PE sides. Bars: 25 nm. Imaging rate: 2 fps. Video plays at 5 fps. Related to **Figure 5**.

Video 5 HS-AFM imaging to investigate the dynamic structures during helical twisting by measuring the variations of MADs in the barely supertwisted and untwisted half helices neighboring a cofilin cluster on the PE and BE sides, respectively, as well as in the normally twisted half helices in the same images. Actin filaments were imaged in an F1 buffer containing 300 nM cofilin and 1 mM ADP, 5 U/ml hexokinase and 10 mM glucose. There was minimal nonspecific binding of cofilin to this lipid membrane surface. Bars: 25 nm. Imaging rate: 2 fps. Video plays at 5 fps. Related to **Figure supplement 8**.

Video 6 HS-AFM imaging to demonstrate the preferentially cooperative binding of cofilin toward a pre-shortened bare half helix resembling C-actin structures along actin filaments. Even with 10 mM Pi, the cofilin cluster could supertwist the half helices in the actin filaments once a critical cluster was established. Bars: 25 nm. Imaging rate: 2 fps. Video plays at 5 fps. Related to **Figure 7**.

2298

Video 7 Cooperative binding of cofilins along actin filaments was greatly inhibited when actin
filaments were strongly immobilized on lipid membrane composed of 1,2-dipalmitoyl-sn-
glycero-3-phosphocholine (DPPC) and 1,2-dipalmitoyl-3-trimethylammonium-propane
(DPTAP) (75/25 wt%). The strong immobilization hindered the torsional motion of actin
filaments during helical twisting, resulting in suppressing cofilin binding and cluster expansion.
Actin filaments were imaged in an F1 buffer containing 2000 nM cofilin and 0.5 mM ATP. Time
label denotes the elapsed time after addition of cofilin. Bars: 25 nm. Imaging rate: 2 fps. Video
plays at 10 fps. Related to **Figure 8A**.

2307

Video 8 Cooperative binding of cofilins along actin filaments was normal when actin filaments
were loosely immobilized on lipid membrane composed of 1,2-dipalmitoyl-sn-glycero-3-
phosphocholine (DPPC) and 1,2-dipalmitoyl-3-trimethylammonium-propane (DPTAP) (90/10
wt%). Actin filaments were imaged in an F1 buffer containing 300 nM cofilin and 0.5 mM ATP.
Time label denotes the elapsed time after addition of cofilin. Bars: 25 nm. Imaging rate: 2 fps.
Video plays at 10 fps. Related to **Figure 8B**.

2314

Video 9 Cooperative binding of cofilins along actin filaments was normal when actin filaments
were loosely immobilized on lipid membrane composed of 1,2-dipalmitoyl-sn-glycero-3-
phosphocholine (DPPC) and 1,2-dipalmitoyl-3-trimethylammonium-propane (DPTAP) (90/10
wt%). Actin filaments were imaged in an F1 buffer containing 300 nM cofilin, 1 mM ADP, 5
U/ml hexokinase, and 10 mM glucose. Time label denotes the elapsed time after addition of
cofilin. Bars: 25 nm. Imaging rate: 2 fps. Video plays at 10 fps. Related to **Figure 8C**.

2321

Video 10 Growth of cofilin cluster along actin filaments was normal when actin filaments were
loosely immobilized on lipid membrane composed of 1,2-dipalmitoyl-sn-glycero-3-
phosphocholine (DPPC) and 1,2-dipalmitoyl-3-trimethylammonium-propane (DPTAP) (90/10
wt%). Even with 10 mM Pi, the cofilin cluster could supertwist the half helices in the actin
filaments once a critical cluster was established. Actin filaments were imaged in an F1 buffer
containing 75 nM cofilin and 1 mM ATP + 10 mM Pi (See Methods). Time label denotes the
elapsed time after addition of cofilin. Bars: 25 nm. Imaging rate: 2 fps. Video plays at 10 fps.
Related to **Figure 8D**.