Single-cell transcriptional changes across the blood stages of artemisinin-resistant K13^{580Y} *Plasmodium falciparum* upon dihydroartemisinin exposure

Cliff I. Oduor[1^], Sean V. Connelly[2^], Clark Cunningham[2], Ian Petrow[1,3], Nazrin Rustamzade[1], Jenna Zuromski[1], Deborah M. Chin[1], Christian P. Nixon[1], Jonathan D. Kurtis[1,3], Jonathan J. Juliano[2,4,5,6+], Jeffrey A. Bailey[1+*]

1. Department of Pathology and Laboratory Medicine, Brown University, Providence, RI, USA

2. Division of Infectious Diseases, Department of Medicine, University of North Carolina, Chapel Hill, NC, USA

3. Center for International Health Research, Providence, RI, USA

4. Curriculum in Genetics and Molecular Biology, School of Medicine, University of North Carolina, Chapel Hill, NC, USA

5. Department of Epidemiology, Gillings School of Global Public Health, University of North Carolina, Chapel Hill, NC, USA

6. Institute for Global Health and Infectious Diseases, University of North Carolina, Chapel Hill, NC, USA

^: Co-first authors

+: Co-senior authors

*: Corresponding author

Jonathan Juliano

110 Mason Farm Rd.

Chapel Hill, NC 27599

919-843-5065

jjuliano@med.unc.edu

ABSTRACT

Artemisinin-based therapies have been central to malaria control, but partial resistance in *Plasmodium falciparum*, driven by mutations in the Kelch13 (K13) gene, threatens these gains. To investigate the molecular basis of this resistance, we applied single-cell RNA sequencing to coisogenic parasite lines, K13 wild-type (K13-C580) and the artemisinin-resistant mutant (K13-580Y), following a 6 hour pulse of dihydroartemisinin (DHA). This approach enabled high-resolution profiling across all intraerythrocytic stages. Both lines exhibited stage-specific transcriptional responses, with the most pronounced changes in ring and trophozoite stages. These included downregulation of metabolic genes and upregulation of stress response pathways, including the unfolded protein response and protein trafficking. Transcriptional signatures consistent with a dormancy-like state were observed, particularly in the downregulation of SERA family genes. Notably, K13-580Y parasites showed a more consistent and robust transcriptional adaptation across stages, suggesting enhanced survival under drug-induced stress. Using MELD, a computational framework for quantifying transcriptional perturbation, we found that K13-580Y parasites exhibited higher DHA likelihood scores across the asexual cycle, indicating broad and sustained transcriptional shifts in response to treatment. Additionally, increased expression of surface proteins such as PfEMP1 and GARP was observed in K13-580Y parasites at baseline and following DHA exposure. Functional assays confirmed that anti-GARP antibodies retained efficacy against both lines, supporting its potential as a therapeutic target. These findings provide a comprehensive view of the cellular responses that may facilitate artemisinin resistance, identify molecular features of dormancy and pathogenesis, and highlight surface proteins like GARP and PfEMP1 as promising intervention targets. This work underscores the power of single-cell approaches to dissect drug responses and guide strategies to overcome resistance in malaria parasites.

AUTHOR SUMMARY (150 words)

Single-cell RNA sequencing enables high-resolution analysis of *Plasmodium falciparum* responses to antimalarial treatment across developmental stages. By comparing artemisinin-sensitive and -resistant coisogenic lines, we identified shared and distinct transcriptional responses to dihydroartemisinin (DHA), including metabolic arrest, stress adaptation, and signatures of dormancy. Using MELD, a computational tool that quantifies transcriptional perturbation, we found that resistant K13-580Y parasites exhibited broader and more sustained shifts in gene expression across the asexual cycle. This suggests a pervasive cellular response to drug pressure in resistant parasites. Notably, increased expression of surface proteins such as PfEMP1 and GARP was observed in K13-580Y parasites, and anti-GARP antibodies retained efficacy across genotypes. These findings highlight potential molecular vulnerabilities and therapeutic targets in artemisinin-resistant parasites and demonstrate the power of single-cell approaches to dissect drug response heterogeneity.

Keywords: *Plasmodium falciparum*, drug resistance, artemisinin, dihydroartemisinin, Kelch13, Single-cell, RNA-seq, Transcriptome.

INTRODUCTION

Malaria remains one of the most common infectious diseases in the world and one of the greatest global health concerns, with an estimated 247 million reported malaria cases and 619,000 malaria-related deaths in 2022 of which 76% were in children under five years (1). *Plasmodium falciparum* (*Pf*) accounts for 99.7% of estimated malaria cases and virtually all the deaths in Africa (1). The World Health Organization recommends artemisinin-based combination therapies (ACTs) for the treatment of uncomplicated falciparum malaria. These combinations rely on the fast-acting properties of the artemisinin derivative to rapidly clear parasites with the longer-acting partner drugs clearing residual parasites, reducing the frequent observation of recrudescent infections when artemisinin drugs are used alone (2). Effective treatment and management of *P. falciparum* is now endangered by emerging partial resistance to artemisinin with multiple validated mutations in Africa, primarily in Rwanda, Uganda, Ethiopia, and Tanzania (3–9). The emergence of artemisinin partial resistance may precede development of partner drug resistance, jeopardizing the considerable gains made against malaria over the past decade. Given the vast majority of disease burden is in Africa, a public health disaster could arise if artemisinin partial resistance spreads across the continent (10, 11).

Resistance to artemisinin (ART) derivatives first emerged in western Cambodia (12). Clinical artemisinin partial resistance (ART-R) is expressed as a reduced rate of parasite clearance (>5 hr parasite half life), residual parasitemia 3 days after treatment, or increase rate of recrudescence (13). This resistance biologically manifests as a decreased susceptibility of ring-stage parasites to ART and is highly attributed to nonsynonymous mutations in the ß-propeller domain of the Kelch13 (K13) protein (7, 14, 15). The identification of *K13* (PF3D7_1343700) gene mutations as a marker of resistance has provided the impetus for a comprehensive series of studies that track and analyze the distribution of K13 mutations in malaria-endemic regions (7, 16). Over 100 different K13 mutations have been identified (17), but thirteen are considered validated resistance markers and 10 are considered candidate resistance markers (18). The K13[580Y] mutation is the most dominant in Southeast Asia (19–21). In Eastern Africa, the mutations K13[561H,675V,469Y,622I] have emerged and are now found in multiple countries (22). The role of K13 mutations in conferring artemisinin partial resistance has been investigated through selection experiments and gene editing (15, 23). While the

degree of resistance is modulated by the genetic background (24), K13 mutants exhibit about tenfold higher

residual viability than their genetically matched wild-type parasites when exposed to dihydroartemisinin (DHA)

at the early ring stage (15), sparking interest in K13's role in artemisinin treatment response.

K13 protein function remains to be fully understood. It consists of 726 amino acids and exhibits sequence

similarity to a class of Kelch/BTB/POZ ubiquitination adaptors (14, 25). K13 protein functions within the

endocytosis pathway required for hemoglobin uptake from the host cell (23), which is critical for parasite

survival and activation of the endoperoxide bridge of artemisinin (26). Resistant K13 mutations in the propeller

are thought to partially reduce normal function, which in turn is thought to diminish hemoglobin endocytosis.

Reduced hemoglobin uptake limits the availability of hemoglobin-derived Fe2+-heme necessary for drug

activation to kill the rings via alkylation of proximal proteins, leading to reduced ART killing (25, 26).

In this study, we explored the single cell transcriptomic profile of K13[580Y] (CamWT_C580Y, MRA-1251) and

K13[C580] (CamWT, MRA-1250) *P. falciparum* coisogenic lines following DHA exposure using a well based single

cell platform, SeqWell (27, 28). The well based single cell platform Smart-seq2 has successfully been used in

the past to study *Plasmodium berghei*, allowing for detection of cells across the complete life cycle, including

the extracellular merozoite stage (29). Single cell transcriptomic profile enabled us to measure transcriptional

effects of DHA treatment effect on plasmodium cell cycle stages. We observed parasite stage specific gene

expression profiles and also potential vulnerabilities of the parasite's response to DHA treatment, such as

elevated GARP surface protein expression, that can be a logical therapeutic target for effective combination

therapies in the face of ART-R.

 RESULTS

 Stage-specific differential expression between untreated K13^{580Y} and K13^{C580}.

To study the transcriptomic profiles of *P. falciparum* cells in response to DHA exposure, we isolated coisogenic

non-synchronized K13^{C580} and K13^{580Y} parasites after 48hrs in culture, and profiled the single cell

transcriptome at different timepoints of treatment with DHA (**Figure 1** and **2**). The transcriptional differences

between K13^{580Y} and K13^{C580} were determined in the absence of the drug by comparing the gene expression

profile of the K13^{580Y} untreated control to K13^{C580} untreated control (**Figure 3**). The genetic difference between

K13^{580Y} and K13^{C580} was verified using molecular inversion probes targeting multiple drug resistant markers

(**Supplemental Methods** and **Supplementary Results**). The coisogenic parasites identities were consistent

with previous genotyping, only differing in K13 (Figure 2C). *K13* gene expression was observed to be highly

expressed in the ring stages of both the K13^{C580} and K13^{580Y} parasites compared to all the other stages (**Figure**

3). Despite the fact that the K13^{580Y} mutation is not known to affect transcription efficiency of the gene, *K13*

gene expression was significantly lower (log2FC=-0.92, *p*-value=2.13e-38) in the K13^{580Y} compared to K13^{C580}

(**Figure 3** and **Supplemental Table S1**). This lower *K13* expression was observed in the rings, and merozoite

stages of K13^{580Y} (**Figure 3**). With the K13^{580Y} mutation partially affecting hemoglobin uptake, elevated

expression of PF3D7_1121600 (*EXP1*) has been linked to the initiation of the compensatory process involved

in minimizing the fitness costs associated with the partial loss of function associated with the K13 mutation

(30). We observed elevated expression of *EXP1* (log2FC=1.25, p-value=6.5E-136) in the K13^{580Y} relative to

K13^{C580} rings.

Impact of DHA exposure on K13^{C580} parasites.

In K13^{C580} parasites, the transcriptional response to DHA dramatically differed based on parasite stages

relative to DMSO (drug vehicle) treated controls. While we observed relative gene expression changes across

all the erythrocytic stages of development (Supplementary Table S2), the ring stage had the greatest gene

expression changes with 188, 270, and 353 genes identified as differentially expressed (DE) (p-value<0.05

and log2FC>0.4) at 2hrs, 4hrs and 6hrs after pulsed DHA exposure, respectively. The genes DE and

downregulated across all the treatment timepoints in the rings were categorically associated with metabolic

processes, such as glycolysis (*GAPDH*) and ATP generation (**Supplemental Table S2**), consistent with an arrest of parasite growth, as previously reported within 18h of DHA exposure (31). We observed the upregulation of stress-response genes such as *HSP70, HSP90, HSP101*, and *NOT4* across all the treatment timepoints (Figure 4). We also observed the elevated expression of genes such as *PTP3, PTP4, REX1,* and *MAHRP1,* which are involved in protein trafficking and export within the parasite (**Figure 4** and **Supplemental Table S2**). The *K13* gene and *Kelch13* interaction candidate (*KIC*) genes (such as *KIC1* and *KIC5*), known to code for proteins that associate with K13, were observed to be upregulated upon DHA treatment, impling that hemoglobin endocytosis is probably enhanced as a stress response and as a result enhancing drug potency within the parasite.

DHA is known to mainly affect phenotypic changes and survival in the ring and trophozoite stages of the malaria parasite (32). In our study, we explored the gene expression changes in the trophozoite stages of the $K13^{C580}$ variant across all time points, but we found very few DE genes. We did observe some expression changes in the merozoite, schizont and gametocyte stages of $K13^{C580}$ upon drug exposure (**Supplemental Table S2** and **Supplemental Figure S2**).

Impact of DHA exposure on $K13^{580Y}$ parasites.

Examining $K13^{580Y}$ after DHA treatment, we observed greater gene expression changes in the ring and trophozoite stages compared to all the other erythrocytic stages. We identified 99, 203, and 211 genes to be DE in the ring stage in response to 2hrs, 4hrs and 6hrs DHA exposure, respectively. In the $K13^{580Y}$ ring stages, downregulated genes included those linked to glycolysis (*GAPDH*) and ATP generation, suggesting a developmental arrest similar to $K13^{C580}$. Across all timepoints of drug exposure, we observed the downregulation of the *GARP* gene in the ring stage of the $K13^{580Y}$ strain (Supplemental Table S3 and Figure 5), but an upregulation of *GARP* in the trophozoite stage. We observed elevated expression of heat shock proteins (*HSP70*, *HSP90* and *HSP101*), upregulation of genes associated with protein trafficking (*PTP3*, *PTP4*, *REX1*, and *MAHRP1*) and also upregulation of *K13* and *KIC* genes (such as *KIC1*, *KIC5*) (**Figure 5** and **Supplemental Table S3**). These findings imply that the gene expression response initiated by DHA on $K13^{580Y}$

involved the downregulation of metabolic process, upregulation of the unfolded protein response, elevated

hemoglobin uptake and increased protein trafficking, which is a similar response initiated by the ring stage of

K13^{C580}, but a more enhanced cytoprotective response.

The trophozoite stage of the K13^{580Y} strain showed many gene expression changes upon drug exposure. We

observed 140, 93, and 222 genes to be DE in the trophozoite stage of K13^{580Y} in response to 2hrs, 4hrs and

6hrs DHA exposure, respectively. Within the trophozoite stage, we observed the upregulation of *PTP3*, *PTP4*,

EXP1 (protein trafficking), *HSP70, HSP90, HSP101* (unfolded protein response), *ApiAP2* and, contrary to the

rings stage, observed the upregulation of *GARP* across all timepoints of DHA exposure. The gene expression

changes in the trophozoite also appear to be associated with the restriction of metabolic processes, initiation of

the unfolded protein response, and elevated protein trafficking upon exposure to 700nM DHA over a 6hr

period. We also did observe some expression changes in the K13^{580Y} strain merozoite, schizont and

gametocyte stages upon drug exposure primarily after 4 to 6hrs of DHA exposure (**Supplemental Table S3**).

K13^{580Y} and K13^{C580} gene expression differences after treatment.

Upon DHA treatment, the transcriptional response appeared to be similar for both K13^{580Y} and K13^{C580} parasite

lines. Both K13^{580Y} and K13^{C580} ring stages upregulated genes associated with unfolded protein response

(UPR) and protein trafficking upon drug exposure, while genes associated with metabolic processes were

downregulated (**Supplemental Table S4**). However, there were are some differences observed between these

coisogenic lines upon DHA exposure, genes within the SERA gene family (PF3D7_0207600 (SERA5),

PF3D7_0207500 (SERA6)) had significant decreased expression (log2FC=-1.9,s, adj-*p*-value = 4.54E-06) in

K13^{580Y} compared to K13^{C580} trophozoites (**Figure 6** and **Supplementary Table S4**). Given SERA family

genes are thought to be important in maintaining the erythrocytic cycle growth (33–35), downregulation of

these genes in K13^{580Y} compared to DHA treated K13^{C580} could contribute to the initiation of parasite dormancy

through suppression of parasite development (Supplemental Table S3).

Elevated *PfGARP* expression in K13^{580Y}.

GARP has recently been identified as a promising therapeutic target based on the impact of anti-GARP antibodies shown to trigger parasite death (36). The exact role and function of GARP still remains to be elucidated, however it has been reported to be restrictively expressed in the early trophozoite stage (37). In this study, we show that the PfGARP gene is expressed in more stages of the parasite's life cycle, unlike previously reported (**Figure 6**). However, the expression level of this gene is variable across the stages, with higher expression between the late ring to the mid trophozoite stages (**Figure 6A**). We also observed variable *GARP* expression between the coisogenic lines used in the study. The $K13^{580Y}$ exhibited significantly higher *GARP* gene expression compared to $K13^{C580}$ (log2FC=3.49, adj-*p*-value<0.00001) (**Figure 6A** and **6B**). Upon DHA treatment of both $K13^{C580}$ and $K13^{580Y}$, we observed that *GARP* expression was significantly increased in the trophozoite stage (**Figure 6B**). These findings and observations imply that GARP may be involved in the parasite's response to stress/drug exposure, and in this case the response seems to be more pronounced in the $K13^{580Y}$ parasite.

Fine grained analysis between $K13^{580Y}$ vs. $K13^{C580}$ allows for quantification of treatment perturbation across the malaria asexual lifecycle

To quantify DHA perturbation across the malaria cell cycle stages, the MELD algorithm was utilized (see **Supplemental Methods**) (38). The output was a continuous measure, where a cell's likelihood score that approaches 1 indicates a cell's transcriptional profile is more similar to the DHA treatment condition, called the DHA likelihood. DHA likelihoods are scores ranging from 0 to 1, i.e. from less to more perturbed by DHA treatment. Pseudotime analysis was performed with slingshot (39), allowing visualization of the distribution of DHA likelihoods across the lifecycle, within each timepoint and strain (**Supplemental Figure S3**). Both lines show similar responses to DHA treatment, where there are separate transcriptional states based on treatment condition across the lifecycle (**Figure 7A**). As expected, DHA treated cells have a higher DHA likelihood than DMSO treated cells (**Figure 7B**). In $K13^{C580}$, DHA likelihoods of the DHA treatment condition increased over the asexual lifecycle at the first two timepoints (2hrs and 4hrs) and remained elevated in the ring stages at 6hrs. In $K13^{580Y}$, the same pattern was present, but with a greater representation of all the asexual stages across timepoints and higher levels of DHA likelihood values in the treatment condition.

When focusing on the ring and trophozoite stages, we isolated cell populations with the highest DHA likelihood within each timepoint using vertex frequency clustering (VFC) (38). We separated these highly enriched clusters of high DHA likelihood cells (**Supplemental Figures S4** and **S5**) for fine grained differential expression analysis in rings and trophozoites.

We compared the expression between the clusters with the highest DHA likelihood between K13^{580Y} and K13^{C580} in the ring and trophozoite stages (**Figure 8** and **Supplemental Figure S6**). Rings and trophozoites in K13^{580Y} showed consistent upregulation of GARP and downregulation of Knob Associated Histidine Rich Protein (KAHRP, PF3D7-0202000) and Ring Infected Erythrocyte Surface Antigen (RESA, PF3D7-0102200) compared to K13^{C580}. To investigate differential gene pathway expression, gene ontology (GO) analysis was performed using the differentially expressed genes in K13^{580Y} vs. K13^{C580} comparison (**Supplemental Figure S7**). In the ring stage, GO terms related to metabolic processes and translation were enriched in upregulated genes. In the trophozoite stage, GO terms related to nuclear regulation were enriched in downregulated genes in timepoints 4 hours and 6 hours.

Anti-GARP kills K13^{580Y} equivalent to K13^{C580} *in vitro*.

GARP was elevated upon DHA treatment of K13^{580Y} compared to K13^{C580}. In order to evaluate if this elevated expression resulted in a functional or aberrant accumulation of the protein in the membrane, we performed growth-inhibition assays (GIAs) using polyclonal mouse anti-GARP antiserum generated by lipid nanoparticle (LNP)-mRNA based vaccination (36). We found statistically significant growth inhibition of both K13^{C580} or K13^{580Y} lines after exposure to anti-GARP (Figure 9),consistent with functionality of the pathway in both lines (**Figure 9**).

DISCUSSION

To date, studies of Plasmodium falciparum transcriptional responses to artemisinin have largely relied on bulk RNA sequencing of synchronized parasites, limiting resolution across developmental stages and obscuring cell-to-cell heterogeneity (13, 24, 31, 40). Here, we leveraged single-cell RNA sequencing to comprehensively profile the transcriptional landscape of artemisinin-sensitive (K13^{C580}) and artemisinin-resistant (K13^{580Y}) coisogenic parasites following dihydroartemisinin (DHA) exposure. This approach revealed stage-specific and strain-specific responses across the asexual blood-stage lifecycle. We identified a shared core response to DHA in both lines, including downregulation of metabolic genes and upregulation of stress response pathways such as the unfolded protein response and protein trafficking. However, resistant parasites exhibited a more consistent and sustained transcriptional shift across stages, including signatures of dormancy and enhanced expression of surface proteins such as PfEMP1 and GARP. Using MELD, we quantified the extent of transcriptional perturbation and found that resistant parasites had higher DHA likelihood scores across the lifecycle, indicating broader and more persistent transcriptional remodeling (see Figures 6, 8 and Supplemental Figure S7). These findings suggest that artemisinin resistance is associated not only with specific gene expression changes but also with a fundamentally altered capacity to respond to drug-induced stress at the single-cell level. Our results provide a high-resolution view of the molecular adaptations that underlie artemisinin resistance and identify potential vulnerabilities for therapeutic targeting.

When comparing K13^{580Y} and K13^{C580} parasites, we found interesting differences in gene expression. GCN5 has recently been found to regulate stress response genes after artemisinin treatment (41) and this gene was upregulated in K13^{580Y} trophozoites at all timepoints compared to K13^{C580} trophozoites (**Figure 8**). Additionally, *FIKK1* and *FIKK4.2* genes were upregulated in K13^{580Y} rings and trophozoites compared to K13^{C580}, genes that were both part of the "artemisinin resistance-associated transcriptional profile" found when analyzing isolates from the Greater Mekong Subregion (GMS) (42). Downregulation of the *SERA5* and 6 genes in K13^{580Y} vs. K13^{C580} upon DHA exposure highlights some of the changes that could lead to parasite quiescence due to slowed development (Figure 8). We also revealed an increased expression of transport associated genes in K13^{580Y} vs. K13^{C580} when exposed to DHA as well as a striking elevated expression of *GARP*, a potential *P.*

falciparum therapeutic target (36), in the trophozoite stage of K13^{580Y} upon DHA exposure, making it a viable therapeutic target candidate against the emergence of ART-R as a result of K13^{580Y} mutation.

Our transcriptomic analysis revealed that untreated K13^{580Y} had lower *K13* gene expression compared to untreated K13^{C580}. K13 is thought to aid hemoglobin endocytosis into the parasite, to be broken down as a source of amino acids to facilitate the assembly of parasite proteins and support the life of the parasite. The presence of the K13^{580Y} mutation is associated with a slow rate of progression of the parasite from the ring to trophozoite to schizont stage compared to K13^{C580} (**Figure 2C**). This deceleration in development in the blood-stage is thought to be as a result of impaired hemoglobin uptake, which is necessary for parasite protein synthesis (23, 43). The K13 function of facilitating hemoglobin entry has been shown to be aided by a complex of other proteins, known as the K13 compartment proteins (KICs) (23). During a stress response, *P. falciparum* has been shown to elevate *K13* gene and protein expression to facilitate hemoglobin entry, thereby availing an amino acid source which will be used to synthesize proteins necessary to control the stress response (31). Parasites exposed to DHA treatment did exhibit elevated expression of *K13* and genes coding for the KIC's mainly at the ring stages of both K13^{C580} and K13^{580Y}. However in K13^{580Y}, *K13* gene expression was lower in most of the parasite stages expressing the gene compared to K13^{C580} (**Figure 6**), despite some of the KIC genes being elevated upon DHA exposure (**Supplementary Table 2** and **3**). This would suggest that altered protein expression of K13 downstream of transcription likely alters parasite DHA sensitivity in K13^{580Y}(23).

Artemisinin and its derivatives are known to exert their antimalarial activity by generating carbon-centered radicals that cause oxidative stress and subsequent protein alkylation (25, 44). Accumulation of the alkylated proteins increases proteotoxic stress within the parasites, causing a phenotype reminiscent of that induced by heat shock (45, 46), hence the observed upregulation of *HSP70* and *HSP90* in both K13^{580Y} and K13^{C580} upon DHA treatment. This finding, together with the elevated expression of genes associated with protein synthesis (FIKK family of genes, *HSP40*) and transcriptional activation of genes (*ApiAP2*), implies the induction of the UPR in both parasite lines upon DHA exposure. We also observed a relatively high number of genes coding for export and trafficking proteins to be upregulated in the rings and trophozoites of both K13^{580Y} and K13^{C580} upon

DHA exposure, which could be as a result of ER stress response (47). Our findings show that both K13^{580Y} and K13^{C580} exhibit a similar gene expression response to DHA exposure. However, K13^{580Y} exhibits enhanced expression of UPR associated genes compared to K13^{C580}, increasing the capacity of K13^{580Y} to quickly repair or degrade proteins damaged by artemisinin. This capacity could minimize the damaging effects caused by the accumulation of DHA in K13^{580Y} parasites. This elevated UPR in K13^{580Y}, coupled with having lower levels of hemoglobin endocytosis and decreasing free heme-mediated artemisinin activation, contributes to the tolerance of K13^{580Y} to artemisinin (48), leading to resistance/delayed killing of K13^{580Y}.

GARP was recently identified as a therapeutic target (36) and previous studies have shown that the expression of *GARP* is highly restricted to the early trophozoite stage of the parasite's life cycle; however, we show that the gene is also expressed in the late ring stages and trophozoites (**Figure 6**). *GARP* shares no homology with proteins of known function, it contains several complex repeat regions and its function remains to be characterized. We observed elevated expression of *GARP* in K13^{580Y} compared to K13^{C580} before treatment, which could imply it may be as a result of compensatory processes initiated by the parasite due to the K13 mutation. While not as elevated, other surface protein transcripts show upregulation suggesting a possible general effect. Given gene expression changes can correlate with function or represent dysregulation (e.g. increased turnover), we examined whether GARP still retained function as a therapeutic target, finding statistically significant anti-GARP killing of both K13^{580Y} and K13^{C580} parasites (Figure 9). This suggests GARP and likely other surface proteins maintain normal function and suggests anti-GARP therapy can be a potential addition to our limited armamentarium as an effective therapy in the face of K13 artemisinin partial resistance mutations.

In this study, we provide a comprehensive single-cell transcriptional map of *Plasmodium falciparum*'s response to artemisinin exposure across the asexual blood-stage lifecycle, revealing both shared and resistance-associated adaptations in these coisogenic parasites. We show that the K13^{580Y} mutant exhibits a broader and more sustained transcriptional response to DHA, including enhanced activation of stress response pathways, metabolic arrest, and signatures of dormancy. Using MELD, we quantified these perturbations and

demonstrated that resistant parasites maintain elevated transcriptional shifts across stages, suggesting a

fundamentally altered capacity to withstand drug pressure. Importantly, we identified consistently elevated

expression of the surface protein GARP in resistant K13^{580Y} parasites, both at baseline and following

treatment, and confirmed its vulnerability to antibody-mediated inhibition. These findings not only deepen our

understanding of the cellular mechanisms underlying artemisinin resistance but also highlight GARP and other

surface proteins as promising therapeutic targets. Our work underscores the power of single-cell approaches

to dissect complex drug responses and provides a framework for identifying molecular vulnerabilities in

resistant malaria parasites.

  METHODS

  Parasites lines and culture. The strains of parasite lines used in this study were MRA-1250 CamWT, an

  artemisinin susceptible fast-clearing Cambodian *P. falciparum* isolate, and MRA-1251 CamWT_C580Y, a

  Cambodian *P. falciparum* isolate which features a single nucleotide substitution leading to a C580Y amino acid

  change (MR4, BEI Resources, Manassas, VA). The *P. falciparum* parasites were maintained at 3% hematocrit

  in human O+ RBCs and *P. falciparum* culture media comprising RPMI 1640 (Thermo Fisher Scientific)

  supplemented with 0.5% (w/v) Albumax II, 50 mg/L hypoxanthine, 0.225% $NaHCO_3$, 25mM HEPES, and 10

  mg/L gentamycin. Parasites were cultured at 37 °C in 5% O_2, 5% CO_2, and 90% N_2 (49). The parasite lines

  were cultured for 48hrs before we began exposing them to DHA and DMSO.

  Drug susceptibility assays. Drug stocks of dihydroartemisinin (DHA) were made in dimethyl sulfoxide

  (DMSO) and aliquots were stored at −20 °C. All drug assays were conducted such that the final DMSO

  concentration was <0.5%. Each parasite line was exposed to 700 nM DHA along with vehicle-treated 0.05%

  DMSO controls in independent experiments for 6hrs (**Figure 1**). Samples from each treated parasite line were

  collected at 2hr intervals for single cell library preparation and sequencing.

  Enrichment of parasite infected RBCs. To prepare the single cell RNAseq libraries, we needed to enrich the

  parasite infected RBCs thus increasing chances of loading and capturing mRNA transcripts from the parasites.

  Our unsynchronized parasite cultures were enriched using SLOPE as previously described (50, 51) with some

  modifications. Briefly, erythrocyte density was measured using a Cellometer Auto T4 (Nexcelom Biosciences,

  Lawrence, MA) and cell density adjusted to 2×10^9 erythrocytes/mL. The 22.5Units of the lytic agent

  streptolysin-O (SLO) per 50µl RBCs was added in a ratio of 2 parts SLO solution to 5 parts erythrocytes.

  Samples were mixed well by pipetting and incubated at room temperature for 6 min. 10 volumes of 1X PBS or

  media (RPMI 1640 HEPES) were added and cells were centrifuged at 2,500xg for 3 min. After removal of the

  supernatant, cells were washed twice more with 1X PBS. Following SLO lysis, cells suspended in 1X PBS

were layered onto a 60% Percoll gradient (Sigma Aldrich, St Louis, MO) and centrifuged at 1,500xg for 10 min. The top layer of Percoll was discarded while the lower cell pellet, rich in infected erythrocytes, was collected and washed twice with 1xPBS and later suspended in 200μl RPMI to be loaded into the SeqWell arrays.

Single-Cell library preparation and sequencing. To capture the transcriptome of *Plasmodium* single cells and generate single cell libraries, we utilized SeqWell, a massively parallel, low-input scRNA-seq platform (27, 28). Briefly, 10-15,000 *Pf* infected RBCs from each time point and treatment condition were loaded onto a functionalized polydimethylsiloxane (PDMS) array preloaded with uniquely barcoded mRNA capture beads (Chemgenes; Macosko-2011-10(V+), Wilmington, MA). After cells had settled into wells, the arrays were then sealed with a hydroxylated polycarbonate membrane with a pore size of 10 nm, facilitating buffer exchange while confining biological molecules (such as DNA and RNA) within each well. Following membrane-sealing, subsequent buffer exchange permitted cell lysis, mRNA transcript hybridization to beads, and bead removal before proceeding with reverse transcription. The obtained bead-bound cDNA product then underwent Exonuclease I treatment to remove excess primer before proceeding with second-strand synthesis, then PCR amplification of the cDNA product. The cDNA products were purified using the SPRIselect beads (Beckman Coulter, Brea, CA), and were run on a Fragment analyzer (Agilent Technologies, Santa Clara, CA) to verify the cDNA fragment sizes to be between 0.7-2kbps. The cDNA libraries were then fragmented and amplified for sequencing using the Illumina Nextera XT DNA Library Preparation kit (Illumina, FC-131-1096, San Diego, CA) using custom primers that enabled the specific amplification of only the 3' ends. The libraries were purified, quantified, and then sequenced on a NextSeq550 system using High output 75 cycle kits (Illumina) at an average of 20,000 reads per cell. Raw read data and processed UMI count matrices from *Pf* MRA1250 and MRA1251 single cell experiments have been deposited and are publicly available through the SRA database (BioProject ID: PRJNA1049964).

Single-cell transcriptome analysis. Raw reads were processed using version 2.3.0 of the Drop-seq pipeline (52), and according to the 'Drop-seq Alignment Cookbook', both found at http://mccarrolllab.com/dropseq/. Demultiplexed FASTQs were aligned to the *P. falciparum* 3D7 reference genome (PlasmoDB v.51) using

STAR aligner. Individual reads were tagged with a 12bp barcode and 9bp unique molecular identifier (UMI) contained in Read 1 of each sequencing fragment. Following alignment, reads were grouped by the 12bp cell barcodes and subsequently collapsed by the 9bp UMI to generate a gene expression count matrix. Gene expression count matrices from *Pf* MRA1250 and *Pf* MRA1251 cell line single-cell experiments (n = 7 per *Pf* cell line) were analyzed and visualized in R (v.4.0.3) using the Seurat package (v.4.0.4) (53). Cells with less than 300 genes and 500 UMIs detected in less than ten cells were excluded from further analysis. We then proceeded to filter out 459 multigene family genes expressed such as *vars (PfEMP1)*, *rifins* and *stevors*, retaining genes present across the different experimental conditions for each parasite line. Library size for each group in the study were internally normalized to 10,000 transcripts, log transformed and regressed on the number of UMIs per cell before dimensionality reduction and clustering. We integrated the datasets by utilizing the canonical correlation analysis ('CCA') in Seurat (54) to identify anchors/shared sources of variations between each treated and untreated groups to learn the parasite stage specific responses to DHA and DMSO treatment. Next, read data were scaled and analyzed via principal component analysis (PCA). The top PCs (n = 20) were dimensionally reduced via uniform manifold approximation (UMAP) (55) and unsupervised clustering was performed at several resolutions to identify parasite development stages and analyze differential gene expression.

A *p*-value cutoff of 0.05 was used to determine significance. For differential gene expression (DGE) between the treatment groups, a Bonferroni correction was applied to generate adjusted *p*-values. DGE analysis was done using the MAST package (56).

Details regarding fine-grained differential expression analysis utilizing MELD is in Supplemental Methods.

Growth inhibition assay on K13^{C580} and K13^{580Y}. K13^{C580} and K13^{580Y} cell lines were synchronized to the ring stage by treatment with 5% sorbitol for at least two successive replication cycles and cultured to the ring stage. Parasites were diluted with O+ RBCs to 0.5-0.75% parasitemia and 2% hematocrit in incomplete media (RPMI 1640 medium containing 25 mm HEPES and 50µg/mL hypoxanthine) supplemented with 10% AlbuMax II, 48mM sodium bicarbonate, and 20µg/mL gentamycin. Anti-PfGARP (10% v/v) or normal mouse sera (10%

13   v/v) was added to the cultures in 96-well round-bottom microtiter plates. After a 72-hour incubation, cultures

14   were stained with Hoechst nucleic acid stain to quantify parasitemia using a CytoFLEX flow cytometer.

16   **Data availability**: All sequence data has been submitted to SRA (PRJNA1049964). Code is available at

17   https://github.com/sconnelly007/K13_mt_v_wt_sc.

Acknowledgments: The following reagents were obtained through BEI Resources, NIAID, NIH: *Plasmodium falciparum*, strains CamWT, MRA-1250 and CamMT, MRA-1251, contributed by David A. Fidock; and *Plasmodium falciparum*, Strain 3D7, MRA-102, contributed by Daniel J. Carucci.

Author contributions

CIO, SVC, CC, JAB, and JJJ designed the research study. CIO, CC, IP, DC, NR, and JZ conducted the experiments. CIO and SVC analyzed the data. CIO, SVC, JAB and JJJ wrote the manuscript. All authors reviewed and revised the manuscript.

Funding: The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. This work was funded by the National Institute for Allergy and Infectious Diseases (K24AI134990 to JJJ; R01AI121558 to JJJ, and R01AI156267 to JAB and JJJ). Funds were received by the Falk Medical Research Trust Transformational Award (JK).

Conflicts of Interest: All authors declare no competing interests.

Ethics Statement: This research does not constitute human subjects research.

REFERENCES

1. Programme GM. 2022. World malaria report 2022. World Health Organization. https://www.who.int/publications/i/item/9789240064898. Retrieved 14 November 2023.

2. Teuscher F, Gatton ML, Chen N, Peters J, Kyle DE, Cheng Q. 2010. Artemisinin☐induced dormancy in plasmodium falciparum: duration, recovery rates, and implications in treatment failure. J Infect Dis 202:1362–1368.

3. Rosenthal PJ. 2021. Has artemisinin resistance emerged in Africa? Lancet Infect Dis 21:1056–1057.

4. Tacoli C, Gai PP, Bayingana C, Sifft K, Geus D, Ndoli J, Sendegeya A, Gahutu JB, Mockenhaupt FP. 2016. Artemisinin Resistance-Associated K13 Polymorphisms of Plasmodium falciparum in Southern Rwanda, 2010-2015. Am J Trop Med Hyg 95:1090–1093.

5. Nyunt MH, Soe MT, Myint HW, Oo HW, Aye MM, Han SS, Zaw NN, Cho C, Aung PZ, Kyaw KT, Aye TT, San NA, Ortega L, Thimasarn K, Bustos MDG, Galit S, Hoque MR, Ringwald P, Han E-T, Kyaw MP. 2017. Clinical and molecular surveillance of artemisinin resistant falciparum malaria in Myanmar (2009-2013). Malar J 16:333.

6. Yasri S, Wiwanitkit V. 2021. Artemisinin resistance: an important emerging clinical problem in tropical medicine. Int J Physiol Pathophysiol Pharmacol 13:152–157.

7. Ashley EA, Dhorda M, Fairhurst RM, Amaratunga C, Lim P, Suon S, Sreng S, Anderson JM, Mao S, Sam B, Sopha C, Chuor CM, Nguon C, Sovannaroth S, Pukrittayakamee S, Jittamala P, Chotivanich K, Chutasmit K, Suchatsoonthorn C, Runcharoen R, Hien TT, Thuy-Nhien NT, Thanh NV, Phu NH, Htut Y, Han K-T, Aye KH, Mokuolu OA, Olaosebikan RR, Folaranmi OO, Mayxay M, Khanthavong M, Hongvanthong B, Newton PN, Onyamboko MA, Fanello CI, Tshefu AK, Mishra N, Valecha N, Phyo AP, Nosten F, Yi P, Tripura R, Borrmann S, Bashraheil M, Peshu J, Faiz MA, Ghose A, Hossain MA, Samad R, Rahman MR, Hasan MM, Islam A, Miotto O, Amato R, MacInnis B, Stalker J, Kwiatkowski DP, Bozdech Z, Jeeyapant A, Cheah PY, Sakulthaew T, Chalk J, Intharabut B, Silamut K, Lee SJ, Vihokhern B, Kunasol C, Imwong M, Tarning J, Taylor WJ, Yeung S, Woodrow CJ, Flegg JA, Das D, Smith J, Venkatesan M, Plowe CV, Stepniewska K, Guerin PJ, Dondorp AM, Day NP, White NJ, Tracking Resistance to Artemisinin Collaboration (TRAC). 2014. Spread of artemisinin resistance in Plasmodium

falciparum malaria. N Engl J Med 371:411–423.

8. Fola AA, Feleke SM, Mohammed H, Brhane BG, Hennelly CM, Assefa A, Crudal RM, Reichert E, Juliano

JJ, Cunningham J, Mamo H, Solomon H, Tasew G, Petros B, Parr JB, Bailey JA. 2023. Plasmodium

falciparum resistant to artemisinin and diagnostics have emerged in Ethiopia. Nat Microbiol 8:1911–1919.

9. Juliano JJ, Giesbrecht DJ, Simkin A, Fola AA, Lyimo BM, Pereus D, Bakari C, Madebe RA, Seth MD,

Mandara CI, Popkin-Hall ZR, Moshi R, Mbwambo RB, Niaré K, MacInnis B, Francis F, Mbwambo D,

Garimo I, Chacky F, Aaron S, Lusasi A, Molteni F, Njau R, Lazaro S, Mohamed A, Bailey JA, Ishengoma

DS. 2023. Country wide surveillance reveals prevalent artemisinin partial resistance mutations with

evidence for multiple origins and expansion of sulfadoxine-pyrimethamine resistance mutations in

northwest Tanzania. bioRxiv.

10. Checchi F, Durand R, Balkan S, Vonhm BT, Kollie JZ, Biberson P, Baron E, Le Bras J, Guthmann JP.

2002. High Plasmodium falciparum resistance to chloroquine and sulfadoxine-pyrimethamine in Harper,

Liberia: results in vivo and analysis of point mutations. Trans R Soc Trop Med Hyg 96:664–669.

11. Roux AT, Maharaj L, Oyegoke O, Akoniyon OP, Adeleke MA, Maharaj R, Okpeku M. 2021. Chloroquine

and Sulfadoxine-Pyrimethamine Resistance in Sub-Saharan Africa-A Review. Front Genet 12:668574.

12. Noedl H, Se Y, Schaecher K, Smith BL, Socheat D, Fukuda MM, Artemisinin Resistance in Cambodia 1

(ARC1) Study Consortium. 2008. Evidence of artemisinin-resistant malaria in western Cambodia. N Engl J

Med 359:2619–2620.

13. Dondorp AM, Nosten F, Yi P, Das D, Phyo AP, Tarning J, Lwin KM, Ariey F, Hanpithakpong W, Lee SJ,

Ringwald P, Silamut K, Imwong M, Chotivanich K, Lim P, Herdman T, An SS, Yeung S, Singhasivanon P,

Day NPJ, Lindegardh N, Socheat D, White NJ. 2009. Artemisinin resistance in Plasmodium falciparum

malaria. N Engl J Med 361:455–467.

14. Ariey F, Witkowski B, Amaratunga C, Beghain J, Langlois A-C, Khim N, Kim S, Duru V, Bouchier C, Ma L,

Lim P, Leang R, Duong S, Sreng S, Suon S, Chuor CM, Bout DM, Ménard S, Rogers WO, Genton B,

Fandeur T, Miotto O, Ringwald P, Le Bras J, Berry A, Barale J-C, Fairhurst RM, Benoit-Vical F,

Mercereau-Puijalon O, Ménard D. 2014. A molecular marker of artemisinin-resistant Plasmodium

falciparum malaria. Nature 505:50–55.

15. Straimer J, Gnädig NF, Witkowski B, Amaratunga C, Duru V, Ramadani AP, Dacheux M, Khim N, Zhang L, Lam S, Gregory PD, Urnov FD, Mercereau-Puijalon O, Benoit-Vical F, Fairhurst RM, Ménard D, Fidock DA. 2015. Drug resistance. K13-propeller mutations confer artemisinin resistance in Plasmodium falciparum clinical isolates. Science 347:428–431.

16. Intharabut B, Kingston HW, Srinamon K, Ashley EA, Imwong M, Dhorda M, Woodrow C, Stepniewska K, Silamut K, Day NPJ, Dondorp AM, White NJ, Tracking Resistance to Artemisinin Collaboration. 2019. Artemisinin Resistance and Stage Dependency of Parasite Clearance in Falciparum Malaria. J Infect Dis 219:1483–1489.

17. Ménard D, Khim N, Beghain J, Adegnika AA, Shafiul-Alam M, Amodu O, Rahim-Awab G, Barnadas C, Berry A, Boum Y, Bustos MD, Cao J, Chen J-H, Collet L, Cui L, Thakur G-D, Dieye A, Djallé D, Dorkenoo MA, Eboumbou-Moukoko CE, Espino F-E-CJ, Fandeur T, Ferreira-da-Cruz M-F, Fola AA, Fuehrer H-P, Hassan AM, Herrera S, Hongvanthong B, Houzé S, Ibrahim ML, Jahirul-Karim M, Jiang L, Kano S, Ali-Khan W, Khanthavong M, Kremsner PG, Lacerda M, Leang R, Leelawong M, Li M, Lin K, Mazarati J-B, Ménard S, Morlais I, Muhindo-Mavoko H, Musset L, Na-Bangchang K, Nambozi M, Niaré K, Noedl H, Ouédraogo J-B, Pillai DR, Pradines B, Quang-Phuc B, Ramharter M, Randrianarivelojosia M, Sattabongkot J, Sheikh-Omar A, Silué KD, Sirima SB, Sutherland C, Syafruddin D, Tahar R, Tang L-H, Touré OA, Tshibangu-wa-Tshibangu P, Vigan-Womas I, Warsame M, Wini L, Zakeri S, Kim S, Eam R, Berne L, Khean C, Chy S, Ken M, Loch K, Canier L, Duru V, Legrand E, Barale J-C, Stokes B, Straimer J, Witkowski B, Fidock DA, Rogier C, Ringwald P, Ariey F, Mercereau-Puijalon O, KARMA Consortium. 2016. A Worldwide Map of Plasmodium falciparum K13-Propeller Polymorphisms. N Engl J Med 374:2453–2464.

18. World Health Organization. 2020. Report on antimalarial drug efficacy, resistance and response: 10 years of surveillance (2010-2019). World Health Organization, Genève, Switzerland.

19. Witkowski B, Amaratunga C, Khim N, Sreng S, Chim P, Kim S, Lim P, Mao S, Sopha C, Sam B, Anderson JM, Duong S, Chuor CM, Taylor WRJ, Suon S, Mercereau-Puijalon O, Fairhurst RM, Menard D. 2013. Novel phenotypic assays for the detection of artemisinin-resistant Plasmodium falciparum malaria in Cambodia: in-vitro and ex-vivo drug-response studies. Lancet Infect Dis 13:1043–1049.

20. Klonis N, Xie SC, McCaw JM, Crespo-Ortiz MP, Zaloumis SG, Simpson JA, Tilley L. 2013. Altered

temporal response of malaria parasites determines differential sensitivity to artemisinin. Proc Natl Acad

Sci U S A 110:5157–5162.

21. Saralamba S, Pan-Ngum W, Maude RJ, Lee SJ, Tarning J, Lindegårdh N, Chotivanich K, Nosten F, Day

NPJ, Socheat D, White NJ, Dondorp AM, White LJ. 2011. Intrahost modeling of artemisinin resistance in

Plasmodium falciparum. Proc Natl Acad Sci U S A 108:397–402.

22. Ishengoma DS, Gosling R, Martinez-Vega R, Beshir KB, Bailey JA, Chimumbwa J, Sutherland C, Conrad

MD, Tadesse FG, Juliano JJ, Kamya MR, Mbacham WF, Ménard D, Rosenthal PJ, Raman J, Tatarsky A,

Tessema SK, Fidock DA, Djimde AA. 2024. Urgent action is needed to confront artemisinin partial

resistance in African malaria parasites. Nat Med 30:1807–1808.

23. Birnbaum J, Scharf S, Schmidt S, Jonscher E, Hoeijmakers WAM, Flemming S, Toenhake CG, Schmitt M,

Sabitzki R, Bergmann B, Fröhlke U, Mesén-Ramírez P, Soares AB, Herrmann H, Bártfai R, Spielmann T.

2020. A Kelch13-defined endocytosis pathway mediates artemisinin resistance in malaria parasites.

Science.

24. Zhu L, Tripathi J, Rocamora FM, Miotto O, van der Pluijm R, Voss TS, Mok S, Kwiatkowski DP, Nosten F,

Day NPJ, White NJ, Dondorp AM, Bozdech Z, Tracking Resistance to Artemisinin Collaboration I. 2018.

The origins of malaria artemisinin resistance defined by a genetic and transcriptomic background. Nat

Commun 9:5158.

25. Tilley L, Straimer J, Gnädig NF, Ralph SA, Fidock DA. 2016. Artemisinin Action and Resistance in

Plasmodium falciparum. Trends Parasitol 32:682–696.

26. Meshnick SR. 2002. Artemisinin: mechanisms of action, resistance and toxicity. Int J Parasitol 32:1655–

1660.

27. Gierahn TM, Wadsworth MH 2nd, Hughes TK, Bryson BD, Butler A, Satija R, Fortune S, Love JC, Shalek

AK. 2017. Seq-Well: portable, low-cost RNA sequencing of single cells at high throughput. Nat Methods

14:395–398.

28. Aicher TP, Carroll S, Raddi G, Gierahn T, Wadsworth MH 2nd, Hughes TK, Love C, Shalek AK. 2019.

Seq-Well: A Sample-Efficient, Portable Picowell Platform for Massively Parallel Single-Cell RNA

Sequencing. Methods Mol Biol 1979:111–132.

29. Howick VM, Russell AJC, Andrews T, Heaton H, Reid AJ, Natarajan K, Butungi H, Metcalf T, Verzier LH, Rayner JC, Berriman M, Herren JK, Billker O, Hemberg M, Talman AM, Lawniczak MKN. 2019. The Malaria Cell Atlas: Single parasite transcriptomes across the complete Plasmodium life cycle. Science 365:eaaw2619.

30. Mesén-Ramírez P, Bergmann B, Elhabiri M, Zhu L, von Thien H, Castro-Peña C, Gilberger T-W, Davioud-Charvet E, Bozdech Z, Bachmann A, Spielmann T. 2021. The parasitophorous vacuole nutrient channel is critical for drug access in malaria parasites and modulates the artemisinin resistance fitness cost. Cell Host Microbe 29:1774–1787.e9.

31. Mok S, Stokes BH, Gnädig NF, Ross LS, Yeo T, Amaratunga C, Allman E, Solyakov L, Bottrill AR, Tripathi J, Fairhurst RM, Llinás M, Bozdech Z, Tobin AB, Fidock DA. 2021. Artemisinin-resistant K13 mutations rewire Plasmodium falciparum's intra-erythrocytic metabolic program to enhance survival. Nat Commun 12:1–15.

32. Xie SC, Dogovski C, Hanssen E, Chiu F, Yang T, Crespo MP, Stafford C, Batinovic S, Teguh S, Charman S, Klonis N, Tilley L. 2016. Haemoglobin degradation underpins the sensitivity of early ring stage Plasmodium falciparum to artemisinins. J Cell Sci 129:406–416.

33. Miller SK, Good RT, Drew DR, Delorenzi M, Sanders PR, Hodder AN, Speed TP, Cowman AF, de Koning-Ward TF, Crabb BS. 2002. A subset of Plasmodium falciparum SERA genes are expressed and appear to play an important role in the erythrocytic cycle. J Biol Chem 277:47524–47532.

34. McCoubrie JE, Miller SK, Sargeant T, Good RT, Hodder AN, Speed TP, de Koning-Ward TF, Crabb BS. 2007. Evidence for a common role for the serine-type Plasmodium falciparum serine repeat antigen proteases: implications for vaccine and drug design. Infect Immun 75:5565–5574.

35. Thomas JA, Collins CR, Das S, Hackett F, Graindorge A, Bell D, Deu E, Blackman MJ. 2016. Development and Application of a Simple Plaque Assay for the Human Malaria Parasite Plasmodium falciparum. PLoS One 11:e0157873.

36. Raj DK, Das Mohapatra A, Jnawali A, Zuromski J, Jha A, Cham-Kpu G, Sherman B, Rudlaff RM, Nixon CE, Hilton N, Oleinikov AV, Chesnokov O, Merritt J, Pond-Tor S, Burns L, Jolly G, Ben Mamoun C,

Kabyemela E, Muehlenbachs A, Lambert L, Orr-Gonzalez S, Gnädig NF, Fidock DA, Park S, Dvorin JD, Pardi N, Weissman D, Mui BL, Tam YK, Friedman JF, Fried M, Duffy PE, Kurtis JD. 2020. Anti-PfGARP activates programmed cell death of parasites and reduces severe malaria. Nature 582:104–108.

37. López-Barragán MJ, Lemieux J, Quiñones M, Williamson KC, Molina-Cruz A, Cui K, Barillas-Mury C, Zhao K, Su X-Z. 2011. Directional gene expression and antisense transcripts in sexual and asexual stages of Plasmodium falciparum. BMC Genomics 12:587.

38. Burkhardt DB, Stanley JS, Tong A, Perdigoto AL, Gigante SA, Herold KC, Wolf G, Giraldez AJ, van Dijk D, Krishnaswamy S. 2021. Quantifying the effect of experimental perturbations at single-cell resolution. Nat Biotechnol 39:619–629.

39. Street K, Risso D, Fletcher RB, Das D, Ngai J, Yosef N, Purdom E, Dudoit S. 2018. Slingshot: cell lineage and pseudotime inference for single-cell transcriptomics. BMC Genomics 19:477.

40. Mok S, Ashley EA, Ferreira PE, Zhu L, Lin Z, Yeo T, Chotivanich K, Imwong M, Pukrittayakamee S, Dhorda M, Nguon C, Lim P, Amaratunga C, Suon S, Hien TT, Htut Y, Faiz MA, Onyamboko MA, Mayxay M, Newton PN, Tripura R, Woodrow CJ, Miotto O, Kwiatkowski DP, Nosten F, Day NPJ, Preiser PR, White NJ, Dondorp AM, Fairhurst RM, Bozdech Z. 2015. Drug resistance. Population transcriptomics of human malaria parasites reveals the mechanism of artemisinin resistance. Science 347:431–435.

41. Rawat M, Kanyal A, Sahasrabudhe A, Vembar SS, Lopez-Rubio J-J, Karmodiya K. 2021. Histone acetyltransferase PfGCN5 regulates stress responsive and artemisinin resistance related genes in Plasmodium falciparum. Sci Rep 11:1–13.

42. Zhu L, van der Pluijm RW, Kucharski M, Nayak S, Tripathi J, White NJ, Day NPJ, Faiz A, Phyo AP, Amaratunga C, Lek D, Ashley EA, Nosten F, Smithuis F, Ginsburg H, von Seidlein L, Lin K, Imwong M, Chotivanich K, Mayxay M, Dhorda M, Nguyen HC, Nguyen TNT, Miotto O, Newton PN, Jittamala P, Tripura R, Pukrittayakamee S, Peto TJ, Hien TT, Dondorp AM, Bozdech Z. 2022. Artemisinin resistance in the malaria parasite, Plasmodium falciparum, originates from its initial transcriptional response. Commun Biol 5:274.

43. Yang T, Yeoh LM, Tutor MV, Dixon MW, McMillan PJ, Xie SC, Bridgford JL, Gillett DL, Duffy MF, Ralph SA, McConville MJ, Tilley L, Cobbold SA. 2019. Decreased K13 Abundance Reduces Hemoglobin

Catabolism and Proteotoxic Stress, Underpinning Artemisinin Resistance. Cell Rep 29:2917–2928.e5.

44. Paloque L, Ramadani AP, Mercereau-Puijalon O, Augereau J-M, Benoit-Vical F. 2016. Plasmodium falciparum: multifaceted resistance to artemisinins. Malar J 15:149.

45. Morano KA, Grant CM, Moye-Rowley WS. 2012. The response to heat shock and oxidative stress in Saccharomyces cerevisiae. Genetics 190:1157–1195.

46. Roti Roti JL. 2008. Cellular responses to hyperthermia (40-46 degrees C): cell killing and molecular events. Int J Hyperthermia 24:3–15.

47. Marti M, Spielmann T. 2013. Protein export in malaria parasites: many membranes to cross. Curr Opin Microbiol 16:445–451.

48. Rosenthal MR, Ng CL. 2020. Plasmodium falciparum Artemisinin Resistance: The Effect of Heme, Protein Damage, and Parasite Cell Stress Response. ACS Infect Dis 6:1599–1614.

49. Schuster FL. 2002. Cultivation of plasmodium spp. Clin Microbiol Rev 15:355–364.

50. Brown AC, Guler JL. 2018. SLOPE: A two-part method for the enrichment of ring stage Plasmodium falciparum parasites. bioRxiv.

51. Brown AC, Moore CC, Guler JL. 2020. Cholesterol-dependent enrichment of understudied erythrocytic stages of human Plasmodium parasites. Sci Rep 10:1–15.

52. Macosko EZ, Basu A, Satija R, Nemesh J, Shekhar K, Goldman M, Tirosh I, Bialas AR, Kamitaki N, Martersteck EM, Trombetta JJ, Weitz DA, Sanes JR, Shalek AK, Regev A, McCarroll SA. 2015. Highly Parallel Genome-wide Expression Profiling of Individual Cells Using Nanoliter Droplets. Cell 161:1202–1214.

53. Butler A, Hoffman P, Smibert P, Papalexi E, Satija R. 2018. Integrating single-cell transcriptomic data across different conditions, technologies, and species. Nat Biotechnol 36:411–420.

54. Stuart T, Butler A, Hoffman P, Hafemeister C, Papalexi E, Mauck WM 3rd, Hao Y, Stoeckius M, Smibert P, Satija R. 2019. Comprehensive Integration of Single-Cell Data. Cell 177:1888–1902.e21.

55. Becht E, McInnes L, Healy J, Dutertre C-A, Kwok IWH, Ng LG, Ginhoux F, Newell EW. 2018. Dimensionality reduction for visualizing single-cell data using UMAP. Nat Biotechnol https://doi.org/10.1038/nbt.4314.

24   56.   Finak G, McDavid A, Yajima M, Deng J, Gersuk V, Shalek AK, Slichter CK, Miller HW, McElrath MJ, Prlic

25   M, Linsley PS, Gottardo R. 2015. MAST: a flexible statistical framework for assessing transcriptional

26   changes and characterizing heterogeneity in single-cell RNA sequencing data. Genome Biol 16:278.

SUPPLEMENTAL CAPTIONS

Supplementary Figure S1- A.) UMAP plot of cell clusters generated before annotating clusters. B.) Heatmap showing top 50 genes enriched in each cell cluster, which identifies stage-specific gene expression profile. The cells in the heatmap are ordered by their developmental progression.

Supplementary Figure S2. UMAP projection plots of K13^{C580} and K13^{580Y} untreated *Pf* parasites.

Supplemental Figure S3. Pseudotime over the asexual cycle, separated by strain. Pseudotime was calculated using slingshot for each strain (top: K13^{C580}, bottom: K13^{580Y}) over the ring, trophozoite, and schizont stages. Each cell is assigned a pseudotime value. Lower values correspond to earlier developmental pseudotime (ring stage) and are colored purple. Pseudotime values then increase through the trophozoite stage (green) and the schizont stage (yellow).

Supplemental Figure S4. Vertex frequency clustering (VFC) results for rings. For each timepoint, the ring stage was sub-clustered using Vertex Frequency Clustering. The first row (A,C,E) of each timepoint is K13^{C580,} while the second row (B,D,F) is K13^{580Y}. For each timepoint, the clusters with the highest DHA likelihood were selected for differential expression. For example, the 2hr timepoint compared VFC cluster 8 in K13^{580Y} vs. VFC cluster 7 in K13^{C580}.

Supplemental Figure S5. Vertex frequency clustering (VFC) results for trophozoites. For each timepoint, the trophozoite stage was sub-clustered using Vertex Frequency Clustering. The first row (A,C,E) of each timepoint is K13^{C580,} while the second row (B,D,F) is K13^{580Y}. For each timepoint, the clusters with the highest DHA likelihood were selected for differential expression. For example, the 2hr timepoint compared VFC cluster 12 in K13^{580Y} vs. VFC cluster 13 in K13^{C580}.

Supplemental Figure S6. Differential expression utilizing MELD and high DHA likelihood subclusters

show similar results to traditional differential expression analysis in Figure 6. Ring stage (A) and

trophozoite stage (B) comparisons are plotted for each timepoint (2hr (left), 4hr (middle), and 6hr (right)).

Below each title, the plots are annotated by the number of differentially expressed genes (DEG) and the cell

number of each strain in the differential expression analysis. The x-axis plots the average log2FC, and the y

axis plots the -log10 BH-adjusted p value of each gene. Each dot is a gene, where grey colors indicate non-

significant genes at an absolute log2FC > 0.3 and a BH-adjusted p value of < 0.001. Green indicates genes

significant with the log2FC threshold, and blue indicates genes only significant by the BH-adjusted p value

threshold. Red indicates genes significant at both thresholds.

Supplemental Figure S7. GO term enrichment analysis of K13^{580Y} vs. K13^{C580} shows terms enriched

related to metabolism and translation in upregulated ring stage genes (A) while terms related to

nuclear regulation were downregulated in trophozoite stages (B). Utilizing differentially expressed genes

that were significant at a BH-adjusted p value < 0.001, genes were categorized as downregulated (average

log2FC < -0.3), neutral (-0.3 < average log2FC < 0.3), or upregulated (average log2FC > 0.3) after comparing

K13^{580Y} vs. K13^{C580}. GO term enrichment was performed (see Methods) at each timepoint, except for the 2hr

trophozoite timepoint, as there were too few differentially expressed genes.

Supplemental Figure S8. Optimization of MELD algorithm over different KNN (y axis) and beta (x axis)

parameters. As described in the Supplemental Methods, MELD was fit to each strain's normalized *cell x gene*

matrix separately and over a range of KNN and beta values. The red dot on each plot is the optimal

parameters used to run MELD for each line.

Supplementary Table S1- Top 50 genes enriched in each *P. falciparum* single cell cluster. Gene

expression profile associated with parasite stages.

Supplementary Table S2- Genes differentially expressed between the DHA treated K13^{C580} and the

DMSO treated K13^{C580} parasite over the course of 6hr DHA treatment.

833 **Supplementary Table S3-Genes differentially expressed between the DHA treated K13^{580Y} and the**

834 **DMSO treated K13^{580Y} parasite over the course of 6hr DHA treatment.**

836 **Supplementary Table S4- Genes differentially expressed between the DHA treated K13^{580Y} and the DHA**

837 **treated K13^{C580} parasite over the course of 6hr DHA treatment.**

Figure 1: Schema of the experimental workflow applied in this study to explore transcriptome changes upon DHA exposure. This schematic diagram illustrates the study design, points of sample collection and treatment, sample processing and single cell library preparation using the SeqWell platform. Both the wildtype (K13^{C580} strain) and mutant (K13^{580Y} mutant strain) *Pf* parasite cell lines were cultured separately for 48hrs. The control (untreated) parasite was collected for library prep. The controls for each parasite strain were then divided into two groups, some treated with 700 nM DHA (lethal dose of DHA) and DMSO. After DHA and DMSO treatment or exposure, samples were collected after every 2 hrs for single cell library preparation using SeqWell.

Figure 2: Single-cell gene expression identifies distinct life-cycle stages. A. UMAP of single-cell transcriptomes sampled from all stages of the life cycle, with cells colored according to their stage. **B**. UMAP representations displaying the expression of parasite stage specific gene expression markers. With this single-cell gene expression profile of the *Pf* parasite, we can identify new unknown stage specific parasite marker genes. **C**. Presence and absence of known drug resistant mutations between the two Cambodian parasite strains (mutant and wildtype). We validated that the only variant difference between the two strains used in this study was the K13^{C580Y} mutation is only present in the mutant strain. **D.** Distribution of parasite developmental stages across all sample groups (untreated controls, DHA and DMSO treated parasites).

Figure 3: Gene expression differences between K13^{580Y} and K13^{C580} parasites. A. *Kelch13* gene expression was observed to be downregulated in the K13^{580Y} parasite population compared to the K13^{C580} (BH-adjusted p-value<0.01 and log2FC=1.5). *PfKelch13* gene expression across most parasite stages (except for the schizonts) in each line, shows *PfKelch13* gene to be highly expressed in the K13^{C580} across all the stages compared to K13^{580Y}. **B**. Volcano plot showing DEGs in the untreated K13^{580Y} compared to untreated K13^{C580} on different parasite stages. Each red dot denotes genes differentially expressed between the K13^{C580} and the K13^{580Y} (such as *GARP* and *MAHRP1* which are upregulated in K13^{580Y} rings compared to K13^{C580} rings).

K13^{C580} Rings

Log$_2$ fold change cutoff, 0.5; p-value cutoff, 10e-3

K13^{C580} Trophozoites

Log$_2$ fold change cutoff, 0.5; p-value cutoff, 10e-3

Figure 4: Gene expression differences associated with K13^{C580} (Rings and Trophozoites) DHA response. Volcano plot showing DEGs in K13^{C580} DHA treated versus K13^{C580} DMSO treated (control) at 3 different timepoints. Each red dot denotes genes passing p-value and fold change difference (BH-adjusted p-value<0.01, average log2 fold-change>0.3) thresholds. Those on the right side of the plots are the upregulated genes induced due to DHA exposure. The red dots on the left side of the plots represent the downregulated genes passing the p-value and fold change difference between the DHA treated parasite and the DMSO treated parasite control (BH-adjusted p-value<0.05, average log2 fold-change<-0.3).

K13^{580Y} Rings

2hrs DHA
DEG = 89

4hrs DHA
DEG = 203

6hrs DHA
DEG = 211

Log$_2$ fold change cutoff, 0.5; p−value cutoff, 10e−3

K13^{580Y} Trophozoites

2hrs DHA
DEG = 140

4hrs DHA
DEG = 93

6hrs DHA
DEG = 222

Log$_2$ fold change cutoff, 0.5; p−value cutoff, 10e−3

Figure 5: Gene expression differences associated with K13^{580Y} (Rings and Trophozoites) response to DHA treatment *in vitro*. Volcano plot showing DEGs in K13^{580Y} mutant treated with DHA versus K13^{580Y} treated with DMSO (control) at 3 different timepoints. Each red dot denotes genes passing p-value and fold change difference (BH-adjusted p-value<0.05, average log2 fold-change>0.3) thresholds. Those on the right are the upregulated genes induced due to DHA exposure. The red dots on the left represent the downregulated genes passing the p-value and fold change difference thresholds between the DHA treated parasite and the DMSO treated parasite control (BH-adjusted p-value<0.01, average log2 fold-change<-0.3).

3
Figure 6: Gene expression differences between DHA treated K13^580Y^ and K13^C580^ parasites. A. *PfGARP* was observed to be
highly expressed in the K13^580Y^ parasite population compared to the K13^C580^ parasite population (BH-adjusted p-value<0.01 and
log2FC=3.8). Stratifying the *PfGARP* expression across the parasite stages in each strain shows *PfGARP* to be highly expressed in
K13^580Y^ across all the stages compared to K13^C580^. **B**. Volcano plot showing DEGs in K13^580Y^ treated with DHA versus K13^C580^ treated
with DHA at 3 different timepoints. Each red dot denotes genes passing p-value and fold change difference (BH-adjusted p-
value<0.001, average log2 fold-change>0.3) thresholds. The dots on the right are the upregulated genes induced due to DHA
0 exposure. The red dots on the left represent the downregulated genes passing the p-value and fold change difference thresholds
1 between the DHA treated parasite and the DMSO treated parasite control (BH-adjusted p-value<0.001, average log2 fold-change<-0.3).

Figure 7. Overall DHA transcriptional response trend is similar, but persistently elevated across the asexual cycle in the K13[580Y] strain. Pseudotime, calculated per strain through slingshot (39), is plotted on the x-axis while the DHA likelihood output from MELD is plotted on the y-axis for each facet of the figure. Cells are colored by stage annotation (A) or treatment condition (B). The left

6   column plots K13^{C580} cells while the right column plots K13^{580Y} cells. Each row is a different timepoint, with the first row being the 2 hour

7   timepoint, middle row being the 4 hour timepoint and the last row being the 6 hour timepoint.

Figure 8. Differential expression results show upregulation of *GARP* and stress response genes (*HSP70*, *GCN5*), perturbations in genes related to protein export (*FIKK* gene family, *EXP1*, *ETRAMP* genes, *KAHRP*, *RESA*), and downregulation of genes related to dormancy (SERA5, SERA6). The union of the top50 differentially expressed genes (by absolute log2FC) per time point between the highest VFC subcluster in $K13^{C580}$ compared with $K13^{580Y}$ in rings (A) and trophozoites (B). For each timepoint in (A) or (B), the top 50 genes by absolute log2FC were identified. The vectors of the top 50 genes were concatenated and filtered to unique genes. The average log2FC per timepoint is plotted, with "----" indicating genes that were not differentially expressed at that timepoint. An asterisk indicates genes that were significantly differentially expressed at an absolute log2FC > 0.3 and a BH-adjusted p-value of < 0.001.

Figure 9: Anti-PfGARP inhibits parasite growth in both artemisinin sensitive and resistant parasites. Artemisinin sensitive (K13^{C580}) and resistant (K13^{580Y}) parasites were synchronized to the ring stage, adjusted to 0.5-0.75% parasitemia and 2% hematocrit and incubated with anti-PfGARP (10% v/v) or normal mouse sera (10% v/v). Parasites were cultured for 72 hours followed by quantification of parasitemia by flow cytometry. Bars represent the mean of three technical replicates and error bars represent mean +/-

4 standard deviation. Results representative of 3 independent experiments (unpaired t-test with bonferroni adjustment, *p < 0.5, **p <
5 0.01).
6