## 2 Bacterial pathogens dynamic during multi-species infections

3    Marie-Anne Barny*, Sylvia Thieffry, Christelle Gomes de Faria, Elisa Thebault and Jacques Pédron

4    *Sorbonne Université, INRAE, IRD, CNRS, UPEC, Institute of Ecology and Environmental Sciences-*

5    *Paris (iEES-Paris), 4 place Jussieu, F-75252 Paris, France*

6    *\*E-mail : marie-anne.barny@sorbonne-universite.fr*

**8    Abstract**

Soft rot *Pectobacteriacea* (SRP) gathers more than 30 bacterial species that collectively rot a wide range

of plants by producing and secreting a large set of plant cell wall degrading enzymes (PCWDEs).

Worldwide potato field surveys identified 15 different SRP species on symptomatic plants and tubers.

The abundance of each species observed during outbreaks varies over space and time and the

mechanisms driving species shift during outbreak are unknown. Furthermore, multi-species infections

are frequently observed and the dynamics of these coinfections are not well understood.

To understand the dynamics of coinfections, we set up 16 different synthetic communities of 6 SRP

strains to mimic coinfections. The bacteria present in each tested community were representative of 2

different species, with 3 strains per species. These communities were inoculated in potato tubers or on

synthetic media and their outcome was followed by amplification and Illumina sequencing of the

discriminatory housekeeping gene *gap*A. We also compared disease incidence and bacterial

multiplication in potato tubers during mixed-species infection and single-species infection. A species

that was unable to induce disease in potato was efficiently maintained and eventually became dominant

in some of the communities tested, indicating that cheating can shape dominant species. Modeling

indicates that the cost of PCWDEs production and secretion, the rate of potato degradation and the

diffusion rate of degraded substrate could favor the cheater species. Interaction outcomes differed

between potato tuber and synthetic medium, highlighting the driving effect of environmental conditions,

with higher antagonistic interactions observed in potato tubers. Antagonistic interactions were strain

specific and not species specific. Toxicity interference was also observed within some communities,

allowing the maintenance of strains otherwise sensitive to toxic compounds. Overall, the results indicate

that intraspecific competition, cooperation through trophic interaction and toxicity interference

contribute to the maintenance of SRP diversity. The implications of these processes for epidemiological

surveillance are discussed.

**Introduction**

Pathogenic bacteria developed specialized weapons to attack their host, subvert host defenses and gain nutrient to develop inside their host. Among pathogens, some are specialists with a narrow host range and infect a single host while others have the capacity to infect many different hosts. A mirrored situation occurred with disease, while many diseases are defined by a single well known pathogenic species, other diseases could be triggered by a complex of bacterial species that share common bacterial weapons and could act either individually and/or collectively during disease development. When a disease is triggered by a species complex many questions concerning the dynamic of the complex are still unsolved. Notably, it is unclear to what extend the different species involved compete, cheat, and cooperate within the symptoms. As well it is also unclear if cheating, competition and cooperation behavior mainly occurs at the strain level or at the species level. Furthermore, as each disease cycle involved many different steps, it is unclear to which extend the competition/cooperation game change with changing environmental conditions and to what extend these interactions are shaping the bacterial diversity of pathogens.

Plant pathogens with a broad host range, such as the soft rot *Pectobacteriacea* (SRP) species complex, provide an interesting model to address these questions. SRP pathogens belong to three genera, *Pectobacterium, Dickeya* and *Musicola*, of the family *Pectobacteriacea* within the order *Enterobacteral*. Taxonomic studies recognize 38 species within the SRP group, of which 23 belong to the genus *Pectobacterium*, 12 belong to the *Dickeya* genus and 2 were formerly assigned to the *Dickeya* genus and now belong to the newly described genus *Musicola* (Charkowski 2018; Hugouvieux-Cotte-Pattat et al. 2021). All SRP attack the host plant by secreting a cocktail of plant cell wall degrading enzymes through the type 2 secretion system (T2SS), resulting in rotting and maceration of plant tissues on which the bacteria actively develop (Charkowski 2018; Hugouvieux-Cotte-Pattat et al. 2014). Due to their collective large host spectrum, SRP plant infections lead to severe economic losses and are collectively recognized as one of the ten most important bacterial plant pathogens (Mansfield et al. 2012; Ma et al. 2007; Portier et al. 2020). Among SRP species, some species like *Pectobacterium brasiliense*, *Pectobacterium carotovorum* or *Pectobacterium versatile* are described as broad range species isolated from a large number of plant species while other like *Pectobacterium parmentieri*, *Pectobacterium atrosepticum* or *Dickeya solani* have been mainly recorded on a single host (Ma et al. 2007; Portier et al. 2020; Hugouvieux-Cotte-Pattat et al. 2023). *P. versatile* is the most frequently isolated taxon among the SRP species, both on plants and in the environment, a characteristic that might be linked with its higher resistance to antibiotics than

other *Pectobacterium* spp. (Royer et al. 2022; Smoktunowicz et al. 2022; Ben Moussa et al.
2022; Portier et al. 2020) but this species is currently considered as a minor pathogen not
responsible for severe outbreaks but frequent companion species during disease or on
symptomless plants (Curland et al. 2021; Ma et al. 2022; Smoktunowicz et al. 2022). Some
newly described species like *Pectobacterium aquaticum* or *Pectobacterium quasiaquaticum*
have been isolated solely from river water and their potential host plants (if any) are currently
unknown raising the possibility that these species act are merely saprophytes on plant debris or
evolved toward hosts alternative to plant, for example as gut species of plant feeding insect or
water microfauna (Pédron et al. 2019; Ben Moussa et al. 2022, 2023). Species belonging to the
*Dickeya* genus are well known to infect ornamentals of high economical value but several
species could also be observed on major crops such as maize (*Dickeya zeae*) rice (*Dickeya*
*oryzae*) or potato (*Dickeya dianthicola* or *D. solani*) where they have been responsible of severe
outbreaks (Hugouvieux-Cotte-Pattat et al. 2023).

Potato production, estimated according to FAO to over 376 million tons worldwide in 2021, is
particularly exposed to SRP species. In potato field, SRP triggers a disease called blackleg. This
name referred to the necrosis observed at the stem basis of infected potato plant. Harvested
potato tubers could also develop a severe rotting, called soft rot, that could destroy the whole
harvest. Given its economic importance, potato is the most intensively surveyed crop attacked
by SRP and 15 different SRP species have been recorded on potato worldwide (Toth et al.
2021). In Europe and North America, *P. atrosepticum* was initially considered to be the primary
pathogen causing blackleg and soft rot in potatoes. However, from the 1970s, *D. dianthicola*
increased in prevalence, until the appearance of *D. solani* in the early 2000s while more recently
the spread of *P. parmentieri*, and *P. brasiliense* was observed (van der Wolf et al. 2021). The
succession of these prevalent species is not yet understood. Notably, aggressiveness of the
species is likely not the main factor explaining the prevalence of species in potato field (de
Werra et al. 2021). Other traits such as survival ability in the environment or
competition/cooperation abilities within the host plant are likely involved in the observed turn
over. Importantly, mix of SRP spp. during infection could represent up to 20% of recorded
symptoms (Ge et al. 2021; Motyka-Pomagruk et al. 2021; de Werra et al. 2021). These mixed
infections probably contribute to the evolution of epidemic clones through HGT transfer of trait
associated to the bacterial fitness (Khayi et al. 2015; Royer et al. 2022). Furthermore mix of
SRP spp. are also frequently reported on asymptomatic potato tubers (Degefu 2021;
Smoktunowicz et al. 2022). These latent infections likely influence the dynamic of disease
through cooperation and competition but this has not been investigated in details.

The aim of the present work was to experimentally assess the fate of mix infection during soft

rot within potato tubers. To understand the dynamic of coinfections, we set up 16 different

synthetic communities of 6 SRP strains to mimic coinfection. Bacteria present in each synthetic

community were representative of 2 different species, with 3 strains per species. These

communities were inoculated on potato tuber or on growth medium and the outcome of each

synthetic community was followed by amplification and Illumina sequencing of the

discriminative housekeeping gene *gap*A (Cigna et al. 2017; Ben Moussa et al. 2022). We also

compared disease rating and bacterial multiplication on potato tuber during coinfection and

single species infection. Overall, the results indicate that competition, cooperation and trophic

interactions shape the bacterial SRP community and allow for the maintenance of SRP

diversity.

**Materials and Methods**

**Strains and media used**

The strain used in this study are described in table S1. For each species, where possible, the

bacterial strains were selected to allow strain discrimination using the gapA barcode (see

below). For some species, such as *D. dianthicola*, *D. solani* and *P. polaris/P. parvum*, we did

not find discriminating strains in our collection (Table S2). This is probably due to the fact that

these latter species are quite homogeneous (Hugouvieux-Cotte-Pattat et al. 2023; Pasanen et al.

2020). The bacterial media used were LB medium without NaCl (hereafter LB: per liter 10 g

tryptone, 5 g yeast extract) and TSB (per liter: 1,7 g casein peptone, 0.3 g soya peptone, 0.3 g

NaCl, 0.25 g $K_2HPO_4$, 0.25 g glucose). When needed, 15 g agar per liter were added for solid

media.

**Inocula preparation**

The bacteria were plated on LB plates and incubated overnight at 28°C. A single colony was

then used to inoculate 2 ml of LB medium which was incubated overnight at 28°C, with

agitation at 120 rpm. This liquid culture (100 µl) was used to inoculate a 10% TSB agar plate,

which was incubated overnight at 28°C. The grown bacterial layer was then suspended in 50

mM phosphate buffer (pH 7) and adjusted to an $OD_{600nm}=1$. The 3 strains of the same species

were then mixed in a 1/1/1 ratio. An exception was the inoculum composed of *P. parvum* (2

strains) and *P. polaris* (1 strain), which were mixed together as the experiment was set up before

the description of the *P. parvum* species, a close relative of *P. polaris* (Pasanen et al. 2020).

The bacterial load of each 3 strains-mixtures was check by serial dilution and plating on TSB

agar plates. The species mixtures were then prepared by mixing two of the prepared strain-

mixtures in volumes 1/1. Each final species-mixture is thus the mixture of 6 strains adjusted to

$DO_{600nm}$=1 in the ratio 1/1/1/1/1/1, where the first strain-mixture corresponds to one of the nine

following species: *P. atrosepticum, P. carotovorum, P. parmentieri, P. versatile, P.*

*polaris/P.parvum, D. dianthicola, D. solani, P. versatile, P. aquaticum*, and the second strain-

mixture corresponds either to the species *P. aquaticum* or to the species *P. versatile*. A 2 ml

aliquot of each time zero species-mix inoculum was stored at -20°C until DNA extraction.

These 18 species-mixtures were then used to inoculate triplicates of 2 ml TSB liquid medium

(1 µl in 2 ml) or 10 potato tubers (10 µl per tuber). The inoculated TSB medium was incubated

at 26°C with shaking during 48h. Serial dilution and plating was performed to check the

bacterial growth at the end of the culture. The bacterial load at time zero (inoculum) and at the

end of the experiment were used to calculate the number of bacterial generation achieved ($\log_{10}$

(cfu end experiment/cfu T0)/$\log_{10}$ (2)). Cultures were then centrifuged and the pellet was kept

at -20°C until DNA extraction.

**In vitro bacterial competition assays**

Pairwise in vitro bacterial competition assays were performed between the *P. brasiliense, P.*

*versatile* and *P. aquaticum* strains (3 strains per species, 81 combinations tested). Briefly,

overnight TSB cultures of each strain were adjusted to an $DO_{600nm} = 0.15$. Samples of 5 µl of

the strain 1 were spotted onto freshly poured agar lawn inoculated with strain 2 (1 ml with 15

ml TSB agar medium) and plates were allowed to grow for 24 h at 28°C. If strain 1 inhibits the

growth of strain 2, a clear halo is observed around the spot of strain 1. Experiment was repeated

twice with similar results.

**Potato inoculation**

Three experiments were performed, the first one was devoted to analyze the mono-species

infection with the mix of the 3 strains/species. The second and third one involved mix of two

different species (6 strains) with either *P. aquaticum* species or *P. versatile* species challenged

by one of the other 8 species.

For each mix, 10 potato tubers were thoroughly washed, and surface disinfected in a 1 %

hypochlorite solution bath prior inoculation and each tuber was punctured with pipette tips

containing 10 µl of the inoculum. The tip was left in the tuber. As a negative control, a pipette

tip with 10µl sterile water was inserted into one tuber. Inoculated tubers were placed on paper

towel in plastic boxes and 50 mL of distilled water was poured into the bottom of the boxes and

the boxes were carefully closed to achieve relative humidity above 90 %. After 5 days of

incubation at 26°C, the tubers were cut and the entire macerating symptom was collected and

weighed. Part of the symptom (50 to 300 mg) was also weighed and suspended in 1 ml of sterile

phosphate buffer in order to count by serial dilution and plating the bacterial load present in the

symptom. To recover the bacteria within the symptoms, the rest of the symptom was suspended

with 7 ml of 50 mM phosphate buffer and stirred for 4 hours at ambient temperature. The tubes

were then left without shaking for 10 minutes to allow the excess starch to settle and 2 ml at the

surface of the suspension was recovered and centrifuged for 10 minutes at 10,000 g. The formed

pellet was kept at -80°C until DNA extraction.

**Statistical analysis**

One specie mix (*P. aquaticum/P. versatile*) was shared between the second and the third

experiments and this allowed to eliminate inter-experiment variation unrelated to the involved

species mix before statistical analysis. To do so we corrected the maceration weight observed

in the third experiment with the following coefficient X = (mean of *P. aquaticum/P.versatile*

10 potato tubers maceration weight of the 2$^{nd}$experiment)/ mean of *P. aquaticum/P.versatile* 10

potato tubers maceration weight results of the 3$^{rd}$ experiment) before statistical analysis.

Statistical analyses were performed in R version 4.0.2 using a Kruskal-Wallis test with p-value

< 0.05 followed by a post-hoc test (Dunn test with Bonferroni correction). Spearman correlation

calculation were performed on line (https://biostatgv.sentiweb.fr/?module=tests/spearman).

**DNA extraction, DNA amplification and sequencing**

For each of the 18 species-mix analyzed, DNA extraction was performed on 10 ml of the time

zero inoculum, on 2 ml of each of the TSB triplicate after 48h of culture and on 2 ml of 3 to 6

potato macerated tubers symptoms. The bacteria were pelleted by centrifugation and DNA

extraction was performed with the Wizard® genomic DNA purification kit (Promega).

Amplification and sequencing were performed at MR DNA (www.mrdnalab.com, Shallowater,

TX, USA). Briefly, the *gap*A 376 partial gene sequence was amplified using PCR primers

*gap*AF376: GCCCGTCTCACAAAGA and *gap*AR: TCRTACCARGAAACCAGTT) with

barcode on the forward primer. PCR using the HotStarTaq Plus Master Mix Kit (Qiagen, USA)

was performed under the following conditions: 94°C for 3 min, followed by 28 cycles of 94°C

for 30 s, 57°C for 40 s and 72°C for 1 min, after which a final elongation step at 72°C for 5 min

was performed. Amplified PCR products were checked in 2 % agarose gel to determine the

success of amplification and the relative intensity of bands. Multiple samples were pooled

together in equal proportions based on their molecular weight and DNA concentrations. Pooled

samples were purified using calibrated Ampure XP beads and used to prepare Illumina DNA

library. Sequencing was performed on a MiSeq sequencer following the manufacturer's

guidelines. Sequence data were processed using MR DNA analysis pipeline (MR DNA,

Shallowater, TX, USA). In summary, sequences were joined, depleted of barcodes and

sequences <150bp or with ambiguous base calls were removed. Reads were filtered based on

Q score and expected error probability and any read with a number of expected errors greater

than 1.0 were discarded.

**Illumina sequencing analysis**

After quality trimming, a total of 2,230,817 and 3,215,158 reads were obtained for *in vitro*

(TSB, 64 samples) and potato tuber (100 samples) assays respectively, with an average of

$34,857 \pm 963$ and $32,152 \pm 7538$ reads per samples respectively.

In order to quantify the distribution of the *Pectobacterium* and *Dickeya* strains after *in vitro*

growth or potato inoculation, reads were aligned to the corresponding amplified sequences of

the *gap*A gene, using the nucleotide-nucleotide Blastn tool (version 2.11.0+, e-value threshold

$10^{-5}$). Only reads with 100% full-length identity were used (547,866 reads, 24.6% of the total

reads and 1,213,517 reads, 37.7% of the total reads for *in vitro* and potato tuber assays

respectively. A few potato tubers were withdrawn of the analysis since unexpected *P.*

*parmentieri* sequences were observed, suggesting an initial presence of this pathogen as latent

species hosted on the tuber prior to the inoculation.

Bacterial distribution analysis was performed at the species level for all assays (sum of all

strains of the same species), and at the strain level when it was possible. In fact, for some

species, certain strains cannot be distinguished due to sequence similarities: *P. versatile*

CFBP6051 and A73.1, all *D. dianthicola* strains, all *D. solani* strains, *P. parmentieri* CFBP1338

and CFBP5382, *P. atrosepticum* CFBP1527 and CFBP1453, *P. carotovorum* CFBP6074 and

CFBP1402, all *D. polaris* strains.

**Model presentation**

The equations of the model are:

$dD_X/dt = k(D_R - D_X) - (gA - \epsilon)D_X X + cX/z + X$

$dD_A/dt = k(D_R - D_A) - g_A D_A A$

$dD_R/dt = -k(2D_R - D_X - D_A) - \lambda D_R$

$dX/dt = (g_A - \epsilon)\,D_X X - mX$

$dA/dt = g_A D_A A - mA$

with X the density of the species that produces enzymes able to degrade the substrate, A the

density of the cheater species that only consumes the degraded substrate without being able to

produce it, $D_X$ is the concentration of degraded substrate in the vicinity of species X (i.e. local

pool of degraded substrate of X), $D_A$ is the corresponding local substrate pool for A, and $D_R$ is

the concentration of degraded substrate in the regional environmental (hereafter regional pool

of degraded substrate), k is the diffusion rate of degraded substrate between local and regional

pools, c the maximum rate of enzymatic substrate degradation, z half-saturation constant of

enzymatic substrate degradation, gA consumption rate of degraded substrate without cost, $\epsilon$

cost of enzyme PCWDEs production and secretion on consumption rate of degraded substrate,

m Mortality rate of X and A, λ Loss of degraded substrate from regional pool. The mathematical

analysis of the model together with the parameter's values used in numerical simulations are

provided in supplementary file S1.

**Search for carbapenem synthesis and resistance**

The beta-lactam carbapenem is synthesized and degraded by the *car*ABCDEF genes while the

*car*GHR genes are involved in resistance to the produced carbapenem (Shyntum et al. 2019).

These genes were blast searched with *car*ABCDEFGHR genes of *P. brasiliense* CFBP6617 as

query among the 27 strains used to build up the synthetic communities using the genoscope

Microbial          Genome          Annotation          &          Analysis          Platform

(https://mage.genoscope.cns.fr/microscope/).

**Results:**

**Set up of the experiment**

A general view of the experimental set up is shown in Figure 1. In brief, each synthetic SRP

community was a mixture of 6 strains belonging to two different species. The mixture of two

species was preferred because the mixture of two different species is often observed in the field

(Degefu 2021; de Werra et al. 2021; Ge et al. 2021; Motyka-Pomagruk et al. 2021;

Smoktunowicz et al. 2022). The addition of three strains per species allows to disentangle the

"species" effects from the "strain" effects. The species involved were 8 species commonly

isolated from potato tubers, *P. versatile, P. atrosepticum, P. carotovorum, P. parmentieri, P.*
*polaris, P. brasiliense, D. solani, D. dianthicola*, and one species *P. aquaticum* that was only
isolated from fresh surface water. Synthetic communities were set up with either *P. versatile* or
*P. aquaticum* facing one of the others 8 species. *P. versatile* and *P. aquaticum* were chosen as
challenging species because of their very different distribution aera. *P. versatile* that occurs on
many different plants, is the main SRP species represented in bacterial collections and is also
frequently isolated from non-host environments, such as surface freshwater highlighting its
ecological success among SRP (Portier et al. 2019; Ben Moussa et al. 2022). In contrast, *P.*
*aquaticum* has not been isolated from plants and is frequently isolated from water (Portier et al.
2020; Ben Moussa et al. 2022). SRP communities were inoculated on TSB medium or on potato
tubers and the proportion of each species at the beginning and end of the experiment was
assessed by Illumina sequencing of the *gap*A discriminative gene marker (Cigna et al. 2017;
Ben Moussa et al. 2022). Symptoms occurring on potato tubers were also recorded for each 6
strains community and compared with symptoms observed with the mix of 3 strains of single-
species inoculation.

285
**Figure 1:** Set up of the experiment.

**Symptoms observation and bacterial multiplication in potato tubers**

Single-species infection resulted in various symptom development in potato tubers depending

on the inoculated species tested. Outliers were frequently observed and there was considerable

variability in response between the 10 inoculated potato tubers (Fig. 2A). The 10 potato tubers

inoculated with *P. aquaticum* showed hardly any symptoms and statistical analysis confirmed

that *P. aquaticum* developed less severe symptoms than *P. versatile, P. parmentieri, P. brasil-*

*iense* and *P. atrosepticum* (Fig. 2A). Bacterial multiplication was assessed and the number of

bacterial generations achieved within each inoculated tuber was calculated (Fig. 2B). It varied

from 0.9 to 13.7 with a median of 8.9 depending on the tuber inoculated. This multiplication

rate correlated with the weight of macerating symptoms (Spearman correlation 0.451, p-value

8.87E-14). After inoculation with *P. aquaticum*, bacterial multiplication within the tubers was

much lower than with other single inoculated single species, confirming *P. aquaticum* poor

ability to rot potato tuber (Fig. 2B) as already described (Ben Moussa et al, 2023). The addition

of either *P. versatile* or *P. aquaticum* to the inoculation mix did not affect the severity of the

symptoms observed compared to the other single species inoculation but *P. aquaticum* (Fig.

2A). Similarly, the severity of rot symptoms was similar for all co-inoculations, except for *P.*

*parmentieri* in competition with *P. versatile*, which showed a significantly higher maceration

rate than *P. aquaticum* in competition with *P. versatile* (Fig. 2A). Bacterial multiplication was

also unaffected by the mixed-species inoculation (Fig. 2B). In the following experiments 3 to 6

tubers from each SRP community were analyzed by amplification and Illumina sequencing of

the *gap*A barcode.

**Figure 2:** Symptoms and bacterial growth of the SRP synthetic communities on potato tubers.

a) Weight of macerate symptom b) bacterial growth within potato tuber. Single-species mixtures (3 strains) are shown in green. Two-species mixtures (6 strains) are shown in blue and yellow for *P. aquaticum*, and *P. versatile* consortia respectively. AQUA : *P. aquaticum* ; ATRO: *P. atrosepticum ;* BRASI: *P. brasiliense* ; CARO: *P. carotovorum*; PARM: *P. parmentieri* ; POL: *P. polaris* ; SOL: *D. solani*; VERS: *P. versatile* . The result of the statistical analysis is represented by the letters (a, b, c) at the top of the boxplots. The bacterial mixtures sharing the same letter(s) are not statistically different from each other (P>0.05 Kruskal Wallis followed by Dunn test with the Bonferroni correction). The strains used in each bacterial mixture are listed in Table S1.

**Species outcome within SRP communities inoculated in TSB medium**

Bacterial multiplication was assessed after 48 hours of growth on TSB medium and the number of bacterial generations obtained was calculated. It ranged from 4.2 to 9 generations and was repeatedly higher in the mixture containing *P. aquaticum* compared to the mixture containing *P. versatile* (Fig. S1). A total of 2,230,817 sequences were obtained after Illumina sequencing,

with an average of 34,857 ± 963 sequences per sample. The 376 pb *gapA* sequence allowed us

to distinguish all the inoculated species and to calculate the percentage of sequences assigned

to *P. aquaticum* or *P. versatile* within each SRP community at time zero and after 48 hours of

growth (Fig 3 A and B). At time zero, *P. aquaticum* or *P. versatile* represented 30% to 60% of

the inoculum in each mix containing another *Pectobacterium* species as expected for an even

inoculation of each strain. However, in the SRP community containing *Dickeya spp.*, *P.*

*aquaticum* and *P. versatile* made up more than 60% of the mixture. This deviation was

particularly important for the mix containing *D. dianthicola* as *P. versatile* and *P. aquaticum*

represented 78% and 79% respectively. A similar deviation was observed after counting the cfu

of each individual 3 species mix at time zero indicative of a higher inoculation of *P. versatile*

and *P. aquaticum* compared to the *D. dianthicola* species in the initial mix (data not shown).

This bias is likely due to the fact that *D. dianthicola* is highly mucoid impacting the optical

density-cfu relation. For all the others synthetic communities tested, no obvious bias was

detected and the strains are correctly equilibrated in the initial mix.

After 48 hours of growth, with an average number of generations of 6.7 after 48 hours of

growth, the initial population size represents only around 1% of the final population size. The

*gap*A barcode allowed to analyzed the proportion of each SRP species within the synthetic

community (Fig. 3 A and B). *P. aquaticum* and *P. versatile* were considered to drastically

outcompete the other SRP species if the mean percentage for the three replicates reached at

least 90% at the end of the experiment. They were considered to be drastically outcompeted by

the other SRP species if their mean percentage for the three replicate was inferior to 10%.

Finally, if the mean percentage of *P. aquaticum* or *P. versatile* was between 10% and 90%, *P.*

*aquaticum* and *P. versatile* were considered to be coexisting with their challenging SRP species.

Using this criterion most of the SRP community (12 out of 16) allowed coexistence of both

species after 2 days of growth in TSB medium (Fig. 3A and B). *P. aquaticum* outcompeted *D.*

*dianthicola* and coexisted with all the other challenging species. *P. versatile* was outcompeted

when co-inoculated with *P. brasiliense, P. carotovorum* or *D. solani* and it coexisted with the

other species at various levels.

**Figure 3:** Species outcome within SRP communities.

Percentage of *P. versatile* or *P. aquaticum* in mixed-species SRP communities after 48h TSB growth (A and B) or 5 days post inoculation within potato tuber (C and D). Grey bars: % of *P. aquaticum* or *P. versatile* within the inocula at time zero, Blue: bars % of *P. aquaticum* or *P. versatile* after 48 hours growth in TSB or 5 days within potato tubers. Three replicates were analysed after 48h growth in TSB and 3 to 5 potato tubers were analysed 5 days post inoculation. Bars : standard deviation. Species's acronym: AQUA : *P. aquaticum* ; ATRO: *P. atrosepticum ;* BRASI: *P. brasiliense* ; CARO: *P. carotovorum*; PARM: *P. parmentieri* ; POL: *P. polaris* ; SOL: *D. solani*; VERS: *P. versatile.*

**Species outcome within SRP communities inoculated within potato tubers**

A total of 3,215,158 sequences were obtained after Illumina sequencing, with an average of 32,152 ± 7538 sequences per sample. We calculated the percentage of reads assigned to *P. aquaticum* or *P. versatile* in the mixed-species SRP communities at time zero and after growth inside the potato tubers (Fig. 3 C and D). Using the same criteria as after TSB growth, coexistence was observed 5 days post infection in 6 out of the 16 mixed-species SRP communities inoculated in potato tubers. Within potato tubers, *P. aquaticum* was outcompeted in presence of *P. carotovorum, P. parmentieri* and *D. solani*, it outcompeted *D. dianthicola* and *P. polaris* and coexisted with *P. atrosepticum, P. brasiliense* and *P. versatile*. *P. versatile* was outcompeted in the presence of *P. parmentieri* and *D. solani*, it outcompeted *P. atrosepticum, D. dianticola* and *P.polaris/P.parvum* and coexisted with *P. aquaticum, P. brasiliense and P. carotovorum.*

The outcome of the competitions in potato tubers was different from that observed after growth

on TSB for 10 out of the 16 SRP communities tested (Table 1). For example, within potato

tubers *P. aquaticum* was outcompeted by *P. carotovorum, P. parmentieri* and *D. solani* while

it coexisted with each of these three species in TSB. Similarly, while *P. versatile* coexisted with

*P. brasiliense* and *P. carotovorum* in potato tubers, it was outcompeted by both species in the

TSB medium. As well, *P. aquaticum* and *P. versatile* were outcompeted by *P. parmentieri* in

potato tubers but both species coexisted with *P. parmentieri* in the TSB medium. Overall, out

of the 16 tested combinations, the two species coexisted twice as much in TSB medium (12/16

SRP communities tested) compared to within potato tuber (6/16 SRP communities tested).

**Table 1:** Comparison of species outcome after growth in TSB or potato tuber.

| Species mix | Potato tuber | TSB |
|---|---|---|
| AQUA-ATRO | 65.02 | 75 |
| AQUA-BRASI | 29.57 | 56.28 |
| AQUA-CARO | 0.77 | 23.8 |
| AQUA-DIAN | 96.81 | 94.88 |
| AQUA-PARM | 0.32 | 75.76 |
| AQUA-POL | 91.37 | 38.39 |
| AQUA-SOL | 9.29 | 40.21 |
| AQUA VERS | 38.29 | 37.14 |
| VERS-ATRO | 97.64 | 77.02 |
| VERS-BRASI | 53.72 | 0.13 |
| VERS-CARO | 37.44 | 6.42 |
| VERS-DIAN | 99.94 | 88.87 |
| VERS-PARM | 1.72 | 73.89 |
| VERS-POL | 99.81 | 72.64 |
| VERS-SOL | 3.23 | 5.93 |
| VERS-AQUA | 88.8 | 45.88 |

Blue: coexistence. Green: *P. versatile or P. aquaticum* dominates. Red: *P. versatile* or *P. aquaticum* was

outcompeted. The number indicated the % mean of *P. aquaticum* or *P. versatile* at the end of the

experiment

**Modelling of cheater persistence**

The SRP species *P. aquaticum* has been isolated from water and is not recorded on plant

(Pédron et al. 2019; Portier et al. 2020). We noticed here that the *P. aquaticum* strains have a

poor capacity to rot potato tubers and multiply within symptoms (Fig. 2). This is a previously

described phenotypic trait that could be linked to a reduced production of PCWDEs (Ben
Moussa et al. 2023). We observed nevertheless, that *P. aquaticum* when co-inoculated with
another species, could be maintained and even become dominant in several synthetic
communities tested (Fig. 3C and D). To investigate the potential mechanisms by which *P.*
*aquaticum* might persist when co-inoculated with other species and become even dominant, we
analysed a model describing the dynamics of two species in competition for one common
resource originating from the degradation of a substrate by the enzymatic activity of only one
of the two species (species X, Fig. 4A, Appendix model). The other species (A) is a cheater as
it does not contribute to the production of degraded substrate, and thus has no cost related to
enzyme production in that context. The model considers heterogeneous distribution of the
degraded substrate, with specific concentrations in the vicinity of each species (species local
pool of degraded substrate, Fig. 4A) resulting from the depletion of degraded substrate through
consumption for both species X and A, and from the creation of degraded substrate by enzyme
production for species X. Degraded substrate diffuses at a given rate between the two species-
specific local pools and a regional pool, determining the interactions between the species A and
X through their indirect effect on the concentration of degraded substrate in the regional pool.
The analysis of the model (Fig. 4 B and C, supplementary file S1) suggests that the cheater
species A can persist when it has sufficiently higher growth rate than the producer species X
(i.e. sufficient cost of enzyme production on the growth rate of species X) and when species X
is sufficiently efficient in producing degraded substrate (i.e. high maximum rate of substrate
degradation). Higher diffusion rate of degraded substrate between the local and regional pools
also favours species A persistence for lower enzyme production cost of species X (Fig. 4B and
C). The cheater species can become dominant (i.e. proportion of abundance of A higher than
0.5) when A has significantly higher growth rate than X (intermediate to high cost of enzyme
production) and when the rate of substrate degradation by X is high, especially when the
diffusion of degraded substrate in the environment are high. However, in such cases the
abundance of X is low (see supplementary file S1) and we are at the limit of the coexistence
domain between X and A. Indeed, as A has relatively very high growth rate in comparison to
X, this can lead to the exclusion of X by A through competition for the consumption of degraded
substrate, further leading to the extinction of A due to the loss of production of degraded
substrate by X (see supplementary file S1).

**Figure 4:** Modeling of cheater abundance.

A: Schematic description of the model analysing the coexistence between a species X that produces

degraded substrate and a cheater species A that only consumes such substrate. The producer species X

and its local pool of degraded substrate are represented by a brown circle and a dark yellow circle

respectively. The cheater species A and its local pool are represented by a grey and a light-yellow circle

respectively. The oval shape corresponds to the regional pool of degraded substrate in the environment.

B and C: effects of the rate of substrate production and the cost of degradation for X on coexistence and

relative abundance of A for two scenarios of flows of degraded substrate in the environment.

**Strains outcome within the SRP communities**

Beyond the species level, the 376 pb *gap*A Illumina sequenced fragment allowed the

differentiation of the 3 inoculated strains of *P. brasiliense* or *P. aquaticum* and one of the three

strains for *P. versatile*, *P. carotovorum*, *P. parmentieri* and *P. atrosepticum* (Table S2). This

allowed us to analyze the fate of these 10 strains within the 16 tested SRP communities within

potato tubers or TSB (Table 2, Fig. S2 and S3). Comparison of the fate of these 10 strains

showed that the outcome was mainly determined at the strain level (Fig. 4, see Fig. S2, Fig. S3

for the 32 tested combinations). In 20 cases, within the 3 strains of a single species, the fate of

the 3 strains was different. For example, within the *P. aquaticum/P. brasiliense* community in

which the six strains could be distinguished, after growth in TSB, one strain belonging to the

*P. aquaticum* species (A212) and two strains belonging to the *P. brasiliense* species (CFBP5381

and CFBP3230) were outcompeted, while two strains of *P. aquaticum* (A101 and A127) and

one strain of *P. brasiliense* (CFBP6617) coexisted (Fig. 5). The result of the same community

grown on potatoes was different, allowing the coexistence of one *P. brasiliense* strain

(CFBP6617) and only one *P. aquaticum* strain (A101) (Fig. 5). Also, within the *P. aquaticum/P.*

*parmentieri* community, the *P. parmentieri* strain CFBP8475 outcompeted the 5 other strains

on potato tubers, but this was not the case on TSB (Fig. 5).

**Figure 5:** Examples of strains outcome within SRP communities.

The proportion of each strain in 3 different mixed-species SRP communities after 48h growth in TSB growth or 5 days post inoculation within potato tubers is indicated in the legend of each graph. *P. aquaticum-P. brasiliense* strains consortium in TSB (A) or within potato tuber (B) *P. versatile - P. brasiliense* strains consortium in TSB (C) or within potato tuber (D). *P. aquaticum - P. parmentieri* strains consortium in TSB (E) or within potato tuber (F). In each graph the first column represents the % of each strain within the inoculum (T0), the second column represents the outcome of the inoculated community after 48h in TSB or 5 days post potato tubers inoculation. Three replicates were analysed after 48h growth in TSB and 3 to 5 potato tubers were analysed 5 days post inoculation. Bars: standard deviation. Species's acronym: AQUA : *P. aquaticum* ; ATRO: *P. atrosepticum ;* BRASI: *P. brasiliense* ; CARO: *P. carotovorum*; PARM: *P. parmentieri* ; POL: *P. polaris* ; SOL: *D. solani*; VERS: *P. versatile.* The names of each strain follow the species'names.

469 The strains had different outcomes when grown on potato tubers and TSB. Overall, the

470 outcompeted strains made up a higher percentage within the potato tuber (66%) than after

471 growth on TSB (32%) (Table 2).

473 **Table 2:** Outcome of individual strains within the tested SRP communities.

| Single strain discriminated with the gapA barcode | nb of tested synthetic communities | potato tubers | | | TSB | | |
|---|---|---|---|---|---|---|---|
| | | >90% | <10% | 10-90% | >90% | <10% | 10-90% |
| VERS CFBP5663 | 9 | 0 | 9 | 0 | 0 | 3 | 6 |
| AQUA A101 | 9 | 0 | 4 | 5 | 0 | 0 | 9 |
| AQUA A127 | 9 | 0 | 7 | 2 | 0 | 2 | 7 |
| AQUA A212 | 9 | 0 | 7 | 2 | 0 | 4 | 5 |
| ATRO CFBP7375 | 2 | 0 | 1 | 1 | 0 | 2 | 0 |
| BRASI CFBP5381 | 2 | 0 | 1 | 1 | 0 | 2 | 0 |
| BRASI CFBP3230 | 2 | 0 | 2 | 0 | 0 | 2 | 0 |
| BRASI CFBP6617 | 2 | 0 | 0 | 2 | 1 | 0 | 1 |
| CARO CFBP2046 | 2 | 0 | 2 | 0 | 0 | 1 | 1 |
| PARM CFBP8475 | 2 | 1 | 0 | 1 | 0 | 0 | 2 |
| Total | 48 | 1 | 33 | 14 | 1 | 16 | 31 |
| % | | 2% | 66% | 28% | 2% | 32% | 62% |

VERS: *P. versatile*; AQUA: *P. aquaticum*; ATRO: *P. atrospecticum*; BRASI: *P. brasiliense*; CARO:
*P. carotovorum*; PARM: *P. parmentieri*.

However, the dominance of a single strain over the other 5 strains of the SRP community was

rarely observed both within tubers and in TSB medium: only the *P. parmentieri* strain

CFBP8575 dominates in potato tubers when associated with *P. aquaticum* and, in TSB medium,

the *P. brasiliense* strain CFBP6617 dominates when associated with *P. versatile* (Table 2, Fig.

5). In the other synthetic communities tested, on TSB or within the potato tubers, coexistence

of at least two strains was observed in 23 of the 32 combinations tested, the remaining 7

combinations were inconclusive because we could not differentiate some strains with the *gap*A

barcode (Fig. S2 and S3).

**Focus on the SRP communities involving *P. brasiliense***

The strain *P. brasiliense* CFBP6617, also called 1692, is a model strain known to produce,

among other toxic compounds, a carbapenem antibiotic (Holden et al. 1998; Shyntum et al.

2019). We set up pairwise *in vitro* competition assays between the 6 strains of *P. brasiliense*

and *P. aquaticum* to check for carbapenem production. This showed that the *P. brasiliense*

strain CFBP6617 inhibited the growth of the other two *P. brasiliense* strains, CFBP3220 and

CFBP5381, and of the two *P. aquaticum* strains, A212 and A127 (Fig. 6). The *P. aquaticum*

strain A101 was not inhibited by the *P. brasiliense* strain CFBP6617. Consistent with this latter

result, the *car*G, *car*H and *car*F genes conferring resistance to carbapenem (Coulthurst et al.

2005) were only detected in the genome of the strain *P. aquaticum* A101. During our pairwise

inhibition tests, we also observed that the *P. aquaticum* strain A127 produces an unknown

compound that inhibits the growth of the strain *P. brasiliense* CFBP6617. The radius of action

of this unknown toxin is smaller than the one induced by the carbapenem production by the *P.*

*brasiliense* strain CFBP6617 (Fig. 6).

**A**

**B**

|  | strain | Spot | |
|---|---|---|---|
|  |  | BRASI 6617 | AQUA A127 |
|  | BRASI 6617 | R | s |
|  | BRASI 3220 | S | s |
|  | BRASI 5381 | S | s |
| **Lawn** | VERS 6051 | R | s |
|  | VERS 5663 | R | R |
|  | VERS A73 | R | s |
|  | AQUA A212 | S | R |
|  | AQUA A127 | S | R |
|  | AQUA A101 | R | R |

**Figure 6:** Pairwise competition assays.

Pairwise competition assays were setup between the 9 strains belonging to the species *P. brasiliense*, *P.*

*aquaticum* and *P. versatile*. A: example of inhibition assay showing that the *P. brasiliense* strain

CFBP3230 (lawn BRASI 3220) is inhibited by the *P. brasiliense* strain CFBP6617 (spot BRASI 6617)

and also by the *P. aquaticum* strain A127 (spot AQUA A127). Note that the radius of inhibition triggered

by the *P. aquaticum* strain A127 is smaller than the one triggered by the *P. brasiliense* strain CFBP6617.

B: inhibition observed for the 81 pairwise tested competitions. R: no inhibition; S: the strain in the lawn

is inhibited by the strain in the spot with a large radius of inhibition; s: the strain in the lawn is inhibited

by the strain in the spot with a small radius of inhibition; BRASI: strains belonging to the *P. brasiliense*

species; AQUA: strains belonging to the *P. aquaticum* species; VERS: strains belonging to *the P.*
*versatile* species.

These observed pairwise inhibitions could probably explain the outcome of the *P. aquaticum/P.*
*brasiliense* community after growth on TSB medium (Fig. 5), as the 3 strains CFBP 6617, A127
and A101 were efficiently maintained after growth on this medium, while the 3 other strains
were outcompeted. The result of the same community on potato tubers was slightly different as
only 2 strains, CFBP6617 and A101, were efficiently maintained (Fig. 5). This could probably
be explained by the fact that the unknown toxic compound that counteract the growth of *P.*
*brasiliense* strain CFBP6617 is not produced by strain A127 or inefficient within potato tuber,
while the carbapenem is still produced within potato tuber. Surprisingly, within the *P.*
*versatile/P. brasiliense* community, we observed that the *P. brasiliense* strain CFBP5381 was
maintained within the potato tuber despite its sensitivity to the carbapenem produced by the *P.*
*brasiliense* strain CFBP6617 (as revealed with the *P.aquaticum/P. brasiliense* SRP community)
(Fig. 5). This suggests that at least one of the two *P. versatile* strains CFBP6051 or A73 still
observed in the potato tubers at the end of the experiment is interfering with the toxic effect of
*P. brasiliense* CFBP6617 to allow the growth of *P. brasiliense* CFBP5381. The mechanism that
allows the *P. versatile* strains to resist to carbapenem toxicity of *P. brasiliense* CFBP6617 is
unknown, as the two strains do not carry the carbapenem resistance genes *car*G, *car*H and *car*F
in their genomes (Coulthurst et al. 2005). Whatever the mechanism involved, it is either not
induced or not efficient in TSB medium, as CFBP6617 dominates over the other 5 strains of
the *P.brasiliense/P. versatile* community TSB medium.

**Discussion:**

The dynamic of microbial communities is an important research field. Theoretical studies
showed that cooperative trophic interaction generally leads to the steady coexistence of distinct
bacterial species, that organize themselves to create a community-level metabolism that best
exploits the nutrients present (Gralka et al. 2020; Taillefumier et al. 2017). This is exemplified
by synthetic metabolic consortia set up for various biotechnological applications that regroup
highly diverse bacteria with different metabolic capacity that could be stably maintained. As
well, metabolic cross feeding could allow maintenance of bacterial species otherwise
outcompeted in natural environment (Franza et al, 2016). The maintenance of microbial
diversity within pathogenic bacterial species complex cannot follow the same rule as, in that

case, the microbes within these species complex are closely phylogenetically related with
highly similar metabolic properties. This metabolic closeness implies a strong substrate overlap
likely associated with strong competition between species within the same niche (Weiss et al.
2022). It is therefore not straightforward to understand which mechanisms allow the
maintenance of diversity within these pathogenic species complexes. The bacterial SRP species
complex is probably one of the most diverse species complexes of plant pathogenic bacteria. It
gathers 38 accepted or proposed species and at least 15 SRP species have been identified on the
potato host, with frequent co-infection with at least 2 species (Toth et al. 2021). The work
presented here takes advantage of *gap*A barcoding Illumina sequencing to analyze the bacterial
dynamics of 16 different synthetic SRP communities involving 6 strains belonging to two
different species.

The outcome of the 16 synthetic communities was highly dependent on environmental
conditions and strongly differed between potato tubers and TSB stirred liquid medium. The
striking outcome differences of the synthetic community in the two different environments
should warns us on the use of synthetic liquid medium to test consortia stability when these
consortia should be used in a different environment. Notably, coexistence between the two
species was twice more frequent in TSB liquid medium than within potato tubers and
outcompeted strains were more numerous within the potato tubers than after growth on TSB
liquid medium. Competition through production of secreted toxic compounds, such as
carbapenem (Coulthurst et al. 2005; McGowan et al. 1997), carotovoricin (Nguyen et al. 1999;
Itoh et al. 1978), bacteriocins (Chuang et al. 2007; Chan et al. 2011; Roh et al. 2010; Grinter et
al. 2012), NRPS-PKS secondary metabolites (Brual et al. 2023; Cheng et al. 2013) have been
described in SRP. The production of many of these compounds is enhanced by plant extracts
or in potato tubers (Bellieny-Rabelo et al. 2019; Mattinen et al. 2007) and this could explain
that more strains are outcompeted during growth in potato tubers. Furthermore, competition
mechanisms relying on cell to cell contact, like the type 6 secretion system (T6SS), are also
described in SRP (Shyntum et al. 2019; Arizala and Arif 2019) but are likely inefficient in liquid
stirred medium. Modelling studies showed that in a structured environment such as that found
in potato tubers, the production of antimicrobial molecules could be beneficial even if the
producer is rare (Chao and Levin, 1981; Durrett and Levin, 1997; Gardner et al., 2004). The
enhanced production of antimicrobial molecules and their higher efficiency within a structured
environment, such as potato tubers, likely explain the higher competition observed within
potato tubers compared to liquid TSB medium.

As the outcome of the community is highly dependent on the environment, it is likely that each

of the communities studied will evolve as the infection progresses in the field under natural

conditions. In particular, each of the analysed communities is likely to evolve differently within

the stem during blackleg infection in the field, as the environmental conditions within the potato

tuber and within the potato stem are drastically different. Contrasting SRP fitness in stem and

tuber has been reported previously (Blin et al. 2021) and further work is required to analyse

how our synthetic consortia will evolve in potato stem.

Our results also illustrate that the competition is strain-specific rather than species-specific, as

in many combinations we observed that the fate of the 3 strains of the same species was

different. This is likely related to the fact that toxin production and toxin resistance are

functional traits associated with strains rather than with species and are generally encoded

within the accessory genomes of each species, as exemplified here by biosynthesis and

resistance to carbapenem (Shyntum et al. 2019). In most combinations, the *P. brasiliense* strain

CFBP6617 outcompeted the other two *P. brasiliense* strains, consistent with the observed

pairwise *in vitro* competition and the presence of the carbapenem cluster in the strain genomes.

However, within the *P. versatile/P. brasiliense* synthetic community, we observed an

interference towards the toxic effect of *P. brasiliense* CFBP6617 on *P. brasiliense* strain

CFBP5381, likely mediated by *P. versatile* strains. This interference may involve the secreted

β-lactamase commonly found in *P. versatile* strains which may inactivates the carbapenem

antibiotic (Royer et al. 2022). Further work is needed to test this hypothesis. *In vitro* pairwise

inhibition indicated that "face-to-face" competition may also occur in some of the synthetic

SRP communities tested, as for example *P. brasiliense* strain CFBP6617 is toxic to *P.*

*aquaticum* strain A127, but the reverse is also true. The complexity of the competitive interplay

between strains is enhanced by the number of toxic compounds and resistance mechanisms

known to be produced by SRP. Theoretical models highlight that chemical warfare between

microbes promote diversity (Kelsic et al. 2015; Czárán et al. 2002). The fact that each SRP

strains carries its own set of toxic compounds and resistance mechanisms could therefore

contribute to the maintenance of multiple strains as observed with the majority of the 16

synthetic communities tested. The complexity of the competition between SRP strains rather

than species probably also explains the temporal and spatial succession of different SRP species

in epidemic outbreaks.

Beyond the competition mechanisms, the virulence factors of SRP may also contribute to the

maintenance of SRP diversity, as the main virulence factors of SRP, i.e. the ability of SRP to

degrade the plant cell wall through the secretion of PCWDEs, can be considered as a public

good used to feed the whole community. In potato tubers, the important release of nutrients

once SRP express their main virulence factors allows the development of associated

commensals (Kõiv et al. 2015). Here, we observed that *P. aquaticum* strains were efficiently

maintained in the population within potato tubers in 5 out of 8 synthetic communities tested.

We also observed that *P. aquaticum* strains were unable to produce soft rot disease symptoms

and to multiply efficiently in the potato tubers when inoculated alone. The inability of *P.*

*aquaticum* strains to produce soft rot symptoms when inoculated alone strongly suggests that

the T2SS and associated virulence factors are not properly induced within the potato tuber and,

as such, *P. aquaticum* does not share the cost of producing virulence factors. However, when

mixed with other species, *P. aquaticum* strains could eventually become dominant, suggesting

that they can behave as cheaters, benefiting greatly from nutrient sharing. Several independent

mechanisms could be proposed to explain this efficient cheating behavior. Firstly, the low

expression of T2SS and the associated virulence factors and the smaller genome of *P.*

*aquaticum* strains compared to that of other SRP species may allow a more rapid multiplication

at the expense of the other SRP species. In support of this hypothesis, we observed here that in

TSB medium the 8 synthetic communities with *P. aquaticum* have a tendency to grow at higher

density in TSB medium than the 8 synthetic communities with *P. versatile*. It has also been

reported that *P. aquaticum* strains have increased mobility compared to that of other SRP

species (Ben Moussa et al. 2023). This may also influence the ability of *P. aquaticum* strains to

access nutrients within the symptoms. *P. aquaticum* has never been isolated from potatoes in

the field or in storage, and their cheating behavior observed here could therefore be considered

an artefact. Nevertheless, the observation of this cheating behavior suggests that other SRP

species could also behave as cheaters in the field, as plants infected with several SRP species

are often observed. This is supported by the fact that the repertoire and regulation of virulence

factors is variable among SRP species (Babujee et al. 2012; Jonkheer et al. 2021; Arizala and

Arif 2019; Li et al. 2018), likely allowing for a trade-off between growth and the ability to

cheat. Interestingly, our modelling suggests that the more efficient the SRP strain is at

decomposing potato tubers, the higher the abundance of the cheaters in the symptoms. This

greater abundance of the cheaters may affect the ability to detect the SRP strain responsible for

the symptoms and increase the risk of not identifying the initial SRP strain responsible for the

epidemic. Therefore, field sampling plans should be calibrated to account for this risk.

In summary, the dynamics of the 16 synthetic SRP communities presented here highlight some

possible mechanisms allowing both the maintenance of high SRP diversity within the SRP

species complex and the regular species shifts responsible for outbreaks observed in the field.

Cooperation for virulence, exemplified by cheating behavior of *P. aquaticum*, and the
complexity of competition between strains, are two mechanisms that likely contribute to the
maintenance of the SRP complex diversity. The fact that competition occurs at the strain level
rather than the species level probably also explains the regular appearance of new species
responsible for epidemic outbreaks. This suggests that deep sequencing or *gap*A barcoding of
plant symptoms are important to understand epidemic dynamic and rapidly identify new strains
responsible for outbreaks in the field.
652

**Authors contributions**
**Jacques Pédron**: Conceptualization (equal); supervision (equal): data curation (lead): formal
analysis (equal), methodology (equal), visualization (equal), writing original draft (equal), writ-
ing-review and editing (equal)
**Christelle Gomes de Faria**: investigation (equal), writing-review and editing (equal)
**Sylvia Thieffry**: Methodology (equal), conceptualization (equal), writing-review and editing
(equal)
**Elisa Thebault**: Supervision (equal), conceptualization (lead), methodology (equal), visualiza-
tion (equal), writing-review and editing (equal).
**Marie-anne Barny**: Funding acquisition (lead), supervision (equal), investigation (equal), vis-
ualization (equal), writing original draft (equal), writing-review and editing (equal)
664

**Acknowledgements:**
We thank Denis Faure and Jérémy Cigna for helpful discussions during the setup of the
experiment. We also thank Angel Alfon for performing the pairwise competition tests and Hajar
Ben Moussa for statistical analysis and vizualisation of Fig. 2. This work was possible due to
funding contract ANR-17-CE32-0004.

**Conflict of interest statement**
The authors declare they have no conflict of interest relating to the content of this article.
673

**Data, script and code availability**
Supplemental data are available in the tables S1 and S2, the figures S1, S2, S3 and the file S1
for model details. Sequenced data and the script used to analyse the sequences. could be found
in the zenodo file DOI 10.5281/zenodo.10212828 (https://zenodo.org/records/10212829). The

scripts used to perform the statistical analysis could be found in the zenodo file

https://doi.org/10.5281/zenodo.10404740.

**Supplementary information**

FigS1: Bacterial multiplication in TBS medium

FigS2: Result at strain level of the 16 synthetic communities after growth on TSB.

FigS3: Result at strain level of the 16 synthetic communities 5 days post inoculation on potato

tubers.

File S1: supplemental modeling

Table S1: strains description

Table S2: Intra-species strain discrimination with the 341 nt gapA barcode

All supplementals are available in the zenodo file

https://zenodo.org/doi/10.5281/zenodo.10886378

**References**:

1. Arizala, D., and Arif, M. 2019. Genome-wide analyses revealed remarkable heterogeneity in pathogenicity determinants, antimicrobial compounds, and CRISPR-Cas systems of complex phytopathogenic genus *Pectobacterium*. Pathogens. 8:247. DOI:10.3390/pathogens8040247

2. Babujee, L., Apodaca, J., Balakrishnan, V., Liss, P., Kiley, P. J., Charkowski, A. O., et al. 2012. Evolution of the metabolic and regulatory networks associated with oxygen availability in two phytopathogenic enterobacteria. BMC Genomics. 13:110. DOI:10.1186/1471-2164-13-110

3. Bellieny-Rabelo, D., Tanui, C. K., Miguel, N., Kwenda, S., Shyntum, D. Y., and Moleleki, L. N. 2019. Transcriptome and comparative genomics analyses reveal new functional insights on key determinants of pathogenesis and interbacterial competition in *Pectobacterium* and *Dickeya* spp. Appl. Environ. Microbiol. 85 DOI: 10.1128/AEM.02050-18

4. Ben Moussa, H., Bertrand, C., Rochelle-Newall, E., Fiorini, S., Pédron, J., and Barny, M.-A. 2022. The diversity of soft rot *Pectobacteriaceae* along the Durance River stream in the south-east of France revealed by multiple seasonal surveys. Phytopathology DOI :10.1094/PHYTO-12-21-0515-R

5. Ben Moussa, H., Pédron, J., Hugouvieux-Cotte-Pattat, N., and Barny, M. 2023. Two species with a peculiar evolution within the genus *Pectobacterium* suggest adaptation to a new environmental niche. Environ. Microbiol. :1462-2920.16479. DOI: 10.1111/1462-2920.16479

6. Blin, P., Robic, K., Khayi, S., Cigna, J., Munier, E., Dewaegeneire, P., et al. 2021. Pattern and causes of the establishment of the invasive bacterial potato pathogen *Dickeya solani* and of the maintenance of the resident pathogen *D. dianthicola*. Mol. Ecol. 30:608–624. • DOI: 10.1111/mec.15751

7. Brual, T., Effantin, G., Baltenneck, J., Attaiech, L., Grosbois, C., Royer, M., et al. 2023. A natural single nucleotide mutation in the small regulatory RNA ArcZ of *Dickeya solani* switches off the antimicrobial activities against yeast and bacteria ed. Pascale Romby. PLOS Genet. 19:e1010725. DOI: 0.1371/journal.pgen.1010725

8. Chan, Y.-C., Wu, J.-L., Wu, H.-P., Tzeng, K.-C., and Chuang, D.-Y. 2011. Cloning, purification, and functional characterization of Carocin S2, a ribonuclease bacteriocin produced by *Pectobacterium carotovorum*. BMC Microbiol. 11:99. DOI: 10.1186/1471-2180-11-99

9. Charkowski, A. O. 2018. The changing face of bacterial soft-rot diseases. Annu. Rev. Phytopathol. 56:269–288. DOI: 10.1146/annurev-phyto-080417-045906

10. Cheng, Y., Liu, X., An, S., Chang, C., Zou, Y., Huang, L., et al. 2013. A Nonribosomal peptide synthase containing a stand-alone condensation domain is essential for

phytotoxin zeamine biosynthesis. Mol. Plant-Microbe Interactions. 26:1294–1301.
DOI:10.1094/MPMI-04-13-0098-R

11. Chuang, D., Chien, Y., and Wu, H.-P. 2007. Cloning and expression of the *Erwinia*
*carotovora* subsp. *carotovora* gene encoding the low-molecular-weight bacteriocin
Carocin S1. J. Bacteriol. 189:620–626. DOI: 10.1128/JB.01090-06

12. Cigna, J., Dewaegeneire, P., Beury, A., Gobert, V., and Faure, D. 2017. A gapA PCR-
sequencing assay for identifying the *Dickeya* and *Pectobacterium* potato pathogens.
Plant Dis. 101:1278–1282. DOI: 10.1094/PDIS-12-16-1810-RE

13. Coulthurst, S. J., Barnard, A. M. L., and Salmond, G. P. C. 2005. Regulation and biosyn-
thesis of carbapenem antibiotics in bacteria. Nat. Rev. Microbiol. 3:295–306.
DOI:10.1038/nrmicro1128

14. Curland, R. D., Mainello, A., Perry, K. L., Hao, J., Charkowski, A. O., Bull, C. T., et al.
2021. Species of *Dickeya* and *Pectobacterium* isolated during an outbreak of blackleg
and soft rot of potato in northeastern and north central united states.
Microorganisms. 9:1733. DOI: 10.3390/microorganisms9081733

15. Czárán, T. L., Hoekstra, R. F., and Pagie, L. 2002. Chemical warfare between microbes
promotes biodiversity. Proc. Natl. Acad. Sci. 99:786–790. DOI: 0.1073/pnas.01239989

16. Degefu, Y. 2021. Co-occurrence of latent *Dickeya* and *Pectobacterium* species in
potato seed tuber samples from northern Finland. Agric. Food Sci..
DOI:10.23986/afsci.101446

17. Ge, T., Ekbataniamiri, F., Johnson, S. B., Larkin, R. P., and Hao, J. 2021. Interaction
between *Dickeya dianthicola* and *Pectobacterium parmentieri* in Potato Infection
under Field Conditions. Microorganisms. 9:316. DOI:
10.3390/microorganisms9020316

18. Gralka, M., Szabo, R., Stocker, R., and Cordero, O. X. 2020. Trophic interactions and
the drivers of microbial community assembly. Curr. Biol. 30:R1176–R1188.
DOI:10.1016/j.cub.2020.08.007

19. Grinter, R., Milner, J., and Walker, D. 2012. Ferredoxin containing bacteriocins sug-
gest a novel mechanism of iron uptake in *Pectobacterium* spp ed. Pierre Cornelis.
PLoS ONE. 7:e33033. DOI: 10.1371/journal.pone.0033033

20. Holden, M. T. G., McGowan, S. J., Bycroft, B. W., Stewart, G. S. A. B., Williams, P., and
Salmond, G. P. C. 1998. Cryptic carbapenem antibiotic production genes are
widespread in *Erwinia carotovora*: facile trans activation by the carR transcriptional
regulator. Microbiology. 144:1495–1508. DOI: 10.1099/00221287-144-6-1495

21. Hugouvieux-Cotte-Pattat, N., des-Combes, C. J., Briolay, J., and Pritchard, L. 2021.
Proposal for the creation of a new genus *Musicola* gen. nov., reclassification of

*Dickeya paradisiaca* (Samson et al. 2005) as *Musicola paradisiaca* comb. nov. and de-
scription of a new species *Musicola keenii* sp. nov. Int. J. Syst. Evol. Microbiol. 71 DOI:
22.735 10.1099/ijsem.0.005037
791

792 23. Hugouvieux-Cotte-Pattat, N., Condemine, G., and Shevchik, V. E. 2014. Bacterial
793   pectate lyases, structural and functional diversity: Bacterial pectate lyases. Environ.
794   Microbiol. Rep. 6:427–440. DOI: 10.1111/1758-2229.12166
795

796 24. Hugouvieux-Cotte-Pattat, N., Pédron, J., and Van Gijsegem, F. 2023. Insight into bio-
797   diversity of the recently rearranged genus *Dickeya*. Front. Plant Sci. 14:1168480. DOI:
798   10.3389/fpls.2023.1168480
799

800 25. Itoh, Y., Izaki, K., and Takahashi, H. 1978. Purification and characterization of a bac-
801   teriocin from *Erwinia carotovora*. J. Gen. Appl. Microbiol. 24:27–
802   39.DOI:10.2323/jgam.24.27
803

804 26. Jonkheer, E. M., Brankovics, B., Houwers, I. M., van der Wolf, J. M., Bonants, P. J. M.,
805   Vreeburg, R. A. M., et al. 2021. The *Pectobacterium* pangenome, with a focus on
806   *Pectobacterium brasiliense*, shows a robust core and extensive exchange of genes
807   from a shared gene pool. BMC Genomics. 22:265. DOI: 10.1186/s12864-021-07583-5
808

809 27. Kelsic, E. D., Zhao, J., Vetsigian, K., and Kishony, R. 2015. Counteraction of antibiotic
810   production and degradation stabilizes microbial communities. Nature. 521:516–519.
811   DOI:10.1038/nature14485
812

813 28. Khayi, S., Blin, P., Jacques Pédron, J., Chong, T.-M., Chan, K.-G., Mohieddine Moumni,
814   M., et al. 2015. Population genomics reveals additive and replacing horizontal gene
815   transfers in the emerging pathogen *Dickeya solani*. BMC Genomics. 16:788.
816   DOI:10.1186/s12864-015-1997-z
817

818 29. Li, X., Ma, Y., Liang, S., Tian, Y., Yin, S., Xie, S., et al. 2018. Comparative genomics of 84
819   *Pectobacterium* genomes reveals the variations related to a pathogenic lifestyle. BMC
820   Genomics. 19 DOI: 10.1186/s12864-018-5269-6
821

822 30. Ma, B., Hibbing, M. E., Kim, H.-S., Reedy, R. M., Yedidia, I., Breuer, J., et al. 2007. Host
823   range and molecular phylogenies of the soft rot Enterobacterial genera
824   *Pectobacterium* and *Dickeya*. Phytopathology. 97:1150–1163. DOI: 10.1094/PHYTO-
825   97-9-1150
826

827 31. Ma, X., Brazil, J., Rivedal, H., Perry, K. L., Frost, K., and Swingle, B. 2022. First Report
828   of *Pectobacterium versatile* causing potato soft rot in Oregon and Washington. Plant
829   Dis. 106:1292. DOI: 10.1094/PDIS-08-21-1635-PDN
830

831 32. Mansfield, J., Genin, S., Magori, S., Citovsky, V., Sriariyanum, M., Ronald, P., et al.
832   2012. Top 10 plant pathogenic bacteria in molecular plant pathology. Mol. Plant
833   Pathol. 13:614–629. DOI: 10.1111/j.1364-3703.2012.00804.x
834

33. Mattinen, L., Nissinen, R., Riipi, T., Kalkkinen, N., and Pirhonen, M. 2007. Host-extract

induced changes in the secretome of the plant pathogenic bacterium
*Pectobacterium atrosepticum* . PROTEOMICS. 7:3527–3537. DOI:
10.1002/pmic.200600759

34. McGowan, S. J., Sebaihia, M., O'Leary, S., Hardie, K. R., Williams, P., Stewart, G. S. A.
B., et al. 1997. Analysis of the carbapenem gene cluster of *Erwinia carotovora* :
definition of the antibiotic biosynthetic genes and evidence for a novel β-lactam
resistance mechanism. Mol. Microbiol. 26:545–556. DOI:10.1046/j.1365-
2958.1997.6001974.x

35. Motyka-Pomagruk, A., Zoledowska, S., Sledz, W., and Lojkowska, E. 2021. The
occurrence of bacteria from different species of *Pectobacteriaceae* on seed potato
plantations in Poland. Eur. J. Plant Pathol. 159:309–325. DOI:10.1007/s10658-020-
02163-x

36. Nguyen, A. H., Tomita, T., Hirota, M., Sato, T., and Kamio, Y. 1999. A simple
purification method and morphology and component analyses for carotovoricin Er, a
Phage-tail-like bacteriocin from the plant pathogen *Erwinia carotovora* Er. Biosci.
Biotechnol. Biochem. 63:1360–1369. DOI: 10.1271/bbb.63.1360

37. Pasanen, M., Waleron, M., Schott, T., Cleenwerck, I., Misztak, A., Waleron, K., et al.
2020. *Pectobacterium parvum* sp. nov., having a *Salmonella* SPI-1-like Type III secre-
tion system and low virulence. Int. J. Syst. Evol. Microbiol. 70:2440–2448. DOI:
38. 10.1099/ijsem.0.004057

39. Pédron, J., Bertrand, C., Taghouti, G., Portier, P., and Barny, M.-A. 2019.
*Pectobacterium aquaticum* sp. nov., isolated from waterways. Int. J. Syst. Evol.
Microbiol. 69:745–751. DOI: 10.1099/ijsem.0.003229

40. Portier, P., Pédron, J., Taghouti, G., Dutrieux, C., and Barny, M.-A. 2020. Updated Tax-
onomy of *Pectobacterium* Genus in the CIRM-CFBP Bacterial Collection: When Newly
Described Species Reveal "Old" Endemic Population. Microorganisms. 8:1441. DOI:
41. 10.3390/microorganisms8091441

42. Portier, P., Pédron, J., Taghouti, G., Fischer-Le Saux, M., Caullireau, E., Bertrand, C., et
al. 2019. Elevation of *Pectobacterium carotovorum* subsp. *odoriferum* to species level
as *Pectobacterium odoriferum* sp. nov., proposal of *Pectobacterium brasiliense* sp.
nov. and *Pectobacterium actinidiae* sp. nov., emended description of *Pectobacterium*
*carotovorum* and description of *Pectobacterium versatile* sp. nov., isolated from
streams and symptoms on diverse plants. Int. J. Syst. Evol. Microbiol. 69:3207–3216.
DOI: 10.1099/ijsem.0.003611

43. Roh, E., Park, T.-H., Kim, M., Lee, S., Ryu, S., Oh, C.-S., et al. 2010. Characterization of
a new bacteriocin, Carocin D, from *Pectobacterium carotovorum* subsp. *carotovorum*
Pcc21. Appl. Environ. Microbiol. 76:7541–7549. DOI: 10.1128/aem.03103-09

44. Royer, G., Dixit, Z., Pédron, J., Pierrat, G., Demontant, V., Berçot, B., et al. 2022.
Genetic and Phenotypic Study of the *Pectobacterium versatile* Beta-Lactamase, the

Enzyme Most Similar to the Plasmid-Encoded TEM-1 ed. Christopher A. Elkins. Appl.
Environ. Microbiol. 88:e00220-22. DOI: 10.1128/aem.00220-22
886

887 45. Shyntum, D. Y., Nkomo, N. P., Shingange, N. L., Gricia, A. R., Bellieny-Rabelo, D., and
888 Moleleki, L. N. 2019. The impact of Type VI secretion system, bacteriocins and
889 antibiotics on bacterial competition of *Pectobacterium carotovorum* subsp.
890 *brasiliense* and the regulation of carbapenem biosynthesis by iron and the ferric-
891 uptake regulator. Front. Microbiol. 10 DOI: 10.3389/fmicb.2019.02379
892

893 46. Smoktunowicz, M., Jonca, J., Stachowska, A., May, M., Waleron, M. M., Waleron, M.,
894 et al. 2022. The international trade of ware vegetables and ornamental plants—an
895 underestimated risk of accelerated spreading of phytopathogenic bacteria in the era
896 of globalisation and ongoing climatic changes. Pathogens. 11:728. DOI:10.3390/pa-
897 thogens11070728
898

899 47. Taillefumier, T., Posfai, A., Meir, Y., and Wingreen, N. S. 2017. Microbial consortia at
900 steady supply. eLife. 6:e22644. DOI: 10.7554/eLife.22644
901

902 48. Toth, I. K., Barny, M., Brurberg, M. B., Condemine, G., Czajkowski, R., Elphinstone, J.
903 G., et al. 2021. *Pectobacterium* and *Dickeya*: environment to disease development. In
904 *Plant Diseases Caused by Dickeya and Pectobacterium Species*, eds. Frédérique Van
905 Gijsegem, Jan M. van der Wolf, and Ian K. Toth. Cham: Springer International
906 Publishing, p. 39–84. http://link.springer.com/10.1007/978-3-030-61459-1_3
907

908 49. Weiss, A. S., Burrichter, A. G., Durai Raj, A. C., Von Strempel, A., Meng, C., Kleigrewe,
909 K., et al. 2022. In vitro interaction network of a synthetic gut bacterial community.
910 ISME J. 16:1095–1109. DOI: 10.1038/s41396-021-01153-z
911

912 50. de Werra, P., Debonneville, C., Kellenberger, I., and Dupuis, B. 2021. Pathogenicity
913 and Relative Abundance of *Dickeya* and *Pectobacterium* species in Switzerland: an
914 epidemiological dichotomy. Microorganisms. 9:2270. DOI:10.3390/microorga-
915 nisms9112270
916

51. van der Wolf, J. M., Acuña, I., De Boer, S. H., Brurberg, M. B., Cahill, G., Charkowski,
A. O., et al. 2021. Diseases Caused by *Pectobacterium* and *Dickeya* Species Around
the World. In *Plant Diseases Caused by Dickeya and Pectobacterium Species*, eds.
Frédérique Van Gijsegem, Jan M. van der Wolf, and Ian K. Toth. Cham: Springer
International Publishing, p. 215–261. http://link.springer.com/10.1007/978-3-030-
61459-1_7
923