1 **NEUROPILIN-1 IS A CO-RECEPTOR FOR NGF AND TRKA-EVOKED PAIN**
2

3 Chloe J. Peach[1,2], Raquel Tonello[1,2], Kimberly Gomez[1,2], Aida Calderon-Rivera[1,2], Marina Rodríguez
4 Sánchez[1,2], Laura Maile[3], Samantha Perez-Miller[1], Ana-Maria Manu[1,2], Hyunggu Hahn[1,2], Alex R.B.
5 Thomsen[1,2], Brian L. Schmidt[1,2,4,5], Steve Davidson[3], May Khanna[1,2], Rajesh Khanna[1,2,6],
6 *Nigel W. Bunnett[1,2,6]
7
8 [1]Department of Molecular Pathobiology, College of Dentistry, New York University; [2]Pain Research Center,
9 New York University; [3]Department of Anesthesiology, College of Medicine, University of Cincinnati,
10 Cincinnati, USA, [4]Translational Research Center, College of Dentistry, New York University; [5]Department
11 of Oral and Maxillofacial Surgery, College of Dentistry, New York University; [6]Department of Neuroscience
12 and Physiology, Neuroscience Institute, Grossman School of Medicine, New York University
13
14 **Keywords**: NGF, NRP1, GIPC1, receptor tyrosine kinase, co-receptor, nociception
15
*Correspondence: Nigel W. Bunnett, PhD. Room 724, 433 1st Avenue, New York, 10010, USA.
Nwb2@nyu.edu

ABSTRACT

Nerve growth factor (NGF) monoclonal antibodies are one of the few patient-validated non-opioid
treatments for chronic pain, despite failing to gain FDA approval. Herein, we demonstrate that neuropilin-1
(NRP1) is a co-receptor for NGF that potentiates tropomyosin-related kinase A (TrkA) signaling. NGF binds
NRP1 with nanomolar affinity. NRP1 and its associated adaptor protein GIPC1 are coexpressed with TrkA
in nociceptors, particularly in human tissues. NRP1 inhibitors prevent NGF-stimulated action potential firing
of human and mouse nociceptors and abrogate NGF-evoked and inflammatory nociception in mice. NRP1
knockdown blunts NGF-stimulated TrkA phosphorylation, kinase signaling and transcription, whereas
NRP1 overexpression enhances signaling. As well as interacting with NGF, NRP1 forms a heteromeric
complex that chaperones TrkA from the biosynthetic pathway to the plasma membrane and signaling
endosomes. NRP1 thereby enhances NGF-induced TrkA dimerization, endocytosis and signaling.
Knockdown of GIPC1, a PDZ-binding protein that scaffolds NRP1 and TrkA cargo to myosin VI, abrogates
NGF-evoked nociception. We identify NRP1 as a previously unrecognized co-receptor necessary for
NGF/TrkA pain signaling by direct NGF binding and chaperoning TrkA to the plasma membrane and
signaling endosomes *via* the adaptor protein GIPC1 - a discovery that offers a long-awaited alternative to
block NGF-evoked pain.

INTRODUCTION

The first discovered growth factor, nerve growth factor (NGF), was identified by its ability to
stimulate growth of sympathetic neurons[1]. Tropomyosin-related kinase A (TrkA), a receptor tyrosine kinase
(RTK), mediates the neurotrophic actions of NGF[2]. NGF/TrkA signalosomes are transported in a retrograde
direction from peripheral nerve terminals to the soma, where they regulate transcription and neuronal
development[3,4]. NGF and TrkA are also implicated in pain[5]. p75[NTR] is a co-receptor for NGF and the
immature pro-peptide[6,7] that activates opposing pro-apoptotic signaling pathways[8]. Although NGF and its
receptors have been intensively studied in the context of neuronal development and pain, mechanistic gaps
in understanding have hampered the successful approval of NGF-directed therapeutics.

Chronic pain afflicts twenty percent of the global population at some point in life, yet is inadequately
treated by opioids and non-steroidal anti-inflammatory drugs, which lack efficacy and have life-threatening
side effects. NGF/TrkA are one of the few non-opioid targets for chronic pain validated in patients. NGF,
which is produced by injured and diseased tissues, activates TrkA on peripheral nociceptors to evoke
hyperexcitability and the expression of neuropeptides and ion channels that mediate pain[5]. The central role
TrkA and NGF in pain is evident in patients with Hereditary Sensory and Autonomic Neuropathy (HSAN)
Type IV and V, where pathological insensitivity to pain results from loss-of-function mutations in TrkA[9] and
NGF[10], respectively. Although monoclonal antibodies (mAbs) are analgesic in osteoarthritic patients[11], they
lack FDA approval due to worsening joint damage in some individuals[12]. The identification of nociceptor-
enriched mediators of NGF-induced pain may facilitate development of non-opioid analgesics, avoiding the
adverse effects of systemic NGF sequestration with mAbs.

Transcriptome analyses have identified proteins that are conserved between rodent and human
nociceptors, including neuropilin-1 (NRP1)[13]. NRP1 is a type I transmembrane protein discovered for its
role in axon guidance[14]. Lacking a catalytic domain, NRP1 does not transduce signals *per se* but rather
acts as a co-receptor for unrelated families of proteins, including vascular endothelial growth factor (VEGF-
A)[15]. NRP1 is a co-receptor with VEGF receptor 2 (VEGFR2) that enhances VEGF-A-induced blood vessel
development. NRP1 overexpression in cancers spurred the development of a mAb,
vesencumab/MNRP1685A, to inhibit tumorigenesis[16]. NRP1 is also implicated in VEGF-A-mediated
pain[17,18]. Since NGF, a major pain mediator, possesses a prospective NRP1 binding motif, we hypothesized
that NRP1 is a co-receptor for NGF and TrkA in nociception.

RESULTS

NRP1 binds human β-NGF

NRP1 is a co-receptor for unrelated families of proteins including neuronal guidance molecules
(semaphorins)[19], growth factors (VEGF-A)[15], cell-penetrating peptides[20], and viruses (SARS-CoV-2)[14].
NRP1 ligands share a C-terminal basic motif (C-end rule, "CendR" motif, R/KXXR/K), which interacts with
extracellular "b" domains of NRP1[15,19]. Inspection of the amino acid sequence of mature NGF identified
prospective 'CendR' R/KxxR/K motifs within the C-terminus that are conserved between rodents and
humans and known to mediate the interaction of other growth factors with NRP1[20] (**Fig. 1A**). Molecular
modeling of βNGF and human NRP1 identified a potential interaction between the CendR motif at the C-
terminus of NGF and the b1b2 domain of NRP1. To examine the potential for productive interactions
between NGF and NRP1, we used restrained protein-protein docking of a human NGF structure terminating
in Arg118 (PDB 4GZ9[21]; **Fig. 1B**) to the AlphaFold-predicted structure of soluble human a1a2b1b2 domains
of NRP1. Our modeling supports the hypothesis that these proteins could interact *via* the canonical CendR
binding mechanism, similar to the interactions observed for structures of VEGF$_{165}$a CendR bound to the b1
domain[22].

To directly investigate whether NGF binds purified NRP1, microscale thermophoresis (MST) was
used to analyze interactions between human NGF and His-tagged human NRP1. NGF interacted with
extracellular NRP1 a1a2b1b2 domains with nanomolar affinity (K_d = 65.64 ± 6.66 nM, n=4, **Fig. 1C**). This
is comparable to published affinities between purified VEGF$_{165}$a and NRP1 (K_d 9 nM[23] to 120 nM[24]).

NGF binding to NRP1 in living cells at 37ºC was analyzed using a modified bioluminescence
resonance energy transfer (BRET)-based assay (**Fig. 1D**). VEGF$_{165}$a (positive control growth factor known
to interact with NRP1) or NGF was genetically fused to an 11 amino acid fragment (HiBiT) of nanoluciferase

(NanoLuc). HiBiT has a high affinity for the complementary LgBiT fragment. HiBiT-tagged growth factors
were expressed in HEK293T cells, secreted into supernatant, and conjugated to recombinant LgBiT to form
full length and catalytically active NanoLuc. Upon addition of furimazine substrate, complemented NanoLuc
emits bioluminescence. This supernatant containing HiBiT-tagged growth factor was incubated with cells
expressing SnapTag-NRP1 labeled with SNAP-Surface® Alexa Fluor® 488 fluorophore (SnapTag-AF488,
+) or lacking fluorophore (-). If proteins are in close proximity (<10 nm), luminescent growth factor can act
as a BRET donor to NRP1. As a positive control, BRET was detected between luminescent $VEGF_{165}a$ and
SnapTag-NRP1 (**Fig. 1E**). Pre-incubation of cells with an excess of $VEGF_{165}a$ (10 nM) abolished the BRET
signal. No BRET signal was detected when luminescent $VEGF_{165}a$ was incubated with cells expressing a
binding-dead NRP1 mutant in the b1 domain[25,26]. Importantly, BRET was detected between NGF-HiBiT and
SnapTag-NRP1, and pre-incubation with unlabeled $VEGF_{165}a$ inhibited this response.

The results from molecular modeling, analysis of interactions between recombinant proteins, and
ligand binding studies in intact cells provide evidence that NGF directly binds NRP1 with nanomolar affinity.

***Ntrk1*/TrkA and *Nrp1*/NRP1 are co-expressed in mouse and human DRG neurons**

To determine whether NRP1 could function as an NGF/TrkA co-receptor in nociceptors, TrkA and
NRP1 were localized in mouse and human dorsal root ganglia (DRG) by immunofluorescence (protein) and
RNAScope® *in situ* hybridization (mRNA). In mice, immunoreactive TrkA was detected in vesicles and
immunoreactive NRP1 was localized at the plasma membrane of the same DRG neurons (**Fig. 2A**). *Ntrk1*
(TrkA) and *Nrp1* (NRP1) mRNAs colocalized in both mouse and human DRG neurons, which were identified
by NeuN immunostaining (**Fig. 2B,C**). In mice, *Ntrk1* mRNA was expressed in 27% and *Nrp1* mRNA was
expressed in 30% of the neurons; *Nrp1* was co-expressed in approximately one third of *Ntrk1*-positive
neurons (**Fig. 2D**). In humans, *Nrp1* was co-expressed in approximately 93% of *Ntrk1*-positive neurons
(**Fig. 2E**). Thus, TrkA and NRP1 are co-expressed in the same neurons in DRG in mouse and human,
where NRP1 is appropriately located to control NGF/TrkA signaling.

NRP1 antagonism abrogates NGF-induced ionic currents in mouse and human DRG neurons

NGF/TrkA signaling promotes pain by kinase-mediated phosphorylation and sensitization of ion
channels, leading to enhanced sensitivity to mechanical, thermal and chemical stimuli, thus activating
voltage-gated ion channels in DRG neurons[5]. Patch-clamp recordings under current-clamp mode were
made to determine whether NRP1 is necessary for NGF-induced excitability and ion channel activity in
dissociated human DRG neurons. Human NGF (50 nM) increased the number of action potentials elicited
by a depolarizing ramp pulse from 0 to 250 pA in 1 s (**Fig. 3A,B**). Pre-incubation with the NRP1 inhibitor
EG00229 (30 µM)[25] prevented NGF-induced hyperexcitability (**Fig. 3A,B**). NGF or EG00229 did not affect
the resting membrane potential (**Fig. 3C**) or the rheobase, the minimum current necessary to elicit an action
potential (**Fig. 3D**). Murine NGF similarly stimulated action potential firing in mouse DRG nociceptors (Fig.
3E-G). Pre-incubation with EG00229 blocked NGF-induced excitability (**Fig. 3E,F**), while NGF has no effect
on resting membrane potential or rheobase (**Fig. 3G,H**).

Since both Ca^{2+} and Na^+ channels have critical roles in controlling the excitability of sensory
neurons, we further tested whether NGF increased the activity of voltage-gated ion channels by making
patch-clamp recordings under the voltage-clamp mode in dissociated mouse DRG. NGF (50 nM) increased
total Ca^{2+} currents (**Fig. 3I**) and current density (**Fig. 3J,K**) by ~50% when compared to vehicle (DMSO)-
treated cells. EG00229 prevented NGF-induced activation of Ca^{2+} currents, but had no effect on Ca^{2+}
currents in unstimulated DRGs (**Fig. 3J,K**). Half-maximal activation and inactivation ($V_{1/2}$) potential, as well
as slope factor values (*k*) for activation and inactivation, were not different between the conditions tested,
except for an ~8-mV hyperpolarizing shift in Ca^{2+} channel activation induced by EG00229 when compared
to DMSO- and NGF-treated DRGs (**Fig. 3L, Supplementary Table 1**). Similarly, EG00229 normalized
NGF-induced increases in Na^+ currents (**Fig. 3M**). NGF caused a two-fold increase in Na^+ current density
(**Fig. 3N**) and peak current density (**Fig. 3O**) when compared to vehicle (DMSO)-treated DRG neurons.
EG00229 abolished these effects of NGF but had no effects on Na^+ currents in unstimulated DRGs (**Fig.
3N,O**). There were no detectable changes in the voltage-dependence of activation and inactivation between
the conditions tested, except for a ~10-mV depolarizing shift in the $V_{1/2}$ of inactivation of EG00229 and NGF-
treated cells compared to control (**Fig. 3P**). These data implicate voltage-gated Ca^{2+} and Na^+ channels as

downstream effectors of NRP1-mediated NGF signaling. Thus, NRP1 is necessary for NGF-induced
sensitization of nociceptors in rodents to humans.

NRP1 antagonism inhibits NGF-induced nociception in mice

NGF injection is known to induce pain in humans and rodents[5]. In addition to sensitizing ion
channels, NGF/TrkA signaling also promotes chronic pain through the upregulation of genes that mediate
pain (*e.g.*, substance P, transient receptor potential vanilloid 1)[5]. To test whether NRP1 is necessary for
NGF-induced nociception, murine NGF (50 ng) was administered to mice by intraplantar (i.pl., 10 µl)
injection (**Fig. 4A**). A human mAb against the b1b2 domain of NRP1 (vesencumab[16], 7 µg/10 µl) or control
IgG (7 µg/10 µl), or small molecule NRP1 inhibitor EG00229 (30 µM/10 µl)[25] or compound 5 (Cpd-5; 30
µM/10 µl)[27] or vehicle (PBS, 10 µl) was co-administered (i.pl.) with NGF. Withdrawal responses of NGF-
injected (ipsilateral) hindpaws to stimulation with von Frey filaments (VFF) and radiant heat were assessed
to evaluate mechanical allodynia and thermal hyperalgesia, respectively. In control experiments, NGF
decreased the withdrawal threshold to VFF stimulation and the latency of withdrawal to thermal stimulation
within 30 min for at least 4 h, indicating mechanical allodynia and thermal hyperalgesia (**Fig. 4B-G**).
Vesencumab and EG00229 strongly inhibited NGF-induced mechanical allodynia for 2 h and Cpd-5
inhibited mechanical allodynia for 1 h (**Fig. 4B-C**). Vesencumab, EG00229 and Cpd-5 also inhibited NGF-
induced thermal hyperalgesia for 1-2 h (**Fig. 4E-G).** None of the inhibitors affected the withdrawal responses
of the ipsilateral paw to mechanical or thermal stimuli in mice that did not receive NGF (**Fig. 4B-G**).

NGF mediates inflammatory pain[5], providing an opportunity to determine whether NRP1 is
necessary for the pronociceptive actions of endogenous NGF. Complete Freund's Adjuvant (CFA, 1 mg/ml,
10 µl, i.pl.) was injected into the hindpaw. Vesencumab mAb (7 µg/10 µl) or IgG Ctrl (7 µg/10 µl), or
EG00229 (30 µM/10 µl) or PBS (10 µl) (i.pl.) was injected into the inflamed paw 48 h after CFA (**Fig. 4H**).
CFA induced sustained mechanical allodynia and thermal hyperalgesia (**Fig. 4I-L**). Vesencumab, but not
EG00229, reduced thermal hyperalgesia 1 h after treatment (**Fig. 4K,L**). Vesencumab inhibited mechanical
allodynia compared to IgG control, with a maximal effect at 1 h, lasting up to 2 h post-injection (**Fig. 4I**).
EG00229 also partially reversed mechanical allodynia compared with vehicle, with a maximum inhibitory
effect at 1 h, lasting for up to 4 h (**Fig. 4J**). Thus, NRP1 is necessary for the pronociceptive actions of
exogenous and endogenous NGF in mice.

NRP1 controls NGF and TrkA kinase signaling

NGF primarily signals through TrkA[28], a 'crab claw'-like RTK where NGF binds to extracellular
immunoglobulin-like domains to induce conformational changes throughout the RTK dimer that initiate auto-
and trans-phosphorylation of intracellular tyrosine residues (*e.g.,* Y490, Y785). To determine whether NRP1
controls NGF-stimulated phosphorylation of TrkA, a proximal and necessary component of TrkA signaling,
dissociated mouse DRG neurons were stained with a phospho-specific antibody to TrkA phosphorylation
at residue Y785 (equivalent to Y794 in mouse TrkA). NGF (100 nM, 15 min) stimulated a ~1.4-fold increase
in the intensity TrkA phospho-Y785 immunofluorescence compared to vehicle (**Fig. 5A,B**). Pre-incubation
with EG00229 (30 µM, 30 min) prevented NGF-stimulated phosphorylation of TrkA but had no effect on the
basal unstimulated phosphorylation. These results support the hypothesis that NRP1 is necessary for NGF-
induced activation of TrkA.

Activated TrkA stimulates extracellular signal-regulated kinase (ERK) phosphorylation, as well as
AKT and phospholipase Cγ. In addition to triggering neuronal development, TrkA-induced ERK signaling
contributes to NGF-induced sensitization of nociceptors and pain[5]. The contribution of NRP1 to ERK
activation was monitored in CAD.a cells, a neuron-like cell line modified from the Cath.a catecholaminergic
cell line obtained from a murine tumor[29]. While CAD.a cells lack TrkA expression, they contain p75[NTR] and
NRP1 (**Supplementary Fig. 1**). To probe NGF/TrkA ERK signaling with high spatial and temporal
resolution, Förster resonance energy transfer (FRET) EKAR biosensors targeted to the cytosol or nucleus
were coexpressed with human TrkA in CAD.a cells. EKAR biosensors contain a reversible substrate
sequence separated by two fluorophores (**Fig. 5C**). NGF activated ERK in the cytosol and nucleus within 5
min for at least 20 min (**Fig. 5D**). NGF-induced ERK activation was concentration-dependent, with a higher
potency for nuclear than cytosolic ERK (**Fig. 5E,F, Supplementary Table 2**). Pre-incubation with EG00229
(30 µM) did not affect NGF activation of cytosolic ERK (**Fig. 5E**), but significantly reduced the potency for

NGF activation of nuclear ERK (**Fig. 5F, Supplementary Table 2**; NGF EC_{50}: DMSO control, ~8 pM;
EG00229 ~98 pM; paired t-test, *P*=0.023).

The contribution of NRP1 to NGF-induced ERK signaling was also investigated by NRP1
overexpression with TrkA in HEK293T cells, which express low levels of these proteins. As in CAD.a cells,
NGF activated ERK in the cytosol and nucleus of HEK293T cells (**Fig. 4G**). While NRP1 co-expression did
not influence cytosolic ERK signaling (**Fig. 5H**), NRP1 expression significantly enhanced activation of
nuclear ERK in response to low (1 pM) NGF concentrations (**Fig. 4I**). The outcomes of nuclear ERK
signaling were studied by expression of a transcriptional luciferase reporter, where a luminescent protein
is produced downstream of an ERK promoter (**Fig. 4C**). NGF stimulated concentration-dependent ERK
transcriptional activity with significantly higher potency in cells overexpressing TrkA and NRP1 compared
to cells expressing TrkA alone (**Fig. 5J, Supplementary Table 2**; NGF EC_{50}: TrkA alone, 214 pM; TrkA +
NRP1, 71 pM; paired t-test, *P*=0.0004). NGF did not stimulate ERK transcriptional activity in cells
expressing NRP1 alone, confirming lack of inherent signaling capability of NRP1 (**Fig. 5J**). These results
support the notion that NRP1 enhances NGF-induced TrkA activation and signaling by pathways that
underpin nociception. We next sought to determine the mechanism of NRP1-mediated potentiation of
NGF/TrkA signaling.

NRP1 is a chaperone that forms a complex with TrkA to control trafficking from the biosynthetic
pathway to the plasma membrane

In addition to directly binding NGF, NRP1 could amplify NGF/TrkA signaling by receptor/co-receptor
interactions with TrkA, akin to its co-receptor function with VEGFR2[30]. Cell surface expression of human
TrkA and NRP1 was measured by specific substrate-based labeling of enzymatic tags genetically fused to
the extracellular N-terminus of either receptor. SnapTag-TrkA and HaloTag-NRP1 were expressed in
HEK293T and CAD.a cells and covalently labeled with membrane-impermeant AlexaFluor® substrates,
thereby selectively labeling receptors transported to the cell surface. SnapTag-TrkA and HaloTag-NRP1
were highly co-localized at the cell surface of HEK293T and CAD.a cells, with the latter also showing a high
level of co-localization in subcellular compartments (**Fig. 6A**). Labeling specificity was confirmed in cells
expressing TrkA or NRP1 alone (**Supplementary Fig. 2**).

To investigate the formation of a heteromeric complex between TrkA and NRP1, BRET was
measured in HEK293T cells expressing NanoLuc-NRP1 and SnapTag-TrkA. Extracellular N-terminal
NanoLuc acts as an energy donor to excite a nearby SnapTag. Measurement of BRET between NanoLuc-
NRP1 and increasing levels of SnapTag-TrkA revealed a hyperbolic relationship, indicative of assembly of
a heteromeric complex between NanoLuc-NRP1 and SnapTag-TrkA (**Fig. 6B,C**). As a positive control, a
hyperbolic BRET signal relationship was also detected between NanoLuc-p75[NTR], which is known to interact
with TrkA[5], and increasing levels of SnapTag-TrkA. In contrast, there was a linear BRET signal between
NanoLuc-NRP1 and increasing levels of SnapTag-calcitonin-like receptor (CALCRL), an unrelated
transmembrane receptor. To verify whether this relationship was observed with protein expression, data
from a representative experiment quantifying cell surface SnapTag labeling also demonstrated a hyperbolic
curve between TrkA and NRP1 (**Fig. 6C**). These results suggest that NRP1 and TrkA co-localize at the
plasma membrane as a heteromeric complex.

Recruiting TrkA to the plasma membrane would enable increased access to extracellular growth
factor. The effect of NRP1 on the cell surface expression of TrkA was determined using the membrane-
impermeant SnapTag fluorophore to selectively label cell surface TrkA (**Fig. 6D**). NRP1 co-expression
increased level of TrkA at the surface in HEK293T cells by 150 ± 43% and of CAD.a cells by 109 ± 4% (**Fig.**
6E-G). Enhanced bystander BRET, which capitalizes on the endogenous affinity of pairs of *Renilla*-tagged
proteins to boost sensitivity[31], was developed to quantify the effects of NRP1 expression on the localization
of TrkA in different subcellular compartments of living cells. TrkA tagged on the C-terminus with Renilla
luciferase (Rluc, BRET) was coexpressed in HEK293T or CAD.a cells with proteins resident of the plasma
membrane (CAAX), early endosomes (Rab5a), recycling endosomes (Rab4a) and the cis-Golgi apparatus
(Giantin) tagged with *Renilla* green fluorescent protein (RGFP) (**Fig. 6H**). In HEK293T cells, NRP1
expression significantly increased BRET between TrkA-Rluc8 and RGFP-CAAX and significantly
decreased BRET between TrkA-Rluc8 and tdRGFP-Rab5a, tdRGFP-Rab4a and tdRGFP-Giantin (**Fig. 6I**).
In CAD.a cells, NRP1 expression did not affect BRET between TrkA-Rluc8 and RGFP-CAAX but
significantly decreased BRET between TrkA-Rluc8 and tdRGFP-Rab5a, tdRGFP-Rab4a and tdRGFP-

Giantin (**Fig. 6J**). These results indicate that NRP1 expression causes a redistribution of TrkA from
subcellular regions involved in receptor recycling or *de novo* export to the plasma membrane, consistent
with a chaperone function. Control studies confirmed NRP1 was successfully expressed using a HaloTag
label, where NRP1 co-expression had no significant effect on overall TrkA expression (**Supplementary**
Fig. 3A-D). These results suggest that NRP1 acts as a chaperone for TrkA.

NRP1 enhances NGF-induced TrkA trafficking and dimerization

Upon NGF stimulation, TrkA traffics to endosomes by clathrin-dependent and -independent
mechanisms, a process known to further enable TrkA signaling through NGF/TrkA signalosomes that are
retrogradely transported in endosomes or multivesicular bodies to the soma of sympathetic neurons
mediate the neurotrophic actions of NGF[3,4]. To determine the effects of NRP1 on NGF-evoked endocytosis
of TrkA, BRET was measured between TrkA-Rluc8 and plasma membrane marker RGFP-CAAX (**Fig. 7A**).
NGF caused a concentration-dependent decrease in TrkA-Rluc8 and RGFP-CAAX BRET (EC$_{50}$ ~870 pM),
with 10-100 nM NGF inducing maximal decrease within 5 min that was sustained for 20 min, consistent
with TrkA endocytosis (**Fig. 7B, Table 2**). Hypertonic (0.45 M) sucrose or the clathrin inhibitor pitstop 2 (30
μM) prevented NGF-induced endocytosis of TrkA (**Fig. 7C**). Co-expression of NRP1 with TrkA enhanced
NGF-stimulated endocytosis of TrkA (**Fig. 7D**). This effect of NRP1 was observed at higher NGF
concentrations (>10 nM), with minimal effect on the potency of NGF-induced endocytosis of TrkA (**Table**
2). NGF also stimulated concentration-dependent removal of TrkA from the plasma membrane of CAD.a
cells (EC$_{50}$ ~490 nM; pEC$_{50}$ = 9.31 ± 0.18, n=4) that was inhibited by hypertonic sucrose and pitstop 2 (**Fig.**
7E-G). Knockdown of endogenous NRP1 with siRNA significantly inhibited NGF-induced endocytosis of
TrkA in CAD.a cells (**Fig. 7G**). NRP1 knockdown was confirmed at a protein level using
immunofluorescence in CAD.a cells, while there was no effect on TrkA expression (**Supplementary Fig.**
3E-F). Thus, whereas NRP1 overexpression enhances agonist-stimulated endocytosis of TrkA in HEK293T
cells, NRP1 knockdown has the opposite effect in CAD.a cells. The results are consistent with a chaperone
role for NRP1 in NGF-stimulated endocytosis of TrkA.

Dimerization of TrkA is pivotal for NGF/TrkA signaling[5]. The contribution of NRP1 to TrkA
oligomerization was evaluated by measuring BRET between NanoLuc-TrkA and SnapTag-TrkA in
HEK293T cells (**Fig. 7H**). Expression of increasing amounts of SnapTag-TrkA in HEK293T cells expressing
a fixed amount of NanoLuc-TrkA produced a hyperbolic BRET signal, indicating oligomerization (**Fig. 7I**).
NGF (30 nM) significantly enhanced this signal, consistent with agonist-evoked TrkA oligomerization. In
cells expressing a fixed ratio of SnapTag-TrkA and NanoLuc-TrkA, NGF stimulated a concentration-
dependent increase in BRET that was maximal after 5 min and sustained for at least 20 min (EC$_{50}$ 15 nM;
Fig. 7J, Table 2). While NGF had a lower potency with respect to dimerization than endocytosis, the kinetics
of TrkA dimerization and endocytosis were similar. NRP1 overexpression enhanced TrkA oligomerization
in response to higher NGF concentrations (>30 nM) (**Fig. 7K**). As dimerization is the first step in RTK
activation, with well-established evidence for TrkA signaling from endosomes, these results provide a
mechanistic insight into the role of NRP1 in controlling TrkA signaling and trafficking.

GIPC1 mediates NRP1/TrkA interactions and NGF-induced pain signaling

While NRP1 lacks intrinsic catalytic activity, the short cytoplasmic C-terminus interacts with G Alpha
Interacting Protein (GAIP) Interacting Protein C-terminus 1 (GIPC1), or synectin, through a PDZ domain[32].
GIPC1 also interacts with the membrane-proximal regions of TrkA[33]. Thus, interaction with GIPC1 could
underpin the co-receptor function of NRP1 in NGF/TrkA-evoked nociception. Analysis by RNAScope® *in*
situ hybridization revealed that *Gipc1* mRNA is expressed by a large proportion of neurons in DRG from
mouse (**Fig. 8A**). In humans, *Gipc1* was expressed by ~85-100% of neurons expressing *Ntrk1* mRNA (**Fig.**
8B,C).

GIPC1 is an intracellular adaptor protein that associates with receptors and channels to regulate
their trafficking by interacting with the inwardly-directed myosin VI motor[34]. To determine whether GIPC1 is
necessary for the TrkA chaperone function of NRP1, BRET between TrkA-Rluc8 and RGFP-CAAX was
measured in HEK293T cells after GIPC1 knockdown. GIPC1 siRNA inhibited NRP1-induced plasma
membrane expression of TrkA (**Fig. 8D**). In CAD.a cells, preincubation with a GIPC1 inhibitor (300 μM
CR1023[35]) or a myosin VI inhibitor (50 μM TIP[36]) reduced NGF-induced TrkA trafficking to the plasma
membrane (**Fig. 8E,F**). Similarly, GIPC1 siRNA inhibited the maximal response of NGF-induced ERK

signaling quantified using the downstream transcriptional reporter in CAD.a cells, with no effect on potency
(**Fig. 8G**) (Ctrl siRNA pEC_{50} = 9.58 ± 0.03, GIPC1 siRNA pEC_{50} = 9.63 ± 0.09, n=5). GIPC1 siRNA also
prevented NGF-induced action potential firing in mouse DRG nociceptors, determined by patch-clamp
recordings (**Fig. 8H,I**). GIPC1 siRNA knockdown was confirmed at the mRNA level in HEK293T cells,
CAD.a cells and DRG (**Supplementary Fig. 4A-C**).

To evaluate the role of GIPC1 in NGF-induced nociceptive behavior, GIPC1 or control siRNA was
administered to mice by intrathecal (i.t.) injection 48 h before NGF intraplantar injection. In mice treated
with control siRNA, NGF (50 ng/10 µl, i.pl.) caused mechanical allodynia and thermal hyperalgesia in the
ipsilateral paw within 30 min for at least 24 h (**Fig. 8J,K**). GIPC1 siRNA prevented NGF-evoked mechanical
allodynia and thermal hyperalgesia (**Fig 8J,K**). NGF or siRNA administration did not affect withdrawal
responses of the contralateral paw to mechanical stimuli (**Fig. 8L**). The nociceptive role of GIPC1 was also
investigated in a preclinical model of inflammatory pain in mice. Control or GIPC1 siRNA was administered
(i.t.) 48 h after CFA (i.pl.). Knockdown of GIPC1 in DRG was confirmed using RNAScope® *in situ*
hybridization (**Supplementary Fig. 4D**). While no effect was observed in the contralateral paw (**Fig. 8O**),
GIPC1 siRNA inhibited CFA-induced mechanical allodynia and thermal hyperalgesia after 24 and 48 h (**Fig.
8M,N**). These results reveal a major role for GIPC1 in NGF-evoked pain.

DISCUSSION

NRP1 is a previously unrecognized co-receptor for NGF and TrkA that is necessary for NGF-
induced nociception, demonstrated in both rodent and human tissue. NRP1 enhances NGF/TrkA signaling,
since NRP1 inhibition abrogates the actions of NGF in intact mice, isolated nociceptors and cell lines,
whereas NRP1 overexpression enhances TrkA signaling and trafficking. The results are consistent with
evidence that NRP1 is a co-receptor for VEGF-mediated angiogenesis using pharmacological[24] or genetic[37]
interventions. NRP1 is proposed to promote NGF/TrkA-mediated pain by at least two mechanisms: as a
coreceptor that directly binds NGF, and as a chaperone that assembles a heterodimeric complex to
enhance TrkA trafficking to and from the plasma membrane (**Fig. 9**).

The coreceptor function of NRP1 is supported by nanomolar affinity binding measurements
between recombinant NRP1 and NGF. Molecular modeling indicates that the "CendR" motif of NGF remains
exposed when interacting with TrkA[28], which could allow assembly of a trimeric NGF/TrkA/NRP1 complex.
Despite the glycosylated nature of NRP1, indirect association of NGF with matrix components is unlikely to
mediate NGF/NRP1 interactions as NGF does not directly interact with the extracellular matrix[38]. NRP1
potentiation of VEGFR2 signaling is linked to growth factor binding kinetics[26], similar to p75[NTR] potentiation
of NGF/TrkA binding kinetics[39]. Further studies of the effects of NRP1 on NGF/TrkA binding kinetics are
warranted. Since the NGF/p75[NTR] binding site[40] does not directly conflict with the proposed NGF/NRP1
binding site, a TrkA/p75[NTR]/NRP1 complex could interact with NGF (*e.g.,* 2:1:1 stoichiometry), although
there could be steric clashes between NRP1 and p75[NTR] for the same binding site on NGF.

The chaperone function of NRP1 is supported by BRET proximity assays showing that NRP1 and
TrkA form a heteromeric complex and that NRP1 re-routes TrkA from the biosynthetic pathway to the
plasma membrane and subsequently signaling endosomes. NRP1 colocalizes with TrkA to a greater extent
than with VEGFR2[30]. Further studies are required to identify potential NRP1/TrkA interaction sites. TrkA
redistribution to the cell surface would amplify exposure to extracellular NGF. Accordingly, NRP1
potentiates the effects of NGF on TrkA dimerization and endocytosis.

GIPC1 is a previously unrecognized mediator of pain. In the context of NGF/TrkA-evoked pain,
GIPC1 interacts with TrkA[33] and NRP1[32], and could thus scaffold TrkA/NRP1 interactions to facilitate NGF
signaling. GIPC1 disruption abrogates NRP1-stimulated translocation of TrkA to the plasma membrane and
inhibits NGF-evoked endocytosis of TrkA and ERK signaling, in line with reports in other systems[41].
Considering GIPC1 regulates the trafficking of many receptors and ion channels, other mechanisms could
mediate the antinociceptive effects of GIPC1 knockdown. Crucially, NRP1 and GIPC1 prominently
colocalize with TrkA in human nociceptors where they could mediate NGF-induced pain. NRP1 may also
serve as a co-receptor for NGF/TrkA signaling from adjacent cells, akin to VEGFR2 modulation from both
'cis' (same cell) and 'trans' (neighboring cell) forms[42]. This intercellular signaling may underlie functional
interactions between nociceptors and adjacent cells that contribute to pain (*e.g.,* Schwann cells).

Electrophysiological and behavioral studies support the essential role of NRP1 and GIPC1 in NGF-
induced pain. Two chemically distinct small molecule NRP1 inhibitors and a mAb against the b1 domain of
NRP1 prevent NGF-evoked mechanical allodynia and thermal hyperalgesia in mice, providing confidence
in specificity. GIPC1 siRNA similarly prevents NGF-induced nociception. NRP1 and GIPC1 inhibition blunts
inflammatory nociception, indicating that NRP1 and GIPC1 mediate the nociceptive action of endogenous
and exogenous NGF. NRP1 is necessary for NGF-induced action potential firing in both mouse and human
DRG neurons, supporting human translation. NRP1 is required for the proximal events of NGF-evoked
nociception (TrkA dimerization, phosphorylation, channel activation), as well as sustained components of
pain signaling (ERK activation, transcription), both of which mediate nociceptor sensitization[43]. Although
TrkA is the principal pronociceptive NGF receptor, p75[NTR] also contributes to NGF-induced pain[44] and the
effect of NRP1 on p75[NTR] warrants further study. NGF/TrkA signalosomes transported from nerve terminals
to the soma have been studied in compartmentalized sympathetic[45] and DRG[46] neuronal cultures, where
retrograde NGF/TrkA signaling controls gene expression during the development of sympathetic neurons.
Whether endosomal signaling of NGF/TrkA contributes to pain, as observed with G protein-coupled
receptors[47], deserves further attention. Additional studies are also required to determine whether NRP1 is
necessary for NGF-induced expression of pronociceptive transmitters and channels, and whether NRP1
mediates the neurotrophic actions of NGF. NRP1 interacts with semaphorins[19] as a co-receptor for plexins,
which regulate TrkA retrograde signaling[48]. Accordingly, NRP1 is also highly expressed in axonal growth
cones[49], as well as small diameter NGF-responsive DRG neurons during neurite sprouting[50]. Thus,
disruption of NRP1 could affect NGF-induced neuronal growth, with implications for the neuroplasticity
associated with chronic pain.

NGF is implicated in neuropathic, surgical and cancer pain[5], highlighting the need to study
contributions of NRP1 and GIPC1 across pain pathologies. Upregulation of NRP1 and GIPC1 in painful
conditions, such as cancer, could amplify NGF/TrkA signaling. Changes in NRP1 expression have been
reported in migraine patients[51]; though presumed to link to vasculature changes through VEGF signaling,
this could mediate migraine pain through NGF[52]. Given the key role of NGF/TrkA in pain and the challenge
of developing TrkA-selective antagonists, NGF mAbs were prioritized for the treatment of pain. While NGF
mAbs provided beneficial pain relief to patients with arthritis, rapidly progressing osteoarthritis observed in
some patients has precluded FDA approval[12]. It is unknown whether NRP1 contributes to NGF signaling in
bone tissue, considering its expression on osteoblasts[53]. Similar to the development of partial or biased
drugs that 'fine-tune' G protein-coupled receptor signaling to reduce adverse effects, targeting NGF
signaling *via* co-receptors/scaffolds could be advantageous for treating patients that suffer from chronic
pain without the side effects of systemic NGF sequestration. Identifying the antinociceptive efficacy of
vesencumab highlights an opportunity to repurpose a biologic developed for cancer for the treatment of
chronic pain, facilitating the development of a non-opioid therapeutic targeting NGF signaling through this
novel co-receptor.

ACKNOWLEDGEMENTS

CJP was funded by the Leon Levy Fellowship in Neuroscience (CP, 2022-2023). Supported by
grants from: National Institutes of Health (NS102722, DK118971, DE026806, DE029951, RM1DE033491
(NWB, BLS), GM147088 (ARBT) and Department of Defense (W81XWH1810431, W81XWH2210239,
NWB, BLS). We thank Laura Kilpatrick and Stephen Hill from the University of Nottingham, and Promega
Corporation for NRP1 cDNA plasmids. We thank Paz Durán for the provision of CAD.a cells.

AUTHOR CONTRIBUTIONS

C.J.P., R.T., K.G., M.R-S., A.C-R., L.M., A-M.M., S.P-M., H.H., A.R.B.T., S.D., B.L.S., M.K., R.K.,
N.W.B. designed and conducted the experiments, performed the analysis, discussed the results, and wrote
the paper.

COMPETING INTERESTS

Nigel W. Bunnett is a founding scientist of Endosome Therapeutics and pH-ARM Therapeutics.
Research in his laboratory is funded in part by Takeda Therapeutics.

MATERIALS AND METHODS

Compounds. Recombinant βNGF (R&D Systems or Alomone Labs) and recombinant NRP1 (R&D
Systems) were reconstituted according to manufacturer's instructions. GIPC1 inhibitors (CR1023 or
negative control) were from Genscript. Myosin VI inhibitor (2,4,6-triiodophenol) was from Thermo Scientific
Chemicals. Unless stated otherwise, *in vitro* studies used human NGF, whereas *ex vivo* and *in vivo* studies
used murine NGF.

cDNAs. SnapTag-TrkA was cloned with a GSSGAIA spacer and an N-terminal interleukin-6 signal
sequence (IL-6-SnapTag-TrkA). IL-6-NanoLuc-TrkA, IL-6-NanoLuc-p75[NTR] and TrkA-HA-Rluc8 were
cloned using Gibson assembly. HiBiT, a short fragment of NanoLuc[54], was cloned on the N-terminus of
VEGF$_{165}$a or C-terminus of βNGF with a GSSG linker. HaloTag-NRP1, SnapTag-NRP1 and NanoLuc-
NRP1 were from Dr. Laura Kilpatrick and Prof. Stephen Hill (University of Nottingham). SnapTag-CALCRL
was from Alex Thomsen (New York University). RGFP-CAAX (prenylation CAAX box of KRas), tdRGFP-
Rab5a, tdRGFP-Rab4a and tdRGFP-Giantin were from M. Bouvier (Université de Montréal). Nuclear and
cytosolic EKAR (CFP/YFP) FRET biosensors were from Addgene.

Cell Culture. HEK293T cells were grown in complete Dulbecco's modified Eagle's medium GlutaMAX^{TM}
supplemented with 10% fetal bovine serum (FBS) and 100 U/mL penicillin-streptomycin. CAD.a cells were
grown in RPMI 1640 containing 4% FBS, 8% horse serum and 100 U/mL penicillin-streptomycin.

siRNAs. ONTARGETplus siRNA targeting human GIPC1 (L-019997-00-0010), murine GIPC1 (L-062534-
00-0005) or nontargeting control (Ctrl) siRNA (D-001810-10-05) were from Dharmacon (**Supplementary
Table 3**).

Transfection of cell lines. HEK293T cells were transfected using polyethylenimine (PEI, Polysciences;
1:6 DNA:PEI) diluted in a 150 mM NaCl solution. CAD.a cells were transfected with Lipofectamine 3000
(according to manufacturer's instructions) diluted in serum-free DMEM.

Animals. Experiments were in accordance with the guidelines recommended by the National Institute of
Health, the International Association for the study of Pain, the National Centre for the Replacement,
Refinement and Reduction of Animals in Research (ARRIVE) guidelines, and were approved by the New
York University Institutional Animal Care and Use Committee. Male C57BL/6 mice (8-10 weeks, Charles
River) were housed four per cage at 22 ± 0.5°C under a controlled 14/10 h light/dark cycle with free access
to food and water. Mice were randomly assigned to experimental groups; group size was based on previous
similar studies. Investigators were blind to treatments.

Molecular docking. The predicted structure of full-length human NRP1 (UniProt O14786) was obtained
from AlphaFold. In the downloaded model, both the transmembrane region and the MAM domains were
folded against the a1a2b1b2 globular domains; therefore, domains were removed from the model prior to
docking, leaving only the a1a2b1b2 domains. The AlphaFold predicted structure of full-length NGF (UniProt
P01138) or the NGF/NRP1 complex was modeled using PIPER[55] from within the Schrödinger Biologics
Suite (Release 2022-1). For PIPER, the NMR structure of the human NGF dimer[56] contains a C-terminal
Arg118. All 20 NMR models were examined and one with the C-terminal residues in an extended
conformation was selected. Initial PIPER docking analysis revealed that the orientation of the a1 domain
potentially inhibited access of the NGF dimer to the b1 domain. Thus, the position of the NRP1 a1 domain
was modified to match that observed in the structure of mouse NRP1[21]. Model rebuilding was performed in
COOT with Ramachandran restraints. Models were prepared using the Schrödinger Suite and PIPER
docking was repeated using standard mode with a single attractive distance restraint that Arg118 (NGF
chain A) and Asp320 (NRP1) sidechains be within 2-4 Å. Other settings were left at default values: standard
mode, a maximum of 70,000 ligand rotations to probe, a maximum of 30 poses returned, and output poses
were refined. All of the 30 returned clusters had Arg118 in or near to the b1 pocket. Only three clusters had
the Arg118 side-chain terminal nitrogens oriented toward the terminal oxygens of Asp320. The largest of
these clusters (117 members, PIPER pose energy = 634 kcal/mol, PIPER pose score -66 kcal/mol) was
selected for visualization of potential interactions.

Microscale thermophoresis. Purified NRP1-His (Bio-Techne, 3870-N1) was fluorescently labeled using
the Monolith Protein Labeling kit RED-Tris-NTA (NanoTemper). 200 nM of NRP-His in Tris 40 mM, NaCl
250 mM and DTT 1 mM was mixed with NT-647-His-labeling dye (30 min, RT). Labeled NRP was
centrifuged (15,000 g, 10 min, 4°C). NRP1-His (15 nM) was mixed with increasing concentrations of
unlabeled NGF (10^{-10} M to 10^{-5} M) in PBS (10 min, RT). Thermophoresis was measured using a Monolith

NT.115 (NanoTemper), using premium MST capillaries at 100% MST power. Data were analyzed with MO
Affinity Analysis software using the K_d model.

Binding using HiBiT-BRET. HEK293T cells were transfected with SnapTag-NRP1 or SnapTag-NRP1
Y297A (3 μg/10 cm dish) using PEI. After 24 h, cells were transferred to 96-well plates. The next day, cells
were incubated with SNAP-Surface® Alexa Fluor® 488 (0.25 μM, 1 h, 37ºC), then washed in HBSS/0.1%
BSA. Separately, HEK293T cells were transfected with IL-6-HiBiT-VEGF$_{165}$a (3 μg/10 cm dish) or IL-6-
NGF-HiBiT (5 μg/dish). After 48 h, medium was replaced with HBSS/0.1% BSA to collect secreted growth
factor (2 h, 37ºC). Supernatant was diluted (1:2) in HBSS/0.1% BSA and incubated with recombinant
HaloTag-LgBiT protein and furimazine (1:50 final dilution for both; Promega Corporation). NRP1-expressing
cells were pre-incubated with vehicle or unlabeled VEGF$_{165}$a (10 nM, 30 min), followed by supernatant with
furimazine (1:2, 15 min, 37ºC). Luminescence and fluorescence emissions were recorded (Synergy Neo2,
Agilent BioTek; BRET1 donor 460 ± 40 nm, acceptor 540 ± 25 nm).

Collection of mouse tissue. Mice were anesthetized (5% isoflurane) and perfused through the ascending
aorta with PBS and then 4% paraformaldehyde in PBS. DRG (vertebrae L4-L5) were removed, fixed in 4%
paraformaldehyde in PBS (1 h, 4°C), cryoprotected in 30% sucrose (24 h, 4°C), and embedded in Optimal
Cutting Temperature compound (Tissue Tek). Frozen sections (10-12 μm) were mounted, dried (15 min)
and stored (-20°C).

Collection of human tissue. For RNAScope® studies, the collection of DRG from deidentified organ
donors was reviewed by the Institutional Review Board of the University of Cincinnati (#00003152, Study
ID 2015-5302) and deemed to be human subjects exempt. Donor information has been provided[57]. DRG
(L4-L5) were collected in the operating room within 90 min of aortic cross clamp and after removal of vital
organs. DRG were placed in N-methyl-D-glucamine-artificial cerebrospinal fluid (4°C), dissected and
immersion fixed in 4% paraformaldehyde in PBS (overnight, 4°C), cryoprotected in 30% sucrose (24 h,
4°C), and embedded in Optimal Cutting Temperature compound. Frozen sections (14-18 μm) were
mounted, dried (15 min) and stored (-80°C).

Immunofluorescence. Mouse tissue was blocked (2% bovine serum albumin, BSA, 0.2% Triton X-100 in
PBS; 1 h, room temperature (RT)). Sections were incubated (4ºC, overnight) with monoclonal rabbit anti-
TrkA (1:50; Millipore-Sigma, SU0354), and monoclonal mouse anti-NRP1 (1:200; Santa Cruz, A-12) or
monoclonal human anti-NRP1 vesencumab (1:500; Invitrogen, MA5-41940). Slides were washed and
incubated with donkey anti-rabbit Alexa Fluor® 488 (1:1000) and either donkey anti-mouse or anti-human
Alexa Fluor® 647 (1:1000, 1 h, RT). Slides were incubated with DAPI (1 μg/ml, 5 min) and mounted
(ProLong® Gold Antifade). Sections were imaged using a Leica SP8 confocal microscope with HCX PL
APO 10x air objective (Leica Microsystems).

RNAScope® *in situ* **hybridization in mouse and human tissue.** RNAScope Multiplex Fluorescent
Reagent v2 Assay (Advanced Cell Diagnostics Inc.) was carried out for fresh-frozen tissue as
recommended by the manufacturer, except for omission of the initial on-slide fixation step with mouse
tissue. Probes to Mm-Nrp1(#471621-C3), Mm-Ntrk1 (#435791-C2), Mm-Gipc1 (#1232971-C1), Hs-Nrp1
(#424361-C1) Hs-Ntrk1 (#402631-C2) and Hs-Gipc1 (#1232961-C3) were used. Mouse tissue was
incubated with Opal 620 (1:1000, FP1495001KT, Akoya Biosciences) and Opal 520 (FP1487001KT)
reagents for fluorescence detection. For human tissue, Tyramide Signal Amplification Plus Cyanine 3
(1:1500) and Cyanine 5 (1:1000) were used to detect the hybridized signals. To detect neurons in mouse
tissue, hybridized slides were blocked and incubated with guinea pig anti-NeuN antibody (1:500, overnight,
4°C; EMD Millipore, N90;). Slides were washed and incubated with goat anti-guinea pig Alexa Fluor® 647
(1:1000, 45 min, RT). Alternatively, neurons were detected using Nissl staining. Slides were washed,
incubated with DAPI and mounted. For mouse tissue, sections were imaged using a Leica SP8 confocal
microscope with HCX PL APO 40x (NA 1.30) oil objective. The percentage of hybridized positive neurons
were quantified and normalized by the total number of neurons. For human tissue, images were captured
as stitched Z stacks using Keyence BZ-X800E fluorescence microscope. Cells positive for Ntrk1, Nrp1 and
Gipc1 were counted to mark positive cells, where 10+ fluorescent puncta were considered to be positive.

Human DRG cultures. Human DRG suspension cells were obtained from AnaBios Corporation. Cells were
recovered with a gentle centrifugation at room temperature (3 min) and resuspension with DMEM/F12
media containing 10% horse serum, 1% penicillin-streptomycin, 25 ng/ml hNGF and 25 ng/ml GDNF. Cells
were seeded on poly-D-lysine (0.1 mg/ml) and laminin (1 mg/ml)-coated 12-mm glass coverslips. Half of
the culture medium was replaced with fresh medium every 3 days. All cultures used within 96 h.

Mouse DRG cultures. C57BL/6 mice (~3-4 weeks) were anaesthetized (5% isoflurane) and decapitated.
Thoracic and lumbar DRG were excised, then ganglia dissociated by collagenase type I (1 mg/ml) and
neutral protease (0.62 mg/ml; 45 min, 37°C) under gentle agitation. Dissociated cells were pelleted and
resuspended in DMEM containing 10% FBS and 1% penicillin-streptomycin. For siRNA transfection, cells
were resuspended in Nucleofector® transfection reagent containing GIPC1 or Ctrl siRNA (500 nM) and a
plasmid encoding the enhanced green fluorescent protein (eGFP, 1 µg). Cells were electroporated using
Amaxa Biosystem (Lonza) with protocol O-003. Neurons were plated on glass coverslips pretreated with
poly-D-lysine (0.1 mg/ml) and laminin (0.01 mg/ml; 30 min). Neurons were used within 48 h and siRNA
transfection verified by eGFP fluorescence.

Whole-cell patch-clamp recordings of evoked action potentials in mouse DRG neurons. For current-
clamp recordings, the external solution contained (mM): 154 NaCl, 5.6 KCl, 2 $CaCl_2$, 1 $MgCl_2$, 10 D-Glucose,
and 8 HEPES (pH 7.4 adjusted with KOH, and mOsm/L= 300). The internal solution comprised (mM): 137
KCl, 10 NaCl, 1 $MgCl_2$, 1 EGTA, and 10 HEPES (pH 7.3 adjusted with KOH, and mOsm/L= 277).
Recordings of action potentials were made at RT in whole-cell patch clamp configuration and current-clamp
mode. DRG neurons with a resting membrane potential (RMP) more hyperpolarized than −40 mV, stable
baseline recordings, and evoked spikes that overshot 0 mV were used for experiments and analysis. The
action potentials were evoked by a ramp pulse from 0-1000 pA in 1 sec for human DRG, or 0-250 pA in 1
sec for mouse DRG. Rheobase was measured by injecting currents from 0 pA with an increment of 50 pA
in 100 ms (human) or 10 pA in 50 ms (mouse). Analyses were performed by using Fitmaster software
(HEKA) and Origin 9.0 software (OriginLab).

Whole-cell patch-clamp recordings of Ca^{2+} and Na^+ currents in mouse DRG neurons. Currents were
recorded using an EPC 10 Amplifier-HEKA (HEKA Elektronik) with Patchmaster software. DRG neurons
were preincubated with EG00229 (30 µM, 30 min) or vehicle (0.1% DMSO) and then challenged with NGF
(50 nM, 30 min). Ca^{2+} currents (ICa^{2+}) and Na^+ (INa^+) currents were recorded as described[17,18]. In brief,
peak Ca^{2+} current was acquired by applying 200-millisecond voltage steps from −70 to +70 mV in 10-mV
increments from a holding potential of −90 mV to obtain the current-voltage (I-V) relation. Peak Na^+ current
was acquired by applying 150-millisecond voltage steps from −70 to +60 mV in 5-mV increments from a
holding potential of −60 mV to obtain the current-voltage (I-V) relation. Capacitive artifacts were fully
compensated, and series resistance was compensated by ~70 %. Recordings made from cells with greater
than a 20% shift in series resistance compensation error were excluded from the analysis. All recordings
were made at RT (~23 °C).

NGF-evoked nociception. Mouse NGF (50 ng/10 µl) was administered by i.pl. injection into the right
hindpaw of mice. EG00229 (MedChemExpress, HY-10799), compound 5[27], a human monoclonal anti-
NRP1 antibody (Vesencumab, Invitrogen, MA5-41940), or IgG control (Ctrl, ThermoFisher 02-6502) was
administered (i.pl.) concomitantly with NGF.

Inflammatory pain. CFA (1 mg/ml) or vehicle (0.9% NaCl) was administered (10 µl, i.pl.) into the right
hindpaw of sedated mice (2% isoflurane). NRP1 inhibitors were injected 48 h after CFA.

Intrathecal administration of siRNA. Mouse GIPC1 or Ctrl siRNA (1.25 µg) was mixed with *in vivo*-jetPEI
transfection reagent (8:1 PEI:DNA; Polyplus, 201-50G) and administered to conscious mice (5 µl, i.t., L4-
L5), 48 h before NGF (i.pl.) or 48 h after CFA (i.pl.). *Gipc1* mRNA in DRG (L4-L5) was analyzed by
RNAScope® *in situ* hybridization, 48 h after siRNA injection.

Nociception assays. Investigators were blinded to treatments. Mechanical allodynia was assessed by
measuring hindpaw withdrawal response to VFF stimulation using the up-and-down method[57]. The
Hargreaves apparatus was used to evaluate hypersensitivity to heat (Ugo Basile)[57].

Phosphorylated TrkA assays. Acutely dispersed mouse DRG neurons were plated onto coverslips pre-
coated with Cell-Tak (30 min, 37°C; CB-40240) in DMEM containing 10% FBS. After 18 h, neurons were
incubated with a monoclonal NGF antibody (1:500, 4 h, 37°C; MA5-41968) in serum-free DMEM to reduce
interference from endogenous NGF. Neurons were washed in HBSS/0.1% BSA, pre-incubated with
EG00229 (30 µM, 30 min) or control, then stimulated with vehicle or murine NGF (100 nM, 15 min, 37°C).
Neurons were fixed with 4% paraformaldehyde in PBS (20 min, 4°C), blocked (2% BSA, 0.2% Triton-X-100
in PBS), and incubated with a rabbit mAb to phosphorylated TrkA Y785/TrkB Y816 (1:500, overnight, 4°C;
Cell Signaling, 4168). Neurons were washed in PBS and incubated with donkey anti-rabbit Alexa Fluor®
488 (1:1:000, 1 h, RT). Slides were incubated with DAPI, mounted then imaged using a Leica SP8 confocal
microscope with HCX PL APO 40x (NA 1.30) oil objective. Mean fluorescence intensity was quantified from

regions of interest drawn using a phase contrast image with the maximum projection from four Z planes
(0.5 µm thickness). Images were quantified from 34-44 individual cells across three independent
experiments using separate mice.

ERK activity FRET biosensor assays. CAD.a cells were serum-starved and transfected with EKAR FRET
biosensors (300 ng/well) and human SnapTag-TrkA (150 ng/well). HEK293T-FLPN cells were transfected
with FRET biosensors (1 µg/10 cm dish), SnapTag-TrkA (1 µg), and pcDNA3.1 or HaloTag-NRP1 (2 µg).
After 24 h, cells were transferred to 96-well plates and serum-starved overnight. On the day of the assay,
cells were washed and incubated in Hanks' buffered saline solution (HBSS, pH 7.4) containing 0.1% BSA.
After 30 min (37°C), FRET was measured at 60 s intervals (CLARIOstar, BMG Labtech). After 5 baseline
reads, cells were stimulated with NGF (0.1 pM-10 nM) or phorbol 12,13-dibutyrate (PDBu, 10 µM). ΔFRET
represents the ratio (YFP/CFP), minus the mean from 5 initial reads and baseline-corrected to vehicle. Area
under the curve (AUC) was determined for each replicate.

ERK transcriptional assays. HEK293T-FLPN cells were transfected with SRE-Luc2P (3 µg/10 cm dish),
SnapTag-TrkA (1 µg) and either pcDNA3.1 or HaloTag-NRP1 (2 µg). After 24 h, cells were transferred to
96-well plates. CAD.a cells were transfected with SRE-Luc2P (250 ng/well, 96-well plate), SnapTag-TrkA
(150 ng/well) and 100 nM siRNA. After 24 h, cells were serum-starved overnight. Cells were then incubated
in HBSS/0.1% BSA and stimulated with NGF (1 pM-10 nM) or PDBu (10 µM). After 5 h stimulation (37°C),
cells were incubated with luciferin (1:2) and lysis buffer (1:5). After 5 min to enable the substrate reaction,
luminescence emissions were recorded (CLARIOstar).

Live imaging. HEK293T cells were transfected with SnapTag-TrkA (1 µg/10cm dish), HaloTag-NRP1 (2
µg) or both TrkA and NRP1. After 24 h, cells were transferred to 35 mm dishes. CAD.a cells were seeded
in 35 mm dishes, serum-starved for 24 h, and then transfected with SnapTag-TrkA (250 µg/35 mm),
HaloTag-NRP1 (500 µg) or both. The next day, proteins were labeled with SNAP-Surface® Alexa Fluor®
488 and membrane-impermeant HaloTag® Alexa Fluor® 660 (0.5 µM, 30 min, 37°C). Cells were washed
in HBSS and imaged on the Leica SP8 confocal.

Receptor-receptor BRET. For endpoint studies, HEK293T cells were transfected with a fixed
concentration of NanoLuc-NRP1, NanoLuc-p75[NTR] or NanoLuc-TrkA (10 ng/well, 96-well plate) and
increasing concentrations of SnapTag-TrkA or SnapTag-CALCRL (0-50 ng/well). For kinetic studies, cells
were transfected with NanoLuc-TrkA (10 ng/well), SnapTag-TrkA (25 ng/well) and either pcDNA3.1 or
HaloTag-NRP1 (50 ng/well). After 48 h, cells were incubated with SNAP-Surface® Alexa Fluor® 488 (0.5
µM, 30 min, 37°C), washed in HBSS before fluorescence emissions were measured (Synergy Neo2). In
some conditions, cells were incubated with NGF (30 nM, 30 min, 37°C) or vehicle. Cells were then incubated
with furimazine (10 µM, 5 min, Promega). For kinetic studies, cells were stimulated with increasing NGF
concentrations (1 pM-100 nM) following 5 baseline reads. Luminescence and fluorescence emissions were
quantified (Synergy Neo2; BRET1 filter).

Quantifying SNAP-Surface® Alexa Fluor® 488. HEK293T cells were transfected with SnapTag-TrkA (25
ng/well, 96-well plate) and pcDNA3.1 or HaloTag-NRP1 (50 ng/well). CAD.a cells were transfected with
SnapTag-TrkA (150 ng/well) and either pcDNA3.1 or HaloTag-NRP1 (300 ng/well). Cells were labeled using
SNAP-Surface® Alexa Fluor® 488 and HaloTag® Alexa Fluor® 660 (0.5 µM, 30 min, 37°C), washed in
HBSS/0.1% BSA and fluorescence emissions were recorded (CLARIOstar).

BRET trafficking. HEK293T cells were transfected with TrkA-HA-Rluc8 (15 ng/well, 96-well plate) and a
fluorescent marker (15 ng/well) for the plasma membrane (RGFP-CAAX), early endosome (tdRGFP-
Rab5a), recycling endosome (tdRGFP-Rab4a) or cis-Golgi apparatus (tdRGFP-Giantin). In some
conditions, cells were co-transfected with pcDNA3.1 or HaloTag-NRP1 (30 ng/well), or 100 nM siRNA.
Alternatively, CAD.a cells were transfected with TrkA-HA-Rluc8 (125 ng/well) in the absence or presence
of HaloTag-NRP1 (250 ng/well), as well as RGFP-CAAX, tdRGFP-Rab5a, tdRGFP-Rab4a or tdRGFP-
Giantin (150 ng/well). After 48 h, cells washed in HBSS/0.1% BSA were pre-incubated (30 min, 37°C) with
hypertonic sucrose (0.45 M), Pitstop 2 (30 µM; Abcam), an inhibitor of GIPC1 (CR1023, 300 µM) or its
negative control (CR2055), or an inhibitor of myosin VI (TIP, 50 µM). Cells were incubated with
coelenterazine purple (25 µg/ml, 10 min, Nanolight). For kinetic studies, 5 baseline reads were measured,
then cells were stimulated with NGF (1 pM-100 nM), using 30 nM NGF for endocytic inhibitors. BRET ratios
were recorded every 45 s (Synergy Neo2; BRET2 filter: donor 410 ± 80 nm, acceptor 515 ± 30 nm). ΔBRET
represents BRET in the presence of agonist, minus the vehicle BRET over time. For endpoint studies

lacking agonist stimulation, BRET ratios were baseline-corrected to TrkA-HA-Rluc8 expression alone
(100%).

Quantitative PCR (qPCR). RNA was isolated from cells or snap-frozen tissues using Direct-zol RNA
MiniPrep Kit (Zymo Research, R2050). cDNA was prepared using MultiScribe Reverse Transcriptase
(Thermo Fisher, 4311235). cDNA was amplified for 40 cycles by qRT-PCR using the QuantStudio 3 Real-
Time PCR System and TaqMan Fast Advanced PCR Mastermix (4444556). Primers to hTrkA
(Hs01021011_m1), mTrkA (Mm01219406_m1), hNgfr (Hs00609976_m1), mNgfr (Mm00446296_m1),
hGIPC1 (Hs00991802_m1), mGIPC1 (Mm00457561_m1), or GAPDH (Mn99999915_g1) were used. The
relative abundance of mRNA was calculated as described[57].

Data analysis. Data presented as mean ± SEM using GraphPad Prism (8.0). Images processed using
ImageJ. Illustrations were produced using Adobe Illustrator, Mol*Viewer or BioRender. Concentration-
response data fit using non-linear regression analysis as log *vs.* response (three parameters) to
determine EC_{50} values. Differences were assessed using paired or unpaired t-test for two comparisons,
and 1- or 2-way ANOVA and Tukey's, Dunnett's or Šidák's post-hoc test for multiple comparisons. *P*<0.05
was considered significant at the 95% confidence level.

FIGURES & FIGURE LEGENDS

Figure 1. NGF directly interacts with NRP1. (**A**) Carboxy-terminal sequences of NGF, highlighting
prospective CendR motifs (R/KxxR/K). Amino acids numbered according to the mature βNGF sequence
(1-120 equivalent to proNGF 122-241). *h, Homo sapiens; r, Rattus norvegicus; m, Mus musculus*. (**B**)
Prospective binding site between human βNGF and NRP1 modelled using protein docking PIPER software.
(**C**) Interaction between purified His-tagged human NRP1 (residues 22-644) and unlabeled human beta-
NGF using MST. Data from 4 independent experiments. Mean±SEM. (**D-E**) Bioluminescence resonance
energy transfer (BRET) to measure proximity (<10 nm) at full-length NRP1 in living HEK293T cells at 37ºC.
Supernatant was collected from cells secreting growth factor (VEGF$_{165}$a or NGF) tagged with HiBiT, a small
portion of nanoluciferase with high affinity. When reconstituted with recombinant LgBiT and luciferase
substrate, this acts as a bioluminescent donor for SnapTag-NRP1 labeled with SNAPTag-Alexa Fluor® 488
(AF488). BRET was compared to negative control (HiBiT/LgBiT only lacking AF488). Cells were pre-
incubated with vehicle or 10 nM unlabeled VEGF$_{165}$a (30 min), followed by luminescent growth factor (15
min, 37ºC). BRET with HiBiT-VEGF$_{165}$a was measured between HiBiT-VEGF$_{165}$a and either SnapTag-
NRP1 (WT) or the known VEGF$_{165}$a binding-dead mutant (Y297A), as well as HiBiT-tagged NGF (**E**). Data
from 4 independent experiments with triplicate wells. F. 1-way ANOVA, Šídák's multiple comparisons.
*$P<0.05$, **$P<0.01$, *** $P<0.001$, ****$P<0.0001$.

640
Figure 2. *Ntrk1*/TrkA and *Nrp1*/NRP1 are coexpressed in DRG. (A) TrkA and NRP1 immunofluorescence
in mouse DRG, showing subcellular TrkA (dashed arrows), and cells coexpressing TrkA and NRP1 (solid
arrows). Scale bar, 50 μm. (**B**) RNAScope® localization of *Ntrk1* and *Nrp1* mRNA in mouse DRG, showing
mRNA expression in the same neuron using NeuN immunofluorescence (arrowheads). Scale bar, 500 μm.
(**C**) RNAScope® localization of *Ntrk1* and *Nrp1* mRNA in human DRG, showing mRNA expression in the
same cell (arrowheads). * Denotes fluorescence due to lipofuscin (*). Nuclei shown in blue. (**D, E**) Quantified
co-expression of *Ntrk1* and *Nrp1* mRNA in mouse DRG (**D**) or human DRG (**E**). Representative images,
n=4-5 mice; n=3 human sections. Hybridized positive cells (%) were normalized to the total number of cells.

649
Figure 3. NRP1 inhibition prevents NGF-mediated increases in neuronal firing and ion channel
currents. Effect of NRP1 inhibitor EG00229 (30 μM, 30 min) on responses of dissociated human (A-D) or
mouse DRG neurons (**E-P**) to NGF (50 nM, 30 min). (**A-B,E-F**) Representative action potential firing evoked
by a depolarizing ramp stimulus (**A,E**), with summary of the number of evoked action potentials (**B,F**). N=7-
10 cells. (**C-D,G-H**) Resting membrane potential (**C,G,** millivolts, mV) and rheobase (**D,H**, picoamperes,
pA). (**E-H**) Representative family of Ca^{2+} current traces recorded from small diameter DRG neurons in
response to depolarization steps from -70 to +70 mV from a holding potential of -90 mV (**E**), with double
Boltzmann fits for current density-voltage curve (**F**), summary of peak calcium current densities (**G**; pA/pF)
and Boltzmann fits for voltage-dependence of activation and inactivation (**H**). N=7-10 cells. (**I-L**)
Representative family of Na^+ current traces, where currents were evoked by 150 ms pulses between −70
and +60 mV, with double Boltzmann fits for current density-voltage curve (**J**), summary of peak sodium
current densities (**K**, pA/pF) and Boltzmann fits for voltage-dependence of activation and inactivation (**L**).
N=10-14 cells. Mean±SEM. *P<0.05, ** P<0.01. B, C, D, G, K. Kruskal-Wallis, Dunn's multiple comparisons.

663
Figure 4. NRP1 inhibition abrogates NGF- and CFA-induced nociception in mice. (**A-G**) NGF-induced
nociception. Effects of an antibody against the b1 domain of NRP1 (**B, E**; 7 μg/10 μl i.pl.), or NRP1 inhibitors
EG00229 (**C, F**; 30 μM/10 μl i.pl.) or Compound 5 (**D, G**; Cpd5, C,F; 30 μM/10 μl i.pl.). After baseline ("B")
measurements, inhibitors were co-injected with murine NGF (50 ng/10 μl, i.pl.). Mechanical allodynia (**B-D**)
and thermal hyperalgesia (**E-G**) were measured. n=5-8 mice per group. **(H-L)** CFA-induced Inflammatory
pain. Effects of anti-NRP1 (**I, K**) or EG00229 (**J, L**). Inhibitors were injected (i.pl.) 48 h after CFA (i.pl.).
Mechanical allodynia (**I, J**) and thermal hyperalgesia (**K, L**) were measured. n=8-9 mice per group.
Mean±SEM. *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001 vs. PBS or IgG Ctrl. B-G, I-L. 2-way ANOVA,
Sídák's multiple comparisons.

Figure 5. NRP1 modulates TrkA-mediated kinase signaling. (A-B) Effect of murine NGF (15 min) and NRP1 inhibitor EG00229 (30 min pre-incubation) on TrkA Y785 staining in mouse DRG neurons. Scale bar, 20 µm. RFU, relative fluorescence units. N=34-44 neurons from 3 independent experiments. (**C**) NGF-induced ERK signaling measured using FRET-based EKAR biosensors (**D-I**) or a downstream luciferase reporter (**J**). (**D-I**) NGF-induced modulation of ERK using biosensors localized to the cytosol (**D, E**) or nucleus (**D, F**) in neuron-like CAD.a cells expressing human TrkA. Kinetics of NGF-induced ERK monitored in CAD.a cells (**D**), comparing increasing NGF concentrations after pre-incubation with EG00229 (30 µM, 30 min) (**E-F**). (**G-I**) ERK signaling in HEK293T cells expressing TrkA alone or transfected with both TrkA and NRP1 (**H, I**). (**J**) Effect of increasing NGF concentrations on ERK transcription in cells expressing TrkA, NRP1 or both (% positive control, 10 µM PDBu). AUC, area under curve. RLU, relative luminescence units. Mean±SEM. Data from 4-8 independent experiments with triplicate wells. *$P<0.05$, ***$P<0.001$, ****$P<0.0001$. B, I, J. 1-way ANOVA, Sídák's multiple comparisons.

Figure 6. TrkA and NRP1 form a heteromeric complex. (**A**) HEK293T or CAD.a cells expressing
SnapTag-TrkA and HaloTag-NRP1 simultaneously labeled with membrane-impermeant substrate
(SNAPTag-Alexa Fluor® 488, HaloTag-Alexa Fluor® 660). Representative images from n=5 independent
experiments. (**B-C**) BRET assays to monitor proximity between NanoLuc-NRP1 or NanoLuc-p75^{NTR} (10 ng)
and increasing SnapTag-TrkA DNA. Negative control, NanoLuc-TrkA and SnapTag-CALCRL.
Representative replicate (**C**) plotting BRET against relative fluorescence units (RFU) to quantify protein
expression. (**D-G**) Isolating cell-surface TrkA, imaged in HEK293T overlayed with phase-contrast imaging
in the absence or presence of NRP1 coexpression (**E**). (**F-G**) Quantification comparing fluorescence without
receptor (-), SnapTag-TrkA alone or SnapTag-TrkA co-transfected with NRP1 in HEK293T (**F**) or CAD.a
(**G**) cells. (**H-J**) BRET between TrkA tagged with *Renilla* luciferase (Rluc8) and *Renilla* green fluorescent
protein (RGFP) tagged markers of the plasma membrane (PM, RGFP-CAAX), early endosome (EE,
tdRGFP-Rab5a), recycling endosomes (RE, tdRGFP-Rab4a) or the cis-Golgi apparatus (tdRGFP-Giantin).
HEK293T cells (**I**) or CAD.a cells (**J**) were transfected with TrkA-Rluc8 in the absence (-) or presence (+)
of NRP1 co-expression. BRET was normalized relative to TrkA-Rluc8 alone (100%). Data from 5-6
independent experiments with triplicate wells. Mean±SEM. Scale bar, 20 μm. F, G. Paired t-test. I, J. 1-way
ANOVA with Šídák's multiple comparisons. *$P<0.05$, **$P<0.01$, ***$P<0.001$, ****$P<0.0001$.

Figure 7. NRP1 modulates NGF-induced TrkA trafficking and RTK oligomerization. (**A-G**) BRET was measured at 37ºC between Rluc8-tagged TrkA and a fluorescent marker of the plasma membrane (RGFP-CAAX). Decreased BRET indicates removal from the plasma membrane (**A**). (**B-D**) Trafficking in HEK293T cells incubated with increasing NGF concentrations (**B**), or pre-incubated for 30 with hypertonic sucrose (0.45 M), clathrin inhibitor pitstop 2 (30 µM) or vehicle (Veh) (**C**). Effect of NRP1 co-expression (**D**). (**E-G**) Trafficking in CAD.a cells expressing human TrkA with increasing NGF concentrations (**E**), pre-incubation with endocytic inhibitors (**F**) with transfected with control or murine NRP1 siRNA (**G**). (**H-K**) BRET measured between NanoLuc- and SnapTag- TrkA to measure RTK oligomerization in HEK293T cells. (**I**) BRET with increasing SnapTag-TrkA and fixed NanoLuc-TrkA (10 ng), after 30 min incubation with vehicle or NGF. (**J-K**) Using a fixed donor:acceptor ratio (1:2.5), oligomerization kinetics in response to increasing NGF concentrations (**J**) in the absence and presence of NRP1 co-expression (**K**). Kinetic measurements were made at 37ºC. Data from 4-6 independent experiments with triplicate wells. AUC, area under curve. Mean±SEM. *$P<0.05$, **$P<0.01$, ***$P<0.001$, ****$P<0.0001$. G, Unpaired t-test. D, I, K. 1-way ANOVA with Sídák's multiple comparisons.

Figure 8. GIPC1 modulates TrkA trafficking, signaling and NGF-induced nociception. (**A-C**) RNAScope® localization of *Gipc1* mRNA in mouse DRG (**A**) or *Ntrk1* and *Gipc1* mRNA in human DRG (**B, C**). Arrows indicate mRNA expression within the same cell. Representative images, n=5 mice; n=3 human sections. Scale bar, 500 μm. (**D**) Effect of GIPC1 siRNA on increased TrkA at the plasma membrane under basal conditions upon NRP1 co-expression in HEK293T cells measured using BRET. (**E-F**) Effect of 30 min pre-incubation of GIPC1 antagonist (300 μM CR1023 or negative control) or myosin VI inhibitor (50 μM 2,4,6-Triiodophenol, TIP) on NGF-induced TrkA-Rluc8 trafficking from a marker of the plasma membrane (RGFP-CAAX) in CAD.a cells. (**G**) Effect of GIPC1 siRNA on NGF-induced downstream ERK transcription in CAD.a cells. RLU, relative luminescence units. Data from 5-6 independent experiments with triplicate wells. (**H-I**) Sample traces of action potential firing in mouse DRG neurons evoked by injecting a 1 s ramp pulse from 0 to 250 pA (**G**), with the number of evoked action potentials (**H**). N=7-10 cells. (**J-O**) Effects of GIPC1 siRNA (i.t.) on NGF (50 ng/10 μl, i.pl.)-induced (**J-L**) or CFA-induced (**M-O**) mechanical allodynia (**J-K,M-N**) and thermal hyperalgesia (**L,O**) in mice, including mechanical allodynia tested in the contralateral hind paw after NGF (**K**) or CFA injection (**N**). N=6-8 mice per group. AUC, area under curve. 'B', basal. Mean±SEM. C,E. 1-way ANOVA, Sídák's multiple comparisons. H, Tukey's multiple comparison. I-O, 2-way ANOVA, Sídák's multiple comparisons. *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001.

Figure 9. Proposed mechanism for NRP1- and GIPC1-mediated NGF/TrkA signaling. NGF released from sites of injury, inflammation or the tumor microenvironment act on TrkA expressed by primary afferent neurons. NRP1 and GIPC1 potentiate kinase signaling, neuronal excitability and nociception. As a co-receptor, NRP1 can directly bind NGF and act as a chaperone, mediated by interactions with adaptor protein GIPC1.

743 **TABLES**

744 **Suppl. Table 1. Gating properties of Ca^{2+} and Na^+ currents recorded from mouse DRG neurons.**

Total Ca^{2+}	DMSO (0.1%)	EG00229 (30 µM)	NGF (50 nM)	EG00229 + NGF
Activation				
$V_{1/2}$	5.793 ± 1.891 (10)	-2.828 ± 0.871 (6)†	5.834 ± 1.995 (9)††	-0.473 ± 1.265 (8)
k	7.756 ± 1.534 (10)	4.230 ± 0.731 (6)	7.972 ± 1.608 (9)	6.428 ± 1.109 (8)
Inactivation				
$V_{1/2}$	-27.372 ± 1.919 (9)	-20.137 ± 0.901 (7)	-23.909 ± 3.080 (9)	-20.166 ± 1.314 (8)
k	-14.944 ± 2.109 (9)	-10.623 ± 0.842 (7)	-15.079 ± 3.305 (9)	-13.176 ± 1.287 (8)

Total Na^+	DMSO (0.1%)	EG00229 (30 µM)	NGF (50 nM)	EG00229 + NGF
Activation				
$V_{1/2}$	-22.436 ± 1.150 (10)	-23.229 ± 0.499 (13)	-23.151 ± 0.418 (14)	-23.600 ± 0.993 (11)
k	5.098 ± 1.014 (10)	3.580 ± 0.434 (13)	3.432 ± 0.362 (14)	5.145 ± 0.877 (11)
Inactivation				
$V_{1/2}$	-48.736 ± 4.219 (10)	-43.306 ± 4.249 (13)	-54.450 ± 2.244 (14)	-38.622 ± 4.013 (11)*
k	-18.568 ± 4.934 (10)	-21.036 ± 5.439 (13)	-14.748 ± 2.243 (14)	-19.732 ± 4.708 (11)

Values are mean±SEM calculated from fits of the data from the indicated number of individual cells (in
parentheses) to the Boltzmann equation; $V_{1/2}$ midpoint potential (mV) for voltage-dependent of activation or
inactivation; k, slope factor. Data were analyzed with 1-way ANOVA with Tukey's multiple comparisons. †
p=0.0115 for $V_{1/2}$ activation of EG00229 vs DMSO (0.1%); †† p=0.0131 for $V_{1/2}$ activation of NGF vs
EG00229; * p= 0.0175 for $V_{1/2}$ inactivation of EG00229 + NGF vs DMSO (0.1%).

Suppl. Table 2. Pharmacological parameters (pEC_{50}) of NGF derived in at human TrkA, in the
absence or presence of NRP1 modulation.

CAD.a cells	Control	NRP1 inhibition (EG00229)
Cytosolic ERK (FRET)	10.64 ± 0.41 (5)	10.53 ± 0.17 (5)
Nuclear ERK (FRET)	11.09 ± 0.33 (5)	10.01 ± 0.38 (5) *
HEK293T cells	**Control**	**NRP1 overexpression (HaloTag)**
Cytosolic ERK (FRET)	9.89 ± 0.21 (5)	9.87 ± 0.10 (4)
Nuclear ERK (FRET)	10.37 ± 0.19 (5)	10.59 ± 0.33 (5)
ERK Transcription (SRE-Luc2P)	9.67 ± 0.48 (8)	10.15 ± 0.57 (8) ***
TrkA-Rluc8 Trafficking (BRET)	9.06 ± 0.10 (6)	9.08 ± 0.13 (6)
TrkA/TrkA Oligomerization (BRET)	7.81 ± 0.07 (4)	7.87 ± 0.04 (5)

752 Derived in HEK293T or neuron-like CAD.a cells transfected with full-length, human TrkA. Values are
753 mean±SEM calculated from non-linear regressions fit of the data from independent replicates (in
754 parentheses), each with triplicate wells. Quantified as the negative antilog of the half-maximal effective
755 concentration (pEC_{50} = -$logEC_{50}$). * $p<0.05$, *** $p<0.001$. Paired t-test.

756 **Suppl. Table 3. Summary of siRNA sequences used in this study**.

Target	Sequence
Control	UGGUUUACAUGUCGACUAA UGGUUUACAUGUUGUGUGA UGGUUUACAUGUUUUCUGA UGGUUUACAUGUUUUCCUA
mNrp1	GAAUUGCUGUGGAUGAUAU AGUAAGAGGUGUCAUCAUU CCACAAGGUUCAUCAGGAU GGAAUGUUCUGUCGCUAUG
hGipc1	GCACUCGGGCUCACCAUCA GGCCGUACCUUCACGCUGA GCAAGGCCUUCGACAUGAU CUGGAGAGUUACAUGGGUA
mGipc1	GCAUCGAGGGCUUCACUAA CGUCGGCCUUUGAGGAGAA GUGGAUGACUUGCUAGAGA GCUGAGGCCUUCCGACUAC

SUPPLEMENTARY FIGURES

Supplementary Figure 1. Characterization of receptor expression in neuron-like CAD.a cells. (**A**)
Quantification of expression of *Ntrk1* mRNA (TrkA) or *Ngfr* (p75NTR) in HEK293T or CAD.a cells determined
by quantitative RT-PCR. Data from 3 independent experiments with triplicate wells. (**B**) Immunofluorescent
(IF) staining of NRP1 in HEK293T or CAD.a cells using vesencumab. Scale bar, 20 µm. Representative
from 4 independent experiments. Mean±SEM. A, 1-way ANOVA, Dunnett's multiple comparisons. * $P<0.05$.

Supplementary Figure 2. Specificity of SnapTag- and HaloTag- labeling in live cells. HEK293T cells (**A**) or CAD.a cells (**B**) were transfected with SnapTag-TrkA, HaloTag-NRP1 or both (1:2 TrkA to NRP1 ratio). Tags were simultaneously labeled with membrane-impermeant substrate (SNAPTag-Alexa Fluor® 488, HaloTag-Alexa Fluor® 660). Representative from n=5 independent experiments. Scale bar, 20 μm.

Supplementary Figure 3. Effect of co-expression or siRNA on TrkA and NRP1 expression. (**A-D**)
HEK293T (***A,C***) or CAD.a (***B,D***) cells expressed TrkA-Rluc8 in the absence or presence of HaloTag-NRP1.
Cells were labeled with membrane-impermeant HaloTag-Alexa Fluor® 660 and emissions were measured
for AlexaFluor660 as relative fluorescence units (RFU) (***A,B***). Alternatively, TrkA-Rluc8 emissions were
measured after incubation with coelenterazine purple to infer TrkA expression as relative luminescence
units (RLU) (***C,D***). (**E-F**) Effect of NRP1 siRNA on expression of NRP1 (***E***) or TrkA (***F***) quantified using IF
staining normalized to nuclei count (DAPI). Data expressed as percentage change over control (100%).
Mean±SEM. A-D, G, H. Unpaired t-test. **$P<0.01$, *** $P<0.001$.

Supplementary Figure 4. Confirmation of GIPC1 knockdown using siRNA. (**A-C**) Quantification of
expression of *Gipc1* mRNA in HEK293T (**A**), CAD.a cells (**B**) or cultured dissociated DRG neurons (**C**) in
the presence of control (Ctrl) or GIPC1 siRNA determined by quantitative RT-PCR. Data from 3-4
independent experiments with triplicate wells. (**D**) RNAScope® localization of *Gipc1* mRNA in murine DRG
after intrathecal injection of control or GIPC1 siRNA. Representative from n=5 mice. A-D. Unpaired t-test.
*$P<0.05$, **$P<0.01$, *** $P<0.001$, **** $P<0.0001$.

REFERENCES

Levi-Montalcini, R. & Hamburger, V. Selective growth stimulating effects of mouse sarcoma on the sensory and sympathetic nervous system of the chick embryo. *J Exp Zool* **116**, 321-361 (1951). https://doi.org:10.1002/jez.1401160206

Klein, R., Jing, S. Q., Nanduri, V., O'Rourke, E. & Barbacid, M. The trk proto-oncogene encodes a receptor for nerve growth factor. *Cell* **65**, 189-197 (1991). https://doi.org:10.1016/0092-8674(91)90419-y

Ye, M., Lehigh, K. M. & Ginty, D. D. Multivesicular bodies mediate long-range retrograde NGF-TrkA signaling. *Elife* **7** (2018). https://doi.org:10.7554/eLife.33012

Grimes, M. L., Beattie, E. & Mobley, W. C. A signaling organelle containing the nerve growth factor-activated receptor tyrosine kinase, TrkA. *Proc Natl Acad Sci U S A* **94**, 9909-9914 (1997). https://doi.org:10.1073/pnas.94.18.9909

Denk, F., Bennett, D. L. & McMahon, S. B. Nerve Growth Factor and Pain Mechanisms. *Annual Review of Neuroscience* **40**, 307-325 (2017). https://doi.org:10.1146/annurev-neuro-072116-031121

Banerjee, S. P., Snyder, S. H., Cuatrecasas, P. & Greene, L. A. Binding of nerve growth factor receptor in sympathetic ganglia. *Proc Natl Acad Sci U S A* **70**, 2519-2523 (1973). https://doi.org:10.1073/pnas.70.9.2519

Johnson, D. *et al.* Expression and structure of the human NGF receptor. *Cell* **47**, 545-554 (1986). https://doi.org:https://doi.org/10.1016/0092-8674(86)90619-7

Hempstead, B. L., Martin-Zanca, D., Kaplan, D. R., Parada, L. F. & Chao, M. V. High-affinity NGF binding requires coexpression of the trk proto-oncogene and the low-affinity NGF receptor. *Nature* **350**, 678-683 (1991). https://doi.org:10.1038/350678a0

Indo, Y. *et al.* Mutations in the TRKA/NGF receptor gene in patients with congenital insensitivity to pain with anhidrosis. *Nat Genet* **13**, 485-488 (1996). https://doi.org:10.1038/ng0896-485

Einarsdottir, E. *et al.* A mutation in the nerve growth factor beta gene (NGFB) causes loss of pain perception. *Hum Mol Genet* **13**, 799-805 (2004). https://doi.org:10.1093/hmg/ddh096

Lane, N. E. *et al.* Tanezumab for the treatment of pain from osteoarthritis of the knee. *N Engl J Med* **363**, 1521-1531 (2010). https://doi.org:10.1056/NEJMoa0901510

Hochberg, M. C. *et al.* When Is Osteonecrosis Not Osteonecrosis?: Adjudication of Reported Serious Adverse Joint Events in the Tanezumab Clinical Development Program. *Arthritis Rheumatol* **68**, 382-391 (2016). https://doi.org:10.1002/art.39492

Ray, P. *et al.* Comparative transcriptome profiling of the human and mouse dorsal root ganglia: an RNA-seq-based resource for pain and sensory neuroscience research. *Pain* **159**, 1325-1345 (2018). https://doi.org:10.1097/j.pain.0000000000001217

Broz, M., Kolarič, A., Jukič, M. & Bren, U. Neuropilin (NRPs) Related Pathological Conditions and Their Modulators. *International Journal of Molecular Sciences* **23**, 8402 (2022).

Soker, S., Takashima, S., Miao, H. Q., Neufeld, G. & Klagsbrun, M. Neuropilin-1 is expressed by endothelial and tumor cells as an isoform-specific receptor for vascular endothelial growth factor. *Cell* **92**, 735-745 (1998). https://doi.org:10.1016/s0092-8674(00)81402-6

Xin, Y. *et al.* Pharmacokinetic and pharmacodynamic analysis of circulating biomarkers of anti-NRP1, a novel antiangiogenesis agent, in two phase I trials in patients with advanced solid tumors. *Clin Cancer Res* **18**, 6040-6048 (2012). https://doi.org:10.1158/1078-0432.CCR-12-1652

Moutal, A. *et al.* SARS-CoV-2 spike protein co-opts VEGF-A/neuropilin-1 receptor signaling to induce analgesia. *Pain* **162**, 243-252 (2021). https://doi.org:10.1097/j.pain.0000000000002097

Gomez, K. *et al.* Neuropilin-1 is essential for vascular endothelial growth factor A-mediated increase of sensory neuron activity and development of pain-like behaviors. *Pain* (2023). https://doi.org:10.1097/j.pain.0000000000002970

Kolodkin, A. L. *et al.* Neuropilin is a semaphorin III receptor. *Cell* **90**, 753-762 (1997). https://doi.org:10.1016/s0092-8674(00)80535-8

Teesalu, T., Sugahara, K. N., Kotamraju, V. R. & Ruoslahti, E. C-end rule peptides mediate neuropilin-1-dependent cell, vascular, and tissue penetration. *Proc Natl Acad Sci U S A* **106**, 16157-16162 (2009). https://doi.org:10.1073/pnas.0908201106

Janssen, B. J. *et al.* Neuropilins lock secreted semaphorins onto plexins in a ternary signaling complex. *Nat Struct Mol Biol* **19**, 1293-1299 (2012). https://doi.org:10.1038/nsmb.2416

22   Parker, M. W., Xu, P., Li, X. & Vander Kooi, C. W. Structural basis for selective vascular
endothelial growth factor-A (VEGF-A) binding to neuropilin-1. *J Biol Chem* **287**, 11082-11089
(2012). https://doi.org:10.1074/jbc.M111.331140
23   Teran, M. & Nugent, M. A. Characterization of receptor binding kinetics for vascular endothelial
growth factor-A using SPR. *Anal Biochem* **564-565**, 21-31 (2019).
https://doi.org:10.1016/j.ab.2018.10.001
24   Pan, Q. *et al.* Neuropilin-1 binds to VEGF121 and regulates endothelial cell migration and
sprouting. *J Biol Chem* **282**, 24049-24056 (2007). https://doi.org:10.1074/jbc.M703554200
25   Jarvis, A. *et al.* Small molecule inhibitors of the neuropilin-1 vascular endothelial growth factor A
(VEGF-A) interaction. *J Med Chem* **53**, 2215-2226 (2010). https://doi.org:10.1021/jm901755g
26   Peach, C. J. *et al.* Real-Time Ligand Binding of Fluorescent VEGF-A Isoforms that Discriminate
between VEGFR2 and NRP1 in Living Cells. *Cell Chem Biol* **25**, 1208-1218 e1205 (2018).
https://doi.org:10.1016/j.chembiol.2018.06.012
27   Perez-Miller, S. *et al.* Novel Compounds Targeting Neuropilin Receptor 1 with Potential To
Interfere with SARS-CoV-2 Virus Entry. *ACS Chem Neurosci* **12**, 1299-1312 (2021).
https://doi.org:10.1021/acschemneuro.0c00619
28   Wehrman, T. *et al.* Structural and mechanistic insights into nerve growth factor interactions with
the TrkA and p75 receptors. *Neuron* **53**, 25-38 (2007).
https://doi.org:10.1016/j.neuron.2006.09.034
29   Qi, Y., Wang, J. K., McMillian, M. & Chikaraishi, D. M. Characterization of a CNS cell line, CAD,
in which morphological differentiation is initiated by serum deprivation. *J Neurosci* **17**, 1217-1225
(1997). https://doi.org:10.1523/JNEUROSCI.17-04-01217.1997
30   Peach, C. J., Kilpatrick, L. E., Woolard, J. & Hill, S. J. Use of NanoBiT and NanoBRET to monitor
fluorescent VEGF-A binding kinetics to VEGFR2/NRP1 heteromeric complexes in living cells. *Br J
Pharmacol* **178**, 2393-2411 (2021). https://doi.org:10.1111/bph.15426
31   Namkung, Y. *et al.* Monitoring G protein-coupled receptor and β-arrestin trafficking in live cells
using enhanced bystander BRET. *Nature Communications* **7**, 12178 (2016).
https://doi.org:10.1038/ncomms12178
32   Cai, H. & Reed, R. R. Cloning and characterization of neuropilin-1-interacting protein: a PSD-
95/Dlg/ZO-1 domain-containing protein that interacts with the cytoplasmic domain of neuropilin-1.
J Neurosci **19**, 6519-6527 (1999). https://doi.org:10.1523/JNEUROSCI.19-15-06519.1999
33   Lou, X., Yano, H., Lee, F., Chao, M. V. & Farquhar, M. G. GIPC and GAIP form a complex with
TrkA: a putative link between G protein and receptor tyrosine kinase pathways. *Mol Biol Cell* **12**,
615-627 (2001). https://doi.org:10.1091/mbc.12.3.615
34   Naccache, S. N., Hasson, T. & Horowitz, A. Binding of internalized receptors to the PDZ domain
of GIPC/synectin recruits myosin VI to endocytic vesicles. *Proc Natl Acad Sci U S A* **103**, 12735-
12740 (2006). https://doi.org:10.1073/pnas.0605317103
35   Muders, M. H. *et al.* Targeting GIPC/synectin in pancreatic cancer inhibits tumor growth. *Clin
Cancer Res* **15**, 4095-4103 (2009). https://doi.org:10.1158/1078-0432.CCR-08-2837
36   Heissler, S. M. *et al.* Kinetic properties and small-molecule inhibition of human myosin-6. *FEBS
Lett* **586**, 3208-3214 (2012). https://doi.org:10.1016/j.febslet.2012.07.014
37   Herzog, B., Pellet-Many, C., Britton, G., Hartzoulakis, B. & Zachary, I. C. VEGF binding to NRP1
is essential for VEGF stimulation of endothelial cell migration, complex formation between NRP1
and VEGFR2, and signaling via FAK Tyr407 phosphorylation. *Mol Biol Cell* **22**, 2766-2776 (2011).
https://doi.org:10.1091/mbc.E09-12-1061
38   Martino, M. M., Briquez, P. S., Ranga, A., Lutolf, M. P. & Hubbell, J. A. Heparin-binding domain of
fibrin(ogen) binds growth factors and promotes tissue repair when incorporated within a synthetic
matrix. *Proc Natl Acad Sci U S A* **110**, 4563-4568 (2013).
https://doi.org:10.1073/pnas.1221602110
39   Mahadeo, D., Kaplan, L., Chao, M. V. & Hempstead, B. L. High affinity nerve growth factor
binding displays a faster rate of association than p140trk binding. Implications for multi-subunit
polypeptide receptors. *J Biol Chem* **269**, 6884-6891 (1994).
40   He, X. L. & Garcia, K. C. Structure of nerve growth factor complexed with the shared neurotrophin
receptor p75. *Science* **304**, 870-875 (2004). https://doi.org:10.1126/science.1095190

41   Lin, D. C. *et al.* APPL1 associates with TrkA and GIPC1 and is required for nerve growth factor-
mediated signal transduction. *Mol Cell Biol* **26**, 8928-8941 (2006).
https://doi.org:10.1128/MCB.00228-06
42   Koch, S. *et al.* NRP1 presented in trans to the endothelium arrests VEGFR2 endocytosis,
preventing angiogenic signaling and tumor initiation. *Dev Cell* **28**, 633-646 (2014).
https://doi.org:10.1016/j.devcel.2014.02.010
43   Malik-Hall, M., Dina, O. A. & Levine, J. D. Primary afferent nociceptor mechanisms mediating
NGF-induced mechanical hyperalgesia. *Eur J Neurosci* **21**, 3387-3394 (2005).
https://doi.org:10.1111/j.1460-9568.2005.04173.x
44   Sung, K. *et al.* Swedish Nerve Growth Factor Mutation (NGF(R100W)) Defines a Role for TrkA
and p75(NTR) in Nociception. *J Neurosci* **38**, 3394-3413 (2018).
https://doi.org:10.1523/JNEUROSCI.1686-17.2018
45   Kuruvilla, R., Ye, H. & Ginty, D. D. Spatially and functionally distinct roles of the PI3-K effector
pathway during NGF signaling in sympathetic neurons. *Neuron* **27**, 499-512 (2000).
https://doi.org:10.1016/s0896-6273(00)00061-1
46   Delcroix, J. D. *et al.* NGF signaling in sensory neurons: evidence that early endosomes carry
NGF retrograde signals. *Neuron* **39**, 69-84 (2003). https://doi.org:10.1016/s0896-6273(03)00397-
0
47   Jensen, D. D. *et al.* Neurokinin 1 receptor signaling in endosomes mediates sustained
nociception and is a viable therapeutic target for prolonged pain relief. *Sci Transl Med* **9** (2017).
https://doi.org:10.1126/scitranslmed.aal3447
48   Yamashita, N., Yamane, M., Suto, F. & Goshima, Y. TrkA mediates retrograde semaphorin 3A
signaling through plexin A4 to regulate dendritic branching. *J Cell Sci* **129**, 1802-1814 (2016).
https://doi.org:10.1242/jcs.184580
49   Estrada-Bernal, A. *et al.* Functional complexity of the axonal growth cone: a proteomic analysis.
PLoS One **7**, e31858 (2012). https://doi.org:10.1371/journal.pone.0031858
50   Reza, J. N., Gavazzi, I. & Cohen, J. Neuropilin-1 is expressed on adult mammalian dorsal root
ganglion neurons and mediates semaphorin3a/collapsin-1-induced growth cone collapse by small
diameter sensory afferents. *Mol Cell Neurosci* **14**, 317-326 (1999).
https://doi.org:10.1006/mcne.1999.0786
51   Gormley, P. *et al.* Meta-analysis of 375,000 individuals identifies 38 susceptibility loci for
migraine. *Nat Genet* **48**, 856-866 (2016). https://doi.org:10.1038/ng.3598
52   Sarchielli, P. & Gallai, V. Nerve growth factor and chronic daily headache: a potential implication
for therapy. *Expert Rev Neurother* **4**, 115-127 (2004). https://doi.org:10.1586/14737175.4.1.115
53   Harper, J., Gerstenfeld, L. C. & Klagsbrun, M. Neuropilin-1 expression in osteogenic cells: down-
regulation during differentiation of osteoblasts into osteocytes. *J Cell Biochem* **81**, 82-92 (2001).
https://doi.org:10.1002/1097-4644(20010401)81:1<82::aid-jcb1025>3.0.co;2-p
54   Dixon, A. S. *et al.* NanoLuc Complementation Reporter Optimized for Accurate Measurement of
Protein Interactions in Cells. *ACS Chem Biol* **11**, 400-408 (2016).
https://doi.org:10.1021/acschembio.5b00753
55   Chuang, G. Y., Kozakov, D., Brenke, R., Comeau, S. R. & Vajda, S. DARS (Decoys As the
Reference State) potentials for protein-protein docking. *Biophys J* **95**, 4217-4227 (2008).
https://doi.org:10.1529/biophysj.108.135814
56   Paoletti, F. *et al.* Endogenous modulators of neurotrophin signaling: Landscape of the transient
ATP-NGF interactions. *Comput Struct Biotechnol J* **19**, 2938-2949 (2021).
https://doi.org:10.1016/j.csbj.2021.05.009
57   Tonello, R. *et al.* The contribution of endocytosis to sensitization of nociceptors and synaptic
transmission in nociceptive circuits. *Pain* **164**, 1355-1374 (2023).
https://doi.org:10.1097/j.pain.0000000000002826