1

**Individual dietary amino acid restrictions induce distinct metabolic and chromatin states**

Spencer A. Haws[1,2], Yang Liu[3,4], Cara L. Green[3,4], Krittisak Chaiyakul[5], Pragyan Mishra[1,2], Reji Babygirija[3,4,6], Eric A. Armstrong[1,2], Anusha T. Mehendale[1,2], Irene M. Ong[5,7,8,9], Dudley W. Lamming[3,4,6,9,10*], and John M. Denu[1,2,9,10*].

[1]Wisconsin Institute for Discovery, University of Wisconsin-Madison, Madison, WI 53715, USA.

[2]Department of Biomolecular Chemistry, University of Wisconsin-Madison, Madison, WI 53706, USA.

[3]Department of Medicine, University of Wisconsin-Madison, Madison, WI 53705, USA.

[4]William S. Middleton Memorial Veterans Hospital, Madison, WI 53705, USA.

[5]Department of Biostatistics and Medical Informatics, University of Wisconsin-Madison, Madison, WI 53705, USA.

[6]Cellular and Molecular Biology Graduate Program, University of Wisconsin-Madison, Madison, WI, 53706, USA.

[7]Department of Obstetrics and Gynecology, University of Wisconsin-Madison, Madison, WI 53705, USA.

[8]University of Wisconsin Center for Human Genomics and Precision Medicine, University of Wisconsin, Madison, WI 53705, USA.

[9]University of Wisconsin Carbone Comprehensive Cancer Center, University of Wisconsin, Madison, WI 53705, USA.

[10]University of Wisconsin Comprehensive Diabetes Center, Madison, WI 53705, USA

*  **Correspondence**

Dudley W. Lamming, PhD
Associate Professor of Medicine
University of Wisconsin-Madison
1685 Highland Ave
MFCB Rm 4147
Madison, WI 53705
dlamming@medicine.wisc.edu
Tel: 608-262-7341

John M. Denu, PhD
Professor of Biomolecular Chemistry
University of Wisconsin-Madison
330 N Orchard St.
Madison, WI 53715
john.denu@wisc.edu
Tel: 608-316-4341

## **Abstract**

Dietary protein and essential amino acid (EAA) restriction promote favorable metabolic reprogramming, although the extent to which shared or EAA-specific mechanisms facilitate diet-associated phenotypes remains unclear. Here, we compared the physiological and molecular effects of dietary methionine, leucine, or isoleucine depletion (Met-D, Leu-D, and Ile-D) in C57BL/6J mice. Each diet elicited responses not phenocopied by mTORC1 inhibition, including reduced fat mass and hepatic amino acid catabolism. Ile-D yielded additional distinct responses, highlighted by histone H2A/H4 hypoacetylation and maintained hepatic acetyl-CoA levels despite downregulated FA β-oxidation. Multi-Omics Factor Analysis of 14,139 data points objectively affirmed Ile-D phenotypes are distinct from Met-D or Leu-D and identified several metabolic and chromatin features as primary discriminators. Metabolic and epigenetic responses to Ile-D were recapitulated *in vitro*, suggesting underlying mechanisms represent fundamental cellular properties. Together, these results demonstrate EAAs can stimulate unique phenotypes and highlight distinct molecular mechanisms by which EAAs may inform metabolic health.

Keywords: methionine, leucine, isoleucine, epigenetics, post-translational modification, protein restriction

## Introduction

Dietary protein is a key regulator of metabolism, health, and longevity ([1]). It has long been suspected that the health benefits of a low protein diet arise not simply due to the consumption of reduced protein, but due to decreased levels of specific amino acids. For many years, attention has focused on the potential role of dietary methionine in regulating metabolism and health. Plant-based protein may have reduced levels of methionine compared to animal proteins, and vegans have lower than normal plasma levels of methionine; thus, many have attributed the health benefits of a vegan diet to decreased consumption of methionine ([2–4]). The potential metabolic benefits of reduced methionine intake are readily observed in rodents, where restriction of methionine extends lifespan, promotes leanness and insulin sensitivity, and promotes the rapid loss of adipose tissue and improved glycemic control in diet-induced obese mice ([5–9]).

However, methionine is only one of the twenty common amino acids, and there is a growing awareness that other dietary essential amino acids have critical roles in regulating metabolism and health. The most well-known of these are the branched-chain amino acids (BCAAs; leucine, isoleucine, and valine), which are elevated in individuals with diabetes, are associated with insulin resistance in humans and rodents, and which promote insulin resistance in rats when supplemented in the diet ([10]). Conversely, dietary restriction of BCAAs promotes glucose tolerance, insulin sensitivity, and reduces adiposity in mice and rats ([11–13]). BCAA intake also regulates the lifespan of mice, with BCAA restriction extending the lifespan of mice and promoting lifelong leanness, and BCAA supplementation shortening lifespan and promoting obesity ([14,15]). Many of the metabolic and anti-aging effects of BCAAs may be mediated by isoleucine, as restriction of isoleucine is sufficient to improve metabolic health, extend lifespan, and rejuvenate aged mouse tissues ([16–19]).

Fluctuations in dietary nutrient consumption have been proposed to regulate physiological responses, in part, via epigenetic mechanisms as central metabolites act as co-substrates for chromatin-modifying enzymes ([20,21]). For example, methionine is essential for the production of S-adenosylmethionine (SAM), which acts as a methyl-donor for methyltransferases. Reduced SAM availability is associated with both global and site-specific decreases in histone methylation post-translational modifications that play critical roles in regulating gene expression and genome stability ([22]). Similarly, BCAAs are catabolized to either acetyl-CoA (leucine), propionyl-CoA (valine), or both (isoleucine), and while having many other biological roles these BCAA-derived catabolites can also be used for the posttranslational modification of histones. Acetyl-CoA is required for histone acetylation, and acetyl-CoA derived from leucine can be utilized by the p300 histone acetyltransferase to acetylate multiple substrates ([23,24]). Isoleucine-derived propionyl-CoA can also be used as a substrate by histone acyltransferases and is required for propionylation of cardiac H3K23 ([25]). The effects of leucine and isoleucine are particularly interesting as recent work has shown that restriction of leucine and isoleucine have dramatically different effects on the metabolic health of mice, with restriction of isoleucine being necessary and sufficient for the metabolic benefits of protein restriction including reduced adipose tissue, increased energy expenditure, and improved glycemic control, while reducing leucine did not ([19]). Further, dietary levels of isoleucine, but not leucine, are associated with body mass index in humans ([19]).

It is thus clear that adaptive mechanisms exist which allow animals to actively respond not just to times
of famine, but to decreased availability of specific essential amino acids. These may occur over different time
periods, including meal to meal variation or simply during the day in response to feeding and circadian rhythm.
Some of these mechanisms could involve the mechanistic Target of Rapamycin Complex 1 (mTORC1), an
evolutionarily conserved sensor of amino acids - including Met, Leu, and Ile - that promotes growth and anabolic
metabolism when amino acids are sufficient ([26]).

Here, we investigate the metabolic effects of restricting methionine, leucine, and isoleucine on the
physiology and the liver metabolome, epigenome, and transcriptome of C57BL/6J male mice. We limited the
treatments to 3 weeks, with the goal of capturing early-stage responses to depletion of EAAs.  We further
examine the potential role of mTORC1 in facilitating these responses by including a treatment group receiving
an inhibitor of mTORC1, rapamycin. By assaying these treatment groups in parallel within a single study, we aim
to quantitatively determine the degree to which dietary EAA restrictions elicit unique and/or shared physiological
and hepatic responses to inform how the specific amino acid profile of dietary proteins plays a key role in
regulating metabolic health at the molecular level.

**Results**

***Essential amino acid restrictions elicit systemic physiologic responses unique from rapamycin***
***treatment***

To investigate how complete dietary restriction of select EAAs subsequently impacts different
physiological measurements, we placed 14-week-old C57BL/6J male mice on one of four AA-defined diets for 3
weeks (**Fig. 1A**). Briefly, we utilized a widely used commercially available AA-defined Control diet in which 15.6%
of calories are derived from AAs. We then generated a series of diets in which either methionine (Met-D), leucine
(Leu-D), or isoleucine (Ile-D) were depleted. All of the diets are isocaloric, with identical levels of fat and
carbohydrate; in each restricted diet, non-EAAs were increased to keep the calories derived from AAs constant
(**Table S1**). In parallel, a subset of Control-fed mice was treated daily with rapamycin in order to assess if
mTORC1 inhibition mimics some or all of the physiological and molecular effects of EAA restriction.

We found that restriction of EAAs, but not rapamycin treatment, led to a steady decrease in total body
weight (**Fig. 1B**). Met-D and Leu-D mice exhibited a similar pattern of lost body weight, weighing 7.5 g and 9.0
g less, respectively, than Control-fed mice following 3 weeks of EAA restriction. Notably, Ile-D mice lost
significantly more body weight than both Met-D and Leu-D mice, weighing on average 12.25 g less than controls
at the study's conclusion. The observed changes in body weight were largely independent of altered food intake
patterns, with food intake being similar or higher in all groups eating an AA-depleted diet from the second week
onwards (**Fig. 1C**). Together, these data suggest 3 weeks of individual EAA restriction is sufficient to induce
whole body weight loss, likely via increased energy expenditure as has been previously reported in similar
systems ([7,19,27,28]).

To assess changes in body weight as alterations in lean and fat mass stores, we analyzed body composition of each mouse at weeks 0 and 3. We found that only control mice retained lean mass over the course of our study. EAA-restricted mice experienced significant losses in lean mass that were greater than those induced by rapamycin treatment (**Fig. 1D**). Losses in lean mass correlated with a significant induction of autophagy in Leu-D mice as determined via decreased quadricep p62 protein expression (**Fig. S1A-B**). Although Leu-D has previously been shown to induce autophagy via mTORC1-inhibition, all mice were harvested following an overnight fast which may have masked autophagy induction relative to controls for Met-D and Ile-D treatment groups ([23]). In addition, EAA restriction induced significant losses in fat mass with a near total loss in detectable fat mass for Ile-D mice (**Fig. 1E**). Loss in fat mass for Leu-D and Ile-D mice was greater in magnitude than the observed losses in lean mass, resulting in an altered body composition consisting of increased lean mass and decreased fat mass percentages (**Figs. 1F-G**). The relative percentages of lean and fat mass were unchanged in Met-D mice, while control and rapamycin treated mice exhibited increased fat mass which is consistent with normal age-associated changes in body composition. Therefore, these data show dietary restriction of individual BCAAs has a greater impact on body composition than Met-D, likely via mTORC1-independent mechanisms.

In addition to altered body composition, dietary EAA restriction has been associated with improved glucose tolerance ([11–13]). Although 3 weeks of complete dietary Met-D was unable to significantly improve glucose tolerance relative to controls, Leu-D and Ile-D mice exhibited improved glucose clearance under glucose tolerance test (GTT) conditions (**Figs. 1H-I**). Unlike EAA restrictions, rapamycin treated mice displayed reduced glucose tolerance as we have previously observed ([29,30]). Collectively, principal component analysis (PCA) of these data show 3 weeks of individual EAA (Met, Leu or Ile) dietary restriction is sufficient to induce whole-organism physiologic changes, with each EAA restriction eliciting distinct phenotypes which were not observed by rapamycin treatment alone (**Figs. 1J**).

### *Specific dietary EAA restriction or rapamycin treatment induces distinct hepatic metabolite profiles*

To determine how specific EAA restrictions influence cellular metabolism, we performed a targeted metabolomics analysis of livers from mice collected following the conclusion of the *in vivo* analysis above, after mice had consumed the indicated diets (or been treated with rapamycin) for 3 weeks (**Table S2**). Hierarchical clustering and principal component analysis (PCA) of treatment groups based on the altered relative abundance of over 100 quantifiable metabolites revealed changes induced by dietary EAA restriction were again more similar to one another than those stimulated by rapamycin treatment (**Figs. 2A-B**). However, unlike the grouping patterns observed in our physiologic measurements, Leu-D mice exhibited a metabolic profile which closely mimicked that of Met-D mice, while Ile-D fed animals were quite distinct from both Met-D and Leu-D-fed, as well as Control-fed animals. This result was largely unexpected as methionine, a non-BCAA, serves in dramatically different metabolic roles and biological pathways than do leucine or isoleucine.

To assess how dietary restriction of EAAs altered specific metabolic pathways in the liver, we assessed the relative changes of key metabolites within and adjacent to glycolysis, the pentose phosphate pathway (PPP), the tricarboxylic acid (TCA) cycle, nucleotide, and redox metabolic networks (**Figs. 2C-F**). For glycolysis, a

trending or significant increase in glucose 6-phosphate (G6P) and dihydroxyacetone phosphate (DHAP) was
observed in all treatment groups although only Ile-D mice displayed an increased abundance of the downstream
metabolite phosphoenolpyruvate (PEP) (**Fig. 2C**). Conversion of PEP to pyruvate appeared uniquely reduced in
Met-D mice as pyruvate and lactate levels were both significantly lower than in control mice. Despite apparent
EAA-specific effects on glycolytic metabolism, all treatment groups exhibited a trending or significant increase in
all measured PPP intermediates with the exception of ribulose 5-phosphate (Ribulose 5P) which was largely
unchanged in response to Met-D but significantly depleted in both Leu-D and Ile-D liver (**Fig. 2C**). Rapamycin
elicited an opposing response in Ribulose 5P availability, with Ribulose 5P being significantly increased followed
by a corresponding decrease in ribose 5-phosphate (Ribose 5P) (**Fig. 2C**). The metabolic changes induced by
Rapamycin are consistent with known role for mTORC1 in positively regulating the translation of ribose-5-
phosphate isomerase A, the enzyme that interconverts ribulose 5-phosphate to ribose 5-phosphate ([31]).

Downstream of glycolysis, all treatments besides Ile-D stimulated a decrease in acetyl-CoA levels relative
to controls (**Fig. 2D**). Similar trends were observed for propionyl- and butyryl-CoA, highlighting a general
decrease in liver short chain acyl-CoA levels for Met-D, Leu-D, and rapamycin treated mice. Despite decreased
acetyl-CoA availability for Met-D, Leu-D, and rapamycin treated mice, all treatment groups exhibited trending or
significant increases in citrate availability (**Fig. 2D**). However, elevated citrate levels were not associated with a
greater abundance of immediate downstream TCA cycle intermediates. Instead, all conditions exhibited similar
or decreased abundance of cis-aconitate, alpha-ketoglutarate (alpha-KG), and/or succinate relative to controls.
2-hydroxyglutarate (2-HG), a metabolite with structural similarities to alpha-KG, was also significantly decreased
in Ile-D mice (**Fig. 2D**). A significant increase in fumarate and malate under Met-D, Leu-D, and Ile-D, but not
rapamycin treatment, mirrored increases in tyrosine and phenylalanine abundance, suggesting the increase in
malate/fumarate is driven by selective amino acid anaplerosis (**Fig. 2D**). Along with tyrosine and phenylalanine,
numerous other amino acids were generally elevated in response to each dietary EAA restriction, but not
rapamycin treatment (**Figs. 2A and 2D**). In fact, the EAA designated for dietary restriction uniformly presented
with elevated hepatic levels. We have previously observed that 5 weeks of Met-D is sufficient to lower liver
methionine levels >50% ([22]), while the current study of dietary EAA restriction (for 3 weeks) shows an increase
in liver methionine, likely reflecting the window of adaptive response where methionine is salvaged from other
tissues.

In addition to central metabolic pathways, other major metabolites showed differential regulation in
response to dietary EAA restrictions. For example, Ile-D led to the accumulation of high energy state adenosine
and guanosine nucleotides (i.e, ATP and GDP) which showed a trending or significant decrease across the other
treatment groups (**Fig. 2E**). Uridine, cytidine, and inosine nucleotides were also elevated in Ile-D mice with no
clear pattern observed in response to other dietary EAA restrictions or rapamycin treatment. Interestingly, the
redox metabolites $NAD^+$/NADH and $NADP^+$/NADPH were significantly reduced across all conditions, including
rapamycin treatment, suggestive of an overall decrease in $NAD^+$ synthesis and/or increase in $NAD^+$ catabolism
(**Fig. 2F**). A decreased ratio of glutathione (GSH) to glutathione disulfide (GSSG) was observed in Met-D and
Leu-D mice with no changes in the relative abundance of either redox metabolite altered in Ile-D or rapamycin
mice. Together, these metabolic analyses show dietary Ile-D has a unique impact on the relative levels of major

metabolic cofactors with dietary Met-D and Leu-D displaying common responses unique from both Ile-D and
rapamycin-treated mice.

***Dietary EAA restrictions stimulate unique Histone PTM patterns across all conditions that are not***
***mimicked by mTORC1 inhibition***

Fluctuations in metabolite availability can regulate cellular processes through numerous, diverse
mechanisms. This includes direct regulation of the epigenome, as the enzymes responsible for adding and
removing epigenetic post-translational modifications (PTMs) rely on metabolic co-substrates to support their
catalytic activity. Notably, all three EAAs investigated in this study are metabolized to such co-substrates. To
determine whether the liver epigenome is uniquely sensitive to dietary EAA restrictions or rapamycin treatment,
we performed LC-MS/MS histone proteomics on liver tissue harvested at the study's completion (**Table S3**).
Hierarchical clustering and PCA analyses based on the altered relative abundance of 93 uniquely modified
histone peptides revealed distinct global histone PTM profiles for all treatment groups (**Figs. 3A-B**).  We
previously investigated the epigenetic responses to Met-D in detail, revealing adaptive chromatin mechanisms
highlighted by preferential H3K9me1 maintenance in the presence of global decreases to higher-state histone
methylations ([22,32]). These trends under Met-D were replicated in the present study (**Fig. 3C-E**).

Unexpectedly, Ile-D and Met-D groups exhibited a highly similar trend in H3K9 methylation states, as
well as increased acetylation on H3K9/K14 and increased propionylation on both H3K9/K14 and H3K18/K23
(**Figs. 3C-F**). All three EAA restrictions led to similar increases in H3 acylation states, but rapamycin treatment
only stimulated a significant increase in H3K9/K14 propionylation. Interestingly, unlike the increase of H3
acylations, Met-D and Ile-D led to significantly decreased levels of acetylation on H4 (**Fig. 3G-J**). The decrease
in H4 acetylation was especially prevalent in Ile-D liver as all acetylated H4 peptide species (i.e., 1ac, 2ac, 3ac,
and 4ac) were reduced under this condition. While Met-D mice displayed decreased liver acetyl-CoA levels, Ile-
D acetyl-CoA levels were unchanged. These data either suggest independent mechanisms dictate H4ac profiles
in response to dietary Met-D/Ile-D, or that H4 acetylation mechanisms are shared but function independent of
whole-cell acetyl-CoA availability. In support of the latter, H4 acetylation levels in Leu-D and rapamycin treated
mice were robust despite decreased liver acetyl-CoA availability (**Fig. 3G-J**). Notably, unlike acetylation levels
on H4, other short chain acylations were largely unchanged across conditions with only Ile-D mice exhibiting
slight increases in butyrylated and crotonylated H4 residues (**Fig. 3K-L**). Together, these data demonstrate that
3-week dietary EAA restrictions or rapamycin treatment have differential effects on the liver epigenome, providing
a potential mechanism driving EAA restriction-specific phenotypes.

***Ile-D induces a robust transcriptional response compared to Met-D, Leu-D, or Rapa treatment***

To investigate how each treatment altered the functional gene expression landscape of the liver, we
performed bulk RNA-sequencing (**Table S4**). PCA of log transformed gene counts per million (CPM) values
revealed identical groupings to those observed from the metabolomics analyses (**Figure 4A**), suggesting a direct

functional link between the changes in transcripts and subsequent metabolic pathways. Also, the similar PCA

groupings of Met-D and Leu-D further suggest that these amino acid deficiencies are sensed through a common

mechanism, distinct from Ile-D. In support of this observation, 45.6% (n = 4088) of all differentially expressed

(DE) genes identified in Ile-D liver were uniquely altered under this condition alone (**Fig. 4B**).   Only 11.7%

(n=590) and 13.2% (n=494) of DE genes in Leu-D and Met-D mice, respectively, were unique to each given

condition (**Fig. 4B**). In fact, a majority of Leu-D or Met-D DE genes were also differentially regulated by at least

one other dietary EAA restriction (i.e., Leu-D = 83%, Met-D = 81.4%). A majority of rapamycin DE genes (77.5%)

were also shared by at least one other EAA restricted condition, although relative to the changes observed in

other conditions, the absolute number of DE genes contributing to this pattern (n=478) was significantly lower

(**Fig. 4B**). The absolute numbers of identified DE genes mirrored the relative magnitude of each response, with

Met-D and Leu-D DE genes eliciting transcriptional responses less significant than Ile-D but greater than

rapamycin treatment (**Figs. 4C-4F**). Together, these data collectively suggest EAA restrictions stimulate distinct

hepatic transcriptional responses from rapamycin treatment.

With EAA transcriptional responses being clearly distinct from those elicited by rapamycin treatment, we

performed Gene Ontology Biological Process (GO-BP) functional annotation of DE genes shared by Met-D, Leu-

D, and Ile-D conditions to identify candidate processes which are broadly responsive to dietary EAA restriction

([33]). This analysis identified 59 significantly enriched GO-BP terms covering cellular functions related to lipid

(e.g., fatty acid and cholesterol metabolic processes) and amino acid (e.g., glutathione and glutamate metabolic

processes) metabolism as well as chromatin regulation (e.g., chromatin remodeling and chromatin looping) and

RNA processing (e.g., rRNA processing and RNA splicing) (**Table S5**).    Notably, the GO-BP term

"transmembrane transport" possessed the second-most significant FDR value and was primarily enriched for

solute carriers (SLCs), including the amino acid transporters SLC25A22 and SLC36A4. These results highlight

how dietary EAA restrictions uniformly stimulate significant responses by nutrient uptake and utilization pathways

in addition to shared aspects of chromatin and RNA regulation.

We next sought to assess the unique functional consequences of each individual EAA restriction or

rapamycin treatment by analyzing condition-specific DE genes via a GO-BP pre-ranked Gene Set Enrichment

Analysis (GSEA) (**Figs. 4G-J, Table S6**) ([34]). Met-D-specific DE genes were assigned to a small number of

significantly enriched gene sets (n=6) which primarily possessed positive NES values and consisted of

membrane organization/cell motility biological processes (**Fig. 4G**). A similar number of significantly enriched

gene sets were identified using Ile-D specific transcripts (n=11) which also predominantly held positive NES

scores. Positive NES gene sets were overwhelmingly associated with diverse intracellular signal transduction

pathways such as the MAPK signaling cascade (**Fig. 4I**). Given the greater magnitude of transcriptional response

identified in Ile-D mice, it was unsurprising to find that Ile-D-specific DE genes were significantly enriched in the

greatest number of GO-BP gene sets (n=74) (**Fig. 4J**). Interestingly, multiple positive NES gene sets were

associated with histone/protein post-translational modifications, providing potential insights into mechanisms

regulating the unique histone PTM profile of Ile-D mice (**Fig. 3A**). Negative NES gene sets were primarily

associated with lipid metabolism and mitochondrial oxidative phosphorylation, suggestive of a compensatory

mechanism aimed towards preserving liver lipid stores in the presence of a near complete loss in total fat mass

(**Fig. 1E**). Lastly, as observed in both metabolomics and histone proteomics datasets, rapamycin-specific
enriched gene sets (n=5) were again distinct from all dietary EAA restrictions and primarily consisted of NES
gene sets related to immunity-associated transcriptional programs (**Fig. 4G**). Therefore, these data collectively
reveal that Ile-D stimulates the largest and most distinct transcriptional response of all treatments while providing
insight into underlying mechanisms supporting this intervention's unique physiologic, metabolic, and epigenetic
phenotypes.

*Multi-Omics Factor Analysis identifies differential histone acetylation and expression of central*
*metabolic genes as defining features of Ile-D*

To generate a quantitative understanding of the molecular responses unique to Ile-D, a Multi-Omics
Factor Analysis (MOFA) was performed (**Extended Data Figure 2**). MOFA is an unsupervised computational
method that co-analyzes datasets to identify a set of Factors which explain the biological and technical variability
between experimental groups ([35]). When applying MOFA to the 14,139 data points collected in this study, 15
distinct factors were identified with a majority of the variability explained by Factors 1 through 3 (**Fig. 5A**). Ile-D
measurements strongly correlated with Factor 1 (r= 0.9) whereas remaining treatment groups possessed either
negligible (Leu-D: r= 0.06, Met-D: r= -0.14) or negative correlations with this Factor (Control: r= -0.4, Rapa: r= -
0.41) (**Fig. 5B**). The percent of total variance contributed by each dataset was greatest in Factor 1, supporting
our observation that Ile-D elicits distinct responses across all measurements taken in this study (**Fig. 5A**).
Additionally, Met-D and Leu-D treatment groups strongly correlated with Factor 2, further suggesting dietary
methionine or leucine restriction may function through similar mechanisms.

Interestingly, the variance contributed by histone PTM measurements was primarily associated with
Factor 1 (10.84%), suggesting differences in histone PTM regulation most strongly distinguish Ile-D from
remaining treatment groups (**Fig. 5A**).  When assessing which histone PTMs assigned to Factor 1 primarily
contributed to this variance, we detected a redistribution in histone acetylation across H3, H4, and H2A isoforms
(**Fig. 5C**). Specifically, we identified hypoacetylation at H4 K5/K8/K12/K16 and H2A1/J/X K5/K9 with
hyperacetylation at H3 K9/K14 and H4 K77ac (**Fig. 5C**). This pattern of acetyl-group redistribution is supported
by the GSEA results generated from Ile-D-specific DE genes which identified the H3K14 acetyltransferase Kat7
as well as the broad histone deacetylases Sirt1 and Hdac4 in the "Histone Modification" gene set (**Fig. 4J**).
Together, these MOFA Factor 1-associated data suggest differential regulation of histone acetylation profiles is
a unique and defining characteristic of Ile-D in the liver.

As the epigenome directly regulates gene expression profiles, we next analyzed Factor 1 Principal
Component Gene Set Enrichment (PCGSE) results to identify which pathways are transcriptionally associated
with this redistribution in histone acetylation. PCGSE of Factor 1 genes identified an upregulation of pathways
linked with RNA metabolism and processing as well as chromatin organization and modification (**Fig. 5D**).
Notably, "Chromatin Modification," "Chromatin Organization," and "Covalent Chromatin Modification" gene sets
included Kat7, Sirt1, and Hdac4, further implicating these chromatin modifiers as mediators of the epigenetic
response to Ile-D. Downregulated Factor 1 gene sets were primarily enriched for organic acid metabolism and

lipid oxidation, two pathways critical for mitochondrial acetyl-CoA production (**Fig. 5E**). The robust transcriptional
downregulation of mitochondrial acetyl-CoA producing pathways were striking as hepatic acetyl-CoA levels were
maintained during Ile-D, suggesting alternative pathways may be engaged to support acetyl-CoA maintenance
under this condition. Together, these MOFA PCGSE results highlight the association of chromatin regulation and
mitochondrial acetyl-CoA metabolism with histone acetylation redistribution in Ile-D liver.
330
331  ***Ile-D induced metabolic and epigenetic remodeling occur rapidly within isolated cells***
332
Although distinct acetyl-CoA metabolism and histone acetylation states are unique features of Ile-D within
the liver, it remains unclear whether these mechanisms require complex inter-cellular/organ signaling or
represent inherent, fundamental intracellular responses. To determine whether isolated cells possess the
inherent capability of initiating these molecular responses, we investigated the metabolic and epigenetic
responses of human liver HepG2 cells over 24 hours of Ile-D (**Fig. 6A, Tables S7 and S8**). *In vitro*, rapid removal
of Ile from culture media led to a dramatic loss in its downstream catabolic product KMVA, which was no longer
detectable after 10 minutes (**Figs. 6B-C**). Importantly, this perturbation had minimal effects on global BCAA
metabolism as leucine, valine, and their metabolic products KICA and KIVA, respectively, exhibited minor
increases in abundance (**Figs. 6B-C**). Similarly to the *in vivo* metabolomics data, Ile-D HepG2 cells were largely
capable of maintaining cellular acetyl-CoA pools (**Fig. 6D**). This maintenance of acetyl-CoA was particularly
striking as propionyl-CoA and succinyl-CoA, downstream acyl-CoA products of isoleucine metabolism, were
rapidly depleted and maintained at low levels over 24 hours of Ile-D (**Fig. 6D**).
*In vivo*, maintained acetyl-CoA levels were accompanied by selective amino acid anaplerosis which
restored TCA cycle intermediate abundance from fumarate to citrate in contrast to the decreased abundance of
intermediates from cis-aconitate to succinate (**Fig. 2D**). *In vitro,* Ile-D induced similar changes in TCA cycle
intermediate abundance, with alpha-KG and succinate both significantly decreased over the initial 30 minutes
(**Fig. 6E**). Fumarate and malate were generally maintained during this period although the abundance of both
metabolites decreased slightly (**Fig. 6E**). Notably, the amino acids aspartate, asparagine, and phenylalanine all
increased in abundance over the initial 30 minutes of Ile-D, which again implicates selective amino acid
anaplerosis in supporting fumarate and malate levels (**Fig. 6F**). However, the relative abundances of all TCA
cycle intermediates recovered to or above pre-restriction levels by 24 hours, coinciding with maintained elevation
of anaplerotic amino acids, revealing that isolated HepG2 cells are capable of restoring TCA cycle metabolism
under prolonged Ile-D *in vitro* (**Fig. 6E-F**). These data indicate TCA cycle metabolism and selective anaplerosis
are similarly responsive to Ile-D both *in vitro* and *in vivo*, with the immediate *in vitro* response (*i.e.*, 10-30 min)
being most similar to 3-week Ile-D mouse liver.
In addition to possessing similar metabolic responses to *in vivo* Ile-D, HepG2 cells also exhibited a similar
reorganization of histone acetylation (**Fig 5A**). Specifically, Ile-D HepG2 cells displayed hypoacetylation at H4
K5/K8/K12/K16 and H2A/X K5/K9 and maintained levels of H3 K9/K14ac (**Fig. 6G**). Changes in histone PTM
abundance were established during the first 10 minutes of Ile-D and held largely constant over the entire 24-
hour time course (**Fig. 6G**). Together, these data illustrate that Ile-D stimulated remodeling of acetyl-CoA and

TCA cycle metabolism, as well as unique regulation of histone acetylation, are inherent, fundamental responses
of isolated cells.

**Discussion**

In this study, we set out to gain new insight into the molecular mechanisms by which dietary EAA
availability regulates physiologic and hepatic responses at the molecular level. We found that individual depletion
of EAAs led to both common and diet-specific changes. Consistent with prior work, individual depletion of Ile,
Leu, or Met led to improved glucose tolerance with losses in fat and lean mass ([12,22,36,37]). Here, EAA restriction-
induced changes in glucose tolerance and body composition were accompanied by dynamic alterations to the
hepatic metabolome, epigenome, and transcriptome. Metabolically, increases in hepatic glycolytic and PPP
intermediates were prominent across EAA restrictions. It has been estimated that as much as ⅓ of incoming
glucose can be consumed by the PPP ([38,39]), suggesting increased glucose flux through the PPP pathway could
promote greater glucose clearance. An increased abundance of free amino acids in the liver, even among the
amino acids being restricted, was identified across all restrictions. An increase in hepatic amino acid levels,
which was accompanied by a progressive decline in lean mass, is consistent with skeletal muscle loss
(autophagy) and the freeing up of AAs for essential protein synthesis and other metabolic processes throughout
the body. Together, these data highlight the trend of liver self-preservation in response to EAA restrictions at the
expense of skeletal muscle and fat stores.

While a subset of physiologic and hepatic molecular responses was shared across EAA restrictions,
unique response patterns were identified. For example, we were surprised to find that leucine and methionine
restriction stimulated highly similar effects in liver metabolite and gene expression programs. Multi-omics factor
analysis (MOFA) objectively identified the similarity between Leu-D and Met-D molecular responses, uniquely
correlating both treatment groups with Factor 2, a Factor with high variance contributions from hepatic metabolite
and transcript measurements (**Fig. 5A-B**). Factor 2 metabolite trends included a general accumulation of amino
acids, which corresponded with the apparent downregulation of amino acid catabolism genes. Decreased acetyl-
CoA abundance was also associated with Factor 2, highlighting the disparate mechanisms by which Met-D/Leu-
D and Ile-D regulate acetyl-CoA metabolism.

Unique from Met-D and Leu-D, Ile-D consistently elicited distinct responses across all physiological and
molecular measurements taken. This distinct response was confirmed via MOFA, which uniquely correlated Ile-
D with Factor 1-associated features. PCGSE of Factor 1 genes identified an upregulation of chromatin regulation
as well as the downregulation of organic acid metabolism and lipid oxidation pathways. This latter difference
between Ile-D animals and Met-D/Leu-D animals was unexpected as the Ile-D liver was uniquely able to maintain
acetyl-CoA levels. These data suggest acetyl-CoA is likely maintained through alternative pathways under Ile-
D. Our metabolomics analysis implicates cytoplasmic acetyl-CoA synthesis via ACLY as one candidate pathway.
Citrate, ACLY's obligatory substrate, shows the greatest magnitude of accumulation in Ile-D liver. Additionally,
our metabolomics data suggest there is a significant increase in TCA cycle anaplerosis via the conversion of
phenylalanine and tyrosine to fumarate, which would provide additional carbons for acetyl-CoA and citrate

synthesis. The enzyme responsible for catalyzing the first step of tyrosine catabolism, TAT, is upregulated in Ile-

D liver (**Table S5**). Importantly, tyrosine and phenylalanine breakdown through the TAT pathway generates

acetoacetate, which can be further metabolized to two molecules of acetyl-CoA independent of ACLY activity

([40]). Very recent work has shown that acetoacetate-derived acetyl-CoA is capable of supporting *de novo*

lipogenesis upon inhibition of the mitochondrial citrate shuttle, highlighting the flexibility of cells to generate

cellular acetyl-CoA through alternative pathways when necessary ([41]). Further work will be needed to

demonstrate that carbon flux through Phe/Tyr fuels acetyl-CoA synthesis when adipose tissue is depleted, as in

Ile-D.

Although acetyl-CoA levels were maintained, the significant redistribution of histone acetylation was

identified as a discriminating feature of Ile-D liver. This included the global hypoacetylation of H2A and H4 with

a corresponding increase in H3K9/K14 acetylation. The opposing trends in H4 and H3 acetylation are consistent

with a previous report in which Hat1 acetylated, nascently translated H4 proteins are subsequently deacetylated

following their transport to the nucleus. The liberated acetyl-groups were then utilized to provide a local co-

substrate supply and support targeted H3K9 acetylation ([42]). Here, consistent with this mechanism, Hat1 and the

histone deacetylases Hdac4 and SIRT1 are specifically upregulated in response to Ile-D. It is interesting to

speculate that more efficient utilization of acetyl-CoA in response to Ile-D functions in cooperation with selective

anaplerosis, acetoacetate catabolism, and/or ACLY-dependent acetyl-CoA production to maintain global

metabolite pools under this condition. If true, such a cooperative mechanism would be required to maintain global

acetyl-CoA availability in the presence downregulated amino acid catabolism, fatty acid metabolism, and organic

acid metabolism as acetyl groups stored on histone proteins are not present in adequate amounts to support

total cellular requirements ([43]). However, H4 deacetylation has been shown to liberate acetyl groups for utilization

outside the nucleus when needed ([44]). Notably, Ile-D HepG2 cells were also largely capable of maintaining acetyl-

CoA levels. This corresponded with a remodeling of amino acid and TCA cycle metabolism as well as differential

acetyl-CoA utilization by histone acetyltransferases, closely mimicking our *in vivo* observations. Therefore, these

data suggest the ability to cooperatively remodel acetyl-CoA metabolism and histone acetylation profiles is an

inherent capability of isolated cells under Ile-D.

In this study, we also assessed the ability of mTORC1 inhibition via rapamycin to replicate the physiologic

and molecular responses to EAA restrictions. Overall, we found minimal overlap among our hepatic

metabolomic, epigenetic, and transcriptomic analyses. In fact, rapamycin treatment elicited relatively muted

responses compared to dietary EAA restrictions across all physiologic and molecular responses assayed in this

study. Together, these observations imply that downregulation of the mTORC1 signaling pathway in response

to select EAA restrictions plays a minor role in facilitating whole organism and hepatic molecular responses.

However, a potential limitation of this conclusion is that there are rapamycin-resistant functions of mTORC1 that

could be engaged by EAA restriction ([45]), making additional work with genetically engineered mouse models of

altered mTORC1 signaling needed to confirm that EAAs largely function through mTORC1-independent

pathways. Our study is also limited by its focused analysis of male C57BL/6J mice, as rapamycin and dietary

EAA restrictions are known to elicit sex-specific phenotypes ([46,47]). Similarly, although all molecular

measurements in this study focused on hepatic responses, the inherent ability of isolated cells to mount a

conserved response to Ile-D was limited to the analysis of a singular, non-primary liver cell line. Future studies

are also needed to assess how the molecular responses to individual EAA restrictions identified here can be

stimulated by low, but still physiologic levels of EAA restriction and to determine the time-dependence of which

these diets influence physiologic and molecular phenotypes.

In conclusion, we have shown that dietary restriction of single EAAs, while having similar effects on

glucose homeostasis, body composition, and general amino acid catabolism, can yield individual responses by

the hepatic metabolome, epigenome, and transcriptome which are independent of mTORC1 activity. Notably,

Ile-D elicited the most significant and unique response, defined by distinct anaplerosis, decreased BCAA and

fatty acid catabolism, increased pentose phosphate pathway and Phe/Tyr catabolism, as well as histone H4/H2A

hypoacetylation and histone H3K9/K14 hyperacetylation. Together, these results highlight the potential

importance of dietary protein EAA composition in regulating metabolic health at the molecular level.

**Methods**

***Animals and Diets***

All procedures were performed in conformance with institutional guidelines and were approved by the

Institutional Animal Care and Use Committee of the William S. Middleton Memorial Veterans Hospital (Madison,

WI, USA). Male C57BL/6J (#000664) mice were obtained from The Jackson Laboratory. All mice were housed

three per cage, maintained at a temperature of approximately 22°C, and health checks were completed on all

mice daily. Studies were commenced when the mice were 14 weeks of age.

Mice were then randomized to receive the indicated diets from Inotiv either the Control (TD.01084), Met-

D (TD.140119), Leu-D (TD.180789) or Ile-D (TD.190879) diets, or the Control diet plus rapamycin (delivered I.P.

in 5% Tween 20, 5% PEG40, 0.9% NaCl, 3% ethanol at 4 mg/kg/day). Full diet descriptions are provided in

**Table S1**. The randomization of mice was performed at the cage level to ensure that all groups had

approximately the same initial starting weight and body composition. Mice were housed in a SPF mouse facility

in static microisolator cages, except when temporarily housed in a Columbus Instruments Oxymax/CLAMS-HC

metabolic chamber system. Mice were housed under a 12:12 h light/dark cycle with free access to food and

water, except where noted in the procedures below.

***In vitro Cell Culture***

Human HepG2 cell lines were initially cultured in EMEM media (Corning, 10-009-CV) containing all amino acids

and 10% FBS at 37°C, 5% $CO_2$, while EMEM media devoid of amino acids (US Biological, M3859-01) was

supplemented with all amino acids except isoleucine to generate isoleucine depleted media for the appropriate

studies. All non-isoleucine amino acids were added according to their replete EMEM concentrations. Cells were

seeded at a seeding density of 0.4 x$10^6$ and 3 x$10^6$ in 6-well plates and 100mm culture dishes for LC-MS

metabolite analysis and histone proteomics respectively. Prior to Ile depletion, complete EMEM media was

replaced with Ile-Replete media, supplemented with 10% dialyzed FBS (Gibco, A3382001) for one hour. This

was followed by rinsing the cells with PBS, pH 7.4, and subsequent addition of Ile-Deplete media (with 10%

dialyzed FBS) for restriction durations of 10 min, 30 min, 1.5 hr, 5 hr, and 24 hr. At the end of each restriction

time point, cells were harvested for metabolites and core histones extraction accordingly.

### *In vivo Procedures*

Glucose tolerance tests were performed by fasting the mice overnight for 16 hours and then injecting glucose (1

g kg$^{-1}$) ([7,48]). Glucose measurements were taken using a Bayer Contour blood glucose meter (Bayer,

Leverkusen, Germany) and test strips. Mouse body composition was determined using an EchoMRI Body

Composition Analyzer (EchoMRI, Houston, TX, USA). For assay of multiple metabolic parameters [$O_2$, $CO_2$, food

consumption, respiratory exchange ratio (RER), energy expenditure] and activity tracking, mice were acclimated

to housing in a Oxymax/CLAMS-HC metabolic chamber system (Columbus Instruments) for ~24 h and data

from a continuous 24 h period was then recorded and analyzed. Food consumption in home cages was

measured by moving mice to clean cages, filling the hopper with a measured quantity of fresh diet in the morning

and measuring the remainder in the morning 3 days later. The amount was adjusted for the number of mice per

cage, the number of days that passed and the relative weights of the mice (i.e., heavier mice were credited with

a larger relative portion of the food intake). Mice were euthanized by cervical dislocation after an overnight (16h)

fast and tissues for molecular analysis were flash-frozen in liquid nitrogen and stored at -80$^{\circ}$C.

### *Immunoblotting*

Tissue samples from muscle (quadriceps, lateral, on femur) were lysed in cold RIPA buffer supplemented with

phosphatase inhibitor and protease inhibitor cocktail tablets (Thermo Fisher Scientific, Waltham, MA, USA) using

a FastPrep 24 (M.P. Biomedicals, Santa Ana, CA, USA) with bead-beating tubes (16466–042) from (VWR,

Radnor, PA, USA) and zirconium ceramic oxide bulk beads (15340159) from (Thermo Fisher Scientific, Waltham,

MA, USA). Protein lysates were then centrifuged at 13,300 rpm for 10 min and the supernatant was collected.

Protein concentration was determined by Bradford (Pierce Biotechnology, Waltham, MA, USA). 24 µg protein

was separated by SDS– PAGE (sodium dodecyl sulfate–polyacrylamide gel electrophoresis) on 10% resolving

gels (ThermoFisher Scientific, Waltham, MA, USA) and transferred to PVDF membrane (EMD Millipore,

Burlington, MA, USA). Imaging was performed using the BioRad Chemidoc Imaging System (primary antibodies:

SQSTM1/p62 (1:1000; American Research Products, #03-GP62-C; host: Rabbit) and GAPDH (1:1000; Cell

Signaling Technology, #5174; host: Guinea Pig). secondary antibodies: Anti-Guinea pig (1:2000; American

Research Products, #90001; host: Goat) and Anti-Rabbit (1:2000; Cell Signaling Technology, #7074; host: Goat).

Quantification was performed by densitometry using NIH ImageJ software.

### *Metabolite Extraction*

For *in vivo* samples, approximately 30 mg of flash-frozen liver tissue was pulverized in 1 mL of -80°C 80:20

MeOH:$H_2O$ extraction solvent using a motorized tissue grinder and stored on dry ice for 5 minutes. Samples

were then centrifuged at 21,100xg at 4°C for 5 minutes. Supernatants were transferred to individual 15 mL

conical tubes after which 0.8 ml of -20°C 40:40:20 ACN:MeOH:$H_2O$ extraction solvent was used to resuspend

the pelleted samples using a motorized tissue grinder. Resuspended samples were stored on wet ice for 5

minutes. Samples were then centrifuged at max speed for 5 minutes at 4°C and supernatants were combined

with those collected after the initial extraction. The 40:40:20 ACN:MeOH:$H_2O$ extraction was then repeated as

previously described. Combined metabolite extracts were centrifuged at max speed for 5 minutes at 4°C to pellet

any potential insoluble debris after which supernatants were transferred to new individual 15 mL conical tubes.

Finally, extracts were dried using a Thermo Fisher Savant ISS110 SpeedVac and resuspended in 150 µL $H_2O$

per 5 mg of the original liver input. Resuspended extracts were centrifuged at max speed for 5 minutes at 4°C

after which supernatants were transferred to glass vials for LC-MS analysis.

For tissue culture samples, HepG2 cells were rapidly washed with 2x with 1 ml of ice-cold PBS pH 7.4

and incubated in 1 ml of -80°C 80:20 MeOH:$H_2O$ extraction solvent for 15 min at -80°C. Cells were scraped and

transferred to a 1.5 ml eppendorf tube, centrifuged at 21,100xg for 5 min at 4°C, after which the supernatant was

transferred to a new 2 ml eppendorf tube and stored on ice. The remaining pellet was subjected to an additional

extraction by vortexing the sample with 0.8 ml of -20°C 40:40:20 ACN:MeOH:$H_2O$ extraction solvent. This sample

was centrifuged at 21,100xg for 5 min at 4°C after which the supernatant was pooled with the 80:20 MeOH:$H_2O$

extraction supernatant. The remaining cell pellet was used for RIPA protein extraction and BCA protein

quantification to generate normalization factors for LC-MS metabolomics data analyses. Pooled supernatants

were dried using a Thermo Fisher Savant ISS110 SpeedVac and resuspended in 100 µL $H_2O$. Resuspended

extracts were centrifuged at max speed for 5 minutes at 4°C after which supernatants were transferred to glass

vials for LC-MS analysis.

***LC-MS Metabolomics***

Prepared metabolite samples were injected in random order onto a Thermo Fisher Scientific Vanquish UHPLC

with a Waters Acquity UPLC BEH C18 column (1.7 µm, 2.1x100mm; Waters Corp., Milford, MA, USA) and

analyzed using a Thermo Fisher Q Exactive Orbitrap mass spectrometer in negative  ionization  mode.  LC

separation was performed over a 25 minute method with a 14.5 minute linear gradient of mobile phase

(buffer A, 97% water with 3% methanol, 10mM tributylamine, and acetic acid-adjusted pH of 8.3) and organic

phase (buffer B, 100% methanol) (0 minute, 5% B; 2.5 minute, 5% B; 17 minute, 95% B; 19.5 minute, 5% B; 20

minute, 5% B; 25 minute,  5%  B,  flow  rate  0.2mL/min).  A quantity of 10 µ of each sample was injected into

the system for analysis. The ESI settings were 30/10/1 for sheath/aux/sweep gas flow rates, 2.50kV for spray

voltage, 50 for S-lens RF level, 350C for capillary temperature, and 300C for auxiliary gas heater temperature.

MS1 scans were operated at resolution = 70,000, scan range = 85-1250m/z, automatic gain control target = 1 x

$10^6$, and 100 ms maximum IT. Raw data files were converted into mzml for metabolite identification and peak

AreaTop quantification using El-MAVEN (v0.12.1-beta) ([49]). Peak AreaTop values were imported into

MetaboAnalyst for statistical analysis (one factor) using default settings ([50]).

### *Transcriptomic Analysis*

RNA was extracted from liver with TRI Reagent (Thermo Fisher, AM9738), then treated with DNase (Invitrogen, 12185010) and purified with RNA purification kit (Ambion, 12183025). The concentration and purity of RNA was determined using a NanoDrop 2000c spectrophotometer (Thermo Fisher Scientific, Waltham, MA) and RNA was diluted to 100-400 ng/µl for sequencing. The RNA was then submitted to the University of Wisconsin-Madison Biotechnology Center Gene Expression Center & DNA Sequencing Facility, and RNA quality was assayed using an Agilent RNA NanoChip. RNA libraries were prepared using the TruSeq Stranded Total RNA Sample Preparation protocol (Illumina, San Diego, CA) with 250ng of mRNA, and cleanup was done using RNA Clean beads (lot #17225200). Reads were aligned to the mouse (*Mus musculus*) with genome-build GRCm38.p5 of accession NCBI:GCA_000001635.7[DL9]  and expected counts were generated with ensembl gene IDs ([51]).

Analysis of significantly differentially expressed genes (DEGs) was completed in R version 4.3.0. (Team, 2017) using *edgeR* ([52]) and *limma* ([53]). Gene names were converted to gene symbol and Entrez ID formats using the *mygene* package. Genes with too many missing values were removed, if genes were present in less than one diet/age group they were removed. To reduce the impact of external factors not of biological interest that may affect expression, data was normalized to ensure the expression distributions of each sample are within a similar range. We normalized using the trimmed mean of M-values (TMM), which scales to library size. Heteroscedasticity was accounted for using the voom function, DEGs were identified using an empirical Bayes moderated linear model, and log coefficients and Benjamini-Hochberg (BH) adjusted p-values were generated for each comparison of interest ([54]) (Benjamini and Hochberg, 1995). DEGs specific to each condition were rank sorted by log2 fold-change and subsequently used in GSEA Preranked analyses with default settings for gene ontology biological process (GO-BP) pathway enrichment ([34]).

### *Histone Extraction and Chemical Derivatization*

For *in vivo* samples, approximately 30 mg of liver tissue was resuspended in 800 µl of ice-cold Buffer A (10 mM Tris-HCl pH 7.4, 10 mM NaCl, and 3 mM MgCl2) supplied with protease and histone deacetylase inhibitors (10 µg/ml leupeptin, 10 µg/ml aprotinin, 100 µM phenylmethylsulfonyl fluoride, 10 mM nicotinamide, 1 mM sodium-butyrate, and 4 µM trichostatin A) followed by 20 strokes of loose- and 20 strokes of tight-pestle homogenization in a 1 mL Wheaton dounce homogenizer and strained through a 100 µM filter before being transferred to a new 1.5 mL Eppendorf tube. Samples were then centrifuged at 800xg for 10 minutes at 4°C to pellet nuclei. Supernatants were transferred to fresh 1.5 mL Eppendorf tubes or discarded. The remaining nuclei pellet was resuspended in 500 µl ice-cold PBS pH 7.4 and centrifuged at 800xg for 10 minutes at 4°C. The supernatant was discarded and nuclei were again washed with 500 µl ice-cold PBS pH 7.4. Next, pelleted nuclei were resuspended in 500 µl of 0.4N $H_2SO_4$ and rotated at 4°C for 4 hours. Samples were centrifuged at 3,400xg for 5 minutes at 4°C to pellet nuclear debris and precipitated non-histone proteins. Supernatants were transferred to new 1.5 mL Eppendorf tubes after which 125 µl of 100% trichloroacetic acid was added and incubated overnight on ice at 4°C. The next day, samples were centrifuged at 3,400xg for 5 minutes at 4°C to pellet precipitated histone proteins. Supernatants were discarded and precipitant was washed with 1 mL ice-cold acetone +0.1%

HCl. Samples were centrifuged at 3,400xg for 2 minutes at 4°C and supernatants were again discarded. This

process was repeated with a 100% ice-cold acetone wash. Residual acetone was allowed to evaporate at room

temperature for 10 minutes after which dried precipitant was resuspended in $H_2O$. Finally, samples were

centrifuged at 21,100xg for 5 minutes at 4°C to pellet any remaining insoluble debris and supernatants containing

purified histone were transferred to new 1.5 mL Eppendorf tubes. Histone extractions from flash frozen HepG2

cell pellets were performed as described above.

To prepare purified histone samples for LC-MS/MS analysis, 5 µg of each sample were diluted with $H_2O$

to a final volume of 9 µl. 1 µl of 1 M triethylammonium bicarbonate was added to each sample to buffer the

solution to a final pH of 7-9. Next, 1 µl of 1:100 d6-acetic anhydride:H2O was added to each sample followed by

a 2-minute room temperature incubation. The reaction was quenched via addition of 1 µl 80 mM hydroxylamine

followed by a 20-minute room temperature incubation. Next, d3-acetylated histones were digested with 0.1 µg

trypsin for 4 hours at 37°C. Upon completion of trypsin digestion, 5 µl of 0.02 M NaOH was added to adjust the

final pH to ~9-10. Prepared histone peptides were then N-terminally modified with 1 µl 1:50 phenyl

isocyanate:ACN for 1-hour at 37°C. Modified peptides were desalted and eluted-off of EmporeC18 extraction

membrane. Eluted samples were dried completely using a Thermo Fisher Scientific Savant ISS110 SpeedVac

and resuspended in 40 µl sample diluent (94.9% H2O, 5% ACN, 0.1% TFA). Resuspended samples were

centrifuged at max speed for 5 min at 4°C after which supernatants were transferred to glass vials for LC-MS/MS

analysis.

**_LC-MS/MS Histone Proteomics_**

Derivatized histone peptides were injected onto a Dionex Ultimate3000 nanoflow HPLC with a Waters nanoEase

UPLC C18 column (100 m x 150 mm, 3 µm) coupled to a Thermo Fisher Q-Exactive mass spectrometer at 700

nL/ min. Aqueous phase (A) consisted of H2O + 0.1% formic acid while the mobile phase (B) consisted of

acetonitrile + 0.1% formic acid (B). Histone peptides were resolved using the following linear gradient: 0 min,

2.0% B; 5 min, 2.0% B; 65 min, 35% B; 67 min, 95% B; 77 min, 95% B; 79 min, 2.0% B; 89 min, 2.0% B. Data

was acquired using data-independent acquisition (DIA) mode. The mass spectrometer was operated with a MS1

scan at resolution = 35,000, automatic gain control target = 1 × 106, and scan range = 390-910 m/z, followed by

a DIA scan with a loop count of 10. DIA settings were as follows: window size = 10 m/z, resolution = 17,500,

automatic gain control target = 1 × 106, DIA maximum fill time = AUTO, and normalized collision energy = 30.

DIA Thermo .raw files were analyzed via the EpiProfile 2.0 AcD3 module ([55]). Subsequent data filtering

(i.e., removing samples with (1) >2 null values for common peptides or (2) >50% CV) and normalization was

performed using our published Histone Analysis Workflow ([56]).

**_Multi-Omics Factor Analysis_**

To integrate transcriptomics, metabolomics, histone proteomics, and physiological data, Multi- Omics Factor

Analysis (MOFA) ([35]) was performed on 60 mice consisting of 13929 transcripts, 103 metabolites, 92 histone

proteins, and 15 phenotypes. The model was trained on 15 factors (minimum explained variance 0.8% in at least

one data type) with default parameters except for likelihoods = "gaussian", maxiter = 1000, convergence_mode

= "slow", and seed = 42. The variance decompositions between factors and datasets were computed. Pearson

correlation coefficients (Freedman et al., 2007) were calculated to show the association factors and five

experimental conditions (control, isoleucine restriction, leucine restriction, methionine restriction, and

rapamycin). To run an enrichment analysis, principal component gene set enrichment (PCGSE) (Frost et al.,

2015) was performed on MSigDB_v6.0_C5_mouse ([57]) gene set consisting of genes annotated by GO Biological

Process terms. All analyses were conducted using R (version 4.2.1) and MOFA2 (version 1.8.0).

*Statistics*

Most statistical analyses were conducted using Prism, version 9 (GraphPad Software Inc., San Diego, CA, USA),

R (version 4.1.0), or MatLab (version R2023b). Tests involving multiple factors were or by one-way ANOVA with

Diet as the categorical variable followed by a Tukey–Kramer or Dunnett's *post hoc* test for multiple comparisons

as appropriate. Data distribution was assumed to be normal but was not formally tested.

**Acknowledgements**

The Lamming lab is supported in part by the NIA (AG056771, AG081482, AG084156, and AG085898), the

NIDDK (DK125859), and startup funds from UW-Madison. The Denu lab is supported in part by the NIH

(GM149279 and DK125859).  The Ong lab is supported in part by the NIH (U19AI104317 and DK125859), and

startup funds from UW-Madison. This research was supported by The New York Stem Cell Foundation, SAH is

a NYSCF - Druckenmiller Fellow. CLG was supported in part by Dalio Philanthropies, a Glenn Foundation

Postdoctoral Fellowship, and by Hevolution Foundation award HF-AGE AGE-009. RB is supported by

F31AG081115. The authors used the UW-Madison Biotechnology Center Gene Expression Center

(RRID:SCR_017757) which is supported in part by the UWCCC (P30CA014520). The CLAMS-HC was

purchased with funds from the U.S. Department of Veterans Affairs (IS1-BX005524), and this work was

supported using facilities and resources from the William S. Middleton Memorial Veterans Hospital. The content

is solely the responsibility of the authors and does not necessarily represent the official views of the NIH. This

work does not represent the views of the Department of Veterans Affairs or the United States Government.

**Conflict of interests**

SAH is a consultant for Galilei Biosciences. JMD is a co-founder of Galilei Biosciences and a consultant for

Evrys Bio. DWL has received funding from, and is a scientific advisory board member of, Aeovian

Pharmaceuticals, which seeks to develop novel, selective mTOR inhibitors for the treatment of various diseases.

**Figure Legends**

**Figure 1: *Physiologic responses to EAA restrictions and rapamycin treatment.*** (*A*) Visual representation

of study design. (*B*) Weekly body weight measurements. (*C*) Bar graph illustrating weekly food consumption

normalized to mouse body weight values. (*D-E*) Bar graphs depicting absolute changes in lean and fat mass
measured via whole-body MRI following three weeks of dietary intervention or rapamycin treatment. (*F-G*) Bar
graphs depicting the body composition percentages of lean and fat following three weeks of dietary intervention
or rapamycin treatment. (*H-I*) Glucose tolerance test curves and calculated area under the curve (AUC) values.
Error bars = SEM; N = 12; * = p-value < 0.05 as measured via Student's t-test.

**Figure 2:** *Alterations in hepatic metabolism following EAA restrictions and rapamycin treatment.* (*A*)
Heatmap with hierarchical clustering of metabolite abundance values. (*B*) PCA analysis of log transformed,
relative metabolite abundance values. (*C-F*) Diagram depicting log2 fold-changes in metabolite abundance
relative to controls within the larger context of their associated metabolic pathways. N = 12 for control, rapamycin,
Met-D, and Ile-D groups; N = 11 for Leu-D; p-values measured using default MetaboAnalyst settings.

**Figure 3:** *Hepatic histone PTM responses to EAA restrictions and rapamycin treatment.* (*A*) Heatmap with
hierarchical clustering of histone peptide stoichiometric abundance values compared to control values. (*B*) PCA
analysis of log transformed, relative histone peptide stoichiometry values. (*C-L*) Box plots depicting deconvoluted
stoichiometry percentages of residue-specific histone PTMs. Error bars = standard deviation; N = 12 for control,
N = 10 for Rapa; N = 7 for Leu-D; N = 6 for Met-D and Ile-D; * = p-value < 0.05 as measured via Student's t-test.

**Figure 4:** *Hepatic transcriptional programs elicited by EAA restrictions and rapamycin treatment.* (*A*) PCA
analysis of log transformed gene CPM values. (*B*) Venn diagram of differentially expressed genes with p<0.05.
(*C-F*) Volcano plots depicting log2 FC and -log10 adj. p-values for all identified genes. Horizontal marker
indicates threshold for statistical significance (p-adj. < 0.05) with significantly changing genes marked in red. (G-
J) Dot plots illustrating enriched Gene Ontology – Biological Process (GO-BP) gene sets as determined via Gene
Set Enrichment Analyses (GSEA) of differentially expressed genes unique to each given condition. Example
GSEA plots highlighting significantly enriched GO-BP gene groups with positive and negative normalized
enrichment scores (NES) are present beneath each corresponding dot plot. N = 12.

**Figure 5:** *Multi-Omics Factor Analysis of whole organism and hepatic molecular measurements.* (*A*)
Heatmap of variance decomposition by factor. (*B*) Plot illustrating Pearson correlation coefficients for control and
treatment groups with 15 MOFA Factors. (*C*) Heatmap with hierarchical clustering of Factor 1-associated histone
peptide stoichiometric abundance values compared to control values. Factor 1-associated histone peptides were
determined using a weight cutoff value of +/- 0.2 which identified the top 18 associated histone peptides. (*D-E*)
Plots depicting a subset of significantly enriched gene sets identified using Factor 1 transcripts. All significantly
enriched gene sets can be found in Extended Data Figure 2. * = p-value < 0.05 as measured via Student's t-test.

**Figure 6:** *Metabolic and epigenetic mechanisms elicited by Ile-D of isolated cells.* (*A*) Schematic depicting
experiment design for *in vitro* Ile-D of HepG2 hepatocellular carcinoma cells. (*B-C*) Plots depicting changes in
metabolite abundance relative to 0 hr control values. (*E-F*) Heatmaps of metabolite abundance values. (*G*)

Heatmap with hierarchical clustering of histone peptide stoichiometric abundance values compared to 0 hr control values. Depicted peptides of interest were originally identified via MOFA in Fig 5. Histone peptides with similar changes in abundance *in vitro* as previously identified *in vivo* are underlined. Error bars = STDEV; N = 4 for metabolite measurements; N = 3 for histone PTM measurements; * = p-value < 0.05 as measured via Student's t-test.

**Extended Data Figure 1:** *Differential p62 expression following dietary EAA depletions.* (*A*) Bar graph depicting muscle (quadricep) p62 protein levels normalized to GAPDH. (*B*) Representative western blot image of those used for the quantification depicted in panel A. Error bars = SEM; N = 12; * = p-value < 0.05 as measured via Student's t-test.

**Extended Data Figure 2:** *Interactive Multi-Omics Factor Analysis Results.* HTML file linking to interactive MOFA results.

**References**

1. Green, C. L., Lamming, D. W. & Fontana, L. Molecular mechanisms of dietary restriction promoting health and longevity. *Nat. Rev. Mol. Cell Biol.* **23**, 56–73 (2022).

2. MacArthur, M. R. *et al.* Total protein, not amino acid composition, differs in plant-based versus omnivorous dietary patterns and determines metabolic health effects in mice. *Cell Metab.* **33**, 1808-1819.e2 (2021).

3. McCarty, M. F., Barroso-Aranda, J. & Contreras, F. The low-methionine content of vegan diets may make methionine restriction feasible as a life extension strategy. *Med. Hypotheses* **72**, 125–128 (2009).

4. Schmidt, J. A. *et al.* Plasma concentrations and intakes of amino acids in male meat-eaters, fish-eaters, vegetarians and vegans: a cross-sectional analysis in the EPIC-Oxford cohort. *Eur. J. Clin. Nutr.* **70**, 306–312 (2016).

5. Miller, R. A. *et al.* Methionine-deficient diet extends mouse lifespan, slows immune and lens aging, alters glucose, T4, IGF-I and insulin levels, and increases hepatocyte MIF levels and stress resistance. *Aging Cell* **4**, 119–125 (2005).

6. Orentreich, N., Matias, J. R., DeFelice, A. & Zimmerman, J. A. Low Methionine Ingestion by Rats Extends Life Span. *J. Nutr.* **123**, 269–274 (1993).

7. Yu, D. *et al.* Short-term methionine deprivation improves metabolic health via sexually dimorphic, mTORC1-independent mechanisms. *FASEB J.* **32**, 3471–3482 (2018).

8. Wanders, D. *et al.* FGF21 Mediates the Thermogenic and Insulin-Sensitizing Effects of Dietary Methionine Restriction but Not Its Effects on Hepatic Lipid Metabolism. *Diabetes* **66**, 858–867 (2017).

9. Sun, L., Sadighi Akha, A. A., Miller, R. A. & Harper, J. M. Life-span extension in mice by preweaning food restriction and by methionine restriction in middle age. *J. Gerontol. A. Biol. Sci. Med. Sci.* **64**, 711–722 (2009).

10. Newgard, C. B. *et al.* A branched-chain amino acid-related metabolic signature that differentiates obese and lean humans and contributes to insulin resistance. *Cell Metab.* **9**, 311–326 (2009).

11. White, P. J. *et al.* Branched-chain amino acid restriction in Zucker-fatty rats improves muscle insulin sensitivity by enhancing efficiency of fatty acid oxidation and acyl-glycine export. *Mol. Metab.* **5**, 538–551 (2016).

12. Cummings, N. E. *et al.* Restoration of metabolic health by decreased consumption of branched-chain amino acids. *J. Physiol.* **596**, 623–645 (2018).

13.    Fontana, L. *et al.* Decreased Consumption of Branched-Chain Amino Acids Improves Metabolic Health.

*Cell Rep.* **16**, 520–530 (2016).

14.    Solon-Biet, S. M. *et al.* Branched chain amino acids impact health and lifespan indirectly via amino acid

balance and appetite control. *Nat. Metab.* **1**, 532–545 (2019).

15.    Richardson, N. E. *et al.* Lifelong restriction of dietary branched-chain amino acids has sex-specific

benefits for frailty and lifespan in mice. *Nat. Aging* **1**, 73–86 (2021).

16.    Yeh, C.-Y. *et al.* Late-life protein or isoleucine restriction impacts physiological and molecular

signatures of aging. *Nat. Aging* **4**, 1760–1771 (2024).

17.    Trautman, M. E. *et al.* Dietary isoleucine content defines the metabolic and molecular response to a

Western diet. *BioRxiv Prepr. Serv. Biol.* 2024.05.30.596340 (2024) doi:10.1101/2024.05.30.596340.

18.    Green, C. L. *et al.* Dietary restriction of isoleucine increases healthspan and lifespan of genetically

heterogeneous mice. *Cell Metab.* **35**, 1976-1995.e6 (2023).

19.    Yu, D. *et al.* The adverse metabolic effects of branched-chain amino acids are mediated by isoleucine

and valine. *Cell Metab.* **33**, 905-922.e6 (2021).

20.    Boon, R., Silveira, G. G. & Mostoslavsky, R. Nuclear metabolism and the regulation of the epigenome.

*Nat. Metab.* **2**, 1190–1203 (2020).

21.    Haws, S. A., Leech, C. M. & Denu, J. M. Metabolism and the Epigenome: A Dynamic Relationship.

*Trends Biochem. Sci.* **45**, 731–747 (2020).

22.    Haws, S. A. *et al.* Methyl-Metabolite Depletion Elicits Adaptive Responses to Support Heterochromatin

Stability and Epigenetic Persistence. *Mol. Cell* **78**, 210-223.e8 (2020).

23.    Son, S. M. *et al.* Leucine Signals to mTORC1 via Its Metabolite Acetyl-Coenzyme A. *Cell Metab.* **29**,

192-201.e7 (2019).

24.    Vermeulen, L. *et al.* Wnt activity defines colon cancer stem cells and is regulated by the

microenvironment. *Nat. Cell Biol.* **12**, 468–476 (2010).

25.    Yang, Z., He, M., Austin, J., Sayed, D. & Abdellatif, M. Reducing branched-chain amino acids improves

cardiac stress response in mice by decreasing histone H3K23 propionylation. *J. Clin. Invest.* **133**, e169399

(2023).

26.    Simcox, J. & Lamming, D. W. The central moTOR of metabolism. *Dev. Cell* **57**, 691–706 (2022).

27.  Cheng, Y. *et al.* Leucine deprivation decreases fat mass by stimulation of lipolysis in white adipose

tissue and upregulation of uncoupling protein 1 (UCP1) in brown adipose tissue. *Diabetes* **59**, 17–25 (2010).

28.  Hope, D. C. D. *et al.* Hypoaminoacidemia underpins glucagon-mediated energy expenditure and weight

loss. *Cell Rep. Med.* **3**, 100810 (2022).

29.  Lamming, D. W. *et al.* Young and old genetically heterogeneous HET3 mice on a rapamycin diet are

glucose intolerant but insulin sensitive. *Aging Cell* **12**, 712–718 (2013).

30.  Lamming, D. W. *et al.* Rapamycin-induced insulin resistance is mediated by mTORC2 loss and

uncoupled from longevity. *Science* **335**, 1638–1643 (2012).

31.  Buj, R. *et al.* Suppression of p16 Induces mTORC1-Mediated Nucleotide Metabolic Reprogramming.

*Cell Rep.* **28**, 1971-1980.e8 (2019).

32.  Haws, S. A. *et al.* Intrinsic catalytic properties of histone H3 lysine-9 methyltransferases preserve

monomethylation levels under low S-adenosylmethionine. *J. Biol. Chem.* **299**, 104938 (2023).

33.  Dennis, G. *et al.* DAVID: Database for Annotation, Visualization, and Integrated Discovery. *Genome*

*Biol.* **4**, P3 (2003).

34.  Subramanian, A. *et al.* Gene set enrichment analysis: a knowledge-based approach for interpreting

genome-wide expression profiles. *Proc. Natl. Acad. Sci. U. S. A.* **102**, 15545–15550 (2005).

35.  Argelaguet, R. *et al.* Multi-Omics Factor Analysis-a framework for unsupervised integration of multi-

omics data sets. *Mol. Syst. Biol.* **14**, e8124 (2018).

36.  Xiao, F. *et al.* Effects of individual branched-chain amino acids deprivation on insulin sensitivity and

glucose metabolism in mice. *Metabolism.* **63**, 841–850 (2014).

37.  Xiao, F. *et al.* Leucine deprivation increases hepatic insulin sensitivity via GCN2/mTOR/S6K1 and

AMPK pathways. *Diabetes* **60**, 746–756 (2011).

38.  Lee, M. H., Malloy, C. R., Corbin, I. R., Li, J. & Jin, E. S. Assessing the pentose phosphate pathway

using [2, 3-13 C2 ]glucose. *NMR Biomed.* **32**, e4096 (2019).

39.  Hostetler, K. Y. & Landau, B. R. Estimation of the pentose cycle contribution to glucose metabolism in

tissue in vivo. *Biochemistry* **6**, 2961–2964 (1967).

40.  Bergstrom, J. D. The lipogenic enzyme acetoacetyl-CoA synthetase and ketone body utilization for

denovo lipid synthesis, a review. *J. Lipid Res.* **64**, 100407 (2023).

41.      Rauckhorst, A. J. *et al.* A hierarchical hepatic de novo lipogenesis substrate supply network utilizing

pyruvate, acetate, and ketones. *Cell Metab.* **37**, 255-273.e6 (2025).

42.      Gruber, J. J. *et al.* HAT1 Coordinates Histone Production and Acetylation via H4 Promoter Binding.

*Mol. Cell* **75**, 711-724.e5 (2019).

43.      Soaita, I. *et al.* Dynamic protein deacetylation is a limited carbon source for acetyl-CoA-dependent

metabolism. *J. Biol. Chem.* **299**, 104772 (2023).

44.      Charidemou, E. *et al.* Hyperacetylated histone H4 is a source of carbon contributing to lipid synthesis.

*EMBO J.* **43**, 1187–1213 (2024).

45.      Thoreen, C. C. & Sabatini, D. M. Rapamycin inhibits mTORC1, but not completely. *Autophagy* **5**, 725–

726 (2009).

46.      Baar, E. L., Carbajal, K. A., Ong, I. M. & Lamming, D. W. Sex- and tissue-specific changes in mTOR

signaling with age in C57BL/6J mice. *Aging Cell* **15**, 155–166 (2016).

47.      Green, C. L. *et al.* Sex and genetic background define the metabolic, physiologic, and molecular

response to protein restriction. *Cell Metab.* **34**, 209-226.e5 (2022).

48.      Bellantuono, I. *et al.* A toolbox for the longitudinal assessment of healthspan in aging mice. *Nat. Protoc.*

**15**, 540–574 (2020).

49.      Agrawal, S. *et al.* El-MAVEN: A Fast, Robust, and User-Friendly Mass Spectrometry Data Processing

Engine for Metabolomics. *Methods Mol. Biol. Clifton NJ* **1978**, 301–321 (2019).

50.      Xia, J., Psychogios, N., Young, N. & Wishart, D. S. MetaboAnalyst: a web server for metabolomic data

analysis and interpretation. *Nucleic Acids Res.* **37**, W652-660 (2009).

51.      Zerbino, D. R. *et al.* Ensembl 2018. *Nucleic Acids Res.* **46**, D754–D761 (2018).

52.      Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for differential

expression analysis of digital gene expression data. *Bioinforma. Oxf. Engl.* **26**, 139–140 (2010).

53.      Ritchie, M. E. *et al.* limma powers differential expression analyses for RNA-sequencing and microarray

studies. *Nucleic Acids Res.* **43**, e47 (2015).

54.      Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and Powerful Approach

to Multiple Testing. *J. R. Stat. Soc. Ser. B Methodol.* **57**, 289–300 (1995).

55.      Yuan, Z.-F. *et al.* EpiProfile 2.0: A Computational Platform for Processing Epi-Proteomics Mass

824             Spectrometry Data. *J. Proteome Res.* **17**, 2533–2541 (2018).

825     56.      Thomas, S. P., Haws, S. A., Borth, L. E. & Denu, J. M. A practical guide for analysis of histone post-

826             translational modifications by mass spectrometry: best practices and pitfalls. *Methods* (2019)

827             doi:10.1016/j.ymeth.2019.12.001.

828     57.      Castanza, A. S. *et al.* Extending support for mouse data in the Molecular Signatures Database

829             (MSigDB). *Nat. Methods* **20**, 1619–1620 (2023).

**Figure 1 - Physiologic responses to EAA restrictions and rapamycin treatment.**

**Figure 1: Physiologic responses to EAA restrictions and rapamycin treatment.** (*A*) Visual representation of study design. (*B*) Weekly body weight measurements. (*C*) Bar graph illustrating weekly food consumption normalized to mouse body weight values. (*D-E*) Bar graphs depicting absolute changes in lean and fat mass measured via whole-body MRI following three weeks of dietary intervention or rapamycin treatment. (*F-G*) Bar graphs depicting the body composition percentages of lean and fat following three weeks of dietary intervention or rapamycin treatment. (*H-I*) Glucose tolerance test curves and calculated area under the curve (AUC) values. Error bars = SEM; N = 12; * = p-value < 0.05 as measured via Student's t-test.

# Figure 2 - Alterations in hepatic metabolism following EAA restrictions and rapamycin treatment.

**Figure 2: Alterations in hepatic metabolism following EAA restrictions and rapamycin treatment.** (*A*) Heatmap with hierarchical clustering of metabolite abundance values. (*B*) PCA analysis of log transformed, relative metabolite abundance values. (*C-F*) Diagram depicting log2 fold-changes in metabolite abundance relative to controls within the larger context of their associated metabolic pathways. N = 12 for control, rapamycin, Met-D, and Ile-D groups; N = 11 for Leu-D;  * = p-value measured using default MetaboAnalyst settings.

**Figure 3 - Hepatic histone PTM responses to EAA restrictions and rapamycin treatment.**

**Figure 3: Hepatic histone PTM responses to EAA restrictions and rapamycin treatment.** (*A*) Heatmap with hierarchical clustering of histone peptide stoichiometric abundance values compared to control values. (*B*) PCA analysis of log transformed, relative histone peptide stoichiometry values. (*C-L*) Box plots depicting deconvoluted stoichiometry percentages of residue-specific histone PTMs relative to controls. Error bars = standard deviation; N = 12 for control, N = 10 for Rapa; N = 7 for Leu-D; N = 6 for Met-D and Ile-D; * = p-value < 0.05 as measured via Student's t-test.

**Figure 4 - Hepatic transcriptional programs elicited by EAA restrictions and rapamycin treatment.**

**Figure 4: Hepatic transcriptional programs elicited by EAA restrictions and rapamycin treatment.** (*A*) PCA analysis of log transformed gene CPM values. (*B*) Venn diagram of differentially expressed genes with p-adj. <0.05. (*C-F*) Volcano plots depicting log2 FC and -log10 adj. p-values for all identified genes. Horizontal marker indicates threshold for statistical significance (p-adj. < 0.05) with significantly changing genes marked in red. (*G-J*) Dot plots illustrating enriched Gene Ontology – Biological Process (GO-BP) gene sets as determined via Gene Set Enrichment Analyses (GSEA) of differentially expressed genes unique to each given condition. Example GSEA plots highlighting significantly enriched GO-BP gene groups with positive and negative normalized enrichment scores (NES) are present beneath each corresponding dot plot. N = 12.

# Figure 5 - Multi-Omics Factor Analysis of whole organism and hepatic molecular measurements.

**Figure 5: Multi-Omics Factor Analysis of whole organism and hepatic molecular measurements.** (*A*) Heatmap of variance decomposition by factor. (*B*) Plot illustrating Pearson correlation coefficients for control and treatment groups with 15 MOFA Factors. (*C*) Heatmap with hierarchical clustering of Factor 1-associated histone peptide stoichiometric abundance values compared to control values. Factor 1-associated histone peptides were determined using a weight cutoff value of +/- 0.2 which identified the top 18 associated histone peptides. (*D-E*) Plots depicting a subset of significantly enriched gene sets identified using Factor 1 transcripts. All significantly enriched gene sets can be found in Extended Data Figure 2. * = p-value < 0.05 as measured via Student's t-test.

**Figure 6 - Metabolic and epigenetic mechanisms elicited by Ile-D of isolated cells.**

**Figure 6: Metabolic and epigenetic mechanisms elicited by Ile-D of isolated cells.** (*A*) Schematic depicting experiment design for in vitro Ile-D of HepG2 hepatocellular carcinoma cells. (*B-C*) Plots depicting changes in metabolite abundance relative to 0 hr control values. (*E-F*) Heatmaps of metabolite abundance values. (*G*) Heatmap with hierarchical clustering of histone peptide stoichiometric abundance values compared to 0 hr control values. Depicted peptides of interest were originally identified via MOFA in Fig 5. Histone peptides with similar changes in abundance *in vitro* as previously identified *in vivo* are underlined. Error bars = STDEV, N = 4 for metabolite measurements; N = 3 for histone PTM measurements; * = p-value < 0.05 as measured via Student's t-test.

**Extended Data Figure 1 - Differential p62 expression following dietary EAA depletions.**

**Extended Data Figure 1: Differential p62 expression following dietary EAA depletions.** (*A*) Bar graph depicting muscle (quadricep) p62 protein levels normalized to GAPDH. (*B*) Representative western blot image of those used for the quantification depicted in panel A. Error bars = SEM; N = 12; * = p-value < 0.05 as measured via Student's t-test.