Population size interacts with reproductive longevity to shape the germline mutation rate

Luke Zhu[1*], Annabel Beichman[2*], and Kelley Harris[2,3**]

[1]Department of Bioengineering, University of Washington

[2]Department of Genome Sciences, University of Washington

[3]Computational Biology Division, Fred Hutchinson Cancer Center

*These authors contributed equally to this work
**Corresponding Author (harriske@uw.edu)

Classification: Biological Sciences; Evolution

Keywords: Mutation rate, generation time, effective population size, nearly neutral theory, mutator allele

Abstract

Mutation rates vary across the tree of life by many orders of magnitude, with lower mutation rates in species that reproduce quickly and maintain large effective population sizes. A compelling explanation for this trend is that large effective population sizes facilitate selection against weakly deleterious "mutator alleles" such as variants that interfere with the molecular efficacy of DNA repair. However, in multicellular organisms, the relationship of the mutation rate to DNA repair efficacy is complicated by variation in reproductive age. Long generation times leave more time for mutations to accrue each generation, and late reproduction likely amplifies the fitness consequences of any DNA repair defect that creates extra mutations in the sperm or eggs. Here, we present theoretical and empirical evidence that a long generation time amplifies the strength of selection for low mutation rates in the spermatocytes and oocytes. This leads to the counterintuitive prediction that the species with the highest germline mutation rates per generation are also the species with most effective mechanisms for DNA proofreading and repair in their germ cells. In contrast, species with different generation times accumulate similar mutation loads during embryonic development. Our results parallel recent findings that the longest-lived species have the lowest mutation rates in adult somatic tissues, potentially due to selection to keep the lifetime mutation load below a harmful threshold.

Significance Statement

All cells accumulate mutations due to DNA damage and replication errors. When mutations occur in germ tissues including sperm, eggs, and the early embryo, they create changes in the gene pool that can be passed down to future generations. Here, we examine how rates of

germline mutations vary within and between mammalian species, and we find that species which reproduce at older ages tend to accumulate fewer mutations per year in their sperm and eggs. This finding suggests that the evolution of humans' long reproductive lifespan created evolutionary pressure to improve the fidelity of DNA maintenance in germ tissues, paralleling the pressure to avoid accumulating too many mutations in the body over a long lifespan.

Introduction

Germline mutation rates vary by orders of magnitude across the tree of life and ultimately limit the adaptability and the complexity of each species (1–5). Low mutation rates may limit the rate of adaptation to new challenges (6–8), while high mutation rates may limit the ability of a well-adapted population to maintain its fitness and dominance (9, 10). Maintenance of a low mutation rate also incurs an energetic cost, requiring investment of resources and genomic real estate in DNA repair machinery and other mutation-avoiding systems (11–14). As organisms get more complex, the possible consequences of a high mutation rate get more complex as well, leading to confusion and debate about which evolutionary forces ultimately shape this important parameter (15–17).

One widely cited model, the drift barrier hypothesis, posits that mutation rate variation is largely driven by differences in effective population size that modulate the efficacy of selection against weakly deleterious alleles (5, 18–20). A "mutator allele" that raises the germline mutation rate is likely to be deleterious given that harmful mutations outnumber beneficial mutations, but since most mutations are neutral or only weakly harmful, a modest increase in the mutation rate is only expected to decrease fitness by a small amount (21, 22). A corollary of the drift barrier hypothesis is that genetic drift likely limits the ability of DNA repair enzymes to function near their biophysical optima, since optimal functioning would require natural selection to weed out mutator alleles that cause very few additional germline mutations each generation and thus have nearly-neutral fitness effects (23). As a result, different nearly-neutral mutator alleles are likely to accumulate over time in each population and species, causing the molecular

efficacy of each DNA repair enzyme to diverge across the tree of life (24, 25). Although there

exists little direct data on the molecular efficacy of DNA repair and how it varies among species,

the predictions of the drift-barrier hypothesis enjoy broad indirect support from mutation rate

data, which are easier (though still expensive) to measure. Across the tree of life, population

size is inversely correlated with the mutation rate per site per generation (26), and a similar

correlation was recently measured using vertebrate mutation rate data alone (27).

In single-celled organisms, there is a fairly direct connection between DNA repair

efficacy and mutation rate per generation (which is the same as the mutation rate per cell

division). Single-celled organisms also exhibit substantial diversity in the architecture of DNA

repair, ranging from the minimalist repair systems of some obligate symbionts (which have very

high mutation rates (28)) to unique genomic proofreading mechanisms in ciliates such as

Paramecium, which have some of the lowest mutation rates known to science (29–31). In

contrast, multicellular eukaryotes have more standardized cellular housekeeping processes but

varied, multi-stage life histories, with each generation involving multiple cell divisions as well as

potentially mutagenic cell states associated with sex and embryonic development (32–34). This

complexity muddies the relationship between the mutation rate and the molecular efficacy of

DNA repair and complicates the interpretation of the correlation between mutation rate and

effective population size. When Bergeron et al. noted that effective population size was

correlated with mutation rate among vertebrates, they noted that a similar amount of vertebrate

mutation rate variation could be explained by generation time: the typical interval between

reproduction events (27). A strong negative correlation between generation time and the

mutation rate per generation was previously inferred from phylogenetic substitution data, and

the etiology of this pattern has been long debated (16, 17, 35). Measurements of mutation rate

variation within human families have made it clear that generation time can influence the

mutation rate independently of molecular DNA repair efficacy: as parents age, their children are

born with more and more mutations (36, 37).

The effect of parental age on the human mutation rate has been well characterized

thanks to the availability of thousands of mutation rate measurements from trios where the ages

of the parents at the birth of the child are known (38). Similar (though smaller) trio datasets have

also been generated for several non-human mammalian species, and all show the same

qualitative pattern of increasing mutation rate as a function of parental age (39–45). These data

show evidence of significant mutation rate differences among species, and they also differ in

estimates of the rate at which mutation rates increase with the ages of the father and mother.

However, the same sample sizes of most non-human mutation rate studies come with high

degrees of statistical uncertainty, and some recent studies of mutation rates in primates and

carnivores have argued that parental age effects in these species are not statistically

distinguishable from each other (40, 44). Instead, they found that mutation rate measurements

from several primate species, as well as the domestic cat, were consistent with a *reproductive*

longevity model where the molecular efficacy of DNA repair is assumed to be invariant among

species and mutation rate differences are instead driven by differences in the timing of puberty

and reproduction.

Here, we study the etiology of vertebrate mutation rate variation by decomposing it into

its three main components: the rate of mutations that accumulate during embryonic

development, the rate of mutations occurring in the gametes per year of adult reproductive life,

and the length of the time elapsed between puberty and reproduction. Embryonic and gamete

mutation rates are molecular parameters that reflect rates of DNA damage and repair in two

very different germ tissues, while the time elapsed between puberty and reproduction is a

demographic parameter that varies due to a combination of biology and environmental

contingency. Extending the theoretical framework of the drift-barrier hypothesis, we separately

model the fitness effects of variation in the embryonic and gamete mutation rates and infer that

the fitness effects of alleles that increase the gamete mutation rate are likely to scale with

generation time. This scaling reverses the direction of one drift-barrier hypothesis prediction,

implying that selection against gamete mutator alleles will be most effective in species with long

generation times, not in species with large effective population sizes that tend to have short

generation times. We test our predictions by estimating gamete and embryonic mutation rates

from published regressions of mutation rate against generation time from eight mammalian

species: consistent with our model, we find that generation time appears to be positively

correlated with the embryonic mutation rate but negatively correlated with the gamete mutation

rate. We go on to show that mutation rate variation among species is broadly consistent with a

"relaxed clock" reproductive longevity model where embryonic mutation rates vary according to

the classic drift-barrier hypothesis predictions but gamete mutation rates are shaped by a

modified drift-barrier model where selection against mutators is intensified by late reproduction.

Results

As generation time increases, mutation rates in the embryo and the gametes trend in opposite directions

129
130
131

Several recent papers have modeled the etiology of germline mutations by first

separating mutations occurring in early development from mutations that occur post-puberty in

the parents' germ cells (40, 46, 47). In this context, Thomas et al. proposed that there is little

variation among mammals in the total mutation load occurring before puberty and in the

mutation rate per year occurring in the gametes after puberty, but that most germline mutation

rate variation is caused by variation in two demographic parameters: the age of puberty and the

time elapsed between puberty and reproduction. To paraphrase the mathematical description of

their model, we will let P denote the age of puberty, A_M and A_P denote maternal and paternal

ages at conception of an offspring, μ_E denote the rate per generation of mutations that

accumulate in the embryo before puberty, and μ_S and μ_O denote the mutation rates per year in

mature spermatocytes and oocytes. In terms of these variables, the germline mutation rate u_g as

a function of parental age is:

$$u_g(A_P, A_M) = \mu_E + (A_P - P) * \mu_S + (A_M - P) * \mu_O \qquad (1)$$

When Thomas, et al. and Wang, et al. published mutation rate data for owl monkeys (40),

rhesus macaques (42), and domestic cats (44), they inferred species-specific values of the

mutation rate parameters μ_E and $\mu_O + \mu_S$ (relatively few germline mutation rate studies currently

have the power to infer μ_O and μ_S separately). However, they also argued that these species-

specific values did not fit the mutation rate data significantly better than a unified model that

uses mutation rate parameters μ_E, μ_O, and μ_S that were previously inferred from human

mutation data. **Figure 1A,B** illustrates how this reproductive longevity model can explain

variation in mutation rates between species, while **Figure 1C** illustrates a contrasting model

where mutation rate variation is driven by variation in the rate parameters μ_E and $\mu_O + \mu_S$.

153
Figure 1: Models of germline mutation rate variation. A. Within species, mutation rates vary
as a function of age at reproduction. Each individual is expected to accumulate an embryonic
mutation load μ_E plus inherit mutations that accumulated in their parents' sperm and eggs at
rate $\mu_O + \mu_S$ each year between puberty and conception. **B.** Two species with different lifespans
and/or different ages of puberty onset may have different distributions of mutation rates despite
similar mutation parameters μ_E and $\mu_O + \mu_S$, as posited in (40, 44). **C.** Two species with similar
lifespans and similar ages of puberty onset might still have different mutation rates due to
genetic differences that affect rates of DNA damage, repair, or proofreading. This type of
mutation rate variation is driven by variation in the parameters μ_E and/or $\mu_O + \mu_S$.
163
Although the constant-rate reproductive longevity model appears to explain much of the

mutation rate variation among humans, owl monkeys, macaques, and domestic cats, Lindsay,

et al. 2019 previously noted that the spermatocyte mutation rate per year was 5-fold higher in

mice compared to humans (48). To formally test whether the Lindsay, et al. mouse data reject a

constant-rate reproductive longevity model, we inferred μ_E and $\mu_O + \mu_S$ from the Lindsay, et al.

mouse DNM dataset. We found that these rate parameters both significantly diverged from their

human counterparts, with disjoint 95% confidence intervals. In mice, the embryonic mutation

rate $\mu_E = 3.75 \times 10^{-9}$ (95% CI 2.89×10^{-9}; 4.6×10^{-9}), while in humans the rate is nearly 2-

fold higher: $\mu_E = 6.35 \times 10^{-9}$ (95% CI 5.47×10^{-9}; 1.21×10^{-8}). Conversely, the mouse

gamete mutation rate $\mu_O + \mu_S = 1.64 \times 10^{-9}$ (95% CI 4.10×10^{-10}; 2.85×10^{-9}), while in

humans, the rate is 5-fold lower, as previously noted: $\mu_O + \mu_S = 3.5 \times 10^{-10}$ (95% CI

3.3×10^{-10}; 3.7×10^{-10}).

To test whether the difference between mouse and human mutation rate parameters is

representative of a broader dependence of these rate parameters on generation time, we

searched the literature for other regressions of mutation rate against parental age that would

permit estimation of μ_E and $\mu_O + \mu_S$ for additional species. We found appropriate data for five

additional primates plus two carnivores, transformed these species-specific regression

parameters into standardized mutation rate units, and compiled these parameters in **Table 1**.

Species	Embryonic mutation rate μ_E (muts/site/ generation)	Mutation rate $\mu_O + \mu_S$ in the gametes after puberty (muts/site/year)	Age of puberty/ first reproduction (years)	Generation time g (years)	Mutation rate $\mu = \mu_E + g \cdot (\mu_O + \mu_S)$ (muts/site/ generation)
Human (38)	6.26e-9 (95% C.I. 5.47e-9, 12.13e-9)	3.5e-10 (95% C.I. 3.3e-10, 3.7e-10)	13	30	1.2e-8
Chimpanzee (39)	5.11e-9	6.25e-10	14	25	1.2e-8

Olive baboon (41)	5.0e-9	1.4e-10	5.4	10	5.6e-9
Rhesus macaque (42)	3.9e-9	4.3e-10	3.5	8	5.8e-9
Owl monkey (40)	4.4e-9	6.6e-10	1	6.6	8.1e-9
Domestic dog (45)	3.75e-9	4.5e-10	1	4	5.1e-9
Domestic cat (44)	5.9e-9	8.2e-10	0.5	3.8	8.6e-9
Mouse (48)	3.75e-9 (95% C.I. 2.89e-9, 4.6e-9)	1.64e-9 (95% C.I. 4.10e-10, 2.85e-9)	0.15	0.75	4.7e-9

183 **Table 1: Regression-based estimates of embryo and gamete mutation rates.** The
184 generation times and ages at first reproduction in the table are drawn from the publications
185 reporting each set of mutation rate data. See Supplementary Methods for a description of how
186 these standardized rates were calculated from each study's reported data.
187

We performed log-log-linear regressions of μ_E, $\mu_O + \mu_S$, and $\mu = \mu_E + g \cdot (\mu_O + \mu_S)$ as

functions of generation time (log-log linear regressions are more appropriate than natural scale

regressions because the distributions of generation times and mutation rate estimates are

closer to lognormal than normal, as shown in **Supplementary Figure 1**). The regression results

demonstrate that μ_E is positively correlated with generation time across these species, though

less correlated with generation time than the raw germline mutation rate μ (**Figure 2A**). In

contrast, the gamete mutation rate $\mu_O + \mu_S$ is inversely correlated with generation time (**Figure**

2B). We performed all three of these regressions using a phylogenetic least squares (PGLS)

approach but found that these traits had no phylogenetic signal across this small dataset

(Pagel's $\lambda = 0$), indicating that standard linear regression is also appropriate (see

Supplementary Table 1 for details). This result echoes recent findings of inverse correlations

between lifespan and somatic mutation rates, a pattern that is hypothesized to result from

selective pressure to moderate cancer risk and age-related decline in long-lived species (49–

51).

202
Figure 2: Variation among mammals in the rates of germline mutations occurring in the
embryo and the gametes. A. Both the early embryonic mutation rate μ_E and the total mutation
rate $\mu = \mu_E + g \cdot (\mu_O + \mu_S)$ are positively correlated with the generation time g as measured by
log-log linear regression. **B**. The mutation rate per year in the spermatocytes and oocytes post-
puberty, $\mu_O + \mu_S$, is negatively correlated with generation time as measured by a log-log linear
regression.
209

210   **A "relaxed clock" reproductive longevity model predicts mutation rate**
211   **variation across the full range of vertebrate lifespans**
212

Figure 2 suggests that μ_E and $\mu_O + \mu_S$ are not invariant among vertebrate species, but

instead depend on generation time due to factors such as cell division rates, environmental

mutagens, or the molecular efficacy of DNA repair. That being said, **Figure 2** contains data from

only a handful of species due to the limited availability of suitable data for directly estimating μ_E

and $\mu_O + \mu_S$. Estimates of the overall germline mutation rate μ are available for many more

species, and we hypothesized that the relationship among generation time, μ_E, and $\mu_O + \mu_S$

might translate into some constraints on the overall relationship between g and μ. Motivated by

this, we developed a test to evaluate the fit of an empirical mutation rate distribution to either a

strict, fixed-rate reproductive longevity model or a "relaxed clock" reproductive longevity model

where μ_E and $\mu_O + \mu_S$ are allowed to vary among species.

To formulate this test, we first approximated Equation (1) as a simple linear function of

parental age by studying the relationship between age at first reproduction (a proxy for the

timing of puberty) and average age at reproduction (a proxy for the generation time g) in a large

set of vertebrate demographic data (52). In the notation of Equation (1), g equals both the

paternal age A_P and the maternal age A_M. We performed a linear regression of the age at first

reproduction (P) against the average age at reproduction (g) and found that P is approximately

equal to 0.42*g across 230 species with generation times ranging from 2 to 52 years ($r = 0.87$,

see **Supplementary Figure 2**). Motivated by this, we further approximated Equation (1) using

the assumption that $p = P/g$ is a constant across species such that $g - P = g \cdot (1 - P/g) = g \cdot$

$(1 - p)$ and

$$u_g(g) = \mu_E + g \cdot (1 - p) \cdot (\mu_S + \mu_O). \tag{2}$$

233   Letting $\mu_E{}^H$, $\mu_S{}^H$ and $\mu_O{}^H$ be values of the embryonic, spermatocytic, and oocytic

234   mutation rates estimated from human data, we substituted these values into (2) to predict

235   mutation rate in the context of a strict reproductive longevity model that predicts the germline

236   mutation rate u_g as a function of mutation rate parameters $\mu_E{}^H$ and $\mu_O{}^H + \mu_S{}^H$:

$$u_g(g) = \mu_E{}^H + g \cdot (1 - p) \cdot (\mu_O{}^H + \mu_S{}^H) \tag{3}$$

237   We then adapted equation (3) to formulate a relaxed-clock reproductive longevity model that

238   allows the rates μ_E and $\mu_O + \mu_S$ to vary as inferred from our meta-analysis in **Figure 2**. To

239   capture variation in u_E as a function of generation time g, we let $u_E{}^{(g)}$ denote the early

240   embryonic mutation rate at a generation time of g and let α denote the slope relating $log\, \mu_E{}^{(g)}$ to

241   $log\, g$. By these definitions,

$$\tag{4}$$

$$log \; \mu_E{}^{(g)} = log \; \mu_E{}^{(1)} + \alpha \; log \; g.$$

242   Exponentiating both sides of Equation (4) yields:

$$\mu_E{}^{(g)} = \mu_E{}^{(1)} g^{\alpha}. \tag{5}$$

243   To capture gamete mutation rate variation in a similar way, we let β denote the slope of the

244   regression relating $log \; (\mu_S{}^{(g)} + \mu_O{}^{(g)})$ to $log \; g$, such that

$$log \; (\mu_S{}^{(g)} + \mu_O{}^{(g)}) = log \; (\mu_S{}^{(1)} + \mu_O{}^{(1)}) + \beta \; log \; g \tag{6}$$

245   and

$$\mu_S{}^{(g)} + \mu_O{}^{(g)} = (\mu_S{}^{(1)} + \mu_O{}^{(1)}) \cdot g^{\beta}. \tag{7}$$

246   Substituting these values into equation (3) yields a prediction of the overall mutation rate:

$$u_g = \mu_E{}^{(1)} g^{\alpha} + g \cdot (1-p) \cdot (\mu_S{}^{(1)} + \mu_O{}^{(1)}) \cdot g^{\beta} \tag{8}$$

247   This simplifies to

$$u_g = \mu_E{}^{(1)} g^{\alpha} + (1-p) \cdot (\mu_S{}^{(1)} + \mu_O{}^{(1)}) \cdot g^{\beta+1}. \tag{9}$$

Equations (3) and (9) make two different concrete predictions about how mutation rates

should vary with generation time among vertebrates. We were able to compare the accuracy of

these predictions using a large vertebrate mutation rate dataset that was recently compiled by

Wang and Obbard (26). As shown in **Figure 3**, the mutation rate per generation curve predicted

by Equation (9) closely approximates the PGLS correlation between mutation rate and

generation time. In contrast, the human constant-rate reproductive longevity model (Equation

(3) with human-trained parameters) overestimates the mutation rates of species with short

generation times. We also substituted mouse mutation rate parameters into (3) and found that

the resulting model fits the mutation rates of short-generation-time vertebrates but

overestimates the mutation rates of species with longer generation times. Both the human and

mouse reproductive longevity models have greater upward concavity than the relaxed clock model: these models predict a relatively constant mutation rate for generation times less than 1 year, which is the generation time range where these models predict that almost all germline mutations occur in the embryo rather than the gametes.

Figure 3: The relaxed-clock reproductive longevity model explains the correlation
between mutation rate and generation time. A dashed line shows the PGLS regression of
mutation rate versus generation time in vertebrates from Wang and Obbard's mutation rate
meta-analysis (26). This is close to the prediction of the relaxed rate reproductive longevity
model fit to the multispecies pedigree data (solid red line). The prediction of the fixed-rate
reproductive longevity model with human parameters (orange dotted line) overestimates the
mutation rates associated with short generation times, while the fixed-rate reproductive
longevity model with mouse parameters (blue dotted line) overestimates mutation rates
associated with long generation times.

276
277

278   ## Long lifespan increases the efficacy of selection for a low mutation
279   ## rates in the gametes as well as the soma

280
So far, we have shown that vertebrate mutation rate variation is well described by a relaxed-

clock reproductive longevity model where the early embryonic mutation rate per generation

increases with reproductive age and the mutation rate per year in the gametes decreases with

reproductive age. We will now go on to show that both the gamete mutation rate $\mu_O + \mu_S$ and

the embryonic mutation rate μ_E appear to be evolving in accord with the predictions of the drift-

barrier hypothesis, with appropriate modification.

The drift barrier hypothesis explains the inverse correlation between mutation rate and

N_e as a consequence of selection against weakly deleterious mutator alleles (19, 53). Mutator

alleles might directly perturb DNA repair or proofreading, or they might indirectly affect the

mutation rate by perturbing a trait like metabolism. Species with larger effective population sizes

are generally better able to eliminate weakly deleterious alleles, while species with small

effective sizes are more likely to retain these alleles as a result of stronger genetic drift (54).

This leads to the prediction that mutator alleles will be more prevalent in low-N_e species, which

also tend to have long generation times (55, 56). The gamete mutation rate $\mu_O + \mu_S$ seems to

contradict this prediction: we can extrapolate from **Figure 2B** that species with the smallest

effective population sizes are somehow the most effective at eliminating gamete mutator alleles.

We can explain this contradiction by looking more closely at how the fitness effect of a mutator

allele is calculated.

Let $S_u^{(g)}$ be the selection coefficient of a mutator allele that creates u additional

mutations per generation. Lynch previously estimated $S_u^{(g)}$ as follows (57): if L is the length of

the diploid genome and each mutation has an expected fitness cost of E[s], then the expected

selective cost of the mutator allele each generation is

$$S_u^{(g)} = -2uLE[s]. \tag{10}$$

303   In a population of effective size N_e, selection is predicted to eliminate mutations for

304   which $|S_u^{(g)}| > 1/(2N_e)$. By this logic, natural selection should eliminate mutators whose per-

305   generation mutation load u mutations per genome per generation satisfies the inequality

$$u > \frac{1/2N_e}{E[s]} = 1/(2N_e E[s]). \tag{11}$$

If we assume that u, N_e, L, and E[s] are essentially independent variables, then as N_e

gets larger, it will get progressively more difficult for a mutator to satisfy inequality (11) and thus

the population should get more effective at purging away mutator alleles. A caveat is that this

argument does not account for statistical dependence among u, N_e, and the generation time g.

We can reasonably assume that u and g are independent when considering a mutator allele that

modifies μ_E, since such a mutator will create the same embryonic mutation load regardless of

when parents reproduce. However, for a mutator allele that alters $\mu_S + \mu_O$ by creating extra

mutations during spermatogenesis or oogenesis, the total mutation load created by the mutator

each generation will scale proportional to g, as illustrated in **Figure 4A**. This will shift the

distribution of mutator allele fitness effects toward more deleterious values in species with long

generation times, an idea that Lindsay et al. previously posited to explain why mice have higher

per-year germline mutation rates than humans do (48). We will refer to such a modifier of $\mu_S +$

μ_O as a "clocklike" mutator, in contrast to a "non-clocklike" mutator that modifies μ_E by a fixed

amount each generation.

For a clocklike mutator that creates k additional mutations per year after puberty, the

total fitness impact $S_k^{(y)}$ per generation will be the proportional to k times the number of years

that elapse between puberty and reproduction in a generation of length g, which is $g(1-p)$. If

the average fitness cost of a single mutation is $E[s]$, then the total fitness impact of the mutator

each generation will be

$$S_k^{(y)} = kg(1-p)E[s]. \tag{12}$$

325   Since $S_k^{(y)}$ is proportional to the generation time g, this implies that as generation time

326   increases, selection against clocklike mutators may get stronger, decreasing the mutation rate

327   per year in the gametes and explaining the trend in **Figure 2B**. In order for the clocklike mutator

328   to persist in the population, it must satisfy the familiar inequality $S_k^{(y)} > 1/(2N_e)$, which will only

329   hold if

$$k > \frac{1/(2N_e)}{g(1-p)E[s]} = 1/(2gN_e(1-p)E[s]). \tag{13}$$

Inequality (11) defines a threshold of near-neutrality for modifiers of μ_E, while (13)

defines a threshold of near-neutrality for modifiers of $\mu_S + \mu_O$. If we ignore $E[s]$ and p,

assuming that these parameters do not vary much among species, then we conclude that the

efficacy of selection against modifiers of μ_E is determined by N_e alone, while the efficacy of

selection against modifiers of $\mu_S + \mu_O$ is determined by the product gN_e. **Figure 4B** summarizes

how g and N_e interact to shape the gamete mutation load.

A. The fitness effect of a clocklike mutator allele should be proportional to generation time

B. A summary of causal relationships among the variables of the clocklike drift barrier model

Figure 4: A model of germline mutation rate variation as a function of generation time,
effective population size, and genetic variation that impacts the mutation rate measured
per year. A. Here, we compare the effects of identical molecular changes occuring in some
human DNA repair gene as well as its mouse homolog. If these mutator alleles produce the
same number of germline mutations per year, the human allele will produce a greater mutation
burden per generation compared to the mouse allele, leading to a greater expected fitness cost
and a larger negative selection coefficient in the long-generation-time species. *Figure credit:*
Natalie Telis. **B.** This diagram summarizes the multiple ways that generation time can affect the

mutation rate, including its direct impact on the number of mutations that accumulate in a
generation and its other impacts on the effective population size and the efficiency of natural
selection. Pink arrows indicate positive correlations (an increase in the upstream variable
causes an increase in the downstream variable), and blue arrows indicate negative correlations
(an increase in the upstream variable causes a decrease in the downstream variable).
Our calculations suggest that the species with the lowest gamete mutation rates with be

the species for which gN_e is the largest. However, the inverse correlation between g and N_e

means that it is not obvious which life history strategies will maximize gN_e. To gain clarity, we

note that the relationship between N_e and g was previously studied in some detail during the

initial development of the nearly neutral theory, since it was needed to explain the consistency

in molecular substitution rates across the tree of life (58, 59). In this context, Chao and Carr

previously measured an inverse log-linear correlation between N_e and g (55). We were able to

reproduce this log-linear relationship in the Wang and Obbard mutation rate data (26)(**Figure**

5A).

The linear relationship $log\ N_e\ =\ \gamma\ log\ g\ +log\ C$ (where γ and C are constants) implies

that $N_e = Cg^\gamma$ and $gN_e = Cg^{1+\gamma}$. This expression might increase or decrease with increasing g

depending whether γ is greater or less than -1, so knowing the value of γ is key to deciding

whether species with long or short generation times are likely to have the lowest mutation rates.

We estimate that $\gamma \approx -0.487$ based on a PGLS regression of log(N_e) against log(g), remarkably

close to the value of -0.5 that Kimura and Ohta originally proposed to reconcile the nearly

neutral theory with the molecular clock model (55, 60). This implies that g$gN_e = g^{1+\gamma} =$

$g^{1-0.487} = g^{0.513}$. As shown in **Figure 5B,** this implies that gN_e behaves approximately like $\sqrt{g}$,

increasing as g increases. Therefore, if we compare fast-reproducing species like mice to

slower-reproducing species like humans, the slower-reproducing species will have smaller

values of N_e but larger values of gN_e, which is the parameter that determines the strength of

selection for a low mutation rate per year in the gametes. **Figure 5C** shows empirically that gN_e

is negatively correlated with the germline mutation rate per year, consistent with the idea that

the parameter gN_e determines the strength of selection against mutator alleles. We can also

see that humans and other long-lived primates have high values of gN_e compared to the short-

lived mouse.

375
Figure 5: The relationship among N_e, generation time, and the strength of selection
against clocklike mutator alleles. A. The parameters $\log(N_e)$ and $\log(g)$ are inversely
correlated in the Wang and Obbard mutation rate data (26). We estimate a slope of -0.487
based on a PGLS regression. **B.** Expected values of N_e and gN_e as functions of g, extrapolated
from the regression line in panel A and converted from log scale to natural scale. Each curve
has been visualized using an arbitrary y-axis scaling, and together they illustrate that gN_e
increases with increasing g even as N_e decreases. **C**. Mutation rate estimates from Bergeron, et
al. confirm that the mutation rate per year decreases as a function of gN_e, as expected if long
generation times dominate the effect of decreasing effective population size to strengthen
selection against clocklike mutator alleles. Note that the long-lived primates have higher values
of gN_e than the short-lived, high-N_e mouse.
387

Discussion

We have introduced a framework for combining two models of mutation rate evolution,

the reproductive longevity model and the drift-barrier model, into a relaxed-clock reproductive

longevity model that explains the nuanced relationship between mutation rate and reproductive

age. The early embryonic mutation rate appears to have been pushed to its lowest levels in

species with the largest effective population sizes, consistent with the predictions of the nearly

neutral theory. In contrast, the gamete mutation rate trends in the opposite direction, achieving

its lowest levels in long-lived animals with small effective population sizes. This is consistent

with our argument that long generation times should intensify the strength of selection against

clocklike mutator alleles, overcoming the tendency of small effective population sizes to dampen

the general effectiveness of selection.

Variation in the gamete mutation rate per year appears to echo patterns of mutation rate

variation in somatic tissues. A recent study of colon crypt mutations found an inverse log-log

linear relationship between lifespan and the mutation rate per year (51), mirroring the correlation

we observe between generation time and the mutation rate in the gametes. In both cases, the

fitness effect of any mutation rate increase becomes compounded over the lifetime of the cell

lineage that is mutating, giving long-lived, late-reproducing organisms a stronger incentive to

preserve genomic integrity (61, 62). In gerontology, this concept is known as the disposable

soma theory (63, 64), and our analysis suggests that a version of this theory is also applicable

to renewing germline tissues. Since the same molecular machinery is ultimately responsible for

safeguarding both germline and somatic DNA, pleiotropy between somatic and germline

mutation rates may amplify differences among species in the strength of selection against

clocklike mutator alleles.

While selection against nearly neutral mutator alleles is a parsimonious explanation for

the observation that longer generation times are associated with higher rates of embryonic

mutations and lower rates of gamete mutations, other explanations are also possible. Later

reproduction is generally associated with a larger body size and longer gestation, either of which

might cause additional mutations to accumulate in the embryonic germline. It is also possible

that the higher gamete mutation rate in fast-reproducing organisms might be driven by biological

factors such as higher metabolism or higher sperm production volume. These alternate

hypotheses may become testable as additional generation-time-calibrated mutation rate

estimates become available. Our theoretical work underscores the value of collecting mutation

rate data in a way that facilitates separate estimation of embryonic and germ cell mutation rates,

whether by sequencing multi-offspring pedigrees (65–67) or using emerging technologies such

as single-cell gamete sequencing (68, 69).

Recent research on de novo mutagenesis has built a multifactorial case that most

mutations are products of DNA damage rather than cell division error (41, 47, 70–72). However,

embryonic mutations might be the exception to this rule if they largely originate during a few

error-prone postzygotic cell divisions. Human and mouse DNM data, which are higher resolution

than the data available for any other species, make it clear that early embryonic cell divisions

have elevated mutation rates (34, 73–76), possibly due to the reliance of this early-stage

embryo on maternal DNA repair prior to the maternal-zygotic transition (73, 74). However, Drost

and Lee have argued that most mammals, including mice and humans, have similar primordial

germ cell developmental trajectories, with similar numbers of cell divisions leading from the

zygote to the germ cells (32). This implies that variation in the rate of embryonic mutations

among mammals is not likely driven by variation in the number of early embryonic cell divisions

but is more likely driven by variation in DNA damage or repair during early development.

Primordial germ cell specification occurs around gastrulation, which takes place between 6 and

9 days of embryonic development in mouse (77) and between 14 and 21 days of embryonic

development in humans (78). It is possible that the slower pace of early development in longer-

lived vertebrates allows more unrepaired DNA damage to accumulate and drives the tendency

of longer-lived vertebrates to have higher rates of early embryonic mutations.

In addition to making testable predictions about the molecular efficacy of DNA repair and

how it varies among species, our model provides a straightforward way to impute the germline

mutation rates of species for which direct measurements are missing. If a species' age of

reproductive maturity and average generation time have both been estimated, Equation (9)

provides a mutation rate estimate that can be used for calibrating phylogenetic trees and

demographic histories. Although such a mutation rate estimate will not be as accurate as a

mutation rate estimated directly from trio sequencing data, it may be more reliable than

attempting to infer the mutation rate from phylogenetic data, which famously overestimated the

human mutation rate by a factor of 2 (79–81) and also reached inaccurate conclusions about

baleen whale mutation rates (82). Our model may even be useful for imputing the mutation rates

of non-mammalian species; for example, the mutation rate of the black abalone is similar to the

mutation rates of vertebrates with similar reproductive lifespans (83). We have not attempted

here to deduce how mutation rates are affected by body size (84), domestication history (85), or

the countless other variables that may affect genomic integrity, but a good model encapsulating

the effects of generation time should improve our power to learn the effects of additional

variables in the future.

Methods

Meta-analysis of mutation rates from mammalian pedigrees

We obtained estimates of the embryonic mutation rate (μ_E) and the gamete mutation

rate per year after puberty ($\mu_O + \mu_S$) from eight mammalian pedigree studies. Each study

performed a regression of mutation rate against paternal and/or maternal age, but the studies

reported the regression results in a variety of different ways. Below we report how each study's

age regression parameters were transformed into estimates of μ_E and $\mu_O + \mu_S$.

Human: Our human mutation parameter estimates are derived from Jonsson, et al. 2017 (38),

Supplementary Table 6, which gives the maternal slope m_s, maternal intercept m_i, paternal

slope p_s, and paternal intercept p_i of the paper's Poisson regression of the dependence of

mutation rate on parental age (maternal and paternal intercepts represent the interpolated

maternal and paternal mutation loads at a reproductive age of zero years). Upper and lower

95% confidence bounds for each of these variables are also given. The accessible haploid

genome size *A* is listed as 2722501677 base pairs in the caption of Supplementary Table 17.

We calculated μ_E , the mutation load at puberty (age 13) and $\mu_O + \mu_S$, the mutation rate per

year in the gametes post puberty, as follows:

473
$$\mu_E = \frac{1}{2A}(m_i + p_i + 13 \cdot (m_s + p_s))$$

474
$$= \frac{1}{2 \cdot 2722501677} \cdot (3.61 + 6.05 + 13 \cdot (0.37 + 1.51)) = 6.26 \cdot 10^{-9} \text{ muts/bp/gen}$$

475
$$\mu_O + \mu_E = \frac{1}{2A}(m_s + p_s) = \frac{1}{2 \cdot 2722501677} \cdot (0.37 + 1.51) = 3.5 \cdot 10^{-10} \text{ muts/bp/year}$$

The upper and lower confidence bounds on μ_E and $\mu_O + \mu_S$ were calculated in the same way

using the upper and lower bounds of the regression parameters.

Chimpanzee: Venn, et al. (39) reported a chimpanzee paternal age effect of 2.95 additional

mutations per site per year and a maternal age effect of zero additional mutations per site per

year (all regression parameter estimates are given in Table S10). They reported a paternal

intercept of -23.8 total mutations per generation and a flat maternal contribution of 6.65

mutations per generation. The earliest reproductive age reported in the data is 14 years, and the

size of the accessible haploid genome is reported to be 2360 megabases. Using these

parameters, we calculated that:

485
$$\mu_E = \frac{-23.8 + 6.65 + 14 \cdot 2.95}{2 \cdot 2360 \cdot 10^6} = 5.11 \cdot 10^{-9} \text{ muts/bp/generation}$$

486
$$\mu_S + \mu_O = \frac{2.95}{2 \cdot 2360 \cdot 10^6} = 6.25 \cdot 10^{-10} \text{ muts/bp/year.}$$

487   **Olive baboon:** Wu, et al. 2020 (41) reported a paternal slope of 0.15 DNMs per genome per

488   year and a maternal slope of 0.65 DNMs per genome per year (see results section "Estimating

489   sex-specific germline mutation rates and age effects"). These values are scaled to a haploid

490   genome size of $2.581 \cdot 10^9$ base pairs, from which we calculate that

491
$$\mu_O + \mu_S = \frac{0.15 + 0.65}{2 \cdot 2.581 \cdot 10^9} = 1.4 \cdot 10^{-10} \text{ muts/bp/year.}$$

To calculate μ_E, we used the regression coefficients reported in S2 Data, Fig 2B. The reported

maternal intercept is 0.23 mutations per genome at a maternal age of 0.55 years, and the

reported paternal intercept is 22.16 mutations per genome at a paternal age of 0.15 years.

Supplementary Table 14 reports an age of male puberty of 5.41 years, so we estimated the

mutation load at puberty by adding the maternal and maternal intercepts to the estimated

maternal and paternal mutation load accumulated in a period of 5 years. Dividing this load by

the genome size, we obtain:

$$\mu_E = \frac{22.16+0.23+5\times(0.15+0.65)}{2\times2.581\times10^9} = 5.0 \cdot 10^{-9} \text{ muts/bp/generation.}$$

Rhesus macaque: Wang, et al. 2020 (42) report a total parental age slope of $4.3 \cdot 10^{-10}$

mutations per site per year and a mutation load at puberty of $3.9 \cdot 10^{-9}$ mutations per site per

generation. We were able to use these values without further transformation. A second linear

model of macaque mutation rate as a function of generation time was generated by Bergeron, et

al. (43), but we chose to use the Wang et al. model for consistency with the pipeline that was

used to generate the owl monkey and domestic cat mutation rate models.

Owl monkey: Equation (2) in Thomas, et al. 2018 (40) reports a parental age slope of $\mu_O +$

$\mu_S = 6.62 \cdot 10^{-10}$ mutations per site per year and y-intercept of $3.74 \cdot 10^{-9}$. We estimate a pre-

puberty mutation load $\mu_E = 4.40 \cdot 10^{-9}$ assuming a generation time of 1 year and adding a year

of gamete mutation accumulation to the *y*-intercept. Since Thomas, et al. report paternal and

maternal generation times of 6.64 and 6.53, we use an owl monkey generation time of 6.6

years.

Domestic cat: Wang et al. 2022 (44) report mutation rates of $\mu_E = 5.9 \times 10^{-9}$ per site per

generation for reproduction at the age of puberty and an overall average mutation rate of

8.6×10^{-9} mutations per site per generation. They assume that puberty occurs at 0.5 years and

report an average reproductive age of 3.8 years in their data. Using these values we calculate

that

$$\mu_O + \mu_S = \frac{8.6*10^{-9}-5.9\cdot10^{-9}}{3.8-0.5} = 8.2 \cdot 10^{-10} \text{ muts/bp/year.}$$

Domestic dog: Figure 2b of Zhang, et al. 2024 (45) shows bar plot representations of the

slopes and intercepts defining the maternal and paternal mutation rates as linear functions of

reproductive age. Since numerical estimates of these parameters are not reported in the text,

we extrapolated them from the bar plot heights. The maternal mutation rate slope and intercept

appear to be 1×10^{-10} and 8×10^{-10} , while the paternal mutation rate slope and intercept

appear to be 3.5×10^{-10} and 2.5×10^{-9}. Assuming an age of 1 year at puberty (which appears

to be the minimum age at first reproduction represented in the dataset) we conclude that:

$$\mu_O + \mu_S \;=\; 1 \cdot 10^{-10} + 3.5 \cdot 10^{-10} = 4.5 \cdot 10^{-10} \text{ muts/bp/year}$$

$$\mu_E \;=\; 8 \cdot 10^{-10} + 2.5 \cdot 10^{-9} + 1 \cdot 4.5 \cdot 10^{-10} = 3.75 \cdot 10^{-9} \text{ muts/bp/generation}$$

Mouse: We downloaded the supplementary mutation data from Lindsay, et al. 2019 (48), which

reports accessible-genome-corrected mutation counts and parental age at conception in weeks

for all of the offspring in their pedigrees. We performed a regression of mutation rate against

parental age and used the results to calculate means and confidence intervals for murine $\mu_O +$

μ_S and μ_E.

Meta-analysis of the correlation between mutation rate per year and

generation time

We used the nucleotide diversity (π),mutation rate (μ) and generation time (g) data

compiled by Wang and Obbard to quantify the relationship between g and N_e. We first estimated

N_e for each species via the formula $N_e = \pi/(4 \cdot \mu)$ (26) (see Data and Code Availability). We

then performed a PGLS regression of mutation rate against $g \cdot N_e$ using the R library caper (86).

Additionally, we estimated Pagel's λ (87) to be 0.92 using caper's maximum likelihood

implementation. λ is commonly used to quantify the amount of phylogenetic signal in the

dataset. It is a scaling parameter applied to internal branch lengths in the phylogenetic tree, and

is typically a value between 0 and 1. $\lambda = 1$ means that the traits being regressed against one

another appear to have evolved according to a Brownian motion evolutionary model and is

interpreted as strong evidence for phylogenetic signal in the dataset, whereas $\lambda = 0$ suggests

that the traits evolved completely independently of the phylogenetic tree structure. See

Supplementary Table 1 for detailed numerical regression results.

Competing Interest Statement

The authors declare no competing interests.

Data and Code Availability

549   The mutation rates, nucleotide diversity, generation time data, and phylogenetic tree utilized in

550   our calculations were originally compiled by Wang and Obbard (26) and are all publicly available

551   at https://github.com/Yiguan/mutation_literature. The code we used to perform this paper's

552   analysis is available at https://github.com/harrispopgen/clocklike-DBH.

Acknowledgements

We thank Joshua Schraiber for manuscript comments, and we thank Michael Lynch and

members of the Harris Lab for helpful advice and discussions. We also thank Natalie Telis for

figure design assistance. K.H. received support from NIH/NIGMS grant R35GM133428, a

Burroughs Wellcome Fund Career Award at the Scientific Interface, a Searle scholarship, a Pew

Scholarship, the Allen Discovery Center for Cell Lineage Tracing, and a Sloan Fellowship.

A.C.B. received additional support from the National Institute of Health (NIH) "Biological

Mechanisms of Healthy Aging" training program (T32 AG066574).

Supplementary Figures

Supplementary Figure 1: Distributions of generation time and mutation rate per generation across species. Data taken from Wang and Obbard (26). Red and black lines correspond to the fitted normal and log-normal distributions, respectively. Lognormal provides a better fit to both the distribution of generation times and the distribution of the mutation rate per generation.

570 **Supplementary Figure 2: Regression of age at first reproduction versus generation time**.
571 Data taken from Pacifici et al. (52). Age at first reproduction is used as a proxy for age at
572 puberty. Age at first reproduction is found to be linear with respect to generation time, with a
573 slope of 0.42.

References

1. E. G. Leigh, The evolution of mutation rates. *Genetics* **73**, Suppl 73:1-18 (1973).

2. C. F. Baer, M. M. Miyamoto, D. R. Denver, Mutation rate variation in multicellular eukaryotes: causes and consequences. *Nat. Rev. Genet.* **8**, 619–631 (2007).

3. J. J. Bull, R. Sanjuán, C. O. Wilke, Theory of Lethal Mutagenesis for Viruses. *J. Virol.* (2007). https://doi.org/10.1128/jvi.01624-06.

4. M. Eigen, P. Schuster, *The Hypercycle: A Principle of Natural Self-Organization* (Springer Science & Business Media, 2012).

5. M. Lynch, *et al.*, Genetic drift, selection and the evolution of the mutation rate. *Nat. Rev. Genet.* **17**, 704–714 (2016).

6. M. A. Bedau, N. H. Packard, Evolution of evolvability via adaptation of mutation rates. *Biosystems* **69**, 143–162 (2003).

7. J. L. Payne, A. Wagner, The causes of evolvability and their evolution. *Nat. Rev. Genet.* **20**, 24–38 (2019).

8. W. Wei, *et al.*, Rapid evolution of mutation rate and spectrum in response to environmental and population-genetic challenges. *Nat. Commun.* **13**, 4752 (2022).

9. M. Lynch, R. Bürger, D. Butcher, W. Gabriel, The Mutational Meltdown in Asexual Populations. *J. Hered.* **84**, 339–344 (1993).

10. C. Zeyl, M. Mizesko, J. A. G. M. De Visser, Mutational Meltdown in Laboratory Yeast Populations. *Evolution* **55**, 909–917 (2001).

11. J. W. Drake, A constant rate of spontaneous mutation in DNA-based microbes. *Proc. Natl. Acad. Sci.* **88**, 7160–7164 (1991).

12. A. S. Kondrashov, Modifiers of mutation-selection balance: general approach and the evolution of mutation rates. *Genet. Res.* **66**, 53–69 (1995).

13. P. D. Sniegowski, P. J. Gerrish, T. Johnson, A. Shaver, The evolution of mutation rates: separating causes from consequences. *BioEssays* **22**, 1057–1066 (2000).

14. M. Lynch, G. K. Marinov, The bioenergetic costs of a gene. *Proc. Natl. Acad. Sci.* **112**, 15690–15695 (2015).

15. L. Bromham, A. Rambaut, P. H. Harvey, Determinants of rate variation in mammalian DNA sequence evolution. *J. Mol. Evol.* **43**, 610–621 (1996).

16. B. Nabholz, S. Glémin, N. Galtier, Strong Variations of Mitochondrial Mutation Rate across Mammals—the Longevity Hypothesis. *Mol. Biol. Evol.* **25**, 120–130 (2008).

17. L. Bromham, The genome as a life-history character: why rate of molecular evolution
varies between mammal species. *Philos. Trans. R. Soc. B Biol. Sci.* **366**, 2503–2513
(2011).

18. M. Kimura, On the evolutionary adjustment of spontaneous mutation rates. *Genet. Res.* **9**,
23–34 (1967).

19. W. Sung, M. S. Ackerman, S. F. Miller, T. G. Doak, M. Lynch, Drift-barrier hypothesis and
mutation-rate evolution. *Proc. Natl. Acad. Sci.* **109**, 18488–18492 (2012).

20. M. Lynch, *et al.*, The divergence of mutation rates and spectra across the Tree of Life.
EMBO Rep. **24**, e57561 (2023).

21. B. H. Good, M. M. Desai, Evolution of Mutation Rates in Rapidly Adapting Asexual
Populations. *Genetics* **204**, 1249–1266 (2016).

22. W. R. Milligan, G. Amster, G. Sella, The impact of genetic modifiers on variation in
germline mutation rates within and among human populations. *Genetics* **221**, iyac087
(2022).

23. M. Lynch, The evolutionary scaling of cellular traits imposed by the drift barrier. *Proc. Natl.*
Acad. Sci. **117**, 10435–10444 (2020).

24. A. H. Sturtevant, Essays on Evolution. I. On the Effects of Selection on Mutation Rate. *Q.*
Rev. Biol. **12**, 464–467 (1937).

25. K. Harris, J. K. Pritchard, Rapid evolution of the human mutation spectrum. *eLife* **6**, e24284
(2017).

26. Y. Wang, D. J. Obbard, Experimental estimates of germline mutation rate in eukaryotes: a
phylogenetic meta-analysis. *Evol. Lett.* **7**, 216–226 (2023).

27. L. A. Bergeron, *et al.*, Evolution of the germline mutation rate across vertebrates. *Nature*
615, 285–291 (2023).

28. N. A. Moran, H. J. McLaughlin, R. Sorek, The Dynamics and Time Scale of Ongoing
Genomic Erosion in Symbiotic Bacteria. *Science* **323**, 379–382 (2009).

29. M. S. Snoke, T. U. Berendonk, D. Barth, M. Lynch, Large Global Effective Population Sizes
in Paramecium. *Mol. Biol. Evol.* **23**, 2474–2479 (2006).

30. W. Sung, *et al.*, Extraordinary genome stability in the ciliate Paramecium tetraurelia. *Proc.*
Natl. Acad. Sci. **109**, 19339–19344 (2012).

31. H. Long, T. G. Doak, M. Lynch, Limited Mutation-Rate Variation Within the Paramecium
aurelia Species Complex. *G3 GenesGenomesGenetics* **8**, 2523–2526 (2018).

32. J. B. Drost, W. R. Lee, Biological basis of germline mutation: Comparisons of spontaneous
germline mutation rates among drosophila, mouse, and human. *Environ. Mol. Mutagen.*
25, 48–64 (1995).

33.   B. Arbeithuber, A. J. Betancourt, T. Ebner, I. Tiemann-Boege, Crossovers are associated
with mutation and biased gene conversion at recombination hotspots. *Proc. Natl. Acad.*
Sci. **112**, 2109–2114 (2015).

34.   Y. S. Ju, *et al.*, Somatic mutations reveal asymmetric cellular dynamics in the early human
embryo. *Nature* **543**, 714–718 (2017).

35.   J. J. Welch, O. R. Bininda-Emonds, L. Bromham, Correlates of substitution rate variation in
mammalian protein-coding sequences. *BMC Evol. Biol.* **8**, 53 (2008).

36.   N. Risch, E. W. Reich, M. M. Wishnick, J. G. McCarthy, Spontaneous mutation and
parental age in humans. *Am. J. Hum. Genet.* **41**, 218 (1987).

37.   A. Kong, *et al.*, Rate of de novo mutations and the importance of father's age to disease
risk. *Nature* **488**, 471–475 (2012).

38.   H. Jónsson, *et al.*, Parental influence on human germline de novo mutations in 1,548 trios
from Iceland. *Nature* **549**, 519–522 (2017).

39.   O. Venn, *et al.*, Strong male bias drives germline mutation in chimpanzees. *Science* **344**,
1272–1275 (2014).

40.   G. W. C. Thomas, *et al.*, Reproductive longevity predicts mutation rates in primates. *Curr.*
Biol. **28**, 3193-3197.e5 (2018).

41.   F. L. Wu, *et al.*, A comparison of humans and baboons suggests germline mutation rates
do not track cell divisions. *PLOS Biol.* **18**, e3000838 (2020).

42.   R. J. Wang, *et al.*, Paternal age in rhesus macaques is positively associated with germline
mutation accumulation but not with measures of offspring sociability. *Genome Res.* **30**,
826–834 (2020).

43.   L. A. Bergeron, *et al.*, The germline mutational process in rhesus macaque and its
implications for phylogenetic dating. *GigaScience* **10** (2021).

44.   R. J. Wang, *et al.*, De novo mutations in domestic cat are consistent with an effect of
reproductive longevity on both the rate and spectrum of mutations. *Mol. Biol. Evol.* **39**,
msac147 (2022).

45.   S.-J. Zhang, *et al.*, Determinants of de novo mutations in extended pedigrees of 43 dog
breeds. [Preprint] (2024). Available at:
https://www.biorxiv.org/content/10.1101/2024.06.04.596747v1 [Accessed 7 November
2024].

46.   G. Amster, G. Sella, Life history effects on the molecular clock of autosomes and sex
chromosomes. *Proc. Natl. Acad. Sci.* **113**, 1588–1593 (2016).

47.   Z. Gao, M. J. Wyman, G. Sella, M. Przeworski, Interpreting the Dependence of Mutation
Rates on Age and Time. *PLOS Biol.* **14**, e1002355 (2016).

48.   S. J. Lindsay, R. Rahbari, J. Kaplanis, T. Keane, M. E. Hurles, Similarities and differences
678   in patterns of germline mutation between mice and humans. *Nat. Commun.* **10**, 4053
679   (2019).

680   49.   B. Milholland, *et al.*, Differences between germline and somatic mutation rates in humans
681   and mice. *Nat. Commun.* **8**, 15183 (2017).

682   50.   L. Zhang, *et al.*, Maintenance of genome sequence integrity in long- and short-lived rodent
683   species. *Sci. Adv.* **7**, eabj3284 (2021).

684   51.   A. Cagan, *et al.*, Somatic mutation rates scale with lifespan across mammals. *Nature* **604**,
685   517–524 (2022).

686   52.   M. Pacifici, *et al.*, Generation length for mammals. *Nat. Conserv.* **5**, 89–94 (2013).

687   53.   M. Lynch, Evolution of the mutation rate. *Trends Genet.* **26**, 345–352 (2010).

688   54.   T. Ohta, The nearly neutral theory of molecular evolution. *Annu. Rev. Ecol. Syst.* **23**, 263–
689   286 (1992).

690   55.   L. Chao, D. E. Carr, The molecular clock and the relationship between population size and
691   generation time. *Evolution* **47**, 688–690 (1993).

692   56.   R. S. Waples, G. Luikart, J. R. Faulkner, D. A. Tallmon, Simple life-history traits explain
693   key effective population size ratios across diverse taxa. *Proc. R. Soc. B Biol. Sci.* (2013).
694   https://doi.org/10.1098/rspb.2013.1339.

695   57.   M. Lynch, The cellular, developmental and population-genetic determinants of mutation-
696   rate evolution. *Genetics* **180**, 933–943 (2008).

697   58.   T. Ohta, Very slightly deleterious mutations and the molecular clock. *J. Mol. Evol.* **26**, 1–6
698   (1987).

699   59.   T. Ohta, H. Tachida, Theoretical study of near neutrality. I. Heterozygosity and rate of
700   mutant substitution. *Genetics* **126**, 219–229 (1990).

701   60.   M. Kimura, Model of effectively neutral mutations in which selective constraint is
702   incorporated. *Proc. Natl. Acad. Sci.* **76**, 3440–3444 (1979).

703   61.   R. Peto, F. J. Roe, P. N. Lee, L. Levy, J. Clack, Cancer and ageing in mice and men. *Br. J.*
704   *Cancer* **32**, 411–426 (1975).

705   62.   M. Tollis, A. M. Boddy, C. C. Maley, Peto's Paradox: how has evolution solved the problem
706   of cancer prevention? *BMC Biol.* **15**, 60 (2017).

707   63.   L. Hayflick, "Current theories of biological aging" in *Biology of Aging and Development*, G.
708   J. Thorbecke, Ed. (Springer US, 1975), pp. 11–19.

709   64.   T. B. L. Kirkwood, R. Holliday, J. Maynard Smith, R. Holliday, The evolution of ageing and
710   longevity. *Proc. R. Soc. Lond. B Biol. Sci.* **205**, 531–546 (1997).

65. R. Rahbari, *et al.*, Timing, rates and spectra of human germline mutation. *Nat. Genet.* **48**,
126–133 (2016).

66. T. A. Sasani, *et al.*, Large, three-generation human families reveal post-zygotic mosaicism
and variability in germline mutation accumulation. *eLife* **8**, e46922 (2019).

67. D. Porubsky, *et al.*, A familial, telomere-to-telomere reference for human de novo mutation
and recombination from a four-generation pedigree. *bioRxiv* 2024.08.05.606142 (2024).
https://doi.org/10.1101/2024.08.05.606142.

68. A. D. Bell, *et al.*, Insights into variation in meiosis from 31,228 human sperm genomes.
Nature **583**, 259–264 (2020).

69. M. D. Neville, *et al.*, Sperm sequencing reveals extensive positive selection in the male
germline. [Preprint] (2024). Available at:
https://www.medrxiv.org/content/10.1101/2024.10.30.24316414v1 [Accessed 7 November
2024].

70. W. S. W. Wong, *et al.*, New observations on maternal age effect on germline de novo
mutations. *Nat. Commun.* **7**, 10486 (2016).

71. Z. Gao, *et al.*, Overlooked roles of DNA damage and maternal age in generating human
germline mutations. *Proc. Natl. Acad. Sci.* **116**, 9491–9500 (2019).

72. N. Spisak, M. de Manuel, W. Milligan, G. Sella, M. Przeworski, The clock-like accumulation
of germline and somatic mutations can arise from the interplay of DNA damage and repair.
PLOS Biol. **22**, e3002678 (2024).

73. M. A. Eckersley-Maslin, C. Alda-Catalinas, W. Reik, Dynamics of the epigenetic landscape
during the maternal-to-zygotic transition. *Nat. Rev. Mol. Cell Biol.* **19**, 436–450 (2018).

74. E. V. Khokhlova, Z. S. Fesenko, J. V. Sopova, E. I. Leonova, Features of DNA Repair in
the Early Stages of Mammalian Embryonic Development. *Genes* **11**, 1138 (2020).

75. H. Lee, *et al.*, Characterization of early postzygotic somatic mutations through multi-organ
analysis. *J. Hum. Genet.* **66**, 777–784 (2021).

76. A. Uchimura, *et al.*, Early embryonic mutations reveal dynamics of somatic and germ cell
lineages in mice. *Genome Res.* **32**, 945–955 (2022).

77. E. S. Bardot, A.-K. Hadjantonakis, Mouse gastrulation: Coordination of tissue patterning,
specification and diversification of cell fate. *Mech. Dev.* **163**, 103617 (2020).

78. R. C. V. Tyser, *et al.*, Single-cell transcriptomic characterization of a gastrulating human
embryo. *Nature* **600**, 285–289 (2021).

79. A. Scally, R. Durbin, Revising the human mutation rate: implications for understanding
human evolution. *Nat. Rev. Genet.* **13**, 745–753 (2012).

80. A. Scally, Mutation rates and the evolution of germline structure. *Philos. Trans. R. Soc. B
Biol. Sci.* **371**, 20150137 (2016).

81.   P. Moorjani, Z. Gao, M. Przeworski, Human Germline Mutation and the Erratic
748   Evolutionary Clock. *PLOS Biol.* **14**, e2000744 (2016).

749   82.   M. Suárez-Menéndez, *et al.*, Wild pedigrees inform mutation rates and historic abundance
750   in baleen whales. *Science* **381**, 990–995 (2023).

751   83.   T. B. Wooldridge, *et al.*, Direct measurement of the mutation rate and its evolutionary
752   consequences in a critically endangered mollusk. [Preprint] (2024). Available at:
753   https://www.biorxiv.org/content/10.1101/2024.09.16.613283v1 [Accessed 7 November
754   2024].

755   84.   A. P. Martin, S. R. Palumbi, Body size, metabolic rate, generation time, and the molecular
756   clock. *Proc. Natl. Acad. Sci.* **90**, 4087–4091 (1993).

757   85.   M. Bosse, H.-J. Megens, M. F. L. Derks, Á. M. R. de Cara, M. A. M. Groenen, Deleterious
758   alleles in the context of domestication, inbreeding, and selection. *Evol. Appl.* **12**, 6–17
759   (2019).

760   86.   D. Orme, *et al.*, CAPER: comparative analyses of phylogenetics and evolution in R.
761   *Methods Ecol. Evol.* **3**, 145–151 (2013).

762   87.   M. Pagel, Inferring the historical patterns of biological evolution. *Nature* **401**, 877–884
763   (1999).