˜**Full:** The evolutionary dynamics of locally adaptive chromosome inversions in *Mimulus guttatus*

Short title: Adaptive inversion evolution in *Mimulus guttatus*

Authors: Leslie M. Kollar[1,2,3,*], Lauren E. Stanley[1,2,3], Sunil K. Kenchanmane Raju[1,4], David B.
Lowry[1,2,3], Chad E. Niederhuth[1,*]

[1]Department of Plant Biology, Michigan State University, East Lansing, MI, 48824, USA

[2]Plant Resilience Institute, Michigan State University, East Lansing, MI, 48824, USA

[3]Program in Ecology, Evolution, and Behavior, Michigan State University, East Lansing, MI, 48824,
USA

[4]Department of Botany and Plant Sciences, University of California Riverside, Riverside, CA, 92521,
USA

* Co-corresponding authors contributed equally

Corresponding Authors:

Chad E. Niederhuth
Email: niederhuth@protonmail.com
PH: (517) 355-4683
612 Wilson Rd.
East Lansing, MI 48824
19
20   Leslie M. Kollar
21 PH: (517) 355-4683
22 Email: lesliemkollar@gmail.com
23   612 Wilson Rd.
24   East Lansing, MI 48824

Abstract

Chromosomal inversion polymorphisms are ubiquitous across the diversity of diploid organisms and play

a significant role in the evolution of adaptations in those species. Inversions are thought to operate as

supergenes by trapping adaptive alleles at multiple linked loci through the suppression of recombination.

While there is now considerable support for the supergene mechanism of inversion evolution, the extent

to which inversions trap pre-existing adaptive genetic variation versus accumulate new adaptive variants

over time remains unclear. In this study, we report new insights into the evolutionary dynamics of a

locally adaptive chromosomal inversion polymorphism (inv_chr8A), which contributes to the

evolutionary divergence between coastal perennial and inland annual ecotypes of the yellow

monkeyflower, *Mimulus guttatus*. This research was enabled by the sequencing, assembly, and annotation

of new annual and perennial genomes of *M. guttatus* using Oxford Nanopore long-read sequencing

technology. In addition to the adaptive inv_chr8A inversion, we identified three other large inversion

polymorphisms, including a previously unknown large inversion (inv_chr8B) nested within the

inv_chr8A. Through population genomic analyses and comparative genomics, we determined that the

nested inv_chr8B inversion is significantly older than the larger chromosomal inversion in which it

resides. We also evaluated key candidate genes involved in gibberellin biosynthesis and anthocyanin

regulation, which we hypothesize to underlie the adaptive phenotypic effects of the inv_chr8A inversion.

Although little evidence was found to suggest that inversion breakpoint mutations drive adaptive

phenotypic effects, our findings support the supergene mechanism of adaptive evolution as a dynamic and

continuous process.

Introduction

Chromosomal inversions have been implicated in evolutionary adaptations since classic studies in

Drosophila found that inversion polymorphisms are frequently correlated with environmental conditions

(Adrion et al., 2015; Dobzhansky, 1951, 1970; Hoffmann & Rieseberg, 2008; Kapun & Flatt, 2019;

Kirkpatrick & Kern, 2012; Villoutreix et al., 2021). Large chromosomal inversions rearrange the structure

of genomes causing major phenotypic consequences (Dobzhansky, 1936; Elgin & Reuter, 2013; Lupski,

1998; Marshall et al., 2008; Muller, 1930; Puig et al., 2015). Inversions strongly suppress genetic

recombination in heterokaryotic individuals (inversion heterozygotes) because recombinant gametes are

unbalanced (Dobzhansky, 1970; Huang & Rieseberg, 2020; Rieseberg, 2001). Researchers have thus

hypothesized that inversions could act as supergenes by holding together haplotype blocks containing

multiple locally adaptive polymorphisms through suppressed recombination (Berdan et al., 2023;

Darlington & Mather, 1950; Dobzhansky, 1970; Kirkpatrick & Barrett, 2016; Kirkpatrick & Barton, 2006;

Kirkpatrick & Kern, 2012; Schwander et al., 2014; Thompson & Jiggins, 2014).

The realization that inversions could spread rapidly as a result of trapping locally adaptive alleles at linked

loci led Kirkpatrick and Barton to state a decade ago: "If the local adaptation mechanism is so powerful,

why are inversions not everywhere? One possibility is that they in fact are, and that their frequency has

been greatly underestimated" (Kirkpatrick & Barton, 2006). Since Kirkpatrick and Barton's 2006 paper, it

has become clear that inversion polymorphisms truly are everywhere and are frequently associated with

evolutionary adaptations. At least 49 unique inversions have been linked to adaptive phenotypic traits in

Anopheles mosquitoes (D. Ayala et al., 2014). Inversions have been shown to control the phenotypes

underlying mimicry in butterflies (Joron et al., 2011; Kunte et al., 2014; Thompson & Jiggins, 2014) as

well as mating and migratory behaviors in birds (Jeong et al., 2022; Lamichhaney et al., 2016; Lundberg et

al., 2023; Tuttle et al., 2016). Shifts in the frequency of inversion polymorphisms have been linked to

adaptation to forest versus prairie habitats in deer mice (Hager et al., 2022; Harringmeyer & Hoekstra,

2022) and to global climate change in *Drosophila* (Anderson et al., 2005; Rane et al., 2015). Inversions

appear to play a prominent role in adaptation and domestication of crops, including maize, wheat, and

peaches (Fang et al., 2012; Pyhäjärvi et al., 2013; Zhang et al., 2015; Zhou et al., 2021). Human and

chimpanzee genomes differ by nine major chromosomal inversions and >100 smaller ones (Marquès-

Bonet et al., 2004; Newman et al., 2005), which may suggest they played a key role in speciation from our closest relatives (F. J. Ayala & Coluzzi, 2005; Navarro & Barton, 2003). Despite the extensive progress in understanding adaptive inversion evolution, there is still much to learn about how inversions evolve and what genes contribute to their overall phenotypic effects (Berdan et al., 2023).

In this study, we evaluated the evolutionary dynamics of multiple chromosomal inversion polymorphism in the yellow monkeyflower, *Mimulus guttatus* (syn. *Erythranthe guttata*). This work builds on the previous discovery of a chromosomal inversion polymorphism on chromosome 8 underlying a quantitative trait locus (*DIV1*) in this system. This inversion has shown to be definitively linked to local adaptation through a field reciprocal transplant experiment (Lowry & Willis, 2010). While circumstantial evidence has been presented for the role of inversions in local adaptation in many other systems (Beardmore et al., 1960; Butlin & Day, 1984; Chouteau et al., 2017; Dobzhansky, T, 1947; Hager et al., 2022; Mérot et al., 2020), this is the only inversion polymorphism that has been definitively linked to local adaptation in the wild (Berdan et al., 2023). The inversion at the *DIV1* locus, which we henceforth refer to as inv_chr8A, has large phenotypic effects on a suite of adaptive traits associated with the transition from an annual to perennial life history, including flowering time, branching, allocation to vegetative growth, and herbivore resistance (Lowry et al., 2019; Lowry & Willis, 2010). The inv_chr8A inversion polymorphism is widespread across the geographic range of *Mimulus guttatus*, with one orientation of the inversion commonly found in perennial populations, and the other in annual populations (Lowry & Willis, 2010). The inversion most strikingly contributes to the divergence of coastal perennial populations, which experience cooler conditions and moisture supplied by summer sea fog, and nearby inland annual populations, which complete their life cycle quickly due to diminishing water availability during summer drought (Hall et al., 2010; Lowry et al., 2008).

Recently, Coughlin and Willis (2019) provided evidence to support the hypothesis that the inv_chr8A trapped pre-existing adaptive alleles in crosses between annual *M. guttatus* and a perennial sister species,

M. tilingii, which share the same ancestral chromosomal inversion orientation. The orientation of

inv_chr8A found in perennial *M. guttatus* is derived, even though perenniality itself is thought to be the

ancestral state in this system (Coughlan & Willis, 2019; Friedman, 2014). Further, Coughlin and Willis

(2019) were able to exploit free recombination within the inv_chr8A region to show that there are at least

two quantitative trait loci (QTL) for morphological divergence in crosses between annual *M. guttatus* and

perennial *M. tilingii*. While this result provides some evidence for the supergene hypothesis, the dynamics

of how the inversion has evolved over time remain and the genes underlying the supergene effects of the

inversion are still unknown. Identification of the candidate genes that cause the phenotypic effects of

inversion polymorphisms in *M. guttatus* have been difficult because available genome assemblies derived

from short read sequencing are fragmented (Hellsten et al., 2013) and were available only for ecotypes

with the annual orientation. As a result, the full complement of genomic DNA and the inversion

breakpoints of the adaptive inv_chr8A and other known inversions have remained unidentified.

Here, we report a set of important new discoveries regarding chromosomal inversion polymorphisms in

M. guttatus, some of which play crucial roles in the local adaptation and differentiation of annual and

perennial ecotypes within this species. This progress was made possible by the sequencing, assembly, and

annotation of two new genomes for coastal perennial and inland annual lines of *M. guttatus* using Oxford

Nanopore technologies that we report here. With these genomes, we were able to overcome previous

challenges and identify the breakpoints of these inversions as well as the full complement of genes within

the inversions, including inv_chr8A. To identify candidate genes for the adaptive phenotypic effects of

the inversions, as well as explore possible phenotypic consequences of the inversion breakpoint mutations

themselves, we combined our new genome assemblies with a set of existing population genetic and gene

expression datasets. We also combined our analyses with a population genomic data set to evaluate

whether particular genes and loci within the adaptive inv_chr8A inversion had undergone recent selective

sweeps in either coastal or inland habitats. Through this process, we discovered another large

chromosomal inversion (inv_chr8B) nested within inv_chr8A. This finding suggests that this supergene

may have begun as a moderate sized inversion that then expanded by being trapped within a larger

inversion. Overall, this study brings us closer to understanding how chromosomal inversions evolve and

why genomes rearrange dramatically over long periods of evolutionary history.

Materials and Methods

Tissue harvesting and DNA extraction

Our genome sequencing efforts focused on one inland annual line, LMC-L1 (Yorktown, CA, 38° 51'

50.3388" N, 123° 5' 2.1012" W), and one coastal perennial line, SWB-S1 (Irish Beach, CA, 39° 2' 9.5388"

N, 123° 41' 25.6812" W). While these lines are significant because they have been used in multiple

studies (Friedman, 2014; Gould et al., 2018; Lowry et al., 2019; Lowry & Willis, 2010), we recognize

that these lines are not representative of all coastal perennial and inland annuals. LMC-L1 was inbred for

6 generations and SWB-S1 was inbred for 14 generations in the Duke University greenhouses and the

Michigan State University growth facilities. The fewer number of generations inbreeding of LMC-L1 is

due to the line having a high level of sterility-based inbreeding depression. Floral buds were collected for

DNA extractions from plants that were grown in growth chambers in the following conditions: 16 hour

day length, 22°C day and 18°C night temperatures, with 60% relative humidity, and a light intensity of

460mE. DNA extracted from this tissue was used for both long read Nanopore and short read Illumina

sequencing. For Nanopore sequencing, nuclei were extracted from the floral bud tissue following Lu et al

(Lu et al., 2017), with minor modifications. We ground 1-1.5 g of floral bud tissue into a coarse powder

in liquid nitrogen. Ground tissue was resuspended in LB01 buffer (15 mM Tris pH 7.5, 2 mM EDTA

(Ethylenediaminetetraacetic acid), 80 mM KCl, 20 mM NaCl, 15 mM 2-Mercaptoethanol, 0.15% Triton-

X 100 and 0.5 mM Spermine) on ice with intermittent mixing. Tissue homogenate was then filtered

through four-layers of miracloth (EMD Millipore Corp) followed by filtration through a 20 μm cell

strainer (pluriStrainer®). Filtrate was carefully pipetted on top of equal volume Density Gradient

Centrifugation Buffer (1.7 M Sucrose, 10 mM Tris-HCl pH 8.0. 2 mM MgCl2, 5 mM 2-Mercaptoethanol,

1 mM EDTA, 0.15% Triton-X 100) in a 50 ml falcon tube and centrifuged at 2500g for 30 mins at 4□.

After centrifugation, the supernatant was decanted and the nuclei pellet was used to isolate high-

molecular weight (HMW) DNA using the Nanobind Plant Nuclei Big DNA kit (Circulomics, Baltimore,

MD, Cat # NB-900-801-01).

Oxford Nanopore PromethION sequencing

Large chromosomal rearrangements can be identified using population genetics methods with short-read

sequencing (Todesco et al., 2020), but detecting the full complement of DNA within inversions and

pinpointing inversion breakpoints requires long-read sequencing. To achieve this, we performed Oxford

Nanopore Technologies (ONT) sequencing. Nanopore libraries were prepared using the Oxford Nanopore

SQK-LSK110 Ligation Sequencing Kit and loaded onto a FLO-PRO002 (vR9.4.1) flow cell. Sequencing

was carried out on a PromethION sequencer (21.02.7) for 96 hours, followed by basecalling with the

Nanopore Guppy basecaller software (v4.3.4). We obtained 12,580,640 (120.74 GB) and 13,106,600

(133.36 GB) nanopore reads for LMC-L1 and SWB-S1, respectively, with median read lengths of 6.1 kb

and 7.7 kb. Quality trimming resulted in 9,255,001 reads for LMC-L1 (337x coverage) and 9,423,361

reads for SWB-S1 (360x coverage), with 73.6% and 71.9% of reads passing filtering for LMC-L1 and

SWB-S1, respectively.

Illumina Sequencing

We polished the genome assembly and performed SNP calling using short-read sequencing. DNA was

extracted from unopened floral buds using a DNeasy Plant Mini Kit (Qiagen) and prepped with the

Illumina TruSeq Nano DNA Library prep kit. Sequencing was conducted on one lane of an Illumina

NovaSeq 6000 S4 flow (2x150) using a NovaSeq V1.5 300-cycle reagent kit. Base calling was done with

Illumina RTA v3.4.4, and the output was demultiplexed and converted to FastQ format using Bcl2fastq

v2.20.0.

Oxford Nanopore PromethION assembly, polishing, and scaffolding

Adapters were removed from the reads using Porechop v0.2.5 (https://github.com/rrwick/Porechop) and

quality filtered using NanoLyse and Nanofilt (-q 0 and -l 300). We removed possible contaminants using

a microbe genebank (genbank-k31.sbt.json). *De novo* genome assembly was conducted with Flye v2.9.1

(Kolmogorov et al., 2020), followed by two rounds of polishing using Racon (v1.5.0) (Vaser et al., 2017)

and a single round of Medaka (https://github.com/nanoporetech/medaka). The genome was finished with

two rounds of Pilon (Walker et al., 2014) polishing using the Illumina sequencing. We checked for

misassembly using Tigment-long (Jackman et al., 2018). Scaffolding into chromosomes was initially

performed using Allmaps (Tang et al., 2015) by combining genetic map data (Lowry & Willis, 2010) and

synteny to the IM62, TOL, and NONTOL genomes (phytozome-next.jgi.doe.gov). Additional contigs

were added to the chromosomes through an iterative process using ragtag (Alonge et al., 2022) and lrscaf

(Qin et al., 2019), combined with manual inspection and correction. Genome assembly statistics were

calculated using assembly-stats (*v1.0.1*, https://github.com/sanger-pathogens/assembly-stats).

Completeness of the genome assembly was verified throughout the pipeline using BUSCO v2.0 (Simão et

al., 2015) and the LTR Assembly Index (LAI) pipeline (Ou et al., 2018).

Gene annotations

Genes were annotated using the MAKER pipeline (Cantarel et al., 2008; Holt & Yandell, 2011). Support

for gene models was provided by transcripts assembled from RNA-seq and protein alignments. Illumina

reads were trimmed for adapter sequences using trimmomatic (v0.39; (Bolger et al., 2014) and aligned to

their respective genome using HISAT2 (v2.2.1; (Kim et al., 2019)). PacBio reads were aligned by

minimap2 (v2.24; (Li, 2018; *Phytozome v13*, n.d.)). Transcripts were assembled by StringTie2 (v2.2.1;

(Kovaka et al., 2019)). For protein alignments, protein sequences from *Arabidopsis thaliana*

(Araport11;(Cheng et al., 2017), *Oryza sativa* (v7; (Kawahara et al., 2013)), *Solanum lycopersicum*

(ITAG4.0; (Hosmani et al., 2019)), and UniProtKB/Swiss-Prot plants (release-2022_02; (Schneider et al.,

2005)) were aligned to each genome using Exonerate protein2genome (Slater & Birney, 2005).

Repeatmasker (v4.1.1; (*RepeatMasker Home Page*, n.d.)) was used to soft-mask the genome using TE

sequences identified by Extensive *de-novo* TE Annotator (EDTA, see TE Annotations). An initial round

of MAKER was run using the soft-masked genome and gff files from the transcriptome assembly and

protein alignment. Annotations from this initial round of MAKER were then used to train SNAP

(v2013_11_29; (Korf, 2004) and AUGUSTUS (v3.4.0; (Stanke & Morgenstern, 2005)). To prevent

overfitting, only 600 randomly sampled MAKER annotations were used to train AUGUSTUS. MAKER

was then run a second time using models from SNAP and AUGUSTUS. To identify putative genes

missed by MAKER in either genome, we next used Liftoff (v1.6.3; (Shumate & Salzberg, 2020)) to

transfer annotations from one genome onto the other. Lifted annotations were then filtered using

GffCompare (v0.11.2; (Pertea & Pertea, 2020)) to remove annotations with significant overlap to existing

MAKER annotations. Remaining lifted annotations with valid open reading frames (ORF;

"valid_ORFs=1" or "valid_ORF=True" in the GFF attributes column) were considered as putative genes,

while those without a valid ORF ("valid_ORFs=0" or "valid_ORF=False" in the GFF attributes column)

were considered putative pseudogenes. This process was repeated using annotations from the reference

IM62 (v2.0; Hellsten et al., 2013) genome annotation. Genes were flagged as potentially misannotated

transposons as described previously (Bowman et al., 2017), with the exclusion of searching of Gypsy

HMM profiles, as we found that this led to too many real genes being excluded. Given the often blurry

line between genes and transposons, we retained these gene models in the annotation and simply flag

them as possible transposons.

Gene Function

Gene functions were assigned by first searching for PFAM domains using InterProScan (v5.57-90.0) and

then identifying the top five hits with DIAMOND BLASTX against Arabidopsis TAIR10 proteins (e-

value cutoff 1e-6). Results were integrated using a custom script from Kevin Childs (available at

https://github.com/niederhuth/mimulus-assembly).

Structural variant and SNP calling and annotation

Structural variants were called via the whole genome alignment approach using MUM&Co (*v3.0.0*,

O'Donnell & Fischer, 2020). Pseudo-chromosome assemblies were aligned to each other, switching the

reference and query, for calling SVs for each line. SVs (inversions, deletions, insertions, and duplications)

were quality filtered to ≥ 50 bp. Because both SWB-S1 and LMC-L1 were inbred lines, we removed

heterozygous variants as those are likely false positives. Lastly, we removed called insertions and

deletions with a continuous strand of "NNN," as these segments represent gaps created by RagTag and

are likely false positives. Structural variants were annotated using BEDtools *intersect* (Quinlan & Hall,

2010) to identify overlapping genes and structural variants.

To call SNPs, we aligned the Illumina sequences to each genome. We quality trimmed using

Trimmomatic v0.39 (Bolger et al., 2014) to a minimum length of 50 bps and a quality of phred33. We

used BWA-MEM2 (Vasimuddin et al., 2019) to align SWB-S1 reads to the inland annual genome

(98.63% alignment rate) and LMC-L1 reads to the coastal perennial (98.3% alignment rate) genome.

SNPs were called using GATK (McKenna et al., 2010). We marked duplicates using PICARDTOOLS'

MarkDuplicates function and then used GATK's *HaplotypeCaller* to call SNPs in individual samples. We

then genotyped the SNPs using GATK's *GenotypeGVCF* and filtered them using *VariantFiltration*.

SNPs were annotated using ANNOVAR (Wang et al., 2010) and filtered to include: frameshift insertions,

frameshift deletions, stop loss, stop gain, and splicing. ANNOVAR (Wang et al., 2010) was used to

identify synonymous and nonsynonymous SNPs. To estimate a rate of synonymous substitutions at 4-fold

degenerate sites, we identified 4-fold degenerate sites using degenotate

(https://github.com/harvardinformatics/degenotate.git). We intersected the synonymous SNPs with the list

of 4-fold degenerate sites using bedtools *intersect (Quinlan & Hall, 2010)*.

We also reanalyzed SNPs from pooled sequencing from Gould et al (Gould et al. 2017) using our genome

assemblies (SI Appendix, supplementary methods 2).

Copy number (CNVs) and presence absence variants (PAVs) were identified with a custom script using

the outputs from GENESPACE. We included the *M. guttatus* IM62 v2 genome (Hellsten et al., 2013) as

well as SWB-S1 and LMC-L1 genomes.

Synteny, PAVs, CNVs, and inversion breakpoints

Our new genome assemblies fully localized breakpoint regions in the inv_chr8A inversion on

chromosome 8. Due to the complex repetitive nature of the breakpoints, we defined them as regions

extending from the last gene of the upstream collinear syntenic block to the second gene of the inverted

block. Similarly, the breakpoint on the opposite end spans from the second-to-last gene of the inverted

block to the first gene of the downstream collinear block. (SI Appendix, Fig. S13). To identify a stringent

set of syntenic genes between ecotypes, we implemented GENESPACE (Lovell et al., 2022), a pipeline

that uses both OrthoFinder (Emms, n.d.; Emms & Kelly, 2019) and blastp. For all estimates of sequence

level differences, we used the first base pair of the first gene of the inverted syntenic block to the last gene

of the inverted syntenic block. In addition to inv_chr8A, we were able to localize the boundaries of

another previously known inversion on chromosome 5 (inv_chr5A), as well as two previously unknown

large inversions and many small inversions. These newly identified inversions include a smaller inversion

nested within inv_chr8A (inv_chr8B) and a large inversion on chromosome 14 (inv_chr14A). For all of

these inversions, the repetitive nature of the region around the inversion breakpoints made it challenging

to define the exact nucleotide of the breakpoint mutation. While we could not identify exact breakpoints,

we could define which genes were closest to both sides of each breakpoint. Expression differences at the

breakpoints were identified by reanalyzing Gould et al (Gould et al. 2018, SI appendix, supplementary

methods 3).

qRT-PCR of *GA20ox2* (AT5G51810, Migut.H00683)

To determine whether expression differences in *GA20ox2* might contribute to phenotypic differences

between ecotypes, we conducted qRT-PCR because this gene is expressed at a level too low to be

analyzed with RNA-seq. We extracted RNA from three biological replicates of leaf tissue and the floral

shoot apex from wild-type LMC-L1 and SWB-S1 plants using the Spectrum Plant Total RNA Kit

(Sigma). cDNA was synthesized for each sample using GoScript Reverse Transcription Mix, Oligo(dT)

(Promega). We performed qRT-PCR using Power SYBR Green PCR master mix (Applied Biosystems)

on a CFX96 touch real-time PCR machine (Bio-Rad). The PCR program was as follows: 40 cycles at

95°C for 15 s and 60°C for 30 s. We determined amplification efficiencies for each primer pair using a

dilution series (1:2, 1:4, 1:8, 1:16) of pooled cDNA samples. We used *UBC*, the ubiquitin-conjugating

enzyme gene, as a reference gene to calculate the relative expression of *GA20ox2* using the formula

$(E_{ref})^{CP(ref)}/(E_{target})^{CP(target)}$ (Pfaffl, 2001). No differential expression was found for leaf tissue and these

results are not reported (two-tailed Student's t-test, $p > 0.05$).

Results

De novo assembly of M. guttatus inland annual and coastal perennial lines

The final, polished genome assemblies were approximately 277 Mbps (335X coverage ONT and 150X

coverage WGS) spanning 1,198 contigs for LMC-L1 (inland annual) and 278 Mbps (365X coverage ONT

and 151X) spanning 707 contigs for SWB-S1 (coastal perennial). Both assemblies demonstrated high

contiguity with a contig N50 of 5.83 Mbps (LMC-L1) and 4.90 Mbps (SWB-S1) and a scaffold N50 of

18.2 Mbps (LMC-L1) and 18.8 Mbps (SWB-S1). Based on genetic maps and synteny, we assembled

93.0% of the LMC-L1 bps and 93.7% of the SWB-S1 sequence into 14 chromosomes (Hellsten et al.,

2013). Our genomes recovered more than 98% of the eudicot set of BUSCO (Simão et al., 2015)

orthologs (eudicots_odb10), an improvement from 96.9% in the previously available IM62 v2 assembly

(Hellsten et al., 2013). The LTR Assembly Index (LAI), an independent assessment of assembly quality

based on the completeness of assembled LTR retrotransposons (Ou et al., 2018), was much higher in both

LMC-L1 (13.47) and SWB-S1 (16.28) than the IM62 v2 reference (8.79), likely indicating more complete

assembly of repetitive regions (S1 Appendix, Table S1).

We annotated a total of 27,583 genes in LMC-L1 and 26,876 genes in SWB, in comparison to 28,140

genes in IM62 v2 (Hellsten et al., 2013) (SI Appendix, Table S1). We compared the LMC-L1 and SWB-

S1 genomes using *GENESPACE* (Lovell et al., 2022), identifying 19,245 syntenic genes between LMC-

L1 and SWB-S1 (SI Appendix, Table S2). Additionally in LMC-L1, we found 5,105 presence absence

variants (PAVs), 3,026 pseudogenes, and 1,411 tandem duplicates. In SWB-S1, we found 4,707 PAVs,

3,110 pseudogenes, and 1,401 tandem duplicates.

Genome wide SNPs and structural variants

To identify larger structural variants (SVs) between the LMC-L1 and SWB-S1 genomes, we used

MUM&CO, a whole genome alignment method (O'Donnell & Fischer, 2020). The number of SVs

between the LMC-L1 and SWB-S1 genomes are reported in SI Appendix, Table S3. Across the genome

there were 12,013 genes associated with structural variants in LMC-L1 (variants affecting genes: 1,379

insertions, 8,493 deletions, 72 duplications, and 2,069 inversions) and 11,582 genes associated with

structural variants in the SWB-S1 (variants affecting genes: 1,375 insertions, 8,300 deletions, 51

duplications, and 1,856 inversions).

To identify SNPs between the LMC-L1 and SWB-S1 accessions, we aligned whole genome sequencing

(WGS) from one accession to the other accession's reference genome. From these alignments, we focused

on the evolution of coding regions by identifying high impact SNPs, which include the following variant

types: stop gain (2,787 in SWB-S1 and 2,638 in LMC-L1), stop loss (723 in SWB-S1 and 700 in LMC-

L1), frameshift deletion (7,299 in SWB-S1 and 7,541 in LMC-L1), frameshift insertion (7,002 in SWB-

S1 and 6,951 in LMC-L1), and splicing (3,986 in SWB-S1 and 3,392 in LMC-L1). We also identified all

other synonymous and nonsynonymous SNPs. All coding region SNPs are reported for both genomes in

SI Appendix, Tables S4 and S5.

Identification of large chromosome inversions and their breakpoints

The assembly of the nanopore long-read genomes allowed us for the first time to identify large

chromosomal inversions from sequencing data alone as well as identify the locations of the breakpoints

mutations that formed these inversions in the first place. The locally adaptive Inv_chr8A was found to

span 6.7 Mb in LMC-L1 and 5.6 Mb in SWB-S1 (Fig. 1A, SI Appendix Fig. S1 and Table S5). In

addition, our analysis led to the discovery of large inversion nested within inv_chr8A. This nested

inversion(inv_chr8B) spans 213,792 bps in LMC-L1 and 261,302 bps in SWB-S1 (Fig. 1B, SI Appendix

Fig. 1 and Table S5). Beyond chromosome 8, we were able to localize a previously identified inversion

on chromosome 5 (inv_chr5A), which is approximately 4.2 Mb long in LMC-L1 and 4.0 Mb in SWB-S1

(Fig. 1A, SI Appendix Fig. 1 and Table S5). While we have successfully identified inv_chr5A from our

genome assemblies, it is located near the end of the chromosome arm and there are gaps within the

inversion in LMC-L1 of the assemblies. Therefore, there is uncertainty about the location of this

inversion's breakpoint toward the end of the chromosome. Finally, we identified a fourth large inversion

on chromosome 14 (inv_chr14A), which was previously unknown. The inv_chr14A inversion has an

approximate size of 2.4 Mb in LMC-L1 and 2.8 Mb in SWB-S1 (Fig. 1A, SI Appendix, Fig 1 Table S5).

Because of the differences in size of the inversion regions in the two genome assemblies, we were curious

whether there were any major differences in gene content between the inversion orientations. Our

analyses revealed striking differences in gene content. Inv_chr8A (including inv_chr8B) had 177 LMC-

L1-specific genes, 125 SWB-S1-specific genes, and 613 genes shared between the two genomes.

Inv_chr8B shares 36 genes between the genomes but has 9 LMC-L1-specific genes and 8 SWB-S1-

specific genes. Inv_chr5A had a total of 113 LMC-L1 specific genes and 90 SWB-S1 specific genes but

shared 396 genes between the LMC-L1 and SWB-S1. Lastly, the chromosome 14 inversion had 247

LMC-L1-specific genes, 308 SWB-S1-specific genes, and 1,133 shared genes. Many of these large gene

content differences are due to PAVs and copy number variants (CNVs). Errors in gene annotations could

also contribute to PAVs and CNVs and thus, our estimated gene content differences.

While previous studies have demonstrated the role of inv_chr8A in local adaptation (Lowry & Willis,

2010), and implicated inv_chr5A (Gould et al., 2017), our improved assemblies now capture the full

complement of genes within these inversions and make it possible to identify candidate genes that could

have driven the geographic distributions of these inversions. In the following section, we highlight

genetic differences between inversion orientations. CNVs, including PAVs, could explain both

differences in inversion size and phenotypic differences. Within inv_chr8A, there were 338 CNVs with

302 being PAVs; in inv_chr8B, there were 19 CNVs with 17 being PAVs. Within inv_chr5A, there were

227 CNVs, with 203 being PAVs. In inv_chr14A, there were 100 CNVs, with 84 being PAVs. Some of

these CNVs in the inversions could be explained by genes becoming pseudogenized within the inversion.

In inv_chr8A, including inv_chr8B, there were 90 pseudogenes in LMC-L1 and 74 pseudogenes in SWB-

S1 (SI Appendix, Table S6). Inv_chr8B included 6 pseudogenes in LMC-L1 and 8 pseudogenes in SWB-

S1 (SI Appendix, Table S6). In inv_chr5A, there were 54 pseudogenes in LMC-L1 and 43 pseudogenes in

SWB-S1 (SI Appendix, Table S6). Lastly, in inv_chr14A there were 22 pseudogenes in LMC-L1 and 19

pseudogenes in SWB-S1 (SI Appendix, Table S6). We report SNPs (all SNPs, high impact SNPs, and

synonymous and nonsynonymous SNPs) falling within the chromosome inversions in SI Appendix,

Tables S4 and S5. These SNPs were identified from WGS comparison of the SWB-S1 and LMC-L1

genomes.

Evolutionary histories of large inversions

To determine which inversion orientations are ancestral, we implemented GENESPACE (Lovell et al.,

2022) using *Mimulus lewisii,* a distant relative, as the reference genome for comparison

(http://mimubase.org/FTP/Genomes/) (SI Appendix, Fig. S2). From this analysis, we reasoned that

inversions sharing the same orientation as *M. lewisii* are in the ancestral orientation. We found the

orientations of inv_ch5A (SI Appendix, Fig. S3), inv_chr8A (SI Appendix, Fig. S4), and inv_chr8B (SI

Appendix, Fig. S4) are ancestral in LMC-L1. SWB-S1 has the ancestral orientation for inv_chr14A (SI

Appendix, Fig. S5). The results for inv_chr8A are consistent with the results found by Coughlan et al.

(Coughlan et al., 2021).

In addition to identifying which inversion orientation is ancestral, we were interested in the relative ages

of inversion polymorphisms. Here, we compared the level of divergence between the LMC-L1 and

SWB-S1 genomes for inverted regions and to the rest of the genome. When an inversion first forms, the

two orientations of an inversion should have similar levels of divergence as the rest of the genome. As an

inversion ages, each orientation should accumulate genetic changes that lead to elevated levels of

divergence. To evaluate the relative ages of each inversion, we aligned Illumina reads from LMC-L1 to

the SWB-S1 genome and aligned reads from SWB-S1 to the LMC-L1 genome. We then compared the

levels of divergence at 4-fold degenerate synonymous sites for each inversion. The genome-wide average

divergence at 4-fold degenerate sites was 0.043, regardless of which genome was used as the reference

(SI Appendix, Table 2). Both the inv_chr5A (LMC-L1 Illumina resequencing: 0.055, SWB-S1 Illumina

resequencing: 0.061) and inv_chr8A (LMC-L1 Illumina resequencing: 0.057, SWB-S1 Illumina

resequencing: 0.058) inversions had elevated divergence relative to the genome-wide average, which

suggests that these inversions have been segregating within this species for a long time. The inv_chr14A

had similar levels of 4-fold degenerate site divergence (LMC-L1 Illumina resequencing: 0.041, SWB-S1

Illumina resequencing: 0.037) as the genome-wide average, suggesting that it is a relatively new

inversion. Interestingly, the smaller nested inversion (inv_chr8B) had very high levels of divergence

(LMC-L1 Illumina resequencing: 0.084 SWB-S1 Illumina resequencing: 0.083, SI Appendix, Table 2),

which suggests that it is much older than any of the other inversions.

In addition to directly comparing divergence between the SWB-S1 and LMC-L1 genomes, we were able

to evaluate whether there were differences in ecotype-wide polymorphism using a previous population

genetic dataset (Gould et al., 2017). That study compared population genetic parameters for whole

genome pooled sequences of 101 coastal perennial accessions versus 92 inland annual accessions (Gould

et al., 2017). We reanalyzed this data using our new LMC-L1 and SWB-S1 genome assemblies (SI

Appendix, Methods). Given the improvements in the genome assemblies over the previous reference

IM62 v2 genome (Hellsten et al., 2013), we anticipated potentially new patterns in sequence diversity and

differentiation. This new analysis also allowed us to evaluate how reference bias in alignments can affect

population genetic summary statistic estimation. Regardless of the reference genome, the inland annual

ecotype had higher nucleotide diversity ($\pi = 0.040$ regardless of reference genome). In contrast, the

coastal ecotype pool differed slightly in diversity based on reference genome ($\pi = 0.032$ aligned to SWB-

S1, $\pi = 0.034$ aligned to LMC-L1). The lower diversity of the coastal ecotype is consistent with prior

studies and the hypothesis that the coastal ecotype was founded through a population bottleneck (Gould et

al., 2017; Lowry et al., 2008) (SI Appendix, Fig. S6).

The population genetic data (Gould et al. 2017) allowed us to better evaluate the recent evolutionary

history of the four large inversions. First, we compared the allelic differentiation (*G* statistic) for each

inversion versus the rest of the genome. Consistent with our analysis of four-fold degenerate divergence

between the SWB-S1 and LMC-L1 genomes, the nested inv_chr8B inversion had the highest level of

allelic differentiation between the ecotypes (Fig. 2, SI Appendix, Fig. S7 and Table 1). The inv_chr8A

and inv_chr5A inversions also had higher allelic divergence than the genome-wide average. Also

consistent with the divergence result, the inv_chr14A inversion had similar levels of allelic differentiation

as the rest of the genome. To evaluate whether any of the inversions had undergone a recent selective

sweep, we compared the within ecotype diversity (π) for the inverted region to the rest of the genome.

The idea here is that a significant reduction of within ecotype diversity for any of the inversions would be

consistent with a recent selective sweep. To simplify this analysis, we calculated the ratio of π (π inland

annual pool/π coastal perennial pool) for windows across the genome. A π-ratio elevated above the

genome-wide average (π-ratio = 1.25) would be consistent with a sweep within the coastal perennial

ecotype, while a reduction in the π-ratio would be consistent with a sweep in the inland annual ecotype.

Overall, the distribution of π-ratios was similar for inv_chr5A, inv_chr8A, and inv_chr14A (Fig. 2, SI

Appendix, Fig. S6 and Table 1). In contrast, the distribution of the π-ratio was significantly elevated for

the nested inv_chr8B inversion, suggesting the possibility of a recent selective sweep in the coastal

perennial ecotype.

Candidate genes for inversion supergene effects

With clear resolution of the breakpoints for inv_chr8A and inv_chr8B, we were able to localize candidate

genes that could underlie the phenotypic effects of inversions. We identified genes affected by divergence

as those with outlier (top 1%) windows for the *G* statistic (explained above). Across the genome, the top

windows for *G* contained 753 LMC-L1 and 635 SWB-S1 genes and/or their promoters and transcription

start site (TSS) regions or 3' untranslated regions (UTR). Many of these outlier genes, promoters and TSS

regions, and 3' UTRs were syntenic between the LMC-L1 and SWB-S1 genomes, including 193 genes,

57 promoters and TSS regions, and 90 3' UTRs. Some of these outlier regions also fall within the

chromosome inversions. Outlier windows falling within genes, promoters and TSS regions, and 3' UTRs

are interesting because these could contribute to phenotypic differences between the ecotypes. While

these results do not suggest a direct cause in phenotypic variation, it does suggest these outlier genes are

differentiated and warrant further investigation.

One such gene is *GA20ox2*, which codes for a structural gene in the gibberellin hormone synthesis

pathway. Foliar application of gibberellin growth hormone (GA3) to coastal perennial plants has been

shown to increase internode elongation and convert branches from vegetative stolons to aerial flowering

branches in a manner that mimics inland annual growth architecture (Lowry et al. 2019). *GA20ox2* falls

within inv_chr8A and may contribute to adaptive divergence between the ecotypes. *GA20ox2*

(MgS1_08g02280) was an outlier for G and π ratio in our reanalysis using the SWB-S1 reference

genome, similar to what was found in Gould et al. (Gould et al., 2017) when using IM62 as the reference

genome. Both MgS1_08g02280 and MgL1_08g0964, the gene names for *GA20ox2* in each of the two

reference genomes, appear to be intact genes with ORFs. This was confirmed by Sanger sequencing, and

both genes are expressed based on qRT-PCR (Fig. 3). In LMC-L1, expression of *GA20ox2* was ~1.7-fold

higher in floral shoot apex compared to SWB-S1.

Another likely set of genes underlying phenotypes that differentiate ecotypes are a tandem array of *R2R3-*

MYB genes within inv_chr8A (Cooley et al., 2011; Lowry et al., 2012). Previous work showed that

several calyx, corolla, and leaf anthocyanin pigmentation traits differentiating the ecotypes also map to

the region containing inv_chr8A (Lowry et al., 2012). It has been difficult to resolve the number of

tandem copies of *R2R3-MYB* with previously available genome assemblies (i.e. IM62, TOL, and

NONTOL; phytozome-next.jgi.doe.gov/). Based on our CNV analysis, LMC-L1 and SWB-S1 have

different numbers of copies of these subgroup six *R2R3-MYB* genes. The LMC-L1 genome has five

partial or complete copies of this gene, with three being flagged as pseudogenes (genes without valid

open reading frames) in our analysis, while the SWB-S1 genome appears to have six partial or complete

copies of this gene, none of which were called as pseudogenes (Fig. 4). Interestingly, the subgroup six

R2R3-MYB genes appear syntenic with conserved directionality in LMC-L1 and SWB-S1, except for

MgL1_08g06980, which is in the opposite orientation of all the other MYBs. This could indicate that it

was duplicated independently. In investigating why some LMC-L1 copies were called as pseudogenes,

we found that MgL1_08g07004 matched perfectly to an annotation in IM62 (H00280), which is not

annotated as an R2R3-MYB gene because it is missing distinguishing domains. However,

MgL1_08g07004 almost perfectly matches MgL1_08g07000, especially in its intronic sequence,

indicating that it is a partially duplicated gene.

Genes near inversion breakpoints were differentially expressed

501
502
The identification of inversion breakpoints allows for the evaluation of whether those breakpoints have

disrupted genes in ways that could contribute to phenotypic effects. If an inversion breakpoint occurs

within a gene, it would disrupt and likely eliminate its function. Breakpoints occurring near genes can

also cause changes in gene expression by disrupting or changing the position of *cis*-regulatory elements.

To establish whether gene expression might have been disrupted by the inversion breakpoint mutations,

we evaluated transcript abundances within the breakpoint regions. We focused on the first two and last

two genes of each inversion. We also evaluated transcript abundance for the first genes on either side just

outside of the inverted region (SI Appendix, Fig. S13). We investigated these breakpoint region genes by

looking at differentially expressed genes from RNAseq data (Gould et al., 2018) generated from a

reciprocal transplant experiment (Fig. 5, SI Appendix, Table S7). We found that a total of 7 genes within

the breakpoint regions across 4 inversions showed expression level differences.

DISCUSSION

While chromosomal inversions are now well recognized as key factors in adaptive evolution, there is still

much to be learned about how they evolve over time and what genetic mechanisms underlie their

important phenotypic effects. In general, our results are consistent with the supergene hypothesis of

inversion evolution and suggest that not only are multiple linked loci within the adaptive inversion

contributing to its evolution, but that some of those loci may continue to evolve in response to local

selective pressures in coastal versus inland habitats. The long-read sequencing assemblies that we report

in this study also allowed us to localize three other large chromosomal inversions, each of which

illustrates a key component of the evolution of inversions within species. This included the identification

of the nested inversion (inv_chr8B) within inv_chr8A, which had not been identified in prior studies.

Beyond chromosome 8, we were able to greatly improve the assembly of the inv_chr5A inversion, which

was first reported in a cross between inland annual and coastal perennial lines (Holeski et al., 2014). The

inv_chr5A inversion is suspected to play a role in adaptation based on elevated allelic differentiation of

this region between coastal perennial and inland annual populations (Gould et al., 2018), which is a

similar pattern to that for the locally adaptive inv_chr8A inversion. The inv_chr14A inversion was

mentioned in Flagel et al. (Flagel et al., 2019) as a 4 MB region of suppressed recombination. We were

able to confirm that this region is in fact an inversion. Based on our analyses, the inv_chr14 inversion was

less differentiated between the ecotypes, suggesting that it may not be as geographically widespread as

the chromosome 5 and 8 inversions. Overall, the results of this study illustrate multiple important aspects

of large inversion evolution which we discuss below.

The evolution of complex nested inversions

The biggest surprise of our study was the discovery of the relatively large inv_chr8B inversion (214-261

kb; depending on genome) nested within the adaptive inv_chr8A inversion. The evolution of nested

inversions is not unprecedented, but is also not generally discussed in the context of locally adaptive

inversions. Instead, nesting and clustering of multiple inversions in close proximity is typically associated

with the evolution of sex chromosomes. For sex chromosomes, the accumulation of successive inversions

and other rearrangements leads to contrasting levels of divergence (i.e. strata) for different regions of sex

chromosomes, depending on the age of each region of suppressed recombination (D. Charlesworth, 2023;

Filatov, 2022; Handley et al., 2004; Olito & Abbott, 2023). This process eventually leads to distinct X/Y

or Z/W chromosomes that no longer contain pseudoautosomal regions that recombine at all at meiosis

when heterozygous (Bergero & Charlesworth, 2009; D. Charlesworth, 2023). Beyond sex chromosomes,

studies in both maize and sunflowers have found evidence for large nested inversions (Dawe, 2022;

Mroczek et al., 2006; Todesco et al., 2020). In maize, nested inversions have been important for the

evolution of chromosomal knobs that spread by meiotic drive (Dawe, 2022; Mroczek et al., 2006). The

significance of the autosomal nested inversions in sunflowers is unclear.

The entire region that includes both inversions on chromosome 8 is known to be involved in local

adaptation to coastal perennial versus inland annual habitats (Lowry & Willis, 2010). Given the

geographic distribution of the larger inversion (inv_chr8A), it is also likely involved in adaptation to

wetter versus drier habitats within inland regions (Lowry & Willis, 2010; Oneal et al., 2014; Twyford &

Friedman, 2015). Therefore, natural selection is intimately involved in the spread of inv_chr8A. The

question remains as to whether the smaller inv_chr8b inversion also confers habitat-dependent fitness

advantages. The inv_chr8B inversion is particularly interesting, as it appears to be an ancient inversion

that was later captured by the larger inv_chr8A inversion. These results suggest the intriguing possibility

that inv_chr8B is itself an adaptive supergene that now contributes important phenotypic effects to the

larger adaptive inv_chr8A supergene.

The role of context-dependent suppressed recombination in the evolution of adaptive chromosomal

inversions

Inversions likely play a major role in adaptation because they suppress recombination in heterokaryotic

individuals, while allowing for free recombination in homokaryotic individuals (Berdan et al., 2023;

Navarro et al., 1997). This dynamic means that locally adaptive alleles at distant loci on the same

chromosome can be maintained in the same haplotypes for populations located in divergent habitats. At

the same time, free recombination within each habitat allows for purging of deleterious and maladapted

alleles for the same genomic region within habitats (Berdan et al., 2021; B. Charlesworth, 2012; Huang et

al., 2022). Further, free recombination in each habitat minimizes the interference of selection to act on

multiple loci within that region through Hill-Robertson effects (Felsenstein, 1974; Hill & Robertson,

1966; Roze & Barton, 2006). Our study, as well as a previous one (Gould et al., 2017), found evidence

that study's GA20ox2 has undergone recent selective sweep within the coastal perennial ecotype, while

the rest of the inv_chr8A region did not. This suggests that these loci can evolve independently of other

parts of the inversion with ecotypes. Likewise, a recent study in sunflowers found that a high level of

homozygosity for inversions minimized the mutational load in inverted regions (Huang et al., 2022). In

both of these systems, adaptive inversion polymorphisms are thought to primarily occur in a homozygous

state in alternative habitats, which facilitates recombination between loci within the inverted region.

Overall, the ability of inversions to purge deleterious mutations and minimize Hill-Robertson effects give

inversions a major evolutionary advantage over other types of long regions of suppressed recombination,

such as pericentromeric regions (Kuhl & Vader, 2019; Wong & Filatov, 2023).

Candidate genes underlying the adaptive inversion's phenotypic effects

While the idea that maintenance of adaptive alleles across multiple loci through suppressed recombination

is a primary reason for the association of inversions with local adaptation, efforts to identify the genes

underlying adaptive phenotypic effects of inversions are still in the initial stages (Berdan et al., 2023).

Genes within this inversion involved in the gibberellin (GA) hormone pathway are of particular interest

because of the discovery that the addition of gibberellin (GA3) to coastal perennial plants leads them to

develop more like inland annual plants by converting vegetative stolons into upcurved flowering branches

(Lowry et al., 2019). Introgressing the inland annual orientation of the inversion into the coastal perennial

background resulted in nearly identical phenotypic effects on branching and internode elongation as

spraying coastal plants with GA3 (Lowry & Willis, 2010). The GA gene, *GA20ox2* is a top candidate

gene for contributing to inv_chr8A's phenotypic effects. Not only were these genes allele frequency

outliers in comparisons of perennial and annual populations, but *GA20ox2* was expressed at a higher level

in the shoot apex of LMC-L1 than SWB-S1. *GA20ox2* expression is tissue- and time-specific and we

were not able to find evidence that it was expressed when analyzing RNA-seq data from leaf tissue

(Gould et al., 2017). In other systems (Andrés et al., 2014), *GA20ox2* is typically expressed in the floral

shoot apex, similar to our findings in *M. guttatus.* Gould et al (2017) collected tissue only from leaves,

which did not detect differential expression of *GA20ox2* in the floral shoot apex. In contrast, we found

higher expression of *GA20ox2* in the floral apex for the inland annual than the coastal perennial line when

we isolated RNA only from the floral shoot apex tissue. This expression pattern is consistent with the

hypothesis that inland annuals flower earlier and have an upright growth architecture due to higher

production of gibberellin hormones (Lowry et al., 2019).

While our study, as well as ongoing work, implicates a role for *GA20ox2* in the phenotypic effects of

inv_chr8A, the supergene hypothesis requires that we identify multiple genes within the inversion that

contribute to adaptive phenotypic divergence. A promising candidate for also contributing to adaptive

phenotypic effects is the cluster of R2R3-MYB genes within inv_chr8A. A previous study mapped

variation in five vegetative anthocyanin traits to the region inv_chr8A (Lowry et al., 2012), and coastal

perennial and inland annual populations differ in anthocyanin pigmentation. While this gene cluster is

promising, it has been difficult to link vegetative anthocyanin polymorphisms to evolutionary adaptations

(Hatier & Gould, 2008; Hughes et al., 2013; Hughes & Lev-Yadun, 2023; Lee et al., 1979; Manetas,

2006; Schaefer & Rolshausen, 2006). In addition to this R2R3-MYB gene cluster, genes within the

inv_chr8B inversion may play a role in the phenotypic effects of the locally adaptive inv_chr8A

inversion.

Little evidence for a role of breakpoint mutation in causing the phenotypic effects of adaptive

inversions

While the supergene hypothesis has been championed by evolutionary biologists in recent years, it is not

the only mechanism by which inversions can contribute to adaptive phenotypic variation. Inversion

mutations can also break genes or alter gene expression through mutations in promoters/enhancers or by

changing the local chromatin architecture near breakpoints (Berloco et al., 2014; Elgin & Reuter, 2013;

Muller, 1930; Puig et al., 2015; Vogel et al., 2009).

In this study, we found no evidence in this study that the large inversions disrupted any genes. Instead,

breakpoints were located primarily in gene-poor, repeat-rich regions of the genome. While we found no

evidence that the large inversions in our study broke any genes, it is possible that inversion mutations

could have resulted in changing gene expression of nearby genes by disrupting promoter/enhancer

elements or changing local chromatin landscapes. Indeed, we did find that many of the genes located in

close vicinity of the inversion breakpoints on chromosome 5 and 8 had significantly different levels of

transcript abundance between SWB-S1 and LMC-L1 plants. From our study, we cannot establish whether

the breakpoint mutation itself caused these differences in gene expression or if that regulatory divergence

evolved later. A recent study in *Drosophila* that created synthetic inversions at the position of naturally

occuring inversion polymorphism did not cause any major effects on gene expression. (Kapun et al.,

2023; Said et al., 2018) found that no greater elevation in gene expression for genes near the breakpoints

of the Drosophila In(3R)Payne inversion versus elsewhere in the inversion. These results collectively

suggest that the evolution gene expression divergence for inversions often happens over time after the

inversion mutation through the gradual accumulation of new mutations.

Authors contribution

L.M.K, D.B.L, and C.E.N designed research; L.M.K., C.E.N, L.E.S. and S.K.K.R performed research,

and L.M.K. and C.E.N. analyzed data; and L.M.K., C.E.N., L.E.S., and D.B.L. wrote the paper.

Data, materials, and software availability

Code for genome assembly can be found at https://github.com/niederhuth/mimulus-assembly and code for

calling structural variants and SNPs, reanalysis of pooled sequencing and RNAseq, and figures can be

found at https://github.com/Kollarlm/Breakpoint-mutations-supergene-effects-and-ancient-nested-

rearrangements-in-yellow-monkey-flower.git. Raw ONT sequencing files are available through the NCBI

Sequence Read Archive under project identifier PRJNA1047892 – accessions SAMN38597945 (LMC-

L1) and SAMN38597946 (SWB-S1). Raw RNA sequencing files are also available through the NCBI

Sequence Read Archive under project identifier PRJNA1054563. The genome assemblies, annotations,

TE annotations, and other supporting documents can be found on Dryad under

DOI:10.5061/dryad.2ngf1vhwj and CoGE XXX.

Acknowledgments

This study was funded by the National Science Foundation (NSF) Plant Genome Research Program

2010769 to LMK and 2109560 to LES, Michigan State University Plant Biology start-up funds to CEN,

and by two National Science Foundation Division of Integrative Organismal Systems Grants to DBL

(IOS-1855927 and IOS-2153100). We would like to thank Oxford Nanopore Technologies and the MSU

Genomics Core for providing reagents and services that generated sequencing data that made this work

possible. We would also like to thank Kevin Childs for guidance on genome annotation, Eleanore Ritter

for guidance in detecting structural variants, and Lane Vitek for lab work.

Conflict of interest statement

The authors declare no conflict of interest

References

Adrion, J. R., Hahn, M. W., & Cooper, B. S. (2015). Revisiting classic clines in Drosophila melanogaster

in the age of genomics. *Trends in Genetics: TIG*, *31*(8), 434–444.

Alonge, M., Lebeigle, L., Kirsche, M., Jenike, K., Ou, S., Aganezov, S., Wang, X., Lippman, Z. B.,

Schatz, M. C., & Soyk, S. (2022). Automated assembly scaffolding using RagTag elevates a new

tomato system for high-throughput genome editing. *Genome Biology*, *23*(1), 258.

Anderson, A. R., Hoffmann, A. A., McKechnie, S. W., Umina, P. A., & Weeks, A. R. (2005). The

latitudinal cline in the In(3R)Payne inversion polymorphism has shifted in the last 20 years in

Australian Drosophila melanogaster populations. *Molecular Ecology*, *14*(3), 851–858.

Andrés, F., Porri, A., Torti, S., Mateos, J., Romera-Branchat, M., García-Martínez, J. L., Fornara, F.,

Gregis, V., Kater, M. M., & Coupland, G. (2014). SHORT VEGETATIVE PHASE reduces

gibberellin biosynthesis at the Arabidopsis shoot apex to regulate the floral transition. *Proceedings*

of the National Academy of Sciences of the United States of America, *111*(26), E2760–E2769.

Ayala, D., Ullastres, A., & González, J. (2014). Adaptation through chromosomal inversions in

Anopheles. *Frontiers in Genetics*, *5*, 129.

Ayala, F. J., & Coluzzi, M. (2005). Chromosome speciation: humans, Drosophila, and mosquitoes.

Proceedings of the National Academy of Sciences of the United States of America, *102 Suppl*

1(Suppl 1), 6535–6542.

Beardmore, J. A., Dobzhansky, T. H., & Pavlovsky, O. A. (1960). An attempt to compare the fitness of

polymorphic and monomorphic experimental populations of Drosophila pseudoobscura. *Heredity*,

14(1), 19–33.

Berdan, E. L., Barton, N. H., Butlin, R., Charlesworth, B., Faria, R., Fragata, I., Gilbert, K. J., Jay, P.,

Kapun, M., Lotterhos, K. E., Mérot, C., Durmaz Mitchell, E., Pascual, M., Peichel, C. L., Rafajlović,

M., Westram, A. M., Schaeffer, S. W., Johannesson, K., & Flatt, T. (2023). How chromosomal

inversions reorient the evolutionary process. *Journal of Evolutionary Biology*.

https://doi.org/10.1111/jeb.14242

Berdan, E. L., Blanckaert, A., Butlin, R. K., & Bank, C. (2021). Deleterious mutation accumulation and

the long-term fate of chromosomal inversions. *PLoS Genetics*, *17*(3), e1009411.

Bergero, R., & Charlesworth, D. (2009). The evolution of restricted recombination in sex chromosomes.

Trends in Ecology & Evolution, *24*(2), 94–102.

Berloco, M., Palumbo, G., Piacentini, L., Pimpinelli, S., & Fanti, L. (2014). Position effect variegation

and viability are both sensitive to dosage of constitutive heterochromatin in Drosophila. *G3* , *4*(9),

1709–1716.

Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina sequence

data. *Bioinformatics* , *30*(15), 2114–2120.

Bowman, M. J., Pulman, J. A., Liu, T. L., & Childs, K. L. (2017). A modified GC-specific MAKER gene

annotation method reveals improved and novel gene predictions of high and low GC content in

Oryza sativa. *BMC Bioinformatics*, *18*(1), 522.

Buchfink, B., Xie, C., & Huson, D. H. (2015). Fast and sensitive protein alignment using DIAMOND.

Nature Methods, *12*(1), 59–60.

Butlin, R. K., & Day, T. H. (1984). The effect of larval competition on development time and adult size in

the seaweed fly, Coelopa frigida. *Oecologia*, *63*(1), 122–127.

Cantarel, B. L., Korf, I., Robb, S. M. C., Parra, G., Ross, E., Moore, B., Holt, C., Sánchez Alvarado, A.,

& Yandell, M. (2008). MAKER: an easy-to-use annotation pipeline designed for emerging model

organism genomes. *Genome Research*, *18*(1), 188–196.

Charlesworth, B. (2012). The effects of deleterious mutations on evolution at linked sites. *Genetics*,

190(1), 5–22.

Charlesworth, D. (2023). [Review of *Why and how do Y chromosome stop recombining?*]. *Journal of*

Evolutionary Biology, *36*(3), 632–636.

Cheng, C.-Y., Krishnakumar, V., Chan, A. P., Thibaud-Nissen, F., Schobel, S., & Town, C. D. (2017).

Araport11: a complete reannotation of the Arabidopsis thaliana reference genome. *The Plant*

Journal: For Cell and Molecular Biology, *89*(4), 789–804.

Chouteau, M., Llaurens, V., Piron-Prunier, F., & Joron, M. (2017). Polymorphism at a mimicry supergene

maintained by opposing frequency-dependent selection pressures. *Proceedings of the National*

Academy of Sciences of the United States of America, *114*(31), 8325–8329.

Cooley, A. M., Modliszewski, J. L., Rommel, M. L., & Willis, J. H. (2011). Gene duplication in Mimulus

underlies parallel floral evolution via independent trans-regulatory changes. *Current Biology: CB*,

21(8), 700–704.

Coughlan, J. M., Brown, M. W., & Willis, J. H. (2021). The genetic architecture and evolution of life-

history divergence among perennials in the Mimulus guttatus species complex. *Proceedings.*

Biological Sciences / The Royal Society, *288*(1948), 20210077.

Coughlan, J. M., & Willis, J. H. (2019). Dissecting the role of a large chromosomal inversion in life

history divergence throughout the Mimulus guttatus species complex. *Molecular Ecology*, *28*(6),

1343–1357.

Darlington, C. D., & Mather, K. (1950). The elements of genetics. *Population*, *5*(2), 386.

Dawe, R. K. (2022). The maize abnormal chromosome 10 meiotic drive haplotype: a review.

Chromosome Research: An International Journal on the Molecular, Supramolecular and

Evolutionary Aspects of Chromosome Biology, *30*(2-3), 205–216.

Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., &

Gingeras, T. R. (2013). STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* , *29*(1), 15–21.

Dobzhansky, T. (1936). Position effects on genes. *Biological Reviews of the Cambridge Philosophical*

Society, *11*(3), 364–384.

Dobzhansky, T. (1947). Adaptive Changes Induced by Natural Selection in Wild Populations of

Drosophila. *Evolution; International Journal of Organic Evolution*, *1*(1/2), 1–16.

Dobzhansky, T. (1951). *Genetics and the Origin of Species (3rd ed.)*. Columbia University Press: New

York, NY, USA.

Dobzhansky, T. (1970). *Genetics of the Evolutionary Process*. Columbia University Press.

Elgin, S. C., & Reuter, G. (2013). *Position-effect variegation, heterochromatin formation, and gene*

silencing in Drosophila. Cold Spring Harb. Perspect. Biol. 5: a017780.

Emms, D. M. (n.d.). Kelly S. 2015a. OrthoFinder: solving fundamental biases in whole genome

comparisons dramatically improves orthogroup inference accuracy. *Genome Biology.*

Emms, D. M., & Kelly, S. (2019). OrthoFinder: phylogenetic orthology inference for comparative

genomics. *Genome Biology*, *20*(1), 238.

Fang, Z., Pyhäjärvi, T., Weber, A. L., Dawe, R. K., Glaubitz, J. C., González, J. de J. S., Ross-Ibarra, C.,

Doebley, J., Morrell, P. L., & Ross-Ibarra, J. (2012). Megabase-scale inversion polymorphism in the

wild ancestor of maize. *Genetics*, *191*(3), 883–894.

Felsenstein, J. (1974). The evolutionary advantage of recombination. *Genetics*, *78*(2), 737–756.

Filatov, D. A. (2022). Recent expansion of the non-recombining sex-linked region on Silene latifolia sex

chromosomes. *Journal of Evolutionary Biology*, *35*(12), 1696–1708.

Flagel, L. E., Blackman, B. K., Fishman, L., Monnahan, P. J., Sweigart, A., & Kelly, J. K. (2019).

GOOGA: A platform to synthesize mapping experiments and identify genomic structural diversity.

PLoS Computational Biology, *15*(4), e1006949.

Friedman, J. (2014). Genetic determinants and epistasis for life history trait differences in the common

monkeyflower, Mimulus guttatus. *The Journal of Heredity*, *105*(6), 816–827.

Gould, B. A., Chen, Y., & Lowry, D. B. (2017). Pooled ecotype sequencing reveals candidate genetic

mechanisms for adaptive differentiation and reproductive isolation. *Molecular Ecology.*

https://doi.org/10.1111/mec.13881

Gould, B. A., Chen, Y., & Lowry, D. B. (2018). Gene regulatory divergence between locally adapted

ecotypes in their native habitats. *Molecular Ecology*, *27*(21), 4174–4188.

Hager, E. R., Harringmeyer, O. S., Wooldridge, T. B., Theingi, S., Gable, J. T., McFadden, S.,

Neugeboren, B., Turner, K. M., Jensen, J. D., & Hoekstra, H. E. (2022). A chromosomal inversion

contributes to divergence in multiple traits between deer mouse ecotypes. *Science*, *377*(6604), 399–

405.

Hall, M. C., Lowry, D. B., & Willis, J. H. (2010). Is local adaptation in Mimulus guttatus caused by trade-

offs at individual loci? *Molecular Ecology*, *19*(13), 2739–2753.

Handley, L.-J. L., Ceplitis, H., & Ellegren, H. (2004). Evolutionary strata on the chicken Z chromosome:

implications for sex chromosome evolution. *Genetics*, *167*(1), 367–376.

Harringmeyer, O. S., & Hoekstra, H. E. (2022). Chromosomal inversion polymorphisms shape the

genomic landscape of deer mice. *Nature Ecology & Evolution*, *6*(12), 1965–1979.

Hatier, J.-H. B., & Gould, K. S. (2008). Foliar anthocyanins as modulators of stress signals. *Journal of*

Theoretical Biology, *253*(3), 625–627.

Hellsten, U., Wright, K. M., Jenkins, J., Shu, S., Yuan, Y., Wessler, S. R., Schmutz, J., Willis, J. H., &

Rokhsar, D. S. (2013). Fine-scale variation in meiotic recombination in Mimulus inferred from

population shotgun sequencing. *Proceedings of the National Academy of Sciences of the United*

States of America, *110*(48), 19478–19482.

Hill, W. G., & Robertson, A. (1966). The effect of linkage on limits to artificial selection. *Genetical*

Research, *8*(3), 269–294.

Hoffmann, A. A., & Rieseberg, L. H. (2008). Revisiting the Impact of Inversions in Evolution: From

Population Genetic Markers to Drivers of Adaptive Shifts and Speciation? *Annual Review of*

Ecology, Evolution, and Systematics, *39*, 21–42.

Holeski, L. M., Monnahan, P., Koseva, B., McCool, N., Lindroth, R. L., & Kelly, J. K. (2014). A high-

resolution genetic map of yellow monkeyflower identifies chemical defense QTLs and

recombination rate variation. *G3* , *4*(5), 813–821.

Holt, C., & Yandell, M. (2011). MAKER2: an annotation pipeline and genome-database management tool

for second-generation genome projects. *BMC Bioinformatics*, *12*, 491.

Hosmani, P. S., Flores-Gonzalez, M., van de Geest, H., Maumus, F., Bakker, L. V., Schijlen, E., van

Haarst, J., Cordewener, J., Sanchez-Perez, G., Peters, S., Fei, Z., Giovannoni, J. J., Mueller, L. A., &

Saha, S. (2019). An improved de novo assembly and annotation of the tomato reference genome

using single-molecule sequencing, Hi-C proximity ligation and optical maps. In *bioRxiv* (p. 767764).

https://doi.org/10.1101/767764

Huang, K., Ostevik, K. L., Elphinstone, C., Todesco, M., Bercovich, N., Owens, G. L., & Rieseberg, L.

H. (2022). Mutation Load in Sunflower Inversions Is Negatively Correlated with Inversion

Heterozygosity. *Molecular Biology and Evolution*, *39*(5). https://doi.org/10.1093/molbev/msac101

Huang, K., & Rieseberg, L. H. (2020). Frequency, origins, and evolutionary role of chromosomal

inversions in plants. *Frontiers in Plant Science*.

https://www.frontiersin.org/articles/10.3389/fpls.2020.00296/full

Hughes, N. M., Carpenter, K. L., & Cannon, J. G. (2013). Estimating contribution of anthocyanin

pigments to osmotic adjustment during winter leaf reddening. *Journal of Plant Physiology*, *170*(2),

230–233.

Hughes, N. M., & Lev-Yadun, S. (2023). Review: Why do some plants have leaves with red or purple

undersides? *Environmental and Experimental Botany*, *205*, 105126.

Jackman, S. D., Coombe, L., Chu, J., Warren, R. L., Vandervalk, B. P., Yeo, S., Xue, Z., Mohamadi, H.,

Bohlmann, J., Jones, S. J. M., & Birol, I. (2018). Tigmint: correcting assembly errors using linked

reads from large molecules. *BMC Bioinformatics*, *19*(1), 393.

Jeong, H., Baran, N. M., Sun, D., Chatterjee, P., Layman, T. S., Balakrishnan, C. N., Maney, D. L., & Yi,

S. V. (2022). Dynamic molecular evolution of a supergene with suppressed recombination in white-

throated sparrows. *eLife*, *11*. https://doi.org/10.7554/eLife.79387

Jones, P., Binns, D., Chang, H.-Y., Fraser, M., Li, W., McAnulla, C., McWilliam, H., Maslen, J.,

Mitchell, A., Nuka, G., Pesseat, S., Quinn, A. F., Sangrador-Vegas, A., Scheremetjew, M., Yong, S.-

Y., Lopez, R., & Hunter, S. (2014). InterProScan 5: genome-scale protein function classification.

Bioinformatics , *30*(9), 1236–1240.

Joron, M., Frezal, L., Jones, R. T., Chamberlain, N. L., Lee, S. F., Haag, C. R., Whibley, A., Becuwe, M.,

Baxter, S. W., Ferguson, L., Wilkinson, P. A., Salazar, C., Davidson, C., Clark, R., Quail, M. A.,

Beasley, H., Glithero, R., Lloyd, C., Sims, S., … ffrench-Constant, R. H. (2011). Chromosomal

rearrangements maintain a polymorphic supergene controlling butterfly mimicry. *Nature*, *477*(7363),

203–206.

Kapun, M., & Flatt, T. (2019). The adaptive significance of chromosomal inversion polymorphisms in

Drosophila melanogaster. *Molecular Ecology*, *28*(6), 1263–1282.

Kapun, M., Mitchell, E. D., Kawecki, T. J., Schmidt, P., & Flatt, T. (2023). An Ancestral Balanced

Inversion Polymorphism Confers Global Adaptation. *Molecular Biology and Evolution*, *40*(6).

https://doi.org/10.1093/molbev/msad118

Kawahara, Y., de la Bastide, M., Hamilton, J. P., Kanamori, H., McCombie, W. R., Ouyang, S.,

Schwartz, D. C., Tanaka, T., Wu, J., Zhou, S., Childs, K. L., Davidson, R. M., Lin, H., Quesada-

Ocampo, L., Vaillancourt, B., Sakai, H., Lee, S. S., Kim, J., Numa, H., … Matsumoto, T. (2013).

Improvement of the Oryza sativa Nipponbare reference genome using next generation sequence and

optical map data. *Rice* , *6*(1), 4.

Kim, D., Paggi, J. M., Park, C., Bennett, C., & Salzberg, S. L. (2019). Graph-based genome alignment

and genotyping with HISAT2 and HISAT-genotype. *Nature Biotechnology*, *37*(8), 907–915.

Kirkpatrick, M., & Barrett, B. (2016). Chromosome inversions, adaptive cassettes and the evolution of

species' ranges. In *Invasion Genetics* (pp. 175–186). John Wiley & Sons, Ltd.

Kirkpatrick, M., & Barton, N. (2006). Chromosome inversions, local adaptation and speciation. *Genetics*,

173(1), 419–434.

Kirkpatrick, M., & Kern, A. (2012). Where's the Money? Inversions, Genes, and the Hunt for Genomic

Targets of Selection. *Genetics*, *190*(4), 1153–1155.

Kolmogorov, M., Bickhart, D. M., Behsaz, B., Gurevich, A., Rayko, M., Shin, S. B., Kuhn, K., Yuan, J.,

Polevikov, E., Smith, T. P. L., & Pevzner, P. A. (2020). metaFlye: scalable long-read metagenome

assembly using repeat graphs. *Nature Methods*, *17*(11), 1103–1110.

Korf, I. (2004). Gene finding in novel genomes. *BMC Bioinformatics*, *5*, 59.

Kovaka, S., Zimin, A. V., Pertea, G. M., Razaghi, R., Salzberg, S. L., & Pertea, M. (2019). Transcriptome

assembly from long-read RNA-seq alignments with StringTie2. *Genome Biology*, *20*(1), 278.

Kuhl, L.-M., & Vader, G. (2019). Kinetochores, cohesin, and DNA breaks: Controlling meiotic

recombination within pericentromeres. *Yeast* , *36*(3), 121–127.

Kunte, K., Zhang, W., Tenger-Trolander, A., Palmer, D. H., Martin, A., Reed, R. D., Mullen, S. P., &

Kronforst, M. R. (2014). doublesex is a mimicry supergene. *Nature*, *507*(7491), 229–232.

Lamichhaney, S., Fan, G., Widemo, F., Gunnarsson, U., Thalmann, D. S., Hoeppner, M. P., Kerje, S.,

Gustafson, U., Shi, C., Zhang, H., Chen, W., Liang, X., Huang, L., Wang, J., Liang, E., Wu, Q., Lee,

S. M.-Y., Xu, X., Höglund, J., … Andersson, L. (2016). Structural genomic changes underlie

alternative reproductive strategies in the ruff (Philomachus pugnax). *Nature Genetics*, *48*(1), 84–88.

Lee, D. W., Lowry, J. B., & Stone, B. C. (1979). Abaxial Anthocyanin Layer in Leaves of Tropical Rain

Forest Plants: Enhancer of Light Capture in Deep Shade. *Biotropica*, *11*(1), 70–77.

Li, H. (2018). Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* , *34*(18), 3094–

3100.

Lovell, J. T., Sreedasyam, A., Schranz, M. E., Wilson, M., Carlson, J. W., Harkess, A., Emms, D.,

Goodstein, D. M., & Schmutz, J. (2022). GENESPACE tracks regions of interest and gene copy

number variation across multiple genomes. *eLife*, *11*. https://doi.org/10.7554/eLife.78526

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. *Genome Biology*, *15*(12), 550.

Lowry, D. B., Cotton Rockwood, R., & Willis, J. H. (2008). Ecological reproductive isolation of coast

and inland races of Mimulus guttatus. In *Evolution* (Vol. 62, Issue 9, pp. 2196–2214).

https://doi.org/10.1111/j.1558-5646.2008.00457.x

Lowry, D. B., Popovic, D., Brennan, D. J., & Holeski, L. M. (2019). Mechanisms of a locally adaptive

shift in allocation among growth, reproduction, and herbivore resistance in Mimulus guttatus.

Evolution, *73*(6), 1168–1181.

Lowry, D. B., Sheng, C. C., Lasky, J. R., & Willis, J. H. (2012). Five anthocyanin polymorphisms are

associated with an R2R3-MYB cluster in Mimulus guttatus (Phrymaceae). *American Journal of*

Botany, *99*(1), 82–91.

Lowry, D. B., & Willis, J. H. (2010). A widespread chromosomal inversion polymorphism contributes to

a major life-history transition, local adaptation, and reproductive isolation. *PLoS Biology*, *8*(9).

https://doi.org/10.1371/journal.pbio.1000500

Lundberg, M., Mackintosh, A., Petri, A., & Bensch, S. (2023). Inversions maintain differences between

migratory phenotypes of a songbird. *Nature Communications*, *14*(1), 452.

Lupski, J. R. (1998). Genomic disorders: structural features of the genome can lead to DNA

rearrangements and human disease traits. *Trends in Genetics: TIG*, *14*(10), 417–422.

Lu, Z., Hofmeister, B. T., Vollmers, C., DuBois, R. M., & Schmitz, R. J. (2017). Combining ATAC-seq

with nuclei sorting for discovery of cis-regulatory regions in plant genomes. *Nucleic Acids Research*,

45(6), e41–e41.

Manetas, Y. (2006). Why some leaves are anthocyanic and why most anthocyanic leaves are red? *Flora -*

Morphology, Distribution, Functional Ecology of Plants, *201*(3), 163–177.

Marquès-Bonet, T., Cáceres, M., Bertranpetit, J., Preuss, T. M., Thomas, J. W., & Navarro, A. (2004).

Chromosomal rearrangements and the genomic distribution of gene-expression divergence in

humans and chimpanzees. *Trends in Genetics: TIG*, *20*(11), 524–529.

Marshall, C. R., Noor, A., Vincent, J. B., Lionel, A. C., Feuk, L., Skaug, J., Shago, M., Moessner, R.,

Pinto, D., Ren, Y., Thiruvahindrapduram, B., Fiebig, A., Schreiber, S., Friedman, J., Ketelaars, C. E.

J., Vos, Y. J., Ficicioglu, C., Kirkpatrick, S., Nicolson, R., … Scherer, S. W. (2008). Structural

variation of chromosomes in autism spectrum disorder. *American Journal of Human Genetics*, *82*(2),

477–488.

McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A., Garimella, K.,

Altshuler, D., Gabriel, S., Daly, M., & DePristo, M. A. (2010). The Genome Analysis Toolkit: a

MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Research*,

20(9), 1297–1303.

Mérot, C., Llaurens, V., Normandeau, E., Bernatchez, L., & Wellenreuther, M. (2020). Balancing

selection via life-history trade-offs maintains an inversion polymorphism in a seaweed fly. *Nature*

Communications, *11*(1), 670.

Mroczek, R. J., Melo, J. R., Luce, A. C., Hiatt, E. N., & Dawe, R. K. (2006). The maize Ab10 meiotic

drive system maps to supernumerary sequences in a large complex haplotype. *Genetics*, *174*(1), 145–

154.

Muller, H. J. (1930). Types of visible variations induced by X-rays in Drosophila. *Journal of Genetics*,

22, 299–334.

Navarro, A., & Barton, N. H. (2003). Chromosomal speciation and molecular divergence--accelerated

evolution in rearranged chromosomes. *Science*, *300*(5617), 321–324.

Navarro, A., Betrán, E., Barbadilla, A., & Ruiz, A. (1997). Recombination and gene flux caused by gene

conversion and crossing over in inversion heterokaryotypes. *Genetics*, *146*(2), 695–709.

Newman, T. L., Tuzun, E., Morrison, V. A., Hayden, K. E., Ventura, M., McGrath, S. D., Rocchi, M., &

Eichler, E. E. (2005). A genome-wide survey of structural variation between human and chimpanzee.

Genome Research, *15*(10), 1344–1356.

O'Donnell, S., & Fischer, G. (2020). MUM&Co: accurate detection of all SV types through whole-

genome alignment. *Bioinformatics* , *36*(10), 3242–3243.

Olito, C., & Abbott, J. K. (2023). The evolution of suppressed recombination between sex chromosomes

and the lengths of evolutionary strata. *Evolution; International Journal of Organic Evolution*, *77*(4),

1077–1090.

Oneal, E., Lowry, D. B., Wright, K. M., Zhu, Z., & Willis, J. H. (2014). Divergent population structure

and climate associations of a chromosomal inversion polymorphism across the Mimulus guttatus

species complex. *Molecular Ecology*, *23*(11), 2844–2860.

Ou, S., Chen, J., & Jiang, N. (2018). Assessing genome assembly quality using the LTR Assembly Index

(LAI). *Nucleic Acids Research*, *46*(21), e126.

Pertea, G., & Pertea, M. (2020). GFF Utilities: GffRead and GffCompare. *F1000Research*, *9*.

https://doi.org/10.12688/f1000research.23297.2

Pfaffl, M. W. (2001). A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic*

Acids Research, *29*(9), e45.

Phytozome v13. (n.d.). Retrieved December 4, 2023, from

https://phytozome.jgi.doe.gov/info/Mguttatusvar_IM767_v1_1

Puig, M., Casillas, S., Villatoro, S., & Cáceres, M. (2015). Human inversions and their functional

consequences. *Briefings in Functional Genomics*, *14*(5), 369–379.

Pyhäjärvi, T., Hufford, M. B., Mezmouk, S., & Ross-Ibarra, J. (2013). Complex patterns of local

adaptation in teosinte. *Genome Biology and Evolution*, *5*(9), 1594–1609.

Qin, M., Wu, S., Li, A., Zhao, F., Feng, H., Ding, L., & Ruan, J. (2019). LRScaf: improving draft

genomes using long noisy reads. *BMC Genomics*, *20*(1), 955.

Quinlan, A. R., & Hall, I. M. (2010). BEDTools: a flexible suite of utilities for comparing genomic

features. *Bioinformatics* , *26*(6), 841–842.

Rane, R. V., Rako, L., Kapun, M., Lee, S. F., & Hoffmann, A. A. (2015). Genomic evidence for role of

inversion 3RP of Drosophila melanogaster in facilitating climate change adaptation. *Molecular*

Ecology, *24*(10), 2423–2432.

RepeatMasker Home Page. (n.d.). Retrieved March 13, 2023, from http://www.repeatmasker.org/

Rieseberg, L. H. (2001). Chromosomal rearrangements and speciation. *Trends in Ecology & Evolution*,

16(7), 351–358.

Roze, D., & Barton, N. H. (2006). The Hill–Robertson Effect and the Evolution of Recombination.

Genetics, *173*(3), 1793–1811.

Said, I., Byrne, A., Serrano, V., Cardeno, C., Vollmers, C., & Corbett-Detig, R. (2018). Linked genetic

variation and not genome structure causes widespread differential expression associated with

chromosomal inversions. *Proceedings of the National Academy of Sciences of the United States of*

America, *115*(21), 5492–5497.

Schaefer, H. M., & Rolshausen, G. (2006). Plants on red alert: do insects pay attention? *BioEssays: News*

and Reviews in Molecular, Cellular and Developmental Biology, *28*(1), 65–71.

Schneider, M., Bairoch, A., Wu, C. H., & Apweiler, R. (2005). Plant protein annotation in the UniProt

Knowledgebase. *Plant Physiology*, *138*(1), 59–66.

Schwander, T., Libbrecht, R., & Keller, L. (2014). Supergenes and complex phenotypes. *Current*

Biology: CB, *24*(7), R288–R294.

Shumate, A., & Salzberg, S. L. (2020). Liftoff: accurate mapping of gene annotations. *Bioinformatics* ,

37(12), 1639–1643.

Simão, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V., & Zdobnov, E. M. (2015). BUSCO:

assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics*

, *31*(19), 3210–3212.

Slater, G. S. C., & Birney, E. (2005). Automated generation of heuristics for biological sequence

comparison. *BMC Bioinformatics*, *6*, 31.

Stanke, M., & Morgenstern, B. (2005). AUGUSTUS: a web server for gene prediction in eukaryotes that

allows user-defined constraints. *Nucleic Acids Research*, *33*(Web Server issue), W465–W467.

Tang, H., Zhang, X., Miao, C., Zhang, J., Ming, R., Schnable, J. C., Schnable, P. S., Lyons, E., & Lu, J.

(2015). ALLMAPS: robust scaffold ordering based on multiple maps. *Genome Biology*, *16*(1), 3.

Thompson, M. J., & Jiggins, C. D. (2014). Supergenes and their role in evolution. *Heredity*, *113*(1), 1–8.

Todesco, M., Owens, G. L., Bercovich, N., Légaré, J.-S., Soudi, S., Burge, D. O., Huang, K., Ostevik, K.

L., Drummond, E. B. M., Imerovski, I., Lande, K., Pascual-Robles, M. A., Nanavati, M., Jahani, M.,

Cheung, W., Staton, S. E., Muños, S., Nielsen, R., Donovan, L. A., … Rieseberg, L. H. (2020).

Massive haplotypes underlie ecotypic differentiation in sunflowers. *Nature*, *584*(7822), 602–607.

Tuttle, E. M., Bergland, A. O., Korody, M. L., Brewer, M. S., Newhouse, D. J., Minx, P., Stager, M.,

Betuel, A., Cheviron, Z. A., Warren, W. C., Gonser, R. A., & Balakrishnan, C. N. (2016).

Divergence and Functional Degradation of a Sex Chromosome-like Supergene. *Current Biology:*

CB, *26*(3), 344–350.

Twyford, A. D., & Friedman, J. (2015). Adaptive divergence in the monkey flower Mimulus guttatus is

maintained by a chromosomal inversion. *Evolution; International Journal of Organic Evolution*,

69(6), 1476–1486.

Vaser, R., Sović, I., Nagarajan, N., & Šikić, M. (2017). Fast and accurate de novo genome assembly from

983 long uncorrected reads. *Genome Research*, *27*(5), 737–746.

984 Vasimuddin, M., Misra, S., Li, H., & Aluru, S. (2019). Efficient Architecture-Aware Acceleration of

985 BWA-MEM for Multicore Systems. *2019 IEEE International Parallel and Distributed Processing*

986 *Symposium (IPDPS)*, 314–324.

987 Villoutreix, R., Ayala, D., Joron, M., Gompert, Z., Feder, J. L., & Nosil, P. (2021). Inversion breakpoints

988 and the evolution of supergenes. *Molecular Ecology*, *30*(12), 2738–2755.

989 Vogel, M. J., Pagie, L., Talhout, W., Nieuwland, M., Kerkhoven, R. M., & van Steensel, B. (2009). High-

990 resolution mapping of heterochromatin redistribution in a Drosophila position-effect variegation

991 model. *Epigenetics & Chromatin*, *2*(1), 1.

992 Walker, B. J., Abeel, T., Shea, T., Priest, M., Abouelliel, A., Sakthikumar, S., Cuomo, C. A., Zeng, Q.,

993 Wortman, J., Young, S. K., & Earl, A. M. (2014). Pilon: an integrated tool for comprehensive

994 microbial variant detection and genome assembly improvement. *PloS One*, *9*(11), e112963.

995 Wang, K., Li, M., & Hakonarson, H. (2010). ANNOVAR: functional annotation of genetic variants from

996 high-throughput sequencing data. *Nucleic Acids Research*, *38*(16), e164.

997 Wong, E. L. Y., & Filatov, D. A. (2023). The role of recombination landscape in species hybridisation

998 and speciation. *Frontiers in Plant Science*, *14*, 1223148.

999 Zhang, Z., Zhu, H., Gill, B. S., & Li, W. (2015). Fine mapping of shattering locus Br2 reveals a putative

1000 chromosomal inversion polymorphism between the two lineages of Aegilops tauschii. *TAG.*

1001 *Theoretical and Applied Genetics. Theoretische Und Angewandte Genetik*, *128*(4), 745–755.

1002 Zhou, H., Ma, R., Gao, L., Zhang, J., Zhang, A., Zhang, X., Ren, F., Zhang, W., Liao, L., Yang, Q., Xu,

1003 S., Otieno Ogutu, C., Zhao, J., Yu, M., Jiang, Q., Korban, S. S., & Han, Y. (2021). A 1.7-Mb

1004 chromosomal inversion downstream of a PpOFP1 gene is responsible for flat fruit shape in peach.

1005 *Plant Biotechnology Journal*, *19*(1), 192–205.

1006

1007

1008

Figures and Tables (In text)

Figure 1. A. Whole genome alignment of all 14 chromosomes between the SWB-S1 and LMC-L1 genomes. Large inversions on chromosomes 5, 8, and 14 are visible in the dotplots, where the teal dots represent the negatively oriented strands while orange represent positively oriented strands. **B.** The inv_chr8B captured within inv_chr8A.

Figure 2. *G* statistic and π ratio for each of the inversions (salmon) compared to the non-inverted (teal) sequence of the same chromosome when aligned to the LMC-L1 Genome **A** and to the SWB-S1 genome **B**.

Reference genome	Location	π IA	π CP	π ratio	G statistic
SWB-S1	Colinear regions of chromosome 5	0.040	0.032	1.26	9.98
SWB-S1	inv_chr5A	0.040	0.029	1.41	20.28
SWB-S1	Colinear regions of chromosome 8	0.040	0.032	1.25	9.90
SWB-S1	inv_chr8A	0.041	0.028	1.48	20.07
SWB-S1	inv_chr8A (without inv_chr8B)	0.041	0.028	1.46	19.57
SWB-S1	inv_chr8B	0.042	0.021	2.02	32.88
SWB-S1	Colinear regions of chromosome 14	0.040	0.032	1.26	10.16
SWB-S1	inv_chr14A	0.041	0.033	1.25	10.09
SWB-S1	genome-wide	0.040	0.032	1.26	10.16
LMC-L1	Colinear regions of chromosome 5	0.043	0.037	1.18	8.19
LMC-L1	inv_chr5A	0.040	0.031	1.28	20.05
LMC-L1	Colinear regions of chromosome 8	0.040	0.033	1.21	10.70
LMC-L1	inv_chr8A	0.040	0.029	1.37	19.64
LMC-L1	inv_chr8A (without inv_chr8B)	0.040	0.030	1.35	18.71
LMC-L1	inv_chr8B	0.042	0.021	1.93	35.76
LMC-L1	Colinear regions of chromosome 14	0.039	0.031	1.23	11.03
LMC-L1	inv_chr14A	0.041	0.034	1.20	9.83
LMC-L1	genome-wide	0.040	0.034	1.20	10.01

Table 1. Comparison of π and *G* statistic across the genome and for each inversion. IA represents the inland annual pool and CP represents the coastal perennial pool.

Genotype	Location	4-fold degenerate SNPs	4-fold degenerate sites	4-fold degenerate SNPs / 4-fold degenerate sites
LMC-L1	Genome-wide	174,745	4,106,234	0.043
LMC-L1	inv_5A	4,497	81,202	0.055
LMC-L1	inv_8A	8,030	139,941	0.057
LMC-L1	inv_8B	636	7,569	0.084
LMC-L1	inv_14A	1,634	39,641	0.041
SWB-S1	Genome-wide	176,470	4,093,796	0.043
SWB-S1	inv_5A	4,772	78,850	0.061
SWB-S1	inv_8A	7,954	136,834	0.058
SWB-S1	inv_8B	630	7,633	0.083
SWB-S1	inv_14A	1,555	42,095	0.037

Table 2. LMC-L1 and SWB-S1 SNPs called from WGS were annotated and filtered to include synonymous SNPs at 4-fold degenerate sites for each inversion and genome-wide. Note SNPs for LMC-L1 were called after aligned WGS data to the SWB-S1 genome and SWB-S1 SNPs were called after aligned WGS data to the LMC-L1 genome.

Figure 3. Quantitative RT-PCR showing expression of *GA20ox2* in wild-type LMC-L1 and SWB-S1 floral shoot apex (three biological replicates). Error bars show standard deviation.

Figure 4. The MYB region along chromosome 8 in LMC-L1 (**A**) and SWB-S1 (**B**). Each arrow (teal in LMC-L1 and orange in SWB-S1) represents a gene and orientation. The gray arrows represent pseudogenes.

Figure 5. A. Differential expression represented in transcripts per million (TPM) in genes at the inversion breakpoints. Note, some plots are nearly identical because they are LMC-L1 (labeled LMC) and SWB-S1 (labeled SWB) orthologs (MgL14g10910 and MgS1_14g08070,

MgS1_05g10260 and MgL1_05g16120). **B.** A diagram depicting the inversion breakpoint hypothesis.