Hyper-specialized bamboo lemurs possess a reduced suite of xenobiotic-metabolizing

cytochrome P450 genes.

Morgan E. Chaney*[1,2], Christina M. Bergey†[2], Anthony J. Tosi†[1]

*: corresponding author, email: morgan.chaney@rutgers.edu

†: these authors contributed equally to this manuscript

1: Department of Anthropology & School of Biomedical Sciences, Kent State University, Kent,

Ohio 44242 USA

2: Department of Genetics, Rutgers, The State University of New Jersey, Piscataway, New Jersey

08854 USA

Introduction

Among mammals, a large proportion of xenobiotic compounds are transformed or

detoxified by enzymes within the cytochrome P450 superfamily. Because of this role that P450s

play in mediating potential environmental toxins and because species vary in their diets and

exposure to such compounds, selective pressure has induced non-random additive and

adaptive changes in genetic sequence and copy number among species (Chaney et al., 2018,

2020; Ingelman-Sundberg, 2022; Kawashima & Satta, 2014; Kirischian et al., 2011; Skopec et al.,

2022). The effect of evolving toxin resistance has been especially well-documented for pest

insect species because of the economic importance of pesticides and insects' evolutionary

responses to those substances (Wondji et al. 2009; Feyereisen 2011; Bass et al. 2014; Wang et

al. 2014; Walsh et al. 2018; Joussen and Heckel 2021). Studies of P450 gene adaptation outside

of pest insects have often focused on dietary xenobiotics or plant secondary metabolites

(PSMs). For example, research on a model system comprising several species of woodrat

(*Neotoma* spp.) has shown that PSMs common in their diet (*e.g.*, alkaloids, terpenoids) are

metabolized especially by CYP2B enzymes and, to a minor extent, other xenobiotic-

metabolizing enzymes (Haley et al., 2007a, 2007b; Orr et al., 2020; Wilderman et al., 2014).

Comparisons between specialist and generalist species of these woodrats have shed

light on the effects that diet can have on P450 evolution. For instance, the generalist woodrat

species (*N. albigula*) was found to have lower copy number but greater sequence diversity of

CYP2B relative to a specialist (*N. stephensi*). This greater diversity of the generalist is consistent

with the broader array of PSMs with which the generalist must contend (Kitanovic et al., 2018).

The increased copy number of the specialist, in contrast, is in line with the prediction that genes

important for coping with the specialist's narrow diet will be subject to *additive* evolutionary

processes (*e.g.*, gene duplication, positive-selection) that will enhance the species' ability to

mitigate a narrow range of metabolically harmful compounds (Dearing et al., 2005; Freeland &

Janzen, 1974). This is taken to an extreme for specialized koala (*Phascolartcos cinereus*), in

which the key xenobiotic metabolizing *CYP2C* subfamily has reached a copy number of 31 via

serial duplication (Johnson et al., 2018).

38   Mammalian dietary specialization has also been linked to patterns of gene loss (Blumer

et al., 2022; Hecker et al., 2019; Itoigawa et al., 2019; Lauterbur, 2019; Marciniak et al., 2021;

Wagner et al., 2022). For example, vampire bats (genera *Desmodus, Diaemus,* and *Diphylla*)

have lost sweet and umami taste receptors (Zhao et al., 2010b; Zhao et al., 2012) and have

experienced reduction of their bitter-taste sense through pseudogenization of *TAS2R* genes

(Hong & Zhao, 2014). Interestingly, losses of gustatory genes have also been found for

cetaceans, which have lost nearly all taste-receptor genes apart from those for salty taste (Feng

et al., 2014), and for pandas (*Ailuropoda melanoleuca*), which have lost their umami taste-

receptors (Zhao et al., 2010a). Theory predicts that genes unimportant to a specialist's dietary

adaptation will be subject to a suite of *subtractive* evolutionary processes (*e.g.,* accumulation of

deleterious mutations, pseudogenization, or loss of protein coding region or entire gene). Gene

loss could either be for adaptive reasons (see Olson, 1999) or because of a release of selective

pressure (see Blumer et al., 2022; Huelsmann et al., 2019). Despite these examples and the

theory surrounding them, subtractive mechanisms such as gene loss have remained relatively

understudied compared to additive mechanisms such as gene duplication and positive or

directional selection.

In summary, specialized folivory is predicted to have two evolutionary effects on a

species' xenobiotic-metabolizing toolkit: First, genes involved in detoxification or avoidance of

xenobiotic toxins that are rich in a folivorous specialist's diet should show signals of selection in

the form of positive copy-number changes or positive selection. Second, genes *not* involved in

such species' defense against toxins in their specialized diet should exhibit signals of negative

copy-number changes and relaxation of selection. Although the first prediction is well-

supported by prior studies of P450 gene evolution, comparatively few studies have investigated

patterns of P450 gene loss or relaxation of selection in specialized folivores.

Bamboo lemurs are an ideal system for testing predictions about P450 gene loss in

specialized folivores due to their narrow diets, which are among the most specialized in the

entire primate order. Despite their diminutive body sizes (0.9-2.6 kg: Tan, 2006), they inhabit an

ecological niche that is often compared with that of the giant panda (Eronen et al., 2017;

McKenney et al., 2017; Tan, 2006) and are the only extant primates known to specialize so

intensely on bamboo or grass species (Table S1). The three sympatric species at Ranomafana

National Park (*i.e.*, *Hapalemur aureus, H. griseus*, and *Prolemur simus*) are the source of much

of the current understanding of bamboo lemur ecology. All three of these populations have

diets that are often predominantly composed of a small number of species of bamboo or liana

(Grassi, 2006; Overdorff et al., 1997; Tan, 1999), and the sympatry of these species has been

understood in terms of niche partitioning, notably by their differential reliance on separate

plant parts from the same species of giant bamboo (Glander et al., 1989). Although other

species outside of Ranomafana (*e.g.*, *H. meridionalis, H. alaotrensis*) do not actually consume

bamboo, they nonetheless eat grasses (Poaceae) or sedges (Cyperaceae) at similarly high

frequencies (Table S1). Thus, regardless of habitat or species, bamboo lemurs subsist on

constrained, highly folivorous diets.

One bamboo PSM is particularly intriguing for its toxicity. Consumption of the specific

bamboo parts consumed by *H. aureus* and *Prol. simus* exposes these lemurs to high levels of

potent cyanogenic glycosides (Ballhorn et al., 2009; Eppley et al., 2017; Glander et al., 1989).

When plant tissue is disrupted by some destructive process like mastication, these glycosides

will be hydrolyzed by the plant cell's own β-glycosidase enzymes to release cyanide (Ballhorn,

2011; Cressey & Reeve, 2019). At Ranomafana, two bamboo lemur species have been

estimated to consume many times the estimated lethal dose of cyanide for mammals of their

body sizes (Ballhorn et al., 2009; Glander et al., 1989). Although cyanide, the primary known

antifeedant of this bamboo, is detoxified by a metabolic pathway that does not include any

cytochrome P450 enzymes (reviewed by Isom et al., 2015), bamboo lemurs likely have to

contend with other PSMs known to be present in bamboo species, such as phenolic and

polyphenolic compounds (Zhu et al. 2018). Thus, we still expect evolution of bamboo lemur

P450 genes to be impacted by their hyper-specialized diet.

For this project, we hypothesized that the highly specialized diet observed for bamboo

lemurs should be associated with a narrower P450 repertoire, streamlined to those subfamilies

that metabolize xenobiotic compounds found in bamboo. Therefore, we predicted that,

compared to other generalist Malagasy strepsirrhines, the bamboo lemur species *Prol. simus*

and *H. griseus* should exhibit a clear pattern of cytochrome P450 (1) gene loss and (2)

signatures of relaxed natural selection, because the specialist diet contains a narrow set of

PSMs, of which the primary toxin (*i.e.*, cyanide), is not even metabolized by P450s. Similar to

previous work (Chaney et al., 2020), we focus on P450 genes in the *CYP1-3* subfamilies. Briefly,

this decision was made because of these subfamilies' monophyletic placement within the

cytochrome P450 superfamily of genes (Kirischian et al., 2011) and the presence of genes

encoding both xenobiotic and endobiotic metabolizers within that gene-clade (see Guengerich,

2015).

Materials and Methods

<u>Generation of *Hapalemur griseus* genome assembly</u>

For this study, we constructed a genome assembly for a male *H. griseus* (named Beavis)

housed at the Duke Lemur Center in Durham, North Carolina. A fresh whole blood sample was

shipped to the Molecular Primatology Laboratory of New York University for processing and

sequencing. High molecular weight (HMW) DNA was extracted from 200 µL of whole blood

using a MagAttract HMW DNA Kit (QIAGEN Sciences Inc., Germantown, MD) following the

manufacturer's protocol. DNA was eluted into a low-bind 1.5 mL tube and stored at 4°C prior

to size selection. In order to obtain DNA fragments roughly 40 kilobases or longer, the extracted

HMW DNA was run through a Blue Pippin DNA Size Selection System (Sage Science Inc., Beverly,

MA) using a 0.75% agarose gel cassette with the R1 size standard. Following this size selection

protocol, the HMW DNA concentration was quantified using a Qubit 2.0 fluorometer (Thermo

Fisher Scientific Inc., Carlsbad, CA) using the broad range standard. The extraction yielded 3.2

ng/µL of HMW DNA.

After extraction and quantification, the HMW DNA sample was sent to NYU Langone

Medical School's Genome Technology Center for quality control tests, which consisted of DNA

quantification and fragment size determination using a Tape Station (Agilent Technologies Inc.,

Santa Clara, CA). After passing this checkpoint, the HMW DNA was prepared for sequencing

using a Chromium Genome Linked Reads Kit (10X Genomics Inc., Pleasanton, CA) and

sequenced on a HiSeq 2500 DNA Sequencing Platform (Illumina Inc., San Diego, CA). The linked-

read output data were assembled using the Supernova assembler (Weisenfeld et al., 2017). We

used the pseudohaplotype output as our genome assembly for all downstream steps (Table 1).

<u>Additional data gathering</u>

In addition to the above-mentioned assembly for *H. griseus*, we mined data from

publicly available genome assemblies for 14 species: *Prolemur simus* (Hawkins et al., 2018),

Lemur catta (Palmada-Flores et al., 2022), *Eulemur flavifrons* and *E. macaco* (Meyer et al. 2015),

Propithecus coquereli (Lowe and Eddy 1997; Guevara et al. 2021), *Indri indri* (accession number:

GCA_004363605.1), *Daubentonia madagascariensis* (accession number: GCA_004027145.1),

Mirza coquereli (accession number: GCA_004024645.1), *Mirza zaza* (Hunnicutt et al., 2020)*,* and

Microcebus murinus (Averdam et al., 2011; Lecompte et al., 2016), as well as the following

additional species of mouse lemur: *Mic. griseorufus, Mic. mittermeieri, Mic. ravelobensis,* and

Mic. tavaratra (Hunnicutt et al., 2020). These assemblies, along with any associated annotation

files, were downloaded locally and formatted into BLAST databases within Geneious Prime,

version 2022.1.1.

We located the loci for all annotated *CYP1-3* homologs in the *L. catta, Prop. coquereli,*

and *Mic. murinus* by using the associated annotation (GFF3) files for each. We defined these

loci by the non-P450 genes that bounded them; therefore, those surrounding genes were used

initially as queries for local BLAST searches. In this way, each locus was linked to two searches

per species. The three reference genomes listed above were used because they are all

members of separate strepsirrhine families (Lemuridae, Indriidae, and Cheirogaleidae,

respectively), and they were each therefore used as a starting point to extract the desired

CYP1-3 genes or loci for confamilial species. Ideally, a pair of BLAST searches would return

results that included the same scaffold. By locating both BLAST hits on each of these scaffolds,

we were able to extract genomic regions that were hypothetically orthologous to those P450

loci in the *L. catta* assembly. After locating the scaffolds in each assembly corresponding to

each P450 locus, we used LASTZ (Harris, 2007) to interrogate the homology of those scaffolds

by aligning them to the confirmed P450 locus from the appropriate confamilial reference

genome. Positive results from these alignments were checked using the Mauve genome aligner

(Darling et al., 2004) on the same sequences. If output from both of these aligners indicated

that the reference had homology with the query scaffold(s), then the annotations from the

reference genome were used to extract the corresponding sequence in the other species'

genome. In this way, we mined the genome assemblies listed above for as many complete P450

genes loci as we could confidently locate.

<u>Inference of gene birth and death</u>

For this first portion of the study, we used only species for which each *CYP1-3* locus

could be wholly collected from a single scaffold or reasonably reconstructed if not found on a

single scaffold using the process described above. In order to model the events of gene birth

and death in this subset of lemur species, our alignment strategy followed a similar workflow as

outlined in previous work with other datasets (Chaney et al., 2018, 2020), but several

modifications were made for this project in order to allow for more standardization and

automation across subfamilies. First, all of the P450 genes were extracted from each species'

locus according to the annotation file associated with its confamilial reference. Then, all of the

genes from a given P450 subfamily were aligned using MAFFT (Katoh & Standley, 2013), and the

resulting alignment was stripped of all sites containing any gaps using trimAl (Capella-Gutierrez

et al., 2009). After the best-fitting nucleotide substitution model was inferred by jModelTest

(Darriba et al., 2012), this stripped alignment was visualized with PhyML 3.0 and the strength of

that resulting phylogenetic tree was tested by comparing it to 1000 bootstrap replicates

(Guindon et al., 2010).

The gene trees constructed with PhyML were then passed to Possvm (Grau-Bové &

Sebé-Pedrós, 2021). This program uses the intrinsic information contained in a phylogram to

infer speciation and gene-duplication events; it does this using the species-overlap algorithm in

the ETE3 toolkit (Huerta-Cepas et al., 2007, 2016). Briefly, this algorithm compares the

intersection of species present in both descendants of an internal node of a tree to the union of

species present in those descendants; using these values, the algorithm computes a species-

overlap score which it then uses to identify each internal node as either a speciation event,

having an overlap score, or a duplication event, having a high overlap score (Huerta-Cepas et

al., 2007). Once the identities of each node were estimated in this way, we then manually

examined each subtree rooted by a node called as a duplication event to infer whether any

gene loss had occurred. This was examined on a case-by-case basis using the reasoning that,

after a duplication event, each descendant of that node should recapitulate the organismal

phylogeny present at the time of duplication. Therefore, any species missing in one of those

subtrees must have lost one of the duplicates born in the earlier duplication event as long as

the subtree in question was well-resolved in terms of bootstrap support. In cases where

multiple species lineages may be absent, we deferred to the parsimonious hypothesis that a

loss event would have occurred prior to the divergence of those lineages, rather than a more

complicated hypothesis that the same paralog had been independently lost in both species

after their split. We visualized the Possvm output using the program Treerecs (Comte et al.,

2020) and then, in some cases, manually modified the depicted gene-evolution scenario in

order to accommodate the Possvm results.

Selection analysis

We used RELAX (Wertheim et al., 2015) to test our hypothesis concerning the relaxing of

selectional pressure on the P450s covered in this project. Each run of this program requires,

first, a codon alignment stripped of any stop codons and, second, a phylogenetic tree that

contains labels for the branches on which *a priori* relaxation of natural selection is hypothesized

to have occurred. For the codon alignments, we used preexisting annotations from each of the

annotated genome assemblies to extract the likely coding DNA sequence (CDS) from each of

the ostensibly complete P450 genes gathered in the first portion of the study. Those

annotations were then leveraged to find the likely CDS in unannotated homologous sequences

from confamilial relatives.

We aligned the CDS from each subfamily in Geneious Prime by translation using the

MAFFT aligner (Katoh & Standley, 2013), and we then checked each alignment by eye. Once the

alignment had been checked by eye and the maximum number of useable sequences were

included, it was exported in FASTA format. The trees used for all runs of RELAX were adapted

from the phylogenies constructed in the earlier part of this study concerned with gene birth

and death. We compared the rows in each subfamily's final CDS alignment to the leaves of the

gene tree constructed earlier with an R script using functions from the *phytools* (Revell, 2012)

and *seqinr* (Charif & Lobry, 2007) R packages, and a new tree was written that contained only

the intersection of these two sets. This filtered gene tree was then viewed with the web-based

phylotree.js platform (Shank et al., 2018), which allowed us to easily label branches on each

tree. For subfamilies where no duplications had been detected in the first part of this study

(*e.g.*, *CYP1B*), only foreground branches were labeled for *Prol. simus, H. griseus*, and any

internal branches exclusive to the clade formed by these two species. In these cases, all

unlabeled branches were used for the estimation of the background *dN/dS* ratios. However, we

followed a more complicated, yet still conservative, strategy when dealing with subfamilies

where multiple orthogroups were detected by Possvm (*e.g.*, *CYP2A*). For these subfamilies, all

non-bamboo lemur branches were marked as background branches except for the deepest

internal branches that separate orthogroups, which were left unlabeled. This allowed us to

exclude those deep portions of the phylogeny, which we imagined may have coincided with

instances of past neofunctionalization following gene duplication. If true, this effect would

probably bias the *dN/dS* distribution of the background branches; thus, we excluded those deep

branches from analysis in the relevant cases. Aside from manually reading the text-file output,

the HyPhy Vision platform (http://vision.hyphy.org/RELAX) was used to generate some of the

graphs made from the output JSON files.

Finally, we performed follow-up tests for diversifying selection for one gene, *CYP2F*, as

initial analyses with RELAX revealed unexpected signatures of selection intensification. We first

used BUSTED (Murrell et al., 2015) as a confirmatory check for any evidence of

diversifying/positive selection in the input alignment, followed by aBSREL (Smith et al., 2018), in

conjunction with the branch-labeling workflow described above, to test for diversifying/positive

selection specifically in the bamboo-lemur branches of the gene tree.

Results

The evidence presented here largely displays a trend in which the common lineage of

bamboo lemurs has a diminished or expected number of gene copies relative to generalist

species, with several xenobiotic-metabolizing P450 subfamilies exhibiting a resolved gene-tree

topology indicating genic loss in one or both of the bamboo lemur species included here.

Because of the other species included in this data set, there are also some insights to be gained

from examining other portions of the gene trees outside of the portions focused on our species

of interest.

<u>Heightened gene loss within the bamboo-lemur lineage</u>

The topologies of the *CYP1A* and *CYP1B* subfamilies show strong bootstrap support and

near-perfect correspondence to the species phylogeny for their member orthologs (Figs. S1 &

S2). The only exception to this derives from the *Propithecus coquereli* lineage, which saw the

duplication of the *CYP1A1* ortholog after the divergence of the *Prop. coquereli* and *Indri indri*

lineages*.*

The shape of the *CYP2A* gene tree implies substantially more genic turnover than either

of the *CYP1* subfamilies (Fig. 1, top-left; Fig. S3). While the *Microcebus murinus* lineage shows a

very straightforward trajectory of no duplications, all other species have either duplicated or

lost orthologs according to the pattern of relatedness displayed in this tree. Due to the mid-

point rooting of this tree by Possvm, it is unclear whether two or three orthogroups are truly

present in this case (see Fig. S3), but we opt for a parsimonious interpretation of two

orthogroups here (see Fig. 1). . Nonetheless, at least one duplication was certainly specific to

the *Prop. coquereli* lineage, depicted here by the common node of *CYP2A13L1* and *CYP2A13L2*.

The lemurid portion of the tree indicates a gene duplication in the common ancestor of all

three species and another duplication that was specific to *L. catta*. Conversely, the topology

displayed here implies genic loss for *E. macaco* and for *Prol. simus.*

The only members of *CYP2B* examined in this study are *CYP2B6* homologs because

another group of orthologs in the *L. catta* genome assembly (*CYP2B11*) did not show a CDS

length that was remotely close to the normal length of a functional primate P450 gene (*i.e.*, *ca.*

1500 nt). In the *CYP2B6* tree (Fig. 1, bottom-left; Fig. S4), we find two major orthogroups among

the five species analyzed. One of these shows a topology that, in itself, appears to recapitulate

the organismal phylogeny with no duplications. However, the other orthogroup exhibits a

signature of interesting dynamism. There are two serial duplication events in the *Prop.*

coquereli lineage (*CYP2B6L1, 3,* and *4P*). In the lemurid portion of this subtree, there are two

more serial duplications prior to the speciation of *L. catta* and *Prol. simus.* The pattern in this

part of the tree most parsimoniously indicates, first, that the bamboo-lemur common ancestor

lost genes orthologous to the *CYP2B6L3* and *CYP2B6L4* genes in the *L. catta* assembly and,

second, that the *H. griseus* lineage further lost its final remaining gene in this orthogroup.

The *CYP2C* genes yielded a phylogeny whose complexity is underscored by the large

number of duplications within it (Fig. 1, right; Fig. S5). Possvm separated this phylogeny into

five coherent orthogroups. The first of these (*i.e.,* OG-0) contained a duplication prior to the

divergence of *L. catta* and the bamboo lemurs. Following the *Prolemur-Hapalemur* divergence,

Prol. simus lost one of these genes but the other paralog was lost by *H. griseus*; thus, both

species retained separate paralogs. OG-1 and OG-2 lack *Prol. simus* genes but do contain genes

belonging to *H. griseus*; this pattern indicates two lineage specific events of gene loss for *Prol.*

simus. While this latter species, the greater bamboo lemur, showed a pattern of complete loss

for these orthogroups, *Mic. murinus* experienced a duplication in OG-2. All lemurid genes were

ostensibly lost from OG-3, while the mouse-lemur lineage saw a single-gene expansion of this

orthogroup. The final clade of genes, OG-4, contained three duplication events and,

parsimoniously, three instances of gene loss. While *L. catta* retained every one of the

duplicated genes from this sequence of events, bamboo lemurs likely lost the ortholog to *L.*

catta CYP2C18L1, and both *Prol. simus* and *H. griseus* further lost one more gene each.

However, these two final losses were lineage-specific, and each involved a separate genic

lineage. Thus, with the exception of genes in one orthogroup (*i.e.*, OG-3), bamboo lemurs

exhibited all of the losses detected in this tree.

Similar to *CYP2C*, the *CYP2D* subfamily also showed an increased amount of evolutionary

dynamism with regard to gene birth and death (Fig. 2, top-left; Fig. S6). Along a basal branch of

the phylogeny, there are two serial duplications, resulting in three *Mic. murinus* homologs

equally related to all lemurid *CYP2D* homologs. Each of the black lemur (*E. macaco*) species has

two very similar *CYP2D* genes which arose from the duplication specific to the *Eulemur* lineage.

Prior to the divergence of the *L. catta* lineage from that of the bamboo lemurs, there is some

uncertainty in these results whether only a single duplication occurred or whether two

occurred. This uncertainty stems from two nodes displaying low bootstrap support (73.0% and

57.5%; see Fig. S6) in the original PhyML tree. The topology displayed in Figure S6 describes two

duplications and parsimoniously implies just as many deletions. In light of the low bootstrap

support for this pattern, however, it is conceivable that the true topology could place the *L.*

catta CYP2D17L2 gene into the clade with the two bamboo-lemur genes. Such a revision would

reduce the number of deletions and duplications. The other duplicated gene in the lemurid

lineage, while secondarily duplicated in the *L. catta* lineage, was lost in the *H. griseus* lineage.

Lastly, no functional homologs for the *CYP2D* subfamily could be retrieved for *H. griseus*

because no open reading frames devoid of stop codons could be found in the extracted

sequence for this species' single *CYP2D* gene. However, this isolated instance may be due to an

assembly artifact, as the *CYP2D* locus was located on the ends of two separate scaffolds in the

original Supernova assembly. In all, the general pattern of this subfamily is similar to that of

CYP2C: evidence of heightened genic turnover with an emphasis on loss in the specific

lineage(s) of the bamboo lemurs.

The *CYP2J* subfamily divided into three orthogroups (Fig. 2, top-right; Fig. S7). The first

of these (OG-0) contains only a single duplication node before the speciation node for the split

of *L. catta* from the bamboo lemurs. The complete lack of any bamboo-lemur genes in this

portion of the tree parsimoniously indicates a loss of that duplicate prior to the *Prolemur-*

Hapalemur divergence. The remaining two *CYP2J* orthogroups (OG-1 and OG-2) arose as the

product of a duplication event prior to the first speciation event of all species involved in this

analysis. While no more duplications were called in the OG1-OG2 clade, several deletions are

implied by the tree's topology. First, one gene appears to have been lost in the *Prop. coquereli*

lineage. Second, another deletion was ostensibly lost in the ancestral lemurid lineage. Last, *H.*

griseus lost one more homolog after that lineage's split from that of *Prol. simus.* Thus, the

CYP2J subfamily displays more of the above-mentioned pattern wherein the bamboo lemurs

stand out for their likelihood of losing P450 genes.

The topologies of the remaining *CYP2* subfamilies (*i.e.*, *E, F, G, R, S, U,* and *W*) show no

evidence of duplication (Figs. S8-S13). Degeneration of one *CYP2F* gene may have occurred in

the *Mic. murinus* lineage (see Fig. S9), but anthropoids and at least some lorisiforms appear to

possess only a single *CYP2F* gene as well (Chaney et al., 2020; Chaney, 2023). Therefore, the

ancestral primate number of *CYP2F* orthologs is uncertain without the inclusion of a more basal

species in the analysis. With this single possible exception, bamboo lemurs have clearly

maintained the same number of gene copies in each of these subfamilies as the other lemur

species analyzed here.

The final subfamily of this part of the analysis further underscores the pattern described

for other subfamilies above (Fig. 2, bottom-left; Fig. S14). While OG-0 within the *CYP3A* tree

possesses only speciation nodes, one loss is evident in the absence of *Prop. coquereli* from this

clade. The other part of the tree exhibits a very complex topology because of the number of

duplication nodes contained within it. Many of these outside of the lemurid portion of the tree

are taxon-specific: *Prop. coquereli* and *Mic. murinus* each experienced two and four exclusive

duplication events, respectively. The lemurid portion of OG-1 shows at least three duplication

events prior to the divergence of *L. catta* and from the ancestral bamboo-lemur lineage. One

more duplication event, leading to the *L. catta* genes *CYP3A4L1* and *CYP3A4L2*, would be most

parsimoniously interpreted as having occurred specifically in ring-tail lemur lineage, but it is not

ruled out that this duplication may have occurred in an ancestral species and been lost by both

bamboo lemur species. Regardless, at least two separate events of genic loss seem to have

occurred in the lineage of the bamboo lemurs. Thus, while *L. catta* appears to exhibit six

separate *CYP3A* homologs, bamboo lemurs possess half that number; furthermore, we

conclude that this pattern is attributable nearly wholly to gene loss by bamboo lemurs rather

than lineage-specific expansion of *CYP3A* for *L. catta.*

<u>Selection relaxed on two important P450 xenobiotic metabolizers but intensified on another</u>

Out of all P450 subfamilies studied here, only *CYP1A* and *CYP2D* showed significant

signals of selectional relaxation within the bamboo-lemur clade (Table S2). Upon comparing the

dN/dS distribution for the reference branches to that of the bamboo-lemur branches, it appears

that the signatures of relaxation in the bamboo-lemur portions of these analyses are due to the

lessening of the pressure for diversifying selection (Fig. 3). The degree of relaxation appears to

differ between these two subfamilies. RELAX's estimate for the selection intensity parameter

(k), while significantly estimated to be less than 1.00, implies only a moderate level of

selectional relaxation for *CYP2D*. On the other hand, in the case of *CYP1A*, RELAX's estimate

for k was especially diminutive, indicating a stronger effect for that subfamily than for *CYP2A13*

and *CYP2D*.

Interestingly, the *CYP2F* subfamily showed significant intensification of selection in the

RELAX models reported here, exhibiting an estimate for k that was significantly higher than 1.00

(Table S3). The *dN/dS* categories within this subfamily's distribution portray a subfamily under

purifying selection for reference branches in the model (Fig. 3). Bamboo lemurs show an

exaggeration of this simple pattern, first, in the elevation of the strength of signals for both

diversifying selection and purifying selection. We followed this RELAX analysis with analyses

using both BUSTED (Murrell et al., 2015) and aBSREL (Smith et al., 2018). BUSTED analyses

found evidence of diversifying selection within the alignment (LRT = 21.47, p = 1.1e-5), and a

subsequent aBSREL analysis found significant signals for diversifying selection along the

terminal branches for *H. griseus CYP2F2* (LRT = 9.46, p = 1.53e-2) and *Prol. simus CYP2F2* (LRT =

85.35, p < 1e-5).

Discussion

In this study, we have demonstrated that the genomes of two bamboo lemur species (*H.*

griseus and *Prol. simus*) harbor, at a conservative maximum, the modal number of cytochrome

P450 genes across the other lemur species included here (Table S1). Indeed, in most cases

where losses of P450 genes were inferred, they are most parsimoniously attributed to a

bamboo lemur-specific branch of the phylogeny in question (see Figs. 1-6). Thus, bamboo

lemurs appear to show a narrower repertoire of xenobiotic-metabolizing P450s insofar as they

have generally fewer genes than other species of lemur, especially in comparison to their

closest relative, *Lemur catta.* Bamboo lemurs also show signals of relaxation of natural selection

for the *CYP1A* and *CYP2D* subfamilies (see Fig. 3), which are known to be involved in xenobiotic

metabolism (Danielson, 2002; Guengerich, 2015).

These results are perhaps best understood in terms of an evolutionary tradeoff on

detoxification strategies (Dearing et al., 2005; Freeland & Janzen, 1974; Ueno, 2001). In a

pivotal paper outlining several nutritional-ecological hypotheses, Freeland and Janzen (1974)

formalized the notion that generalist and specialist herbivores should differ inversely with

regard to the breadth of their arsenals for the metabolism of toxic plant secondary metabolites.

In their view, herbivores should possess a comparatively wide array of detoxification

mechanisms to handle low doses of a commensurately broad variety of toxins, and specialist

herbivores should, conversely, possess a smaller array of such mechanisms able to handle large

doses of a narrow variety of toxins (Dearing et al., 2005; Freeland & Janzen, 1974; Marsh et al.,

2006). The general pattern of copy-number differences and the more limited result of

selectional relaxation shown in this study seems to support this hypothesis in the case of

bamboo lemurs, alongside this taxon's high degree of specialization (see Table S1) and the

stupendous amounts of cyanide consumed by them (Ballhorn et al., 2009; Glander et al., 1989).

Despite bamboo lemurs' observed amount of gene loss for some xenobiotic-

metabolizing P450 genes, the *CYP2F* subfamily contrasted with other P450 subfamilies by

having not only signals of selection intensification but also a clear signal of diversifying

selection. Although this P450 subfamily is most highly expressed in the respiratory tract,

hepatic expression has been demonstrated, albeit at lower levels (Duff et al., 2015; Fagerberg

et al., 2014; Yue et al., 2014). Based on these results, we speculated that CYP2F must

metabolize a PSM present in bamboo. Human CYP2F1 and its model-rodent homologs can

activate coumarins through epoxidation reactions, leading to cytotoxicity and necrosis at

varying species-dependent levels (Born et al, 2002; Vassalo et al., 2004). Similarly, CYP2F

enzymes have been linked with bioactivation of the toxin naphthalene, also through an

oxidation reaction (Lanza et al., 1999; Li et al., 2011). These biochemical associations are

notable here because chromatographic assays and microbiome studies of various species of

bamboo have implied or directly indicated the presence of both naphthalene (Li et al., 2002;

Takahashi et al., 2010) and coumarins (Coffie et al., 2014) in those bamboo species (reviewed

by Gagliano et al., 2022). Generalizing from these findings, we would hypothesize, first, that

naphthalene and coumarins are present in Malagasy bamboo consumed by bamboo lemurs

and, second, that bamboo lemurs would be susceptible to toxicity from either of these

compounds or their metabolite oxides, which are toxic in themselves (napthalene: see Yost et

al., 2021; coumarins: see Foroozesh et al., 2019). Thus, both species' *CYP2F2* orthologs may

have experienced selective pressure for altering its translated enzyme's affinity or efficiency for

either naphthalene or coumarins.

The difference between bamboo lemurs and the sifaka *Propithecus coquereli* stands out

as a second confirmatory signal of Freeland and Janzen's (1974) tradeoff hypothesis because,

while both of these taxa are generally classified as folivorous, they differ in their amount of

dietary specialization. As mentioned previously, all species of bamboo lemur that have been

studied are heavily reliant on either grasses (usually bamboo) or sedges, and very few plants,

often only a single species, are required to describe a large majority of their diets (see Table

S1). Conversely, sifakas (*Propithecus* spp.) consume much more varied foods (Hemingway,

1998; Norscia et al., 2006; Powzyk & Mowry, 2003; Rowe et al., 2016), seasonally trending

toward gramnivory or frugivory depending on the species in question (Irwin 2006). Despite this

dietary diversity, sifakas do possess clear gastrointestinal adaptations related to hindgut

fermentation, such as a greatly elongated intestine and an expanded, haustrated cecum

(Campbell et al., 2000). For this reason, sifakas are often described as *anatomical folivores*, with

the implication that they are more behaviorally flexible with their diet choice than one might be

led to believe upon viewing their gastrointestinal anatomy. Meanwhile, bamboo lemurs of all

three major species possess a relatively simpler gastrointestinal anatomy, with an overall

relatively shorter intestine and a shorter, though completely haustrated, colon (Campbell et al.

2000; Yamashita et al. 2008). All of this information serves to position sifakas and bamboo

lemurs on opposite sides of a specialist-generalist continuum among herbivores.

Such dietary differences between these two taxa helps to clarify an ultimate explanation

for the opposite pattern of gene birth and death seen in the *CYP2B6* gene cluster for bamboo

lemurs and *Prop. coquereli* (see Fig. 1). This is also true, but to a less pronounced degree, for

the *CYP2A* subfamily (see Fig. 1). In both of these cases, bamboo lemurs exhibit clade-specific

gene losses and *Prop. coquereli* shows an opposite signal of lineage-specific gene duplications.

Indeed, (Guevara et al. 2021), utilizing this same genome assembly for *Prop. coquereli,*

independently found that these two P450 subfamilies had been enriched in so-called "rapidly

evolving genes," or genes with increased *dN/dS* ratios according to PAML branch-site models

(Yang, 2007). Because both *CYP2A* and *CYP2B* subfamilies are involved in the metabolism of

xenobiotic compounds (Guengerich 2015; Orr et al. 2020), at least these two P450 subfamilies

stand out as candidate gene groups supporting Freeland and Janzen's (1974) prediction, that

specialist and generalist herbivores should differ oppositely in the diversity of their means for

detoxifying plant secondary metabolites. This interesting natural experiment, of comparing

bamboo lemurs and sifakas (or indriids in general), deserves future investigation.

A second line of inquiry is the contrast between *Lemur catta* and the bamboo lemurs.

The number of xenobiotic-metabolizing P450 genes in the *L. catta* genome was always greater

than or equal to those in either bamboo lemur genome. The highly varied and very seasonal

diet documented for *L. catta* (Jolly, 1966; LaFleur, 2012; Sauther, 1992; Sauther et al., 1999;

Simmen, Sauther, et al., 2006; Soma, 2006) stands out as a potential explanation for this large

number of detoxification mechanisms. Indeed, Sauther et al. (1999) argue that the best dietary

classification for *L. catta* should be "opportunistic omnivores" (p. 122) because of the breadth

of this species' dietary repertoire and the drastic spikes of certain foods based on their seasonal

availability. This species also appears to be relatively inured to plant secondary metabolites

based on the high tannin and alkaloid content of their foods (Gould et al., 2009; Simmen,

Sauther, et al., 2006) and their relatively high rejection threshold for the bitter alkaloid quinine

(Simmen, Peronny, et al., 2006). This constellation of information portrays a pattern of an

animal that is exposed to a broad array of plant secondary metabolites and, perhaps because of

the annual variety of their foods, exposed to them at relatively low levels.

While *L. catta* possesses a sizeable number of xenobiotic-metabolizing P450 genes in

comparison to both species of bamboo lemur (see Table S1), these genes are not the products

of species-specific duplications, as is plain from the phylogenetic pattern of relatedness among

all the genes falling into the clade of ringtail and bamboo lemurs. Instead, many of these

duplicated genes seem to have arisen during the internode of a lemurid common ancestor, or

that of a common ancestor of the *Hapalemur* and *Lemur* genera, and then subsequently lost by

one or both of the bamboo lemurs species included here. This result accords well with

experimental work on bitter taste receptors (TAS2R16), which shows that the species of

bamboo lemur included here (as well as *H. aureus*) show a decreased sensitivity to several

bitter β-glycoside compounds (Itoigawa et al., 2021). Revealingly, however, the missense

mutations that most altered the response of the TAS2R16 protein were separate for *Prol. simus*

and *Hapalemur* spp., and ancestral reconstructions of the last common ancestors for both the

genus *Hapalemur* and for all bamboo lemurs showed a more sensitive response to these bitter

compounds than any of the present-day species-specific TAS2R16 isoforms (Itoigawa et al.,

2021). Given this earlier work and the pattern of P450 gene birth and death presented here, a

generalist dietary pattern is clearly the best estimate for an ancestral ecology of the

Hapalemur-Lemur last common ancestor. Furthermore, the complex pattern of loss and

retention of separate P450 paralogs by *Prol. simus* and *H. griseus* seems to agree with the

conclusion of Itoigawa et al. (2021) that these species' varying, but in all cases intense,

specialization on cyanogenic bamboo may have evolved in parallel rather than what would be

predicted based on parsimony, namely that this specialization would have evolved in the

common ancestor of all extant bamboo-lemur species.

On a methodological note, the pattern of gene gain and loss reported here is certainly

not the conclusion that would be reached by using an approach where the gene-count table is

parsimoniously compared to an organismal phylogeny to estimate ancestral states of gene

counts, as with the commonly used program for studying gene-family evolution, CAFE (De Bie

et al., 2006). This software program takes phylogeny into account only insofar as the species

themselves are concerned but does not include any information about the genes' own

phylogenetic relatedness, and for this reason, we see it as an undesirable alternative to

examining a phylogeny built from the genes themselves. While CAFE is very useful in its ability

to be easily automated, manual examinations of gene trees, or perhaps the use of programs

such as Possvm (Grau-Bové & Sebé-Pedrós, 2021) that make direct use of gene trees via the

species-overlap method (Huerta-Cepas, et al., 2007, 2016), would yield more accurate ancestral

reconstructions of gene copy-number differences among species.

A lingering question remains as to what compounds, or families of compounds, are

leading to the evolutionary expansions and contractions on display in this study. However,

there is currently a clear disconnect in the literature that inhibits any discussion between those

who have precise knowledge of P450 substrate compounds (*e.g.,* Guengerich 2015) and those

who perform phytochemical assays on foods of wild primates (*e.g.,* Eppley et al. 2017; Thurau

et al. 2021; Windley et al. 2022). The disconnect arises from differing levels of scale on which

each of these two groups operates: pharmacologists often use precise molecules (*e.g.,*

benzo[a]pyrene, aflatoxin B1) in their experiments and parlance, while phytochemists and

primatologists speak in the more general terms of compound families (*e.g.,* alkaloids,

condensed tannins). This nomenclatural gulf between these two camps is a foggy one, and it

serves to cloud any inferences that may be gained about the functions of free-ranging, non-

model primate species.

Here, we have found that bamboo lemurs have, on average, fewer *CYP1-3* gene copies

than other lemurs, and this is especially true when they are viewed alongside their closest

relative, the ringtail lemur. Viewed another way, bamboo lemurs have, at most, the same

number of P450 genes per subfamily as *L. catta*, and they often have fewer such genes. The

retention of these specific P450 homologs was not parsimonious when compared with the

organismal phylogeny, and the complexity of this pattern may imply some information about

bamboo-lemur evolution in general, namely that their specialization on cyanogenic bamboo

may have evolved in parallel rather than as a sort of ecological synapomorphy early after the

divergence of these two lineages. Only two P450 subfamilies (*CYP1A* and *CYP2D*), both of which

are linked with xenobiotic metabolism in humans (Guengerich, 2015), showed a statistical

decrease in the intensity of selection on the CDS for the genes within those subfamilies. As in

previous work on the evolution of the cytochrome P450 superfamily of genes, this topic

provides an interesting, complex, and sometimes unexpected window into biological evolution.

Future work that includes other species of bamboo lemur (*e.g.*, *H. alaotrensis, H. aureus*) will

doubtless only clarify that window.

Acknowledgments

This project was supported by NSF BCS1919857 to AJT and MEC and NSF

BCS1717188/BCS1718715/BCS1718339 to CMB and AJT. Transfer of the *Hapalemur griseus*

sample from Duke University to Kent State University was coordinated by   Erin Ehmke;

this is Duke Lemur Center publication number XXXX. We are thankful to Roy Heath and Joshua

Talbott for access to and support with high-power computing at Kent State University. Andrew

Burrell and Cody Ruiz were instrumental in extracting the HMW-DNA and preparing the library

for sequencing the *H. griseus* genome. MEC is grateful to Caylee Heiremans and Peter Chaney

for their support during and beyond this work.

References

Albalat, R., & Cañestro, C. (2016). Evolution by gene loss. *Nature Reviews Genetics*, *17*(7), 379–391. https://doi.org/10.1038/nrg.2016.39

Averdam, A., Kuschal, C., Otto, N., Westphal, N., Roos, C., Reinhardt, R., & Walter, L. (2011). Sequence analysis of the grey mouse lemur (Microcebus murinus) MHC class II DQ and DR region. *Immunogenetics*, *63*(2), 85–93. https://doi.org/10.1007/s00251-010-0487-3

Ballhorn, D. J., Kautz, S., & Rakotoarivelo, F. P. (2009). Quantitative Variability of Cyanogenesis in *Cathariostachys madagascariensis*—the Main Food Plant of Bamboo Lemurs in Southeastern Madagascar. *American Journal of Primatology*, *71*(4), 305–315. https://doi.org/10.1002/ajp.20653

Ballhorn, D. J., Rakotoarivelo, F. P., & Kautz, S. (2016). Coevolution of Cyanogenic Bamboos and Bamboo Lemurs on Madagascar. *PLoS ONE*, *11*(8), e0158935. https://doi.org/10.1371/journal.pone.0158935

Bass, C., Puinean, A. M., Zimmer, C. T., Denholm, I., Field, L. M., Foster, S. P., Gutbrod, O., Nauen, R., Slater, R., & Williamson, M. S. (2014). The evolution of insecticide resistance in the peach potato aphid, *Myzus persicae*. *Insect Biochemistry and Molecular Biology*, *51*(1), 41–51. https://doi.org/10.1016/j.ibmb.2014.05.003

Blumer, M., Brown, T., Freitas, M. B., Destro, A. L., Oliveira, J. A., Morales, A. E., Schell, T., Greve, C., Pippel, M., Jebb, D., Hecker, N., Ahmed, A. W., Kirilenko, B. M., Foote, M., Janke, A., Lim, B. K., & Hiller, M. (2022). Gene losses in the common vampire bat illuminate molecular adaptations to blood feeding. *Science Advances*, *8*(12). https://doi.org/10.1126/sciadv.abm6494

Born, S. L., Caudill, D., Fliter, K. L., & Purdon, M. P. (2002). Identification of the Cytochromes P450

That Catalyze Coumarin 3,4-Epoxidation and 3-Hydroxylation. *Drug Metabolism and*

Disposition, *30*(5), 483–487. https://doi.org/10.1124/dmd.30.5.483

Burns, J. J. (1957). Missing Step in Man, Monkey and Guinea Pig required for the Biosynthesis of L-

Ascorbic Acid. *Nature*, *180*(4585), 553–553. https://doi.org/10.1038/180553a0

Campbell, J. L., Eisemann, J. H., Williams, C. V., & Glenn, K. M. (2000). Description of the

gastrointestinal tract of five Lemur species: *Propithecus tattersalli, Propithecus verreauxi*

coquereli, Varecia variegata, Hapalemur griseus, and *Lemur catta*. *American Journal of*

Primatology, *52*(3), 133–142. https://doi.org/10.1002/1098-2345(200011)52:3<133::AID-

AJP2>3.0.CO;2-#

Capella-Gutierrez, S., Silla-Martinez, J. M., & Gabaldon, T. (2009). trimAl: a tool for automated

alignment trimming in large-scale phylogenetic analyses. *Bioinformatics*, *25*(15), 1972–1973.

https://doi.org/10.1093/bioinformatics/btp348

Chaney, M. E. (2023). *Primates, poison, and cytochrome P450: Evolutionary dynamism of the* CYP1-3

gene families within the primate order [Ph.D. dissertation]. Kent State University.

Chaney, M. E., Piontkivska, H., & Tosi, A. J. (2018). Retained duplications and deletions of *CYP2C*

genes among primates. *Molecular Phylogenetics and Evolution*, *125*, 204–212.

https://doi.org/10.1016/j.ympev.2018.03.037

Chaney, M. E., Romine, M. G., Piontkivska, H., & Tosi, A. J. (2020). Diversifying selection detected in

only a minority of xenobiotic-metabolizing *CYP1-3* genes among primate species. *Xenobiotica*,

50. https://doi.org/10.1080/00498254.2020.1785580

560 Charif, D., & Lobry, J. R. (2007). *SeqinR 1.0-2: A Contributed Package to the R Project for Statistical*

Computing Devoted to Biological Sequences Retrieval and Analysis (pp. 207–232).

https://doi.org/10.1007/978-3-540-35306-5_10

Coffie, G. Y., Antwi-Boasiako, C., & Darkwa, N. A. (2014). Phytochemical constituents of the leaves of

three bamboo (Poaceae) species in Ghana. *Journal of Pharmacognosy & Phytochemistry*, *2*(6),

34–38.

Comte, N., Morel, B., Hasić, D., Guéguen, L., Boussau, B., Daubin, V., Penel, S., Scornavacca, C., Gouy,

M., Stamatakis, A., Tannier, E., & Parsons, D. P. (2020). Treerecs: An integrated phylogenetic

tool, from sequences to reconciliations. *Bioinformatics*, *36*(18), 4822–4824.

https://doi.org/10.1093/bioinformatics/btaa615

Danielson, P. B. (2002). The Cytochrome P450 Superfamily: Biochemistry, Evolution and Drug

Metabolism in Humans. *Current Drug Metabolism*, *3*(6), 561–597.

https://doi.org/10.2174/1389200023337054

Darling, A. C. E., Mau, B., Blattner, F. R., & Perna, N. T. (2004). Mauve: Multiple Alignment of

Conserved Genomic Sequence With Rearrangements. *Genome Research*, *14*(7), 1394–1403.

https://doi.org/10.1101/gr.2289704

Darriba, D., Taboada, G. L., Doalla, R., & Posada, D. (2012). jModelTest 2: More models, new

heuristics and parallel computing. *Nature Methods*, *9*(8), 772.

Darwin, C. R. (1859). *On the origin of species by means of natural selection*. John Murray.

De Bie, T., Cristianini, N., Demuth, J. P., & Hahn, M. W. (2006). CAFE: A computational tool for the

study of gene family evolution. *Bioinformatics*, *22*(10), 1269–1271.

https://doi.org/10.1093/bioinformatics/btl097

Dearing, M. D., Foley, W. J., & McLean, S. (2005). The influence of plant secondary metabolites on

the nutritional ecology of herbivorous terrestrial vertebrates. *Annual Review of Ecology,*

Evolution, and Systematics, *36*(May), 169–189.

https://doi.org/10.1146/annurev.ecolsys.36.102003.152617

Duff, M. O., Olson, S., Wei, X., Garrett, S. C., Osman, A., Bolisetty, M., Plocik, A., Celniker, S. E., &

Graveley, B. R. (2015). Genome-wide identification of zero nucleotide recursive splicing in

Drosophila. *Nature*, *521*(7552), 376–379. https://doi.org/10.1038/nature14475

Eppley, T. M., Donati, G., Ramanamanjato, J. B., Randriatafika, F., Andriamandimbiarisoa, L. N.,

Rabehevitra, D., Ravelomanantsoa, R., & Ganzhorn, J. U. (2015). The use of an invasive species

habitat by a small folivorous primate: Implications for lemur conservation in Madagascar. *PLoS*

ONE, *10*(11), e0140981. https://doi.org/10.1371/journal.pone.0140981

Eppley, T. M., Tan, C. L., Arrigo-Nelson, S. J., Donati, G., Ballhorn, D. J., & Ganzhorn, J. U. (2017). High

Energy or Protein Concentrations in Food as Possible Offsets for Cyanide Consumption by

Specialized Bamboo Lemurs in Madagascar. *International Journal of Primatology*, *38*, 881–899.

https://doi.org/10.1007/s10764-017-9987-4

Gronen, J. T., Zohdy, S., Evans, A. R., Tecot, S. R., Wright, P. C., & Jernvall, J. (2017). Feeding Ecology

and Morphology Make a Bamboo Specialist Vulnerable to Climate Change. *Current Biology*,

27(21), 3384–3389. https://doi.org/10.1016/j.cub.2017.09.050

Eschler, B. M., Pass, D. M., Willis, R., & Foley, W. J. (2000). Distribution of foliar formylated

phloroglucinol derivatives amongst Eucalyptus species. *Biochemical Systematics and Ecology*.

https://doi.org/10.1016/S0305-1978(99)00123-4

Lagerberg, L., Hallström, B. M., Oksvold, P., Kampf, C., Djureinovic, D., Odeberg, J., Habuka, M.,

Tahmasebpoor, S., Danielsson, A., Edlund, K., Asplund, A., Sjöstedt, E., Lundberg, E., Szigyarto,

C. A.-K., Skogs, M., Takanen, J. O., Berling, H., Tegel, H., Mulder, J., … Uhlén, M. (2014). Analysis

of the human tissue-specific expression by genome-wide integration of transcriptomics and

antibody-based proteomics. *Molecular & Cellular Proteomics : MCP*, *13*(2), 397–406.

https://doi.org/10.1074/mcp.M113.035600

Feyereisen, R. (2011). Arthropod CYPomes illustrate the tempo and mode in P450 evolution.

Biochimica et Biophysica Acta, *1814*(1), 19–28. https://doi.org/10.1016/j.bbapap.2010.06.012

Field, L. M., & Devonshire, A. L. (1998). Evidence that the *E4* and *FE4* esterase genes responsible for

insecticide resistance in the aphid *Myzus persicae* (Sulzer) are part of a gene family. *The*

Biochemical Journal, *330*, 169–173.

Foroozesh, M., Sridhar, J., Goyal, N., & Liu, J. (2019). Coumarins and P450s, Studies Reported to-

Date. *Molecules*, *24*(8), 1620. https://doi.org/10.3390/molecules24081620

Freeland, W. J., & Janzen, D. H. (1974). Strategies in Herbivory by Mammals: The Role of Plant

Secondary Compounds. *The American Naturalist*, *108*(961), 269–289.

Gagliano, J., Anselmo-Moreira, F., Sala-Carvalho, W. R., & Furlan, C. M. (2022). What is known about

the medicinal potential of bamboo? *Advances in Traditional Medicine*, *22*(3), 467–495.

https://doi.org/10.1007/s13596-020-00536-5

Glander, K. E., Wright, P. C., Seigler, S. S., Randrianasolo, V., & Randrianasolo, B. (1989).

Consumption of cyanogenic bamboo by a newly discovered species of bamboo lemur. *American*

Journal of Primatology, *19*, 119–124.

Gould, L., Constabel, P., Mellway, R., & Rambeloarivony, H. (2009). Condensed tannin intake in

spiny-forest-dwelling *Lemur catta* at Berenty Reserve, Madagascar, during reproductive

periods. *Folia Primatologica*, *80*(4), 249–263. https://doi.org/10.1159/000252584

Grassi, C. (2006). Variability in Habitat, Diet, and Social Structure of Hapalemur griseus in

Ranomafana National Park, Madagascar. *American Journal of Physical Anthropology*, *131*, 50–

63. https://doi.org/10.1002/ajpa.20423

Grau-Bové, X., & Sebé-Pedrós, A. (2021). Orthology Clusters from Gene Trees with Possvm.

Molecular Biology and Evolution, *38*(11), 5204–5208. https://doi.org/10.1093/molbev/msab234

Guengerich, F. P. (2015). Human Cytochrome P450 Enzymes. In P. R. Ortiz de Montellano (Ed.),

Cytochrome P450: Structure, Mechanism, and Biochemistry (4th ed., pp. 523–785). Springer

International Publishing. https://doi.org/10.1007/978-3-319-12108-6_9

Guevara, E. E., Webster, T. H., Lawler, R. R., Bradley, B. J., Greene, L. K., Ranaivonasy, J., Ratsirarson,

J., Harris, R. A., Liu, Y., Murali, S., Raveendran, M., Hughes, D. S. T., Muzny, D. M., Yoder, A. D.,

Worley, K. C., & Rogers, J. (2021). Comparative genomic analysis of sifakas (*Propithecus*) reveals

selection for folivory and high heterozygosity despite endangered status. *Science Advances*,

7(17), 1–13. https://doi.org/10.1126/sciadv.abd2274

Guindon, S., Dufayard, J.-F., Lefort, V., Anisimova, M., Hordijk, W., & Gascuel, O. (2010). New

Algorithms and Methods to Estimate Maximum-Likelihood Phylogenies: Assessing the

Performance of PhyML 3.0. *Systematic Biology*, *59*(3), 307–321.

Haley, S. L., Lamb, J. G., Franklin, M. R., Constance, J. E., & Dearing, M. D. (2007a). Xenobiotic

metabolism of plant secondary compounds in juniper (*Juniperus monosperma*) by specialist and

generalist woodrat herbivores, genus Neotoma. *Comparative Biochemistry and Physiology, Part*

C: Toxicology and Pharmacology, *146*(4), 552–560. https://doi.org/10.1016/j.cbpc.2007.06.007

Haley, S. L., Lamb, J. G., Franklin, M. R., Constance, J. E., & Dearing, M. D. (2007b). Xenobiotic

metabolism of plant secondary compounds in oak (*Quercus agrifolia*) by specialist and

generalist woodrat herbivores, genus *Neotoma*. *Journal of Chemical Ecology*, *33*(11), 2111–

2122. https://doi.org/10.1007/s10886-007-9371-5

Harris, R. S. (2007). *Improved Pairwise Alignment of Genomic DNA*. The Pennsylvania State

University.

Hawkins, M. T. R., Culligan, R. R., Frasier, C. L., Dikow, R. B., Hagenson, Ryan., Lei, Runhua., & Louis,

E. E. (2018). Genome sequence and population declines in the critically endangered greater

bamboo lemur (Prolemur simus) and implications for conservation. *BMC Genomics*, *19*(1), 445.

https://doi.org/10.1186/s12864-018-4841-4

Hecker, N., Sharma, V., & Hiller, M. (2019). Convergent gene losses illuminate metabolic and

physiological changes in herbivores and carnivores. *Proceedings of the National Academy of*

Sciences of the United States of America, *116*(8), 3036–3041.

https://doi.org/10.1073/pnas.1818504116

Hemingway, C. A. (1998). Selectivity and variability in the diet of Milne-Edwards' sifakas (*Propithecus*

diadema edwardsi): Implications for folivory and seed-eating. *International Journal of*

Primatology, *19*(2), 355–377. https://doi.org/10.1023/A:1020344018670

Hodgson, J. A., Pickrell, J. K., Pearson, L. N., Quillen, E. E., Prista, A., Rocha, J., Soodyall, H., Shriver,

M. D., & Perry, G. H. (2014). Natural selection for the Duffy-null allele in the recently admixed

people of Madagascar. *Proceedings of the Royal Society B: Biological Sciences*, *281*(1789).

https://doi.org/10.1098/rspb.2014.0930

Hornung, T. C., & Biesalski, H.-K. (2019). Glut-1 explains the evolutionary advantage of the loss of

endogenous vitamin C-synthesis. *Evolution, Medicine, and Public Health*, *2019*(1), 221–231.

https://doi.org/10.1093/emph/eoz024

Huelsmann, M., Hecker, N., Springer, M. S., Gatesy, J., Sharma, V., & Hiller, M. (2019). Genes lost

during the transition from land to water in cetaceans highlight genomic changes associated

with aquatic adaptations. *Science Advances*, *5*(9). https://doi.org/10.1126/sciadv.aaw6671

Huerta-Cepas, J., Dopazo, H., Dopazo, J., & Gabaldón, T. (2007). The human phylome. *Genome*

Biology, *8*(6), R109. https://doi.org/10.1186/gb-2007-8-6-r109

Huerta-Cepas, J., Serra, F., & Bork, P. (2016). ETE 3: Reconstruction, Analysis, and Visualization of

Phylogenomic Data. *Molecular Biology and Evolution*, *33*(6), 1635–1638.

https://doi.org/10.1093/molbev/msw046

Hunnicutt, K. E., Tiley, G. P., Williams, R. C., Larsen, P. A., Blanco, M. B., Rasoloarison, R. M.,

Campbell, C. R., Zhu, K., Weisrock, D. W., Matsunami, H., & Yoder, A. D. (2020). Comparative

Genomic Analysis of the Pheromone Receptor Class 1 Family (V1R) Reveals Extreme Complexity

in Mouse Lemurs (Genus, Microcebus) and a Chromosomal Hotspot across Mammals. *Genome*

Biology and Evolution, *12*(1), 3562–3579. https://doi.org/10.1093/gbe/evz200

Ingelman-Sundberg, M. (2022). Cytochrome P450 polymorphism: From evolution to clinical use.

Advances in Pharmacology, *95*, 393–416. https://doi.org/10.1016/bs.apha.2022.04.004

Irwin, M. T. (2006). Ecologically enigmatic lemurs: The sifakas of the eastern forests (*Propithecus*

candidus, P. diadema, P. edwardsi, P. perrieri and *P. tattersalli*). In L. Gould & M. L. Sauther

(Eds.), *Lemurs: Ecology and Adaptation* (pp. 305–326). Springer.

Itoigawa, A., Fierro, F., Chaney, M. E., Lauterbur, M. E., Hayakawa, T., Tosi, A. J., Niv, M. Y., & Imai, H.

(2021). Lowered sensitivity of bitter taste receptors to β-glucosides in bamboo lemurs: An

instance of parallel and adaptive functional decline in TAS2R16? *Proceedings of the Royal*

Society B: Biological Sciences, *288*(1948). https://doi.org/10.1098/rspb.2021.0346

Itoigawa, A., Hayakawa, T., Suzuki-Hashido, N., & Imai, H. (2019). A natural point mutation in the

bitter taste receptor TAS2R16 causes inverse agonism of arbutin in lemur gustation.

Proceedings of the Royal Society B, *286*(1904), 20190884.

https://doi.org/10.1098/rspb.2019.0884

Johnson, R. N., O'Meally, D., Chen, Z., Etherington, G. J., Ho, S. Y. W., Nash, W. J., Grueber, C. E.,

Cheng, Y., Whittington, C. M., Dennison, S., Peel, E., Haerty, W., O'Neill, R. J., Colgan, D., Russell,

T. L., Alquezar-Planas, D. E., Attenbrow, V., Bragg, J. G., Brandies, P. A., … Belov, K. (2018).

Adaptation and conservation insights from the koala genome. *Nature Genetics*.

https://doi.org/10.1038/s41588-018-0153-5

Jolly, A. (1966). *Lemur Behaviour*. University of Chicago Press.

Joussen, N., & Heckel, D. G. (2021). Saltational evolution of a pesticide-metabolizing cytochrome

P450 in a global crop pest. *Pest Management Science*, *77*(7), 3325–3332.

https://doi.org/10.1002/ps.6376

Katoh, K., & Standley, D. M. (2013). MAFFT multiple sequence alignment software version 7:

Improvements in performance and usability. *Molecular Biology and Evolution*, *30*(4), 772–780.

https://doi.org/10.1093/molbev/mst010

Kawashima, A., & Satta, Y. (2014). Substrate-dependent evolution of cytochrome P450: Rapid

turnover of the detoxification-type and conservation of the biosynthesis-type. *PLoS ONE*, *9*(6).

https://doi.org/10.1371/journal.pone.0100059

Kirischian, N., McArthur, A. G., Jesuthasan, C., Krattenmacher, B., & Wilson, J. Y. (2011). Phylogenetic

and functional analysis of the vertebrate cytochrome P450 2 family. *Journal of Molecular*

Evolution, *72*(1), 56–71. https://doi.org/10.1007/s00239-010-9402-7

Kitanovic, S., Orr, T. J., Spalink, D., Cocke, G. B., Schramm, K., Wilderman, P. R., Halpert, J. R., &

Dearing, M. D. (2018). Role of Cytochrome P450 2B Sequence Variation and Gene Copy Number

in Facilitating Dietary Specialization in Mammalian Herbivores. *Molecular Ecology*, *12*(10),

3218–3221. https://doi.org/10.1111/mec.14480

Lachapelle, M. Y., & Drouin, G. (2011). Inactivation dates of the human and guinea pig vitamin C

genes. *Genetica*, *139*(2), 199–207. https://doi.org/10.1007/s10709-010-9537-x

LaFleur, M. (2012). *Ecology of ring-tailed lemurs (Lemur catta) at the Tsimanampetsotsa National*

Park, Madagascar: Implications for Female Dominance and the Evolution of Lemur Traits.

University of Colorado Boulder.

Lanza, D. L., Code, E., Crespi, C. L., Gonzalez, F. J., & Yost, G. S. (1999). Specific dehydrogenation of 3-

methylindole and epoxidation of naphthalene by recombinant human CYP2F1 expressed in

lymphoblastoid cells. *Drug Metabolism and Disposition: The Biological Fate of Chemicals*, *27*(7),

798–803.

Lauterbur, M. E. (2019). In extremis: *Breaking coalescent models in small populations and concerted*

convergence of mammalian cyanide adaptation. Stony Brook University.

Lauterbur, M. E., Tongasoa, L., Peralta, J., Jacox, A., Concheiro-Guisan, M., & Wright, P. C. (2017).

Peeing Poison: The Biochemistry of Bamboo Lemur Cyanide Survival. *Integrative and*

Comparative Biology, *57*, E96.

Lecompte, E., Crouau-Roy, B., Aujard, F., Holota, H., & Murienne, J. (2016). Complete mitochondrial

genome of the gray mouse lemur, Microcebus murinus (Primates, Cheirogaleidae).

Mitochondrial DNA Part A, *27*(5), 3514–3516. https://doi.org/10.3109/19401736.2015.1074196

Li, L., Wei, Y., Van Winkle, L., Zhang, Q.-Y., Zhou, X., Hu, J., Xie, F., Kluetzman, K., & Ding, X. (2011).

Generation and Characterization of a *Cyp2f2* -Null Mouse and Studies on the Role of CYP2F2 in

Naphthalene-Induced Toxicity in the Lung and Nasal Olfactory Mucosa. *Journal of*

Pharmacology and Experimental Therapeutics, *339*(1), 62–71.

https://doi.org/10.1124/jpet.111.184671

Li, S., Tian, Y., Sun, M., Liu, J., Bai, Y., Liu, X., & Guo, Y. (2022). Characterization of Key Aroma

Compounds in Fermented Bamboo Shoots Using Gas Chromatography-Olfactometry-Mass

Spectrometry, Odor Activity Values, and Aroma Recombination Experiments. *Foods*, *11*(14),

2106. https://doi.org/10.3390/foods11142106

Lowe, T. M., & Eddy, S. R. (1997). tRNAscan-SE: A Program for Improved Detection of Transfer RNA

Genes in Genomic Sequence. *Nucleic Acids Research*, *25*(5), 955–964.

https://doi.org/10.1093/nar/25.5.955

Marciniak, S., Mughal, M. R., Godfrey, L. R., Bankoff, R. J., Randrianatoandro, H., Crowley, B. E.,

Bergey, C. M., Muldoon, K. M., Randrianasy, J., Raharivololona, B. M., Schuster, S. C., Mahli, R.

S., Yoder, A. D., Louis, E. E., Kistler, L., & Perry, G. H. (2021). Evolutionary and phylogenetic

insights from a nuclear genome of the extinct giant koala lemur *Megaladapis edwardsi*.

Proceedings of the National Academy of Sciences of the United States of America, *118*(26),

e2022117118. https://doi.org/10.1101/2020.10.16.342907

Marsh, K. J., Wallis, I. R., McLean, S., Sorensen, J. S., & Foley, W. J. (2006). Conflicting demands on

detoxification pathways influence how common brushtail possums choose their diets. *Ecology*,

87(8), 2103–2112. https://doi.org/10.1890/0012-9658(2006)87[2103:CDODPI]2.0.CO;2

McKenney, E. A., Maslanka, M., Rodrigo, A., & Yoder, A. D. (2017). Bamboo Specialists from Two

Mammalian Orders (Primates, Carnivora) Share a High Number of Low-Abundance Gut

Microbes. *Microbial Ecology*, 1–13. https://doi.org/10.1007/s00248-017-1114-8

Meyer, W. K., Venkat, A., Kermany, A. R., Van De Geijn, B., Zhang, S., & Przeworski, M. (2015).

Evolutionary history inferred from the de novo assembly of a nonmodel organism, the blue-

eyed black lemur. *Molecular Ecology*, *24*(17), 4392–4405. https://doi.org/10.1111/mec.13327

Mutschler, T. (1999). Folivory in a small-bodied lemur: The nutrition of the Alaotran gentle lemur

(Hapalemur griseus alaotrensis). In B. Rakotosamimanana, H. Rasamimanana, J. U. Ganzhorn, &

S. M. Goodman (Eds.), *New Directions in Lemur Studies* (pp. 221–240). Kluwer

Academic/Plenum Publishers.

Murrell, B., Weaver, S., Smith, M. D., Wertheim, J. O., Murrell, S., Aylward, A., Eren, K., Pollner, T.,

Martin, D. P., Smith, D. M., Scheffler, K., & Kosakovsky Pond, S. L. (2015). Gene-Wide

Identification of Episodic Selection. *Molecular Biology and Evolution*, *32*(5), 1365–1371.

https://doi.org/10.1093/molbev/msv035

Norscia, I., Carrai, V., & Borgognini-Tarli, S. M. (2006). Influence of dry season and food quality and

quantity on behavior and feeding strategy of *Propithecus verreauxi* in Kirindy, Madagascar.

International Journal of Primatology, *27*(4), 1001–1022. https://doi.org/10.1007/s10764-006-

9056-x

Olson, M. V. (1999). When Less Is More: Gene Loss as an Engine of Evolutionary Change. *American*

Journal of Human Genetics, *64*(1), 18–23. https://doi.org/10.1086/302219

Orr, T. J., Kitanovic, S., Schramm, K. M., Skopec, M. M., Wilderman, P. R., Halpert, J. R., & Dearing, M.

D. (2020). Strategies in herbivory by mammals revisited: The role of liver metabolism in a

juniper specialist (*Neotoma stephensi*) and a generalist (*Neotoma albigula*). *Molecular Ecology*,

29(9), 1674–1683. https://doi.org/10.1111/mec.15431

Overdorff, D. J., Strait, S. G., & Telo, A. (1997). Seasonal variation in activity and diet in a small-

bodied folivorous primate, *Hapalemur griseus*, in southeastern Madagascar. *American Journal*

of Primatology, *43*(3), 211–223. https://doi.org/10.1002/(SICI)1098-2345(1997)43:3<211::AID-

AJP2>3.0.CO;2-#

Palmada-Flores, M., Orkin, J. D., Haase, B., Mountcastle, J., Bertelsen, M. F., Fedrigo, O., Kuderna, L.

F. K., Jarvis, E. D., & Marques-Bonet, T. (2022). A high-quality, long-read genome assembly of

the endangered ring-tailed lemur (*Lemur catta*). *GigaScience*, *11*, 1–7.

https://doi.org/10.1093/gigascience/giac026

Pass, G. J., McLean, S., Stupans, I., & Davies, N. (2001). Microsomal metabolism of the terpene 1,8-

cineole in the common brushtail possum (*Trichosurus vulpecula*), koala (*Phascolarctos*

cinereus), rat and human. *Xenobiotica*, *31*(4), 205–221.

Pauling, L. (1970). Evolution and the Need for Ascorbic Acid. *Proceedings of the National Academy of*

Sciences of the United States of America, *67*(4), 1643–1648.

https://doi.org/10.1073/pnas.67.4.1643

Powzyk, J. A., & Mowry, C. B. (2003). Dietary and Feeding Differences between Sympatric

Propithecus diadema diadema and *Indri indri*. *International Journal of Primatology*, *24*(6), 1143–

1162. https://doi.org/10.1023/B:IJOP.0000005984.36518.94

Revell, L. J. (2012). phytools: An R package for phylogenetic comparative biology (and other things).

Methods in Ecology and Evolution, *3*(2), 217–223. https://doi.org/10.1111/j.2041-

210X.2011.00169.x

Rowe, N., Wright, P. C., & Jacobs, R. (2016). Sifakas, Indris, and Wooly Lemurs. In N. Rowe & M.

Myers (Eds.), *All the World's Primates* (pp. 87–88). Pogonias Press.

Sauther, M. L. (1992). *Effect of reproductive state, social rank and group size on resource use among*

free-ranging ringtailed lemurs (Lemur catta) of Madagascar. Washington University, St. Louis.

Sauther, M. L., Sussman, R. W., & Gould, L. (1999). The socioecology of the ringtailed lemur: Thirty-

five years of research. *Evolutionary Anthropology*, *8*(4), 120–132.

https://doi.org/10.1002/(SICI)1520-6505(1999)8:4<120::AID-EVAN3>3.0.CO;2-O

Shank, S. D., Weaver, S., & Kosakovsky Pond, S. L. (2018). phylotree.js - a JavaScript library for

application development and interactive data visualization in phylogenetics. *BMC*

Bioinformatics, *19*(1), 276. https://doi.org/10.1186/s12859-018-2283-2

Sharma, V., Hecker, N., Roscito, J. G., Foerster, L., Langer, B. E., & Hiller, M. (2018). A genomics

approach reveals insights into the importance of gene losses for mammalian adaptations.

Nature Communications, *9*(1), 1–9. https://doi.org/10.1038/s41467-018-03667-1

Simão, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V., & Zdobnov, E. M. (2015). BUSCO:

assessing genome assembly and annotation completeness with single-copy orthologs.

Bioinformatics, *31*(19), 3210–3212. https://doi.org/10.1093/bioinformatics/btv351

Simmen, B., Peronny, S., Jeanson, M., Hladik, A., & Marez, A. (2006). Diet quality and taste

perception of plant secondary metabolites by *Lemur catta*. In A. Jolly, R. W. Sussman, N.

Koyama, & H. Rasamimanana (Eds.), *Ringtailed Lemur Biology* (pp. 160–183). Springer

Science+Business Media.

Simmen, B., Sauther, M. L., Soma, T., Rasamimanana, H., Sussman, R. W., Jolly, A., Tarnaud, L., &

Hladik, A. (2006). Plant species fed on by *Lemur catta* in gallery forests of the southern domain

of Madagascar. In A. Jolly, R. W. Sussman, N. Koyama, & H. Rasamimanana (Eds.), *Ringtailed*

Lemur Biology (pp. 55--68). Springer Science+Business Media, LLC.

Skopec, M. M., Halpert, J. R., & Dearing, M. D. (2022). Mammalian cytochrome P450 biodiversity:

Physiological importance, function, and protein and genomic structures of cytochromes P4502B

in multiple species of woodrats with different dietary preferences. *Advances in Pharmacology*,

95, 107–129. https://doi.org/10.1016/bs.apha.2022.05.002

Smith, M. D., Wertheim, J. O., Weaver, S., Murrell, B., Scheffler, K., & Kosakovsky Pond, S. L. (2018).

Less Is More: An Adaptive Branch-Site Random Effects Model for Efficient Detection of Episodic

Diversifying Selection. *Molecular Biology and Evolution*, *32*(5), 1342–1353.

https://doi.org/10.1093/molbev/msv022

Soma, T. (2006). Tradition and novelty: *Lemur catta* feeding strategy on introduced tree species at

Berenty Reserve. In A. Jolly, R. W. Sussman, N. Koyama, & H. Rasamimanana (Eds.), *Ringtailed*

Lemur Biology (pp. 141–159). Springer Science+Business Media.

Sorensen, J. S., & Dearing, M. D. (2003). Elimination of plant toxins by herbivorous woodrats:

Revisiting an explanation for dietary specialization in mammalian herbivores. *Oecologia*, *134*(1),

88–94. https://doi.org/10.1007/s00442-002-1085-3

Tan, C. L. (1999). Group composition, home range size, and diet of three sympatric bamboo lemur

species (genus *Hapalemur*) in Ranomafana National Park, Madagascar. *International Journal of*

Primatology, *20*(4), 547–566. https://doi.org/10.1023/A:1020390723639

Takahashi, T., Mizui, K., & Miyazawa, M. (2010). Volatile compounds with characteristic odour in

moso-bamboo stems (*Phyllostachys pubescens* Mazel ex Houz. De ehaie). *Phytochemical*

Analysis, *21*(5), 489–495. https://doi.org/10.1002/pca.1224

Tan, C. L. (2006). Behavior and ecology of gentle lemurs (Genus *Hapalemur*). In L. Gould & M. L.

Sauther (Eds.), *Lemurs: Ecology and Adaptation* (pp. 369–382). Springer Science+Business

Media, LLC.

Thurau, E. G., Rahajanirina, A. N., & Irwin, M. T. (2021). Condensed tannins in the diet of folivorous

diademed sifakas and the gap between crude and available protein. *American Journal of*

Primatology, *83*(3), 1–14. https://doi.org/10.1002/ajp.23239

Ueno, Y. (2001). How do we eat? Hypothesis of foraging strategy from the viewpoint of gustation in

primates. In T. Matsuzawa (Ed.), *Primate Origins of Human Cognition and Behavior* (pp. 104–

111). Springer Japan.

Vassallo, J. D., Hicks, S. M., Born, S. L., & Daston, G. P. (2004). Roles for Epoxidation and

Detoxification of Coumarin in Determining Species Differences in Clara Cell Toxicity.

Toxicological Sciences, *82*(1), 26–33. https://doi.org/10.1093/toxsci/kfh237

Vivekanandhan, P., Thendralmanikandan, A., Kweka, E. J., & Mahande, A. M. (2021). Resistance to

temephos in *Anopheles stephensi* larvae is associated with increased cytochrome P450 and α-

esterase genes overexpression. *International Journal of Tropical Insect Science*, *41*(4), 2543–

2548. https://doi.org/10.1007/s42690-021-00434-6

Vontas, J., Katsavou, E., & Mavridis, K. (2020). Cytochrome P450-based metabolic insecticide

resistance in Anopheles and Aedes mosquito vectors: Muddying the waters. *Pesticide*

Biochemistry and Physiology, *170*(April), 104666.

https://doi.org/10.1016/j.pestbp.2020.104666

Wagner, F., Ruf, I., Lehmann, T., Hofmann, R., Ortmann, S., Schiffmann, C., Hiller, M., Stefen, C., &

Stuckas, H. (2022). Reconstruction of evolutionary changes in fat and toxin consumption reveals

associations with gene losses in mammals: A case study for the lipase inhibitor PNLIPRP1 and

the xenobiotic receptor NR1I3. *Journal of Evolutionary Biology*, *35*(2), 225–239.

https://doi.org/10.1111/jeb.13970

Walsh, T. K., Joussen, N., Tian, K., McGaughran, A., Anderson, C. J., Qiu, X., Ahn, S.-J., Bird, L., Pavlidi,

N., Vontas, J., Ryu, J., Rasool, A., Barony Macedo, I., Tay, W. T., Zhang, Y., Whitehouse, M. E. A.,

Silvie, P. J., Downes, S., Nemec, L., & Heckel, D. G. (2018). Multiple recombination events

between two cytochrome P450 loci contribute to global pyrethroid resistance in Helicoverpa

armigera. *PLOS ONE*, *13*(11), e0197760. https://doi.org/10.1371/journal.pone.0197760

Wang, B., Shahzad, M. F., Zhang, Z., Sun, H., Han, P., Li, F., & Han, Z. (2014). Genome-wide analysis

reveals the expansion of Cytochrome P450 genes associated with xenobiotic metabolism in rice

striped stem borer, *Chilo suppressalis*. *Biochemical and Biophysical Research Communications*,

443(2), 756–760. https://doi.org/10.1016/j.bbrc.2013.12.045

Weisenfeld, N. I., Kumar, V., Shah, P., Church, D. M., & Jaffe, D. B. (2017). Direct determination of

diploid genome sequences. *Genome Research*, *27*(5), 757–767.

https://doi.org/10.1101/gr.214874.116

Wertheim, J. O., Murrell, B., Smith, M. D., Pond, S. L. K., & Scheffler, K. (2015). RELAX: Detecting

relaxed selection in a phylogenetic framework. *Molecular Biology and Evolution*, *32*(3), 820–

832. https://doi.org/10.1093/molbev/msu400

Wilderman, P. R., Jang, H.-H., Malenke, J. R., Salib, M., Angermeier, E., Lamime, S., Dearing, M. D., &

Halpert, J. R. (2014). Functional characterization of cytochromes P450 2B from the desert

woodrat Neotoma lepida. *Toxicology and Applied Pharmacology*, *274*(3), 393–401.

https://doi.org/10.1016/j.taap.2013.12.005

Windley, H. R., Starrs, D., Stalenberg, E., Rothman, J. M., Ganzhorn, J. U., & Foley, W. J. (2022). Plant

secondary metabolites and primate food choices: A meta-analysis and future directions.

American Journal of Primatology, *84*(8), 1–18. https://doi.org/10.1002/ajp.23397

Wondji, C. S., Irving, H., Morgan, J., Wondji, C. S., Irving, H., Morgan, J., Lobo, N. F., Collins, F. H.,

Hunt, R. H., Coetzee, M., Hemingway, J., & Ranson, H. (2009). Two duplicated P450 genes are

associated with pyrethroid resistance in *Anopheles funestus*, a major malaria vector. *Genome*

Research, *19*, 452–459. https://doi.org/10.1101/gr.087916.108

Yamashita, N., Tan, C. L., & Vinyard, C. J. (2008). A comparison of gastrointestinal tracts of sympatric

Hapalemur species. *American Journal of Physical Anthropology*, *Suppl.*, 226.

Yamashita, N., Tan, C. L., Vinyard, C. J., & Williams, C. (2010). Semi-Quantitative Tests of Cyanide in

Foods and Excreta of Three Hapalemur Species in Madagascar. *American Journal of*

Primatology, *72*, 56–61. https://doi.org/10.1002/ajp.20751

Yang, Z. (2007). PAML 4: Phylogenetic Analysis by Maximum Likelihood. *Molecular Biology and*

Evolution, *24*(8), 1586–1591. https://doi.org/10.1093/molbev/msm088

Yost, E. E., Galizia, A., Kapraun, D. F., Persad, A. S., Vulimiri, S. V., Angrish, M., Lee, J. S., & Druwe, I. L.

(2021). Health Effects of Naphthalene Exposure: A Systematic Evidence Map and Analysis of

Potential Considerations for Dose–Response Evaluation. *Environmental Health Perspectives*,

129(7). https://doi.org/10.1289/EHP7381

Yue, F., Cheng, Y., Breschi, A., Vierstra, J., Wu, W., Ryba, T., Sandstrom, R., Ma, Z., Davis, C., Pope, B.

D., Shen, Y., Pervouchine, D. D., Djebali, S., Thurman, R. E., Kaul, R., Rynes, E., Kirilusha, A.,

Marinov, G. K., Williams, B. A., … Ren, B. (2014). A comparative encyclopedia of DNA elements

in the mouse genome. *Nature*, *515*(7527), 355–364. https://doi.org/10.1038/nature13992

Zhao, H., Xu, D., Zhang, S., & Zhang, J. (2012). Genomic and Genetic Evidence for the Loss of Umami

Taste in Bats. *Genome Biology and Evolution*, *4*(1), 73–79. https://doi.org/10.1093/gbe/evr126

Zhao, H., Yang, J.-R., Xu, H., & Zhang, J. (2010). Pseudogenization of the Umami Taste Receptor Gene

Tas1r1 in the Giant Panda Coincided with its Dietary Switch to Bamboo. *Molecular Biology and*

Evolution, *27*(12), 2669–2673. https://doi.org/10.1093/molbev/msq153

Zhu, L., Yang, Z., Yao, R., Xu, L., Chen, H., Gu, X., Wu, T., & Yang, X. (2018). Potential Mechanism of

Detoxification of Cyanide Compounds by Gut Microbiomes of Bamboo-Eating Pandas. *MSphere*,

3(3). https://doi.org/10.1128/mSphere.00229-18

<u>Data Accessibility and Benefit-sharing</u>

FASTQ reads from the genomic sequencing of *Hapalemur griseus* have been deposited

in the NCBI SRA (accession number XXXXX), and the alignments and trees necessary to

reproduce this work are available on Dryad.

<u>Author Contributions</u>

MEC and AJT conceptualized the project. MEC collected and analyzed the data and

wrote the first draft of the paper, with guidance from AJT and CMB. All authors edited

subsequent drafts and approved of the final submission.

Table 1: Select statistics from de novo assembly of *Hapalemur griseus* (Beavis) using the

Supernova assembler.

Raw coverage	Contig N_{50}	Scaffold N_{50}	Est. genome size	BUSCOs recovered (Simão et al. 2015)
69.49x	277.73 kb	20.47 Mb	2.61 Gb[1]	3927/4104 (95.7%)

Figure 1: Evolution of three groups of *CYP2A, CYP2B6,* and *CYP2C* gene groups. Notes labels

with squares denote inferred gene-duplication events, while exes denote inferred gene losses.

For the sake of clarity, orthogroups are displayed as separate gene trees in this figure and color

coded to match orthogroup labels in Figures S3-S5. Genes labeled with asterisks are genes that

were originally placed in standalone orthogroups by Possvm but have been assigned to another

orthogroup here.

Figure 2: Evolution of *CYP2D, CYP2J,* and *CYP3A* gene subfamilies. Labeling conventions follow

those described for Figure 1.

Figure 3: *dN/dS*-based selection analyses of *CYP1A, CYP2D,* and *CYP2F* gene subfamilies. The

dashed gray lines in the bar graphs represent the points at which *dN/dS* (*i.e., ω*) = 1.00, which is

predicted by neutral evolution. Values less than 1.00 increasingly indicate purifying selection,

while *ω* > 1.00 indicate diversifying selection. The *ω* distributions of *CYP1* and *CYP2D* show

patterns of relaxed selection, while that for *CYP2F* indicates intensification of selection. Follow-

up HyPhy analysis using aBSREL (bottom-right quadrant) revealed a signal of diversifying

selection on the two terminal bamboo-lemur branches in the *CYP2F2* portion of this gene tree.