Title: Application of slow-controlled release fertilizer coordinates the carbon flow in carbon-nitrogen metabolism to effect rice quality.

Zhengrong Jiang[1, 2, 3], Qiuli Chen[1], Dun Liu[1, 2], Weike Tao[1, 2], Shen Gao[1, 2], Jiaqi Li[1, 2], Chunhao Lin[1, 2], Meichen Zhu[1, 2], Yanfeng Ding[1, 2], Weiwei Li[1, 2], Ganghua Li[1, 2], Soulaiman Sakr[3], Xueli Hong[1,4*]

[1] Sanya Institure of Nanjing Agriculture, Jiangsu Collaborative Innovation Center for Modern Crop Production, Key Laboratory of Crop Physiology Ecology and Production Management, Nanjing Agricultural University, Sanya, 572000, China

[2] China- Kenya belt and road joint laboratory on crop molecular biology

[3] Institut Agro, University of Angers, INRAE, IRHS, SFR QUASAV, Angers, 49000, France

[4] Institute of Agricultural Resources and Environment, Jiangsu Academy of Agricultural Sciences, Nanjing, 210095, China

*Corresponding author. E-mail: xuelihong@jaas.ac.cn, Tel: +86(025)-84391521.

E-mail address:

Zhengrong Jiang: zr.jiang@outlook.com

Qiuli Chen: 2019101047@njau.edu.cn

Dun Liu: 2020101041@stu.njau.edu.cn

Weike Tao: 2020201022@stu.njau.edu.cn

Shen Gao: g18862326620@163.com

Jiaqi Li: 2021101031@stu.njau.edu.cn

Chunhao Lin: 2021101030@stu.njau.edu.cn

Meichen Zhu: ZMC910824@163.com

Yanfeng Ding: dingyf@njau.edu.cn

Weiwei Li：li1990@njau.edu.cn

Ganghua Li: lgh@njau.edu.cn

Soulaiman Sakr: soulaiman.sakr@agrocampus-ouest.fr

Abstract:

Slow-controlled release fertilizers are experiencing a popularity in rice cultivation due to their effectiveness in yield and quality with low environmental costs. However, the underlying mechanism by which these fertilizers regulate grain quality remains inadequately understood. This study investigated the effects of five fertilizer management practices on rice quality in a two-year field experiment: CK, conventional fertilization, and four applications of slow-controlled release fertilizer (UF, urea formaldehyde; SCU, sulfur-coated urea; PCU, polymer-coated urea; BBF, controlled-release bulk blending fertilizer). In 2020 and 2021, the yields of UF and SCU groups showed significant decreases when compared to conventional fertilization, accompanied by a decline in nutritional quality. Additionally, PCU group exhibited poorer cooking and eating qualities. However, BBF group achieved increases in both yield (10.8 t hm^{-2} and 11.0 t hm^{-2}) and grain quality reaching the level of CK group. The sufficient nitrogen supply in both the PCU and BBF groups during the grain-filling stage led to a greater capacity for the accumulation of proteins and amino acids in the PCU group compared to starch accumulation. Intriguingly, BBF group showed better carbon-nitrogen metabolism than that of PCU group. The optimal nitrogen supply present in BBF group suitable boosted the synthesis of amino acids involved in the glycolysis/ tricarboxylic acid cycle, thereby effectively coordinating carbon-nitrogen metabolism. The application of the new slow-controlled release fertilizer, BBF, is advantageous in regulating the carbon flow in the carbon-nitrogen metabolism to enhance rice quality.

Key words: rice, slow-controlled release fertilizer, carbon, nitrogen, quality.

1. Introduction

Rice (*Oryza sativa* L.) is a crucial staple food crop globally, but its production is challenged by two key factors: grain development and grain quality (Thakur et al., 2011; Wang and Peng, 2017; Custodio et al., 2019). With the rapid economic development, there is a growing demand for high-quality rice in developing countries across Asia (Adjao and Staatz, 2015; Concepcion et al., 2015). Grain quality characteristics are essential for determining market value and include factors such as nutritional value, physical appearance, cooking and sensory properties, and milling recovery (Fitzgerald et al., 2009). Fertilization plays a critical role in regulating the grain quality of rice as it affects the content and composition of starch and protein in the grain (Gu et al., 2015). Over the past few decades, conventional fertilizers in rice cultivation face challenges such as low efficiency, high wastage, and labor-intensive application methods, causing environmental pollution and economic losses (Chen et al., 2014). To address these issues, the application of slow-controlled release fertilizers has been proven to be a more efficient method of fertilization, supplying crops with required nutrients in a single basal application throughout the growth period (Naz and Sulaiman, 2016; Qiong et al., 2021). The slow-controlled release fertilizer has a dramatic influence on optimizing grain quality and nitrogen use efficiency, and are widely adopted to minimize economic losses and reduce environmental pollution (Ke et al., 2017; Gao et al., 2021).

A large number of slow-controlled release fertilizers, including urea formaldehyde (UF), sulfur-coated urea (SCU), polymer-coated urea (PCU), and controlled-release bulk blending fertilizer (BBF), have been developed to meet rice's nutrient demands, enhance yield and quality (Wu et al., 2021). UF and SCU are slow-release fertilizers made from easily decomposable organic materials. UF fertilizer is produced by combining urea with formaldehyde, resulting in quick nitrogen release during the early growth stage but limited release during the middle and late stages (Shaviv, 2001). In contrast, SCU fertilizer involves coating urea granules with a layer of sulfur, enabling the nutrients to be slowly released over time; however, the release rate is insufficient during the middle and late stages (Xu et al., 2016). PCU and BBF are two types of controlled-release fertilizers that provide a regulated-release of nutrients to plant. The PCU fertilizer is an enhanced version of SCU, designed to minimize the impact of microorganisms on nutrient release through coating urea granules with a polymer material. BBF is created by blending various types of slow-release and controlled-release fertilizers, allowing for a consistent release of nitrogen nutrients throughout the entire rice growth cycle (Ye et al., 2013). Given the growing demand for grain

quality and efficient fertilizer use, further research is needed to examine the relationship between grain

quality and the application of slow-controlled release fertilizers.

Grain filling is vital for rice quality and is highly influenced by fertilizer use. Accumulation of lipids,

starch, and storage proteins during grain filling stage is crucial for grain quality, and is strongly influenced

by nitrogen availability (Jiang et al., 2021; Wang et al., 2021b; Xu et al., 2021). The improper use of

fertilizer can negatively impact the physical appearance of rice grain, leading to the formation of

chalkiness as a result of an unbalanced carbon-nitrogen metabolism (Zhao et al., 2019; Wang et al., 2021b).

The carbon-to-nitrogen (C: N) balance in rice grains refers to the optimal ratio of protein and starch in the

grain, which is critical for determining the nutritional value and other rice qualities (Dou et al., 2017). The

changes in nitrogen and carbon supply can alter numerous central metabolites involved in carbon and

nitrogen metabolism in parallel (Gao et al., 2020). To support the synthesis of amino acids and ensure

adequate growth in rice grain, an appropriate nitrogen content is needed to drive the production of carbon

skeletons and provide enough amino acids (Gutiérrez et al., 2007; Wang et al., 2022a).

The inorganic nitrogen of soil is absorbed by roots and can then be assimilated into glutamine and

glutamate in grain, which serves as a source of organic nitrogen (Huang et al., 2021). Glutamine synthetase

(GS), encoded by $OsGS2$, is a crucial enzyme for assimilating NH_4^+ in rice. Maintaining a stable GS level

is essential for maintaining the carbon-nitrogen balance during rice growth (Bao et al., 2015). The key

enzymes of the GS and GOGAT (glutamate synthase) cycle play key roles in nitrogen assimilation of rice

grain (Luo et al., 2018; Poucet et al., 2021). Additionally, nitrogen assimilation requires energy and carbon

skeletons, which are derived from sucrose, glucose, other glycolysis-derived carbohydrates (pyruvate) and

the tricarboxylic acid (TCA)-derived organic acids (eg., cetoglutarate and oxaloacetate) (Huang et al.,

2021). The carbon and nitrogen metabolites are also involved in the oxidative pentose phosphate pathway

(OPPP), which is crucial for generating the energy and reducing power required for rice grain growth (Li et

al., 2016; Yang et al., 2016; Wang et al., 2021a). The TCA cycle is essential for providing 2-oxoglutarate,

which is necessary for the production of glutamate and glutamine involved in nitrogen assimilation

(Nunes-Nesi et al., 2010). The intermediates of the glycolysis pathway, like pyruvate, play an important

role in the TCA cycle and nitrogen assimilation in rice (Masumoto et al., 2010).

The primary product of photosynthesis, sucrose, plays a crucial role in supplying the carbon skeletons

required for synthesizing a variety of macromolecules, such as proteins, nucleic acids, lipids, and starch

(Jiang et al., 2021). This is facilitated by the transport pathways of *OsCIN4* (invertase) and *OsSUT2*

(sucrose transporter) during grain filling (Wang et al., 2008; Deng et al., 2021). The major function of

SuSase is to breakdown sucrose to supply carbohydrates for the process of OPPP and glycolysis/TCA

cycle, and the enzymes AGPase, GBSS and SBE play major roles in catalyzing the synthesis of starch to

promote grain filling (Wang et al., 2021a; Xu et al., 2022). However, the mechanisms linking carbon and

nitrogen metabolism to rice grain growth under the influence of slow-controlled release fertilizer are not

yet fully understood.

In this study, a two-year field experiment was conducted to examine the impact of various slow-controlled

release fertilizers on grain quality of rice. To gain a deeper understanding of the nitrogen assimilation

process in rice grain under the influence of slow-controlled release fertilizers, we also analyzed the

underlying carbon-nitrogen mechanisms during grain filling. Our findings will provide a solid foundation

for improving the quality and yield of rice through the use of slow-controlled release fertilizers.

2.   Materials and methods

2.1 Study site and N sources

Field experiments were conducted over two rice growing seasons (2020 and 2021) in Yanling Town of

Danyang city, Jiangsu Province, China (31°54′31″N, 119′28′21″E). The meteorological data came from the

meteorological station (Watch Dog 2900ET, SPECTRUM, USA) installed 100 m from the experimental

station (Figure.S1). The soil properties were classified as Orthic Acrisol (Falsone, 2013). The soil had the

following characteristics: pH = 6.4, organic matter = 18.32 g·kg^{-1}, total N = 1.27 g·kg^{-1}, Olsens-P = 16.85

mg·kg^{-1} and NH$_4$OAc-extractable K = 139.63 mg·kg^{-1}. The slow-controlled release fertilizers used in this

study were: UF (35% N), SCU (37% N), PCU (43% N), and BBF (40% N). The UF, SCU, PCU and BBF

were provided by Hanfeng Slow Release Fertilizer Co., Ltd. (Jiangsu, China). Conventional urea (46% N)

fertilizers were used as CK group for comparison.

2.2 Experimental design

For analyzing effects of different controlled-release fertilizers on grain quality of rice, the experiment was

conducted by rice cultivation of Ningjing 8th (*Oryza sativa* L.), a high-yielding japonica rice cultivar bred

by Nanjing Agricultural University (NAU). This experiment included treatments as follows: a conventional fertilization (CK, four spilt applications of urea), and a single basal application of slow-controlled release fertilizer (SCU, PCU, UF, and BBF). The N release rate of the slow-controlled release fertilizer was shown in Figure. 1, which was determined by buried bag method (Yang et al., 2011). The conventional urea fertilization was split among basic fertilizer, tiller fertilizer, spikelet-promoting fertilizer, and spikelet-developing fertilizer. Each treatment was applied nitrogen at 210 kg·ha^{-1}, and for all treatments, P and K fertilizers were applied as basal dressings at 135 kg·ha^{-1} (P$_2$O$_5$) and 216 kg·ha^{-1} (K$_2$O). The field plots, measuring 7.2 m × 20 m, were arranged in a randomized complete block design with three replicates per treatment. In each replicate, a total of approximately 1600 panicles with similar growth patterns were labeled and monitored for their flowering date. This was done in September, when about 50% panicles had fully emerged from the flag leaf sheath.

2.3 Sampling and analysis

2.3.1 Harvesting and grain quality measurements

At maturity stage, all necessary plants (excluding edge-row plants) from each plot were harvested. The grain yield was determined by multiplying each yield component, which had been measured after removing impurities and adjusting moisture content to 14.5%. Before harvesting, the number of effective tillers per hill was determined by selecting 50 plants from each plot. At maturity, approximately 30 marked plants were used to analyze the 1000-grain weight, seed setting rate, and grain number per panicle.

At maturity, the rice grain quality was analyzed. For rice milling quality, the brown rice yield and milled yield were analyzed according to the method of rice measurement standards (NY147-88; Ministry of Agriculture, PR China, 1988). To test the appearance quality of rice, the chalk characteristics of brown rice were determined by the cleanliness test-bed according to Tang et al. (2019). Rice cooking and eating quality was analyzed by the Rapid Visco Analyser (RVA, Starchmaster 2, Perten Instruments of Australia Pty Ltd., Sydney, New South Wales, Australia) to obtain profile characteristics according to Standard Method AACC61-02. Viscous profile characteristics of RVA were expressed as peak viscosity, hot viscosity, cool viscosity, breakdown viscosity, setback (difference between final viscosity and peak viscosity), and consistency. Rice taste characteristics were analyzed by an STA-1A device (Satake Corp., Hiroshima, Japan) according to the method of Jin et al. (2004).

2.3.2 Grain weight and grain growth rate

To analyze the grain filling process, 90 tagged panicles of each replicate plot were sampled every 7 days post anthesis (DPA) from 7 to 42 DPA (7, 14, 21, 28, 35, and 42 DPA). All spikelets were deactivated at 105°C for 60 min and then dried at 80°C to determine the grain dry weight (DW). Grain filling processes were fit to Richards's growth equation (Richards, 1959).

$$W = \frac{A}{(1+Be^{-kt})^{1/N}} \quad (1) \qquad R = \frac{AkBe^{-kt}}{N(1+Be^{-kt})^{(N+1)/N}} \quad (2)$$

Grain weight (mg), W; Grain filling rate, R; Final grain weight (mg), A; Time after anthesis (days), t; and B, k, and N are coefficients established from the regression of the equation.

2.3.3 Analyze of sugar, starch and storage protein composition

To analyze the sugar and starch content, all samples were first ground into a fine powder and then passed through a 100-mesh sieve. The sugar and starch were extracted using a modified version of the method described by Yang et al. (2001). Approximately 100 mg of the ground sample was mixed with 8 mL of 80% (v/v) ethanol at 80°C for 30 min. After cooling, the tube was centrifuged at 5000 g for 15 min, and the supernatant was collected. The extraction process was repeated three times, and the sugar extract was diluted with distilled water to a final volume of 50 mL. After the sugar extraction, the residues in the tubes were dried at 80°C to extract starch using $HCLO_4$ according to Yang's method. The amylose and amylopectin contents were measured using the Megazyme Amylose/Amylopectin Assay Kit (Megazyme International Ireland Ltd, Bray, Ireland).

The total protein content was measured using an AutoKjeldahl Unit K-370 nitrogen analyzer (Büchi Labortechnik, Flawil, Switzerland) with AACC International Approved Method 46-30.01. Different protein components including albumin, globulin, prolamin, and glutelin were extracted in a stepwise manner using different solvents such as distilled water, dilute hydrochloric acid, ethanol, and dilute alkali. The glutelin content was determined using the biuret colorimetric method, while the remaining protein components were measured using the Coomassie brilliant blue colorimetric method.

2.3.4 Measurements of amino acid composition

To hydrolyze the rice powder samples, approximately 10 mg of the sample was mixed with 1 mL of 6 N

HCl (Sigma, USA) in a 2 mL screw-cap tube. After heating at 110°C for 24 h, the samples were further

treated for 6 h at 65°C. The resulting residue was dissolved in 1 mL of Na-S™ buffer, and the mixture was

centrifuged at $1600 \times g$ for 10 min. The supernatant was filtered through a 0.45 μm nylon membrane

syringe filter (Pall Life Sciences, USA), and about 10 nmol of L-(+)-norleucine (Wako Pure Chemicals,

Japan) was added. Amino acid analysis was performed using a Hitachi L-8900 amino acid analyzer

(Hitachi Corp, Japan) based on the national standard of the People's Republic of China (Ministry of

Agriculture PR China, 1988). The amino acids tested included aspartic acid (Asp), glutamic acid +

glutamine (Glu), glycine (Gly), histidine (His), isoleucine (Ile), leucine (Leu), lysine (Lys), phenylalanine

(Phe), serine (Ser), threonine (Thr), tyrosine (Tyr), and valine (Val). Total amino acid content was the sum

of all amino acid content in the same period. All experiments were conducted at least three biological

replicates in each sample, and the HPLC data were normalized to the level of L- (+)-norleucine per sample.

2.3.5   Enzyme extraction and analysis

For enzyme extraction, the grains collected at 7, 14, 21, 28, 35, and 42 DPA were ground into fine powder

in liquid nitrogen. The activities of several enzymes including sucrose synthase (SuSase), adenine

diphosphoglucose pyrophosphorylase (AGPase), starch branching enzyme (SBE), and granule-bound

starch synthase (GBSS) were determined using the method described by Yang et al. (2003). In addition, the

activities of glutamine synthetase (GS) and glutamate synthetase (GOGAT) were analyzed according to the

protocols outlined by Tang et al. (2018).

2.3.6   RNA extraction and qRT-PCR

Total RNA was extracted from frozen rice grains using a previously described method (Yasuda et al.,

2005). Gene transcription levels of the relevant genes were analyzed using RNase-free DNase I treatment,

cDNA synthesis, and quantitative real-time polymerase chain reaction (qRT-PCR). The RNA-prep pure

PLANT Kit (DP432, Tiangen Biotek, Beijing, China) was used to isolate total RNA from the rice grains,

which was then reverse-transcribed into first-strand cDNA using the Prime-Script-TM RT Reagent Kit

(RR036, Takara, Kyoto, Japan). The quantitative real-time polymerase chain reaction was performed using

the ABI 7300 sequencer and SYBR Premix Ex Taq-TM (RR420, Takara, Kyoto, Japan) according to the

manufacturer's protocol. Each sample was replicated three times to test the expression of genes (*OsSUT2*,

OsCIN4, and *OsGS2*). The primers used in this study are listed in the Supporting Information (Table S1).

2.4 Statistical analysis

Excel 2019 (Microsoft Corp. Redmond, WA, USA), SPSS version 20.0. (SPSS Statistics, SPSS Inc.,

Chicago, USA), and Origin 2021 (OriginLab, Northampton, MA, USA) were used for data visualization.

Data was performed the analysis by using two-way analysis of variance (ANOVA) and principal

component analysis (PCA). Means were compared based on the least significant difference at $P = 0.05$

$(LSD_{0.05})$.

3. Results

3.1 Influence of slow-controlled release fertilizer on rice yield

Compared to the conventional fertilizer's yield (10.2 t hm^{-2} and 11.2 t hm^{-2}) in 2020 and 2021, the rice

yields decreased to 8.4 t hm^{-2} and 9.7 t hm^{-2} after application of urea formaldehyde fertilizer (UF), in

consistent with the decline in yield (9.6 t hm^{-2} and 8.6 t hm^{-2}) of sulfur-coated urea fertilizer (SCU) (Table

1). The groups of PCU and BBF showed similar or even higher yield parameters compared with CK group

(Table 1). In 2020, the yield of the BBF group reached 10.8 t hm^{-2}, marking the highest yield among all

types of fertilizers. Similarly, in 2021, the BBF yield increased to 11.0 t hm^{-2}, significantly surpassing the

yields of the UF and SCU groups.

The application of UF, SCU, PCU, and BBF all resulted in a significant improvement in grain weight

compared with the CK group in 2019 and 2020 (Table 1). The differences in grain weight among the

various fertilizer group were even more conspicuous in 2021, particularly with the application of BBF and

UF significantly promoting grain weight accumulation in comparison to CK group (Figure 2).

Concurrently, the grain filling rate of BBF and UF groups was significantly higher than that of other

treatments at the middle grain-filling stage (14 DPA). These results demonstrate that the use of suitable

slow-controlled release fertilizers, such as BBF, can enhance grain growth and rice yield.

3.2 Effects of slow-controlled release fertilizer on rice quality

In 2020 and 2021, the application of BBF significantly improved the physical appearance, cooking and

eating qualities, as well as the nutritional value of rice, reaching levels comparable to those of the CK

group (Tables 2-3). Nevertheless, the taste values for the PCU group were noticeably lower at 65 and 73 in

2020 and 2021, respectively, compared to the CK group's values of 72.7 and 75 (Table 2). Additionally, the

PCU group displayed notably lower cool viscosity and breakdown than the CK group (Table 2). The taste

value is critical for cooking and eating qualities and is tightly correlated with the RVA value (Kesarwani et

al., 2016). These results indicate that the poor taste value and RVA value resulted in poor cooking and

eating qualities in PCU group. The composition of protein and amino acids greatly contributes to the

nutritional value of rice grain (Amagliani et al., 2017). Compared to the CK group, the application of UF

and SCU resulted in a lower content of nitrogen metabolites in rice grains (Table 3), indicating that the use

of UF and SCU was unfavorable to the nutritional value of rice grain.

3.3 Carbon metabolism of rice grain

The carbon metabolism pathway plays a crucial role in rice quality by facilitating the synthesis of starch

and protein (Dai et al., 2013; Jiang et al., 2021). The sugar content of all slow-controlled release fertilizer

treatments reached or even exceeded the level of CK group at grain-filling stage in 2021 (Figure. 3), while

the PCU group showed significantly lower accumulation of starch and amylopectin compared to the CK

treatment during grain-filling period (Table 3, Figure. 3). The application of slow-controlled release

fertilizers obviously affected the activity of key enzymes involved in carbon-nitrogen metabolism in rice

grains (Table S2). The PCU group did not exhibit the lowest enzyme activities related to carbon

metabolism (SuSase, AGPase, GBSS, and SBE) in grains among all treatments, but the UF group

demonstrated lower enzyme activities during grain filling (Figure. 4). However, the starch content in the

UF group was not the lowest (Figure. 4). The carbon metabolite content and the related key enzyme

activity of BBF were neither the highest nor the lowest among these treatments (Figures. 3-4).

3.4 Nitrogen metabolism of rice grain

The formation of amino acids and proteins through nitrogen metabolism process is crucial for the growth

of rice grains (Yamakawa and Hakata, 2010). During grain filling, the UF group had significantly lower

accumulation of total protein and amino acid in the grain compared to the CK group, whereas the PCU

group's content reached or surpassed the level of CK group in both 2020 and 2021 (Figure. 5). Relative to

the CK group, the UF group exhibited significantly lower activities in key enzymes (GS and GOGAT)

associated with nitrogen assimilation, whereas the PCU group demonstrated comparable activities in GS

and GOGAT to those of the CK group (Figure. 6). During grain filling, the key metabolites and enzyme

activities associated with nitrogen metabolism in the grain of both BBF and SCU groups were not

significantly higher than those in the PCU group, nor were they lower than those in the UF group (Figures.

5-6).

3.5 The crosstalk of carbon and nitrogen metabolism in rice grain

In the dynamic analysis, profiles of sugar-unloading pathway and carbon-nitrogen metabolism were tested

during grain filling (Figure. 7). During grain filling, the gene expression of *OsSUT2* and *OsCIN4*, which

are related sugar unloading (Ji et al., 2005; Jiang et al., 2021), did not significantly decrease in the groups

of UF, PCU and BBF compared to the CK group. This suggested that the application of these fertilizers did

not limit the supply of sucrose. The PCU group showed the strongest accumulation of 12 amino acids in

rice grain compared to other slow-controlled release fertilizers, and reached the level of the CK group

(Figures. 7 and S2). Meanwhile, the UF group showed the lowest accumulation of amino acids among all

fertilizer treatments. The accumulation of the BBF group was not higher than that of the PCU group, but

not lower than the UF group. The *OsGS2* has a significant impact on both the nitrogen transportation in

rice and the metabolism of amino acids in grains (Wang et al., 2022b). During grain filling, the gene

expression of *OsGS2* was higher in the groups of PCU and BBF than that of UF and SCU groups. Thus,

the increased nitrogen supply in groups of PCU and BBF might enhance nitrogen assimilation and led to

an increase in amino acid synthesis of rice grains at the grain filling stage (Table S3, Figures. 1 and S2).

4.   Discussion

4.1 Rice yield and quality are improved by the application of optimal slow-controlled release fertilizer

Compared to the conventional fertilizer, the UF and SCU groups exhibited a significantly lower number of

spikelets per panicle and area, which notably inhibited rice yield (Table 1). Applying an optimal nitrogen

source close to the grain-filling stage can effectively promote yield formation (Nasielski and Deen, 2019).

Compared to the UF and SCU fertilizers, the PCU and BBF fertilizers consistently released nitrogen

throughout the growth stage, providing more nitrogen for panicle differentiation and grain filling (Figure.

1). Intriguingly, the rice yield of PCU and BBF groups did not show significant differences compared to

CK group, and the application of slow-controlled release fertilizers obviously improved grain weight

(Table 1, Figure 2). In 2020, the BBF group achieved the highest yield of 10.8 t hm^{-2} among all fertilizers

(Table 1). In 2021, the BBF yield increased to 11.0 t hm^{-2}, significantly surpassing the UF and SCU

groups' yields (Table 1).

Variations in nitrogen supply significantly impact the distribution and buildup of starch and protein during

grain filling, ultimately affecting grain quality (Wang et al., 2021b; Xu et al., 2021). Compared to

conventional fertilizer, the application of UF and SCU fertilizers led to a significant reduction in nutritional

value, while the PCU group exhibited lower cooking and eating quality (Tables 2-3). The low nutritional

value observed in the UF and SCU groups might be attributed to insufficient nitrogen supply during the

grain filling stage (Figure. 1), based on the report that inadequate nitrogen reduced synthesis of protein and

amino acids (Wang et al., 2021b).   In addition, the cooking and eating quality of rice can be adversely

reduced by high nitrogen application (Shi et al., 2023), the low quality of the PCU group might be

attributed to the release of excess nitrogen during the grain filling stage (Table 2, Figure. 1). Conversely,

the application of BBF fertilizer led to good grain quality in terms of physical appearance, cooking and

eating qualities, and nutritional value (Tables 2-3). This was achieved due to its optimal nitrogen release

during grain filling (Figure. 1), with each quality-related index reaching those of CK group and surpassing

other fertilizer treatments (Tables 2-3). Hence, to ensure a high rice yield and quality, it is crucial to

employ an optimal slow-controlled release fertilizer, such as BBF.

4.2 The carbon and nitrogen balance in rice grains is influenced by the type of slow-controlled

release fertilizer

The accumulation of starch, protein, and amino acids determines grain quality, and this process requires

significant amounts of carbon and nitrogen metabolites (Yamakawa and Hakata, 2010; Jiang et al., 2021;

Liu et al., 2021). The application of slow-controlled release fertilization did not hinder sucrose

accumulation but led to a decrease in starch and amylopectin content in the PCU group compared to the

CK group (Table 2, Figure. 3). Meanwhile, the PCU group showed no reduction in enzyme activity related

to carbon metabolism (Figure. 4). In contrast, the UF group decreased the activities of SuSase, AGPase,

and SBE, but did not show lower starch and amylopectin content compared to the PCU group (Table 2,

Figures. 3-4). Carbon metabolism is crucial in nitrogen metabolism as it provides the carbon skeletons

necessary for amino acid synthesis, including pyruvate, oxaloacetate, and alpha-ketoglutarate (Xu et al.,

2012). Thus, the complex results observed in this study might be linked to the interplay between carbon

and nitrogen metabolism (Baunsgaard et al., 2005; Seung et al., 2013; MacNeill et al., 2017).

The PCU and BBF fertilizers had a higher nitrogen-release ability during grain filling compared to UF and

SCU fertilizers (Ye et al., 2013; Wu et al., 2021). Intriguingly, the application of UF resulted in a

significant reduction in accumulation of protein and amino acids compared to conventional fertilizer,

whereas PCU fertilizers achieved levels similar to those of the CK group during grain filling (Figure. 5).

The GS/GOGAT cycle plays a key role in nitrogen assimilation (Gao et al., 2019; Hou et al., 2019). During

grain filling, the PCU group showed increased activity in GS and GOGAT enzymes, reaching levels

comparable to those of the CK group, whereas the UF group showed significantly lower activity in these

enzymes (Figures. 5-6). These results indicated that increased nitrogen supply mainly leads to increased

synthesis of protein and amino acids in rice grains rather than increased accumulation of storage starch.

However, the BBF group showed a better partition in starch and protein content, which was neither the

highest nor the lowest compared to others (Figures. 3-6). This phenomenon may be related to the

appropriate nitrogen release ability of BBF fertilizer during grain filling (Figure. 1). Despite the complex

relationship between protein production and carbon-nitrogen metabolism, the dynamic changes in protein

synthesis are significantly related to the application of slow-controlled release fertilizers (Table S3, Figure.

S2).

4.3 Key steps for promoting carbon-nitrogen crosstalk in developing grain under the application of

slow-controlled release fertilizer

Maintaining a balance between carbon and nitrogen metabolism is crucial for grain filling of rice, as

nitrogen, functioning as a signaling molecule, influences plant metabolism and physiology through

changes in gene expression (Xin et al., 2019). During the grain-filling stage, rice grain undergoes dynamic

metabolic adjustments to meet its nutritional demands (Jiang et al., 2022a). An analysis of the primary

changes in amino acid metabolism revealed clear differences in the carbon and nitrogen metabolism among

various slow-controlled release fertilizers (Figure. 7).

The expression of genes (*OsSUT2* and *OsCIN4*) related to sucrose unloading (Sekhar et al., 2021), as well

as sucrose content, in the UF, PCU, and BBF groups were not lower than that of CK group (Figure. 3 and

7), suggesting that the availability of carbohydrates was not restricted by these application of slow-

controlled release fertilizers. N-nutrient/metabolites and carbon metabolism are intricately interconnected

(Yamakawa and Hakata, 2010). Carbon metabolism plays a critical role in incorporating nitrogen into cell metabolism, primarily through the oxidative pentose phosphate pathway (OPPP) and glycolysis/ the tricarboxylic acid (TCA) cycle (Bussell et al., 2013; Yang et al., 2016; Wang et al., 2021a; Jiang et al., 2022b). When comparing various types of slow-controlled release fertilizers, it was found that the application of PCU and BBF fertilizers led to a higher capacity for sucrose unloading and nitrogen assimilation (Figures. 7 and S2). In the metabolic analysis, the increased carbon flux was primarily directed towards the glycolysis and TCA cycle for amino acid synthesis in the PCU and BBF groups (Figure. 6 and S2). As the increase of nitrogen supply in PCU and BBF (Figure. 1), the GS/GOGAT cycle was significantly promoted in rice grain at the filling stage (Table S2, Figures. 7 and S2). The PCU group demonstrated significantly higher enzyme activity in carbon and nitrogen metabolism and a lower starch content in rice grain, compared to the UF and SCU groups (Figures. 3-6). A high supply of nitrogen can increase the sugar unloading and metabolic utilization (Zhang et al., 2021). Consequently, the PCU group's high capacity for nitrogen assimilation lead to efficient utilization of carbohydrates for amino acid synthesis during grain filling. Interestingly, the BBF group showed a relatively balanced performance in key enzymes of carbon and nitrogen metabolism, as well as in the levels of starch, protein, and amino acids (Figures. 3-6). These results suggest that a suitable supply of nitrogen promotes both grain filling and maintains an optimal carbon-nitrogen state in the rice grain. Therefore, the application of slow-controlled release fertilizer like BBF is a practical solution for manipulating the balance between carbon and nitrogen in rice.

Overall, these observations suggest that an increase in nitrogen supply not only enhances nitrogen metabolism in grain, but also improves carbon flow (Table S2, Figures. 7 and S2). Our study proposes a model in which nitrogen supply regulates the networks of carbon-nitrogen metabolism, and the regulation of starch and protein synthesis under high nitrogen conditions is linked to nitrogen assimilation and the glycolysis/TCA cycle (Table S3, Figure. 7 and S2).

4.4 Conclusions

The application of BBF fertilizer not only increased rice yield but also enhanced grain quality compared to other slow-release fertilizers, such as UF, SCU, and PCU. Moreover, it has the potential to reach or even exceed the levels achieved with conventional fertilization. This can be primarily attributed to BBF

fertilizer's superior regulation of the carbon-nitrogen balance in rice grains compared to other slow-
controlled release fertilizers. The application of BBF fertilizer appropriately increases nitrogen assimilation
(amino acid synthesis and nitrogen transport) in the grain by modulating carbon flow in the carbon
metabolism of grain (e.g., glycolytic metabolism and the TCA cycle). Our research provides new insight
into the relationship among the nitrogen supply, metabolism pathway of carbon and nitrogen, and grain
growth to finely promote grain quality under the application of slow-controlled release fertilizer in rice.

Abbreviations:
CK: Conventional fertilization
UF: urea formaldehyde fertilizer
SCU: sulfur-coated urea fertilizer
PCU: polymer-coated urea fertilizer
BBF: controlled-release bulk blending fertilizer
TCA: the tricarboxylic acid
NUE: nitrogen use efficiency
GS: Glutamine synthetase
GOGAT: glutamate synthase
OPPP: oxidative pentose phosphate pathway
SuSase: Sucrose Synthase
AGPase: ADP-glucose pyrophosphorylase
GBSS: granule-bound starch synthase
SBE: starch branching enzyme
Glu: glutamine
Gly: glycine
His: histidine
Ile: isoleucine
Leu: leucine
Lys: lysine
Phe: phenylalanine
Ser: serine
Thr: threonine
Tyr: tyrosine
Val: valine
DPA: Days post-anthesis
RVA: rapid viscosity analyzer
DW: Dry weight.

Declarations

Ethics approval and consent to participate

Not applicable

Availability of Supporting Data

The datasets generated and/or analysed during the current study are available from the corresponding author on reasonable request.

Competing interest

The authors declare no conflicts of interest.

Funding

This work was funded by the National Key Research and Development Program of China (No. 2021YFD1700803), and the Province Key Research and Development Program of Jiangsu, China (D21YFD17008).

Author Contributions:

ZJ and QC: Conceptualization, Software, Formal analysis, Investigation, Writing - Original Draft and Visualization; DL, WT, SG, and JL: Validation and Investigation. CL and MZ: Data Curation; YD, and WL: Resources and Supervision; GL and SS: Conceptualization and Revising - Original Draft; LX: Draft revision, Project administration and Funding acquisition. All authors read and approved the final manuscript.

Acknowledgments

The authors thank Maria-Dolores Pérez-Garcia, and Laurent Ogé for kindly providing guidance and Gaoya Sun for providing technical assistance.

References

Adjao, R., Staatz, J., 2015. Asian rice economy changes and implications for sub-Saharan Africa. Global Food Security 5, 50-55.

Amagliani, L., O'Regan, J., Kelly, A., O'Mahony, J., 2017. Composition and protein profile analysis of rice protein ingredients. Journal of Food Composition and Analysis 59, 18-26.

Bao, A., Zhao, Z., Ding, G., Shi, L., Xu, F., Cai, H., 2015. The stable level of glutamine synthetase 2 plays an important role in rice growth and in carbon-nitrogen metabolic balance. International Journal of Molecular Science 16, 12713-12736.

Baunsgaard, L., Lütken, H., Mikkelsen, R., Glaring, M.A., Pham, T.T., Blennow, A., 2005. A novel isoform of glucan, water dikinase phosphorylates pre-phosphorylated α-glucans and is involved in starch degradation in Arabidopsis. The Plant Journal 41, 595-605.

Bussell, J., Keech, O., Fenske, R., Smith, S., 2013. Requirement for the plastidial oxidative pentose phosphate pathway for nitrate assimilation in Arabidopsis. The Plant Journal 75, 578-591.

Chen, X., Cui, Z., Fan, M., Vitousek, P., Zhao, M., Ma, W., Wang, Z., Zhang, W., Yan, X., Yang, J., Deng, X., Gao, Q., Zhang, Q., Guo, S., Ren, J., Li, S., Ye, Y., Wang, Z., Huang, J., Tang, Q., Sun, Y., Peng, X., Zhang, J., He, M., Zhu, Y., Xue, J., Wang, G., Wu, L., An, N., Wu, L., Ma, L., Zhang, W., Zhang, F., 2014. Producing more grain with lower environmental costs. Nature 514, 486-489.

Concepcion, J., Ouk, M., Zhao, D., Fitzgerald, M., 2015. The need for new tools and investment to improve the accuracy of selecting for grain quality in rice. Field Crops Research 182, 60-67.

Custodio, M., Cuevas, R., Ynion, J., Laborte, A., Velasco, M., Demont, M., 2019. Rice quality: How is it defined by consumers, industry, food scientists, and geneticists? Trends in Food Science & Technology 92, 122-137.

Dai, Z., Léon, C., Feil, R., Lunn, J., Delrot, S., Gomès, E., 2013. Metabolic profiling reveals coordinated switches in primary carbohydrate metabolism in grape berry (*Vitis vinifera* L.), a non-climacteric fleshy fruit. Journal of Experimental Botany 64, 1345-1355.

Deng, Y., Yu, Y., Hu, Y., Ma, L., Lin, Y., Wu, Y., Wang, Z., Wang, Z., Bai, J., Ding, Y., Chen, L., 2021. Auxin-mediated regulation of dorsal vascular cell development may be responsible for sucrose phloem unloading in large panicle rice. Frontiers in Plant Science 12, 630997.

Dou, Z., Tang, S., Li, G., Liu, Z.H., Ding, C., Chen, L., Wang, S., Ding, Y., 2017. Application of nitrogen fertilizer at heading stage improves rice quality under elevated temperature during grain-filling stage. Crop Science 57.

Falsone, G., 2013. Keys to Soil Taxonomy 2010.

Fitzgerald, M., McCouch, S., Hall, R., 2009. Not just a grain of rice: the quest for quality. Trends in Plant Science 14, 133-139.

Gao, Y., Song, X., Liu, K., Li, T., Zheng, W., Wang, Y., Liu, Z., Zhang, M., Chen, Q., Li, Z., Li, R., Zheng, L., Liu, W., Miao, T., 2021. Mixture of controlled-release and conventional urea fertilizer application changed soil aggregate stability, humic acid molecular composition, and maize nitrogen uptake. Science of The Total Environment 789, 147778.

Gao, Y., Xu, Z., Zhang, L., Li, S., Wang, S., Yang, H., Liu, X., Zeng, D., Liu, Q., Qian, Q., Zhang, B., Zhou, Y., 2020. *MYB61* is regulated by *GRF4* and promotes nitrogen utilization and biomass production in rice. Nature Communications 11, 5219.

Gao, Z., Wang, Y., Chen, G., Zhang, A., Yang, S., Shang, L., Wang, D., Ruan, B., Liu, C., Jiang, H., 2019. The indica nitrate reductase gene *OsNR2* allele enhances rice yield potential and nitrogen use efficiency. Nature Communications 10, 5207.

Gu, J., Chen, J., Chen, L., Wang, Z., Zhang, H., Yang, J., 2015. Grain quality changes and responses to nitrogen fertilizer of japonica rice cultivars released in the Yangtze River Basin from the 1950s to 2000s. The Crop Journal 3, 285-297.

Gutiérrez, R., Lejay, L., Dean, A., Chiaromonte, F., Shasha, D., Coruzzi, G., 2007. Qualitative network models and genome-wide expression data define carbon/nitrogen-responsive molecular machines in Arabidopsis. Genome Biology 8, 1-13.

Hou, W., Xue, X., Li, X., Khan, M.R., Yan, J., Ren, T., Cong, R., Lu, J., 2019. Interactive effects of nitrogen and potassium on: Grain yield, nitrogen uptake and nitrogen use efficiency of rice in low potassium fertility soil in China. Field Crops Research 236, 14-23.

Huang, X., Zhang, Y., Wang, L., Dong, X., Hu, W., Jiang, M., Chen, G., An, G., Xiong, F., Wu, Y., 2021. *OsDOF11* affects nitrogen metabolism by sucrose transport signaling in rice (*Oryza sativa* L.). Frontiers in plant science 12, 703034.

Ji, X., Raveendran, M., Oane, R., Ismail, A., Lafitte, R., Bruskiewich, R., Cheng, S., Bennett, J., 2005. Tissue-specific expression and drought responsiveness of cell-wall invertase genes of rice at flowering. Plant Molecular Biology 59, 945.

Jiang, Z., Chen, Q., Chen, L., Liu, D., Yang, H., Xu, C., Hong, J., Li, J., Ding, Y., Sakr, S., Liu, Z., Jiang, Y., Li, G., 2022a. Sink strength promoting remobilization of non-structural carbohydrates by activating sugar signaling in rice stem during grain filling. International Journal of Molecular Science 23, 4864.

Jiang, Z., Chen, Q., Chen, L., Yang, H., Zhu, M., Ding, Y., Li, W., Liu, Z., Jiang, Y., Li, G., 2021. Efficiency of sucrose to starch metabolism is related to the initiation of inferior grain filling in large panicle rice. Frontiers in plant science 12.

Jiang, Z., Wang, M., Nicolas, M., Ogé, L., Pérez-Garcia, M.-D., Crespel, L., Li, G., Ding, Y., Le Gourrierec, J., Grappin, P., Sakr, S., 2022b. Glucose-6-phosphate dehydrogenases: the hidden players of plant physiology. 23, 16128.

Ke, J., He, R., Hou, P., Ding, C., Ding, Y., Wang, S., Liu, Z., Tang, S., Ding, C., Chen, L., Li, G., 2018. Combined controlled-released nitrogen fertilizers and deep placement effects of N leaching, rice yield and N recovery in machine-transplanted rice. Agriculture, Ecosystems & Environment 265, 402-412.

Ke, J., Xing, X., Li, G., Ding, Y., Dou, F., Wang, S., Liu, Z., Tang, S., Ding, C., Chen, L., 2017. Effects of different controlled-release nitrogen fertilisers on ammonia volatilisation, nitrogen use efficiency and yield of blanket-seedling machine-transplanted rice. Field Crops Research 205, 147-156.

Kesarwani, A., Chiang, P., Chen, S., 2016. Rapid visco analyzer measurements of japonica rice cultivars to study interrelationship between pasting properties and farming system. International Journal of Agronomy 2016, 3595326.

Kurasawa, H., Kanauti, Y., Takei, K., Ogawa, S., Okabe, T., Hayakawa, T., Igaue, I., Chemistry, B., 1972. Correlation analysis between eating quality, rheological property and amylose content of starch. Agricultural and Biological Chemistry 36, 1809-1813.

Li, M., Li, D., Feng, F., Zhang, S., Ma, F., Cheng, L., 2016. Proteomic analysis reveals dynamic regulation of fruit development and sugar and acid accumulation in apple. Journal of Experimental Botany 67, 5145-5157.

Liu, K., Li, T., Chen, Y., Huang, J., Qiu, Y., Li, S., Wang, H., Zhu, A., Zhuo, X., Yu, F., Zhang, H., Gu, J., Liu, L., Yang, J., 2020. Effects of root morphology and physiology on the formation and regulation of large panicles in rice. Field Crops Research 258, 107946.

Liu, W., Yin, T., Zhao, Y., Wang, X., Wang, K., Shen, Y., Ding, Y., Tang, S., 2021. Effects of high temperature on rice grain development and quality formation based on proteomics comparative analysis under field warming. Frontiers in plant science 12, 746180.

Luo, H., Wang, Z., Chen, Y., Zheng, A., Chen, Y., Du, P., Mao, T., Meng, S., Tang, X., 2018. The effects of
different temperatures on the biosynthesis of grain protein in rice at filling stage. Applied Ecology and
Environmental Research 16, 8017-8027.

MacNeill, G., Mehrpouyan, S., Minow, M., Patterson, J., Tetlow, I., Emes, M., 2017. Starch as a source, starch
as a sink: the bifunctional role of starch in carbon allocation. Journal of Experimental Botany 68, 4433-4453.

Masumoto, C., Miyazawa, S.-I., Ohkawa, H., Fukuda, T., Taniguchi, Y., Murayama, S., Kusano, M., Saito, K.,
Fukayama, H., Miyao, M., 2010. Phosphoenolpyruvate carboxylase intrinsically located in the chloroplast of
rice plays a crucial role in ammonium assimilation. Proceedings of the National Academy of Sciences 107,
5226-5231.

Nasielski, J., Deen, B., 2019. Nitrogen applications made close to silking: Implications for yield formation in
maize. Field Crops Research 243, 107621.

Naz, M.Y., Sulaiman, S.A., 2016. Slow release coating remedy for nitrogen loss from conventional urea: a
review. Journal of Controlled Release 225, 109-120.

Ning, P., Yang, L., Li, C., Fritschi, F.B., 2018. Post-silking carbon partitioning under nitrogen deficiency
revealed sink limitation of grain yield in maize. Journal of Experimental Botany 69, 1707-1719.

Nunes-Nesi, A., Fernie, A.R., Stitt, M., 2010. Metabolic and signaling aspects underpinning the regulation of
plant carbon nitrogen interactions. Molecular Plant 3, 973-996.

Poucet, T., González-Moro, M.B., Cabasson, C., Beauvoit, B., Gibon, Y., Dieuaide-Noubhani, M., Marino, D.,
2021. Ammonium supply induces differential metabolic adaptive responses in tomato according to leaf
phenological stage. Journal of Experimental Botany 72, 3185-3199.

Richards, F., 1959. A flexible growth function for empirical use. Journal of Experimental Botany 10, 290-301.

Sekhar, S., Kumar, J., Mohanty, S., Mohanty, N., Panda, R.S., Das, S., Shaw, B.P., Behera, L., 2021.
Identification of novel QTLs for grain fertility and associated traits to decipher poor grain filling of basal
spikelets in dense panicle rice. Scientific Reports 11, 13617.

Seung, D., Thalmann, M., Sparla, F., Abou Hachem, M., Lee, S.K., Issakidis-Bourguet, E., Svensson, B.,
Zeeman, S.C., Santelia, D., 2013. Arabidopsis thaliana *AMY3* is a unique redox-regulated chloroplastic α-
amylase. Journal of Biological Chemistry 288, 33620-33633.

Shaviv, A., 2001. Advances in controlled-release fertilizers. Advances in Agronomy 71, 1-49.

Sheehy, J.E., Dionora, M.J.A., Mitchell, P.L., 2001. Spikelet numbers, sink size and potential yield in rice. Field
Crops Research 71, 77-85.

Shi, S., Zhang, G., Chen, L., Zhang, W., Wang, X., Pan, K., Li, L., Wang, J., Liu, J., Cao, C., Jiang, Y., 2023.
Different nitrogen fertilizer application in the field affects the morphology and structure of protein and starch in
rice during cooking. Food Research International 163, 112193.

Tang, S., Chen, W., Liu, W., Zhou, Q., Zhang, H., Wang, S., Ding, Y., 2018. Open-field warming regulates the
morphological structure, protein synthesis of grain and affects the appearance quality of rice. Journal of Cereal
Science 84, 20-29.

Tang, S., Zhang, H., Liu, W., Dou, Z., Zhou, Q., Chen, W., Wang, S., Ding, Y.J.F.C., 2019. Nitrogen fertilizer at
heading stage effectively compensates for the deterioration of rice quality by affecting the starch-related
properties under elevated temperatures. Food Chemistry 277, 455-462.

Thakur, A.K., Rath, S., Patil, D.U., Kumar, A., 2011. Effects on rice plant morphology and physiology of water
and associated management practices of the system of rice intensification and their implications for crop
performance. Paddy and Water Environment 9, 13-24.

Wang, E., Wang, J., Zhu, X., Hao, W., Wang, L., Li, Q., Zhang, L., He, W., Lu, B., Lin, H., Ma, H., Zhang, G.,
He, Z., 2008. Control of rice grain-filling and yield by a gene with a potential signature of domestication. Nature

Genetics 40, 1370-1374.

Wang, F., Peng, S.B., 2017. Yield potential and nitrogen use efficiency of China's super rice. Journal of
Integrative Agriculture 16, 1000-1008.

Wang, M., Oge, L., Perez Garcia, M.D., Launay-Avon, A., Clement, G., Le Gourrierec, J., Hamama, L., Sakr, S.,
2022a. Antagonistic Effect of Sucrose Availability and auxin on rosa axillary bud metabolism and signaling,
based on the transcriptomics and metabolomics analysis. Frontiers in Plant Science 13, 830840.

Wang, M., Perez-Garcia, M.D., Daviere, J.M., Barbier, F., Oge, L., Gentilhomme, J., Voisine, L., Peron, T.,
Launay-Avon, A., Clement, G., Baumberger, N., Balzergue, S., Macherel, D., Grappin, P., Bertheloot, J.,
Achard, P., Hamama, L., Sakr, S., 2021a. Outgrowth of the axillary bud in rose is controlled by sugar
metabolism and signalling. Journal of Experimental Botany 72, 3044-3060.

Wang, W., Wang, W., Pan, Y., Tan, C., Li, H., Chen, Y., Liu, X., Wei, J., Xu, N., Han, Y., Gu, H., Ye, R., Ding,
Q., Ma, C., 2022b. A new gain-of-function *OsGS2*/*GRF4* allele generated by CRISPR/Cas9 genome editing
increases rice grain size and yield. The Crop Journal 10, 1207-1212.

Wang, X., Wang, K., Yin, T., Zhao, Y., Liu, W., Shen, Y., Ding, Y., Tang, S., 2021b. Nitrogen fertilizer regulated
grain storage protein synthesis and reduced chalkiness of rice under actual field warming. Frontiers in Plant
Science 12, 715436.

Wu, Q., Wang, Y., Ding, Y., Tao, W., Gao, S., Li, Q., Li, W., Liu, Z., Li, G., 2021. Effects of different types of
slow- and controlled-release fertilizers on rice yield. Journal of Integrative Agriculture 20, 1503-1514.

Xin, W., Zhang, L., Zhang, W., Gao, J., Yi, J., Zhen, X., Li, Z., Zhao, Y., Peng, C., Zhao, C., 2019. An integrated
analysis of the rice transcriptome and metabolome reveals differential regulation of carbon and nitrogen
metabolism in response to nitrogen availability. International Journal of Molecular Science 20, 2349.

Xu, C., Yang, F., Tang, X., Lu, B., Li, Z., Liu, Z., Ding, Y., Ding, C., Li, G., 2021. Super rice with high sink
activities has superior adaptability to low filling stage temperature. Frontiers in Plant Science 12, 729021.

Xu, C., Lu, B., Liang, L., Yang, F., Ding, C., Yan, F., Zhou, Y., Jiang, Z., Liu, Z., Ding, Y., Li, W., Li, G., 2022.
Delayed sowing date improves rice cooking and taste quality by regulating the quantity and quality of grains
located on secondary branches. Agronomy Basel 12, 1316.

Xu, G., Fan, X., Miller, A.J., 2012. Plant nitrogen assimilation and use efficiency. Annual review of plant
biology 63, 153-182.

Xu, X., Ma, H., Nin, Y., Wang, J., Zhang, Y., 2016. Effects of slow-released nitrogen fertilizers with different
application patterns on crop yields and nitrogen fertilizer use efficiency in rice-wheat rotation system. Journal of
Plant Nutrition and Fertilizer Science 22, 307-316.

Yamakawa, H., Hakata, M., 2010. Atlas of rice grain filling-related metabolism under high temperature: joint
analysis of metabolome and transcriptome demonstrated inhibition of starch accumulation and induction of
amino acid accumulation. Plant and Cell Physiology 51, 795-809.

Yang, J., Zhang, J., Wang, Z., Zhu, Q., 2001. Activities of starch hydrolytic enzymes and sucrose-phosphate
synthase in the stems of rice subjected to water stress during grain filling. Journal of Experimental Botany 52,
2169-2179.

Yang, J., Zhang, J., Wang, Z., Zhu, Q., Liu, L., 2003. Activities of enzymes involved in sucrose-to-starch
metabolism in rice grains subjected to water stress during filling. Field Crops Research 81, 69-81.

Yang, X., Nian, J., Xie, Q., Feng, J., Zhang, F., Jing, H., Zhang, J., Dong, G., Liang, Y., Peng, J., Wang, G.,
Qian, Q., Zuo, J., 2016. Rice Ferredoxin-dependent glutamate synthase regulates nitrogen–carbon metabolomes
and is genetically differentiated between japonica and indica subspecies. Molecular Plant 9, 1520-1534.

Yang, Y., Zhang, M., Zheng, L., Cheng, D., Liu, M., Geng, Y., 2011. Controlled release urea improved nitrogen
use efficiency, yield, and quality of wheat. Agronomy Journal 103, 479-485.

Yasuda, H., Tada, Y., Hayashi, Y., Jomori, T., Takaiwa, F., 2005. Expression of the small peptide GLP-1 in
transgenic plants. Transgenic Research 14, 677-684.

Ye, Y., Liang, X., Chen, Y., Liu, J., Gu, J., Guo, R., Li, L., 2013. Alternate wetting and drying irrigation and
controlled-release nitrogen fertilizer in late-season rice. Effects on dry matter accumulation, yield, water and
nitrogen use. Field Crops Research 144, 212-224.

Zhang, L., Sun, S., Liang, Y., Li, B., Ma, S., Wang, Z., Ma, B., Li, M., 2021. Nitrogen levels regulate sugar
metabolism and transport in the shoot tips of crabapple plants. Frontiers in plant science 12, 626149.

Zhao, M., Zhao, M., Gu, S., Sun, J., Ma, Z., Wang, L., Zheng, W., Xu, Z., 2019. DEP1 is involved in regulating
the carbon–nitrogen metabolic balance to affect grain yield and quality in rice (*Oriza sativa* L.). Plos one 14,
e0213504.

Jin, Z., Jiang W., Chin, J., Koh, H., 2004. Analysis on the combining ability of taste meter value and starch RVA
properties in Indica rice. Europe PMC 30, 1210-1214.

Zhou, C., Huang, Y., Jia, B., Wang, Y., Wang, Y., Xu, Q., Li, R., Wang, S., Dou, F., 2018. Effects of cultivar,
nitrogen rate, and planting density on rice-grain quality. Agronomy 8, 246.

Tables:

Table 1. Agronomic traits at the maturity stage under different types of slow-controlled release fertilizer in 2020 and 2021.

Table 2. Effects of different types of slow-controlled release fertilizer on grain quality of rice in 2020 and 2021.

Table 3. Carbon and nitrogen metabolites at the maturity stage under different types of slow-controlled release fertilization in 2020 and 2021.

Figures:

Figure 1. Nitrogen release rate (A) and appearance feature (B) of different types of slow-controlled release fertilizer.

Figure 2. Dynamics of rice grain growth after the heading stage under different types of slow-controlled release fertilizer in 2020 and 2021.

Figure 3. Effects of different types of slow-controlled release fertilizer on the changes in carbohydrates in rice grain during the filling stage in 2020 and 2021.

Figure 4. Effects of different types of slow-controlled release fertilizer on the activity of starch synthesis-related enzymes in rice grain during the filling stage in 2020 and 2021.

Figure 5. Effects of different types of slow-controlled release fertilizer on the content of total protein and amino acids in rice grain during the filling stage in 2020 and 2021.

Figure 6. Effects of different types of slow-controlled release fertilizer on changes in nitrogen metabolism-related enzyme activities in rice grain during the filling stage in 2020 and 2021.

Figure 7. Metabolic profiling corresponding to the accumulation of starch and amino acids in rice grains during the filling stage in 2021.

Year	Fertilizer type	Yield (t hm^{-2})	Panicles (No./m^2)	Spikelets (No./panicle)	Spikelets (No./hm^2)	Seed setting rate (%)	Grain weight (mg)
	CK	10.2ab	314.4ab	129.0a	4.1a	93.9a	26.9c
	UF	8.4c	286.2c	112.3b	3.2c	94.9a	27.6ab
2020	SCU	9.6b	305.5abc	117.5b	3.6bc	95.4a	28.0a
	PCU	10.2ab	293.3bc	131.8a	3.9ab	94.9a	27.8a
	BBF	10.8a	325.3a	131.2a	4.3a	93.6a	27.2bc
	CK	11.2a	323.0a	137.9a	4.4a	96.8a	26.1c
	UF	9.7b	335.7a	110.5c	3.7b	93.6a	28.3a
2021	SCU	8.6b	343.3a	94.2c	3.2c	94.9a	28.1ab
	PCU	11.1a	368.3a	116.0abc	4.3a	96.3a	27.2b
	BBF	11.0a	355.7a	117.3ab	4.2a	95.6a	27.5ab
F-Value	T	**	ns	**	**	ns	**
	Y	**	**	*	*	ns	ns
	T×Y	**	ns	ns	**	ns	*

656 CK, conventional fertilization with four spilt applications of urea; UF, urea formaldehyde; SCU, sulfur-
657 coated urea; PCU, polymer-coated urea; BBF, controlled-release bulk blending fertilizer; Values with
658 different letters in the same column are significantly different at $P \leq 0.05$ by the LSD test;
659 **and*significant at 0.01and 0.05 probability level respectively; ns, not significant.

Table 2. Effects of different types of slow-controlled release fertilizer on grain quality of rice in 2020 and 2021.

Year	Fertilizer type	Brown rice rate (%)	Milled rice rate (%)	Head rice rate (%)	Chalky kernel rate (%)	Chalky area (%)	Chalkiness (%)	Taste value
2020	CK	85.06a	74.91a	70.19a	39.67a	22.52ab	8.88ab	72.7a
	UF	84.23b	74.5a	69.22a	34.17a	20.80b	7.17b	73.0a
	SCU	84.63ab	74.78a	69.8a	34.83a	21.72ab	7.61ab	72.3a
	PCU	84.9a	75.01a	70.45a	41.00a	26.00a	10.64a	65.0b
	BBF	84.8a	75.3a	70.46a	34.83a	20.46b	7.11b	71.3a
2021	CK	84.10a	72.53a	68.47a	37.46ab	24.65b	9.18b	75.00a
	UF	83.82a	73.46a	69.25a	32.99b	21.71c	7.15d	76.67a
	SCU	83.72a	74.23a	71.64a	36.98ab	22.56c	8.34bc	76.00a
	PCU	83.46a	72.79a	69.31a	42.41a	27.07a	11.47a	73.00b
	BBF	83.8a	73.64a	69.97a	34.80a	22.31c	7.75cd	76.33a

Year	Fertilizer type	Peak viscosity (cP)	Hot viscosity (cP)	Cool viscosity (cP)	Breakdown (cP)	Setback (cP)	Consistency (cP)	Pasting temperature (°C)	Peak time (min)
2020	CK	3270b	2249ab	2820ab	1021b	450bc	570ab	72.1a	6.6a
	UF	3390a	2187b	2858ab	1204a	532a	671a	71.9a	6.4a
	SCU	3342ab	2209ab	2913a	1133ab	429c	704a	71.9a	6.5a
	PCU	3170c	2298a	2745c	872c	425c	447b	72.1a	6.6a
	BBF	3277b	2194b	2773bc	1082b	504ab	579ab	72.0a	6.4a
2021	CK	3527c	2372a	3051a	1355d	476d	679a	75.0a	6.2a
	UF	3695a	2334a	3060a	1559a	601c	726a	74.2a	6.0a
	SCU	3711a	2281a	2991a	1464b	720a	711a	73.9a	6.1a
	PCU	3529c	2152b	2837b	1303e	692ab	685a	74.4a	6.1a
	BBF	3628b	2302a	3007a	1403c	622bc	705a	74.2a	6.1a

666　CK, conventional fertilization with four spilt applications of urea; UF, urea formaldehyde; SCU, sulfur-coated urea; PCU, polymer-coated urea; BBF, controlled-
667　release bulk blending fertilizer; Values with different letters in the same column are significantly different with $P \leq 0.05$.

Table 3. Carbon and nitrogen metabolites at the maturity stage under different types of slow-controlled release fertilization in 2020 and 2021.

Year	Fertilizer type	Protein (%)	Starch （%）	Amino acids (%)	Albumin (%)	Globulin (%)	Gliadin (%)	Glutenin (%)	Amylose （%）	Amylopectin （%）
2020	CK	8.42a	81.36a	7.16b	0.48a	0.55b	0.67bc	6.71ab	11.89a	69.47a
	UF	7.66b	82.51a	7.21b	0.47a	0.49c	0.62d	6.08c	11.89a	70.62a
	SCU	7.85b	82.27a	6.62c	0.46a	0.51c	0.65c	6.24bc	12.12a	70.16a
	PCU	8.86a	79.35b	8.05a	0.49a	0.63a	0.70ab	7.05a	12.09a	67.26b
	BBF	8.39a	81.28a	7.37b	0.50a	0.64a	0.71a	6.54abc	12.09a	69.19a
2021	CK	9.21a	78.77ab	8.26a	0.51a	0.50a	0.75a	7.46a	9.26b	69.51a
	UF	8.24b	79.08a	7.71b	0.46b	0.45b	0.62c	6.71b	9.25b	69.83a
	SCU	8.78ab	78.46ab	7.85b	0.45b	0.47ab	0.65c	7.20ab	9.90a	68.56b
	PCU	8.78ab	77.52c	8.15a	0.53a	0.48ab	0.75a	7.02ab	10.03a	67.49c
	BBF	8.45b	78.24bc	7.77b	0.47b	0.47ab	0.69b	6.82ab	9.06b	69.18ab
F-Value	T	**	**	**	**	**	**	**	**	**
	Y	**	**	**	ns	**	**	**	**	ns
	T×Y	Ns	ns	**	*	**	**	ns	*	ns

CK, conventional fertilization with four spilt applications of urea; UF, urea formaldehyde; SCU, sulfur-coated urea; PCU, polymer-coated urea; BBF, controlled-release bulk blending fertilizer; Values with different letters in the same column are significantly different at $P \leq 0.05$ by the LSD test; **and*significant at 0.01and 0.05 probability level respectively; ns, not significant.

Figure 1. Nitrogen release rate (A) and appearance feature (B) of different types of slow-controlled release fertilizer. CK, conventional fertilization with four spilt applications of urea; UF, urea formaldehyde; SCU, sulfur-coated urea; PCU, polymer-coated urea; BBF, controlled-release bulk blending fertilizer.

Figure 2. Dynamics of rice grain growth after the heading stage under different types of slow-controlled release fertilizer in 2020 and 2021. CK, conventional fertilization with four spilt applications of urea; UF, urea formaldehyde; SCU, sulfur-coated urea; PCU, polymer-coated urea; BBF, controlled-release bulk blending fertilizer.

694 **Figure 3. Effects of different types of slow-controlled release fertilizer on the changes in carbohydrates in rice grain during the filling stage in 2020 and**
695 **2021.**

696 CK, conventional fertilization with four spilt applications of urea; UF, urea formaldehyde; SCU, sulfur-coated urea; PCU, polymer-coated urea; BBF, controlled-
697 release bulk blending fertilizer; Significant differences at each time point are indicated by different letters ($P < 0.05$) as determined by Duncan's test.

698
699
Figure 4. Effects of different types of slow-controlled release fertilizer on the activity of starch synthesis-related enzymes in rice grain during the filling stage
in 2020 and 2021. CK, conventional fertilization with four spilt applications of urea; UF, urea formaldehyde; SCU, sulfur-coated urea; PCU, polymer-coated urea;
BBF, controlled-release bulk blending fertilizer; Significant differences at each time point are indicated by different letters ($P < 0.05$) as determined by Duncan's test.

Figure 5. Effects of different types of slow-controlled release fertilizer on the content of total
protein and amino acids in rice grain during the filling stage in 2020 and 2021. CK,
conventional fertilization with four spilt applications of urea; UF, urea formaldehyde; SCU, sulfur-
coated urea; PCU, polymer-coated urea; BBF, controlled-release bulk blending fertilizer; Profiling
of total amino acids presented as a heat map calculated by $\log_2$ fold change (red meaning
increased and blue meaning decreased); Significant differences at each time point are indicated by
different letters ($P < 0.05$) as determined by Duncan's test.

Figure 6. Effects of different types of slow-controlled release fertilizer on changes in nitrogen

metabolism-related enzyme activities in rice grain during the filling stage in 2020 and 2021.

CK, conventional fertilization with four spilt applications of urea; UF, urea formaldehyde; SCU,

sulfur-coated urea; PCU, polymer-coated urea; BBF, controlled-release bulk blending fertilizer;

Significant differences at each time point are indicated by different letters ($P < 0.05$) as determined

by Duncan's test.

Figure 7. Metabolic profiling corresponding to the accumulation of starch and amino acids
in rice grains during the filling stage in 2021. CK, conventional fertilization with four spilt
applications of urea; UF, urea formaldehyde; SCU, sulfur-coated urea; PCU, polymer-coated urea;
BBF, controlled-release bulk blending fertilizer; DPA, days post anthesis; Values of the heat map
calculated by $\log_2$ fold change (red meaning increased and blue meaning decreased).

Supporting Information

Article title: **Application of slow-controlled release fertilizer coordinates the carbon flow in**
carbon-nitrogen metabolism to effect rice quality.

Authors: Zhengrong Jiang, Qiuli Chen, Dun Liu, Weike Tao, Shen Gao, Jiaqi Li, Chunhao Lin,
Meichen Zhu, Yanfeng Ding, Weiwei Li, Ganghua Li, Soulaiman Sakr, Xueli Hong*

The following Supporting Information is available for this article:

Table S1. Sequences of primers for Actin and genes for qRT-PCR.

Table S2. Principal components analysis about the effect of slow-controlled release fertilizer on
key enzymes activities related carbon-nitrogen metabolism in rice grain at filling stage in 2020
and 2021.

Table S3. Principal components analysis of comprehensive index and loading matrix of each
component for amino acids accumulation under the application of slow-controlled release
fertilizers in 2020 and 2021.

Figure S1. Daily temperature (line) and precipitation (bars) during the rice growth seasons in
2020 and 2021.

Figure S2. Principal component analysis about the effect of slow-controlled release fertilizer
application on the accumulation of amino acids in rice grains at the filling stage in 2020 and 2021.

Table S1. Sequences of primers for Actin and genes for qRT-PCR.

Gene	Forward primer 5′ → 3'	Reverse primer 5′ → 3'
Actin	CAATCGTGAGAAGATGACCC	GTCCATCAGGAAGCTCGTAGC
OsSUT2	TCTTTTATCGGTGGGCTGGT	TTGCAAAGAATGGCCGACAA
OsCIN4	CATGTTTGTGCCGGATACTG	TGCCATCACCATCTAACCAA
OsGS2	AGGATCGGACAAATCGTTTGG	GCATGACCTCTCCATTTGTTCC

755 **Table S2. Principal components analysis about the effect of slow-controlled release fertilizer**
756 **on key enzymes activities related carbon-nitrogen metabolism in rice grain at filling stage in**
757 **2020 and 2021.**

Traits	PC1	PC2	PC3	PC4	PC5	PC6
SuSase	0.44	-0.24	-0.12	-0.34	0.61	-0.50
ADPGase	0.41	-0.42	0.17	0.61	-0.37	-0.34
GBSS	0.44	-0.22	0.31	0.07	0.31	0.75
SBE	0.44	-0.09	-0.32	-0.56	-0.61	0.14
GS	0.37	0.53	-0.61	0.43	0.14	0.10
GOGAT	0.33	0.66	0.62	-0.13	-0.11	-0.20
Eigenvalue	4.62	0.88	0.28	0.11	0.08	0.03
Contributive ratio (%)	76.93	14.74	4.66	1.86	1.27	0.54
Cumulative (%)	76.93	91.67	96.33	98.19	99.46	100.00

Table S3. Principal components analysis of comprehensive index and loading matrix of each component for amino acids accumulation under the application of slow-controlled release fertilizers in 2020 and 2021.

Traits	PC1	PC2	PC3	PC4	PC5	PC6	PC7	PC8	PC9	PC10	PC11	PC12
Asp	0.30	0.13	0.14	-0.49	0.22	-0.16	0.35	0.28	-0.16	-0.57	-0.08	-0.05
Thr	0.28	-0.39	0.08	0.08	0.00	0.62	-0.09	0.17	-0.50	-0.04	0.30	0.06
Ser	0.25	0.31	-0.47	0.19	0.21	0.48	0.17	0.08	0.50	-0.13	0.05	0.06
Glu	0.34	0.26	-0.02	-0.24	0.17	0.04	0.28	-0.07	-0.25	0.73	-0.21	0.01
Gly	0.35	-0.08	0.09	-0.21	-0.09	0.17	-0.32	-0.49	0.16	-0.11	-0.22	-0.59
Val	0.36	-0.14	0.12	-0.20	-0.10	-0.04	-0.27	-0.20	0.23	-0.04	-0.17	0.77
Ile	0.32	-0.32	0.09	0.01	0.06	-0.26	-0.19	0.61	0.40	0.28	0.13	-0.22
Leu	0.30	0.39	-0.15	-0.04	-0.28	-0.29	-0.15	-0.15	-0.12	-0.01	0.71	-0.01
Tyr	0.27	-0.02	-0.55	0.31	-0.24	-0.26	-0.16	0.17	-0.36	-0.13	-0.46	-0.02
Phe	0.13	0.45	0.56	0.31	-0.48	0.19	0.04	0.24	0.04	-0.04	-0.21	-0.02
Lys	0.20	0.14	0.29	0.52	0.67	-0.21	-0.20	-0.17	-0.13	-0.09	0.02	0.02
His	0.26	-0.40	0.08	0.33	-0.21	-0.18	0.68	-0.31	0.13	-0.03	0.08	-0.02
Eigenvalue	6.57	1.47	0.99	0.89	0.74	0.47	0.29	0.19	0.19	0.08	0.07	0.04
Contributive ratio (%)	54.75	12.25	8.29	7.44	6.19	3.93	2.45	1.59	1.55	0.70	0.55	0.32
Cumulative (%)	54.75	67.00	75.29	82.73	88.92	92.84	95.29	96.89	98.44	99.13	99.68	100.00

Figure S1. Daily temperature (line) and precipitation (bars) during the rice growth seasons in 2020
and 2021.

Figure S2. Principal component analysis about the effect of slow-controlled release fertilizer
application on the accumulation of amino acids in rice grains at the filling stage in 2020 and 2021.
CK, conventional fertilization with four spilt applications of urea; UF, urea formaldehyde; SCU, sulfur-
coated urea; PCU, polymer-coated urea; BBF, controlled-release bulk blending fertilizer.